



Cite this: *RSC Adv.*, 2018, 8, 30080

High throughput mass spectrometry-based characterisation of *Arabidopsis thaliana* group H glycosyltransferases†

Aishat Akere,^a Qian Liu,^b Shibo Wu,^a Bingkai Hou^b and Min Yang^{a*}

In this report, we cloned and characterised four members of group H glycosyltransferases (GTs) by studying their substrate specificities and kinetics. The formation of products and possible glycosylation position was confirmed using MS/MS. The results revealed that 76E1 and 76E5 have broader donor specificity, including UDP-glucose (UDPGlc), UDP-galactose (UDPGal) and UDP-*N*-acetylglucosamine (UDPGlcNAc) with various flavonoids as acceptor substrates. Pseudo-single substrate kinetics data showed a relatively low K_M , indicating a high affinity for substrate UDPGlc and also supported that 76E5 is more of a galactosyl and *N*-acetylglucosamine transferase. Sequence alignment and site-directed mutagenesis studies indeed suggested that serine is a crucial residue in the UDPGlcNAc and UDPGal activity.

Received 8th May 2018
 Accepted 6th August 2018

DOI: 10.1039/c8ra03947j

rsc.li/rsc-advances

Introduction

Glycosylation is a widespread modification of plant secondary metabolites involved in various functions such as the regulation of hormone homeostasis, the detoxification of xenobiotics and the biosynthesis and storage of secondary compounds. In plants, these reactions are controlled by a specific subclass of the glycosyltransferase family. Family 1 plant UDP dependent glycosyltransferases (UGTs) are responsible for the transfer of sugar to many bioactive natural products¹ such as alkaloids, terpenoids, flavonoids and phenylpropanoids. UDP activated sugars are mainly used as donor compounds by UGTs.² Generally, transfer of sugar molecules can be either to macromolecules (such as proteins, lipids) or small molecules (such as plant secondary metabolites – natural products, oligosaccharides).³ However, the products of glycosylation catalysed by UGTs are glycosides of small molecules.⁴

Numerous pharmacological activities have been reported for natural products containing sugar residues. This is because glycosylation improves their stability, solubility in water and bioavailability; hence it has become an important process in drug research and development.⁵ In addition, through this process, glycosides have become attractive compounds used as nutraceuticals and food additives.³ Leaves and seeds of *Digitalis purpurea* and *Digitalis lanata* are the most significant sources of

cardiac steroids important in the treatment of various heart conditions. The aglycone steroid however, binds and dissociates too quickly from the receptor. Attachment of the sugar moiety to the aglycone steroid improves its binding stability to receptor, though the sugar itself has no cardiac activity. Hence, sugar addition enhanced the pharmacokinetic behaviour of the aglycone steroid, leaving us with low dissociation potent glycosides.⁶ New and effective methods to produce glycosylated natural products have been studied over the years.⁷ Since glycosyltransferases are important biocatalysts in sugar addition, it is imperative to study UGTs more in depth.

Presently, 122 UGT encoding genes are found in the model plant *Arabidopsis thaliana*.⁸ Phylogenetic analysis of conserved amino acids in *Arabidopsis* UGTs led to the classification into 14 groups⁹ (ESI Fig. 1†). Although these enzymes have been studied for many years, to date only a few have been characterized in planta. Of the 19 genes found in group H, only 2 are fully characterised.⁸ Varieties of glycosides found in plants *in vivo* signify that plant UGTs can glycosylate diverse compounds.¹⁰ Although UGTs were believed to be highly specific in substrate recognition *in vivo*, reports have disclosed broader range promiscuity against both substrates.^{11–17} This substrate recognition includes recognition of both UDP sugar donor as well as acceptor.¹ Generally, UGTs are highly specific for sugar donor although UDP-glucose is most commonly used.¹⁸ The screening studies^{11–17} which looked into substrate specificity within *Arabidopsis thaliana* phylogenetic groups found that UGTs of the same group glycosylate acceptors belonging to very different compound classes. These early studies challenged efforts to solely base substrate specificity on phylogeny. Although biochemical characterisation of the substrate specificity is quite challenging as it entails testing diverse substrates to fully understand specificity of individual UGTs, it will advance

^aThe School of Pharmacy, University College London, 29-39 Brunswick Square, London WC1N 1AX, UK. E-mail: min.yang@ucl.ac.uk

^bKey Laboratory of Plant Cell Engineering and Germplasm Innovation, Ministry of Education of China, School of Life Sciences, Shandong University, Jinan, 250100, China

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c8ra03947j



prediction of substrate specificity based on phylogeny.^{1,19} Comprehensive facts on the *in vivo* and *in vitro* activity and specificity of UGTs are important in the design of enhanced UGTs with desired properties.²⁰

In recent times, the importance of glycosyltransferases (GTs) has been in the spotlight. However, insufficient biochemical data on individual member enzyme has hindered further research into their functional understanding. Identifying and biochemically characterising some GT genes has helped in studying mechanism of glycosylation *in planta*.²¹ Biochemical data on substrate preferences has been relied on for prediction of substrate specificity based on phylogeny. Till date, most of *Arabidopsis thaliana* UGTs substrates remain unknown.²²

Here, we report the substrate specificity and kinetic analysis of four *Arabidopsis thaliana* group H UGTs (76E1, 76E2, 76E5 and 76D1). Substrate screening showed that the UGTs added sugar to acceptors from different classes of secondary metabolites ranging from flavonoids, cinnamic acids, coumarins to alkaloids. They also recognised varying donor compounds, with all accepting UDPGlc. In addition, 76E1 and 76E5 recognise UDPGal and UDPGlcNAc. Using tandem mass spectrometry, likely glycosylation positions of the acceptors were proposed. Further kinetic study measured each enzyme binding affinity with donor substrates, indicating their donor preferences and catalytic efficiencies. Knowledge of the activity and specificity of UGTs would provide a basis for their potential use in the design of improved UGTs to produce bioactive molecules for pharmaceutical purpose.

Experimental

(a) Gene cloning procedure

Total RNA was extracted from the 14d seedlings of the *Arabidopsis* using the TRIzol reagent (Takara, Japan). First-strand cDNA was synthesized using PrimerScript RT reagent Kit with gDNA Eraser (Takara, Japan), pooled cDNA was used as PCR template. The cDNA sequences of *UGT76E1*, *UGT76E2*, *UGT76E5* and *UGT76D1* were amplified by *TransStart Fast Pfu* DNA Polymerase (*TRANS*, China). A *Bam*HI and *Xho*I restriction site were included in the sequence of forward primer and the reverse primer used for PCR of *UGT76E1* and *UGT76E2*. The PCR primers for *UGT76E1* cloning are: forward GGATCCATGGAA-GAACTAGGAGTGAAG and reverse CTCGAGCTA-CATGAAATTCATCATT; the PCR primers for *UGT76E2* cloning are: forward GGATCCATGGAGGAAAACAAGTGAAG, reverse CTCGAGTCACATGGAATTAACAAAGTC. The *Bam*HI and *Eco*RI were included in the sequence of forward primer and the reverse primer used for PCR of *UGT76E5*. The PCR primers for *UGT76E5* cloning are: forward GGATCCATGGAGAAAAATGCA-GAGAAG, reverse GAATTCTCAAGTATTTCTATACTCTGC. The *Bam*HI and *Sac*I were included in the sequence of forward primer and the reverse primer used for PCR of *UGT76D1*. The primers for *UGT76D1* cloning are: forward CGGGATCCATGG-CAGAGATTCGCCAG, reverse CGAGCTCTCATTGTTCTGTC-CAATTTGCATC. The amplified products were cloned into the corresponding sites of the Blunt simple cloning vector, previously digested with appropriate restriction enzymes. After

sequencing and alignment with reference sequence of the four genes shown in the *Arabidopsis* information resource, the right clones were chosen for further cloning. Lastly, the cDNA fragments of the target genes were inserted and ligated into the pGEX-3H plasmid previously digested with appropriate restriction enzymes, respectively. The resulting recombinant plasmids were transformed into competent *E. coli* BL cells.

(b) Expression and purification of AtUGT76E1, AtUGT76E2 and AtUGT76E5

The recombinant plasmids were transformed into BL1 (DE3) cells for protein expression. Thereafter, they were verified by DNA sequencing *via* sanger sequencing (Source Bioscience Ltd). The bacterial cells were grown at 37 °C while being shaken at 200 rpm to an OD₆₀₀ of 0.6–0.8. Isopropyl 1-thiogalactopyranoside (IPTG) was added to the cell culture to a final concentration of 0.1 mM to induce protein expression. The culture was grown overnight at 20 °C and collected by centrifugation at 4 °C. The GST-tagged recombinant protein was purified by affinity chromatography (columns) and quantified using the Bradford assay according to the standard procedure provided by the suppliers. Cell lysate before purification and purified protein were analysed on sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) gels to verify protein production (ESI Fig. 2a–d†).

(c) *In vitro* UGT reaction assay

The UGT enzyme assay included the following components: 1 mM Tris 1 mM, MgCl₂ (pH 8.0), 10 mM UDP-sugars, 10 mM acceptor compounds and purified target proteins. For the acceptor screening, the 42 compounds screened are shown in the acceptor library (ESI Table 1†). The 7 sugar donor compounds used in the donor screening are also indicated in Fig. 2. The reaction mixture was incubated at 37 °C for 3 and 15 hours at different times. The reaction was terminated with acetonitrile and centrifuged to remove proteins. The supernatant was then analysed with LCMS. Glycosylated products were identified by their molecular weights, and these target compounds were subsequently fragmented using MS/MS for confirmation.

(d) UDP-Glo™ glycosyltransferase assay – for kinetics

Kinetic study was performed using the UDP-Glo™ glycosyltransferases Assay kit following the procedure provided by the supplier. The UGT enzyme assay mixture for kinetics contained Tris buffer (50mM Tris, 5 mM MgCl₂, pH 7.5), kaempferol (for 76E1 and 76E5) and quercetin (for 76E2 and 76D1). Kaempferol and quercetin were fixed at 20 μM. The donor substrates concentration; UDP Glc (five concentrations) was varied from 10–500 μM while UDP Gal and GlcNAc were varied from 0.1–10 μM. The UGT reaction was carried out at 25 °C and terminated using the UDP-Glo assay detection buffer. This kit detects UDP generated after UDP-sugar transfer by converting UDP to light (measured in relative luminescence units) in a luciferase type reaction. A linear standard curve using 0–25 μM UDP was performed (ESI Fig. 17†). Following the kit's protocol, the UGT



assay was combined in a ratio 1 : 1 (10 μ l : 10 μ l) with the UDP-Glo™ detection reagent in individual wells (white flat bottom 384-well plate [Corning]). Luminescence signal was measured after 1 hour incubation using a Pherastar microplate reader (BMG Labtech).

(e) HPLC-MS/MS

Samples were analysed with Agilent 6400 triple quadrupole mass spectrometer coupled with HPLC system using phenomenex-C18 column (50 \times 4.6 mm, kinetex 5u, 100A). The solvents used were as follows: (A) HPLC grade water containing 0.1% formic acid and (B) acetonitrile containing 0.1% formic acid. For the MS scan, both positive and negative spectra were obtained and run at a flow rate of 0.4 ml min⁻¹ in isocratic mode (10% A and 90% B). The injection volume was 10 μ l, detection wavelength at 260 nm and column temperature of 20 °C. Other details include start and end mass of 100 and 1000 respectively, scan time of 500 and cell accelerator voltage of 7. The MS/MS (product ion scan) had a flow rate 0.5 ml min⁻¹ at gradient mode (70% A: 30% B for 1 min, 55% A: 45% B for 1.50 min and lastly 70% A: 30% B for 2.50 min). The injection volume was maintained at 10 μ l and only negative spectra were obtained here. The precursor ions were fragmented for confirmation within the range of *m/z* 100–100. Scan segment details are as follows: scan time 500 fragmentor 135, collision energy 15 and cell accelerator voltage 7. All analyses were done in duplicates.

(f) Site directed mutagenesis

Primer design was done using NEBaseChanger v1.2.7 (<http://nebasechanger.neb.com/>) and the primers synthesized by Eurofin Genomics Ltd. Templates DNAs were extracted using QIAprep Spin miniprep kit protocol. Q5 Site-directed mutagenesis kit protocol (from NewEngland Biolabs) was observed to make mutants from the wild type template DNAs. Mutagenesis was done in three stages of exponential amplification (polymerase chain reaction), digestion and transformation. Mutant plasmid DNAs were subsequently extracted *via* miniprep. The mutant DNAs were sent for sanger sequencing (SourceBio-Science) to confirm mutations. The primer sequences are: 76E2N320S: Forward primer AGAGGAATTCAGTAGGTTGGTTTC, reverse primer GGTAAGGACTCTGTCCATTC; 76E1S318N: forward primer GGAGGAATTC AATAGGTTGGTTT, reverse primer GGTAAGGACTCTGTCCATTC; 76E5S311N: forward primer AGTGGAAGTCAATAAGATTGTCTC, reverse primer GGCATTGACTCTGTACCG. All WT and mutant sequences were confirmed using BLAST (ESI Fig. 11–17†).

(g) Sequence alignments

The amino acid sequences of the wild type UGTs were aligned using clustal omega by European Bioinformatics Institute (<https://www.ebi.ac.uk/Tools/msa/clustalo/>). The mutant and wild type DNA sequences were compared using NCBI nucleotide BLAST (National Centre for Biotechnology Institute).

Results and discussion

(a) MS based HTS method

The genes encoding 76E family and 76D1 were expressed as recombinant proteins in *Escherichia coli* and purified to electrophoretic homogeneity. *In vivo*, the functions of these UGTs are not known except for 76D1 whose functions have been recently published.²³ The Green-Amber-Red 'GAR' screen^{11,24} was developed to probe potential substrates, *in vitro*, using a large acceptor and nucleotide-sugar donor library (Fig. 2, ESI Table 1†). A virtual colour, reflecting DNA micro-array practice, of Green-Amber-Red (GAR, green indicates positive, amber means not clear and red shows negative) is assigned reflecting the presence of selected ions of mass. This procedure using liquid chromatography triple quadrupole Mass Spectrometry (LC-MS/MS) was composed of two parts; a general MS scan to identify desired product peaks such as $[M - H]^-$ or $[M + Cl]^-$ in negative mode and $[M + H]^+$ or $[M + Na]^+$ in positive mode first, and then a MS/MS scan to confirm the presence of acceptor precursor which indicates the loss of the sugar moiety to confirm the reaction product (Fig. 1a). This method, similar to the previous NLCIT,¹¹ could be used to determine the glycosylation site. No difference was noticed in the reaction products from 3 h and 15 h assay incubation. In addition, pseudo single substrate enzyme kinetics was measured by varying one substrate concentrations while saturating the other substrate using a commercial UDP-Glo® assay. An appropriate enzyme concentration was obtained prior to the kinetic study with fixed substrate concentrations (UDPGlc at 100 μ M and KMP at 20 μ M) while enzyme concentration varied. The formation of UDP was monitored in the function of time and initial rates were obtained. Kinetics parameters such as V_{max}/K_{cat} and K_M were generated using either Lineweaver–Burk or Michaelis–Menten equation (ESI Fig. 13–16†).

(b) GAR substrate specificity

(1) Acceptor screening result. The enzyme activity towards acceptor compounds with UDPGlc as the donor using above MS/MS method was summarised in this GAR mode (Fig. 1a). The MS and MS/MS spectra were indicated in ESI Fig. 3–6.†

In this study, we have reported the acceptor substrate profile of UGTs 76E1, 76E2, 76E5 and 76D1 (Fig. 1b). Of the forty-two acceptor substrates screened, 76E1 recognised nine flavonoids and one coumarin. 76E2 only accepted flavonoids as substrates while 76E5 on the other hand, recognised one benzoic acid, one coumarin and two flavonoids. Similarly, 76D1 accepted two flavonoids and one benzoic acid. Although there seem to exist a flavonoid trend, recognition of benzoic acid by 76E5 and 76D1 and non-recognition of other acceptor substrate class reinforces the established finding that substrate recognition cannot be predicted solely based on phylogenetic grouping.¹

Our results showed that although acceptor compounds from varying classes were recognised, flavonoids appeared to be a common factor. According to Parajuli *et al.*,^{25,26} expanded substrate study of a glycosyltransferase *AtUGT89C1* revealed that it recognised five different classes of flavonoids and varying






Fig. 1 MS based method and GAR screening result. (a) The MS/MS method to determine the formation of product using an example of UDPGlcNAc with kaempferol catalyzed by UGT76E5 (inset: reaction scheme and the fragment of product, product structure for illustration purpose only). (b) GAR acceptor screening result of enzymes (c) structures of the compounds showed positive reaction in the acceptor screening.

NDP-sugars. Another *Arabidopsis thaliana* glycosyltransferases, AtUGT78D1 glycosylated several compounds of structural types; flavonoids, flavonoid glycosides and others.²⁶ Furthermore, Jones *et al.*¹⁰ reported flavonoid glucoside/rhamnoside production from UGT73C6. All these studies corroborated that AtUGTs seemed to generally favour flavonoids *O*-glycosylation. Flavonoid glycosides possess promising beneficial health activities²⁷ and regioselective glycosyltransferases offer to overcome limitations in their pharmacological potency.²⁸ This implies that UGTs tested in group H may be useful in producing useful flavonoids glycosides.

(2) **Donor screening result and glycosylation position.** The donor screening was performed with two acceptors, kaempferol and quercetin which were confirmed to be the substrate from the acceptor result (Fig. 1). The MS and MS/MS spectra were indicated in the ESI 7–10.[†] The spectral data is summarised in the GAR screen below (Fig. 2). ESI Table 2[†] highlighted the precursor and product ions used for MS/MS screen. It was noticed that UDPGlc was the most favourite donor but some enzymes did utilise more than one donor compound. 76E1 and 76E5 used UDP Gal and UDPGlcNAc in addition to UDP Glc. 76D1 was reported to use UDPGlc (and UDPXyl) in plant defence

in vivo by glucosylation of DHBAs.¹⁸ This explains why 76D1 only recognised UDPGlc of all donor sugar compounds we screened.

Flavonoids have multiple glycosylation position which can change the product physical and biological activity. Mass spectra have been found to give information on structural characterization, especially glycosylation position. Most sugar molecules in flavonoid *O*-glycosides bind at 3-, 7- or 4' positions (Fig. 1c), leaving others such as 5-OH which show very low activity to glycosylation.²⁹ Differentiating between 3-, 7- and 4' – positions can be quite problematic; however, MS has shown advantage to provide possible solutions. In the MS2 fragment of a glycosylated flavonoid, radical aglycone ions ($[M-H^+-H^+-Gly]^-$, 284/300) are found to be more abundant for flavonol 3-*O* glycosides than regular aglycone ions ($[M-H^+-Gly]^-$, 285/301) which leaves 284 (kaempferol) or 300 (quercetin) instead of 285 or 301 (Fig. 3a). The formation of a radical aglycone ion is due to the homolytic cleavage of the glycosidic bond between the *O*-linked sugar and the aglycone moiety.²⁹

The presence of peak 179 (designated as the diagnostic ion, Fig. 3b) in retrocyclisation pathway involving loss of B-ring indicates specificity of 3' or 4' derivatives in negative ion ESI-MS/MS.³⁰ Fig. 3 showed a few other examples of using the above




Fig. 2 Donor screening result. (A) UDPGlc, (B) UDPGal, (C) UDPGlcNAc, (D) GDPFuc, (E) GDPMan, (F) GDPGlc and (G) UDPMan.






Fig. 3 The MS/MS method to determine the formation of product and potentially glycosylation position. (a) An example to show the 3-O glycosylation – the formation of radical aglycone (283.8). (b) An example to show the glycosylation on 4'-O position – the formation of aglycone (284.9) and the presence of fragment of 179.0. (c) An example to show the 7-O glycosylation – the formation of aglycone (300.8) only.

rule to probe the glycosylation position. For UDPGlcNAc, peak 285 was more prominent, indicating glycosylation may not be at 3-O position. It will either be glycosylation at 7-O or 4'-O position. The presence of peak 179, a diagnostic ion for 4'-O glycosylation suggests that it might be kaempferol 4'-O glucoside (Fig. 3b). Another example showing a glycosylation position was seen in a quercetin glycoside. Here, the regular aglycone ion peak 301 is more prominent than radical aglycone ion 300, showing the glycosylation was not at the 3-O position. Since peak 179 is absent here, we infer that the product was a quercetin 7-O glucoside (Fig. 3c).

From the acceptor screening (Fig. 2) it indicated that 76E1 can be more flexible in accommodating acceptors (either flat or curved). The result indicates the glycosylation position can be on 7-OH but no further information can be assured without NMR data. Interestingly, 76E1 is also very flexible in donor binding pocket with three (3) UDP-sugars used as substrate, but all on the 7-O position. 76E2 performed similarly to 76E1, glucosylating quercetin at 7-OH position. Similar flexibility was demonstrated for 76E5 in its donor binding pocket as 76E1 although 76E1 can utilise little more acceptors than 76E5. When UDPGlc was used as donor, it glycosylated kaempferol at 3-OH but with UDPGlcNAc as donor, the glycosylation position is on 4' or 7-OH. The acceptor screening prediction is challenging as it involves non-flavonoids (Table 1).

(3) Protein sequence analysis and mutation studies. Identification of residues responsible for protein function *via* protein sequence analysis is an important and widely studied concern. This is because it guides experimental analysis and gives insight to function prediction. Multiple sequence alignment (MSA) can expose sites that are conserved which can indicate residues of functional importance involved in binding.³¹ Comparing the MSA of the four studied UGTs showed few differences which seems to be in line with the donor recognised (Fig. 4). Serine (at points 311 and 318 in 76E1 and 76E5 respectively) is present in both UGTs recognising more donor compounds; UDP Gal and UDPGlcNAc. This residue is absent in 76D1 and 76E2 with no UDP Gal and UDPGlcNAc activity. However, 76D1 and 76E2 both has asparagine (N) at this point (circled in Fig. 4a). In our previous research (unpublished), we observed UDP Gal and UDPGlcNAc activity for 76E11 and 76E12, both of which contain serine at the same point as shown in Fig. 4b. The presence of serine at this point in two of studied UGTs and also in two other previously screened UGTs with

UDPGal and UDPGlcNAc activity from the same family, and its absence in two UGTs with no UDPGal and UDPGlcNAc activity could point at something important. Mutation of serine (S) to asparagine (N) at this site in 76E1 and 76E5 was done to establish the importance of serine in the enzyme substrate recognition, particularly if UDPGal and UDPGlcNAc activity will be lost, reduced or retained. In addition, mutation of asparagine (N) to serine (S) was done to observe if UDPGal and UDPGlcNAc activity will be acquired. Hence, mutants 76E1 S318N, 76E5 S311N and 76E2 N320S were made.

Mass spectra from UGT assay catalysed by mutant enzymes 76E1 S318N, 76E5 S311N and 76E2 N320S are shown in ESI Fig. 18–24. † Total loss of glycosylation activity (UDPGlc, UDPGal and UDPGlcNAc) was observed in 76E1 S318N and 76E5 S311N as summarised in Fig. 4c. This indeed demonstrates the strong link between these activities and serine at this point. More interestingly, UDPGal and UDPGlcNAc activity was acquired by 76E2 N320S; the WT enzyme 76E2 only has UDPGlc activity (Fig. 4c). Although the exact mechanism by which serine acts is still unclear, this finding confirms our hypothesis that serine at this point is important in the UDPGlcNAc and UDPGal activity.

(4) Kinetics data. As shown in the plots Michaelis–Menten and Lineweaver–Burk (Table 2, ESI Fig. 26–29†), under the current condition, all the enzymes showed relatively low K_M at μM scale. A low K_M indicates that an enzyme requires a small amount of substrate to become saturated. Hence, the maximum velocity is reached at relatively low substrate concentrations kinetics data revealed relatively low K_M values, suggesting

Table 1 Enzymes, substrates and glycosylation positions

Enzymes	Donor	Acceptor	Peaks used in identification	Glycosylation position
76E1	UDPGlc	Kaempferol	285	7-OH
	UDPGlc	Quercetin	301	7-OH
	UDPGal	Kaempferol	285	7-OH
	UDPGlcNAc	Quercetin	301	7-OH
76E2	UDPGlc	Quercetin	301	7-OH
76E5	UDPGlc	Kaempferol	284	3-OH
	UDPGlc	Quercetin	300	3-OH
	UDPGal	Kaempferol	284	3-OH
	UDPGal	Quercetin	300	3-OH
	UDPGlcNAc	Kaempferol	285, 179	4'-OH
76D1	UDPGlcNAc	Quercetin	301	7-OH
76D1	UDPGlc	Quercetin	301	7-OH






Fig. 4 Sequence alignment of UDPGlcNAc glycosyltransferases and mutant activities. (a) Sequence alignment showing serine thought to be involved in *O*-galactosylation and *O*-GlcNAcylation. (b) Sequence alignment showing serine in other Group H UGTs, thought to be involved in *O*-galactosylation and *O*-GlcNAcylation, (c) mutants' donor activity results: comparison between wild type enzymes 76E5, 76E1 and 76E2 and mutant enzymes 76E5 S311N, 76E1 S318N and 76E2 N320S activity with donor compounds (A) UDPGlc, (B) UDPGal, (C) UDPGlcNAc. 76E2 N320S mutant newly acquired UDPGal and UDPGlcNAc activities.

Table 2 Kinetics study of GTs

Enzyme	Substrate varied	K_M (μM)	K_{cat} (s^{-1})	K_{cat}/K_M ($\text{s}^{-1} \text{mM}^{-1}$)
76E1	UDP glucose	10.5 ± 1.38	0.00832	0.79
76E2	UDP glucose	8.75 ± 1.38	0.0107	1.23
76E5	UDP glucose	8.79 ± 1.64	0.0194	2.21
	UDP galactose	0.57 ± 0.19	0.0130	22.8
76D1	UDPN-acetyl glucosamine	0.78 ± 0.28	0.0166	21.3
	UDP glucose	25.77 ± 8.82	0.00476	0.185

enzymes have good affinity for the substrates UDPGlc, UDPGal and UDPGlcNAc. All 76E1, E2 and E5 enzymes share very similar $K_{M\text{-UDPGlc}}$ values ($\sim 10 \mu\text{M}$), which is a measure of enzymes' substrate preference and catalytic efficiency. However, 76E5 showed strong preferences to both UDPGal and UDPGlcNAc, with low sub μM K_M and much higher K_{cat}/K_M ($>20 \text{ s}^{-1} \text{mM}^{-1}$). This kinetic data strongly suggests that 76E5 is probably more of UDPGlcNAc and UDPGal transferase than UDPglucosyltransferase. This is probably also the case *in vivo*. 76D1 has much high K_M values ($>25 \mu\text{M}$) and low K_{cat}/K_M ($0.185 \text{ s}^{-1} \text{mM}^{-1}$). 76C1 and 76C2 are the only group H members previously characterised, which are plant hormone GTs. The kinetic parameters are with respect to the plant hormone substrates; hence unsuitable for comparison with our studied UGTs.³² There is no other kinetics data available for group H members for comparison. When compared with VvGT1 (K_{cat}/K_M $0.124 \text{ s}^{-1} \text{mM}^{-1}$),¹⁰ our UGTs K_{cat}/K_M values are in the same range for substrate UDPGlc. 76E5 has a far higher K_{cat}/K_M with substrates UDP Gal and UDPGlcNAc (>2000 fold) than VvGT1; affirming that UDP Gal and UDPGlcNAc seems to be natural substrates to 76E5. Functional studies for 76E5 *in vivo* will reveal more in the near future.

Conclusion

Here, we have reported the substrate specificities and kinetics of four UGTs in group H of *Arabidopsis thaliana* (76E1, 76E2, 76E5 and 76D1). These UGTs added sugars to acceptors from different classes of secondary metabolites ranging from flavonoids, cinnamic acids, coumarins to alkaloids. They also

recognised varying donors tested, with all recognising UDP glucose and 76E1 and 76E5 binding to UDP-galactose and UDPGlcNAc in addition. Using tandem mass spectrometry, the likely glycosylation positions on the acceptors were proposed. Further kinetic study measured each enzyme binding affinity with donor substrates, indicating their donor preferences and catalytic efficiencies. The distinctive amino acid sequences of both UGTs recognising UDP Gal and UDP GlcNAc was observed. Mutagenesis highlighted residue serine, which seem to be important for UDPGlcNAc and UDPGal activity. Knowledge of the activity and specificity of UGTs would provide a basis for their potential use to design improved UGTs to produce bioactive molecules for pharmaceutical purposes.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

This work was funded by the Nigerian Government, under the Presidential Special Scholarship Scheme for Innovation and Development (PRESSID) supervised by the Federal Scholarship Board (FSB).

References

- 1 S. A. Osmani, S. Bak and B. L. Moller, *Phytochemistry*, 2009, **70**, 325–347.



- 2 E. K. Lim and D. J. Bowles, *EMBO J.*, 2004, **23**, 2915–2922.
- 3 F. De Bruyn, J. Maertens, J. Beauprez, W. Soetaert and M. De Mey, *Biotechnol. Adv.*, 2015, **33**, 288–302.
- 4 D. Bowles, E. K. Lim, B. Poppenberger and F. E. Vaistij, *Annu. Rev. Plant Biol.*, 2006, **57**, 567–597.
- 5 G. Huang, M. Lv, J. Hu, K. Huang and H. Xu, *Mini-Rev. Med. Chem.*, 2016, **12**, 1013–1016.
- 6 V. Kren and L. Martinkova, *Curr. Med. Chem.*, 2001, **8**, 1303–1328.
- 7 G. Huang and X. Mei, *Curr. Drug Targets*, 2014, **15**, 780–784.
- 8 P. M. Coutinho and B. Henrissat, Carbohydrate-Active Enzymes server at URL, <http://afmb.cnrs-mrs.fr/~cazy/CAZY/index.html>, <http://afmb.cnrs-mrs.fr/CAZY/index.html>.
- 9 J. Ross, Y. Li, E. Lim and D. J. Bowles, *Genome Biol.*, 2001, **2**, REVIEWS3004.
- 10 P. Jones, B. Messner, J. Nakajima, A. R. Schaffner and K. Saito, *J. Biol. Chem.*, 2003, **278**, 43910–43918.
- 11 S. M. Patel, M. Fuente, S. Ke, A. Guimaraes, A. O. Oliyide, X. Ji, P. Stapleton, A. Osbourn, Y. Pan, D. J. Bowles, B. G. Davis, A. Schatzlein and M. Yang, *Chem. Commun.*, 2011, **47**, 10569–10571.
- 12 M. Nielsen, M. M. D. Suits, M. Yang, C. S. Barry, C. Martinez-Fleites, L. E. Tailford, J. E. Flint, C. Dumon, B. G. Davis, H. J. Gilbert and G. J. Davies, *J. Biol. Chem.*, 2011, **286**, 15155–15164.
- 13 M. Yang, G. J. Davies and B. G. Davis, *Angew. Chem., Int. Ed.*, 2007, **46**, 3885–3888.
- 14 D. Bolam, S. Roberts, M. Proctor, J. Turkenburg, E. Dodson, C. Martinez-Fleites, M. Yang, B. G. Davis, G. Davies and H. Gilbert, *Proc. Natl. Acad. Sci. U. S. A.*, 2007, **13**, 5336–5341.
- 15 W. Offen, C. Martinez-Fleites, M. Yang, E. Kiat-Lim, B. G. Davis, C. A. Tarling, C. M. Ford, D. J. Bowles and G. J. Davies, *EMBO J.*, 2006, **25**, 1396–1405.
- 16 M. Yang, M. Brazier, R. Edwards and B. G. Davis, *ChemBioChem*, 2005, **6**, 346–357.
- 17 J. Flint, E. Taylor, M. Yang, D. N. Bolam, L. E. Tailford, C. Martinez-Fleites, E. J. Dodson, B. G. Davis, H. J. Gilbert and G. J. Davies, *Nat. Struct. Mol. Biol.*, 2005, **12**, 608–614.
- 18 L. V. Modolo, J. W. Blount, L. Achnine, M. A. Naoumkina, X. Wang and R. A. Dixon, *Plant Mol. Biol.*, 2007, **64**, 499–518.
- 19 T. Vogt and P. Jones, *Trends Plant Sci.*, 2000, **5**, 380–386.
- 20 D. Bowles, J. Isayenkova, E. K. Lim and B. Poppenberger, *Curr. Opin. Plant Biol.*, 2005, **8**, 254–263.
- 21 T. Mo, X. Liu, Y. Y. Liu, X. H. Wang, L. Zhang, J. Wang, Z. X. Zhang, S. P. Shi and P. F. Tu, *RSC Adv.*, 2016, **6**, 84616–84626.
- 22 V. Veljanovski and C. P. Constabel, *Phytochemistry*, 2013, **91**, 148–157.
- 23 X. X. Huang, G. Q. Zhu, Q. Liu, L. Chen, Y. J. Li and B. K. Hou, *Plant Physiol.*, 2018, **176**, 3103–3119.
- 24 M. Yang, M. Brazier, R. Edwards and B. G. Davis, *ChemBioChem*, 2005, **6**, 346–357.
- 25 P. Parajuli, R. P. Pandey, N. T. H. Trang, T. J. Oh and J. K. Sohng, *Carbohydr. Res.*, 2015, **418**, 13–19.
- 26 P. Parajuli, R. P. Pandey, S. Darsandhari, Y. I. Park and J. K. Sohng, *J. Carbohydr. Chem.*, 2016, **35**, 367–377.
- 27 H. Cao, X. Chen, A. R. Jassbi and J. Xiao, *Biotechnol. Adv.*, 2015, **33**, 214–223.
- 28 K. Yonekura-Sakakibara, T. Tohge, R. Niida and K. Saito, *J. Biol. Chem.*, 2007, **282**, 14932–14941.
- 29 F. Cuyckens and M. Claeys, *J. Mass Spectrom.*, 2004, **39**, 1–15.
- 30 N. Fabre, I. Rustan, E. Hoffmann and J. Quetin-Leclercq, *J. Am. Soc. Mass Spectrom.*, 2001, **12**, 707–715.
- 31 J. A. Capra and M. Singh, *Bioinformatics*, 2007, **23**, 1875–1882.
- 32 B. Hou, E. K. Lim, G. S. Higgins and D. J. Bowles, *J. Biol. Chem.*, 2004, **279**, 47822–47832.

