

Table 1. Demographic characteristics of women at booking

	Total (n=107)	Deflated* (n=42)	Inflated† (n=54)	Unknown (n=11)
Age				
Mean (y) +/- SD	31.8 +/- 4.9	32.5 +/- 4.5	31.3 +/- 4.8	31.2 +/- 6.5
≥35y, n= (%)	33 (30.8)	14 (33.3)	15 (27.8)	4 (36.4)
≥40y, n= (%)	8 (7.5)	3 (7.1)	4 (7.4)	1 (9.1)
Missing, n=	0	0	0	0
Parity				
Nulliparity, n= (%)	50 (46.7)	19 (45.2)	27 (50.0)	4 (36.4)
Missing, n=	0	0	0	0
Ethnicity				
White, n= (%)	100 (93.5)	40 (95.2)	50 (92.6)	10 (90.9)
Asian, n= (%)	4 (3.7)	2 (4.8)	2 (3.7)	0 (0.0)
Black African, n= (%)	1 (0.9)	0 (0.0)	0 (0.0)	1 (9.1)
Mixed, other, n= (%)	2 (1.9)	0 (0.0)	2 (1.9)	0 (0.0)
Missing, n=	0	0	0	0
Weight at booking				
Mean (kg) +/-SD	101.7 +/- 20.9	101.7 +/- 21.0	101.2 +/- 20.7	103.6 +/- 23.6
Missing, n=	1	1	0	0
Booking BMI				
Mean (kg/m ²) +/- SD	36.3 +/- 7.2	36.3 +/- 7.4	36.1 +/- 7.1	37.7 +/- 7.6
Missing, n=	1	1	0	0
Employment status				
Unemployed, n= (%)	26 (25.0)	10 (24.4)	16 (30.2)	0 (0.0)
Missing, n=	3	1	1	1
Smoking status				

Yes, n= (%)	19 (17.8)	4 (9.5)	10 (18.5)	5 (45.5)
Missing, n=	0	0	0	0
Diabetic status[‡]				
Diabetes, n= (%)	7 (6.5)	2 (4.8)	4 (7.4)	1 (0.9)
Missing, n=	0	0	0	0
Hypertensive status[§]				
Hypertensive, n= (%)	11 (10.3)	3 (7.1)	8 (14.8)	0 (0.0)
Missing, n=	0	0	0	0

*Pregnant women who underwent band deflation pre- or during pregnancy. †Pregnant women who had band inflation maintained throughout pregnancy. ‡Pre-existing diabetes. §Pre-existing hypertensive disease