

**Self-reported sensory impairments and changes in cognitive performance: a longitudinal
6-year follow-up study of English community-dwelling adults aged ≥50 years**

Authors:

Ann E.M. Liljas¹, PhD

Kate Walters¹, PhD

Cesar de Oliveira², PhD

S. Goya Wannamethee¹, PhD

Sheena E. Ramsay³, PhD

Livia A. Carvalho⁴, PhD

¹Department of Primary Care and Population Health, University College London, London, UK

²Department of Epidemiology and Public Health, University College London, London, UK

³Institute of Health & Society, Newcastle University, Newcastle-upon-Tyne, UK

⁴Department of Clinical Pharmacology, Queen Mary University of London, London, UK

Corresponding author: Dr Ann E.M. Liljas, UCL Department of Primary Care & Population Health, University College London, Rowland Hill Street, London NW3 2PF, UK. E-mail: ann.liljas.13@ucl.ac.uk Telephone: +44(0)207 830 2393 Fax: +44(0)20 7794 1224

Funding

This work was supported by the National Institute for Health Research, School for Primary Care Research (538469 to A.L.); the National Institute on Aging USA (5 R01 AG017644-19 to C.O.); UK Medical Research Council (G1002391 to S.R.); Dunhill Medical Trust (R396/1114 to S.R.); British Heart Foundation (RG/10/001/28296 to L.C.); and the UK

Medical Research Council (RG71546 to L.C.). The opinions expressed in the paper are those of the authors and not necessarily those of the funding bodies.

Acknowledgement

The English Longitudinal Study of Ageing was developed by a team of researchers based at the University College London, National Centre for Social Research and the Institute for Fiscal Studies. The data were collected by the National Centre for Social Research. The funding is provided by the National Institute of Aging in the United States, and a Consortium of UK Government Departments Coordinated by the Economic and Social Research Council (ESRC). The developers and funders of the English Longitudinal Study of Ageing and the UK Data Archive do not bear any responsibility for the analyses or interpretations presented here.

Conflict of interest

All authors declare that there are no competing interests.

Author contribution

LC, KW, SR, SGW and AL planned the study. AL and CO prepared the data. AL and LC developed the statistical model. AL did the statistical analysis with input from LC. AL wrote the manuscript with input from KW, LC, CO, SR and SGW. AL, LC and KW provided input to the interpretation of the results, and SR, SGW and CO critically revised the manuscript.

All authors have approved the final submitted version.

ABSTRACT

Objective: To investigate the influence of single and dual sensory impairments prospectively on cognition in adults aged ≥ 50 years.

Methods: Community-dwelling English adults ($n=4621$) followed-up from 2008 to 2014. Self-reported hearing and vision were collected in 2008. Change in cognitive performance on working memory and executive function between 2008 and 2014.

Results: Compared to good hearing and good vision respectively, poor hearing and poor vision were associated with worse cognitive function (hearing: unstandardized coefficient $B=0.83$, 95%CI 0.29-1.37; vision: $B=1.61$, 95%CI 0.92-2.29 adjusted for age, sex, baseline cognition).

Compared to no sensory impairment, dual sensory impairment was associated with worse cognition ($B=2.30$, 95%CI 1.21-3.39 adjusted for age, sex, baseline cognition). All associations remained after further adjustment for sociodemographic characteristics, lifestyle factors, chronic conditions, falls, mobility, depression and lack of companionship.

Discussion: The findings are important as age-related sensory impairments are often preventable or modifiable, which may prevent or delay cognitive impairment.

Key words: ageing, hearing loss, vision loss, dual sensory impairment, cognition

INTRODUCTION

Many populations worldwide including the population of England are ageing due to increased life expectancy (ONS (Office for National Statistics), 2015; WHO (World Health Organization), 2015). Advanced age increases the risk of health problems including age-related loss of hearing and vision (Khaw, 1997). Hearing impairment is estimated to affect one in five (19%) adults aged 51-80 years in England and Wales (Akeroyd, Foreman & Holman, 2014). Among older adults aged 60 years and over 11% have a vision impairment (RNIB (Royal National Institute of Blind People), 2013). Experiencing both hearing and vision impairment (dual sensory impairment) is estimated to affect at least 3% of the older population (Heine & Browning, 2015). The number of older adults affected by sensory impairments is furthermore likely to increase as the population ages (Gopinath et al., 2009; Helzner et al., 2005). Both hearing impairment and vision impairment have been associated with chronic diseases and disability (Crews & Campbell, 2004; Liljas et al., 2016b, 2016a, 2016c; West et al., 1997), age-related problems known for reducing the chances of good health, wellbeing and independent living in later life (Campbell et al., 1999). This makes age-related sensory impairments an important public health concern. Another major health issue in later life is cognitive impairment, a key contributor to disability and dependence in older age (Lee et al., 2014; Mograbi et al., 2014). The prevalence of cognitive impairment is increasing in England due to an ageing population and increasing longevity (ONS (Office for National Statistics), 2016).

Several cross-sectional studies have shown associations of impairments in hearing and vision with cognitive impairment (Anstey, Lord & Williams, 1997; F. R. Lin, 2011; F. R. Lin et al., 2011; Lindenberger & Baltes, 1994; Tay et al., 2006). There is also evidence from longitudinal studies reporting increased risks of incident cognitive impairment in those with hearing impairment after adjustment for sociodemographic characteristics and CVD-related measures

(Fischer et al., 2016; F. R. Lin et al., 2013). However, other factors such as depression, social isolation and mobility limitations were not considered in these studies. A previous study investigating impairments of hearing and vision with incident cognitive impairment found that hearing impairment and, in particular, vision impairment were associated with cognitive decline at 6-year follow-up (Valentijn et al., 2005). The results were, however, only adjusted for age, sex and education. Other longitudinal studies have demonstrated that vision impairment, but not hearing impairment, was associated with an increased risk of incident cognitive impairment (Anstey, Luszcz & Sanchez, 2001; M. Y. Lin et al., 2004), suggesting that vision impairment more than hearing impairment predicts cognitive decline. It has furthermore been suggested that the relationship between sensory impairments and subsequent cognitive impairment might not be unique to one sensory function (Fischer et al., 2016). There has been little research on combined hearing and vision impairment (dual sensory impairment) and subsequent cognitive impairment (Heine & Browning, 2015). One study in women found a relationship between dual sensory impairment and incident cognitive impairment (M. Y. Lin et al., 2004), however, another study in both women and men did not observe an association between dual sensory impairment and cognitive decline (Hong et al., 2016). Therefore, this study, in a nationally representative cohort of English women and men aged ≥ 50 years, aims to examine the influence of single and dual sensory functioning on cognitive function at 6-year follow-up on adjustment for a range of possible covariates, including baseline cognitive functioning.

METHODS

Study design and participants

This study uses data from the English Longitudinal Study of Ageing (ELSA). ELSA is a prospective study of a nationally representative sample of men and women aged ≥ 50 years who

participated in the Health Survey for England in 1998, 1999 or 2001 (Marmot et al., 2015). Since 2002, participants have been followed-up every two years for an interview on health and lifestyle and every four years for a physical examination. This study sample is restricted to the 4621 participants (62% of respondents aged ≥ 50 years in 2008) who undertook the cognitive tests in 2008 and 2014 and provided data on sensory function and covariates in 2008 (derivation of study sample outlined in Figure 1). All participants provided informed consent and ethical approval for ELSA was obtained from the Multicentre Research and Ethics Committee.

Hearing impairment

Hearing function was assessed by asking participants to rate their hearing (using a hearing aid if they use one) as excellent, very good, good, fair or poor. Reporting excellent, very good or good hearing was classified as having good hearing and this group formed the reference group. Experiencing fair or poor hearing was considered as poor hearing. The self-reported question used has previously been shown to be accurate when compared against objectively measured hearing (Ferrite, Santana & Marshall, 2011; Gibson et al., 2014).

Vision impairment

Vision function was measured using a validated self-reported question previously demonstrating a significant association with objectively measured eyesight, asking participants whether their eyesight was excellent, very good, good, fair or poor using glasses or corrective lens if they normally do so (Zimdars, Nazroo & Gjonca, 2012). Good vision was defined as reporting excellent, very good or good eyesight and was used as the reference group. Reporting fair or poor eyesight was classified as poor vision.

Dual sensory impairment

Being classified as having both poor hearing and poor vision was defined as having dual sensory impairment. Participants with no sensory impairment acted as the reference group because, as presented above, single sensory impairment is associated with adverse health outcomes. Having no impairment was considered as a more suitable reference to allow for understanding of the magnitude of the impact of having dual impairment compared to the general population.

Assessment of cognitive performance

Participants' cognitive status was first assessed in 2008 and then again in 2014 which allowed for participants to be followed-up for 6 years. Cognition was based on a modified version of the cognitive score developed by Batty et al.(2016) referring to working memory and executive function (Zaninotto & Batty, 2018). Working memory included immediate and delay recall tests of 10 nouns presented to the participants who were asked to recall as many words as possible immediately after the list was read, and then again after an approximately 5-minute delay during which they completed other survey questions (range 0-20). Executive function was ascertained using a word-finding task asking participants to name as many different animals as possible in 1 minute (range 0-60). In the original score, executive function also included a letter cancellation task however such data were not available in 2014. Similar to the calculations of the original score, the results from the three cognitive tests available were summed, providing a cognitive score (range 0-80) with a lower score indicating worse cognitive function. Change in cognitive function was calculated by subtracting baseline scores from the scores at follow-up.

Covariates

Possible confounders considered in the analyses included age, sex, wealth, educational qualification, smoking, alcohol, physical activity, cardiovascular disease (CVD), diabetes and hypertension. Wealth was based on total net non-pension wealth (financial, housing and physical wealth) of the household presented by quintiles. Educational qualification was defined as having an intermediate or higher qualification compared to no qualification. Smoking was defined as reporting being a current smoker or current non-smoker. Alcohol consumption was based on frequency of consumption of all kind of alcoholic drinks in the last 12 months and grouped into ‘daily’, ‘frequently’ (once or twice a week or more, but not every day), ‘rarely’ (once or twice a month/once every couple of months) and ‘never’. Physical activity was based on frequency and intensity in exercise by asking participants how often they engage in vigorous, moderate and mild exercise (more than once a week, once a week, one to three times a month, hardly ever or never). Participants who hardly ever or never engaged in vigorous, moderate and mild activity were classified as sedentary. Engaging in mild activity one to three times a month, once a week or more than once a week, or engaging in moderate activity one to three times a month was classified as low activity. Participants engaging in moderate activity once a week or more than once a week or vigorous activity one to three times a month were classified as being moderately active. Undertaking vigorous activity once a week or more than once a week was classified as high activity. Objective data on height and weight were used to calculate body mass index (BMI). Self-reported doctor-diagnosed CVD (myocardial infarction, angina and/or stroke), diabetes and hypertension were analysed dichotomously. Other important factors potentially associated with sensory impairments and cognitive impairment that were considered included history of falls, mobility limitations, depression and lack of companionship. History of falls was based on participants reporting falling in the last 12 months. Mobility limitations referred to reporting problems walking 100 yards and/or climbing one flight of stairs. Depression was based on the validated 8-item version of CES-D (Radloff,

1977). Scoring positively on 4 or more of the 8 items were classified as having depression. Feeling lack of companionship ‘some of the time’ or ‘often’ were combined and compared to feeling no lack of companionship and compared to feeling no lack of companionship.

Statistical analyses

Linear regression was used to assess longitudinal associations between impairments in hearing and vision (individually and combined) in 2008 with a continuous measure of the outcome variable cognition in 2014. The regression models provided unstandardized coefficients B (the adjusted mean difference in the cognitive measure between those who did and those who did not report sensory impairments) with 95% confidence intervals (CI). Each domain of the cognitive score was furthermore tested individually. Sensory impairment (single/dual) was coded as 0 and no sensory impairment coded as 1. The statistical analyses were adjusted for age, sex and cognitive function at baseline as well as covariates significantly associated with sensory impairments in this study sample (Table 1) and in previous research (Crews & Campbell, 2004; Liljas et al., 2016c). All variables were entered as categorical variables except for age and BMI which were entered as continuous variables. All analyses were carried out using SPSS (Version 22, IBM, Armonk, New York, USA).

RESULTS

A total of 4621 participants (55% women) aged ≥ 50 years (mean age 64.9 years (SD 8.3) were included. One in five (19%) self-reported poor hearing and 10% self-reported poor vision. Dual sensory impairment was prevalent in 179 participants (5% of 3641 subjects who had no sensory impairment or dual sensory impairment). On the cognitive scale ranging from 0-80 with higher scores demonstrating better cognitive function, average performance of all participants was 32.8 (SD 8.3) in 2008 and 31.4 (SD 9.6) in 2014.

Table 1 presents the characteristics of all participants in 2008 (baseline) for hearing impairment and vision impairment. Compared to participants with good hearing, those with poor hearing had significantly lower scores on cognitive function in both 2008 and 2014. Poor hearing was associated with being older, male, less wealthy, having no educational qualification, less physically active, chronic conditions including hypertension, CVD and diabetes, a history of falls, mobility limitations and depression. Participants with poor hearing were more likely to consume alcohol daily but also more likely to never drink, compared to participants with good hearing. Similarly, individuals with poor vision performed worse on cognition in 2008 and in 2014 than those with good vision. Poor vision was also associated with advanced age, being female, less wealth, no educational qualification, lower alcohol consumption, less physically active, BMI ≥ 30 , chronic conditions, falls, mobility limitations, depression and lack of companionship. Table 2 shows the characteristics of 3641 subjects who had no sensory impairment (n=3462) or dual sensory impairment (n=179). In comparison to participants with no sensory impairment, those with dual sensory impairment had lower scores on cognitive function in both 2008 and 2014 and were less wealthy, had no educational qualification, lower alcohol consumption, more likely to smoke, less physically active, chronic conditions, falls, mobility limitations, depression and lack of companionship.

Table 3 presents the findings from the linear regression models investigating whether impairments in hearing and vision influence cognitive function at 6-year follow-up. The findings showed that both hearing impairment and vision impairment were associated with worse cognitive performance at 6-year follow-up (adjusted for age, sex and cognitive function at baseline: hearing impairment unstandardized coefficient B=0.83, 95% CI 0.29, 1.37, p<0.01; vision impairment unstandardized coefficient B=1.61, 95% CI 0.92, 2.29, p<0.01). The

associations remained after further adjustment for wealth, educational qualification, alcohol, smoking, physical activity, obesity, CVD, diabetes, hypertension, falls, mobility, depression and lack of companionship with stronger associations observed for vision impairment (unstandardized coefficient $B=0.93$, 95% CI 0.22, 1.64, $p=0.01$) than for hearing impairment (unstandardized coefficient $B=0.57$, 95% CI 0.03, 1.12, $p=0.04$). Similarly, compared to participants with no sensory impairment, individuals with dual sensory impairment were more likely to demonstrate worse cognitive performance at 6-year follow-up (adjusted for age, sex and cognitive function at baseline unstandardized coefficient $B=2.30$, 95% CI 0.96, 3.13) and the association remained after further adjustment for covariates (unstandardized coefficient $B=1.59$, 95% CI 0.36, 2.58). As previous literature has suggested potential differences in outcomes between men and women with sensory impairments (Murphy & Gates, 1997; West et al., 1997), we tested for an interaction with gender, and this was non-significant.

Supplementary analyses of each cognitive domain part of the cognitive score (Table S1) showed that poor hearing was associated with lower scores on immediate and delayed recall at 6-year follow-up. Poor hearing was not associated with lower scores on executive functioning at follow-up after adjustment for covariates. At 6-year follow-up, poor vision was associated with lower scores on executive functioning and immediate recall but not delayed recall after adjustment for covariates. Dual sensory impairment was associated with lower scores on all three cognitive domains at follow-up and the associations remained after adjustment for covariates.

DISCUSSION

This study investigated the relationships of hearing impairment, vision impairment and dual sensory impairment with change in cognitive performance at 6-year follow-up in English adults

aged ≥ 50 years. The results show that in this ageing population, poor hearing and poor vision, individually and combined, are associated with worse cognitive performance at 6-year follow-up. The associations observed remained after adjustment for a wide range of covariates including sociodemographic characteristics, lifestyle factors, chronic conditions, falls, mobility, depression and lack of companionship.

Our study findings add to current literature on the relationships of impairments in hearing and vision with cognitive function as this is one of the very first studies examining dual sensory impairment and cognitive function longitudinally. Only two previous studies have investigated this relationship prospectively, reporting inconsistent findings; one study demonstrated an association between dual sensory impairment and incident risk of cognitive impairment at 4-year follow-up (M. Y. Lin et al., 2004). However, that study was in women only ($n=6112$). The other study did not observe an association between dual sensory impairment and cognitive decline at 5- and 10-year follow-up, possibly due to lack of statistical power (93 (2.5%) of 3654 participants reported dual sensory impairment at baseline; OR 1.41, 95% CI 0.54–3.72 and OR 1.15, 95% CI 0.28–4.73, respectively) (Hong et al., 2016). In our study we demonstrated a relationship between dual sensory impairment and cognitive decline at 6-year follow-up in both women and men. The findings of the current study also contribute to existing literature on sensory impairments and subsequent cognitive impairment as it showed such relationships even after adjustment for covariates including falls, mobility, depression and lack of companionship, factors not adjusted for in previous studies (Anstey, Luszcz & Sanchez, 2001; F. R. Lin, 2011; F. R. Lin et al., 2013; Lindenberger & Baltes, 1994; Tay et al., 2006; Valentijn et al., 2005). The dual effect on cognitive function appears to be additive, i.e above and beyond the presence of cognitive impairment alone, which is consistent with other evidence from the literature (Guthrie et al., 2018). It remains, however, unclear whether the relationship between

sensory impairments and cognitive impairment is direct or indirect. A direct causal relationship might exist through poor sensory function reducing the opportunities to cognitive stimulation, leading to cognitive deterioration caused by cerebral atrophy (Lindenberger & Baltes, 1994). Alternatively, a direct causal relationship may be explained by poor sensory function requiring more cognitive resources to interpret information perceived, resulting in less cognitive capacity available for other cognitively demanding tasks (Baltes & Lindenberger, 1997). The associations observed between sensory impairments and worse cognitive performance on adjustment for a range of covariates support the hypotheses of a direct causal relationship.

The relationship between sensory impairments and worse cognitive performance could be due to shared age-related factors including degeneration of central nervous structures (Lindenberger & Baltes, 1994), or CVD (Crews & Campbell, 2004; Dregan, Stewart & Gulliford, 2013). While our study showed a relationship of sensory impairments with worse cognitive performance after adjustment for CVD and CVD-related conditions such as hypertension, diabetes, and CVD risk factors including smoking, a higher BMI and physical activity, there may be residual (unmeasured) confounding. There could also be psychosocial factors such as depression and social isolation linking vision impairment to poor future cognitive performance (Barnes et al., 2004; Heine & Browning, 2004). In our study we further explored the independent effect of sensory impairments on cognitive performance after adjustment for the psycho-social factors depression and lack of companionship. However, the measures available with sufficient data may have incompletely captured these domains. Other aspects including anxiety, participation in social activities and subjective feelings of loneliness may also be important. It is also possible that the relationship is due to underlying mechanisms such as inflammation (Peracino & Pecorelli, 2016).

Strengths and limitations

The major strengths of this study are that it is based on data of older English adults from a large population-based cohort. The subjects were followed-up 6 years later for changes in cognitive function and the models were adjusted for a wide range of potential covariates.

Limitations include that hearing impairment and vision impairment were self-reported rather than objectively measured. However, the questions used have been validated against objective measures (Ferrite, Santana & Marshall, 2011; Gibson et al., 2014; Zimdars, Nazroo & Gjonca, 2012), and the prevalence rates of sensory impairments reported are similar to national estimates (Akeroyd, Foreman & Holman, 2014; RNIB (Royal National Institute of Blind People), 2013). Sensory function was assessed at baseline only and data on the primary cause of and change in sensory function were not available. Also, data on type of and frequency of use of glasses / lenses and hearing aids were not available. Furthermore, the differences in cognitive performance associated with sensory impairments were fairly small and may not be clinically relevant.

A modified version rather than the original cognitive score by Batty et al.(2016) was used. Whilst the original cognitive score included three domains of cognitive function; working memory, executive function and processing speed, data on processing speed were not collected in 2014 and hence not available for the analyses conducted in this study. The working memory tests asking the participants to recall 10 common nouns required some degree of hearing to complete. Miscommunication was minimised by verbal information being provided face-to-face in a quiet environment by experienced examiners accustomed to working with older adults. The list of words used for the memory tests was furthermore presented by a recorded computer voice and the volume was adjusted prior to the test if necessary (Marmot et al., 2015).

Nevertheless, supplementary analyses of the individual cognitive domains showed that poor hearing was associated with immediate and delayed recall (domains requiring adequate hearing) but not with executive functioning (no hearing required), and difficulty in initial hearing of the words may have impacted on their performance in the tests of recall. However, no potential study participants reported being unable to undertake the recall tests due to deafness. The measurements of cognition (naming animals and recall of words) did not require adequate eyesight. The exclusion of participants who had incomplete data on sensory impairments, cognition and covariates raise the possibility of a selection bias toward healthier subjects. In keeping with most longitudinal cohort studies, we observed that indeed 1937 subjects with baseline measures eligible for participation in the study lost to follow-up were more likely to be older ($p<0.01$), less wealthy ($p<0.01$) and poorer health including more likely to be a current smoker ($p<0.01$) and having CVD ($p<0.01$), depression ($p<0.01$) and mobility limitations ($p<0.01$). Thus, the associations between sensory impairments and worse cognitive function observed in our study sample of a “younger” and “healthier” population with complete data might have been even stronger in a sample that included the non-respondents too. Study limitations also include several unmeasured and incompletely addressed factors of potential importance (e.g. anxiety and low social engagement) that may have confounded the relationship of impairments in hearing and/or vision and cognitive decline. The study was furthermore carried out in a population of ‘younger old’ adults (average 64.9 years) predominantly of white English ethnic origin. It may therefore not be appropriate to extrapolate our findings to other older populations.

CONCLUSIONS

In our study, ageing adults with individual and combined impairments in hearing and vision had greater risks of worse cognitive performance at 6-year follow-up compared to those with

good sensory function. Sensory impairments can often be prevented or modified and targeting sensory impairments in ageing adults could have potential to prevent or delay cognitive impairment. This is of importance to reduce the risk of cognitive impairment, a key contributor to disability, dependency and mortality in England.

Table 1. Age, sex, sociodemographic characteristics, lifestyle factors, comorbidities, falls, mobility limitations, depression, lack of companionship and cognitive function by hearing function and vision function in a cohort of 4621 English men and women aged 50 years and over in 2008 (baseline)

	Overall	Good hearing	Poor hearing	p-value	Good vision	Poor vision	p-value
<i>Totals, (n)%</i>	4621 (100)	3761 (81)	860 (19)		4143 (90)	478 (10)	
<i>Covariates</i>							
Age, mean (SD)	64.9 (8.3)	64.4 (8.1)	67.1 (8.6)	<0.01	64.7 (8.1)	67.3 (9.5)	<0.01
Male gender, (n)%	2100 (45)	1590 (42)	510 (59)	<0.01	1906 (46)	194 (41)	0.01
Wealth, (n)%							
1 (lowest)	636 (14)	487 (13)	149 (17)	0.01	494 (12)	142 (30)	<0.01
2	829 (18)	667 (18)	162 (19)		737 (18)	92 (19)	
3	935 (20)	761 (20)	174 (20)		843 (20)	92 (19)	
4	1030 (22)	857 (23)	173 (20)		956 (23)	74 (16)	
5 (highest)	1191 (26)	989 (26)	202 (24)		1113 (27)	78 (16)	
No educational qualification, (n)%	1015 (22)	787 (21)	228 (27)	<0.01	857 (21)	158 (34)	<0.01
Alcohol, (n)%							
Daily	763 (17)	604 (16)	159 (19)	0.01	706 (17)	57 (12)	<0.01
Frequently	2223 (48)	1845 (49)	378 (44)		2015 (49)	208 (44)	
Rarely	859 (19)	707 (19)	152 (18)		770 (19)	89 (19)	
Never	776 (17)	605 (16)	171 (20)		652 (16)	124 (26)	
Smoker, (n)%	602 (13)	480 (13)	122 (14)	0.14	508 (12)	94 (20)	<0.01
Levels of physical activity, (n)%							
Sedentary	174 (4)	127 (3)	47 (6)	<0.01	130 (3)	44 (9)	<0.01
Low	568 (12)	433 (12)	135 (16)		459 (11)	109 (23)	
Moderate	2327 (50)	1895 (50)	432 (50)		2086 (50)	241 (51)	
High	1551 (34)	1305 (35)	246 (29)		1468 (35)	83 (17)	
Body mass index (BMI), mean (SD)	28.3 (5.2)	28.3 (5.3)	28.4 (4.8)	0.55	28.3 (5.2)	28.9 (5.6)	0.01
Hypertension, (n)%	1873 (41)	1479 (39)	394 (46)	<0.01	1623 (39)	250 (52)	<0.01
CVD, (n)%	538 (12)	390 (10)	148 (17)	<0.01	441 (11)	97 (20)	<0.01
Diabetes, (n)%	384 (8)	293 (8)	91 (11)	0.01	322 (8)	62 (13)	<0.01
History of falls, (n)%	819 (18)	615 (16)	204 (24)	<0.01	682 (17)	137 (29)	<0.01
Mobility limitations, (n)%	1408 (31)	1168 (28)	240 (50)	<0.01	1063 (28)	345 (40)	<0.01
Depression, (n)%	540 (12)	421 (11)	119 (14)	0.02	441 (11)	99 (21)	<0.01
Lack of companionship, (n)%	1603 (35)	1296 (35)	307 (36)	0.24	1383 (34)	220 (47)	<0.01
<i>Outcomes</i>							
Cognitive function, mean (SD) in 2008	32.8 (8.3)	33.2 (8.2)	31.2 (8.5)	<0.01	33.1 (8.1)	30.6 (9.1)	<0.01
Cognitive function, mean (SD) in 2014	31.4 (9.6)	32.0 (9.5)	29.1 (9.9)	<0.01	31.8 (9.5)	27.9 (10.0)	<0.01

Table 2. Age, sex, sociodemographic characteristics, lifestyle factors, comorbidities, falls, mobility limitations, depression, lack of companionship and cognitive function in a cohort of 3641 English men and women aged 50 years and over with no sensory impairment versus dual sensory impairment in 2008 (baseline)

	Overall	No sensory impairment	Dual sensory impairment	p-value
<i>Totals, (n)%</i>	3641 (100)	3462 (95)	179 (5)	
<i>Covariates</i>				
Age, mean (SD)	64.5 (8.1)	64.3 (7.9)	68.9 (9.5)	<0.01
Male gender, (n)%	1568 (43)	1482 (43)	86 (48)	0.17
Wealth, (n)%				
1 (lowest)	453 (12)	399 (12)	54 (30)	<0.01
2	645 (18)	610 (18)	35 (20)	
3	737 (20)	703 (20)	34 (19)	
4	841 (23)	812 (24)	29 (16)	
5 (highest)	965 (27)	938 (27)	27 (15)	
No educational qualification, (n)%	767 (21)	698 (20)	69 (39)	<0.01
Alcohol, (n)%				
Daily	589 (16)	568 (16)	21 (12)	<0.01
Frequently	1783 (49)	1710 (49)	73 (41)	
Rarely	684 (19)	651 (19)	33 (18)	
Never	585 (16)	533 (15)	52 (29)	
Smoker, (n)%	452 (12)	419 (12)	33 (18)	0.01
Levels of physical activity, (n)%				
Sedentary	123 (3)	103 (3)	20 (11)	<0.01
Low	402 (11)	363 (11)	39 (22)	
Moderate	1834 (50)	1744 (50)	90 (50)	
High	1282 (35)	1252 (36)	30 (17)	
Body mass index (BMI), mean (SD)	28.3 (5.3)	28.2 (5.3)	28.8 (5.3)	0.18
Hypertension, (n)%	1427 (39)	1328 (38)	99 (55)	<0.01
CVD, (n)%	379 (10)	336 (10)	43 (24)	<0.01
Diabetes, (n)%	279 (8)	255 (7)	24 (13)	<0.01
History of falls, (n)%	582 (16)	530 (15)	52 (29)	<0.01
Mobility limitations, (n)%	1033 (28)	928 (27)	105 (59)	<0.01
Depression, (n)%	396 (11)	359 (10)	37 (21)	<0.01
Lack of companionship, (n)%	1242 (34)	1159 (34)	83 (47)	<0.01
<i>Outcomes</i>				
Cognitive function, mean (SD) in 2008	33.1 (8.2)	33.3 (8.1)	28.3 (8.8)	<0.01
Cognitive function, mean (SD) in 2014	31.9 (9.5)	32.2 (9.4)	25.5 (9.2)	<0.01

Table 3. Unstandardized coefficients B with 95% confidence intervals (CI) for relationships of vision impairment, hearing impairment and dual sensory impairment at baseline in 2008 with cognitive function at 6 years of follow up in 2014

Hearing impairment and cognitive function (n=4621)	Unstandardized coefficient B (95% CI)	p-value
Model 1 (M1): adjusted for age, sex, baseline cognitive function	0.83 (0.29, 1.37)	<0.01
Model 2 (M2): M1 + wealth, education	0.73 (0.19, 1.27)	0.01
Model 3 (M3): M2 + alcohol, smoking, physical activity, BMI	0.66 (0.12, 1.20)	0.02
Model 4 (M4): M3 + CVD, diabetes, hypertension	0.64 (0.10, 1.18)	0.02
Model 5 (M5): M4 + falls, mobility	0.59 (0.05, 1.14)	0.03
Model 6 (M6): M5 + depression, lack of companionship	0.57 (0.03, 1.12)	0.04
Vision impairment and cognitive function (n=4621)	Unstandardized coefficient B (95% CI)	p-value
Model 1 (M1): adjusted for age, sex, baseline cognitive function	1.61 (0.92, 2.29)	<0.01
Model 2 (M2): M1 + wealth, education	1.19 (0.49, 1.88)	<0.01
Model 3 (M3): M2 + alcohol, smoking, physical activity, BMI	0.99 (0.29, 1.69)	0.01
Model 4 (M4): M3 + CVD, diabetes, hypertension	0.96 (0.26, 1.66)	0.01
Model 5 (M5): M4 + falls, mobility	0.94 (0.24, 1.64)	0.01
Model 6 (M6): M5 + depression, lack of companionship	0.93 (0.22, 1.64)	0.10
Dual sensory impairment and cognitive function (n=3641)	Unstandardized coefficient B (95% CI)	p-value
Model 1 (M1): adjusted for age, sex, baseline cognitive function	2.30 (1.21, 3.39)	<0.01
Model 2 (M2): M1 + wealth, education	1.86 (0.77, 2.95)	<0.01
Model 3 (M3): M2 + alcohol, smoking, physical activity, BMI	1.67 (0.57, 2.76)	<0.01
Model 4 (M4): M3 + CVD, diabetes, hypertension	1.64 (0.55, 2.74)	<0.01
Model 5 (M5): M4 + falls, mobility	1.51 (0.41, 2.61)	0.01
Model 6 (M6): M5 + depression, lack of companionship	1.59 (0.47, 2.71)	0.01

Acknowledgement

The English Longitudinal Study of Ageing was developed by a team of researchers based at the University College London, National Centre for Social Research and the Institute for Fiscal Studies. The data were collected by the National Centre for Social Research. The funding is provided by the National Institute of Aging in the United States, and a Consortium of UK Government Departments Coordinated by the Economic and Social Research Council (ESRC). The developers and funders of the English Longitudinal Study of Ageing and the UK Data Archive do not bear any responsibility for the analyses or interpretations presented here.

Declaration of Conflicting Interests

None.

Funding

Information on funding would compromise the anonymity of the authors and has therefore been withheld.

REFERENCES

- Akeroyd, M. A., Foreman, K. & Holman, J. A. (2014). Estimates of the number of adults in England, Wales, and Scotland with a hearing loss. *International Journal of Audiology*, 53, (1), 60-1. doi:10.3109/14992027.2013.850539
- Anstey, K. J., Lord, S. R. & Williams, P. (1997). Strength in the lower limbs, visual contrast sensitivity, and simple reaction time predict cognition in older women. *Psychology and Aging*, 12, (1), 137-44.
- Anstey, K. J., Luszcz, M. A. & Sanchez, L. (2001). Two-year decline in vision but not hearing is associated with memory decline in very old adults in a population-based sample. *Gerontology*, 47, (5), 289-93. doi:10.1159/000052814
- Baltes, P. B. & Lindenberger, U. (1997). Emergence of a powerful connection between sensory and cognitive functions across the adult life span: a new window to the study of cognitive aging? *Psychol Aging*, 12, (1), 12-21.
- Barnes, L. L., Mendes de Leon, C. F., Wilson, R. S., Bienias, J. L. & Evans, D. A. (2004). Social resources and cognitive decline in a population of older African Americans and whites. *Neurology*, 63, (12), 2322-6.
- Batty, G. D., Deary, I. J. & Zaninotto, P. (2016). Association of Cognitive Function With Cause-Specific Mortality in Middle and Older Age: Follow-up of Participants in the English Longitudinal Study of Ageing. *Am J Epidemiol*, 183, (3), 183-90. doi:10.1093/aje/kwv139
- Campbell, V. A., Crews, J. E., Moriarty, D. G., Zack, M. M. & Blackman, D. K. (1999). Surveillance for sensory impairment, activity limitation, and health-related quality of life among older adults—United States, 1993-1997. *MMWR CDC Surveillance Summaries*, 48, (8), 131-56.
- Crews, J. E. & Campbell, V. A. (2004). Vision impairment and hearing loss among community-dwelling older Americans: implications for health and functioning. *American Journal of Public Health*, 94, (5), 823-9.
- Dregan, A., Stewart, R. & Gulliford, M. C. (2013). Cardiovascular risk factors and cognitive decline in adults aged 50 and over: a population-based cohort study. *Age Ageing*, 42, (3), 338-45. doi:10.1093/ageing/afs166
- Ferrite, S., Santana, V. S. & Marshall, S. W. (2011). Validity of self-reported hearing loss in adults: performance of three single questions. *Rev Saude Publica*, 45, (5), 824-30.
- Fischer, M. E., Cruickshanks, K. J., Schubert, C. R., Pinto, A. A., Carlsson, C. M., Klein, B. E., Klein, R. & Tweed, T. S. (2016). Age-Related Sensory Impairments and Risk of Cognitive Impairment. *J Am Geriatr Soc*, 64, (10), 1981-87. doi:10.1111/jgs.14308
- Gibson, W.K., Cronin, H., Kenny, R. A. & Setti, A. (2014). Validation of the self-reported hearing questions in the Irish Longitudinal Study on Ageing against the Whispered Voice Test. *BMC Res Notes*, 7, 361. doi:10.1186/1756-0500-7-361
- Gopinath, B., Rochtchina, E., Wang, J. J., Schneider, J., Leeder, S. R. & Mitchell, P. (2009). Prevalence of age-related hearing loss in older adults: Blue Mountains Study. *Archives of Internal Medicine*, 169, (4), 415-6. doi:10.1001/archinternmed.2008.597
- Guthrie, D. M., Davidson, J. G. S., Williams, N., Campos, J., Hunter, K., Mick, P., Orange, J. B., Pichora-Fuller, M. K., Phillips, N. A., Savundranayagam, M. Y. & Wittich, W. (2018). Combined impairments in vision, hearing and cognition are associated with greater levels of functional and communication difficulties than cognitive impairment alone: Analysis of interRAI data for home care and long-term care recipients in Ontario. *PLoS One*, 13, (2), e0192971. doi:10.1371/journal.pone.0192971
- Heine, C. & Browning, C. J. (2004). The communication and psychosocial perceptions of older adults with sensory loss: a qualitative study. *Ageing and Society*, 24, (1), 113-30.
- Heine, C. & Browning, C. (2015). Dual Sensory Loss in Older Adults: A Systematic Review. *Gerontologist*, 55, (5), 913-28. doi:10.1093/geront/gnv074

- Helzner, E. P., Cauley, J. A., Pratt, S. R., Wisniewski, S. R., Zmuda, J. M., Talbott, E. O., de Rekeneire, N., Harris, T. B., Rubin, S. M., Simonsick, E. M., Tylavsky, F. A. & Newman, A. B. (2005). Race and sex differences in age-related hearing loss: the Health, Aging and Body Composition Study. *Journal of the American Geriatrics Society*, 53, (12), 2119-27. doi:10.1111/j.1532-5415.2005.00525.x
- Hong, T., Mitchell, P., Burlutsky, G., Liew, G. & Wang, J. J. (2016). Visual Impairment, Hearing Loss and Cognitive Function in an Older Population: Longitudinal Findings from the Blue Mountains Eye Study. *PLoS One*, 11, (1), e0147646. doi:10.1371/journal.pone.0147646
- Khaw, K. T. (1997). Epidemiological aspects of ageing. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 352, (1363), 1829-35. doi:10.1098/rstb.1997.0168
- Lee, M., Saver, J. L., Hong, K. S., Wu, Y. L., Liu, H. C., Rao, N. M. & Ovbiagele, B. (2014). Cognitive impairment and risk of future stroke: a systematic review and meta-analysis. *CMAJ: Canadian Medical Association Journal*, 186, (14), E536-46. doi:10.1503/cmaj.140147
- Liljas, A. E., Wannamethee, S. G., Whincup, P. H., Papacosta, O., Walters, K., Iliffe, S., Lennon, L. T., Carvalho, L. A. & Ramsay, S. E. (2016a). Sensory Impairments and Cardiovascular Disease Incidence and Mortality in Older British Community-Dwelling Men: A 10-Year Follow-Up Study. *J Am Geriatr Soc*, 64, (2), 442-4. doi:10.1111/jgs.13975
- Liljas, A. E., Wannamethee, S. G., Whincup, P. H., Papacosta, O., Walters, K., Iliffe, S., Lennon, L. T., Carvalho, L. A. & Ramsay, S. E. (2016b). Hearing impairment and incident disability and all-cause mortality in older British community-dwelling men. *Age Ageing*, 45, (5), 662-7. doi:10.1093/ageing/afw080
- Liljas, A. E., Wannamethee, S. G., Whincup, P. H., Papacosta, O., Walters, K., Iliffe, S., Lennon, L. T., Carvalho, L. A. & Ramsay, S. E. (2016c). Socio-demographic characteristics, lifestyle factors and burden of morbidity associated with self-reported hearing and vision impairments in older British community-dwelling men: a cross-sectional study. *J Public Health (Oxf)*, 38, (2), e21-8. doi:10.1093/pubmed/fdv095
- Lin, F. R. (2011). Hearing loss and cognition among older adults in the United States. *Journals of Gerontology. Series A: Biological Sciences and Medical Sciences*, 66, (10), 1131-6. doi:10.1093/gerona/glr115
- Lin, F. R., Ferrucci, L., Metter, E. J., An, Y., Zonderman, A. B. & Resnick, S. M. (2011). Hearing loss and cognition in the Baltimore Longitudinal Study of Aging. *Neuropsychology*, 25, (6), 763-70. doi:10.1037/a0024238
- Lin, F. R., Yaffe, K., Xia, J., Xue, Q. L., Harris, T. B., Purchase-Helzner, E., Satterfield, S., Ayonayon, H. N., Ferrucci, L. & Simonsick, E. M. (2013). Hearing loss and cognitive decline in older adults. *JAMA Intern Med*, 173, (4), 293-9. doi:10.1001/jamainternmed.2013.1868
- Lin, M. Y., Gutierrez, P. R., Stone, K. L., Yaffe, K., Ensrud, K. E., Fink, H. A., Sarkisian, C. A., Coleman, A. L. & Mangione, C. M. (2004). Vision impairment and combined vision and hearing impairment predict cognitive and functional decline in older women. *Journal of the American Geriatrics Society*, 52, (12), 1996-2002. doi:10.1111/j.1532-5415.2004.52554.x
- Lindenberger, U. & Baltes, P. B. (1994). Sensory functioning and intelligence in old age: a strong connection. *Psychology and Aging*, 9, (3), 339-55.
- Marmot, M., Oldfield, Z., Clemens, S., Blake, M., Phelps, A., Nazroo, J., Steptoe, A., Rogers, N. & Banks, J. (2015) *English Longitudinal Study of Ageing: Waves 0-6, 1998-2013*. [online text], UK Data Service <<http://dx.doi.org/10.5255/UKDA-SN-5050-10>>
- Mograbi, D. C., Faria Cde, A., Fichman, H. C., Paradela, E. M. & Lourenco, R. A. (2014). Relationship between activities of daily living and cognitive ability in a sample of older adults with heterogeneous educational level. *Ann Indian Acad Neurol*, 17, (1), 71-6. doi:10.4103/0972-2327.128558
- Murphy, M. P. & Gates, G. A. (1997). Hearing Loss: Does Gender Play a Role? *Medscape Womens Health*, 2, (10), 2.

- ONS (Office for National Statistics). (2015). Ageing of the UK population. Retrieved from <http://webarchive.nationalarchives.gov.uk/20160105160709/http://www.ons.gov.uk/ons/relop/pop-estimate/population-estimates-for-uk--england-and-wales--scotland-and-northern-ireland/mid-2014/sty-ageing-of-the-uk-population.html>
- ONS (Office for National Statistics). (2016). Deaths registered in England and Wales 2015. Retrieved from
<https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/deathsregisteredinenglandandwalesseriesdr/2015>
- Peracino, A. & Pecorelli, S. (2016). The Epidemiology of Cognitive Impairment in the Aging Population: Implications for Hearing Loss. *Audiol Neurotol*, 21, (S1), 3-9.
- Radloff, L.S. (1977). The CES-D Scale: A Self-Report Depression Scale for Research in the General Population. *Applied Psychological Measurement*, 1, (3), 385-401.
- RNIB (Royal National Institute of Blind People). (2013). Sightloss UK 2013. The latest evidence. Retrieved from https://www.rnib.org.uk/sites/default/files/Sight_loss_UK_2013.pdf
- Tay, T., Wang, J. J., Kifley, A., Lindley, R., Newall, P. & Mitchell, P. (2006). Sensory and cognitive association in older persons: findings from an older Australian population. *Gerontology*, 52, (6), 386-94. doi:10.1159/000095129
- Valentijn, S. A., van Boxtel, M. P., van Hooren, S. A., Bosma, H., Beckers, H. J., Ponds, R. W. & Jolles, J. (2005). Change in sensory functioning predicts change in cognitive functioning: results from a 6-year follow-up in the maastricht aging study. *J Am Geriatr Soc*, 53, (3), 374-80. doi:10.1111/j.1532-5415.2005.53152.x
- West, S. K., Munoz, B., Rubin, G. S., Schein, O. D., Bandeen-Roche, K., Zeger, S., German, S. & Fried, L. P. (1997). Function and visual impairment in a population-based study of older adults. The SEE project. *Salisbury Eye Evaluation. Investigative Ophthalmology and Visual Science*, 38, (1), 72-82.
- WHO (World Health Organization). (2015). World report on ageing and health. Retrieved from http://apps.who.int/iris/bitstream/10665/186463/1/9789240694811_eng.pdf?ua=1
- Zaninotto, P. & Batty, G. D. (2018). Cognitive function trajectories and their determinants in older people: 8 years of follow-up in the English Longitudinal Study of Ageing. doi:10.1136/jech-2017-210116
- Zimdars, A., Nazroo, J. & Gjonca, E. (2012). The circumstances of older people in England with self-reported visual impairment: A secondary analysis of the English Longitudinal Study of Ageing (ELSA). *British Journal of Visual Impairment*, 30, (1), 22-30. doi:10.1177/0264619611427374