


Natural cytotoxic diterpenoids, a potential source of drug leads for melanoma therapy.

Jose M. Prieto^{*,a,b}, and Dâmaris Silveira^b

1. ^a Department of Pharmaceutical and Biological Chemistry, UCL School of Pharmacy, London, UK; ^b Department of Pharmacy, Faculty of Health Sciences, Universidade de Brasília, Brasília, Brazil.



Abstract: Diterpenes present complex structure and due to their unique carbon skeleton and interesting biological activities, have been the focus of continuous studies for the development of new anticancer agents. Phorbol esters have been known for their activity against skin malignancies since ancient times. Taxol was first studied in melanoma cells, and recently ingenol mebutate has been approved for the chemoprevention of melanoma in actinic keratosis patients. Therefore, there is scope for research on this class of compounds. We here aim to review the relevant original research reporting on isolated diterpenes with cytotoxic and/or antitumoral activities upon melanoma cells. By collating and discussing the implications of past and current developments on diterpenes, we hope to steer further interest on this field and facilitate the drug discovery activities of the scientific community towards finding potential alternatives to current melanoma chemotherapy.

Keywords: Melanoma, Diterpenes, Cytotoxicity, Cell Migration, Cell Invasion.

1 INTRODUCTION

Terpenoids are a large class of compounds presenting both structurally and stereochemically very diverse skeletons. Typical structures are originated from the C₅ isoprene units and are classified as hemiterpenes (C₅), monoterpenes (C₁₀), sesquiterpenes (C₁₅), diterpenes (C₂₀), sesterpenes (C₂₅), triterpenes (C₃₀), and tetraterpenes (C₄₀). They have been -together with alkaloids- one of the most successful sources for anticancer drugs. This review focuses on the diterpenes, which arise from geranylgeranyl diphosphate (GGPP) and comprise more than 18,000 different natural compounds presenting 126 different carbon skeletons [1]. Such array of structures and functionalities provides a pool of interesting molecules for therapeutic purpose.

The screening program run by the US National Cancer Institute in the 70s and 80s reached its climax with the discovery of taxol from *Taxus brevifolia* Nutt. (Taxaceae) which took the clinical world by storm and put natural products back as a valid source of medicines. This compound was isolated in 1971 by Wall and co-workers from Pacific Yew extracts guided by their powerful cytotoxic activity against a melanoma cell line. Eventually, taxol -now known as paclitaxel- was approved in the USA by the FDA in 1993, not for the treatment of melanoma but mainly ovarian and breast cancers. Since then, melanoma treatment keeps revolving on the use of synthetic chemotherapy drugs such as dacarbazine and temozolamine, alone or in combination with either interleukin-2 (IL-2) or monoclonal antibodies (vemurafenib, ipilimumab, dabrafenib, and trametinib). These therapies would need to be

improved for long-term efficacy and decreased toxicity [2].

The relationship between diterpene-rich plants and skin tumours has been exploited since ancient times. The spurges (a group of *Euphorbia* species) have been known since ancient times for their topical pro-inflammatory activity externally and cathartic effects when taken inwardly. Their characteristic milky sap is loaded with diterpenes. Its careful application onto warts removes them very effectively, but it is not devoid of local adverse effects, usually acute painful oedema. Chronic application of naturally occurring diterpenes in spurges, notably phorbol esters, is known to both induce cancerous changes on normal cells and differentiation in cancer cells.

The recent approval of the diterpenes derivative ingenol mebutate (Picato®) for the treatment of actinic keratosis -a skin condition which leads to melanoma in a very high proportion of patients- has renewed the interest on diterpenes as a source of anticancer drugs with particular activities on skin malignancies.

We here aim to review relevant original research reporting on isolated diterpenes with cytotoxic and/or antitumor activities upon melanoma cells. By collating and discussing the implications of past and current developments on diterpenes, we hope to steer further interest on this field and facilitate the drug discovery activities of the scientific community towards finding new avenues to current melanoma chemotherapy. Pubmed and Scopus have been searched for 'MELANOMA' and 'DITERPEN' to allow for diterpenes and diterpenoids to be retrieved.

All relevant papers to the review, regardless date of publication were used.

2 MELANOMA: PHYSIOPATHOLOGICAL AND MOLECULAR ASPECTS

Skin cancer is classified into non-melanoma skin cancer (NMSC) and melanoma. Melanoma originates from pigment-producing cells called melanocytes which produce melanin, a substance that is responsible for skin colour. While NMSC is the most common form, melanoma is so aggressive that causes approximately 80% of all deaths from skin cancer [3,4]. Melanoma pathogenesis goes through two major stages: the first is melanocytic transformation (melanogenesis) and the second is the spread of melanoma to other organs (metastasis).

Although patterns of UV exposure and personal characteristics greatly influence melanogenesis, this eventually occurs as a result of the accumulation of genetic changes. The most important oncogenes responsible for melanoma malignancy are: BRAF [5], NRAS [6], and mutant NF1 which are found in up to 50%, 20% and 14% of melanoma patients, respectively [7]. This led to a genomic classification consisting of four subtypes based on the pattern of the most prevalent significantly mutated genes: mutant BRAF, mutant RAS, mutant neurofibromin 1 (NF1), and Triple-WT (wild-type) [8]. We may soon be adding other molecular oncogenes to the list such as the tandem cyclin-dependent kinase 4 (CDK4) / cyclin D1 (CCND1): melanomas with wild-type BRAF or N-RAS frequently had increases in the number of copies of these genes [7].

BRAF and NRAS mutations seem to be mutually exclusive in the majority of melanomas, but mutant NRAS can drive a dual pathway signalling if the tumour suppressor PTEN is depleted. Then both the MAPK and the PI3K-AKT signalling pathways are activated enough to support the survival of the cancerous cell (see Fig 1) [9].

The identification of these targets facilitated the development of drugs with specific inhibitory activity upon BRAF and NRAS. Patients enjoy longer expectancy of life with such treatments, but resistance emerges very fast [10]. It looks like the above oncotargets cooperate with each other some combination therapies may be a better approach. In drug-resistant melanomas, the mutant BRAF seems to cooperate with NF1 loss to continue driving melanoma development through the abrogation of oncogene-induced senescence [11]. Furthermore, recent evidence points towards increased expression of the Microphthalmia-associated transcription factor (MITF) to contribute to resistance to BRAF pathway inhibition [12]. Although coming last in the

percentage of cases within melanoma patients with NF1-mutated subtype seems to have a higher mutational burden and strongest UV mutation signature than any of the previous ones [13]. NF1 mutant melanomas –characterised by a high level of CRAF expression and a differential MAPK activation- [14]

Another strategy is enhancing the natural cell defences by upregulating key tumour suppressors in melanoma such as p53, phosphatase and tensin homolog (PTEN), and p14-ARF. There is compelling evidence that the physiological regulation of p53 function depends on MDM2 - itself the product of a p53- inducible gene. The two molecules are linked to each other through an ‘autoregulatory negative feedback loop’ maintaining low cellular p53 levels in the absence of stress. DNA damage induces rapid MDM2 phosphorylation, which is thought to contribute to the p53 activation mechanism, in order to keep MDM2 localised in the nucleolus so that it cannot degrade p53 a 15-kDa protein called p14ARF [15]. Its effect blocks the G1 and G2 phases in the cell cycle and inhibits the growth of abnormal cells by indirectly activating p53. Therefore p14ARF could be considered as important as p53 as a key tumour suppressor in melanoma [16]. It is affected by several proteins but also small xenobiotics/molecules such as the flavonoid apigenin, thus proving that natural small molecules may be viable co-adjuvants in the battle against cancer [17].

A further pathway enhancing cell growth and proliferation without the involvement of the MAPK pathway is the reduction of retinoblastoma (RB) protein family activity [18]. This pathway is a less common one which results in the activation of the cell cycle regulators -and melanoma onto targets as previously explained- Cyclin D1 or CDK4 and initiate clonal expansion. However, it links with the senescence pathway which is central in melanoma. One of the key protein effectors here is p16-CDKN2A acting together with telomere length control. This protein inhibits the activity of cyclin D / CDK4 complex and blocks the hyperphosphorylation of RB to keep E2F transcription factors bound to RB.

Conversely, inactivation of p16-CDKN2A by mutation deletion or promoter silencing causes phosphorylation of RB protein which leads to the release of E2F allowing cell cycle progression. Clearly, this signalling system is the key regulator of melanocyte senescence, and its disruption enables cells to avoid senescence and causes uncontrolled cell growth as well as the aggressiveness of transformed melanocytes [19–22].

Resistance to apoptosis is required for the development of melanoma which is associated with mutations of p53, a major tumour suppressor gene.

P53, p14-CDKN2A and Bcl-2 proteins are the major proteins that are involved in this pathway [22]. As such apoptosis serves as a prophylactic mechanism to cancer development, cell death, the characteristic mechanisms of programmed cell death include chromatin condensation, membrane blebbing, and cell fragmentation, where the fragments are phagocytised by the neighbouring inflammatory cells [22]. These cellular changes occur because of the activity of caspase, an aspartate-specific cysteine protease, which is an essential element in promoting cell death. The caspase cascade system is required for initiation, transduction and amplification of apoptotic signals. It is regulated by several proteins and messengers such as Bcl-2 and Ca^{+2} and activated either through extracellular signals (extrinsic pathway) or intracellular signal (intrinsic pathway). The Bcl-2 family of proteins regulates apoptosis by controlling mitochondrial permeability. Failing that, Cytochrome C leaks into the mitochondrion and activates the caspases which cleave specific cellular substrates to effect cellular death [23].

Metastasis is the second stage in melanoma progression spread of cells from the primary tumour to distant locations where they form a secondary tumour. Tumour metastasis is responsible for 90% of cancer deaths [24]. In metastasis, several signal-transduction pathways are involved, enabling cancer cells to proliferate, remodel their neighbouring environment, invade and migrate to distant locations, and differentiate. Under normal skin condition, melanocytes remain under the control of keratinocytes and only proliferate after keratinocyte stimulation. However, deregulated tight junctions between melanocytes and their neighbouring stromal cells allow melanocytes to escape the tight control by adjacent keratinocytes [19–22].




Fig 1. Scheme of the main pathways, mutations / oncogenes and key tumor suppressors in melanoma.

3 DITERPENES FROM PLANTS

3.1 Phytane

Phytol (3,7,11,15-tetramethyl-2-hexadecen-1-ol) (**01**) is one of the simplest diterpenes. It is a reduced form of geranylgeraniol, which forms the lipophilic side chain of chlorophylls, therefore is produced by almost all photosynthetic organisms including plants, marine animals such as corals and bacteria [25]. It boasts an enormous variety of biological activities, ranging from antimicrobial and cytotoxic actions to immune-modulating properties [26]. It stimulates ROS production in B16F10 murine melanoma cells. This radical burst reduces melanogenesis *via* phosphorylation of the extracellular signal-regulated protein kinase (ERK) which in turn induces the proteasomal degradation of MITF [27].

Geranylgeranylacetone (**02**) (also known as all *E*-treprenone) is a pharmaceutical drug used for the treatment of gastric ulcers in Japan and other South-Asian countries. It was shown that it also induces apoptosis in human melanoma cells *via* the intrinsic pathway [28]. Although treprenone is not a naturally occurring diterpene, can be synthesised from natural compounds such as acyclic sesquiterpenes and polyprenylacetic acids [29,30].

3.2 Labdanes

Forskolin (**03**) is a labdane diterpene that is produced by the Indian Coleus plant (*Plectranthus barbatus* Andrews; Lamiaceae). It can influence the regulation of adenylate cyclase in murine melanoma tumour cell clones in a similar manner than melanocyte-stimulating hormone (MSH): both produce a significantly higher accumulation of intracellular cyclic adenosine 3', 5' monophosphate (cAMP) in strongly metastatic clones than in weakly metastatic B16 tumour cell clones. cAMP-dependent molecular processes are required for the expression of B16 melanoma experimental metastatic potential [31-33], but forskolin seems not to act as a hormone [34]. Clock genes are suppressed in B16 cells and tumours, but forskolin restores circadian rhythmicity and

subsequently cell cycle gene expression resulting in fewer cells in S phase and more in G1 phase. Accordingly, both *in vitro* proliferation and *in vivo* tumour growth slow after forskolin treatment [35].

Triptolide (**04**), from *Tripterygium wilfordii* Hook. f. (Celastraceae) shows promising activities both *in vitro* and *in vivo*. It induces cell arrest at S phase in A375.S2 human melanoma cells *via* the inhibition of cyclin E and CDC25A. Triptolide triggers mitochondrial-dependent apoptosis with caspase [36]. It also impairs DNA damage repair in A375.S2 [37]. Additionally, it inhibits migration and invasion in B16F10 cells by a Nuclear Factor kappaB (NF-κB) dependent pathway [38]. Triptolide decreases circulating T cells and plasma levels of IL-10, TGF-β, and VEGF in C57BL/6 mice bearing B16-F10 [39].

Andrographolide (**05**), from *Andrographis paniculata* (Burm.f.) Nees (Acanthaceae) exerted direct cytotoxicity on melanoma cells by cell-cycle arrest at G0/G1 phase through induction of cell-cycle inhibitory protein p27 and decreased expression of cyclin-dependent kinase 4 (CDK4). Moreover, andrographolide is able to increase the cytotoxic activity of lymphocytes against cancer cells by increased proliferation of lymphocytes and production of interleukin-2, enhanced synthesis of the tumour necrosis factor-alpha, and increased CD marker expression. These *in vitro* activities were confirmed in two *in vivo* syngenic xenograft models using B16F0 melanoma cells [40]. The activity of andrographolide has stimulated researchers to design semisynthetic derivatives which may be more easily protected in patents such as 3,14,19-tripropionylandrographolide (SRS06) (**06**). This compound targets NF-κB both by reducing its gene expression and reducing its DNA binding activity to p65 [41]. Moreover, it also suppresses melanin synthesis through the Akt/GSK3β/β-catenin signal pathway [42] and inhibits melanoma tumour growth by inactivating the TLR4/NF-κB signalling pathway [43].




Fig 2. Phytane and Labdane-type diterpenes with activity on melanoma cells

3.3 Pimaranes

The tubers of *Icacina* species (Icacinaceae) are a rich source of diterpenoids with (9 β H)-17-norpimarane and (9bH)-pimarane structures, the latest being a very rare structural class. 17-hydroxyicacinol (**07**), icacinol (**08**), and humirianthol (**09**), as well as two (9bH)-17-norpimaranes, humirianthenolide C (**10**) and icacenone (**11**), and the icacinlactones A–G (**12–18**) exhibited cytotoxic activity against MDA-MB-435 in the low micromolar range with humirianthenolide C being the most potent of all, [44–46].

3.4 Scopadulanes

The genus *Calceolaria* (Calceolariaceae), native of South America, have yielded several new diterpenes with the scopadulane skeleton, such as thyriflorin A (**19**). Its semisynthetic derivative, demalonyl thyriflorin A (**20**) is cytotoxic to melanoma A375 cells at 6.25 μ M by inducing an apoptotic response accompanied by the reduction of Heat Shock Protein 70 (Hsp70) expression and reactive oxygen species (ROS) production. It inhibits the caspase cascade at increased concentrations (25–50 μ M) an effect still accompanied by ROS increase, probably switching its mechanisms from apoptosis to necrosis [47].




Fig 3. Pimarane-type and Scopadulane-type diterpenes with activity on melanoma cells

3.5 Abietane

Carnosol (**21**), a well-known constituent of *Rosmarinus officinalis* L. (Lamiaceae) extracts, is a phenolic diterpene with antioxidant and anticarcinogen properties in melanoma cells [48,49].

Other researchers have also found that a carnosol-rich extract from *Callicarpa longissima* (Hemsl.) Merr. (Lamiaceae) inhibits melanogenesis in B16F10 melanoma cells [50]. It inhibits B16/F10 cell migration and invasion *in vitro* by differentially

inhibiting MMP-9 mRNA levels over those of MMP-2. Moreover, carnosol was able to inhibit the phosphorylation of extracellular signal-regulated kinase (ERK) 1/2, AKT, p38, JNK and prevented the activation of the transcription factors NF- κ B and c-Jun [48]. Furthermore, carnosol is both a potent genoprotective agent against radiation in normal skin cells and a sensitising agent, increasing the cellular death in the melanoma cell line B16F10 [51].

Carnosic acid (**22**) is another natural abietane diterpene found in rosemary and common sage in quantities up to 1.5 - 2.5% respectively. It may act as a transcriptional inducer of the NQO1 gene, and therefore suppresses the cytotoxic effect of inhibitors of this enzyme such as rhododendrol [52]. On the one hand, it suppresses the phosphorylation of Src, FAK, and AKT as well as the secretion of MMP-9, tissue inhibitor of metalloproteinase 1 (TIMP-1), urokinase plasminogen activator (uPA), and vascular cell adhesion molecule (VCAM)-1. On the other hand, increased the secretion of TIMP-2 at 10 μ M. Remarkably, it both suppresses the mesenchymal markers such as N-cadherin and induces epithelial markers such as E-cadherin thus potentially reverting the epithelial-mesenchymal transition thus hampering B16F10 cell migration [53].

Parvifloron D (**23**), a natural diterpene isolated from *Plectranthus ecklonii* Benth (Lamiaceae) presents broad but not selective cytotoxicity towards many human tumour cells, being able to induce cell death on human A375 and mouse B16V5 cell lines among others [54].

Although sharing commonalities with other abietane diterpenes, 11,12,16-trihydroxy-2-oxo-5-methyl-10-demethyl-abieta-1[10],6,8,11,13-pentene (**24**) isolated from *Premna serratifolia* L. (Lamiaceae) appears to be a novel compound based on a new diterpene skeleton with cytotoxic activity on B16 cells, presenting a promising $IC_{50} = 5 \mu$ M [55].

Several diterpenes containing an abietane skeleton were isolated from roots of *Peltodon longipes* A. St. Hill. Ex Benth (Lamiaceae) and evaluated about cytotoxicity against the human melanoma cancer cell line MV-3. Among them, the compounds 7-acetoxyroyleanone (**25**) and horminone (**26**) presented the highest activities with IC_{50} values of 7.4 and 16.7 μ M, respectively [56].

3.6 Halimanes

Compounds containing halimane skeleton can be found in different plant species from several families, as well as in marine organisms and microorganism. Biogenetically speaking they are a small group of diterpenes closely related to both labdane and clerodane diterpenoids [1]. One such compound, (5R,8R,9S,13R)-halim-1,10-ene-15,16-

diol (**27**), was isolated from *Vellozia kolbekii* Alves (Velloziaceae) and showed cytotoxicity against three human cancer cell lines, including MDA-MB-435 by a mechanism of action involving cell cycle arrest at the S/G2M phases [57].

3.7 Ingenol

The diterpene ingenol (**28**) was first isolated in 1968 by Opferkuch & Hecker. These authors were searching for the “*Euphorbia* factors” (skin irritant and tumour-promoting principles) present in *Euphorbia ingens* E. Mey. (Euphorbiaceae). Phorbol 12-myristate 13-acetate (PMA) is the reference in this category. It is a naturally occurring protein kinase C activator diterpene in many *Croton* species, currently classified as a Category 2 carcinogenic chemical hazard. However, they also isolated some “non-irritant” esters of ingenane-type polyfunctional diterpene alcohols devoid of “reasonable tumorigenic activity in mouse (*i.e.* about 1/10 less irritant than PMA). These authors could elucidate the structure through X-ray crystallography in 1970 and described it as endowed with a unique macrocyclic core. The development of this chemopreventative drug has its origins in the use of *Euphorbia peplus* L. (Euphorbiaceae) in Australian folk medicine for the treatment of actinic keratoses and skin cancer. Actinic keratosis is a precancerous condition which if untreated, usually leads to a melanoma [58].

Pharmacological work carried out during the 1990s showed that ingenol is endowed with both antitumour and *in vitro* anti-HIV activities. Mechanistic studies of its effects on actinic keratosis unveiled a dual mechanism of action including rapid necrosis of the affected skin cells and a specific neutrophil-mediated, antibody-dependent cellular cytotoxicity. Destruction of actinic keratosis lesions is accomplished in 2 or 3 days only, and the subsequent immune-mediated response prevents the development of any residual dysplastic epidermal cells. Direct isolation of ingenol mebutate from *E. peplus* is relatively inefficient (*c.a.* 1 mg of ingenol mebutate per Kg of plant) but still commercially viable so it was chosen as the herbal drug for pharmaceutical production. Partial or total synthesis has been extensively explored as a means to obtain both the active principle and other derivatives with the improved therapeutic profile. In 2012 the diterpene ingenol mebutate (**29**) (commercialised as Picato[®]) was finally approved for the topical treatment of actinic keratosis. The first post-marketing studies on the safety and tolerability of Picato[®] in the treatment of actinic keratosis seem to be very favourable with mild to moderate adverse effects (*i.e.* erythema, flaking/scaling and crusting) and resolve quickly. Further studies to apply it to the treatment of

superficial basal cell carcinomas with a 0.05% gel have been successful [59], so it looks like ingenol mebutate is poised to be a drug of choice for the treatment of superficial melanoma in the years to come.

The tetrahydroingenol diterpene 7,13-diacetyl-5-angeloyl-20-nicotinyl-3-propionyl-1,2,6,7-tetrahydroingenol (DANPT) (**30**), isolated from *Euphorbia erythradenia* Boiss. (Euphorbiaceae), is cytotoxic against A375 and HMCB cells ($IC_{50} < 20 \mu M$). It induced G2/M arrest associated with down-regulation of cyclin B and Cdk-1 and subsequent up-regulation of p53 and p21. Its apoptotic activity occurs with an increased Bax/Bcl-2 ratio, activation of caspase-3, ROS production and loss of mitochondrial membrane potential (ΔY_m) [60].

3.8 Ent-Kauranes

The epimer of kaurenoic acid (EKA) (**31**) isolated from the medicinal plant *Croton antisiphiliticus* Mart. (Euphorbiaceae) inhibited mitochondrial viability and induced apoptosis in B16 with an IC_{50} value of $60 \mu g/mL$ but was not specific when compared with 3T3 control cells.

Kaurenic acid (**32**), isolated from *Espeletia semiglobulata* Cuatrec (Asteraceae), presented potent cytotoxicity against B16-F1 melanoma cells, with an IC_{50} value of $0.8 \mu M$, two orders of magnitude lower than taxol ($IC_{50} = 19 \mu M$). Moreover, kaurenic acid was able to, decrease tumour development in C57BL/6 male mice transplanted with B16-F1 cells, with an inhibition rate of 70 and 80 % at 1.0 and 20.0 mg/kg, respectively, with taxol showing a rate of 54%, at 15 mg/kg [61].

The diterpenoid oridonin (**33**), isolated from *Isodon rubescens* (Hemsl.) H. Hara (Lamiaceae) (previously known as *Rabdosia rubescens*) induced apoptotic effects to human melanoma A375-S2 cells ($IC_{50} = 15 \mu M$) by induction of caspases-3 and -8, increased expression of Bax without affecting Bcl-2, as well as the release of cytochrome C and downstream activation of caspase-9. It shown weak cytotoxicity against PBMC thus pointing out to a good therapeutic window. By contrast, ponidicin (**34**), which is present in the same plant and is structurally very close, was not as active against these melanoma cells and markedly inhibited the growth of PBMC under the same conditions. However, treatment with increased concentrations of oridonin ($137 \mu M$) for 12 h, exerted cytotoxicity by necrosis as measured by an


LDH activity-based assay. These results may be interpreted in the context of convergent apoptotic and necrotic pathways in cancer cells [62].

Further work established that oridonin induced A375-S2 cell apoptosis by activating p53 and ERK pathways in parallel, which explains the apoptotic cell death [63]. Recently its activity upon uveal cells of melanoma origin was also confirmed. The authors found a significantly increased expression of the proapoptotic Bcl-2 family protein Bim and this mechanism was confirmed by knockdown by small interfering RNA. Additionally, oridonin suppressed Fatty Acid Synthase (FAS) expression in uveal melanoma cells, also confirmed by enforced FAS expression by insulin partially rescuing the cells from oridonin-induced apoptosis, showing that inhibition of FAS also contributed to oridonin-mediated apoptosis [64].


3.9 Daphnane

The daphnane-type diterpene daphneresiniferins A (**35**) and B (**36**), yuanhuacine (**37**), yuanhuadine (**38**), yuanhuahine (**39**), genkwadaphnin (**40**), genkwanine A (**41**), genkwanine F (**42**), and genkwanine H (**43**), were isolated from the methanol extract of the flower buds of *Daphne genkwa* Siebold & Zucc. (Thymelaeaceae). All compounds were assayed for both their cytotoxicity and inhibitory effects of the melanogenesis in α -MSH-activated B16 melanoma cells, with IC_{50} values in the range 0.06 - $9.0 \mu M$, while arbutin and Kojic acid, used as controls presented IC_{50} values of 140 and $39 \mu M$, respectively. On the other hand, only daphneresiniferin B showed significant cytotoxicity on B16 melanoma cells, with an IC_{50} value of $6.6 \mu M$ [65].


Gnidimacrin (**44**), isolated from roots of *Stellera chamaejasme* L (Thymelaeaceae), a plant used in Traditional Chinese Medicine (TCM), was active *in vivo* on B16 bearing mice at doses of $0.06 \mu g/kg/day$ [66]. Hirsein A (**45**) and B (**46**), and gnidilatin (**47**) from *Thymelaea hirsuta* (L.) Endl. were capable of lowering melanin content in B16 cells without reduction in cell viability. Hirsein A, at $0.1 \mu M$, reducing the melanin content by 37%, while hirsein B reduced the melanin content by 26 %. The process involves the downregulation of the MITF gene, an important oncotarget in melanoma [67–69].




(21) camosol




(22) carnosic acid




(23) parvifloron D




(24) 11,12,16-trihydroxy-2-oxo-5-methyl-10-demethyl-abieta-1[10],6,8,11,13-pentene




(25) 7a-acetoxyroyleanone




(26) horminone




(27) (5R,8R,9S,13R)-halim-1,10-ene-15,16-diol



(28) ingenol



(29) ingenol mebutato



(30) 7,13-diacetyl-5-angeloyl-20nicotinyl-3-propionyl-1,2,6,7-tetrahydroingenol (DANPT)

Fig 4. Abietane-type, Halimane-type and Ingenol diterpenes with activity on melanoma cells




Fig 5. Kaurane-type and Daphnane-type diterpenes with activity on melanoma cells

3.10 Neoclerodane

Four neoclerodane-type diterpenes, isolated from *Ajuga decumbens* Thunb. (Lamiaceae), namely, ajugacumbin K (**48**), ajugacumbin L (**49**),


ajugacumbin M (**50**), and ajugacumbin N (**51**) were evaluated about cytotoxicity on B16 cell line. All compounds presented cytotoxicity on B16, with an IC₅₀ value ranging from 24.2 to 44.5 μM [70].

3.10 Others


Hypoestoxide (**52**), a verticillane diterpene found in *Hypoestes rosea* P. Beauv. (Acanthaceae), presented potent nonsteroidal anti-inflammatory effects, and exhibited highly powerful activity against B16 melanoma growth in C57BL/6 mice. At a low maximal effective dose of 5 mg/kg, hypoestoxide was able to induce significant *in vivo* antitumor activity that was better than or comparable with most of the standard chemotherapeutic antiangiogenic agents

cortisone acetate, vincristine, bleomycin, adriamycin, 5-fluorouracil, cyclophosphamide, and etoposide, except vincristine [71].


The compound vouacapane-6 α ,7 β ,14 β ,19-tetraol (**53**), isolated from *Pterodon pubescens* Benth. seeds presented *in vitro* cytotoxicity on SK MEL 37 melanoma cells in a dose-dependent way, with IC₅₀ values of 32 μ M, similar to doxorubicin (IC₅₀ = 35 μ M) under identical assay conditions [72].




(48) ajugacumbin K




(49) ajugacumbin L



(50) ajugacumbin M



(51) ajugacumbin N



(52) hypoestoxide (HE)


(53) vouacapane-6 α ,7 β ,14 β ,19-tetraol

Fig 6. Miscellaneous plant-derived diterpenes with activity on melanoma cells

4 DITERPENES FROM MARINE ORGANISMS

4.1 Cembranolides

Cembrane-type diterpenoids can be found in terrestrial and marine organisms, e.g. soft corals [73,74] and plants [75,76].

Sinularin (**46**), isolated from the soft coral *Sinularia flexibilis* Quoy and Gaimard (Alcyoniidae)

inhibited A2058 melanoma cell proliferation (IC₅₀ = 5 μ g/mL), migration, and G2/M phase arrest. The apoptotic effect was evidenced by proteomic data and Western blot displaying the levels of several associated proteins including annexin A1, voltage - dependent anion - selective channel protein 1 and prohibitin (upregulated), heat shock protein 60, heat shock protein beta-1, and peroxiredoxin-2 (downregulated) in A2058 melanoma cells exposed to

sinularin. It increased the expression of p53, cleaved-caspase-3, cleaved-caspase-8, cleaved-caspase-9, p21, and Bax, and decreased expression of Bcl-2 [77].

Sarcophytol A (**47**) and sarcophine (**48**), from the soft coral *Sarcophyton glaucum* Quoy and Gaimard (Alcyoniidae), are naturally occurring cembranolides with potent chemopreventive activity [78]. Their low yield in nature sparked a long-term interest in the synthesis of analogues with similar properties [78–84].

4.2 Norcembranoids

From *Sinularia gardneri* (Pratt) (Alcyoniidae) was obtained, besides other compounds,

a diterpene with norcembranoid skeleton, singardin (**49**), that presented cytotoxicity against MEL-28, in the concentration of 5 µg/mL [85].

4.3 Biscembranoids

Sarcophytolide M (**50**) was active in the low micromolar range to SK-Mel2, followed by sarcophytolide N (**51**), sarcophytolide I (**52**), lobophytone J (**53**) sarcophytolide L (**54**) lobophytone O (**55**), lobophytone U (**56**), methyl totuoate A (**57**) methyl tortuoate B (**58**) and methyl sartortuoate (**59**) all isolated from the Vietnamese soft coral *Sarcophyton pauciplicatum* (Alcyoniidae) [86].




Fig 7. Cembrane-type diterpenes from marine organisms with activity on melanoma cells.

4.4 Furanoditerpenes

The activity of the marine furanoditerpenes, spongiatriol (**60**) and epispongiatriol (**61**), was evaluated in MM96E and MM96L - two sublines of human melanoma cells derived from the same metastatic lesion. The two strains were very similar although MM96E presented higher tyrosinase activity and lower expression of alkaline phosphatase than MM96L. Spongiatriol caused dendritic morphology in a proportion of MM96L cells whilst epispongiatriol was able to induce a marked G2/M arrest in MM96E cells. Treatment with cytostatic doses of both diterpenes for 72 h inhibited B8G3 expression and tyrosinase activity with no significant effect on the expression of tyrosinase antigen, with MM96L cells more affected than MM96E. It is believed that these subtle differences in human melanoma phenotypes, is correlated with the expression of different neurotransmitter receptors [87].

4.5 Enhygromic Acids

Enhygromic acid (**62**), isolated from a marine myxobacterium, *Enhygromyxa* sp. exhibited cytotoxicity against B16 melanoma cells [88].

4.6 Spatanes

Spatane diterpenoids isolated from the brown marine algae *Stoechospermum marginatum* (C. Agardh) Kuntzing, such as 5(R),19-diacetoxy-15,18(R and S)-dihydrospata-13,16(E)-diene (**63**), induce apoptosis in B16F10 melanoma cells more selectively than in THP1, U937, COLO205, and HL60 cells. The generation of ROS induces unfavourable changes in the Bax/Bcl-2 ratio, eventually disrupting the mitochondrial $\Delta\Psi_m$. Cytochrome C leaks to the cytoplasm and activates the caspase-mediated apoptotic pathway. Also, there is a deregulation of the PI3K/AKT signalling pathway. Cell arrest occurs at

the S-phase arrest in the cell cycle. This diterpene effectively inhibited tumour growth in C57BL/6 mice bearing B16F10 melanoma without apparent toxic effects [89].

4.7 Lissoclimides

Chlorinated lissoclimide-type diterpenes can be isolated from the mollusc species *Pleurobranchus* (Pleurobranchidae). These diterpenes are presumably metabolites of a *Lissoclinum* species of sea squirts (family Didemnidae) on which the molluscs use to feed. Chlorolissoclimide (**64**) and dichlorolissoclimide (**65**) were found to be potent cytotoxins in the National Cancer Institute's screening panel of 60 tumour cell lines and showed some selectivity for melanomas.

4.8 Briaranes

Diterpenes with briarane skeleton can be found in soft corals, nudibranchs, and sponges [90]. Tubiporein (**67**), isolated from soft coral *Tubipora* sp, was able to inhibit the growth of B-16 mouse melanoma cells at the $IC_{50} = 2 \mu\text{g/mL}$ [91].

5 DITERPENES FROM FUNGI

Sphaeropsidin A (**66**) is a fungal bioactive secondary metabolite, with a rearranged primarane skeleton with $\text{Na}^+/\text{K}^+ / 2\text{Cl}^-$ cotransporter and/or anion exchanger inhibitor activities. It can overcome apoptosis as well as multidrug resistance in SKMEL-28 and mouse B16F10 by inducing a marked and rapid cellular shrinkage related to the loss of intracellular Cl^- and the decreased HCO_3^- concentration in the culture supernatant [92].




Fig 8. Other miscellaneous diterpenes from marine organisms and fungi with activity on melanoma cells.

6 COMBINATION THERAPIES

The management of patients with metastatic malignant melanoma remains difficult. Conventional chemotherapy has been disappointingly ineffective. Early work in going beyond cytotoxic monotherapy saw taxanes combined with DTIC, temozolomide, cisplatin, carboplatin and tamoxifen. The results are suggesting that they are at least as effective as various other combination regimens [93].

The combination of recombinant human fibroblast interferon (INF- δ) and mezerein (**68**) (itself a PKC activator) results in a synergistic antiproliferative effect on human melanoma cells accompanied by an increase in melanin synthesis [94]. Mezerein was first described as a major toxic principle of *Daphne mezereum* L. (Thymelaeaceae) [95].

One molecular target for this activity is human gamma interferon (IFN- γ) binding and Class I HLA and melanoma-associated antigen (MAA) expression in the HO-1 human melanoma cell line. Studies using HO-1 demonstrated the superiority of the combination Mezerein + IFN- δ . This synergistic combination also increased the expression of HLA Class I antigens but decreased the expression of the high molecular weight-melanoma associated antigen (HMW-MAA) [94].

In vitro combinations of sphaeropsidin A (Sph A) (**66**) with either cisplatin or temozolomide, showed that combining 4 μ M Sph A with 75 μ M cisplatin for 72 h had a synergistic cytotoxic effect independent of BRAF or NRAS mutations [96].

Tanshinone (**69**) isolated from *Radix Astragali* acts synergistically with carboplatin to inhibit B16 cell growth after excessive UV radiation by inducing apoptosis [97].

A combination therapy that uses both triptolide (**04**) and VNP20009 (an antitumour factor

from *Salmonella*) significantly improved the tumour colonisation of VNP20009 by reducing both the number of infiltrated neutrophils and the expression of VEGF synergistically, thus retarding the growth of the melanoma [98].




Fig 9. Examples of diterpenes can act synergistically with drugs used in melanoma therapy

7 FUTURE DIRECTIONS

The commercialisation of Paclitaxel (Taxol®) and docetaxel (Taxotere®) did not discourage researchers to continue designing and synthesising new taxane analogues, and some 9-beta-dihydrobaccatin-9,10-acetals exhibited potent antitumor effects against B16 melanoma BL6 *in vivo* by both iv and p.o. administration [99]. Similarly, the synthesis of derivatives from the above reviewed natural diterpenes is a logical ‘next step’ which is actively pursued by a number of groups working on the ingenol skeleton [100,101], leelamine [102], andrographolide [103], the cembranolides from the soft coral *Sarcophyton glaucum* [78–84]. This approach depends on a reliable source for the core skeleton, and diterpenes are notoriously minor compounds in most cases. Cell culture in bioreactors may afford a more reliable and sustainable supply such in the case of the bioactive abietane diterpenes from *Salvia sclarea* (Lamiaceae) which synthesis *in vitro* can be by elicited transcriptional reprogramming of their hairy roots [106]. Total synthesis is always complicated and

usually the last resort but nevertheless possible and always with the advantage of giving a total control on the quality, purity and supply of the drug. Earlier developments tapped into close precursors in high quantities such as communic acid (**70**) –up to 50% w/w in the cones of some conifers- to synthesise podolactone-type diterpenes. In this manner, Barrero and co-workers obtained some bioactive derivatives with one-digit micromolar to submicromolar range IC_{50} s (LL-Z1271 α , **71**) [107,108].

However, modern combinatorial metabolic engineering of microbial or plant hosts, using the enzymes of the biosynthetic pathways, offers an alternative environmentally benign access to structurally complex diterpenoids with specific and novel stereochemistry, which defines their therapeutic value. Using such approach, the group of Prof Hamberger achieved the stereoselective biosynthesis of over 50 diterpene skeletons, including natural variants and new enantiomeric or diastereomeric counterparts and demonstrated its scalable biotechnological production in engineered strains of *Saccharomyces cerevisiae* [109].




Fig. 10. Podolactone-type diterpenes derived from communic acids

8 CONCLUDING REMARKS

Diterpenoids can still be a source of lead compounds for future treatments either preventative or curative in melanoma. Given the enormous diversity of this class of compounds, and that much of them are still to be discovered in marine organisms or fungi, the availability of data seems scarce. Moreover, not all reports on diterpenoids show activity against melanoma, for example, some jatrophone diterpenoid esters [110] and even – a powerful and well-established DNA polymerase inhibitor – and a series of semisynthetic derivatives were not of significant activity on melanoma cells [111]. There is a need to understand the structure-activity relationship when more expedite procedures for the synthesis of these molecules become mainstream as reviewed in the previous section.

Still what we already know shows great potential. The most studied compounds are the labdanes which have given several potential blockbusters: forskolin, triptolide, carnolic acid, carnosol and andrographolide have both potentials for industrial sourcing and a well-known favourable safety profiles. It is andrographolide which is well ahead of the race with several developers putting forward patented derivatives for serious clinical development. The long time necessary for the development and approval of anticancer compounds

makes any forecast difficult, but the case of ingenol is an enormous encouragement for researchers in the field. There is still room for natural products to beat purely synthetic drugs providing new mechanisms of action and better treatments. They are just waiting for us out there.

CONFLICT OF INTEREST

Authors declare no conflict of interest.

ACKNOWLEDGEMENTS

This work has been possible thanks to a Visiting Scientist Grant (‘Pesquisador Visitante’) awarded to the correspondence author by The Brazilian National Council for Scientific and Technological Development (CNPq).

9 REFERENCES

- [1] Roncero AM, Tobal IE, Moro RF, Marcos IS. Halimane diterpenoids: sources, structures, nomenclature and biological activities. *Nat Prod Rep* 2018; 35,955.
- [2] Palathinkal DM, Sharma TR, Koon HB, Bordeaux JS. Current Systemic Therapies for Melanoma. *Dermatologic Surg* 2014;40(9):948–63.

- [3] Kwasniak LA, Garcia-Zuazaga J. Basal cell carcinoma: evidence-based medicine and review of treatment modalities. *Int J Dermatol* 2011;50(6):645–58.
- [4] Kuphal S, Bosserhoff A. Recent progress in understanding the pathology of malignant melanoma. *J Pathol* 2009;219(4):400–9.
- [5] Davies H, Bignell GR, Cox C, Stephens P, Edkins S, Clegg S, et al. Mutations of the BRAF gene in human cancer. *Nature* 2002;417(6892):949–54.
- [6] Milagre C, Dhomen N, Geyer FC, Hayward R, Lambros M, Reis-Filho JS, et al. A mouse model of melanoma driven by oncogenic KRAS. *Cancer Res* 2010;70(13):5549–57.
- [7] Curtin JA, Fridlyand J, Kageshita T, Patel HN, Busam KJ, Kutzner H, et al. Distinct Sets of Genetic Alterations in Melanoma. *N Engl J Med* 2005;353(20):2135–47.
- [8] Cancer Genome Atlas Network TCGA. Genomic Classification of Cutaneous Melanoma. *Cell* 2015;161(7):1681–96.
- [9] Arkenau H-T, Kefford R, Long G V. Targeting BRAF for patients with melanoma. *Br J Cancer* 2011;104(3):392–8.
- [10] Gray-Schopfer V, Wellbrock C, Marais R. Melanoma biology and new targeted therapy. *Nature* 2007;445(7130):851–7.
- [11] Gibney GT, Smalley KSM. An Unholy Alliance: Cooperation between BRAF and NF1 in Melanoma Development and BRAF Inhibitor Resistance. *Cancer Discov* 2013;3(3):260–3.
- [12] Müller J, Krijgsman O, Tsoi J, Robert L, Hugo W, Song C, et al. Low MITF/AXL ratio predicts early resistance to multiple targeted drugs in melanoma. *Nat Commun* 2014;5:5712.
- [13] Cirenajwis H, Lauss M, Ekedahl H, Törngren T, Kvist A, Saal LH, et al. NF1-mutated melanoma tumors harbor distinct clinical and biological characteristics. *Mol Oncol* 2017;11(4):438–51.
- [14] Kiuru M, Busam KJ. The NF1 gene in tumor syndromes and melanoma. *Lab Invest* 2017;97(2):146–57.
- [15] Moll UM, Petrenko O. The MDM2-p53 interaction. *Mol Cancer Res* 2003;1(14):1001–8.
- [16] Llanos S, Clark PA, Rowe J, Peters G. Stabilization of p53 by p14ARF without relocation of MDM2 to the nucleolus. *Nat Cell Biol* 2001;3(5):445–52.
- [17] Agrawal A, Yang J, Murphy RF, Agrawal DK. Regulation of the p14ARF-Mdm2-p53 pathway: An overview in breast cancer. *Exp Mol Pathol* 2006;81(2):115–22.
- [18] Berger MF, Garraway LA. Applications of genomics in melanoma oncogene discovery. *Hematol Oncol Clin North Am* 2009;23(3):397–414, vii.
- [19] Guterres AN, Herlyn M, Villanueva J. Melanoma. In: eLS. Chichester, UK: John Wiley & Sons, Ltd; 2018. page 1–10.
- [20] Rees JL. Melanoma: What Are the Gaps in Our Knowledge? *PLoS Med* 2008;5(6):e122.
- [21] Palmieri G, Capone M, Ascierto M, Gentilcore G, Stroncek DF, Casula M, et al. Main roads to melanoma. *J Transl Med* 2009;7(1):86.
- [22] Bogenrieder T, Herlyn M. The molecular pathology of cutaneous melanoma. *Cancer Biomarkers* 2011;9(1–6):267–86.
- [23] Shi Y. Mechanisms of caspase activation and inhibition during apoptosis. *Mol Cell* 2002;9(3):459–70.
- [24] Bravo-Cordero JJ, Hodgson L, Condeelis J. Directed cell invasion and migration during metastasis. *Curr Opin Cell Biol* 2012;24(2):277–83.
- [25] van den Brink DM, Wanders RJA. Phytanic acid: production from phytol, its breakdown and role in human disease. *Cell Mol Life Sci* 2006;63(15):1752–65.
- [26] Islam MT, Ali ES, Uddin SJ, Shaw S, Islam MA, Ahmed MI, et al. Phytol: A review of biomedical activities. *Food Chem Toxicol* 2018;121:82–94.
- [27] Ko G-A, Cho SK. Phytol suppresses melanogenesis through proteasomal degradation of MITF via the ROS-ERK signaling pathway. *Chem Biol Interact* 2018;286:132–40.
- [28] Jo AR, Jeong H-S, Kim M-K, Yun H-Y, Baek KJ, Kwon NS, et al. Geranylgeranylacetone induces apoptosis via the intrinsic pathway in human melanoma cells. *Biomed Pharmacother* 2016;82:15–9.
- [29] Nigmatov AG, Serebryakov P. Synthesis of derivatives of polyprenylacetic acids and related isoprenoids (review). *Pharm Chem J* 1990;24(4):278–86.
- [30] Chen H, Hu Z. Improved Synthesis of Teprenone. *CHINESE J Pharm* 2004;35(8):449–50.
- [31] Sheppard JR, Lester B, Doll J, Buscarino C, Gonzales E, Corwin S, et al. Biochemical regulation of adenylate cyclase in murine melanoma clones with different metastatic properties. *Int J cancer* 1986;37(5):713–22.
- [32] Lester BR, Stadel JM, Buscarino C, Sheppard J, Greig RG, Poste G. Reconstitution of the Gs

- protein from B16 melanoma clones of high and low experimental metastatic potential into S49 cyc-membranes. *Biochem Biophys Res Commun* 1987;147(1):443–51.
- [33] Ormerod EJ, Hart IR. Different growth responses to agents which elevate cAMP in human melanoma cell lines of high and low experimental metastatic capacity. *Clin Exp Metastasis* 1989;7(1):85–95.
- [34] Gerst JE, Sole J, Mather JP, Salomon Y. Regulation of adenylate cyclase by beta-melanotropin in the M2R melanoma cell line. *Mol Cell Endocrinol* 1986;46(2):137–47.
- [35] Kiessling S, Beaulieu-Laroche L, Blum ID, Landgraf D, Welsh DK, Storch K-F, et al. Enhancing circadian clock function in cancer cells inhibits tumor growth. *BMC Biol* 2017;15(1):13.
- [36] Hung F-M, Chen Y-L, Huang A-C, Hsiao Y-P, Yang J-S, CHUNG M-T, et al. Triptolide induces S phase arrest via the inhibition of cyclin E and CDC25A and triggers apoptosis via caspase- and mitochondrial-dependent signaling pathways in A375.S2 human melanoma cells. *Oncol Rep* 2013;29(3):1053–60.
- [37] Chueh F-S, Chen Y-L, Hsu S-C, Yang J-S, Hsueh S-C, Ji B-C, et al. Triptolide induced DNA damage in A375.S2 human malignant melanoma cells is mediated via reduction of DNA repair genes. *Oncol Rep* 2013;29(2):613–8.
- [38] Jao H-Y, Yu F-S, Yu C-S, Chang S-J, Liu K-C, Liao C-L, et al. Suppression of the migration and invasion is mediated by triptolide in B16F10 mouse melanoma cells through the NF-kappaB-dependent pathway. *Environ Toxicol* 2016;31(12):1974–84.
- [39] Liu B, Zhang H, Li J, Lu C, Chen G, Zhang G, et al. Triptolide Downregulates Treg Cells and the Level of IL-10, TGF- β , and VEGF in Melanoma-Bearing Mice. *Planta Med* 2013;79(15):1401–7.
- [40] Rajagopal S, Kumar RA, Deevi DS, Satyanarayana C, Rajagopalan R. Andrographolide, a potential cancer therapeutic agent isolated from *Andrographis paniculata*. *J Exp Ther Oncol* 2003;3(3):147–58.
- [41] Lim JCW, Jeyaraj EJ, Sagineedu SR, Wong WSF, Stanslas J. SRS06, a New Semisynthetic Andrographolide Derivative with Improved Anticancer Potency and Selectivity, Inhibits Nuclear Factor-kB Nuclear Binding in the A549 Non-Small Cell Lung Cancer Cell Line. *Pharmacology* 2015;95(1–2):70–7.
- [42] Zhu P-Y, Yin W-H, Wang M-R, Dang Y-Y, Ye X-Y. Andrographolide suppresses melanin synthesis through Akt/GSK3 β / β -catenin signal pathway. *J Dermatol Sci* 2015;79(1):74–83.
- [43] Zhang Q-Q, Zhou D-L, Ding Y, Liu H-Y, Lei Y, Fang H-Y, et al. Andrographolide inhibits melanoma tumor growth by inactivating the TLR4/NF- κ B signaling pathway. *Melanoma Res* 2014;24(6):545–55.
- [44] Zhao M, Onakpa MM, Chen W-L, Santarsiero BD, Swanson SM, Burdette JE, et al. 17-Norpimaranes and (9 β H)-17-Norpimaranes from the Tuber of *Icacina trichantha*. *J Nat Prod* 2015;78(4):789–96.
- [45] Zhao M, Onakpa MM, Santarsiero BD, Chen W-L, Szymulanska-Ramamurthy KM, Swanson SM, et al. (9 β H)-Pimaranes and Derivatives from the Tuber of *Icacina trichantha*. *J Nat Prod* 2015;78(11):2731–7.
- [46] Monday Onakpa M, Zhao M, Gödecke T, Chen W-L, Che C-T, Santarsiero BD, et al. Cytotoxic (9 β H)-Pimarane and (9 β H)-17-Norpimarane Diterpenes from the Tuber of *Icacina trichantha*. *Chem Biodivers* 2014;11(12):1914–22.
- [47] Cardile V, Avola R, Graziano ACE, Piovano M, Russo A. Cytotoxicity of demalonyl thrysiflorin A, a semisynthetic labdane-derived diterpenoid, to melanoma cells. *Toxicol Vitr* 2018;47(2018):274–80.
- [48] Huang S-C, Ho C-T, Lin-Shiau S-Y, Lin J-K. Carnosol inhibits the invasion of B16/F10 mouse melanoma cells by suppressing metalloproteinase-9 through down-regulating nuclear factor-kappa B and c-Jun. *Biochem Pharmacol* 2005;69(2):221–32.
- [49] Cattaneo L, Cicconi R, Mignogna G, Giorgi A, Mattei M, Graziani G, et al. Anti-Proliferative Effect of *Rosmarinus officinalis* L. Extract on Human Melanoma A375 Cells. *PLoS One* 2015;10(7):e0132439.
- [50] Yamahara M, Sugimura K, Kumagai A, Fuchino H, Kuroi A, Kagawa M, et al. *Callicarpa longissima* extract, carnosol-rich, potently inhibits melanogenesis in B16F10 melanoma cells. *J Nat Med* 2016;70(1):28–35.
- [51] Alcaraz M, Achel DG, Olivares A, Olmos E, Alcaraz-Saura M, Castillo J. Carnosol, radiation and melanoma: a translational possibility. *Clin Transl Oncol* 2013;15(9):712–9.
- [52] Okubo A, Yasuhira S, Shibazaki M, Takahashi K, Akasaka T, Masuda T, et al. NAD(P)H dehydrogenase, quinone 1 (NQO1), protects melanin-producing cells from

- cytotoxicity of rhododendrol. *Pigment Cell Melanoma Res* 2016;29(3):309–16.
- [53] Park S, Song H, Sung M-K, Kang Y-H, Lee K, Park J. Carnosic Acid Inhibits the Epithelial-Mesenchymal Transition in B16F10 Melanoma Cells: A Possible Mechanism for the Inhibition of Cell Migration. *Int J Mol Sci* 2014;15(7):12698–713.
- [54] Silva CO, Molpeceres J, Batanero B, Fernandes AS, Saraiva N, Costa JG, et al. Functionalized diterpene parvifloron D-loaded hybrid nanoparticles for targeted delivery in melanoma therapy. *Ther Deliv* 2016;7(8):521–44.
- [55] Habtemariam S, Varghese GK. A Novel Diterpene Skeleton: Identification of a Highly Aromatic, Cytotoxic and Antioxidant 5-Methyl-10-demethyl-abietane-type Diterpene from *Premna serratifolia*. *Phyther Res* 2015;29(1):80–5.
- [56] Fronza M, Murillo R, Ślusarczyk S, Adams M, Hamburger M, Heinzmann B, et al. In vitro cytotoxic activity of abietane diterpenes from *Peltodon longipes* as well as *Salvia multiorrhiza* and *Salvia sahendica*. *Bioorg Med Chem* 2011;19(16):4876–81.
- [57] Silva CG, Santos Júnior HM, Barbosa JP, Costa GL, Rodrigues FAR, Oliveira DF, et al. Structure Elucidation, Antimicrobial and Cytotoxic Activities of a Halimane Isolated from *Vellozia kolbekii* Alves (Velloziaceae). *Chem Biodivers* 2015;12(12):1891–901.
- [58] Opferkuch HJ, Hecker E. New diterpenoid irritants from *euphorbia ingens*. *Tetrahedron Lett* 1974;15(3):261–4.
- [59] Izzi S, Sorgi P, Piemonte P, Carbone A, Frascione P. Successfully treated superficial basal cell carcinomas with ingenol mebutate 0.05% gel: Report of twenty cases. *Dermatol Ther* 2016;29(6):470–2.
- [60] Fallahian F, Ghanadian M, Aghaei M, Zarei SM. Induction of G2/M phase arrest and apoptosis by a new tetrahydroingenol diterpenoid from *Euphorbia erythradenia* Bioss. in melanoma cancer cells. *Biomed Pharmacother* 2017;86:334–42.
- [61] Sosa-Sequera MC, Chiurillo M, Moscoso J, Dolinar J, Suarez O, Neira N, et al. Kaurenic acid: Evaluation of the in vivo and in vitro antitumor activity on murine melanoma. *Indian J Pharmacol* 2011;43(6):683–8.
- [62] Nicotera P, Leist M, Single B, Volbracht C. Execution of Apoptosis: Converging or Diverging Pathways? *Biol Chem* 1999;380(9):1035–40.
- [63] Zhang C-L, Wu L-J, Zuo H-J, Tashiro S, Onodera S, Ikejima T. Cytochrome c release from oridonin-treated apoptotic A375-S2 cells is dependent on p53 and extracellular signal-regulated kinase activation. *J Pharmacol Sci* 2004;96(2):155–63.
- [64] Gu Z, Wang X, Qi R, Wei L, Huo Y, Ma Y, et al. Oridonin induces apoptosis in uveal melanoma cells by upregulation of Bim and downregulation of Fatty Acid Synthase. *Biochem Biophys Res Commun* 2015;457(2):187–93.
- [65] Bang KK, Yun C-Y, Lee C, Jin Q, Lee JW, Jung S-H, et al. Melanogenesis inhibitory daphnane diterpenoids from the flower buds of *Daphne genkwa*. *Bioorg Med Chem Lett* 2013;23(11):3334–7.
- [66] Yoshida M, Feng W, Saijo N, Ikekawa T. Antitumor activity of daphnane-type diterpene gnidimacrin isolated from *Stellera chamaejasme* L. *Int J Cancer* 1996;66(2):268–73.
- [67] Matsuyama K, Miyamae Y, Sekii Y, Han J, Abderrabba M, Morio T, et al. Effect of Mediterranean Medicinal Plant Extracts on Melanogenesis Regulation. *J Arid L Stud* 2009;19(1):387–90.
- [68] Villareal MO, Han J, Ikuta K, Isoda H. Daphnane diterpene hirsein B downregulates melanogenesis in B16 murine melanoma cells by cAMP pathway inhibition. *BMC Proc* 2011;5 Suppl 8(Suppl 8):P68.
- [69] Villareal MO, Han J, Ikuta K, Isoda H. Mechanism of Mitf inhibition and morphological differentiation effects of hirsein A on B16 melanoma cells revealed by DNA microarray. *J Dermatol Sci* 2012;67(1):26–36.
- [70] Chen H, Tang B-Q, Chen L, Liang J-Y, Sun J-B. Neo-clerodane diterpenes and phytoecdysteroids from *Ajuga decumbens* Thunb. and evaluation of their effects on cytotoxic, superoxide anion generation and elastase release in vitro. *Fitoterapia* 2018;129:7–12.
- [71] Ojo-Amaize EA, Nchekwube EJ, Cottam HB, Bai R, Verdier-Pinard P, Kakkanaiah VN, et al. Hypoestoxide, a natural nonmutagenic diterpenoid with antiangiogenic and antitumor activity: possible mechanisms of action. *Cancer Res* 2002;62(14):4007–14.
- [72] Vieira CR, Marques MF, Soares PR, Matuda L, de Oliveira CMA, Kato L, et al. Antiproliferative activity of *Pterodon pubescens* Benth. seed oil and its active principle on human melanoma cells. *Phytomedicine* 2008;15(6–7):528–32.

- [73] Zhao M, Cheng S, Yuan W, Xi Y, Li X, Dong J, et al. Cembranoids from a Chinese Collection of the Soft Coral *Lobophytum crassum*. *Mar Drugs* 2016;14(6).
- [74] Tani K, Kamada T, Phan C-S, Vairappan CS. New cembrane-type diterpenoids from Bornean soft coral *Nephthea* sp. with antifungal activity against *Lagenidium thermophilum*. *Nat Prod Res* 2018;1–7.
- [75] He X-F, Hou X-D, Ren X, Guo K, Li X-Z, Yan Z-Q, et al. Two new cembranic diterpenoids from the flowers of *Nicotiana tabacum* L. *Phytochem Lett* 2016;15:238–44.
- [76] Luo P, Yin Z-Y, Sun Z-J, Chen H, Li C-J, Zhou H-H, et al. Cembrane-type diterpenoids from *Macaranga pustulata*. *Fitoterapia* 2018;129:162–6.
- [77] Su T-R, Lin J-J, Chiu C-C, Chen JY-F, Su J-H, Cheng Z-J, et al. Proteomic investigation of anti-tumor activities exerted by sinularin against A2058 melanoma cells. *Electrophoresis* 2012;33(7):1139–52.
- [78] Ne'eman I, Fishelson L, Kashman Y. Sarcophine—a new toxin from the soft coral *Sarcophyton glaucum* (Alcyonaria). *Toxicon* 1974;12(6):593–IN6.
- [79] Kundoor V, Zhang X, Khalifa S, Fahmy H, Dwivedi C, Kundoor V, et al. A Possible Mechanism of Action of the Chemopreventive Effects of Sarcotriol on Skin Tumor Development in CD-1 Mice. *Mar Drugs* 2006;4(4):274–85.
- [80] Szymanski PT, Kuppast B, Ahmed SA, Khalifa S, Fahmy H. Sarcophine-Diol, a Skin Cancer Chemopreventive Agent, Inhibits Proliferation and Stimulates Apoptosis in Mouse Melanoma B16F10 Cell Line. *Mar Drugs* 2011;10(12):1–19.
- [81] Szymanski PT, Muley P, Ahmed SA, Khalifa S, Fahmy H. Sarcophine-Diol Inhibits Expression of COX-2, Inhibits Activity of cPLA2, Enhances Degradation of PLA2 and PLC γ 1 and Inhibits Cell Membrane Permeability in Mouse Melanoma B16F10 Cells. *Mar Drugs* 2012;10(12):2166–80.
- [82] Kundoor V, Zhang X, Khalifa S, Fahmy H, Drugs CD-M, 2006 U. A possible mechanism of action of the chemopreventive effects of sarcotriol on skin tumor development in CD-1 mice. mdpi.com
- [83] Zhang X, Kundoor V, Khalifa S, Zeman D, Fahmy H, Dwivedi C. Chemopreventive effects of sarcophine-diol on skin tumor development in CD-1 mice. *Cancer Lett* 2007;253(1):53–9.
- [84] Szymanski PT, Muley P, Ahmed SA, Khalifa S, Fahmy H, Szymanski PT, et al. Sarcophine-Diol Inhibits Expression of COX-2, Inhibits Activity of cPLA2, Enhances Degradation of PLA2 and PLC γ 1 and Inhibits Cell Membrane Permeability in Mouse Melanoma B16F10 Cells. *Mar Drugs* 2012;10(12):2166–80.
- [85] El Sayed KA, Hamann MT. A New Norcembranoid Dimer from the Red Sea Soft Coral *Sinularia gardineri*. *J Nat Prod* 1996;59(7):687–689.
- [86] Nam NH, Tung PT, Ngoc NT, Hanh TTH, Thao NP, Thanh N Van, et al. Cytotoxic Biscembranoids from the Soft Coral *Sarcophyton pauciplicatum*. *Chem Pharm Bull (Tokyo)* 2015;63(8):636–40.
- [87] Takahashi H, Schumann R, Quinn R, Briscoe TA, Parsons PG. Isomers of a marine diterpene distinguish sublines of human melanoma cells on the basis of apoptosis, cell cycle arrest and differentiation markers. *Melanoma Res* 1992;1(5–6):359–66.
- [88] Tomura T, Nagashima S, Yamazaki S, Iizuka T, Fudou R, Ojika M. An Unusual Diterpene—Enhygromic Acid and Deoxyenhygrolides from a Marine *Myxobacterium*, *Enhygromyxa* sp. *Mar Drugs* 2017;15(4):109.
- [89] Velatooru LR, Baggu CB, Janapala VR. Spatane diterpinoid from the brown algae, *Stoechospermum marginatum* induces apoptosis via ROS induced mitochondrial mediated caspase dependent pathway in murine B16F10 melanoma cells. *Mol Carcinog* 2016;55(12).
- [90] Sung P-J, Sheu J-H, Xu J-P. Survey of Briarane-Type Diterpenoids of Marine Origin. *Heterocycles* 2002;57(3):535.
- [91] Natori T, Kawai H, Fusetani N. Tubiporein, a novel diterpene from a Japanese soft coral *Tubipora* sp. *Tetrahedron Lett* 1990;31(5):689–90.
- [92] Mathieu V, Chantôme A, Lefranc F, Cimmino A, Miklos W, Paulitschke V, et al. Sphaeropsidin A shows promising activity against drug-resistant cancer cells by targeting regulatory volume increase. *Cell Mol Life Sci* 2015;72(19):3731–46.
- [93] Gogas H, Bafaloukos D, Bedikian AY. The role of taxanes in the treatment of metastatic melanoma. *Melanoma Res* 2004;14(5):415–20.
- [94] Fisher PB, Hermo H, Solowey WE, Dietrich MC, Edwalds GM, Weinstein IB, et al. Effect of recombinant human fibroblast interferon and mezerein on growth, differentiation, immune interferon binding and tumor

- associated antigen expression in human melanoma cells. *Anticancer Res* 1986;6(4):765–74.
- [95] Ronlán A, Wickberg B. The structure of mezerein, a major toxic principle of *Daphne mezereum* L. *Tetrahedron Lett* 1970;(49):4261–4.
- [96] Ingels A, Dinhof C, Garg AD, Maddau L, Masi M, Evidente A, et al. Computed determination of the in vitro optimal chemocombinations of sphaeropsidin A with chemotherapeutic agents to combat melanomas. *Cancer Chemother Pharmacol* 2017;79(5):971–83.
- [97] Wu J, Xu H, Zhang L, Zhang X. Radix Astragali and Tanshinone Help Carboplatin Inhibit B16 Tumor Cell Growth. *Technol Cancer Res Treat* 2016;15(4):583–8.
- [98] Chen J, Qiao Y, Tang B, Chen G, Liu X, Yang B, et al. Modulation of *Salmonella* Tumor-Colonization and Intratumoral Anti-angiogenesis by Triptolide and Its Mechanism. *Theranostics* 2017;7(8):2250–60.
- [99] Takeda Y, Uoto K, Iwahana M, Jimbo T, Nagata M, Atsumi R, et al. New highly active taxoids from 9 beta-dihydrobaccatin-9,10-acetals. Part 5. *Bioorg Med Chem Lett* 2004;14(12):3209–15.
- [100] Liang X, Grue-Sørensen G, Månsson K, Vedsø P, Soor A, Stahlhut M, et al. Syntheses, biological evaluation and SAR of ingenol mebutate analogues for treatment of actinic keratosis and non-melanoma skin cancer. *Bioorg Med Chem Lett* 2013;23(20):5624–9.
- [101] Grue-Sørensen G, Liang X, Månsson K, Vedsø P, Dahl Sørensen M, Soor A, et al. Synthesis, biological evaluation and SAR of 3-benzoates of ingenol for treatment of actinic keratosis and non-melanoma skin cancer. *Bioorg Med Chem Lett* 2014;24(1):54–60.
- [102] Gowda R, Inamdar GS, Kuzu O, Dinavahi SS, Krzeminski J, Battu MB, et al. Identifying the structure-activity relationship of leelamine necessary for inhibiting intracellular cholesterol transport. *Oncotarget* 2017;8(17):28260–77.
- [103] Dey SK, Bose D, Hazra A, Naskar S, Nandy A, Munda RN, et al. Cytotoxic Activity and Apoptosis-Inducing Potential of Di-spiropyrrolidino and Di-spiropyrrolidino Oxindole Andrographolide Derivatives. *PLoS One* 2013;8(3):e58055.
- [104] Fahmy H, Zjawiony J, Konoshima T, drugs HT-M, 2006 undefined. Potent skin cancer chemopreventing activity of some novel semi-synthetic cembranoids from marine sources. *Mar. Drugs* 2006; 4(3):28-36.
- [105] Fahmy H, Khalifa S, Konoshima T, Drugs JZ-M, 2004 undefined. An improved synthesis of 7, 8-epoxy-1, 3, 11-cembratriene-15R (α), 16-diol, a cembranoid of marine origin with a potent cancer chemopreventive activity. *Mar. Drugs* 2004;2(1):1-7.
- [106] Vaccaro MC, Mariaevelina A, Malafronte N, De Tommasi N, Leone A. Increasing the synthesis of bioactive abietane diterpenes in *Salvia sclarea* hairy roots by elicited transcriptional reprogramming. *Plant Cell Rep* 2017;36(2):375–86.
- [107] Barrero AF, Arseniyadis S, Quílez del Moral JF, Herrador MM, Valdivia M, Jiménez D. First synthesis of the antifungal oidiolactone C from trans-communic acid: cytotoxic and antimicrobial activity in podolactone-related compounds. *J Org Chem* 2002;67(8):2501–8.
- [108] Barrero AF, Sánchez JF, Elmerabet J, Jiménez-González D, Macías FA, Simonet AM. Enantiospecific syntheses of the potent bioactives nagilactone F and the mould metabolite LL-Z1271 α an evaluation of their allelopathic potential. *Tetrahedron* 1999;55(23):7289–304.
- [109] Andersen-Ranberg J, Kongstad KT, Nielsen MT, Jensen NB, Pateraki I, Bach SS, et al. Expanding the Landscape of Diterpene Structural Diversity through Stereochemically Controlled Combinatorial Biosynthesis. *Angew Chemie Int Ed* 2016;55(6):2142–6.
- [110] Kayser O, Kiderlen AF, Bertels S, Siems K. Antileishmanial activities of aphidicolin and its semisynthetic derivatives. *Antimicrob Agents Chemother* 2001;45(1):288–92.
- [111] Liu LG, Tan RX. New jatrophane diterpenoid esters from *Euphorbia turczaninowii*. *J Nat Prod* 2001;64(8):1064–8.