

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

Interpersonal Problems in Borderline Personality Disorder: Associations with Mentalizing,
Emotion Regulation, and Impulsiveness

Sebastian Euler

University of Basel, Switzerland

Tobias Nolte, Matthew Constantinou, and Julia Griem

University College London

Personality and Mood Disorder Research Network

P. Read Montague

Virginia Tech Carilion Research Institute

Peter Fonagy

University College London

Author Note

Sebastian Euler, Department of Psychiatry, University of Basel, and University College London; Tobias Nolte, Research Department of Clinical, Educational and Health Psychology, University College London; Matthew Constantinou, Research Department of Clinical, Educational and Health Psychology, University College London; Julia Griem, University College London, Research Department of Clinical, Educational and Health Psychology; P. Read Montague, Human Neuroimaging Lab, Virginia Tech Carilion Research Institute, and University College London; Peter Fonagy, Research Department of Clinical, Educational and Health Psychology, University College London.

[Add acknowledgements].

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

P. Read Montague is in receipt of and supported by a Wellcome Trust Principal Research Fellowship (091188/Z/10/Z), which also supported the study funding. Peter Fonagy is in receipt of a National Institute for Health Research (NIHR) Senior Investigator Award (NF-SI-0514-10157). Peter Fonagy was in part supported by the NIHR Collaboration for Leadership in Applied Health Research and Care (CLAHRC) North Thames at Barts Health NHS Trust. The views expressed are those of the authors and not necessarily those of the Wellcome Trust, the NHS, the NIHR or the UK Department of Health.

Correspondence concerning this article should be addressed to Sebastian Euler, Universitäre Psychiatrische Kliniken (UPK) Basel, Wilhelm Klein-Strasse 27, CH-4002 Basel, Switzerland. E-mail: sebastian.euler@upkbs.ch

Abstract

Interpersonal problems are a core symptom of borderline personality disorder (BPD). This study investigated the relationship between emotion dysregulation, impulsiveness, and impaired mentalizing in the context of predicting interpersonal problems in BPD. Two-hundred and ten patients with BPD completed the Difficulties in Emotion Regulation Scale (DERS), Barratt Impulsiveness Scale (BIS-11), Reflective Functioning Questionnaire (RFQ), and Inventory of Interpersonal Problems (IIP-32). We conducted three path models, with either mentalizing, emotion regulation, or impulsiveness as the exogenous variable. Emotion dysregulation and attentional impulsiveness predicted interpersonal problems directly, whereas hypomentalizing predicted interpersonal problems only indirectly throughout emotion dysregulation and attentional impulsiveness. Our results suggest that these domains contribute significantly to interpersonal problems in BPD. Moreover, hypomentalizing might impact on interpersonal problems via its effect on impulsiveness and emotion regulation. The study argues that focussing on emotion regulation and mentalizing in BPD treatments might have inter-linked beneficial effects on interpersonal problems.

Keywords: borderline personality disorder, interpersonal problems, mentalizing, reflective function, impulsiveness, emotion regulation

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

Borderline personality disorder (BPD) is a well-established and valid psychiatric diagnosis with biological and developmental roots (Chanen & Kaess, 2012) and a lifetime prevalence of 2–6% (Grant et al., 2008). Research has demonstrated favorable treatment response and improved symptomatic outcome (Cristea et al., 2017; Zanarini, Frankenburg, Reich, & Fitzmaurice, 2012). Despite this, patients with BPD often experience severe and persisting long-term interpersonal impairments (Choi-Kain, Zanarini, Frankenburg, Fitzmaurice, & Reich, 2010; Liebke et al., 2017; Skodol et al., 2005).

Interpersonal problems in BPD are defined by unstable and intense relationships with an alternation between idealization and devaluation, as well as efforts to avoid abandonment and high interpersonal sensitivity (American Psychiatric Association, 2013; Domes, Schulze, & Herpertz, 2009). Interpersonal problems have been suggested as the most characteristic and discriminative feature of the disorder (Fossati et al., 1999; Gunderson, 2007; Johansen, Karterud, Pedersen, Gude, & Falkum, 2004). Moreover, other cardinal symptoms of BPD such as anger, affective instability, suicidal behavior and impulsiveness mainly manifest within interpersonal contexts (Sharp, 2016). Evidence has shown that interpersonal problems underlie subjective burden and behavioral deviations (e.g., self-harm and violence), and that they are related to neurobiological alterations (King-Casas et al., 2008; Lis & Bohus, 2013; Stepp, Smith, Morse, Hallquist, & Pilkonis, 2012). Consequently, improvement of interpersonal relations is a core aim of effective therapies for BPD (Cristea et al., 2017).

One core domain of BPD which contributes to interpersonal problems is emotion dysregulation (Linehan, Davison, Lynch, & Sanderson, 2006). Emotion dysregulation in BPD comprises a lack of awareness and tolerance, inappropriate management and regulation of emotions, and disturbed behavior under distress (for review see Carpenter & Trull, 2013). It is widely accepted as a major component of the biosocial model, as well as being a central clinical feature of the disorder (Glenn & Klonsky, 2009; Putnam & Silk, 2005; Siever,

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

Torgersen, Gunderson, Livesley, & Kendler, 2002). Compared to other domains of BPD, affective instability and anger showed the lowest remission rates after 2 years, and affective problems (e.g., chronic anger) displayed higher recurrence rates in the long-term (McGlashan et al., 2005; Zanarini, Frankenburg, Reich, & Fitzmaurice, 2016).

The influence of emotion dysregulation on interpersonal problems has been assessed empirically. Affective instability was the only significant predictor of BPD negative relationship scores 2 years later (Tragesser, Solhan, Schwartz-Mette, & Trull, 2007) and was related to social maladjustment (Bagge et al., 2004). Emotion dysregulation upheld negative relationships over a 12-month period (Stepp et al., 2014). Interestingly, the association between overall BPD symptomatology and interpersonal problems has been shown to be fully mediated by emotion regulation (Herr, Rosenthal, Geiger, & Erikson, 2013). Finally, improvement in emotion regulation was associated with better treatment outcome such as improved interpersonal functioning (McMain, Guimond, Streiner, Cardish, & Links, 2012). In line with these findings, interpersonal problems improved during dialectical behavior therapy (DBT; Linehan, 1993a; 1993b), which explicitly focuses on the training of emotion regulation skills (McMain et al., 2013).

Emotion dysregulation is also closely related to impulsiveness, another core domain of BPD (Cackowski et al., 2014; Gratz et al., 2009; van Zutphen, Siep, Jacob, Goebel, & Arntz, 2015). For instance, self-destructive behavior is seen as an impulsive attempt to regulate emotional distress (Linehan, 1993a; Terzi et al., 2017; Tragesser et al., 2007). This indicates the presence of a phenomenological overlap between emotion dysregulation and impulsiveness. Nonetheless, both domains also contribute to the clinical manifestation of BPD as individual factors (Fossati, Gratz, Maffei, & Borroni, 2013, 2014; Tragesser & Robinson, 2009). Impulsiveness is defined as a predisposition to react rapidly and unplanned to internal or external stimuli without considering the negative consequences (Stanford et al.,

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

2009). Impulsiveness underlies self-harm and suicidal behavior (Paris, 2005; Tragesser & Robinson, 2009), and has been highlighted as a core domain of BPD with a critical long-term impact (Crowell, Beauchaine, & Linehan, 2009; Links, Heslegrave, & van Reekum, 1999; McGlashan et al., 2005; Siever et al., 2002). Impulsiveness is also related to interpersonal problems (Fossati et al., 1999; Koenigsberg et al., 2001) in terms of negative relationships (Tragesser & Robinson, 2009) and a higher number of stressful interpersonal life events (Powers, Gleason, & Oltmanns, 2013).

In addition to emotion dysregulation and impulsiveness, an impairment in mentalizing, that is the capacity to comprehend one's own and others' behavior in terms of intentional mental states, has been suggested as a further core domain of BPD (Fonagy, Luyten, & Allison, 2015).

Mentalizing is explicitly defined as a self-other-related construct. Impairments in mentalizing thus appear important for understanding deficits in interpersonal functioning in patients with BPD (Choi-Kain & Gunderson, 2008). For instance, the elevated sensitivity to interpersonal rejection found in patients with BPD is considered a failure of adequately mentalizing others' intentions (Euler et al., 2018; Herpertz & Bertsch, 2014). In line with this, empirical evidence suggests that a low mentalizing capacity is associated with interpersonal problems (Berenson et al., 2016; De Meulemeester, Lowyck, Vermote, Verhaest, & Luyten, 2017). Furthermore, the focus on mentalizing in mentalization-based treatment (MBT; Bateman & Fonagy, 2004) has been shown to significantly improve interpersonal problems (Bateman & Fonagy, 2009).

In sum, emotion dysregulation, impulsiveness, and impaired mentalizing are suggested as core domains of BPD, each with strong links to interpersonal problems. Given their unique but also overlapping contributions, the aim of this cross-sectional study was to determine how emotion regulation, impulsiveness and mentalizing relate to interpersonal

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

problems in BPD when they are considered simultaneously rather than individually. By using three path analytic models, we aimed to identify direct effects and reciprocal indirect effects of emotion dysregulation, impulsiveness, and impaired mentalizing on interpersonal problems.

Methods

Participants and Procedure

Two hundred and ten adult patients who met the DSM-IV criteria for BPD were selected from a larger study investigating social exchanges in BPD and antisocial personality disorder (ASPD, 80% female; $M_{age} = 32$, $SD_{age} = 10$; “Probing Social Exchanges – A Computational Neuroscience Approach to the Understanding of Borderline and Antisocial Personality Disorder, approved by the Research Ethics Committee for Wales, 12/WA/0283). Patients with a pre-existing clinical diagnosis of BPD or ASPD were referred by psychiatrists, care coordinators, and trainee clinical psychologists within personality disorder services of seven London NHS Mental Health Trusts across five London boroughs. Patients were interviewed by experienced and trained clinical psychologists using the Structured Clinical Interview for DSM-IV Axis II Diagnoses (First, Gibbon, Spitzer, Williams, & Benjamin, 1997). To ensure the reliability of the ratings, consensus conferences were regularly held and supervised by senior researchers. The SCID-II has shown adequate inter-rater and internal consistency reliability for the diagnosis of BPD in several studies (e.g., Lobbestael, Leurgans, & Arntz, 2011; Maffei et al., 1997).

A BPD diagnosis and English-language fluency were the inclusion criteria. Individuals with recent psychotic episodes, severe learning disabilities, or current or past neurological disorders or traumas were excluded. 63 of 210 BPD patients included in the current analysis (30%) met the DSM-IV criteria for ASPD which is similar to other clinical

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

samples (Widiger & Trull, 1993; Zanarini et al., 1998). Table 1 presents the sample's demographics. SCID-II interviews and self-report measures were completed within two to three assessments.

Measures

Inventory of Interpersonal Problems-32 (IIP-32). Interpersonal problems were measured using the IIP-32 (Horowitz, Alden, Wiggins, & Pincus, 2000). The IIP-32 consists of 32 items rated on a five-point Likert scale (*not at all to extremely*) with higher scores reflecting greater interpersonal problems. Items span a range of social behaviors that people find too hard to engage in (e.g., hard to make friends) or use too much (e.g., argue with other people too much). Similar to past studies with BPD samples, we used total scores ($\alpha = .75$) to measure the overall amount of interpersonal dysfunction (Bateman & Fonagy, 2009; Bohus et al., 2004).

Difficulties in Emotion Regulation Scale (DERS). The DERS is a 36-item questionnaire assessing problems in multiple domains of emotion regulation, including emotional awareness, emotional acceptance, engagement in goal-directed behavior, disengagement from emotionally charged impulsive behavior, and access to effective emotion regulation strategies (Gratz & Roemer, 2004). Items are rated on a five-point Likert scale (*Almost Never to Almost Always*) with higher scores reflecting greater emotion regulation difficulties. Clinical samples, including patients with BPD, show problems across a range of emotion-regulation domains, so we used total scores ($\alpha = .99$) to measure the extent of general problems in emotion regulation (Gratz, Rosenthal, Tull, Lejuez, & Gunderson, 2006; Tull, Barrett, McMillan, & Roemer, 2007).

Barratt Impulsiveness Scale-11 (BIS-11). The BIS-11 is a 30-item measure of impulsiveness across attentional (e.g., "I [do not] concentrate easily"), motor (e.g., "I act on the spur of the moment"), and future-oriented (e.g., "I [do not] plan for the future") domains

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

(Patton, Stanford, & Barratt, 1995). Items are rated on a four-point Likert scale (*Rarely/Never* to *Almost Always/Always*). The BIS-11 was designed to be multi-dimensional (Stanford et al., 2009). Since impulse control difficulties are also quantified in the DERS total score, BIS subscales were used in the current analysis with the intention to reduce the potential overlap between both measures. Past research in BPD samples supports the use of its subscales (Berlin, Rolls, & Iversen, 2005; Clarkin, Levy, Lenzenweger, & Kernberg, 2013; Fossati et al., 2004). We used the attention ($\alpha = .66$), motor ($\alpha = .67$), and non-planning subscales ($\alpha = .69$), with higher scores reflecting greater impulsiveness.

Reflective Functioning Questionnaire (RFQ). The RFQ is a 54-item measure of reflective functioning, the operationalized form of mentalizing (Fonagy et al., 2016). The RFQ assesses a subject's mentalizing capacity based on response patterns to questions concerning mental processes with respect to oneself and others. All items are scored on a 7-point Likert-type scale, ranging from "*completely disagree*" to "*completely agree*". Two validated subscales capture biases in mental-state reflection. These reflect a two-dimensional model of the process of mentalizing. The first subscale is *certainty about mental states* (RFQc), which positively scores for agreement with improbable (i.e., excessive) mental-state knowledge (e.g., "I know exactly what my close friends are thinking") and disagreement with probable mental-state knowledge (e.g., "I don't always know why I do what I do"). Items are subsequently rescored so that high scores on this scale reflect excessive reflective functioning, i.e., *hypermentalizing*. The second subscale refers to *uncertainty about mental states* (RFQu), and positively scores for agreement with deficient mental state reflection (e.g., "People's thoughts are a mystery to me") and disagreement with probable mental state reflection (e.g., "Understanding what's on someone else's mind is never difficult for me"). Items are subsequently rescored so that high scores on this scale reflect a lack of knowledge about mental states, i.e., *hypomentalizing*.

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

Both subscales showed excellent reliability (α RFQc = .87, α RFQu = .87)

Statistical Analysis

We used path analytic models to explore how emotion dysregulation, impulsiveness, and reflective functioning interact in their relationship with interpersonal problems. The statistical aim of a path analysis is to identify relationships among variables (Loehlin, 1998). Here, the path analytic models allowed the identification of indirect effects of emotion dysregulation, impulsiveness, and reflective functioning via each other in relation to interpersonal problems, rather than the unique contribution of each variable.

We performed three saturated path models with either DERS, BIS (CI, MI, and NP) or RFQ (RFQc and RFQu) as exogenous variables, and the remaining scales as endogenous variables (see figures 1-3). Interpersonal problems were always treated as the criterion variable. In a saturated model, all paths (i.e., unidirectional relationships) are specified.

Before estimating the path models, the DERS, RFQc, and RFQu were square-root transformed due to skewness and the robust maximum likelihood estimator was used in all analyses. Missing data were infrequent (30% of cases accounted for 3% of missing item-level observations) and missing completely at random (MCAR) according to Little's MCAR test ($\chi^2(7,075) = 7,177, p > .05$). Missing data were thus handled with multiple imputation in order to minimize bias associated with traditional deletion techniques (Enders, 2017; Graham, 2009). Thirty data sets were imputed at the item level and aggregated using Rubin's rules.

Results

All path models accounted for 36% of the variance in interpersonal problems, and were adjusted for covariates, including age, sex, years in education, and medication status (see table 3 for coefficients). Moreover, the endogenous variables that directly predicted interpersonal problems were allowed to correlate with each other.

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

Figure 1 shows the standardized estimates for a path model in which uncertainty and certainty in mental-state understanding were exogenous variables. Uncertainty in mental-state understanding ($\beta = .09, p > .05, 95\% \text{ CI } [-.09, .27]$) and certainty in mental-state understanding ($\beta = .07, p > .05, 95\% \text{ CI } [-.07, .20]$) did not directly predict interpersonal problems. Instead, greater uncertainty in mental-state understanding directly predicted more emotion-regulation difficulties ($\beta = .54, p < .001, 95\% \text{ CI } [.40, .67]$), higher attentional impulsiveness ($\beta = .45, p < .001, 95\% \text{ CI } [.28, .61]$), higher motor impulsiveness ($\beta = .37, p < .001, 95\% \text{ CI } [.22, .53]$), and higher non-planning ($\beta = .27, p < .01, 95\% \text{ CI } [.11, .44]$), while greater certainty in mental-state understanding directly predicted only higher motor impulsiveness ($\beta = .20, p < .05, 95\% \text{ CI } [.04, .36]$). In turn, more emotion-regulation problems ($\beta = .31, p < .001, 95\% \text{ CI } [.16, .46]$) and higher attentional impulsiveness ($\beta = .33, p < .001, 95\% \text{ CI } [.18, .47]$) directly predicted greater interpersonal problems in this model.

Therefore, significant indirect effects were found between uncertainty in mental-state understanding and interpersonal problems via emotion regulation problems ($\beta = .17, p < .001, 95\% \text{ CI } [.07, .26]$) and attentional impulsiveness ($\beta = .15, p < .001, 95\% \text{ CI } [.06, .23]$). In other words, the association between uncertainty in mental-state understanding and interpersonal problems was partially explained by problems in emotion regulation and attentional impulsiveness. There were no significant indirect effects between certainty in mental-state understanding and interpersonal problems.

We also estimated the model with the DERS (see Figure 2) or BIS subscales (see Figure 3) as exogenous variables. As an exogenous variable, the DERS strongly and significantly directly predicted interpersonal problems ($\beta = .31, p < .001, 95\% \text{ CI } [.16, .46]$). It also directly predicted uncertainty in mental-state understanding ($\beta = .54, p < .001, 95\% \text{ CI } [.42, .65]$), certainty in mental-state understanding ($\beta = -.22, p < .01, 95\% \text{ CI } [-.36, -.07]$),

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

attentional impulsiveness ($\beta = .49, p < .001, 95\% \text{ CI } [.36, .61]$), motor impulsiveness ($\beta = .32, p < .01, 95\% \text{ CI } [.19, .45]$), and non-planning ($\beta = .27, p < .001, 95\% \text{ CI } [.12, .42]$). The only significant indirect effect was found via attentional impulsiveness ($\beta = .16, p < .001, 95\% \text{ CI } [.08, .24]$); therefore, the association between emotion regulation and interpersonal problems was mainly direct.

In the model using BIS subscales as exogenous variables (see Figure 3), the attentional impulsiveness subscale positively and significantly directly predicted interpersonal problems ($\beta = .33, p < .001, 95\% \text{ CI } [.18, .47]$), uncertainty in mental-state understanding ($\beta = .25, p < .01, 95\% \text{ CI } [.09, .40]$), and emotion-regulation problems ($\beta = .36, p < .001, 95\% \text{ CI } [.24, .49]$). The motor impulsiveness subscale did not significantly predict any variable. The non-planning subscale did not directly predict interpersonal problems ($\beta = -.11, p > .05, 95\% \text{ CI } [-.27, .04]$), however it did directly predict greater uncertainty in mental state understanding ($\beta = .20, p < .05, 95\% \text{ CI } [.04, .36]$), and lower certainty in mental state understanding ($\beta = -.28, p < .001, 95\% \text{ CI } [-.42, -.13]$). The only significant indirect effects were found between attentional impulsiveness and interpersonal problems via emotion-regulation problems ($\beta = .11, p < .01, 95\% \text{ CI } [.04, .17]$).

Discussion

The current study aimed to investigate how emotion regulation, impulsiveness, and mentalizing relate to interpersonal problems in BPD when they are considered simultaneously rather than as unique contributors. Path analyses supported the notion that hypomentalizing, emotion dysregulation, and attentional impulsiveness were associated with interpersonal problems. When emotion dysregulation and attentional impulsiveness were placed as exogenous variables, they each directly predicted interpersonal problems. They also indirectly predicted interpersonal problems via each other, respectively. When acting as the

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

exogenous variable, hypomentalizing revealed no direct effect on interpersonal problems but indirectly predicted interpersonal problems via emotion dysregulation and attentional impulsiveness.

Our findings support the idea that emotion regulation and impulsiveness have important, yet unique roles in predicting interpersonal problems in BPD (e.g. Herr et al., 2013; Koenigsberg et al., 2001; Tragesser & Robinson, 2009). Nonetheless, the reciprocal indirect effects of emotion dysregulation and attentional impulsiveness in their influence on interpersonal problems support the suggestion of a fundamental interplay of the two traits (Turner, Sebastian, & Tüscher, 2017). It seems important that particularly attentional impulsiveness, reflecting attentional problems (i.e., problems “focusing on the task at hand”) and cognitive instability (i.e., “thought insertions and racing thoughts”), interacts with emotion dysregulation in its significance towards the manifestation of interpersonal problems. This finding extends previous knowledge about the particular role attentional impulsiveness plays in emotion dysregulation in BPD (Cackowski et al., 2014; Putnam & Silk, 2005). It further fits well with the biosocial model of the disorder which postulates that the reciprocal interaction between emotion dysregulation and impulsiveness is the primary cause of interpersonal problems in BPD (Crowell et al., 2009; Schmahl et al., 2014).

Overall, attentional impulsiveness, emotion dysregulation, and hypomentalizing were interrelated in their association with interpersonal problems in BPD. This is in line with the theory that hyperarousal in patients with BPD reduces inhibitory control (i.e., attentional/cognitive), which in turn is associated with inducing premature automatic processes of mentalizing (i.e., “non-mentalistic modes”) and impaired emotion regulation (Luyten & Fonagy, 2015; Nolte et al., 2013; Nolte, Campbell, & Fonagy, 2018, in press; Putnam & Silk, 2005). Consequently, interpersonal relationships are further compromised (Fonagy, Luyten, & Bateman, 2015)

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

Although these findings do not permit a causal attribution, the unique indirect association between hypomentalizing and interpersonal problems suggests that interpersonal problems induced by hypomentalizing might manifest through emotion dysregulation and attentional impulsiveness. This finding empirically substantiates the idea that “the defining characteristics of BPD – emotion dysregulation, impulsivity, interpersonal dysfunction” are “rooted in an instability of the reflective, regulatory capacities that mentalizing affords” (Fonagy, Luyten, & Bateman, 2015, p. 381). Our study extends this perspective empirically by showing that problems in mentalizing might manifest as interpersonal problems via difficulties in emotion processing. This finding therefore further strengthens the idea of mentalizing as a translational mechanism of change in treatment of BPD (Kramer, 2018; Sharp & Kalpakci, 2015).

Hypermentalizing, that is the cognitive over-attribution of mental states, did not predict interpersonal problems. This is in line with previous evidence which showed *hypomentalizing* as the major problem in adult patients with BPD (Badoud et al., 2017; Brüne, Walden, Edel, & Dimaggio, 2016; De Meulemeester, Vansteelandt, Luyten, & Lowyck, 2018; Lowyck et al., 2016; Perroud et al., 2017). Interestingly however, in adolescents with borderline traits, hypermentalizing rather than hypomentalizing appeared to be the significant clinical issue (Sharp et al., 2013). Further research is thus needed to explore these partly contradictory results. This also applies to the validity of the RFQc subscale since it seems insufficiently clear whether it measures adaptive or maladaptive aspects of mentalizing (De Meulemeester et al., 2018). The question of RFQc validity is further underlined by our results. These revealed a negative relationship between hypermentalizing and emotion dysregulation. Inconsistencies with previous studies concerning its association with impulsiveness further question this subscale validity: hypermentalizing predicted only motor impulsiveness in our study, contradicting the findings of Perroud et al. (2017). They

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

yielded a correlation of hypermentalizing as measured by the RFQc with all subscales of the BIS. Therefore, the importance of verifying the RFQc subscale as suitable for assessing mentalizing problems is evident.

Clinical Implications

Our findings are consistent with previous research and theories on the clinical phenomenology of BPD. They support the inclusion of both emotional lability and impulsiveness as diagnostic signatures in the hybrid model of the DSM-5, including the alternative model in Section III (criterion B; American Psychiatric Association, 2013). Simultaneously, the dimensional approach (criterion A) is also supported, since the assessment of functional impairments in self and interpersonal domains significantly overlaps with problems in mentalizing (Jeung & Herpertz, 2014). The indirect effects of emotion regulation, (attentional) impulsiveness, and hypomentalizing on interpersonal problems further suggest that focusing on these domains in psychotherapy of BPD might be promising to induce beneficial effects on interpersonal problems.

Limitations

In the current study, interpersonal problems were classified as symptoms whereas emotion dysregulation, impulsiveness, and mentalizing were conceptualized as potential underlying domains. We need to consider that suggestions to categorize personality disorder features as impairments, traits, and symptoms remain provisional (Herpertz et al., 2017). Critically stated, impulsiveness in terms of impulsive behavior may alternatively be described as a symptom of BPD, whereas it is also seen as a component of emotion dysregulation (e.g. within the DERS). However, in our study, we focus on impulsiveness as a bio-socially based psychological domain (American Psychiatric Association, 2013; Crowell et al., 2009; Stanford et al., 2009). To minimize the shared variance between both domains, we utilized the BIS subscales instead of the total score.

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

Only a third of the variance in interpersonal problems was explained by our measures. Emotion dysregulation, impulsiveness, and reflective functioning are closely related constructs, and although the path analysis was useful in uncovering indirect effects on interpersonal problems, it is likely that this truncated some of our findings. This has to be taken into account when interpreting our results. In this regard, other domains linked to interpersonal problems in BPD (e.g., identity diffusion; De Meulemeester et al., 2017) should be included in further investigations to explain more variance.

We have to acknowledge the comorbidity of ASPD in 30% of the patients in our sample. This ratio is similar to past studies reflecting clinical reality as previously shown (e.g., 30% reported by Widiger & Trull 1993 and 22.7% by Zanarini et al. 1998) . However, we recommend a comparison of BPD patients with other PD samples (e.g. ASPD) in future studies to confirm the specificity of our results for BPD.

Our conclusions have to be drawn tentatively because the study relies on self-reports, which are susceptible to biases (Kramer, 2017). The IIP, the DERS, and the BIS are, however, widely accepted and have been applied in many studies with BPD (Gratz, Moore, & Tull, 2016; McFarquhar, Luyten, & Fonagy, 2018; Rufino, Ellis, Clapp, Pearte, & Fowler, 2017; Sebastian, Jacob, Lieb, & Tuscher, 2013). The RFQ has been developed more recently and has been used with adolescents (RFQ-Youth; Ha, Sharp, Ensink, Fonagy, & Cirino, 2013; Sharp et al., 2013) and in some adult populations (Badoud et al., 2017; De Meulemeester et al., 2018; Perroud et al., 2017). However, its construct validity is challenged, especially with respect to the RFQc scale. Further investigations are thus needed to confirm if the RFQ is suitable for assessing problems in mentalizing.

In general, a combination of self-report and behavioural tasks, such as the Movie for the Assessment of Social Cognition (Dziobek et al., 2006), the emotional Paced Auditory Serial Addition Task (Gratz et al., 2006) for emotion regulation, and the Balloon Analogue

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

Risk Task (Lejuez et al., 2002) for impulse control might provide a more comprehensive assessment of the psychological domains used in this study.

Conclusion

The current study confirms that emotion dysregulation, attentional impulsiveness, and hypomentalizing are inter-linked intrinsic components of interpersonal problems in BPD. As such, it provides empirical evidence for the interplay of these domains, which are well captured in recent diagnostic approaches and highlighted within current effective psychotherapies for BPD. By extending previous research on the implication of impaired mentalizing for patients with BPD, the results suggest that interpersonal problems induced by hypomentalizing might manifest throughout emotion dysregulation and attentional impulsiveness.

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington: American Psychiatric Pub.
- Badoud, D., Prada, P., Nicastro, R., Germond, C., Luyten, P., Perroud, N., & Debbané, M. (2017). Attachment and reflective functioning in women with borderline personality disorder. *Journal of Personality Disorders*. doi:10.1521/pedi_2017_31_283
- Bagge, C. L., Nickell, A., Stepp, S. D., Durrett, C., Jackson, K., & Trull, T. J. (2004). Borderline personality disorder features predict negative outcomes 2 years later. *Journal of Abnormal Psychology, 113*(2), 279-288.
- Bateman, A. W., & Fonagy, P. (2004). Mentalization-based treatment of BPD. *Journal of Personality Disorders, 18*(1), 36-51.
- Bateman, A. W., & Fonagy, P. (2009). Randomized controlled trial of outpatient mentalization-based treatment versus structured clinical management for borderline personality disorder. *American Journal of Psychiatry, 166*(12), 1355-1364.
- Berenson, K. R., Dochat, C., Martin, C. G., Yang, X., Rafaeli, E., & Downey, G. (2016). Identification of mental states and interpersonal functioning in borderline personality disorder. *Personality Disorders*. doi:10.1037/per0000228
- Berlin, H. A., Rolls, E. T., & Iversen, S. D. (2005). Borderline personality disorder, impulsivity, and the orbitofrontal cortex. *American Journal of Psychiatry, 162*(12), 2360-2373.
- Bohus, M., Haaf, B., Simms, T., Limberger, M. F., Schmahl, C., Unckel, C., . . . Linehan, M. M. (2004). Effectiveness of inpatient dialectical behavioral therapy for borderline personality disorder: A controlled trial. *Behaviour Research & Therapy, 42*(5), 487-499.
- Brüne, M., Walden, S., Edel, M.-A., & Dimaggio, G. (2016). Mentalization of complex emotions in borderline personality disorder: The impact of parenting and exposure to trauma on the performance in a novel cartoon-based task. *Comprehensive Psychiatry, 64*, 29-37.

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

- Cackowski, S., Reitz, A.-C., Ende, G., Kleindienst, N., Bohus, M., Schmahl, C., & Krause-Utz, A. (2014). Impact of stress on different components of impulsivity in borderline personality disorder. *Psychological Medicine, 44*(15), 3329-3340.
- Carpenter, R. W., & Trull, T. J. (2013). Components of emotion dysregulation in borderline personality disorder: A review. *Current Psychiatry Reports, 15*(1), 335-343.
doi:10.1007/s11920-012-0335-2
- Chanen, A. M., & Kaess, M. (2012). Developmental pathways to borderline personality disorder. *Current Psychiatry Reports, 14*(1), 45-53. doi:10.1007/s11920-011-0242-y
- Choi-Kain, L. W., & Gunderson, J. G. (2008). Mentalization: Ontogeny, assessment, and application in the treatment of borderline personality disorder. *American Journal of Psychiatry, 165*(9), 1127-1135.
- Choi-Kain, L. W., Zanarini, M. C., Frankenburg, F. R., Fitzmaurice, G. M., & Reich, D. B. (2010). A longitudinal study of the 10-year course of interpersonal features in borderline personality disorder. *The Journal of Personality Disorders, 24*(3), 365-376.
doi:10.1521/pedi.2010.24.3.365
- Clarkin, J. F., Levy, K. N., Lenzenweger, M. F., & Kernberg, O. F. (2013). Evaluating three treatments for borderline personality disorder: A multiwave study. *Focus, 11*(2), 269-276.
- Cristea, I. A., Gentili, C., Cotet, C. D., Palomba, D., Barbui, C., & Cuijpers, P. (2017). Efficacy of psychotherapies for borderline personality disorder: A systematic review and meta-analysis. *JAMA Psychiatry, 74*(4), 319-328. doi:10.1001/jamapsychiatry.2016.4287
- Crowell, S. E., Beauchaine, T. P., & Linehan, M. M. (2009). A biosocial developmental model of borderline personality: Elaborating and extending linehan's theory. *Psychological Bulletin, 135*(3), 495-510.
- De Meulemeester, C., Lowyck, B., Vermote, R., Verhaest, Y., & Luyten, P. (2017). Mentalizing and interpersonal problems in borderline personality disorder: The mediating role of identity diffusion. *Psychiatry Research, 258*, 141-144.
- De Meulemeester, C., Vansteelandt, K., Luyten, P., & Lowyck, B. (2018). Mentalizing as a mechanism of change in the treatment of patients with borderline personality disorder: A

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

- parallel process growth modeling approach. *Personality Disorders: Theory, Research, and Treatment*, 9(1), 22-29.
- Domes, G., Schulze, L., & Herpertz, S. C. (2009). Emotion recognition in borderline personality disorder - a review of the literature. *Journal of Personality Disorders*, 23(1), 6-19.
doi:10.1521/pedi.2009.23.1.6
- Dziobek, I., Fleck, S., Kalbe, E., Rogers, K., Hassenstab, J., Brand, M., . . . Convit, A. (2006). Introducing MASC: A movie for the assessment of social cognition. *Journal of Autism and Developmental Disorders*, 36(5), 623-636.
- Enders, C. K. (2017). Multiple imputation as a flexible tool for missing data handling in clinical research. *Behaviour Research and Therapy*, 98, 4-18.
- Euler, S., Wrege, J., Busmann, M., Sollberger, D., Lang, U. E., & M., W. (2018). Exclusion-proneness in borderline personality disorder impairs alliance in mentalization-based group therapy. *Frontiers in Psychology*, 9(824).
- First, M. B., Gibbon, M., Spitzer, R. L., Williams, J. B., & Benjamin, L. S. (1997). *Structured clinical interview for DSM-IV axis II personality disorders, (SCID-II)*. Washington, D.C.: American Psychiatric Press.
- Fonagy, P., Luyten, P., & Allison, E. (2015). Epistemic petrification and the restoration of epistemic trust: A new conceptualization of borderline personality disorder and its psychosocial treatment. *The Journal of Personality Disorders*, 29(5), 575-609.
doi:10.1521/pedi.2015.29.5.575
- Fonagy, P., Luyten, P., & Bateman, A. W. (2015). Translation: Mentalizing as treatment target in borderline personality disorder. *Personality Disorders: Theory, Research, and Treatment*, 6(4), 380-392.
- Fonagy, P., Luyten, P., Moulton-Perkins, A., Lee, Y.-W., Warren, F., Howard, S., . . . Lowyck, B. (2016). Development and validation of a self-report measure of mentalizing: The reflective functioning questionnaire. *PLoS One*, 11(7), e0158678. doi:10.1371/journal.pone.0158678

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

- Fossati, A., Barratt, E. S., Carretta, I., Leonardi, B., Grazioli, F., & Maffei, C. (2004). Predicting borderline and antisocial personality disorder features in nonclinical subjects using measures of impulsivity and aggressiveness. *Psychiatry Research, 125*(2), 161-170.
- Fossati, A., Gratz, K. L., Maffei, C., & Borroni, S. (2013). Emotion dysregulation and impulsivity additively predict borderline personality disorder features in Italian nonclinical adolescents. *Personality and Mental Health, 7*(4), 320-333.
- Fossati, A., Gratz, K. L., Maffei, C., & Borroni, S. (2014). Impulsivity dimensions, emotion dysregulation, and borderline personality disorder features among Italian nonclinical adolescents. *Borderline Personality Disorder and Emotion Dysregulation, 1*(5). doi:10.1186/2051-6673-1-5
- Fossati, A., Maffei, C., Bagnato, M., Donati, D., Namia, C., & Novella, L. (1999). Latent structure analysis of DSM-IV borderline personality disorder criteria. *Comprehensive Psychiatry, 40*(1), 72-79.
- Glenn, C. R., & Klonsky, E. D. (2009). Emotion dysregulation as a core feature of borderline personality disorder. *Journal of Personality Disorders, 23*(1), 20-28.
- Graham, J. W. (2009). Missing data analysis: Making it work in the real world. *Annual Review of Psychology, 60*, 549-576.
- Grant, B. F., Chou, S. P., Goldstein, R. B., Huang, B., Stinson, F. S., Saha, T. D., . . . Ruan, W. J. (2008). Prevalence, correlates, disability, and comorbidity of DSM-IV borderline personality disorder: Results from the wave 2 national epidemiologic survey on alcohol and related conditions. *The Journal of Clinical Psychiatry, 69*(4), 533-545.
- Gratz, K. L., Moore, K. E., & Tull, M. T. (2016). The role of emotion dysregulation in the presence, associated difficulties, and treatment of borderline personality disorder. *Personality Disorders: Theory, Research, and Treatment, 7*(4), 344-353. doi:10.1037/per0000198
- Gratz, K. L., & Roemer, L. (2004). Multidimensional assessment of emotion regulation and dysregulation: Development, factor structure, and initial validation of the difficulties in emotion regulation scale. *Journal of Psychopathology and Behavioral Assessment, 26*(1), 41-54.

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

- Gratz, K. L., Rosenthal, M. Z., Tull, M. T., Lejuez, C. W., & Gunderson, J. G. (2006). An experimental investigation of emotion dysregulation in borderline personality disorder. *Journal of Abnormal Psychology, 115*(4), 850-855.
- Gratz, K. L., Tull, M. T., Reynolds, E. K., Bagge, C. L., Latzman, R. D., Daughters, S. B., & Lejuez, C. W. (2009). Extending extant models of the pathogenesis of borderline personality disorder to childhood borderline personality symptoms: The roles of affective dysfunction, disinhibition, and self-and emotion-regulation deficits. *Development and Psychopathology, 21*(4), 1263-1291.
- Gunderson, J. G. (2007). Disturbed relationships as a phenotype for borderline personality disorder. *The American Journal of Psychiatry, 164*(11), 1637-1640.
doi:10.1176/appi.ajp.2007.07071125
- Ha, C., Sharp, C., Ensink, K., Fonagy, P., & Cirino, P. (2013). The measurement of reflective function in adolescents with and without borderline traits. *Journal of Adolescence, 36*(6), 1215-1223.
- Herpertz, S. C., & Bertsch, K. (2014). The social-cognitive basis of personality disorders. *Current Opinion in Psychiatry, 27*(1), 73-77. doi:10.1097/ycp.0000000000000026
- Herpertz, S. C., Huprich, S. K., Bohus, M., Chanen, A., Goodman, M., Mehlum, L., . . . Sharp, C. (2017). The challenge of transforming the diagnostic system of personality disorders. *Journal of Personality Disorders, 31*(5), 577-589.
- Herr, N. R., Rosenthal, M. Z., Geiger, P. J., & Erikson, K. (2013). Difficulties with emotion regulation mediate the relationship between borderline personality disorder symptom severity and interpersonal problems. *Personality and Mental Health, 7*(3), 191-202.
- Horowitz, L. M., Alden, L. E., Wiggins, J. S., & Pincus, A. L. (2000). *Inventory of interpersonal problems (IIP-32/IIP-64)*. London: Psychological Corporation.
- Jeung, H., & Herpertz, S. C. (2014). Impairments of interpersonal functioning: Empathy and intimacy in borderline personality disorder. *Psychopathology, 47*(4), 220-234.
- Johansen, M., Karterud, S., Pedersen, G., Gude, T., & Falkum, E. (2004). An investigation of the prototype validity of the borderline DSM-IV construct. *Acta Psychiatrica Scandinavica, 109*(4), 289-298.

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

- King-Casas, B., Sharp, C., Lomax-Bream, L., Lohrenz, T., Fonagy, P., & Montague, P. R. (2008). The rupture and repair of cooperation in borderline personality disorder. *Science*, *321*(5890), 806-810.
- Koenigsberg, H. W., Harvey, P. D., Mitropoulou, V., New, A. S., Goodman, M., Silverman, J., . . . Siever, L. J. (2001). Are the interpersonal and identity disturbances in the borderline personality disorder criteria linked to the traits of affective instability and impulsivity? *Journal of Personality Disorders*, *15*(4), 358-370.
- Kramer, U. (2017). Personality, personality disorders, and the process of change. *Psychotherapy Research*, 1-13. doi:10.1080/10503307.2017.1377358
- Kramer, U. (2018). Mechanisms of change in treatments of personality disorders: Introduction to the special section. *Journal of Personality Disorders*, *32*(Supplement), 1-11.
- Lejuez, C. W., Read, J. P., Kahler, C. W., Richards, J. B., Ramsey, S. E., Stuart, G. L., . . . Brown, R. A. (2002). Evaluation of a behavioral measure of risk taking: The Balloon Analogue Risk Task (BART). *Journal of Experimental Psychology: Applied*, *8*(2), 75-84.
- Liebke, L., Bungert, M., Thome, J., Hauschild, S., Gescher, D. M., Schmahl, C., . . . Lis, S. (2017). Loneliness, social networks, and social functioning in borderline personality disorder. *Personality Disorders: Theory, Research, and Treatment*, *8*(4), 349-356.
- Linehan, M. M. (1993a). *Cognitive-behavioral treatment of borderline personality disorder*. New York: Guilford Press.
- Linehan, M. M. (1993b). *Skills training manual for treating borderline personality disorder*. New York: Guilford Press.
- Linehan, M. M., Davison, G. C., Lynch, T. R., & Sanderson, C. (2006). Technique factors in treating personality disorders. In L. G. Castonguay & L. E. Beutler (Eds.), *Principles of therapeutic change that work* (pp. 239-252). Oxford: Oxford University Press.
- Links, P. S., Heslegrave, R., & van Reekum, R. (1999). Impulsivity: Core aspect of borderline personality disorder. *Journal of Personality Disorders*, *13*(1), 1-9.
- Lis, S., & Bohus, M. (2013). Social interaction in borderline personality disorder. *Current Psychiatry Reports*, *15*(2), 338-345. doi:10.1007/s11920-012-0338-z

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

- Lobbestael, J., Leurgans, M., & Arntz, A. (2011). Inter-rater reliability of the Structured Clinical Interview for DSM-IV Axis I disorders (SCID I) and Axis II disorders (SCID II). *Clinical Psychology & Psychotherapy, 18*(1), 75-79.
- Loehlin, J. C. (1998). *Latent variable models: An introduction to factor, path, and structural analysis*. Mahwah, NJ: Lawrence Erlbaum Associates Publishers.
- Lowyck, B., Luyten, P., Vanwalleghem, D., Vermote, R., Mayes, L. C., & Crowley, M. J. (2016). What's in a face? Mentalizing in borderline personality disorder based on dynamically changing facial expressions. *Personality Disorders: Theory, Research, and Treatment, 7*(1), 72-79.
- Luyten, P., & Fonagy, P. (2015). The neurobiology of mentalizing. *Personality Disorders, 6*(4), 366-379. doi:10.1037/per0000117
- Maffei, C., Fossati, A., Agostoni, I., Barraco, A., Bagnato, M., Deborah, D., . . . Petrachi, M. (1997). Interrater reliability and internal consistency of the structured clinical interview for DSM-IV axis II personality disorders (SCID-II), version 2.0. *Journal of Personality Disorders, 11*(3), 279-284.
- McFarquhar, T., Luyten, P., & Fonagy, P. (2018). Changes in interpersonal problems in the psychotherapeutic treatment of depression as measured by the Inventory of Interpersonal Problems: A systematic review and meta-analysis. *Journal of Affective Disorders, 226*, 108-123.
- McGlashan, T. H., Grilo, C. M., Sanislow, C. A., Ralevski, E., Morey, L. C., Gunderson, J. G., . . . Bender, D. (2005). Two-year prevalence and stability of individual DSM-IV criteria for schizotypal, borderline, avoidant, and obsessive-compulsive personality disorders: Toward a hybrid model of axis II disorders. *American Journal of Psychiatry, 162*(5), 883-889.
- McMain, S. F., Guimond, T., Streiner, D. L., Cardish, R. J., & Links, P. S. (2012). Dialectical behavior therapy compared with general psychiatric management for borderline personality disorder: Clinical outcomes and functioning over a 2-year follow-up. *American Journal of Psychiatry, 169*(6), 650-661.

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

- McMain, S. F., Links, P. S., Guimond, T., Wnuk, S., Eynan, R., Bergmans, Y., & Warwar, S. (2013). An exploratory study of the relationship between changes in emotion and cognitive processes and treatment outcome in borderline personality disorder. *Psychotherapy Research, 23*(6), 658-673.
- Nolte, T., Bolling, D. Z., Hudac, C., Fonagy, P., Mayes, L. C., & Pelphrey, K. A. (2013). Brain mechanisms underlying the impact of attachment-related stress on social cognition. *Frontiers in Human Neuroscience, 7*(816). doi:10.3389/fnhum.2013.00816
- Nolte, T., Campbell, C., & Fonagy, P. (2018, in press). Social communicative processes in severe personality disorder. In V. Bizzari, J. Gonçalves, & J. G. Pereira (Eds.), *The Psychotherapy-Neurobiology-Pharmacology Intervention Triangle*. Malaga: Vernon Press.
- Paris, J. (2005). The development of impulsivity and suicidality in borderline personality disorder. *Development and Psychopathology, 17*(4), 1091-1104.
- Patton, J. H., Stanford, M. S., & Barratt, E. S. (1995). Factor structure of the Barratt Impulsiveness Scale. *Journal of Clinical Psychology, 51*(6), 768-774.
- Perroud, N., Badoud, D., Weibel, S., Nicastro, R., Hasler, R., Küng, A. L., . . . Aubry, J. M. (2017). Mentalization in adults with attention deficit hyperactivity disorder: Comparison with controls and patients with borderline personality disorder. *Psychiatry Research, 256*, 334-341.
- Powers, A. D., Gleason, M. E. J., & Oltmanns, T. F. (2013). Symptoms of borderline personality disorder predict interpersonal (but not independent) stressful life events in a community sample of older adults. *Journal of Abnormal Psychology, 122*(2), 469-474.
- Putnam, K. M., & Silk, K. R. (2005). Emotion dysregulation and the development of borderline personality disorder. *Development and Psychopathology, 17*(4), 899-925.
- Rufino, K. A., Ellis, T. E., Clapp, J., Pearte, C., & Fowler, J. C. (2017). Variations of emotion dysregulation in borderline personality disorder: A latent profile analysis approach with adult psychiatric inpatients. *Borderline Personality Disorder and Emotion Dysregulation, 4*(1), 17-26.
- Schmahl, C., Herpertz, S. C., Bertsch, K., Ende, G., Flor, H., Kirsch, P., . . . Schneider, M. (2014). Mechanisms of disturbed emotion processing and social interaction in borderline personality

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

- disorder: State of knowledge and research agenda of the German clinical research unit. *Borderline Personality Disorder and Emotion Dysregulation*, 1(12), 1-17.
- Sebastian, A., Jacob, G., Lieb, K., & Tuscher, O. (2013). Impulsivity in borderline personality disorder: A matter of disturbed impulse control or a facet of emotional dysregulation? *Current Psychiatry Reports*, 15(2), 339-347.
- Sharp, C. (2016). Current trends in BPD research as indicative of a broader sea-change in psychiatric nosology. *Personality Disorders: Theory, Research, and Treatment*, 7(4), 334-343.
- Sharp, C., Ha, C., Carbone, C., Kim, S., Perry, K., Williams, L., & Fonagy, P. (2013). Hypermentalizing in adolescent inpatients: Treatment effects and association with borderline traits. *Journal of Personality Disorders*, 27(1), 3-18.
- Sharp, C., & Kalpakci, A. (2015). Mentalization in borderline personality disorder: From bench to bedside. *Personality Disorders: Theory, Research, and Treatment*, 6(4), 347-355.
- Siever, L. J., Torgersen, S., Gunderson, J. G., Livesley, W. J., & Kendler, K. S. (2002). The borderline diagnosis III: Identifying endophenotypes for genetic studies. *Biological Psychiatry*, 51(12), 964-968.
- Skodol, A. E., Oldham, J. M., Bender, D. S., Dyck, I. R., Stout, R. L., Morey, L. C., . . . Grilo, C. M. (2005). Dimensional representations of DSM-IV personality disorders: Relationships to functional impairment. *American Journal of Psychiatry*, 162(10), 1919-1925.
- Stanford, M. S., Mathias, C. W., Dougherty, D. M., Lake, S. L., Anderson, N. E., & Patton, J. H. (2009). Fifty years of the Barratt Impulsiveness Scale: An update and review. *Personality and Individual Differences*, 47(5), 385-395.
- Stepp, S. D., Scott, L. N., Morse, J. Q., Nolf, K. A., Hallquist, M. N., & Pilkonis, P. A. (2014). Emotion dysregulation as a maintenance factor of borderline personality disorder features. *Comprehensive Psychiatry*, 55(3), 657-666.
- Stepp, S. D., Smith, T. D., Morse, J. Q., Hallquist, M. N., & Pilkonis, P. A. (2012). Prospective associations among borderline personality disorder symptoms, interpersonal problems, and aggressive behaviors. *Journal of Interpersonal Violence*, 27(1), 103-124.

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

- Terzi, L., Martino, F., Berardi, D., Bortolotti, B., Sasdelli, A., & Menchetti, M. (2017). Aggressive behavior and self-harm in borderline personality disorder: The role of impulsivity and emotion dysregulation in a sample of outpatients. *Psychiatry Research, 249*, 321-326.
- Tragesser, S. L., & Robinson, R. J. (2009). The role of affective instability and UPPS impulsivity in borderline personality disorder features. *Journal of Personality Disorders, 23*(4), 370-383.
- Tragesser, S. L., Solhan, M., Schwartz-Mette, R., & Trull, T. J. (2007). The role of affective instability and impulsivity in predicting future BPD features. *Journal of Personality Disorders, 21*(6), 603-614.
- Tull, M. T., Barrett, H. M., McMillan, E. S., & Roemer, L. (2007). A preliminary investigation of the relationship between emotion regulation difficulties and posttraumatic stress symptoms. *Behavior Therapy, 38*(3), 303-313.
- Turner, D., Sebastian, A., & Tüscher, O. (2017). Impulsivity and cluster B personality disorders. *Current Psychiatry Reports, 19*(3), 1-15. doi:10.1007/s11920-017-0768-8
- van Zutphen, L., Siep, N., Jacob, G. A., Goebel, R., & Arntz, A. (2015). Emotional sensitivity, emotion regulation and impulsivity in borderline personality disorder: A critical review of fMRI studies. *Neuroscience and Biobehavioral Review, 51*, 64-76.
doi:10.1016/j.neubiorev.2015.01.001
- Widiger, T. A., & Trull, T. J. (1993). Borderline and narcissistic personality disorders *Comprehensive handbook of psychopathology* (pp. 371-394). Boston, MA.: Springer.
- Zanarini, M. C., Frankenburg, F. R., Dubo, E. D., Sickel, A. E., Trikha, A., Levin, A., & Reynolds, V. (1998). Axis II comorbidity of borderline personality disorder. *Comprehensive Psychiatry, 39*(5), 296-302.
- Zanarini, M. C., Frankenburg, F. R., Reich, D. B., & Fitzmaurice, G. (2012). Attainment and stability of sustained symptomatic remission and recovery among patients with borderline personality disorder and axis II comparison subjects: A 16-year prospective follow-up study. *American Journal of Psychiatry, 169*(5), 476-483.

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

Zanarini, M. C., Frankenburg, F. R., Reich, D. B., & Fitzmaurice, G. M. (2016). Fluidity of the subsyndromal phenomenology of borderline personality disorder over 16 years of prospective follow-up. *American Journal of Psychiatry*, *173*(7), 688-694.

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

Table 1

Sample Demographics and Univariate Scale Descriptives

Demographic	Mean or <i>n</i>	SD or %	Range
Age (years)	32	10	18–58
Sex (female)	167	80%	
Ethnicity			
White ^a	156	75%	
Black/Black British ^b	17	8%	
Asian/British Asian ^c	17	8%	
Mixed	19	9%	
Employment Status			
Employed ^d	55	26%	
Unemployed	124	60%	
Student/Apprenticeship	24	12%	
Retired/Carer	4	2%	
Household Income (per annum)			
<£10k	103	51%	
£10-35k	72	36%	
>£35k	25	13%	
Medication Status (on medication)	133	63%	
Years in Education	14	4	4–26
BPD Symptom Count (SCID-II)	6.8	1.0	5–9
Scale			
IIP-32	68.1	20.4	17-162
DERS	138.0	22.9	14-178
BIS AI	22.4	4.3	2-32
BIS MI	28.2	5.4	17-42
BIS NP	30.9	5.4	16-43
RFQu	29.8	13.9	0-73
RFQc	12.8	10.8	0-57

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

Note. SD = standard deviation. SCID-II = Structured Clinical Interview for DSM-IV Axis II Disorders; BPD = Borderline Personality Disorder; IIP-32 = Inventory of Interpersonal Problems-32; DERS = Difficulties in Emotion Regulation Scale; BIS = Barratt Impulsiveness Scale; AI = Attentional Impulsiveness; MI = Motor Impulsiveness; NP = Non-planning Impulsiveness; RFQc = Reflective Functioning Questionnaire Certainty Subscale; RFQu = Reflective Functioning Questionnaire Uncertainty Subscale.

^aWhite = British, Irish, or Other. ^bBlack = Caribbean, African, or Other. ^cAsian = Indian, Pakistani, Bangladeshi, Chinese, or Other. ^dEmployed = full-time, part-time, self-employed or casual work.

Figure 1. Path diagram of a model directly predicting interpersonal problems (IIP) from uncertainty (RFQu) and certainty (RFQc) in mental-state understanding, and indirectly via emotion regulation (DERS), attentional (BIS AI), motor (BIS MI), and non-planning (BIS NP) impulsiveness. Standardized coefficients are shown; significant coefficients are in bold.

* $p < .05$. ** $p < .01$. *** $p < .001$.

Figure 2. Path diagram of a model directly predicting interpersonal problems (IIP) from difficulties in emotion regulation (DERS), and indirectly via certainty (RFQc) and uncertainty (RFQu) in mental-state understanding, attentional (BIS AI), motor (BIS MI), and non-planning (BIS NP) impulsiveness. Standardized coefficients are shown; significant coefficients are in bold. * $p < .05$. ** $p < .01$. *** $p < .001$.

Figure 3. Path diagram of a model directly predicting interpersonal problems (IIP) from attentional (BIS AI), motor (BIS MI), and non-planning (BIS NP) impulsiveness, and indirectly via certainty (RFQc) and uncertainty (RFQu) in mental-state understanding and

PREDICTORS OF INTERPERSONAL PROBLEMS IN BPD

difficulties in emotion regulation (DERS). Standardized coefficients are shown; significant coefficients are in bold. * $p < .05$. ** $p < .01$. *** $p < .001$.