

Little adaptive potential of a threatened passerine

Pierre de Villemereuil^{1,2,4,*}, Alexis Rutschmann¹, Kate D. Lee¹, John G. Ewen³,
Patricia Brekke^{3,†}, and Anna W. Santure^{1,†}

¹*School of Biological Sciences, University of Auckland, Auckland, NZ*

²*Centre d'Écologie Fonctionnelle et Évolutive, UMR 5175, 1919 Route de Mende, 34293 Montpellier
Cedex 5, France*

³*Institute of Zoology, Zoological Society of London, Regents Park, London, UK*

⁴*Lead Contact*

***Correspondence:** *Pierre de Villemereuil, E-mail: bonamy@horus.ens.fr*

†*These are joint last authors*

Summary

Threatened species face numerous threats, including future challenges triggered by global change. A possible way to cope with these challenges is through adaptive evolution, which requires adaptive potential. Adaptive potential is defined as the genetic variance needed to respond to selection and can be assessed either on adaptive traits or fitness [1]. However, a lack of high quality data has made it difficult to rigorously test adaptive potential in threatened species, leading to controversy over its magnitude [1–3]. Here we assess the adaptive potential of a threatened New Zealand passerine (the hihi, *Notiomystis cincta*) based on two populations: (i) the sole remaining natural population, on the island of Te Hauturu-o-Toi and (ii) a reintroduced population with a long-term dataset (intensively monitored for 20 years) based on the island of Tiritiri Matangi. We use molecular information (reduced representation genome sequencing, on both populations), as well as long-term phenotypic and fitness data from the Tiritiri Matangi population to find: (i) a lack of molecular genetic diversity at a genome-wide level in both populations, (ii) low heritability of traits under selection and (iii) negligible additive genetic variance of fitness in the Tiritiri Matangi population. In combination, these results support a lack of

17 adaptive potential in this threatened species. We discuss our findings within the context
18 of other passerines and methods for assessing adaptive potential; and the impact of these
19 results on conservation practice, for the hihi and species of conservation concern in general.

20 **Keywords:** adaptation, *Notiomystis cincta*, heritability, quantitative genetics, conservation genet-
21 ics, conservation biology

22 Results

23 The hihi (*Notiomystis cincta*, Du Bus, 1839) is an endemic New Zealand bird and the sole
24 representative of the *Notiomystidae* family [4]. Once common across the North Island, it is now
25 naturally occurring only on the island of Te Hauturu-o-Toi (Little Barrier Island, population
26 estimated around 1,000 - 3,000 individuals). As part of a concerted conservation effort, the hihi
27 has been successfully reintroduced to six mammalian predator-free sanctuaries, one of which is
28 situated on the island of Tiritiri Matangi (founded in 1995 by 53 reintroduced individuals and
29 currently stable at around 100-175 breeding individuals). Since its establishment all breeding
30 events have been intensively monitored, and every individual is systematically sampled and
31 nestlings banded and morphological traits measured [5], making this one of the largest and most
32 comprehensive long-term datasets of any threatened species. There is no dispersal in or out of
33 either of the studied islands.

34 Molecular diversity

35 Genomic sequence data from restriction site associated DNA (RAD) sequencing, a form of
36 reduced representation genomic sequencing, revealed low genetic diversity in hihi. Sequence
37 data was derived from 26 individuals from Te Hauturu-o-Toi and five individuals from Tiritiri
38 Matangi populations. The proportion of polymorphic sites within the RAD sequences was 0.36%
39 ($SE \pm 0.0091\%$) using both populations. Polymorphic sites were scarcer in Tiritiri Matangi
40 (0.22%, $SE \pm 0.0048\%$), when compared to the source population of Te Hauturu-o-Toi (0.34%,
41 $SE \pm 0.0047\%$), which might originate from the smaller sample size in Tiritiri Matangi, and/or the
42 bottleneck [6] when Tiritiri Matangi was established. The nucleotide diversity (π) was estimated
43 at 0.00095 ($SE \pm 4.64 \times 10^{-6}$) and was also lower in Tiritiri Matangi (0.00088, $SE \pm 4.35 \times 10^{-6}$)

44 than in Te Hauturu-o-Toi (0.00095 , $SE \pm 1.40 \times 10^{-5}$). Both of these molecular diversity metrics
45 were lower than any comparable (i.e. 'sequence-based') estimates for passerine birds we have
46 been able to identify in the literature [see ??; 7–11]. Note that while the only threatened species
47 (the Florida scrub-jay) included in our comparison has the lowest nucleotide diversity of the
48 other species, the hihi still has a diversity 2.6 times lower.

49 **Quantitative genetics of traits**

50 We performed a quantitative genetics analysis on eleven commonly measured avian traits that
51 were either directly measured or derived from the pedigree data in Tiritiri Matangi: three
52 morphological traits (mass, tarsus length and head-to-bill length, all measured at 21 days of
53 age, just before fledging), three life-history traits (longevity, probability of being recruited and
54 age at first reproduction) and five breeding traits (lay date, number of eggs laid, hatching and
55 fledgling success and time from egg laying to fledgling). To identify the traits under selection, we
56 computed the linear and non-linear selective gradients [12, 13]. We then used 'animal models',
57 a type of generalised linear mixed model, and pedigree information, accounting for confounding
58 effects and using appropriate statistical distributions for each trait, to compute the additive
59 genetic variance (V_A) and heritability (h^2) of all the traits. Estimates are provided on the
60 observed data scale, *sensu* [14]. Finally, we tested the ability of the sample size of our dataset
61 to correctly estimate a *small, but substantial* signal for the adaptive traits and fitness with a
62 simulation analysis. Detailed results are available in Table S1, Table S2 and Figure S3 of the
63 Supplementary Information.

64 **Selection gradients** All traits but time to fledge and age at first reproduction were under
65 selection, with a significant linear and/or non-linear standardised selection gradient (?? and
66 Table S1). The linear selection gradient was positive for all traits except for lay date (and age
67 at first reproduction, although the linear gradient was not significant) for which a negative
68 selection gradient was estimated. Except for adult longevity, all non-linear selection gradients
69 were negative, which suggests stabilising selection is operating, particularly when the parameter
70 was significant.

71 **Quantitative genetics of the studied traits** Estimated posterior modes for heritabilities and
72 additive genetic coefficients of variation of the three morphological traits were small (??, see also
73 Table S2 and Figure S2, fledgling mass: $h^2 = 0.0329$, additive genetic coefficient of variation
74 (CV_A) = 3.5%; tarsus length: $h^2 = 0.123$, $CV_A = 1.7\%$ and head-bill length: $h^2 = 0.0581$,
75 $CV_A = 1.2\%$). In contrast to the morphological traits, all but one (lay date) life-history and
76 breeding traits had an estimated posterior mode for the heritability below 10^{-3} (see ??, Table S2
77 and Figure S2). These extremely low estimates of heritability were driven by very low estimated
78 values of the additive genetic variance, with all lower bounds of the 95% credible interval below
79 10^{-7} . Although the signal for a non-zero additive genetic variance for lay date was stronger, the
80 point estimate (posterior mode $h^2 = 0.0289$) was still extremely low and the 95% credible interval
81 lower bound close to zero (see Table S2). Our simulation analysis (Figure S3) demonstrated
82 our data structure would be able to detect heritabilities as low as 0.1 with confidence for these
83 traits.

84 **Additive genetic variance of fitness**

85 The most direct measure of adaptive potential is the additive genetic variance of fitness. Here,
86 we used an animal model with a zero-inflated Poisson distribution to compute the additive
87 genetic variance of lifetime reproductive success of the individuals. This analysis suggests a
88 very low adaptive potential in hihi, with extremely small additive genetic variances of both the
89 zero-inflated and Poisson component of our model, resulting in extremely small heritabilities and
90 evolvabilities (see Table 1). The zero-inflated component had a relatively large population mean
91 of 0.759, which agrees well with the observed probability of not being recruited into the breeding
92 population (0.724). Our simulation analysis (see Figure S3) placed these estimates rather below
93 what would be considered a *small, but substantial* signal for additive genetic variance of fitness.
94 They also demonstrated an upward bias in the posterior median estimation for the evolvability
95 of the Poisson part, meaning that the true value of I_A is likely to lie between the posterior mode
96 (0.000587) and posterior median (0.01) shown in Table 1.

Discussion

Lack of adaptive potential in the hihi in comparison to other species The three measures of adaptive potential we employed here all support a lack of general and adaptive genetic diversity in the hihi. The levels of molecular genetic diversity found in this study are consistent with a depleted diversity compared to other passerine species (??). Comparing individual heterozygosity in pairs of threatened and non-threatened bird species, [15] similarly found a diversity roughly two to five times lower in threatened birds. Our average level of individual heterozygosity (0.00069, see Figure S1) falls well within the range observed by [15] for threatened bird species (0.0004 – 0.00091). More generally, when compared to estimates available in the database from [16], our nucleotide diversity seems typical of threatened species (average nucleotide diversity of 0.00115 for the 33 species classified as vulnerable, endangered or critically endangered), although the majority of such species are not passerines and might differ due to other features (such as overall fecundity) given the larger phylogenetic divergence. All of the traits studied here displayed small to inconsequential levels of additive genetic variance and heritability. The analysis of two morphological traits (tarsus length and head-bill length) yielded small heritabilities, with two other traits (body mass and lay date) having at best ambiguous support for heritabilities away from zero. The life-history and breeding traits (except lay date) were all characterised by extremely low posterior heritabilities, with all below a value of 0.05. Our simulation analysis confirms that our sample size is sufficient to accurately estimate low heritabilities, and that the majority of our heritabilities are well below what might be considered *small, but substantial* ($h^2 = 0.1$), and hence that adaptive potential is very limited. Furthermore, these estimates remain extremely low in the context of typical heritability values for other passerines found in the current literature. For example, using bird species data available from [17] to compute an average heritability value, we found estimates larger than those found in this study for body mass ($h^2 = 0.42$, 98% of the values above ours), tarsus length ($h^2 = 0.59$, 96%) and lay date ($h^2 = 0.149$, 90%). The additive genetic variance and related standardised measures were also very small for fitness. Only two other studies [18] used non-Gaussian animal models to measure and report the additive genetic variance of fitness [19, 20]. While they report variances on the latent scale, this is equivalent to our evolvability estimate [14]: the estimates

126 for song sparrow (*Melospiza melodia*) of [20] are considerably greater than ours (1.72 for males
127 and 2.01 for females), while the estimates for American red squirrel (*Tamiasciurus hudsonicus*)
128 of [19] are more comparable to ours (from 0.004 to 0.017, but note that their credible intervals
129 have much higher lower and upper bounds). While the additive genetic variance of relative
130 fitness is more related to an *instantaneous* rate of adaptation, rather than a potential for *future*
131 adaptation, it is one of the best proxies to predict the ability of a species to face environmental
132 challenges [18, 21], especially for threatened species such as hihi that are already shifted away
133 from their optimal habitat [22]. The lack of adaptive potential and reduced genetic diversity are
134 likely to be as a result of the bottleneck experienced during the collapse of the species, as well as
135 an extended period where the species consisted of a single population. The hihi was widespread
136 across the North Island of New Zealand before its rapid decline following European colonisation
137 with extirpation from the “mainland” by 1885 [6]. The range of the species was then restricted
138 to a sole surviving population on Te Hauturu-o-Toi (3083 Ha) for more than a century. Despite
139 changes in population size on Te Hauturu-o-Toi, hihi are thought to have retained a relatively
140 stable level of genetic diversity. This is primarily due to the largely promiscuous mode of mating
141 of the hihi, with an extremely high rate of extra-pair paternity [23, 24] and high natal dispersal
142 [25]. The remnant population is also currently estimated to be large and stable [26] and no trace
143 of a recent bottleneck was found using microsatellite data [6]. However, information about past
144 hihi diversity and the composition of Te Hauturu-o-Toi at the time of collapse are unfortunately
145 harder to infer.

146 **Relationship between molecular genetic diversity and adaptive potential** The lack of
147 molecular genetic diversity in the hihi differs from previous conclusions of [6], who found a
148 high genetic diversity, based on microsatellites, in the reintroduced populations and a small
149 reduction in genetic diversity following reintroduction events. This difference is likely because
150 [6] report *relative* rather than *absolute* diversity, i.e. here we include monomorphic sites. Our
151 approach allows for a direct between-species comparison of the levels of polymorphism and
152 related *per-site* nucleotide diversity. Nucleotide diversity will also be relatively slow to recover
153 from erosion due to a low per-nucleotide mutation rate compared to microsatellite markers. In
154 contrast, microsatellites provide information on the distribution of polymorphism (i.e. population
155 structure) rather than its actual level (although number of alleles can be used for this, genotyped

156 microsatellites are usually chosen to be polymorphic in the first place, leading to ascertainment
157 bias, [27]). Average *per-site* nucleotide diversity is also a more convincing molecular proxy of
158 adaptive potential, being closer to the definition of the *per-locus* additive genetic variance [28].
159 A meta-analysis [29] demonstrated little connection between adaptive potential and diversity
160 at molecular markers, which we suggest may be an artifact of ascertainment bias (e.g. using
161 *relative*, rather than *absolute*, diversity). While relative diversity might still be a useful measure
162 to study e.g. inbreeding issues [30–32], we thus advocate for the use of diversity estimates based
163 on sequence polymorphism as these are more closely related to adaptive potential.

164 **Consequences for conservation** Our results suggest that hihi currently do not have the
165 adaptive potential to evolve in response to the environmental threats the species is exposed
166 to (and, possibly, will be exposed to in the future), despite strong signs of selection detected
167 for all but two studied traits (see ?? and [22] for a detailed analysis of lay date). Despite
168 this, most hihi populations are currently showing good demographic health with supportive
169 management (including predator control and supplementary feeding). In particular, the hihi
170 population on Tiritiri Matangi is one of the most productive and currently being used as a
171 source of birds for translocation to other populations [5]. However, our results raise a number
172 of questions about the future management of threatened populations. A fundamental objective
173 of conservation management is to maximize the number and size of populations as quickly as
174 possible, thus avoiding heightened extinction risk stemming from stochastic processes in small
175 populations. Recovering species whose populations have become chronically small will likely
176 face similar problems of low adaptive potential, as we have reported for the hihi. However, the
177 management objective of increasing population size and numbers will ultimately also create
178 conditions which will allow regeneration of adaptive potential, albeit over a long period of
179 time. Theoretical models indicate that the level of genetic variance recovered over time will
180 depend on the population effective size N_e , although the time to equilibrium is also of N_e
181 generations [33, 34]. A reassuring property of the regeneration of genetic variation, however,
182 is that it is gradual (what is gained is gained, as long as conditions are stable) and quicker
183 at the beginning of the process [33]. Nonetheless, when managing rapid recovery from small
184 population sizes, conservation managers should be cautious of the unintentional genetic legacy
185 effects that may ultimately reduce population viability; i.e. increased inbreeding and genetic

186 drift. Both inbreeding accumulation and drift may be promoted, for example, by prioritising a
187 few prolific breeders in a chronically small population in contrast to managing for wider founder
188 representation [35, 36]. The costs and benefits of alternative recovery strategies (combining
189 genetic and non-genetic elements) should be projected as probabilistic extinction risks at suitably
190 long time scales to optimize management effort. Once populations have been recovered to a
191 larger size they will still likely require management support to protect or mitigate environmental
192 pressure in until such time as adaptive potential has been restored.

193 **Acknowledgements** We extend many thanks to the volunteers, past students and Department
194 of Conservation staff who have contributed to monitoring the Tiritiri Matangi hihi population
195 and to the Hihi Recovery Group, Department of Conservation, and Supporters of Tiritiri Matangi
196 for maintaining such a long term vision in monitoring and management of this population.
197 We acknowledge Ngati Manuhiri as Mana Whenua and Kaitiaki of Te Hauturu-o-Toi and its
198 taonga, including hihi. We thank three anonymous reviewers for their insightful comments
199 that improved the manuscript. We are very grateful to Nancy Chen and Charles Perrier
200 for calculating and providing diversity estimates for Florida scrub-jay and for the tit species
201 respectively. Finally, many thanks to Rachel Tucker and Jon Slate at the NERC Biomolecular
202 Analysis Facility, University of Sheffield, for completing the DNA extractions, and to Johanna
203 Nielsen and Kang-Wook Kim for help with microsatellite genotyping. PdV, AWS, KDL, PB
204 and JGE were supported by a Marsden Grant (UOA1408) awarded to AWS from the New
205 Zealand Royal Society Te Aparangi. AWS was also supported by a New Zealand National
206 Science Challenge Biological Heritage Project Grant, Project 1.4, and PB and the genotyping
207 were also supported by an AXA Fellowship. AR was supported by a grant from the Faculty
208 Research Development Fund of the University of Auckland.

209 **Author contributions** AWS and PB conceived of the study. JGE developed the demographic
210 dataset and supervised the data collection. PB developed the microsatellite dataset, supervised
211 the genotyping and performed the pedigree reconstruction. KDL performed the genome assembly.
212 PdV designed and conducted the analysis of the data, with advice from all other authors. PdV
213 wrote the paper, with input from all other authors.

214 **Ethical statement** Permissions to conduct research and collect blood samples on Tiritiri
215 Matangi and Te Hauturu-o-Toi were granted by the New Zealand Department of Conservation,
216 permit numbers 36186-FAU, 15073-RES, 24128-FAU, 13939-RES and 44300-FAU.

217 **Declaration of Interests** The authors declare no competing interests.

218 **References**

- 219 1 Hoffmann, A. A., Sgr, C. M., and Kristensen, T. N. (2017). Revisiting adaptive potential, population
220 size, and conservation. *Trends Ecol. Evol.* 32.7, 506–517.
- 221 2 Willi, Y., Buskirk, J. V., and Hoffmann, A. A. (2006). Limits to the adaptive potential of small
222 populations. *Annu. Rev. Ecol. Evol. Syst.* 37.1, 433–458.
- 223 3 Wood, J. L. A., Yates, M. C., and Fraser, D. J. (2016). Are heritability and selection related to
224 population size in nature? Meta-analysis and conservation implications. *Evol. Appl.* 9.5, 640–657.
- 225 4 Driskell, A., Christidis, L., Gill, B. J., Boles, W. E., Barker, F. K., and Longmore, N. W. (2007).
226 A new endemic family of New Zealand passerine birds: adding heat to a biodiversity hotspot. *Aust.*
227 *J. Zool.* 55.2, 73.
- 228 5 Armstrong, D. P. and Ewen, J. G. (2013). Consistency, continuity and creativity: long-term studies
229 of population dynamics on Tiritiri Matangi Island. *N. Z. J. Ecol.* 37.3, 288.
- 230 6 Brekke, P., Bennett, P. M., Santure, A. W., and Ewen, J. G. (2011). High genetic diversity in
231 the remnant island population of hihi and the genetic consequences of re-introduction. *Mol. Ecol.*
232 20.1, 29–45.
- 233 7 Chen, N., Hout, C. V. V., Gottipati, S., and Clark, A. G. (2014). Using Mendelian inheritance to
234 improve high-throughput SNP discovery. *Genetics* 198.3, 847–857.
- 235 8 Perrier, C., Campo, A. L. del, Szulkin, M., Demeyrier, V., Gregoire, A., and Charmantier, A.
236 (2018a). Great tits and the city: Distribution of genomic diversity and geneenvironment associations
237 along an urbanization gradient. *Evol. Appl.* 11.5, 593–613.
- 238 9 Perrier, C., Delahaie, B., and Charmantier, A. (2018b). Heritability estimates from genomewide
239 relatedness matrices in wild populations: Application to a passerine, using a small sample size.
240 *Mol. Ecol. Resour.* 18.4, 838–853.

- 241 10 Shultz, A. J., Baker, A. J., Hill, G. E., Nolan, P. M., and Edwards, S. V. (2016). SNPs across time
242 and space: population genomic signatures of founder events and epizootics in the House Finch
243 (*Haemorhous mexicanus*). *Ecol. Evol.* 6.20, 7475–7489.
- 244 11 Dutoit, L., Burri, R., Nater, A., Mugal, C. F., and Ellegren, H. (2017). Genomic distribution and
245 estimation of nucleotide diversity in natural populations: perspectives from the collared flycatcher
246 (*Ficedula albicollis*) genome. *Mol. Ecol. Resour.* 17.4, 586–597.
- 247 12 Lande, R. and Arnold, S. J. (1983). The measurement of selection on correlated characters.
248 *Evolution* 37.6, 1210–1226.
- 249 13 Morrissey, M. and Goudie, I. B. J. (2016). Analytical results for directional and quadratic selection
250 gradients for log-linear models of fitness functions. *bioRxiv*, 040618.
- 251 14 de Villemereuil, P., Schielzeth, H., Nakagawa, S., and Morrissey, M. B. (2016). General methods
252 for evolutionary quantitative genetic inference from generalised mixed models. *Genetics* 204.3,
253 1281–1294.
- 254 15 Li, S. et al. (2014). Genomic signatures of near-extinction and rebirth of the crested ibis and other
255 endangered bird species. *Genome Biology* 15.12, 557.
- 256 16 Robinson, J. A., Ortega-Del Vecchyo, D., Fan, Z., Kim, B. Y., vonHoldt, B. M., Marsden, C. D.,
257 Lohmueller, K. E., and Wayne, R. K. (2016). Genomic flatlining in the endangered island fox.
258 *Current Biology* 26.9, 1183–1189.
- 259 17 Postma, E. (2014). Four decades of estimating heritabilities in wild vertebrate populations:
260 Improved methods, more data, better estimates. *Quantitative Genetics in the Wild*. Ed. by A.
261 Charmantier, D. Garant, and L. E. B. Kruuk. Oxford (UK): Oxford University Press, pp. 16–33.
- 262 18 Hendry, A. P., Schoen, D. J., Wolak, M. E., and Reid, J. M. (2018). The contemporary evolution
263 of fitness. *Annu. Rev. Ecol. Evol. Syst.* 49.1, 457–476.
- 264 19 McFarlane, S. E., Gorrell, J. C., Coltman, D. W., Humphries, M. M., Boutin, S., and McAdam,
265 A. G. (2014). Very low levels of direct additive genetic variance in fitness and fitness components
266 in a red squirrel population. *Ecology and Evolution* 4.10, 1729–1738.
- 267 20 Wolak, M. E., Arcese, P., Keller, L. F., Nietlisbach, P., and Reid, J. M. (2018). Sex-specific additive
268 genetic variances and correlations for fitness in a song sparrow (*Melospiza melodia*) population
269 subject to natural immigration and inbreeding. *Evolution* 72.10, 2057–2075.
- 270 21 Shaw, R. G. (2018). From the past to the future: considering the value and limits of evolutionary
271 prediction. *The American Naturalist*.

- 272 22 de Villemereuil, P., Rutschmann, A., Ewen, J. G., Santure, A. W., and Brekke, P. (2018a). Can
273 threatened species adapt in a restored habitat? No expected evolutionary response in lay date for
274 the New Zealand hihi. *Evol. Appl.*
- 275 23 Ewen, J. G., Armstrong, D., and Lambert, D. M. (1999). Floater males gain reproductive success
276 through extrapair fertilizations in the stitchbird. *Anim. Behav.* 58.2, 321–328.
- 277 24 Brekke, P., Ewen, J. G., Clucas, G., and Santure, A. W. (2015). Determinants of male floating
278 behaviour and floater reproduction in a threatened population of the hihi (*Notiomystis cincta*).
279 *Evol. Appl.* 8.8, 796–806.
- 280 25 Richardson, K. M. and Ewen, J. G. (2016). Habitat selection in a reintroduced population: social
281 effects differ between natal and post-release dispersal. *Anim. Conserv.* 19.5, 413–421.
- 282 26 IUCN (2016). *Notiomystis Cincta: BirdLife International: The IUCN Red List of Threatened*
283 *Species 2017: E.T22704154A118814893*.
- 284 27 Hutter, C. M., Schug, M. D., and Aquadro, C. F. (1998). Microsatellite variation in *Drosophila*
285 *melanogaster* and *Drosophila simulans*: a reciprocal test of the ascertainment bias hypothesis.
286 *Mol. Biol. Evol.* 15.12, 1620–1636.
- 287 28 Lynch, M. and Walsh, B. (1998). *Genetics and Analysis of Quantitative Traits*. Sunderland,
288 Massachussets (US): Sinauer Associates.
- 289 29 Mittell, E. A., Nakagawa, S., and Hadfield, J. D. (2015). Are molecular markers useful predictors
290 of adaptive potential? *Ecol. Lett.* 18.8, 772–778.
- 291 30 Keller, L. F. and Waller, D. M. (2002). Inbreeding effects in wild populations. *Trends Ecol. Evol.*
292 17.5, 230–241.
- 293 31 Frankham, R. (2015). Genetic rescue of small inbred populations: meta-analysis reveals large and
294 consistent benefits of gene flow. *Mol. Ecol.* 24.11, 2610–2618.
- 295 32 Chen, N., Cosgrove, E. J., Bowman, R., Fitzpatrick, J. W., and Clark, A. G. (2016). Genomic
296 consequences of population decline in the endangered Florida scrub-jay. *Current Biology* 26.21,
297 2974–2979.
- 298 33 Lynch, M. and Hill, W. G. (1986). Phenotypic evolution by neutral mutation. *Evolution* 40.5,
299 915–935.
- 300 34 Houle, D. (1989). The maintenance of polygenic variation in finite populations. *Evolution* 43.8,
301 1767–1780.
- 302 35 Lacy, R. C. (1989). Analysis of founder representation in pedigrees: Founder equivalents and
303 founder genome equivalents. *Zoo Biol.* 8.2, 111–123.

- 304 36 Haig, S. M., Ballou, J. D., and Derrickson, S. R. (1990). Management options for preserving
305 genetic diversity: reintroduction of Guam rails to the wild. *Conserv. Biol.* 4.3, 290–300.
- 306 37 Castro, I., Minot, E. O., and Alley, J. C. (1994). Feeding and breeding behaviour of hihi or
307 stitchbirds *Notiomystis cincta* recently transferred to Kapiti Island, New Zealand, and possible
308 management alternatives. *Reintroduction Biology of Australian and New Zealand Fauna*. Ed. by
309 M. Serena. Chipping Norton, NSW: Surrey Beatty & Sons Pty Ltd, pp. 121–128.
- 310 38 Ewen, J. G. and Armstrong, D. P. (2000). Male provisioning is negatively correlated with attempted
311 extrapair copulation frequency in the stitchbird (or hihi). *Anim. Behav.* 60.4, 429–433.
- 312 39 Brekke, P., Cassey, P., Ariani, C., and Ewen, J. G. (2013). Evolution of extreme-mating behaviour:
313 patterns of extrapair paternity in a species with forced extrapair copulation. *Behav. Ecol. Sociobiol.*
314 67.6, 963–972.
- 315 40 Richardson, D. S., Jury, F. L., Blaakmeer, K., Komdeur, J., and Burke, T. (2001). Parentage as-
316 signment and extra-group paternity in a cooperative breeder: the Seychelles warbler (*Acrocephalus*
317 *sechellensis*). *Mol. Ecol.* 10.9, 2263–2273.
- 318 41 Brekke, P., Dawson, D. A., Horsburgh, G. J., and Ewen, J. G. (2009). Characterization of
319 microsatellite loci in the hihi *Notiomystis cincta* (*Notiomystidae, Aves*). *Mol. Ecol. Resour.* 9.4,
320 1255–1258.
- 321 42 Brekke, P., Wang, J., Bennett, P. M., Cassey, P., Dawson, D. A., Horsburgh, G. J., and Ewen, J. G.
322 (2012). Postcopulatory mechanisms of inbreeding avoidance in the island endemic hihi (*Notiomystis*
323 *cincta*). *Behav. Ecol.* 23.2, 278–284.
- 324 43 Andrews, S. (2014). *FASTQC: A Quality Control Tool for High Throughput Sequence Data*.
- 325 44 Bolger, A. M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina
326 sequence data. *Bioinformatics* 30.15, 2114–2120.
- 327 45 Luo, R. et al. (2012). SOAPdenovo2: an empirically improved memory-efficient short-read de novo
328 assembler. *GigaScience* 1.1, 18.
- 329 46 Baird, N. A., Etter, P. D., Atwood, T. S., Currey, M. C., Shiver, A. L., Lewis, Z. A., Selker, E. U.,
330 Cresko, W. A., and Johnson, E. A. (2008). Rapid SNP discovery and genetic mapping using
331 sequenced RAD markers. *PLoS ONE* 3.10, e3376.
- 332 47 Catchen, J. M., Amores, A., Hohenlohe, P., Cresko, W., and Postlethwait, J. H. (2011). Stacks:
333 Building and Genotyping Loci De Novo From Short-Read Sequences. *G3* 1.3, 171–182.
- 334 48 Li, H. and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
335 transform. *Bioinformatics* 25.14, 1754–1760.

- 336 49 Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association mapping
337 and population genetical parameter estimation from sequencing data. *Bioinforma. Oxf. Engl.*
338 27.21, 2987–2993.
- 339 50 Danecek, P. et al. (2011). The variant call format and VCFtools. *Bioinformatics* 27.15, 2156–2158.
- 340 51 Hadfield, J. D. (2010). MCMC methods for multi-response generalized linear mixed models: the
341 MCMCglmm R package. *J. Stat. Softw.* 33.2, 1–22.
- 342 52 Heidelberg, P. and Welch, P. D. (1981). A spectral method for confidence interval generation
343 and run length control in simulations. *Commun. ACM* 24.4, 233–245.
- 344 53 Plummer, M., Best, N., Cowles, K., and Vines, K. (2006). CODA: Convergence diagnosis and
345 output analysis for MCMC. *R News* 6.1, 7–11.
- 346 54 Low, M. and Prt, T. (2009). Patterns of mortality for each life-history stage in a population of
347 the endangered New Zealand stitchbird. *J. Anim. Ecol.* 78.4, 761–771.
- 348 55 de Villemereuil, P., Morrissey, M. B., Nakagawa, S., and Schielzeth, H. (2018b). Fixed effect
349 variance and the estimation of repeatabilities and heritabilities: Issues and solutions. *J. Evol. Biol.*
350 31.4, 621–632.
- 351 56 Houle, D. (1992). Comparing evolvability and variability of quantitative traits. *Genetics* 130.1,
352 195–204.
- 353 57 Hansen, T. F., Plabon, C., and Houle, D. (2011). Heritability is not evolvability. *Evol. Biol.* 38.3,
354 258.

Figure 1: Nucleotide diversity (π , red) and proportion of polymorphic sites (blue) for the hihi populations (Both pop: Both populations; Tiri. Mat.: Tiritiri Matangi; Hau.-o-Toi: Te Hauturu-o-Toi) along with estimates for other species (Florida scrub-jay *Aphelocoma coerulescens*, blue tit *Cyanistes caeruleus*, great tit *Parus major*, collared flycatcher *Ficedula albicollis*, house finch *Haemorhous mexicanus*, purple finch *Haemorhous purpureus*, Cassin's finch *Haemorhous cassinii*) identified in the literature [N. Chen, pers. comm. for the Florida Scrub-Jay data, C. Perrier, pers. comm. for the tit data, 7–11]. Nucleotide diversity for the collared flycatcher is a weighted average of autosomes and the Z-chromosome estimates from [11], considering the latter represents 6% of the genome. Error bars symbolising the standard error (computed using leave-one-out jackknife on the individuals) are provided for the hihi only. See more information on individual heterozygosity in Figure S1.

Figure 2: Point estimate and 95% credible/confidence intervals (CI) of the heritability (top-panel, point estimate is the posterior mode), linear (middle-panel) and non-linear standardised selection gradients (bottom-panel) for each of the studied traits. The selection gradients for the probability of recruitment are not calculated, as the fact that all non-recruits have a fitness of 0 creates numerical complications. Sample sizes are (from left to right) 2,098, 2,098, 2,098, 2,288, 2,183, 1,371, 581, 1,259, 1,383, 1,358 and 1,244. More details are available in Figure S2, Figure S3, Table S1, Table S2 and Table S3.

355

356

Table 1: Quantitative genetic analysis of fitness (lifetime reproductive success) using a zero-inflated Poisson. The results are shown separately for the zero-inflated (binomial) and the Poisson part of the model. All statistics (Pop. Mean : population mean, V_P : phenotypic variance, V_A : additive genetic variance, h^2 : heritability and I_A : evolvability) were computed on the observed data scale. Estimates are reported as posterior mode (posterior median) [95% credible interval].

Model part	Pop. Mean	V_P	V_A	h^2	I_A
Zero-infl. part	0.759 (0.76) [0.68 – 0.82]	0.183 (0.18) [0.15 – 0.22]	1.29E-5 (6E-4) [1.4E-11 – 0.0038]	0.00011 (0.0033) [7.8E-11 – 0.022]	3.21E-5 (0.001) [2.3E-11 – 0.0065]
Poisson part	8.14 (8.4) [5.9 – 12]	151 (146) [49 – 479]	0.0078 (0.73) [2.3E-10 – 5.7]	4.97E-5 (0.0045) [1.3E-12 – 0.034]	0.000587 (0.01) [3E-12 – 0.077]

358 **STAR★Methods**

359 **Contact for reagent and resource sharing**

360 Further information and requests for resources and reagents should be directed to and will be fulfilled
361 by the Lead Contact, Pierre de Villemereuil (bonamy@horus.ens.fr).

362 **Experimental model and subject details**

363 The remnant hihi population on Te Hauturu-o-Toi (Little Barrier Island; 3612'S, 17505'E) is largely
364 left unmanaged beyond ensuring the island stays mammalian predator free. Monitoring is limited. In
365 contrast, the hihi population on Tiritiri Matangi (3636'S, 17453'E) is intensively managed and has been
366 closely monitored since its establishment in 1995. All birds are individually identifiable (colour rings
367 added almost exclusively as nestlings) and all nesting attempts known, as hihi nest almost exclusively
368 in nest boxes. Dispersal is not possible between the populations. Hihi feed on a mix of fruits, nectar
369 and small invertebrates [37], but are also provided with supplementary food (20% by mass sugar water)
370 on Tiritiri Matangi. Hihi usually reproduce in their first year, during the austral spring and summer
371 [September to February, 37]. Females lay clutches ranging from three to five eggs and can produce
372 multiple clutches within a season although normally only one or two are successful. Despite males
373 providing around 30% of the care during rearing [38], extra-pair paternity in this species is widespread.
374 Around 60% of chicks within a brood are sired by extra-pair males [39]. Prior to whole genome and
375 RAD sequencing, hihi blood samples stored in ethanol were extracted at the Natural Environmental
376 Research Council Biomolecular Analysis Facility (Sheffield, United Kingdom) using an ammonium
377 acetate protocol [40], and quantified using a DNA fluorometer (Hoefer DynaQuant200) after being
378 assessed for quality on an agarose gel.

379 **Method details**

380 **Pedigree reconstruction** Systematic blood-sampling and genotyping at 18 microsatellite markers and
381 two sex-specific loci for Tiritiri Matangi individuals began in 2004 [41], allowing us to reconstruct the
382 paternities accounting for possible extra-pair copulations, following [42]. Because blood sampling was
383 only initiated in the 2003/2004 breeding season, information relating to the genetic sire of individuals
384 born previously is missing. For these individuals, we considered the information as missing, rather

385 than using the social sire.

386 **Genome assembly** Low coverage whole genome sequencing of ten birds (a subset of the samples used
387 in RAD sequencing below) was used to assemble a draft hihi genome, with seven of the samples sourced
388 from Te Hauturu-o-Toi and the remainder from Tiritiri Matangi. Samples were multiplexed and two
389 PCR-free DNA libraries were prepared by New Zealand Genomics Limited and used to generate 100bp
390 paired-end illumina reads over two lanes of Illumina HiSeq sequencing. This resulted in a total of
391 879,894,554 reads with a median of 89,508,541 per sample. Sequence quality was assessed using FastQC
392 [43]. Adapters and poor quality reads were removed with Trimmomatic-0.33 [44] under strict conditions
393 (ILLUMINACLIP:TruSeq3-PE-2.fa:2:30:10, LEADING:3, TRAILING:3, SLIDINGWINDOW:4:20,
394 MINLEN:70, CROP:110); over-represented reads identified in FastQC were also removed by appending
395 them to the TruSeq3-PE-2.fa file. Sample 10, an individual from Te Hauturu-o-Toi and the sample
396 with the most reads (126,397,278), was used to run SOAPdenovo2 version 1.5.14 [45] at kmer sizes
397 ranging from 25 - 95. The optimum kmer length of 36 bases was determined by N50 value and
398 length of assembly and was used in subsequent assemblies. Each of the samples was assembled using
399 SOAPdenovo2 with kmer 36 and insert size 210. Following assembly, sample 10 was also assessed to
400 have the highest quality assembly, with a total genome size of 1,002,019,675 bases, an N50 of 1928,
401 and 3,024,992 contigs ranging in length from 32 to 24,819 bases, and consequently this assembly was
402 chosen as the draft hihi reference genome.

403 **RAD sequencing** We used RAD sequencing [46] to obtain sequences from 26 individuals from Te
404 Hauturu-o-Toi and 5 individuals from Tiritiri Matangi. The unbalanced sampling is explained by the
405 fact that the RAD sequencing was initially used to detect SNPs to design a SNP chip, targeting the
406 diversity in the Te Hauturu-o-Toi source population. Samples were standardised to \simeq 50ng/uL and
407 then sent to Floragenex Inc. for RAD sequencing following Baird et al. [46]. Extracted DNA was
408 digested with SbfI, and barcodes and adaptors ligated. One sample was replicated, and a *Saccharomyces*
409 *bayanus* (yeast) control sample included, for a total of 33 independently indexed libraries. Libraries
410 were pooled for sequencing across two lanes of Illumina HiSeq 2000, generating 257,833,998 reads of
411 101 bases. The reads were processed and cleaned using the STACKS software [47] (`process_radtags`
412 with options `-e sbfI -c -q -s 15`). They were then mapped on the reference genome (Sample 10
413 above) using the `mem` procedure of the Burrows-Wheeler alignment tool [`bwa` with default options, 48].
414 Alignments were filtered for being mapped with a quality of 60. They were then stacked using the
415 `mpileup` of the `bcftools` suite [again using default parameters 49]. Sites with phred-scale quality

416 scores below 20 were discarded. Individual genotypes with read depth below 10 or allelic depth below
417 5 were considered as missing. Only sites without missing values were kept for each of the sub-samples
418 (total: 753,920 sites; only Te Hauturu-o-Toi: 855,845 sites; or only Tiritiri Matangi: 2,793,318). Our
419 results however are unchanged when using only sites with no missing values or allowing 30% of missing
420 values (maximum relative change of 5%).

421 **Phenotypic information** During the breeding season, all active nest boxes are inspected every few
422 days. The identity of the occupying male and female are recorded, along with the date of egg laying,
423 hatching and fledging, with the corresponding number of eggs/chicks each time. We used these data
424 to derive the number of eggs laid, lay date, hatching success (proportion of eggs hatched), fledging
425 success (proportion of fledged individuals) and time to fledge (time elapsed between lay date and
426 fledging date). Nestling morphological measures are taken (mass, tarsus length and head-to-bill length)
427 and individuals colour-ringed at 21 days of age. We used the pedigree information to derive the
428 following life-history traits: probability of being recruited into the breeding population, adult longevity
429 (number of years survived after the first breeding season post-fledging), and age at first reproduction.
430 Fitness was measured as the total number of offspring fledged over an individual's lifetime (lifetime
431 reproductive success, 0 for individuals that never bred) or from each clutch (number of fledglings) for
432 the breeding data. We removed individuals that were still alive and breeding, and therefore without a
433 complete lifetime reproductive success measure, from the individual fitness measures. Individual and
434 breeding data were collected from the 1995/1996 to 2014/2015, and from the 1997/1998 to 2014/2015
435 breeding seasons respectively.

436 **Quantification and statistical analysis**

437 **Genetic analysis** From the RAD sequencing data, the average *per-site* nucleotide diversity and
438 proportion of polymorphic sites were computed using VCFtools [50].

439 **Quantitative genetics models** To estimate quantitative genetic parameters on the Tiritiri Matangi
440 population, we used the phenotypic and pedigree information to run generalised linear mixed models,
441 known as animal models, using the R package MCMCg1mm [51]. For all traits, the fixed effects of
442 sex, fledgling mass (for individual data, i.e. morphological and life-history traits), clutch number
443 in the season, dam age and laying date (for breeding data) and clutch size (for both) were, when
444 relevant, tested for significance (using the pMCMC value inferred by MCMCg1mm) and included in the

445 mixed model if significant (see Table S2). For the individual traits, random effects included additive
446 genetic effect (using the pedigree information), identity of the dam, identity of the social sire, year
447 (i.e. year corresponding to the start of breeding season) and month of birth. For the breeding traits,
448 random effects included additive genetic effect, identity of the breeding female (accounting for repeated
449 measures), identity of the male mate and year (see above) of the breeding event. The error distributions
450 (Gaussian, Poisson with a log link, or binomial with a probit link) were chosen to fit each trait (see
451 Table S2). The number of iterations and the thinning interval were chosen for each model so as to
452 ensure that the MCMC effective sample sizes for all parameters of the model were above 1,000; and
453 were increased if this was not the case. As a result, our minimum MCMC effective sample size is 2,130.
454 Burn-in was set to a minimum of 3,000 iterations and convergence was checked using the Heidelberger
455 and Welch [52] convergence test as implemented in the `coda` R package [53]. Burn-in was increased
456 until convergence was reached. Whenever possible, we performed father-son, mother-daughter and
457 parent-offspring regressions. For breeding traits, only mother-daughter regressions were possible. Such
458 regressions were not possible for the probability of recruitment and hatching and fledgling success, as
459 there was no variation in the parent population since, by definition, all were recruited. Finally, we
460 performed a quantitative genetic analysis of fitness (measured as lifetime reproductive success as stated
461 above), in order to measure the adaptive potential in its strictest sense (i.e. standardised measure of
462 additive genetic variance of fitness). To do so, we used a zero-inflated Poisson model with independent
463 parameters on the latent traits corresponding to the zero-inflated binomial and Poisson processes.
464 The zero-inflation is mainly due to low survival to reproduction [54]. A model including genetic and
465 environmental correlations between the two latent traits did not yield significant correlations.

466 **Computation of the quantitative genetic parameters** From the output of the animal models, quan-
467 titative genetic parameters (population mean, phenotypic variance V_P , additive genetic variance V_A and
468 heritability h^2) were computed using the `QGglmm` R package [14], using the relevant error distribution
469 for each trait and integrating over the posterior distribution of each parameter. The variances from all
470 random effects were used in the computation of the total phenotypic variance. Note also that the values
471 computed for V_P and thus h^2 account for the variance explained by fixed effects [55]. Where relevant
472 for the trait (i.e. not for non-Gaussian traits nor for laying date, for which the population mean is too
473 arbitrary for CV_A to be meaningful), the coefficients of variation of the additive genetic variance (CV_A)
474 were computed as the square-root of the additive genetic variance divided by the population mean,
475 multiplied by 100. For fitness (lifetime reproductive success), we also estimated the evolvability I_A (V_A
476 divided by the squared population mean) as a standardised measure of adaptive potential [56, 57].

477 **Estimation of selection** Selection gradients were computed for each trait, centred to a mean of 0 and
478 scaled to a variance of 1. We used a Poisson generalised linear model (as implemented in the `glm` R
479 function of the `stats` package) to account for the skewed and discrete nature of fitness and computed
480 the gradients and their standard-errors based on [13].

481 **Simulation analysis** In order to assess the ability of our sampling design (i.e., sample size, along with
482 the pedigree data structure) to estimate *small, but substantial* heritabilities, we performed a simulation
483 analysis. We used the pedigree of the Tiritiri Matangi hihi population to simulate traits according to
484 five different scenarios: (i) the trait is individual-based (all fledglings are measured once) and normally
485 distributed, typical of e.g. tarsus length; (ii) the trait is breeding-based (only breeding females are
486 measured, but with 3 measures *per* female) and normally distributed, typical of e.g. laying date; (iii)
487 the trait is individual-based (all individuals are measured once) and binary (typical of e.g. recruitment
488 status); (iv) the trait is breeding-based and binary (i.e. composed of zeros and ones, with 3 measures
489 per female), typical of e.g. hatching success and (v) a zero-inflated Poisson trait with latent mean
490 and variance comparable to our fitness trait. Because heritability below an arbitrary threshold of 0.1
491 would be regarded as small (close to 0.1) to inconsequential (close to 0), we decided to use 0.1 as
492 the simulated heritability. For the binary trait, parameters were set so that the heritability on the
493 observed data scale (rather than the latent scale or liability) was 0.1. Since scenario (v) was about
494 detecting additive genetic variance of fitness and the heritability is not the best measure for this trait
495 [57], we used an evolvability (or additive genetic variance of relative fitness) of 0.01 for the Poisson
496 process, which corresponds to a heritability of 0.04. Since the meaning of evolvability for binomial
497 traits is less obvious, we kept a heritability of 0.1 for this (part of the) trait in scenario (v), which
498 would correspond to an evolvability of 0.03. As a result of using the pedigree from our study, the
499 sample sizes of each simulated scenario closely followed ours (see simulation code hosted in Dryad).
500 Additionally, in order to follow the structure of our fitted models and account for the precision lost by
501 adding random effects, we simulated the effects of social sire, dam and year of birth for scenario (i),
502 (iii) & (v) and of individual ID (permanent environment), mate ID and year of breeding for scenarios
503 (ii) & (iv). All of the simulated effects were added as random effects in the fitted models. The variance
504 of those random effects were set equal to 1, with a residual variance of 1.5 for the Gaussian trait and a
505 “residual” variance of 1 (as it is fixed for `MCMCg1mm`) for binary traits. Each scenario consisted of 100
506 replicates, fitted in `MCMCg1mm`, for which we computed the posterior mode, median and 95% credible
507 interval of the estimated heritability.

508 **Data and Software availability**

509 Phenotypic data, the pedigree and R code is available online on the UCL Discovery repository (DOI:
510 10.14324/000.ds.10065966). Hihi are of cultural significance to the indigenous people of New Zealand,
511 the Mori, and are considered a taonga (treasured) species. For this reason, the genotypes, raw reads,
512 assembled genome and RAD-seq reads for hihi will be made available by request on the recommendation
513 of the iwi (tribe) that affiliates as kaitiaki (guardians) for hihi. To obtain up-to-date contact details for
514 the iwi, please contact Dr Anna Santure: a.santure@auckland.ac.nz