

Homocysteine and Cognition: A Systematic Review of 111 studies.

AUTHORS:

Esther Setién-Suero ^{a,b,c*+}

Marta Suárez-Pinilla ^{d+}

Paula Suárez-Pinilla ^{a, b,c}

Benedicto Crespo-Facorro ^{a, b, c}

Rosa Ayesa-Arriola ^{a, b, c}

Affiliation/Location of Work:

^a University Hospital Marqués de Valdecilla. Department of Psychiatry, School of Medicine, University of Cantabria, Santander, Spain

^b CIBERSAM, Biomedical Research Network on Mental Health Area, Madrid, Spain.

^c IDIVAL, Valdecilla Biomedical Research Institute, Santander, Spain.

^d Sackler Centre for Consciousness Science. Sussex University. United Kingdom.

Manuscript: 5,821 words; Figures: 7; Supplementary figures: 4; Tables: 4; References: 165

+ These authors have equally contributed to this work

*Corresponding Author:

Esther Setién Suero

Unidad de Investigación Psiquiatría

Hospital Universitario Marqués de Valdecilla

Avda. Valdecilla, SN. 39008

Santander. Spain. Tel: +34-942-203826

E-mail: setiensuero@hotmail.com

HIGHLIGHTS

There is a positive trend between cognitive decline and increased Hcy plasma concentrations.

Vitamin supplementation fails to reduce the cognitive impairments once they appear.

Earlier detection of elevated Hcy levels may prevent cognitive impairment and dementia.

ABSTRACT

Background: Elevated plasma homocysteine (Hcy) levels have been associated with cognitive dysfunction in a wide range of conditions. The aim of this review is to establish which cognitive domains and populations are the most affected.

Methods: We systematically review the literature and consider all articles that showed any relationship between the plasma Hcy levels and scores achieved on cognitive performance tests in both, the general population and patients suffering from central nervous system disorders and other diseases.

Results: We found 111 pertinent articles. There were 24 cohort studies, 18 randomized trials, 21 transverse case-control studies, and 48 cross-sectional studies. This review reveals a positive trend between cognitive decline and increased plasma Hcy concentrations in general population and in patients with cognitive impairments. Executive functions and processing speed are the most often affected domains.

Discussion: Further investigations are warranted to clarify this relationship. Treatment with vitamin supplementation often fails to reduce the cognitive impairments. Earlier detection of the elevated Hcy levels may be an effective intervention to prevent cognitive impairment and dementia.

Keywords: Plasma Hcy, Hyperhomocysteinemia, Cognition, Cognitive Impairment, Cognitive Domains.

1. INTRODUCTION

Homocysteine (Hcy) is a non-protein, neurotoxic, sulfur-containing amino acid that originates from methionine metabolism. Alternatively, methionine is an amino acid precursor of peptides and proteins that plays an important role in the metabolism and transfer of methyl groups (1, 2). There is a close relationship between the plasma Hcy and folate, B6 and B12 vitamin levels, which act as coenzymes of the methionine and Hcy metabolism. Deficiencies in folate, and B6 and B12 are potential factors that contribute to increase the plasma concentrations of Hcy (3).

The reference values for plasma Hcy range from 5 to 15 $\mu\text{mol/L}$ (4). Hyperhomocysteinemia (HHcy) occurs at concentrations over 15 $\mu\text{mol/L}$ in fasting serum. Moreover, HHcy has been classified as moderate (plasma total Hcy concentrations of 15-30 $\mu\text{mol/L}$), intermediate (plasma total Hcy concentrations of 31-100 $\mu\text{mol/L}$), or severe (plasma total Hcy concentrations 100 $\mu\text{mol/L}$) (5).

The total concentration of Hcy in plasma is the result of a complex interaction between multiple genetic and environmental factors. Physiological determinants such as age, gender, ethnicity, menopause and pregnancy should be considered. Thus, the Hcy levels show a progressive increase with age. Moreover, higher total Hcy levels have been found in males compared to females; and postmenopausal women have shown elevated Hcy levels compared to the pre-menopausal women. Additionally, the plasma Hcy levels are generally lower in the black and Asian population (6). Some genetic factors that predispose an individual to HHcy are mutations in methionine synthase, cystathionine b synthase (CBS) and variants of the methylene tetrahydrofolate reductase (MTHFR) gene (7). Lifestyle determinants, such as smoking, alcohol and coffee consumption, and a lack of physical exercise or poor nutrition, may also contribute to the Hcy plasma levels (8, 9). Finally, it is worth knowing that some determinants related to sample collection and storage may also alter the Hcy concentrations (10).

There is growing evidence that higher Hcy levels are involved in age-related cognitive deficits and various types of central nervous system (CNS) disorders (11), including Alzheimer's disease (AD) (12), Parkinson disease (PD) (13), multiple sclerosis (MS) (14), cerebrovascular diseases and strokes (15). The relationship between HHcy and age-related degeneration has also been highlighted (16). Furthermore, elevated Hcy levels may influence the cognitive impairment that is commonly observed among patients with psychiatric disorders (17), such as affective disorders (18) or schizophrenia (SCZ) (19). In most of these cases, the associations remain strong, even after adjusting for vitamin status or renal function.

The CNS disorders related to HHcy may be explained by three possible mechanisms (20, 21). First, by promoting oxidative stress, Hcy may exhibit direct neurotoxicity, alter neurotransmission and induce neuronal excitotoxicity, apoptosis, β - amyloid accumulation and hyperactivation of the NMDA receptors (22, 23). Second, it has also been noted that a metabolic disruption in Hcy and the one-carbon metabolism may alter neurotransmitter synthesis. Finally, the association of homocysteinemia with occlusive vascular disease may be mediated through damage to the blood vessel wall or impaired blood coagulation.

By combining these results with the hypothesis that HHcy is a modifiable risk factor for cognitive dysfunctions, the aim of the present review is to identify which cognitive domains, if any, and populations are related to the increased Hcy levels in plasma.

2. MATERIALS AND METHODS

2.1. Comprehensive search of the literature

A systematic review of the literature was conducted through an electronic search of scientific journals in English and Spanish from the Medline database, via PubMed and Embase. The following key terms were used for the search: "homocysteine" and "cognition". We found 474 articles, and 373 articles were published in the last ten years, until the end of December 2015. In addition, the references from these articles that met the established inclusion criteria were also included.

2.2. Eligibility and study selection

The full text of each potentially eligible article was read by two researchers (one medical doctor and one psychologist) before a final decision was reached about its inclusion in the present review. For inclusion, we considered all of the original articles that addressed the influence of the plasma Hcy levels on cognitive functions in healthy subjects and patients suffering for any disease or condition. The studies could report their results quantitatively or qualitatively. The studies that were not related to Hcy levels and cognitive functions were excluded, even when they were found by the search terms. Non-original articles, such as reviews, meta-analyses, case series, letters to the editor, comments and case reports, were also excluded. We also rejected those studies where Hcy levels were measured in different fluids, such as cerebrospinal fluid. Finally, animal research articles were not included, as our purpose was to identify a relationship between HHcy and the cognitive domains in humans. Disagreements were resolved by consensus.

2.3. Data extraction

Data extraction was classified according to the design of the selected studies. The articles were divided into four blocks based on their design: case-control studies, cohort studies, randomized clinical trials (RCTs) and cross-sectional case studies. A database was created and the common variables of each article were extracted: sample size (N) of the reference population in the study, Hcy levels in plasma, and the neurocognitive tests used to measure the cognitive functions and their scores, when possible, among others. The time of follow-up was also specified for the cohort study block. For the randomized controlled trials, the treatment received in each branch was also noted.

2.4. Data analysis

In regard to the qualitative approach, because of the heterogeneity within each block and to provide a better analysis, the studies were compared and grouped into categories regarding each clinical condition: Alzheimer's and cognitive impairment, Parkinson's disease, schizophrenia, bipolar disorder, geriatric depression, eating disorders, cardiovascular diseases, cerebrovascular diseases, multiple sclerosis, other diseases, and the general population.

A quantitative approach was performed when estimates of association with the standard errors were reported or could be calculated through confidence intervals or p values. All the analyses were performed using Stata 12 (StataCorp, College Station, Tex., USA). The calculated effect sizes (ES) were: standardized mean differences, for group comparisons (i.e. RCT, and Hcy concentration difference between cases and controls), and linear regression coefficient, to show the influence of Hcy variations in

cognitive scores (i.e. cohort studies, cross sectional case studies and case control studies).

Fixed and random effects were used to combine effect sizes, but only random effects are presented in the manuscript. These results were pooled using the inverse variance method based on the DerSimonian and Laird random-effects model [a]. Overall and population subgroup meta-analyses were performed when the possible.

Heterogeneity was assessed using Cochran's Q and I^2 statistics [b, c, d]. An influence analysis was performed across individual studies by successively subtracting each study in order to know the effect size variation, considering the largest studies or studies which could contribute to the heterogeneity because of their special conditions or conducted in specific populations. Publication bias was assessed using the funnel plot method by a visual inspection [e].

3. RESULTS

Of the 373 articles that were selected based on the search criteria, 127 items were chosen as potentially relevant after reading the abstracts. From them, seven studies were excluded because they did not identify a direct relationship between the blood Hcy levels and cognitive functions. From these 120 articles, 11 were excluded because they overlapped with other included studies in the place and time of performance (16, 24-33). Finally, two additional articles were added from the references as they met the established selection criteria (Figure 1).

A total of 111 articles were included in the systematic review. Tables I to IV show the distribution of the articles according to their design: 24 cohort studies, 18 randomized double-blind trials, 21 transverse case-control studies and 48 cross-sectional case studies (Tables I-IV).

3.1. Alzheimer's and cognitive impairment

Together, the eight case-control studies showed significant associations between the plasma Hcy levels and cognitive functions in population with AD and/or cognitive impairment. In four studies, this association was determined using the Mini Mental State Examination (MMSE) test. Overall, an inverse correlation was found between the MMSE scores and the plasma Hcy levels (34-37). Similarly, Kim and colleagues showed a significant negative relationship between the plasma Hcy levels and the scores of the Word List Memory and Constructional Recall (WLMCR) test in the AD group, after adjusting for several covariates. However, this association did not remain significant in either the mild cognitive impairment (MCI) group or in the control group (38). Additionally, Faux and colleagues found a negative trend between the global cognitive scores and plasma Hcy levels; a negative correlation between the plasma Hcy levels and different types of memory was also found (39). Irizarry and colleagues demonstrated a significant relationship between the plasma Hcy levels and cognitive function

only in the group of patients with PD without dementia, while no correlations were shown in the dementia group (40). Finally, in the population over 75, Ford and colleagues observed that a two-fold increase in the plasma Hcy concentrations resulted in a 36% increase in cognitive impairment (41).

Eight case studies with cross sectional analyses of cases included people suffering from AD or cognitive impairment. Four of them, showed an inverse association between the plasma Hcy levels and the MMSE scores (42-45). However, this association did not remain significant when one of the studies adjusted the results by age and gender (42). Likewise, Nilson and colleagues observed that the predictive value of the Hcy levels on cognitive function was reduced (5%) when age was introduced as covariate in the linear regression model (43). Additionally, Adunsky and colleagues found that the cognitive state of patients in the tertile with the highest plasma Hcys levels was significantly impaired compared to the other groups (44). Bonetti and colleagues showed that hyperhomocysteinemia was associated with a significantly greater risk of dementia (45). Similarly, two other studies stated that high plasma Hcy levels are a risk factor for MCI (46, 47). Nevertheless, Manders and colleagues have found contradictory results in institutionalized patients, and did not find any relationship between the plasma Hcy levels and cognitive function (48). Finally, one study found several significant negative associations between the plasma Hcy levels and scores measuring the domain of visuoconstructive abilities (49).

With regard to the four selected RCTs, three found that although vitamin supplementation may reduce the plasma Hcy levels, there was no effect regarding the cognitive tests (50-52). However, De Jager and colleagues observed that executive functions benefit from vitamin supplementation compared with the placebo. There were significant differences in semantic memory and global cognitive function (MMSE) in patients with plasma Hcy levels over 13.1 $\mu\text{mol/liter}$ who received supplements compared to the placebo group (53).

Five cohort studies were included in the AD and/or cognitive impairment group, and the results were disparate. Thus, Tu and colleagues found an association between the reduced Cognitive Abilities Screening Instrument (CASI) and abstract thinking scores and the Hcy levels during six months of follow-up (54), although there were no significant correlations between the cognitive scores and the Hcy levels at the baseline and at the end of the studies. Similarly, Rozzini and colleagues showed an association between HHcy and the progression of memory impairment (55). Moreover, Oulhaj and colleagues found a concentration dependent relationship between the baseline Hcy levels and the rate of decline in the patients' Cambridge Cognition Examination (CAMCOG) scores, where the higher the Hcy levels, the faster the decline. This relationship was significant in patients aged over 75 years who had not previously suffered a stroke (12). However, in Siuda et al., 2009, there was no relationship between the Hcy levels and the evolution in cognitive functions during one-year follow-up, even though the baseline HHcy were higher in the MCI group compared to the healthy individuals (56). Reitz et al., 2009, did not find any associations between the Hcy levels and the prevalence of MCI in either the transverse or longitudinal analyses after adjusting for ethnicity and the ApoE- ϵ 4 genotype (57).

3.2. Parkinson's disease

Regarding the five case-control studies, three did not observe significant relationships between the plasma Hcy levels and cognitive functions (58-60). In contrast, Zoccolella and colleagues showed a significant association between the Hcy levels and the prevalence of dementia in PD (61). In addition, in the study by Ozer and colleagues, the group of PD patients with Hcy levels $> 14 \mu\text{mol/liter}$ had a worse performance in frontal and memory tests (62).

Two case studies with cross sectional analyses of cases were included. In Annanmaki et al. 2008, there was no association between the cognitive function scores and the plasma Hcy

levels (63). Conversely, the study of Martin et al. 2010 showed higher Hcy levels in PD patients with cognitive impairment and levodopa treatment (13).

Two RCTs explored the relationship between the Hcy levels and cognition in PD. Thus, Barone and colleagues showed that elevated Hcy levels were associated with greater differences in rivastigmine-treated patients than normal/low Hcy levels. The group of rivastigmine-treated PD patients presented marked or moderate improvement in their cognitive outcomes after 24 weeks of treatment compared to the placebo group, whereas no between-group differences were found in patients with low or normal Hcy levels (64). Similarly, Litvinenko and colleagues found that patients who were treated with memantine had higher scores on the MMSE, Alzheimer's Disease Assessment Scale-cognitive subscale (ADAS-cog), clock drawing test and Frontal Assessment Battery (FAB) compared to the control group by the end of study at week 24 (65).

3.3. Schizophrenia

Three studies explored the relationship between the plasma Hcy levels and cognitive function in patients with SCZ. The case-control study showed that patients had a statistically significant increase in their plasma Hcy levels. In addition, patients with SCZ exhibited a significantly increase in cognitive deficits. However, no relationship between the Hcy levels and cognitive impairment was detected (66). Moreover, the cross sectional case study did not find associations between the Hcy levels and cognitive test performance in patients with SCZ (67). A randomized clinical trial observed that the Hcy levels were decreased in patients with Hcy > 15 $\mu\text{mol/liter}$ and vitamin therapy compared to the placebo group. The overall neuropsychological test results, and the Wisconsin Card Sort test (WCST) results in particular, were significantly improved after vitamin treatment compared to the placebo (19).

3.4. Bipolar disorder

Three articles contemplated the relationship between the Hcy levels and cognition in euthymic patients with BD. A trend between higher Hcy levels and worse cognitive functioning was noticed. A cross-sectional case study in euthymic patients with BD demonstrated that male bipolar subjects showed higher average Hcy levels than a comparison group of normal subjects, and the authors also found that poorer functioning on a task of executive function (WCST) was related to higher Hcy levels (68). There were two articles with a case-control design based on euthymic patients with BD and healthy controls. In Dittman et al., 2008, the Hcy levels were significantly higher for patients than the controls. Stepwise regression analyses revealed a significant and independent association of the Hcy levels with verbal learning, delayed memory and executive function in the patient group (18). On the other hand, Dias and colleagues showed that the Hcy levels for the euthymic patients and healthy controls were similar. Patients with higher Hcy levels performed significantly worse on all neurocognitive tests, and there was a significant association between the Hcy levels and the number of perseverations on the Stroop Color Tests (SCT) and the number of moves on Hanoi Tower Test (ToH); however, a linear regression model revealed that Hcy was not a significant predictor of neurocognitive test performance (69).

3.5. Geriatric depression

We included two case-control articles that studied cognition and Hcy levels in major depression, with contradictory results. Alexopoulos and colleagues found that depressed patients with high levels of Hcy performed better in language processing and processing speed than patients with Hcy concentrations $\leq 11.7 \mu\text{mol/liter}$, whereas the results were opposite in the healthy control group (70). However, Ford and colleagues found that elevated plasma Hcy levels were associated with weaker performance in tests of immediate and delayed memory and global cognitive performance compared to those with normal Hcy levels, which was independent of the presence of major depression (71).

3.6. Eating disorders

Two cross-sectional studies indicated that higher Hcy levels were related with increased cognitive impairment in patients with anorexia and bulimia nervosa. Thus, Frieling and colleagues showed a significant positive association between long term and short term memory and the Hcy levels (72). Similarly, Welheim et al. 2010, found a significant association between low Hcy levels and cognitive deficits (73). In contrast, a RCT using folic acid versus placebo showed that the supplemented group exhibited significantly increased serum and red blood cell folate levels, reduced Hcy levels and significantly improved most of the cognitive test scores, whereas the placebo group showed no significant changes in any of the evaluated variables (74).

3.7. Cardiovascular diseases

The opposite results were found in the cross sectional case studies of population with cardiovascular diseases. Two of the studies found no significant association between cognition and the Hcy levels after adjusting for demographic and medical factors (75, 76). Nevertheless, one study found a significant association between HHcy and worse cognitive performance after adjusting for age, gender, premorbid intellectual coefficient and white matter lesions (77).

Regarding the three cohort studies of patients with cardiovascular diseases, two of the studies observed a significant relationship between the Hcy levels and cognition during follow-up. Narayan and colleagues showed a positive association between higher Hcy levels and cognitive impairment in three domains (executive function, episodic memory and processing speed). The associations remained significant during a 44-month follow-up. The association with executive function did not remain significant when folate was included as a covariate in the regression model. (78). Similarly, Jochemsen et al. 2013, observed a significant association between higher Hcy concentrations and worse executive function in patients over 65 during a

longitudinal study of 3.9 years. The study highlighted that elevated Hcy levels were related to the progression of ventricular enlargement and increased the risk for a decrease in executive functioning in older persons (79). However, Silbert and colleagues showed no association between the Hcy levels and cognition at any time during a one-year follow-up, but their Hcy levels were strongly associated with age and left ventricular function (80).

3.8. Cerebrovascular diseases

In the group of cerebrovascular diseases, the results were again contradictory. Of the two articles with cross sectional analyses of cases, one study found no differences between HHcy and cognitive performance (81). Conversely, another study showed a significant inverse relationship between the plasma Hcy levels and MMSE scores (82). One study using a transverse case-control design found that the Hcy levels were strongly correlated with cognitive decline in patients with small vessel disease (83). The controversies persisted in two cohort studies of populations with cerebrovascular diseases. One study found no relationship between the plasma Hcy levels and cognitive results after 27 months of follow-up (84). On the other hand, Newman and colleagues considered Hcy to be an independent predictor of less successful cognitive recovery in patients after stroke (85).

3.9. Kidney diseases

No significant relationship between the Hcy levels and cognitive impairment was detected in patients suffering from kidney disease (86-88). In a RCT with B vitamin supplementation versus placebo, the treatment decreased the Hcy levels by 26.7%. However, the treatment did not improve the initial cognitive outcomes or affect the subsequent cognitive status one year later (88).

3.10. Multiple sclerosis

A case-control study showed that patients with MS had higher plasma levels of Hcy and, consequently, HHcy was associated with cognitive impairment (14). However, a longitudinal study showed that the MS patients and healthy controls had similar Hcy levels. Patients with higher Hcy levels had lower cognitive scores, as measured with the Paced Auditory Serial Addition Test (PASAT). During a four-year follow-up, higher Hcy levels were associated with decreased cognitive scores in the PASAT (89).

3.11. Other diseases

In adolescents with epilepsy, there was no relationship between the plasma Hcy levels and cognitive scores, even though the Hcy levels were significantly higher in patients with epilepsy compared to healthy subjects (90). Similarly, there was no association between the plasma Hcy levels and the MMSE or SKT scores in a sample of geriatric patients with multiple co-morbid conditions (91). There was also no correlation between the Hcy levels and cognitive results in patients with obstructive apnea sleep syndrome (92). However, in a sample of patients with multiple system atrophy, a significantly negative correlation appeared between Hcy levels and punctuation in the MMSE (93).

3.12. General population

Twenty-four articles based on the general population used a cross sectional design. From these, 17 identified significant associations between the plasma Hcy concentrations and cognitive scores. A negative association between the MMSE scores and plasma Hcy levels was found in five studies (94-98). Furthermore, two more studies found a significant relationship under special conditions, including people over 60 years (99) and in subjects who were Apo E- ϵ 1 carriers (100). Specifically, negative associations between cognitive performance and Hcy levels were found in psychomotor speed, coordination, manual dexterity, verbal memory and learning (101), processing speed and spatial vision (102), visual and verbal memory (103), delayed memory and attention (104), and executive functions and language (105). One study

showed that the Brief Cognitive Assessment Tool (BCAT) scores were significantly impaired in people with homocysteinemia over 16 $\mu\text{mol/liter}$, specifically in the domains of perceptual speed, mental arithmetic efficacy, spatial vision and working memory (106). There were two articles conducted on post-menopausal women, which showed a significant association between higher plasma Hcy levels and worse cognition performance. In the first study, the association was significant when the working memory and verbal memory scores were grouped in a global score (107). In the other one, the global score was based in scores of five cognitive domains (108). In a sample of right-handed people, processing speed was also associated with the Hcy levels after adjusting for age, gender and educational level. The signification disappeared when the white matter volume was included in the regression analyses as a covariate (109). Finally, the relationship between cholesterol levels and cognitive function depends upon Hcy levels; in participants with normal Hcy levels, an inverse U-shaped relationship between the total cholesterol level and cognitive score was found, indicating that both low and high cholesterol levels were associated with lower cognitive scores. In participants with high Hcy levels, no significant association was found between cholesterol and cognition (110). In seven cross-sectional studies, there were no association between the plasma Hcy levels and cognitive performance (111-117). In one study, although the Hcy concentration was not associated with any particular domain, it was associated with global cognition. However this association did not remain significant after adjusting for the white matter volume and brain infarcts, although the significance persisted when adjusted for the total brain volume (116). Another study found no association between cognition and the Hcy levels in subjects aged 65 years and older when results were adjusted for the folate levels (117).

From the nine randomized clinical trials in the general population, eight showed that although vitamin supplementation reduced the Hcy levels, this did not impact the cognitive test scores (118-125). However, improved cognitive performance was observed in one article where the

vitamin supplementation reduced the Hcy levels. This trial was conducted with middle- and older-aged people with HHcy (126).

Finally, 13 cohort studies were conducted with the general population. From these, there were significant relationships between the Hcy levels and cognitive functions during the transverse analyses in three studies. However, during the follow-up, the Hcy concentrations were not predictors of cognitive impairment (127-129). Nevertheless, Ganguli et al., 2014, found no associations between the Hcy levels and cognitive functions during transverse analyses but there was a significant association between the Hcy levels and the executive function domain during four years of follow-up (130). In addition, five studies demonstrated a significant association between the plasma Hcy levels and cognitive function during some point of the longitudinal design (131-135). Hence, Van den Kommer and colleagues showed that higher Hcy levels at baseline were negatively associated with prolonged lower cognitive function and a faster rate of decline in information processing speed and fluid intelligence (135). Additionally, Hooshmand and colleagues found that higher plasma Hcy levels at the onset of the study were associated with poorer performance in global cognition, episodic memory, executive functions and verbal expression at a seven-year follow-up (134). These results may be comparable with the study of Strand et al. 2013, which was performed in 15-month-old children. In this study, each 2-fold increase in the Hcy concentration was associated with a two-point decrease in the mental development index score (133). Interestingly, Sharma et al., 2014, found that the plasma Hcy levels increased proportionately with the duration of stay at high altitude and hypobaric hypoxia. Moreover, the Hcy levels were inversely correlated with the MMSE scores during across the entire follow-up period (132). Moreover, the study of de Whalley et al., 2014, supported an association between HHcy and late onset dementia during a five-year follow-up. The association between Hcy and dementia was independent of the plasma folate, B12 and antioxidant micronutrient concentrations (131). Finally, in 2014, Samaras and colleagues, found no association between the plasma Hcy levels

and global cognitive function over a two-year study, but the Hcy levels were predictors of an inverse association with executive function (136). On the contrary, Tucker and colleagues showed no association between the plasma Hcy levels and cognitive scores over four years in a male sample, after adjusting for the folate and B6 and B12 vitamin concentrations (137). Furthermore, two more studies observed no association between increased Hcy levels and changes in cognitive performance (138, 139); one of them was conducted with university students, and the time of follow-up was six hours (138).

3.13 Quantitative approach

3.13.1. Case control studies

Fifteen studies comparing Hcy levels in patients versus healthy population reflected that Hcy concentration is significantly higher in the group of patients. This fact is particularly interesting because there is included a wide range of diverse diseases (AD, PD, BD, small vessel disease and MS). However, it is worth noting that there were some diseases that contribute to the ES only with one study. In regard to the PD group, levodopa administration is a confusion factor as the treatment always increases the Hcy levels (Figure 2).

On the other hand, three studies could be included for the case analyses of patients with AD and PD. We found that higher level of Hcy was associated with worse MMSE results in both groups (ES=-0.34 [-0.63, -0.05]) (Figure 3).

3.13.2. Cross sectional case studies

Thirteen cross sectional case studies were included in the quantitative analyses. In this group, we also have found a negative association between plasma Hcy levels and MMSE scores (ES=-0.19 [-0.27, -0.10]) (Figure 4). The negative associations remained significant when subgroup analyses were performed, therefore, there were not significant differences across diverse types of population (Figure 5).

No significant associations were found when Hcy levels were compared with scores of BNT (ES=0.02 [-0.02, 0.05]), TMT (ES=0.02 (-0.05, 0.10)), and RAVLT scores (ES=0.02 [-0.02, 0.05]). It is worth noting that only a small number of studies could be included for each case (Supplementary figures 1-3).

3.13.3. RCT studies

Five RCT articles were included in the quantitative approach. Overall, there was no association between Hcy level and MMSE score (ES= 0.04 [-0.06, 0.14]) (Figure 6). There was neither association between Hcy and RAVLT scores in the analysis that could be performed with three studies (ES=-0.04 -0.08, 0.15)](Supplementary figure 4)

3.13.4. Cohort studies

There was no significant association in six cohort articles that were included for a quantitative exploration of the association between MMSE scores (ES=-0.04 [-0.14, 0.15]) (Figure 7). Subgroup results were no relevant because of the small number of cohort articles for each population that could be included in the analyses.

4. DISCUSSION

Overall, our findings suggest a possible role for the plasma Hcy levels in the development or the progression of cognitive dysfunctions. More than half of the included articles (N=59) show some form of significant association between the plasma Hcy levels and cognitive functions, even when the analyses were adjusted for the relevant covariates. However, it remains unclear whether the high Hcy concentrations are responsible for cognitive impairments, and its etiopathogenic mechanism has not been conclusively identified.

The included articles used several different instruments to measure cognitive impairment. However, the MMSE test is one of the most commonly used tests. The MMSE is frequently used as screening test for global cognition (140). The results of this systematic review include multiple studies with significant associations between the plasma Hcy concentrations and MMSE scores. Overall, the associations between the Hcy levels and cognition were negative. Although this instrument provides information on global cognitive impairment, the MMSE test does not allow us to identify the domains that are the most altered. In many other articles, more complete cognitive batteries demonstrate that the processing speed and the executive functions were the domains that were most affected by the increased plasma Hcy concentrations. Frequently, different types of memory, such as visual memory, short-term memory and explicit long term-memory, are also altered in patients with high Hcy levels. The relationship between the plasma Hcy levels and the scores representing such domains is inverse, similar to the MMSE test. Therefore, higher blood Hcy levels are associated with a worse performance in several cognitive tests.

Interestingly, there are three studies showing a positive association between cognition and the Hcy concentrations (70, 72, 73). Two of them were performed in patients with eating disorders, and found a positive association between the elevated Hcy levels and memory test scores (72), with a better performance in SCT and SKT tests (73). These results note a beneficial

effect of high Hcy levels on memory, selective attention and processing speed in eating disorder patients. This fact may indicate that the link between the elevated Hcy levels and cognition originating from a nutritional deficiency may be different than the relationship between the high Hcy levels and cognition in the elderly. Another study with similar findings was conducted with aging patients with depression (70). In this case, the elevated Hcy concentrations were associated with a better performance in the processing speed and language processing tests. As is highlighted, there are certain results indicating that major depressive disorder may be related to a perturbation in glutamate metabolism (141, 142). Glutamate metabolism is decreased in certain areas of the cortex and Hcy may actually replacing glutamate at excitatory synapses, and, attenuate the cognitive deficits caused by glutamate deficiency (143-147). This theory could be similar in eating disorder patients, where there is also a glutamate reduction (148).

Most of the cohort studies showed an association between the Hcy concentrations and the evolution of the cognitive decline, particularly in the studies concerning the general population, AD and cognitive impairment, cardiovascular disease, cerebrovascular disease and the general conditions associated with aging. This fact may be partially explained by suggesting cumulative effect of Hcy in the elderly, with endothelial injury and atherosclerosis (149). Although some of the studies show no significant association during the follow-up, the association is significant during some of the transverse analyses.

Moreover, in the transverse group, the domains that are most affected by the elevated Hcy levels are executive functions, different types of memory, processing speed and global cognition. It is notorious that Hcy levels are generally higher in the group of patients compared with the healthy population. This fact is particularly interesting because there is included a wide range of diverse diseases (AD, PD, BD, small vessel disease and MS). However, it is worth noting that there were some diseases that contribute to the ES only with one study. In regard

to the PD group, levodopa therapy is a confusion factor as the treatment always increases the Hcy levels (f)

Nevertheless, many of the RCTs find no evidence of B vitamin supplementation and cognitive improvement, although there was evidence that B vitamins may reduce the plasma Hcy levels to the standard values. From the 15 RCTs, the treatment with B vitamin reduced the Hcy levels in all cases, but only six studies indicated an improvement in the patients' scores on different cognitive tests (19, 53, 64, 65, 74, 126). There are two possible theories for these observations. On one hand, the effects of Hcy in the brain may be irreversible and therefore, the effects of vitamin supplementation may be limited once the pathological changes appeared (150). On the other hand, the cognitive impairment may not be caused by the elevated Hcy levels but could be produced by common mechanisms of both conditions (151, 152). Both theories could explain why the vitamin supplementation could reduce the Hcy levels, but could not reduce or prevent the cognitive impairments. It is debatable whether the earlier detection of HHcy and related factors may prevent the neurodegeneration associated with aging and dementia (153). However, as a primary preventative measure, screening for Hcy levels may protect patients from other devastating effects, such as atherosclerosis (154), hypertension (155), thrombus (156), aging-related oxidation (157) and osteoporosis (158), which are commonly associated with higher Hcy concentrations.

This review is subject to some limitations that are worth noting. The reasons why certain cognitive domains are more affected by the high Hcy concentrations in some studies but not in others are not clear. These discrepancies may be due to the differences in the cognitive batteries between studies and those inherent to the administration, as well as to the methodology employed during the experiments (159). Moreover, as noted above, the Hcy concentrations are a complex function resulting from the interaction between genetic and environmental factors (10). Furthermore, the heterogeneity of cognitive testing used in the

studies limited the comparison and pooling of data. Therefore, this review could not elucidate the relative effect of elevated homocysteine in some cognitive domains, and instead was necessarily limited our meta-analyses to only the most commonly measured outcomes. The quantitative approach was also limited by the fact that not all the articles explained their results using an association measure and a discrete number of study could be included in the meta-analysis. The meta-analytic results only represented a subsample of the literature summarized in the present review. Finally, it is worth noting that the small sample size ($N \leq 26$) of the three articles with a positive relationship between cognition and the Hcy levels should be considered when interpreting the results presented here. Among the strengths of this review is the fact that, to our knowledge, it is the largest systematic effort to summarize the data on the associations between Hcy and cognition in different populations.

In conclusion, further investigations are warranted to clarify the relationship between the plasma Hcy levels and cognitive function. Treatment with vitamin supplementation often fails to reduce the cognitive impairment once it appears. Earlier detection of the elevated Hcy levels before the decline in cognitive function may be an effective intervention to prevent cognitive impairment and dementia.

REFERENCES

1. Menéndez Cabezas A, Fernández-Britto Rodríguez JE. Metabolismo de la homocisteína y su relación con la aterosclerosis. *Rev Cubana Invest Biomed.* 1999;18(3):155-68.
2. Rudiger H, Jaenicke L. The biosynthesis of methionine. *Mol Cell Biochem.* 1973 Jun 27;1(2):157-68.
3. Fowler B. Homocysteine: overview of biochemistry, molecular biology, and role in disease processes. *Semin Vasc Med.* 2005 May;5(2):77-86.
4. Cotton F, Wautrecht JC, Lechevin V, Macours P, Thiry P, Gervy C, et al. Reference intervals for plasma homocysteine by the AxSYM immunoassay after collection in fluoride tubes. *Clin Chem.* 2003 Feb;49(2):315-7.
5. Christopher R, Nagaraja D, Shankar SK. Homocysteine and cerebral stroke in developing countries. *Curr Med Chem.* 2007;14(22):2393-401.
6. Refsum H, Smith AD, Ueland PM, Nexø E, Clarke R, McPartlin J, et al. Facts and recommendations about total homocysteine determinations: an expert opinion. *Clin Chem.* 2004 Jan;50(1):3-32.
7. Eghlim FF, Ashavaid TF, Nair KG. Genetic determinants of hyperhomocysteinemia in atherosclerosis. *Indian J Clin Biochem.* 2006 Sep;21(2):4-11.
8. Jacobsen DW. Homocysteine and vitamins in cardiovascular disease. *Clin Chem.* 1998 Aug;44(8 Pt 2):1833-43.
9. Finkelstein JD. Methionine metabolism in mammals. *J Nutr Biochem.* 1990 May;1(5):228-37.
10. De Luis DA, Fernández N, Aller R. Homocysteine, metabolism and dietary factors. *Endocrinol Nutr.* 2004;51(8):458-63.
11. Troen A, Rosenberg I. Homocysteine and cognitive function. *Semin Vasc Med.* 2005 May;5(2):209-14.
12. Oulhaj A, Refsum H, Beaumont H, Williams J, King E, Jacoby R, et al. Homocysteine as a predictor of cognitive decline in Alzheimer's disease. *Int J Geriatr Psychiatry.* 2010 Jan;25(1):82-90.
13. Martin-Fernandez JJ, Carles-Dies R, Canizares F, Parra S, Aviles F, Villegas I, et al. [Homocysteine and cognitive impairment in Parkinson's disease]. *Rev Neurol.* 2010 Feb 1-15;50(3):145-51.
14. Russo C, Morabito F, Luise F, Piromalli A, Battaglia L, Vinci A, et al. Hyperhomocysteinemia is associated with cognitive impairment in multiple sclerosis. *J Neurol.* 2008 Jan;255(1):64-9.
15. Bostom AG, Rosenberg IH, Silbershatz H, Jacques PF, Selhub J, D'Agostino RB, et al. Nonfasting plasma total homocysteine levels and stroke incidence in elderly persons: the Framingham Study. *Ann Intern Med.* 1999 Sep 7;131(5):352-5.
16. Kado DM, Karlamangla AS, Huang MH, Troen A, Rowe JW, Selhub J, et al. Homocysteine versus the vitamins folate, B6, and B12 as predictors of cognitive function and decline in older high-functioning adults: MacArthur Studies of Successful Aging. *Am J Med.* 2005 Feb;118(2):161-7.
17. Moustafa AA, Hewedi DH, Eissa AM, Frydecka D, Misiak B. Homocysteine levels in schizophrenia and affective disorders-focus on cognition. *Front Behav Neurosci.* 2014;8:343.
18. Dittmann S, Seemüller F, Grunze HC, Schwarz MJ, Zach J, Fast K, et al. The impact of homocysteine levels on cognition in euthymic bipolar patients: a cross-sectional study. *J Clin Psychiatry.* 2008 Jun;69(6):899-906.
19. Levine J, Stahl Z, Sela BA, Ruderman V, Shumaico O, Babushkin I, et al. Homocysteine-reducing strategies improve symptoms in chronic schizophrenic patients with hyperhomocysteinemia. *Biol Psychiatry.* 2006 Aug 1;60(3):265-9.

20. Selhub J, Bagley LC, Miller J, Rosenberg IH. B vitamins, homocysteine, and neurocognitive function in the elderly. *Am J Clin Nutr.* 2000 Feb;71(2):614S-20S.
21. Mattson MP, Shea TB. Folate and homocysteine metabolism in neural plasticity and neurodegenerative disorders. *Trends Neurosci.* 2003 Mar;26(3):137-46.
22. Lipton SA, Rosenberg PA. Excitatory amino acids as a final common pathway for neurologic disorders. *N Engl J Med.* 1994 Mar 3;330(9):613-22.
23. Sánchez-Cuevas M, Jiménez-Reséndiz SP, Morgado-Vázquez JS. La Homocisteína: un aminoácido neurotóxico. *REB.* 2009;28(1):3-8.
24. van Raamt AF, Kalmijn S, Mali WP, van Zandvoort MJ, van der Graaf Y. Homocysteine level and cognitive function in patients with arterial disease: the second manifestations of ARTERial Disease Study. *J Am Geriatr Soc.* 2006 Apr;54(4):575-9.
25. Elias MF, Robbins MA, Budge MM, Elias PK, Brennan SL, Johnston C, et al. Homocysteine, folate, and vitamins B6 and B12 blood levels in relation to cognitive performance: the Maine-Syracuse study. *Psychosom Med.* 2006 Jul-Aug;68(4):547-54.
26. Dittmann S, Seemuller F, Schwarz MJ, Kleindienst N, Stampfer R, Zach J, et al. Association of cognitive deficits with elevated homocysteine levels in euthymic bipolar patients and its impact on psychosocial functioning: preliminary results. *Bipolar Disord.* 2007 Feb-Mar;9(1-2):63-70.
27. Chee MW, Chen KH, Zheng H, Chan KP, Isaac V, Sim SK, et al. Cognitive function and brain structure correlations in healthy elderly East Asians. *Neuroimage.* 2009 May 15;46(1):257-69.
28. Cheng DM, Jiang YG, Huang CY, Kong HY, Pang W, Yang HP. Polymorphism of MTHFR C677T, serum vitamin levels and cognition in subjects with hyperhomocysteinemia in China. *Nutr Neurosci.* 2010 Aug;13(4):175-82.
29. Nilsson K, Gustafson L, Hultberg B. Plasma homocysteine, brain imaging and cognition in older patients with mental illness. *Int J Geriatr Psychiatry.* 2012 Apr;27(4):409-14.
30. Slawek J, Roszmann A, Robowski P, Dubaniewicz M, Sitek EJ, Honczarenko K, et al. The impact of MRI white matter hyperintensities on dementia in Parkinson's disease in relation to the homocysteine level and other vascular risk factors. *Neurodegener Dis.* 2013;12(1):1-12.
31. Lipnicki DM, Sachdev PS, Crawford J, Reppermund S, Kochan NA, Trollor JN, et al. Risk factors for late-life cognitive decline and variation with age and sex in the Sydney Memory and Ageing Study. *PLoS ONE.* 2013;8(6):e65841.
32. Huang CW, Chang WN, Huang SH, Lui CC, Chen NC, Chang YT, et al. Impact of homocysteine on cortical perfusion and cognitive decline in mild Alzheimer's dementia. *Eur J Neurol.* 2013 Aug;20(8):1191-7.
33. Wu MK, Lu YT, Huang CW, Lin PH, Chen NC, Lui CC, et al. Clinical Significance of Cerebrovascular Biomarkers and White Matter Tract Integrity in Alzheimer Disease: Clinical correlations With Neurobehavioral Data in Cross-Sectional and After 18 Months Follow-ups. *Medicine (Baltimore).* 2015 Jul;94(28):e1192.
34. Hernanz A, De la Fuente M, Navarro M, Frank A. Plasma aminothiols compounds, but not serum tumor necrosis factor receptor II and soluble receptor for advanced glycation end products, are related to the cognitive impairment in Alzheimer's disease and mild cognitive impairment patients. *Neuroimmunomodulation.* 2007;14(3-4):163-7.
35. Gorgone G, Ursini F, Altamura C, Bressi F, Tombini M, Curcio G, et al. Hyperhomocysteinemia, intima-media thickness and C677T MTHFR gene polymorphism: a correlation study in patients with cognitive impairment. *Atherosclerosis.* 2009 Sep;206(1):309-13.
36. Leparo O, Alajbegovic A, Zaciragic A, Nakas-Icindic E, Valjevac A, Leparo D, et al. Elevated serum homocysteine level is not associated with serum C-reactive protein in patients with probable Alzheimer's disease. *J Neural Transm.* 2009 Dec;116(12):1651-6.

37. Lildballe DL, Fedosov S, Sherliker P, Hin H, Clarke R, Nexø E. Association of cognitive impairment with combinations of vitamin B(1)(2)-related parameters. *Clin Chem*. 2011 Oct;57(10):1436-43.
38. Kim G, Kim H, Kim KN, Son JI, Kim SY, Tamura T, et al. Relationship of cognitive function with B vitamin status, homocysteine, and tissue factor pathway inhibitor in cognitively impaired elderly: a cross-sectional survey. *J Alzheimers Dis*. 2013;33(3):853-62.
39. Faux NG, Ellis KA, Porter L, Fowler CJ, Laws SM, Martins RN, et al. Homocysteine, vitamin B12, and folic acid levels in Alzheimer's disease, mild cognitive impairment, and healthy elderly: baseline characteristics in subjects of the Australian Imaging Biomarker Lifestyle study. *J Alzheimers Dis*. 2011;27(4):909-22.
40. Irizarry MC, Gurol ME, Raju S, Diaz-Arrastia R, Locascio JJ, Tennis M, et al. Association of homocysteine with plasma amyloid beta protein in aging and neurodegenerative disease. *Neurology*. 2005 Nov 8;65(9):1402-8.
41. Ford AH, Flicker L, Hankey GJ, Norman P, van Bockxmeer FM, Almeida OP. Homocysteine, methylenetetrahydrofolate reductase C677T polymorphism and cognitive impairment: the health in men study. *Mol Psychiatry*. 2012 May;17(5):559-66.
42. Li L, Cao D, Desmond R, Rahman A, Lah JJ, Levey AI, et al. Cognitive performance and plasma levels of homocysteine, vitamin B12, folate and lipids in patients with Alzheimer disease. *Dement Geriatr Cogn Disord*. 2008;26(4):384-90.
43. Nilsson K, Gustafson L, Hultberg B. Plasma homocysteine and cognition in elderly patients with dementia or other psychogeriatric diseases. *Dement Geriatr Cogn Disord*. 2010;30(3):198-204.
44. Adunsky A, Arinzon Z, Fidelman Z, Krasniansky I, Arad M, Gepstein R. Plasma homocysteine levels and cognitive status in long-term stay geriatric patients: a cross-sectional study. *Arch Gerontol Geriatr*. 2005 Mar-Apr;40(2):129-38.
45. Bonetti F, Brombo G, Magon S, Zuliani G. Cognitive Status According to Homocysteine and B-Group Vitamins in Elderly Adults. *J Am Geriatr Soc*. 2015 Jun;63(6):1158-63.
46. Kim J, Park MH, Kim E, Han C, Jo SA, Jo I. Plasma homocysteine is associated with the risk of mild cognitive impairment in an elderly Korean population. *J Nutr*. 2007 Sep;137(9):2093-7.
47. Sachdev PS, Lipnicki DM, Crawford J, Reppermund S, Kochan NA, Trollor JN, et al. Risk profiles of subtypes of mild cognitive impairment: the sydney memory and ageing study. *J Am Geriatr Soc*. 2012 Jan;60(1):24-33.
48. Manders M, Vasse E, de Groot LC, van Staveren WA, Bindels JG, Blom HJ, et al. Homocysteine and cognitive function in institutionalised elderly A cross-sectional analysis. *Eur J Nutr*. 2006 Mar;45(2):70-8.
49. Sala I, Belen Sanchez-Saudinos M, Molina-Porcel L, Lazaro E, Gich I, Clarimon J, et al. Homocysteine and cognitive impairment. Relation with diagnosis and neuropsychological performance. *Dement Geriatr Cogn Disord*. 2008;26(6):506-12.
50. Aisen PS, Schneider LS, Sano M, Diaz-Arrastia R, van Dyck CH, Weiner MF, et al. High-dose B vitamin supplementation and cognitive decline in Alzheimer disease: a randomized controlled trial. *JAMA*. 2008 Oct 15;300(15):1774-83.
51. Kwok T, Lee J, Law CB, Pan PC, Yung CY, Choi KC, et al. A randomized placebo controlled trial of homocysteine lowering to reduce cognitive decline in older demented people. *Clin Nutr*. 2011 Jun;30(3):297-302.
52. Kobe T, Witte AV, Schnelle A, Lesemann A, Fabian S, Tesky VA, et al. Combined omega-3 fatty acids, aerobic exercise and cognitive stimulation prevents decline in gray matter volume of the frontal, parietal and cingulate cortex in patients with mild cognitive impairment. *Neuroimage*. 2015 Oct 1.
53. de Jager CA, Oulhaj A, Jacoby R, Refsum H, Smith AD. Cognitive and clinical outcomes of homocysteine-lowering B-vitamin treatment in mild cognitive impairment: a randomized controlled trial. *Int J Geriatr Psychiatry*. 2012 Jun;27(6):592-600.

54. Tu MC, Huang CW, Chen NC, Chang WN, Lui CC, Chen CF, et al. Hyperhomocysteinemia in Alzheimer dementia patients and cognitive decline after 6 months follow-up period. *Acta Neurol Taiwan*. 2010 Sep;19(3):168-77.
55. Rozzini L, Chilovi BV, Bertolotti E, Conti M, Del Rio I, Trabucchi M, et al. Angiotensin converting enzyme (ACE) inhibitors modulate the rate of progression of amnesic mild cognitive impairment. *Int J Geriatr Psychiatry*. 2006 Jun;21(6):550-5.
56. Siuda J, Gorzkowska A, Patalong-Ogiewa M, Krzystanek E, Czech E, Wiechula B, et al. From mild cognitive impairment to Alzheimer's disease - influence of homocysteine, vitamin B12 and folate on cognition over time: results from one-year follow-up. *Neurol Neurochir Pol*. 2009 Jul-Aug;43(4):321-9.
57. Reitz C, Tang MX, Miller J, Green R, Luchsinger JA. Plasma homocysteine and risk of mild cognitive impairment. *Dement Geriatr Cogn Disord*. 2009;27(1):11-7.
58. Bialecka M, Kurzawski M, Roszmann A, Robowski P, Sitek EJ, Honczarenko K, et al. Association of COMT, MTHFR, and SLC19A1(RFC-1) polymorphisms with homocysteine blood levels and cognitive impairment in Parkinson's disease. *Pharmacogenet Genomics*. 2012 Oct;22(10):716-24.
59. Rodriguez-Oroz MC, Lage PM, Sanchez-Mut J, Lamet I, Pagonabarraga J, Toledo JB, et al. Homocysteine and cognitive impairment in Parkinson's disease: a biochemical, neuroimaging, and genetic study. *Mov Disord*. 2009 Jul 30;24(10):1437-44.
60. Camicioli RM, Bouchard TP, Somerville MJ. Homocysteine is not associated with global motor or cognitive measures in nondemented older Parkinson's disease patients. *Mov Disord*. 2009 Jan 30;24(2):176-82.
61. Zoccolella S, dell'Aquila C, Abruzzese G, Antonini A, Bonuccelli U, Canesi M, et al. Hyperhomocysteinemia in levodopa-treated patients with Parkinson's disease dementia. *Mov Disord*. 2009 May 15;24(7):1028-33.
62. Ozer F, Meral H, Hanoglu L, Aydemir T, Yilsen M, Cetin S, et al. Plasma homocysteine levels in patients treated with levodopa: motor and cognitive associations. *Neurol Res*. 2006 Dec;28(8):853-8.
63. Annanmaki T, Pessala-Driver A, Hokkanen L, Murros K. Uric acid associates with cognition in Parkinson's disease. *Parkinsonism Relat Disord*. 2008 Nov;14(7):576-8.
64. Barone P, Burn DJ, van Laar T, Hsu C, Poewe W, Lane RM. Rivastigmine versus placebo in hyperhomocysteinemic Parkinson's disease dementia patients. *Mov Disord*. 2008 Aug 15;23(11):1532-40.
65. Litvinenko IV, Odinak MM, Mogil'naya VI, Perstnev SV. Use of memantine (akatinol) for the correction of cognitive impairments in Parkinson's disease complicated by dementia. *Neurosci Behav Physiol*. 2010 Feb;40(2):149-55.
66. Ayesa-Arriola R, Perez-Iglesias R, Rodriguez-Sanchez JM, Mata I, Gomez-Ruiz E, Garcia-Unzueta M, et al. Homocysteine and cognition in first-episode psychosis patients. *Eur Arch Psychiatry Clin Neurosci*. 2012 Oct;262(7):557-64.
67. Gonzalez-Liencreces C, Tas C, Brown EC, Erdin S, Onur E, Cubukcoglu Z, et al. Oxidative stress in schizophrenia: a case-control study on the effects on social cognition and neurocognition. *BMC Psychiatry*. 2014;14:268.
68. Osher Y, Bersudsky Y, Silver H, Sela BA, Belmaker RH. Neuropsychological correlates of homocysteine levels in euthymic bipolar patients. *J Affect Disord*. 2008 Jan;105(1-3):229-33.
69. Dias VV, Brissos S, Cardoso C, Andreatza AC, Kapczinski F. Serum homocysteine levels and cognitive functioning in euthymic bipolar patients. *J Affect Disord*. 2009 Mar;113(3):285-90.
70. Alexopoulos P, Topalidis S, Irmisch G, Prehn K, Jung SU, Poppe K, et al. Homocysteine and cognitive function in geriatric depression. *Neuropsychobiology*. 2010;61(2):97-104.
71. Ford AH, Flicker L, Singh U, Hirani V, Almeida OP. Homocysteine, depression and cognitive function in older adults. *J Affect Disord*. 2013 Nov;151(2):646-51.

72. Frieling H, Roschke B, Kornhuber J, Wilhelm J, Romer KD, Gruss B, et al. Cognitive impairment and its association with homocysteine plasma levels in females with eating disorders - findings from the HEaD-study. *J Neural Transm*. 2005 Nov;112(11):1591-8.
73. Wilhelm J, Muller E, de Zwaan M, Fischer J, Hillemecher T, Kornhuber J, et al. Elevation of homocysteine levels is onl 1. Menéndez Cabezas A, Fernández-Britto Rodríguez JE. *Metabolismo de la homocisteina y su relación con la aterosclerosis*. *Rev Cubana Invest Biomed*. 1999;18(3):155-68.
2. Rudiger H, Jaenicke L. The biosynthesis of methionine. *Mol Cell Biochem*. 1973 Jun 27;1(2):157-68.
3. Fowler B. Homocysteine: overview of biochemistry, molecular biology, and role in disease processes. *Semin Vasc Med*. 2005 May;5(2):77-86.
4. Cotton F, Wautrecht JC, Lechevin V, Macours P, Thiry P, Gervy C, et al. Reference intervals for plasma homocysteine by the AxSYM immunoassay after collection in fluoride tubes. *Clin Chem*. 2003 Feb;49(2):315-7.
5. Christopher R, Nagaraja D, Shankar SK. Homocysteine and cerebral stroke in developing countries. *Curr Med Chem*. 2007;14(22):2393-401.
6. Refsum H, Smith AD, Ueland PM, Nexø E, Clarke R, McPartlin J, et al. Facts and recommendations about total homocysteine determinations: an expert opinion. *Clin Chem*. 2004 Jan;50(1):3-32.
7. Eghlim FF, Ashavaid TF, Nair KG. Genetic determinants of hyperhomocysteinemia in atherosclerosis. *Indian J Clin Biochem*. 2006 Sep;21(2):4-11.
8. Jacobsen DW. Homocysteine and vitamins in cardiovascular disease. *Clin Chem*. 1998 Aug;44(8 Pt 2):1833-43.
9. Finkelstein JD. Methionine metabolism in mammals. *J Nutr Biochem*. 1990 May;1(5):228-37.
10. De Luis DA, Fernández N, Aller R. Homocysteine, metabolism and dietary factors. *Endocrinol Nutr*. 2004;51(8):458-63.
11. Troen A, Rosenberg I. Homocysteine and cognitive function. *Semin Vasc Med*. 2005 May;5(2):209-14.
12. Oulhaj A, Refsum H, Beaumont H, Williams J, King E, Jacoby R, et al. Homocysteine as a predictor of cognitive decline in Alzheimer's disease. *Int J Geriatr Psychiatry*. 2010 Jan;25(1):82-90.
13. Martín-Fernández JJ, Carles-Dies R, Canizares F, Parra S, Aviles F, Villegas I, et al. [Homocysteine and cognitive impairment in Parkinson's disease]. *Rev Neurol*. 2010 Feb 15;50(3):145-51.
14. Russo C, Morabito F, Luise F, Piromalli A, Battaglia L, Vinci A, et al. Hyperhomocysteinemia is associated with cognitive impairment in multiple sclerosis. *J Neurol*. 2008 Jan;255(1):64-9.
15. Bostom AG, Rosenberg IH, Silbershatz H, Jacques PF, Selhub J, D'Agostino RB, et al. Nonfasting plasma total homocysteine levels and stroke incidence in elderly persons: the Framingham Study. *Ann Intern Med*. 1999 Sep 7;131(5):352-5.
16. Kado DM, Karlamangla AS, Huang MH, Troen A, Rowe JW, Selhub J, et al. Homocysteine versus the vitamins folate, B6, and B12 as predictors of cognitive function and decline in older high-functioning adults: MacArthur Studies of Successful Aging. *Am J Med*. 2005 Feb;118(2):161-7.
17. Moustafa AA, Hewedi DH, Eissa AM, Frydecka D, Misiak B. Homocysteine levels in schizophrenia and affective disorders-focus on cognition. *Front Behav Neurosci*. 2014;8:343.
18. Dittmann S, Seemüller F, Grunze HC, Schwarz MJ, Zach J, Fast K, et al. The impact of homocysteine levels on cognition in euthymic bipolar patients: a cross-sectional study. *J Clin Psychiatry*. 2008 Jun;69(6):899-906.

19. Levine J, Stahl Z, Sela BA, Ruderman V, Shumaico O, Babushkin I, et al. Homocysteine-reducing strategies improve symptoms in chronic schizophrenic patients with hyperhomocysteinemia. *Biol Psychiatry*. 2006 Aug 1;60(3):265-9.
20. Selhub J, Bagley LC, Miller J, Rosenberg IH. B vitamins, homocysteine, and neurocognitive function in the elderly. *Am J Clin Nutr*. 2000 Feb;71(2):614S-20S.
21. Mattson MP, Shea TB. Folate and homocysteine metabolism in neural plasticity and neurodegenerative disorders. *Trends Neurosci*. 2003 Mar;26(3):137-46.
22. Lipton SA, Rosenberg PA. Excitatory amino acids as a final common pathway for neurologic disorders. *N Engl J Med*. 1994 Mar 3;330(9):613-22.
23. Sánchez-Cuevas M, Jiménez-Reséndiz SP, Morgado-Vázquez JS. La Homocisteína: un aminoácido neurotóxico. *REB*. 2009;28(1):3-8.
24. van Raamt AF, Kalmijn S, Mali WP, van Zandvoort MJ, van der Graaf Y. Homocysteine level and cognitive function in patients with arterial disease: the second manifestations of ARTERial Disease Study. *J Am Geriatr Soc*. 2006 Apr;54(4):575-9.
25. Elias MF, Robbins MA, Budge MM, Elias PK, Brennan SL, Johnston C, et al. Homocysteine, folate, and vitamins B6 and B12 blood levels in relation to cognitive performance: the Maine-Syracuse study. *Psychosom Med*. 2006 Jul-Aug;68(4):547-54.
26. Dittmann S, Seemuller F, Schwarz MJ, Kleindienst N, Stampfer R, Zach J, et al. Association of cognitive deficits with elevated homocysteine levels in euthymic bipolar patients and its impact on psychosocial functioning: preliminary results. *Bipolar Disord*. 2007 Feb-Mar;9(1-2):63-70.
27. Chee MW, Chen KH, Zheng H, Chan KP, Isaac V, Sim SK, et al. Cognitive function and brain structure correlations in healthy elderly East Asians. *Neuroimage*. 2009 May 15;46(1):257-69.
28. Cheng DM, Jiang YG, Huang CY, Kong HY, Pang W, Yang HP. Polymorphism of MTHFR C677T, serum vitamin levels and cognition in subjects with hyperhomocysteinemia in China. *Nutr Neurosci*. 2010 Aug;13(4):175-82.
29. Nilsson K, Gustafson L, Hultberg B. Plasma homocysteine, brain imaging and cognition in older patients with mental illness. *Int J Geriatr Psychiatry*. 2012 Apr;27(4):409-14.
30. Slawek J, Roszmann A, Robowski P, Dubaniewicz M, Sitek EJ, Honczarenko K, et al. The impact of MRI white matter hyperintensities on dementia in Parkinson's disease in relation to the homocysteine level and other vascular risk factors. *Neurodegener Dis*. 2013;12(1):1-12.
31. Lipnicki DM, Sachdev PS, Crawford J, Reppermund S, Kochan NA, Trollor JN, et al. Risk factors for late-life cognitive decline and variation with age and sex in the Sydney Memory and Ageing Study. *PLoS ONE*. 2013;8(6):e65841.
32. Huang CW, Chang WN, Huang SH, Lui CC, Chen NC, Chang YT, et al. Impact of homocysteine on cortical perfusion and cognitive decline in mild Alzheimer's dementia. *Eur J Neurol*. 2013 Aug;20(8):1191-7.
33. Wu MK, Lu YT, Huang CW, Lin PH, Chen NC, Lui CC, et al. Clinical Significance of Cerebrovascular Biomarkers and White Matter Tract Integrity in Alzheimer Disease: Clinical correlations With Neurobehavioral Data in Cross-Sectional and After 18 Months Follow-ups. *Medicine (Baltimore)*. 2015 Jul;94(28):e1192.
34. Hernanz A, De la Fuente M, Navarro M, Frank A. Plasma aminothiols, but not serum tumor necrosis factor receptor II and soluble receptor for advanced glycation end products, are related to the cognitive impairment in Alzheimer's disease and mild cognitive impairment patients. *Neuroimmunomodulation*. 2007;14(3-4):163-7.
35. Gorgone G, Ursini F, Altamura C, Bressi F, Tombini M, Curcio G, et al. Hyperhomocysteinemia, intima-media thickness and C677T MTHFR gene polymorphism: a correlation study in patients with cognitive impairment. *Atherosclerosis*. 2009 Sep;206(1):309-13.

36. Lepara O, Alajbegovic A, Zaciragic A, Nakas-Icindic E, Valjevac A, Lepara D, et al. Elevated serum homocysteine level is not associated with serum C-reactive protein in patients with probable Alzheimer's disease. *J Neural Transm*. 2009 Dec;116(12):1651-6.
37. Lildballe DL, Fedosov S, Sherliker P, Hin H, Clarke R, Nexø E. Association of cognitive impairment with combinations of vitamin B(1)(2)-related parameters. *Clin Chem*. 2011 Oct;57(10):1436-43.
38. Kim G, Kim H, Kim KN, Son JI, Kim SY, Tamura T, et al. Relationship of cognitive function with B vitamin status, homocysteine, and tissue factor pathway inhibitor in cognitively impaired elderly: a cross-sectional survey. *J Alzheimers Dis*. 2013;33(3):853-62.
39. Faux NG, Ellis KA, Porter L, Fowler CJ, Laws SM, Martins RN, et al. Homocysteine, vitamin B12, and folic acid levels in Alzheimer's disease, mild cognitive impairment, and healthy elderly: baseline characteristics in subjects of the Australian Imaging Biomarker Lifestyle study. *J Alzheimers Dis*. 2011;27(4):909-22.
40. Irizarry MC, Gurol ME, Raju S, Diaz-Arrastia R, Locascio JJ, Tennis M, et al. Association of homocysteine with plasma amyloid beta protein in aging and neurodegenerative disease. *Neurology*. 2005 Nov 8;65(9):1402-8.
41. Ford AH, Flicker L, Hankey GJ, Norman P, van Bockxmeer FM, Almeida OP. Homocysteine, methylenetetrahydrofolate reductase C677T polymorphism and cognitive impairment: the health in men study. *Mol Psychiatry*. 2012 May;17(5):559-66.
42. Li L, Cao D, Desmond R, Rahman A, Lah JJ, Levey AI, et al. Cognitive performance and plasma levels of homocysteine, vitamin B12, folate and lipids in patients with Alzheimer disease. *Dement Geriatr Cogn Disord*. 2008;26(4):384-90.
43. Nilsson K, Gustafson L, Hultberg B. Plasma homocysteine and cognition in elderly patients with dementia or other psychogeriatric diseases. *Dement Geriatr Cogn Disord*. 2010;30(3):198-204.
44. Adunsky A, Arinzon Z, Fidelman Z, Krasniansky I, Arad M, Gepstein R. Plasma homocysteine levels and cognitive status in long-term stay geriatric patients: a cross-sectional study. *Arch Gerontol Geriatr*. 2005 Mar-Apr;40(2):129-38.
45. Bonetti F, Brombo G, Magon S, Zuliani G. Cognitive Status According to Homocysteine and B-Group Vitamins in Elderly Adults. *J Am Geriatr Soc*. 2015 Jun;63(6):1158-63.
46. Kim J, Park MH, Kim E, Han C, Jo SA, Jo I. Plasma homocysteine is associated with the risk of mild cognitive impairment in an elderly Korean population. *J Nutr*. 2007 Sep;137(9):2093-7.
47. Sachdev PS, Lipnicki DM, Crawford J, Reppermund S, Kochan NA, Trollor JN, et al. Risk profiles of subtypes of mild cognitive impairment: the sydney memory and ageing study. *J Am Geriatr Soc*. 2012 Jan;60(1):24-33.
48. Manders M, Vasse E, de Groot LC, van Staveren WA, Bindels JG, Blom HJ, et al. Homocysteine and cognitive function in institutionalised elderly A cross-sectional analysis. *Eur J Nutr*. 2006 Mar;45(2):70-8.
49. Sala I, Belen Sanchez-Saudinos M, Molina-Porcel L, Lazaro E, Gich I, Clarimon J, et al. Homocysteine and cognitive impairment. Relation with diagnosis and neuropsychological performance. *Dement Geriatr Cogn Disord*. 2008;26(6):506-12.
50. Aisen PS, Schneider LS, Sano M, Diaz-Arrastia R, van Dyck CH, Weiner MF, et al. High-dose B vitamin supplementation and cognitive decline in Alzheimer disease: a randomized controlled trial. *JAMA*. 2008 Oct 15;300(15):1774-83.
51. Kwok T, Lee J, Law CB, Pan PC, Yung CY, Choi KC, et al. A randomized placebo controlled trial of homocysteine lowering to reduce cognitive decline in older demented people. *Clin Nutr*. 2011 Jun;30(3):297-302.
52. Kobe T, Witte AV, Schnelle A, Lesemann A, Fabian S, Tesky VA, et al. Combined omega-3 fatty acids, aerobic exercise and cognitive stimulation prevents decline in gray matter volume of the frontal, parietal and cingulate cortex in patients with mild cognitive impairment. *Neuroimage*. 2015 Oct 1.

53. de Jager CA, Oulhaj A, Jacoby R, Refsum H, Smith AD. Cognitive and clinical outcomes of homocysteine-lowering B-vitamin treatment in mild cognitive impairment: a randomized controlled trial. *Int J Geriatr Psychiatry*. 2012 Jun;27(6):592-600.
54. Tu MC, Huang CW, Chen NC, Chang WN, Lui CC, Chen CF, et al. Hyperhomocysteinemia in Alzheimer dementia patients and cognitive decline after 6 months follow-up period. *Acta Neurol Taiwan*. 2010 Sep;19(3):168-77.
55. Rozzini L, Chilovi BV, Bertoletti E, Conti M, Del Rio I, Trabucchi M, et al. Angiotensin converting enzyme (ACE) inhibitors modulate the rate of progression of amnesic mild cognitive impairment. *Int J Geriatr Psychiatry*. 2006 Jun;21(6):550-5.
56. Siuda J, Gorzkowska A, Patalong-Ogiewa M, Krzystanek E, Czech E, Wiechula B, et al. From mild cognitive impairment to Alzheimer's disease - influence of homocysteine, vitamin B12 and folate on cognition over time: results from one-year follow-up. *Neurol Neurochir Pol*. 2009 Jul-Aug;43(4):321-9.
57. Reitz C, Tang MX, Miller J, Green R, Luchsinger JA. Plasma homocysteine and risk of mild cognitive impairment. *Dement Geriatr Cogn Disord*. 2009;27(1):11-7.
58. Bialecka M, Kurzawski M, Roszmann A, Robowski P, Sitek EJ, Honczarenko K, et al. Association of COMT, MTHFR, and SLC19A1(RFC-1) polymorphisms with homocysteine blood levels and cognitive impairment in Parkinson's disease. *Pharmacogenet Genomics*. 2012 Oct;22(10):716-24.
59. Rodriguez-Oroz MC, Lage PM, Sanchez-Mut J, Lamet I, Pagonabarraga J, Toledo JB, et al. Homocysteine and cognitive impairment in Parkinson's disease: a biochemical, neuroimaging, and genetic study. *Mov Disord*. 2009 Jul 30;24(10):1437-44.
60. Camicioli RM, Bouchard TP, Somerville MJ. Homocysteine is not associated with global motor or cognitive measures in nondemented older Parkinson's disease patients. *Mov Disord*. 2009 Jan 30;24(2):176-82.
61. Zoccolella S, dell'Aquila C, Abruzzese G, Antonini A, Bonuccelli U, Canesi M, et al. Hyperhomocysteinemia in levodopa-treated patients with Parkinson's disease dementia. *Mov Disord*. 2009 May 15;24(7):1028-33.
62. Ozer F, Meral H, Hanoglu L, Aydemir T, Yilsen M, Cetin S, et al. Plasma homocysteine levels in patients treated with levodopa: motor and cognitive associations. *Neurol Res*. 2006 Dec;28(8):853-8.
63. Annanmaki T, Pessala-Driver A, Hokkanen L, Murros K. Uric acid associates with cognition in Parkinson's disease. *Parkinsonism Relat Disord*. 2008 Nov;14(7):576-8.
64. Barone P, Burn DJ, van Laar T, Hsu C, Poewe W, Lane RM. Rivastigmine versus placebo in hyperhomocysteinemic Parkinson's disease dementia patients. *Mov Disord*. 2008 Aug 15;23(11):1532-40.
65. Litvinenko IV, Odinak MM, Mogil'naya VI, Perstnev SV. Use of memantine (akatinol) for the correction of cognitive impairments in Parkinson's disease complicated by dementia. *Neurosci Behav Physiol*. 2010 Feb;40(2):149-55.
66. Ayesa-Arriola R, Perez-Iglesias R, Rodriguez-Sanchez JM, Mata I, Gomez-Ruiz E, Garcia-Unzueta M, et al. Homocysteine and cognition in first-episode psychosis patients. *Eur Arch Psychiatry Clin Neurosci*. 2012 Oct;262(7):557-64.
67. Gonzalez-Liencre C, Tas C, Brown EC, Erdin S, Onur E, Cubukcoglu Z, et al. Oxidative stress in schizophrenia: a case-control study on the effects on social cognition and neurocognition. *BMC Psychiatry*. 2014;14:268.
68. Osher Y, Bersudsky Y, Silver H, Sela BA, Belmaker RH. Neuropsychological correlates of homocysteine levels in euthymic bipolar patients. *J Affect Disord*. 2008 Jan;105(1-3):229-33.
69. Dias VV, Brissos S, Cardoso C, Andrezza AC, Kapczinski F. Serum homocysteine levels and cognitive functioning in euthymic bipolar patients. *J Affect Disord*. 2009 Mar;113(3):285-90.
70. Alexopoulos P, Topalidis S, Irmisch G, Prehn K, Jung SU, Poppe K, et al. Homocysteine and cognitive function in geriatric depression. *Neuropsychobiology*. 2010;61(2):97-104.

71. Ford AH, Flicker L, Singh U, Hirani V, Almeida OP. Homocysteine, depression and cognitive function in older adults. *J Affect Disord.* 2013 Nov;151(2):646-51.
72. Frieling H, Roschke B, Kornhuber J, Wilhelm J, Romer KD, Gruss B, et al. Cognitive impairment and its association with homocysteine plasma levels in females with eating disorders - findings from the HEaD-study. *J Neural Transm.* 2005 Nov;112(11):1591-8.
73. Wilhelm J, Muller E, de Zwaan M, Fischer J, Hillemacher T, Kornhuber J, et al. Elevation of homocysteine levels is only partially reversed after therapy in females with eating disorders. *J Neural Transm.* 2010 Apr;117(4):521-7.
74. Loria-Kohen V, Gomez-Candela C, Palma-Milla S, Amador-Sastre B, Hernanz A, Bermejo LM. A pilot study of folic acid supplementation for improving homocysteine levels, cognitive and depressive status in eating disorders. *Nutr Hosp.* 2013 May-Jun;28(3):807-15.
75. Gunstad J, Bausserman L, Paul RH, Tate DF, Hoth K, Poppas A, et al. C-reactive protein, but not homocysteine, is related to cognitive dysfunction in older adults with cardiovascular disease. *J Clin Neurosci.* 2006 Jun;13(5):540-6.
76. Ford AH, Garrido GJ, Beer C, Lautenschlager NT, Arnolda L, Flicker L, et al. Homocysteine, grey matter and cognitive function in adults with cardiovascular disease. *PLoS ONE.* 2012;7(3):e33345.
77. Kloppenborg RP, Nederkoorn PJ, van der Graaf Y, Geerlings MI. Homocysteine and cerebral small vessel disease in patients with symptomatic atherosclerotic disease. The SMART-MR study. *Atherosclerosis.* 2011 Jun;216(2):461-6.
78. Narayan SK, Saxby BK, Firbank MJ, O'Brien JT, Harrington F, McKeith IG, et al. Plasma homocysteine and cognitive decline in older hypertensive subjects. *Int Psychogeriatr.* 2011 Dec;23(10):1607-15.
79. Jochemsen HM, Kloppenborg RP, de Groot LC, Kampman E, Mali WP, van der Graaf Y, et al. Homocysteine, progression of ventricular enlargement, and cognitive decline: the Second Manifestations of ARterial disease-Magnetic Resonance study. *Alzheimers Dement.* 2013 May;9(3):302-9.
80. Silbert B, Evered L, Scott DA, McCutcheon C, Jamrozik K. Homocysteine and C-reactive protein are not markers of cognitive impairment in patients with major cardiovascular disease. *Dement Geriatr Cogn Disord.* 2008;25(4):309-16.
81. Wong A, Mok V, Fan YH, Lam WW, Liang KS, Wong KS. Hyperhomocysteinemia is associated with volumetric white matter change in patients with small vessel disease. *J Neurol.* 2006 Apr;253(4):441-7.
82. Tay SY, Ampil ER, Chen CP, Auchus AP. The relationship between homocysteine, cognition and stroke subtypes in acute stroke. *J Neurol Sci.* 2006 Dec 1;250(1-2):58-61.
83. Pavlovic AM, Pekmezovic T, Obrenovic R, Novakovic I, Tomic G, Mijajlovic M, et al. Increased total homocysteine level is associated with clinical status and severity of white matter changes in symptomatic patients with subcortical small vessel disease. *Clin Neurol Neurosurg.* 2011 Nov;113(9):711-5.
84. Rowan EN, Dickinson HO, Stephens S, Ballard C, Kalaria R, Kenny RA. Homocysteine and post-stroke cognitive decline. *Age Ageing.* 2007 May;36(3):339-43.
85. Newman GC, Bang H, Hussain SI, Toole JF. Association of diabetes, homocysteine, and HDL with cognition and disability after stroke. *Neurology.* 2007 Nov 27;69(22):2054-62.
86. Kurella Tamura M, Xie D, Yaffe K, Cohen DL, Teal V, Kasner SE, et al. Vascular risk factors and cognitive impairment in chronic kidney disease: the Chronic Renal Insufficiency Cohort (CRIC) study. *Clin J Am Soc Nephrol.* 2011 Feb;6(2):248-56.
87. Troen AM, Scott TM, D'Anci KE, Moorthy D, Dobson B, Rogers G, et al. Cognitive dysfunction and depression in adult kidney transplant recipients: baseline findings from the FAVORIT Ancillary Cognitive Trial (FACT). *J Ren Nutr.* 2012 Mar;22(2):268-76 e1-3.
88. Brady CB, Gaziano JM, Cxypoliski RA, Guarino PD, Kaufman JS, Warren SR, et al. Homocysteine lowering and cognition in CKD: the Veterans Affairs homocysteine study. *Am J Kidney Dis.* 2009 Sep;54(3):440-9.

89. Teunissen CE, Killestein J, Kragt JJ, Polman CH, Dijkstra CD, Blom HJ. Serum homocysteine levels in relation to clinical progression in multiple sclerosis. *J Neurol Neurosurg Psychiatry*. 2008 Dec;79(12):1349-53.
90. Di Rosa G, Lenzo P, Parisi E, Neri M, Guerrera S, Nicotera A, et al. Role of plasma homocysteine levels and MTHFR polymorphisms on IQ scores in children and young adults with epilepsy treated with antiepileptic drugs. *Epilepsy Behav*. 2013 Dec;29(3):548-51.
91. Hengstermann S, Laemmler G, Hanemann A, Schweter A, Steinhagen-Thiessen E, Lun A, et al. Total serum homocysteine levels do not identify cognitive dysfunction in multimorbid elderly patients. *J Nutr Health Aging*. 2009 Feb;13(2):121-6.
92. Sales LV, Bruin VM, D'Almeida V, Pompeia S, Bueno OF, Tufik S, et al. Cognition and biomarkers of oxidative stress in obstructive sleep apnea. *Clinics (Sao Paulo)*. 2013 Apr;68(4):449-55.
93. Chen D, Wei X, Zou J, Wang R, Liu X, Xu X, et al. Contra-Directional Expression of Serum Homocysteine and Uric Acid as Important Biomarkers of Multiple System Atrophy Severity: A Cross-Sectional Study. *Front Cell Neurosci*. 2015;9:247.
94. Aleman A, Muller M, de Haan EH, van der Schouw YT. Vascular risk factors and cognitive function in a sample of independently living men. *Neurobiol Aging*. 2005 Apr;26(4):485-90.
95. Ravaglia G, Forti P, Maioli F, Brunetti N, Martelli M, Servadei L, et al. Serum C-reactive protein and cognitive function in healthy elderly Italian community dwellers. *J Gerontol A Biol Sci Med Sci*. 2005 Aug;60(8):1017-21.
96. Wright CB, Sacco RL, Rundek T, Delman J, Rabbani L, Elkind M. Interleukin-6 is associated with cognitive function: the Northern Manhattan Study. *J Stroke Cerebrovasc Dis*. 2006 Jan-Feb;15(1):34-8.
97. Koike T, Kuzuya M, Kanda S, Okada K, Izawa S, Enoki H, et al. Raised homocysteine and low folate and vitamin B-12 concentrations predict cognitive decline in community-dwelling older Japanese adults. *Clin Nutr*. 2008 Dec;27(6):865-71.
98. Allam M, Fahmy E, Elatti SA, Amer H, Abo-Krysha N, El-Sawy E. Association between total plasma homocysteine level and cognitive functions in elderly Egyptian subjects. *J Neurol Sci*. 2013 Sep 15;332(1-2):86-91.
99. Elias MF, Sullivan LM, D'Agostino RB, Elias PK, Jacques PF, Selhub J, et al. Homocysteine and cognitive performance in the Framingham offspring study: age is important. *Am J Epidemiol*. 2005 Oct 1;162(7):644-53.
100. Elias MF, Robbins MA, Budge MM, Elias PK, Dore GA, Brennan SL, et al. Homocysteine and cognitive performance: modification by the ApoE genotype. *Neurosci Lett*. 2008 Jan 3;430(1):64-9.
101. Schafer JH, Glass TA, Bolla KI, Mintz M, Jedlicka AE, Schwartz BS. Homocysteine and cognitive function in a population-based study of older adults. *J Am Geriatr Soc*. 2005 Mar;53(3):381-8.
102. Feng L, Ng TP, Chuah L, Niti M, Kua EH. Homocysteine, folate, and vitamin B-12 and cognitive performance in older Chinese adults: findings from the Singapore Longitudinal Ageing Study. *Am J Clin Nutr*. 2006 Dec;84(6):1506-12.
103. Chin AV, Robinson DJ, O'Connell H, Hamilton F, Bruce I, Coen R, et al. Vascular biomarkers of cognitive performance in a community-based elderly population: the Dublin Healthy Ageing study. *Age Ageing*. 2008 Sep;37(5):559-64.
104. Agrawal A, Ilango K, Singh PK, Karmakar D, Singh GP, Kumari R, et al. Age dependent levels of plasma homocysteine and cognitive performance. *Behav Brain Res*. 2015 Apr 15;283:139-44.
105. West RK, Beerli MS, Schmeidler J, Mitchell DB, Carlisle KR, Angelo G, et al. Homocysteine and cognitive function in very elderly nondemented subjects. *Am J Geriatr Psychiatry*. 2011 Jul;19(7):673-7.

106. Kong HY, Cheng DM, Pang W, Sun SD, Liu J, Huang CY, et al. Homocysteine levels and cognitive function scores measured with MMSE and BCAT of middle-aged and elderly subjects in Tianjin City. *J Nutr Health Aging*. 2013;17(6):527-32.
107. Clark MS, Guthrie JR, Dennerstein L. Hyperhomocysteinemia is associated with lower performance on memory tasks in post-menopausal women. *Dement Geriatr Cogn Disord*. 2005;20(2-3):57-62.
108. Raszewski G, Lorocho M, Owoc A, Lukawski K, Filip R, Bojar I. Homocysteine and cognitive disorders of postmenopausal women measured by a battery of computer tests--central nervous system vital signs. *Arch Womens Ment Health*. 2015 Aug;18(4):623-30.
109. Feng L, Isaac V, Sim S, Ng TP, Krishnan KR, Chee MW. Associations between elevated homocysteine, cognitive impairment, and reduced white matter volume in healthy old adults. *Am J Geriatr Psychiatry*. 2013 Feb;21(2):164-72.
110. Cheng Y, Jin Y, Unverzagt FW, Su L, Yang L, Ma F, et al. The relationship between cholesterol and cognitive function is homocysteine-dependent. *Clin Interv Aging*. 2014;9:1823-9.
111. Fischer P, Zehetmayer S, Bauer K, Huber K, Jungwirth S, Tragl KH. Relation between vascular risk factors and cognition at age 75. *Acta Neurol Scand*. 2006 Aug;114(2):84-90.
112. de Lau LM, Refsum H, Smith AD, Johnston C, Breteler MM. Plasma folate concentration and cognitive performance: Rotterdam Scan Study. *Am J Clin Nutr*. 2007 Sep;86(3):728-34.
113. Xiu LL, Lee MS, Wahlqvist ML, Chen RC, Huang YC, Chen KJ, et al. Low and high homocysteine are associated with mortality independent of B group vitamins but interactive with cognitive status in a free-living elderly cohort. *Nutr Res*. 2012 Dec;32(12):928-39.
114. Moorthy D, Peter I, Scott TM, Parnell LD, Lai CQ, Crott JW, et al. Status of vitamins B-12 and B-6 but not of folate, homocysteine, and the methylenetetrahydrofolate reductase C677T polymorphism are associated with impaired cognition and depression in adults. *J Nutr*. 2012 Aug;142(8):1554-60.
115. Strain JJ, McSorley EM, van Wijngaarden E, Kobrosly RW, Bonham MP, Mulhern MS, et al. Choline status and neurodevelopmental outcomes at 5 years of age in the Seychelles Child Development Nutrition Study. *Br J Nutr*. 2013 Jul 28;110(2):330-6.
116. Tangney CC, Aggarwal NT, Li H, Wilson RS, Decarli C, Evans DA, et al. Vitamin B12, cognition, and brain MRI measures: a cross-sectional examination. *Neurology*. 2011 Sep 27;77(13):1276-82.
117. Vidal JS, Dufouil C, Ducros V, Tzourio C. Homocysteine, folate and cognition in a large community-based sample of elderly people--the 3C Dijon Study. *Neuroepidemiology*. 2008;30(4):207-14.
118. Lewerin C, Matousek M, Steen G, Johansson B, Steen B, Nilsson-Ehle H. Significant correlations of plasma homocysteine and serum methylmalonic acid with movement and cognitive performance in elderly subjects but no improvement from short-term vitamin therapy: a placebo-controlled randomized study. *Am J Clin Nutr*. 2005 May;81(5):1155-62.
119. McMahon JA, Green TJ, Skeaff CM, Knight RG, Mann JI, Williams SM. A controlled trial of homocysteine lowering and cognitive performance. *N Engl J Med*. 2006 Jun 29;354(26):2764-72.
120. Eussen SJ, Ueland PM, Clarke R, Blom HJ, Hoefnagels WH, van Staveren WA, et al. The association of betaine, homocysteine and related metabolites with cognitive function in Dutch elderly people. *Br J Nutr*. 2007 Nov;98(5):960-8.
121. Macpherson H, Ellis KA, Sali A, Pipingas A. Memory improvements in elderly women following 16 weeks treatment with a combined multivitamin, mineral and herbal supplement: A randomized controlled trial. *Psychopharmacology (Berl)*. 2012 Mar;220(2):351-65.
122. Pipingas A, Camfield DA, Stough C, Scholey AB, Cox KH, White D, et al. Effects of multivitamin, mineral and herbal supplement on cognition in younger adults and the contribution of B group vitamins. *Hum Psychopharmacol*. 2014 Jan;29(1):73-82.

123. van der Zwaluw NL, Dhonukshe-Rutten RA, van Wijngaarden JP, Brouwer-Brolsma EM, van de Rest O, In 't Veld PH, et al. Results of 2-year vitamin B treatment on cognitive performance: secondary data from an RCT. *Neurology*. 2014 Dec 2;83(23):2158-66.
124. Dangour AD, Allen E, Clarke R, Elbourne D, Fletcher AE, Letley L, et al. Effects of vitamin B-12 supplementation on neurologic and cognitive function in older people: a randomized controlled trial. *Am J Clin Nutr*. 2015 Sep;102(3):639-47.
125. Harris E, Macpherson H, Pipingas A. Improved blood biomarkers but no cognitive effects from 16 weeks of multivitamin supplementation in healthy older adults. *Nutrients*. 2015 May;7(5):3796-812.
126. Cheng D, Kong H, Pang W, Yang H, Lu H, Huang C, et al. B vitamin supplementation improves cognitive function in the middle aged and elderly with hyperhomocysteinemia. *Nutr Neurosci*. 2014 Jun 18.
127. Mooijaart SP, Gussekloo J, Frolich M, Jolles J, Stott DJ, Westendorp RG, et al. Homocysteine, vitamin B-12, and folic acid and the risk of cognitive decline in old age: the Leiden 85-Plus study. *Am J Clin Nutr*. 2005 Oct;82(4):866-71.
128. Clarke R, Birks J, Nexo E, Ueland PM, Schneede J, Scott J, et al. Low vitamin B-12 status and risk of cognitive decline in older adults. *Am J Clin Nutr*. 2007 Nov;86(5):1384-91.
129. Brown B, Huang MH, Karlamangla A, Seeman T, Kado D. Do the effects of APOE-epsilon4 on cognitive function and decline depend upon vitamin status? *MacArthur Studies of Successful Aging. J Nutr Health Aging*. 2011 Mar;15(3):196-201.
130. Ganguli M, Fu B, Snitz BE, Unverzagt FW, Loewenstein DA, Hughes TF, et al. Vascular risk factors and cognitive decline in a population sample. *Alzheimer Dis Assoc Disord*. 2014 Jan-Mar;28(1):9-15.
131. Whalley LJ, Duthie SJ, Collins AR, Starr JM, Deary IJ, Lemmon H, et al. Homocysteine, antioxidant micronutrients and late onset dementia. *Eur J Nutr*. 2014 Feb;53(1):277-85.
132. Sharma VK, Das SK, Dhar P, Hota KB, Mahapatra BB, Vashishtha V, et al. Domain specific changes in cognition at high altitude and its correlation with hyperhomocysteinemia. *PLoS ONE*. 2014;9(7):e101448.
133. Strand TA, Taneja S, Ueland PM, Refsum H, Bahl R, Schneede J, et al. Cobalamin and folate status predicts mental development scores in North Indian children 12-18 mo of age. *Am J Clin Nutr*. 2013 Feb;97(2):310-7.
134. Hooshmand B, Solomon A, Kareholt I, Rusanen M, Hanninen T, Leiviska J, et al. Associations between serum homocysteine, holotranscobalamin, folate and cognition in the elderly: a longitudinal study. *J Intern Med*. 2012 Feb;271(2):204-12.
135. van den Kommer TN, Dik MG, Comijs HC, Jonker C, Deeg DJ. Homocysteine and inflammation: predictors of cognitive decline in older persons? *Neurobiol Aging*. 2010 Oct;31(10):1700-9.
136. Samaras K, Lutgers HL, Kochan NA, Crawford JD, Campbell LV, Wen W, et al. The impact of glucose disorders on cognition and brain volumes in the elderly: the Sydney Memory and Ageing Study. *Age (Dordr)*. 2014 Apr;36(2):977-93.
137. Tucker KL, Qiao N, Scott T, Rosenberg I, Spiro A, 3rd. High homocysteine and low B vitamins predict cognitive decline in aging men: the Veterans Affairs Normative Aging Study. *Am J Clin Nutr*. 2005 Sep;82(3):627-35.
138. Alexopoulos P, Lehl S, Richter-Schmidinger T, Kreusslein A, Hauenstein T, Bayerl F, et al. Short-term influence of elevation of plasma homocysteine levels on cognitive function in young healthy adults. *J Nutr Health Aging*. 2010 Apr;14(4):283-7.
139. Tangney CC, Tang Y, Evans DA, Morris MC. Biochemical indicators of vitamin B12 and folate insufficiency and cognitive decline. *Neurology*. 2009 Jan 27;72(4):361-7.
140. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975 Nov;12(3):189-98.

141. Sanacora G, Gueorguieva R, Epperson CN, Wu YT, Appel M, Rothman DL, et al. Subtype-specific alterations of gamma-aminobutyric acid and glutamate in patients with major depression. *Arch Gen Psychiatry*. 2004 Jul;61(7):705-13.
142. Pittenger C, Sanacora G, Krystal JH. The NMDA receptor as a therapeutic target in major depressive disorder. *CNS Neurol Disord Drug Targets*. 2007 Apr;6(2):101-15.
143. Hasler G, van der Veen JW, Tumonis T, Meyers N, Shen J, Drevets WC. Reduced prefrontal glutamate/glutamine and gamma-aminobutyric acid levels in major depression determined using proton magnetic resonance spectroscopy. *Arch Gen Psychiatry*. 2007 Feb;64(2):193-200.
144. Auer DP, Putz B, Kraft E, Lipinski B, Schill J, Holsboer F. Reduced glutamate in the anterior cingulate cortex in depression: an in vivo proton magnetic resonance spectroscopy study. *Biol Psychiatry*. 2000 Feb 15;47(4):305-13.
145. Pfliederer B, Michael N, Erfurth A, Ohrmann P, Hohmann U, Wolgast M, et al. Effective electroconvulsive therapy reverses glutamate/glutamine deficit in the left anterior cingulum of unipolar depressed patients. *Psychiatry Res*. 2003 Apr 1;122(3):185-92.
146. Michael N, Erfurth A, Ohrmann P, Arolt V, Heindel W, Pfliederer B. Metabolic changes within the left dorsolateral prefrontal cortex occurring with electroconvulsive therapy in patients with treatment resistant unipolar depression. *Psychol Med*. 2003 Oct;33(7):1277-84.
147. Michael N, Erfurth A, Ohrmann P, Arolt V, Heindel W, Pfliederer B. Neurotrophic effects of electroconvulsive therapy: a proton magnetic resonance study of the left amygdalar region in patients with treatment-resistant depression. *Neuropsychopharmacology*. 2003 Apr;28(4):720-5.
148. Ohrmann P, Kersting A, Suslow T, Lalee-Mentzel J, Donges US, Fiebich M, et al. Proton magnetic resonance spectroscopy in anorexia nervosa: correlations with cognition. *Neuroreport*. 2004 Mar 1;15(3):549-53.
149. Wu J, Uchino M, Sastry SM, Schaumberg DA. Age-related macular degeneration and the incidence of cardiovascular disease: a systematic review and meta-analysis. *PLoS One*. 2014;9(3):e89600.
150. Garcia A, Zanibbi K. Homocysteine and cognitive function in elderly people. *CMAJ*. 2004 Oct 12;171(8):897-904.
151. McCaddon A. Homocysteine and cognition--a historical perspective. *J Alzheimers Dis*. 2006 Aug;9(4):361-80.
152. McCaddon A. Homocysteine and cognitive impairment; a case series in a General Practice setting. *Nutr J*. 2006;5:6.
153. Wong CW. Vitamin B12 deficiency in the elderly: is it worth screening? *Hong Kong Med J*. 2015 Apr;21(2):155-64.
154. Falk E, Zhou J, Moller J. Homocysteine and atherothrombosis. *Lipids*. 2001;36 Suppl:S3-11.
155. Viridis A, Ghiadoni L, Salvetti G, Versari D, Taddei S, Salvetti A. Hyperhomocyst(e)inemia: is this a novel risk factor in hypertension? *J Nephrol*. 2002 Jul-Aug;15(4):414-21.
156. Harpel PC, Zhang X, Borth W. Homocysteine and hemostasis: pathogenic mechanisms predisposing to thrombosis. *J Nutr*. 1996 Apr;126(4 Suppl):1285S-9S.
157. Heil SG, De Vriese AS, Kluijtmans LA, Mortier S, Den Heijer M, Blom HJ. The role of hyperhomocysteinemia in nitric oxide (NO) and endothelium-derived hyperpolarizing factor (EDHF)-mediated vasodilatation. *Cell Mol Biol (Noisy-le-grand)*. 2004 Dec;50(8):911-6.
158. Herrmann M, Peter Schmidt J, Umanskaya N, Wagner A, Taban-Shomal O, Widmann T, et al. The role of hyperhomocysteinemia as well as folate, vitamin B(6) and B(12) deficiencies in osteoporosis: a systematic review. *Clin Chem Lab Med*. 2007;45(12):1621-32.
159. Ferne T, Rupp AA. A synthesis of 15 years of research on DIF in language testing: Methodological advances, challenges, and recommendations. *Language Assessment Quarterly*. 2007;4:113-48.

- a. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; 7: 177-188
- b. Cochran WG. The Combination of Estimates from Different Experiments. *Biometrics*. 1954; 10: 101-129
- c. Julian P. T. Higgins; and Simon G. Thompson. Quantifying heterogeneity in a meta-analysis. *Statist. Med.* 2002; 21:1539–1558
- d. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003 Sep 6;327(7414):557-60.
- e. Sterne JAC, Egger M, Moher D: Addressing reporting biases in: Higgins JPT, Green S (eds.): *Cochrane Handbook for Systematic reviews of interventions*, Chichester, Wiley, 2008. 297-334.
- f. Postuma RB, Lang AE. Homocysteine and levodopa: should Parkinson disease patients receive preventative therapy?. *Neurology*. 2004 Sep 14;63(5):886-91.




Figure 1. Flow diagram selection of study process




Figure 2. Case-control studies. Comparison of plasma Hcy levels between controls and patients. Abbreviations: CI= Confidence intervals; ES= Effect size; MS= Multiple sclerosis; N= sample size; SVD=Small vessel disease.




Figure 3. Case-control studies. Association between plasma Hcy levels and Mini Mental State Examination (MMSE) scores. Abbreviations: CI= confidence intervals; ES= Effect size; N= sample size




Figure 4. Cross-sectional studies. Association between plasma Hcy levels and Mini Mental State Examination (MMSE) scores in the whole sample. Abbreviations: CI= confidence intervals; CVD= cardiovascular disease; ES= Effect size; MSA= Multiple System Atrophy; N= sample size




Figure 5. Cross-sectional case studies. Association between plasma Hcy levels and Mini Mental State Examination (MMSE) scores sample divided by different diseases. Abbreviations: CI= confidence intervals; CVD= cardiovascular disease; ES= Effect size; MSA= Multiple System Atrophy; N= sample size




Figure 6. Randomize clinical trial (RCT) studies . Association between plasma Hcy levels and Mini Mental Examination (MMSE) scores. Abbreviations: CI= confidence intervals; ES= Effect size; N= sample size.




Figure 7. Cohort studies. Association between plasma Hcy levels and Mini Mental Examination (MMSE) scores. Abbreviations: CI= confidence intervals; CVD= cardiovascular disease; ES= Effect size; N= sample size

Table 1. Homocysteine and Cognition: Cohort studies

Population	Article	N	Age (years) mean (SD)	Follow-up	Hcy (μmol/l) mean (SD)	Neurocognitive tests
General	Tucker KL et al. 2005	321	67 (7)	3y	11 (5)	MMSE, Backward DS from WAIS-R, CERAD, Developmental test of VMI
	Moolijaart SP et al. 2005	559	85	4y	--	MMSE, Stroop test, LDCT, WLT
	Clarke R et al. 2007	2741	75.7 (7)	10y	14.5 (6.3)	MMSE
	Tangney CC et al. 2009	516	80 (6)	6y	11.5 (4.8)	MMSE, EBMT, SDMT
	Alexopoulos P et al. 2010	100	Group A: 23.6 (5.5) Group B: 21.6 (2.6)	6h	8.9 (3.2) 10.1 (3.5)	SKT, Stroop test, Short Test for General Intelligence
	Van den Kommer TM et al. 2010	1257	75.4 (6.6)	6y	13.6*	MMSE, AVLT, RCPM, Alphabet Coding Task-15
	Brown B et al. 2011	499	Group A: 74.3 (2.8) Group B: 74.3 (2.8)	7y	12.4 (5) 11.4 (4)	DRST, Similarities of WAIS-R
	Hooshmand B et al. 2012	274	70.1	7y	12.2*	MMSE, LDST, PPT, Stroop Test, Immediate Word Recall Test, Fluency test.
	Strand TA et al. 2013	650	Group A: 14.8 (2) m Group B: 15 (2) m	4m	14.7*	Bayley/Scales of Infant Development II
	Samaras K et al. 2014	880	Group A: 78.4 (4.7) No DM: 78.4 (4.8)	2y	12.6 (3.9) 11.1 (3.9)	Digit Symbol Coding, Logical Memory and Block Design from WAIS-III, TMT, RAVLT, BVRT, VFT, BNT
	Sharma VK et al. 2014	727	27.3 (2.3)	18m	8.6 (0.8)	MMSE, MSCST, RSPM, CDT, BGT, SDLT, Stroop Test, TMT, VFT
	Whalley LG et al. 2014	201	Group A: 77.2 (0.8) Group B: 77.1 (0.7) Group C: 77.1 (0.6)	5y	--	MMSE
	Ganguli M et al. 2014	559	77.6 (7.4)	4y	65% >10	--
Alzheimer and cognitive decline	Rozzini L et al. 2006	74	Group A: 69.1 (7.1) Group B: 73.9 (7.3)	1y	--	MMSE, CDR, ADAS-cog
	Sluda J et al. 2009	55	--	1y	--	--
Cardiovascular disease	Reitz C et al. 2009	678	77.4 (5.8)	5.2y	16.8 (7.9)	MMSE, BNT, COWAT, BDAE, Similarities from WAIS-R, MDRS, RDT, BVRT, SRT
	Tu MC et al. 2010	92	73.8 (9.4)	6.3m	12.9 (7.3)	MMSE, CDR, CASI
	Oulhaj A et al. 2010	97	71.9 (8.2)	1.5-9.5y	14.2 (4.8)	CAMCOG
	Silbert B et al. 2008	264	67.8 (7.7)	1y	10.1*	CERAD Auditory Verbal Learning Test, D5ST, TMT, COWAT, VFT, GPT, NART
	Narayan SK et al. 2011	182	80 (4)	44m	--	CDR, NART, TMT, VFT
	Jochimsen HM et al. 2013	416	58 (10)	3.9y	HHcy: 22 (4) No HHcy: 12 (3)	RAVLT, ROCF, VE of TEA, BSAT, VFT, DART
	Newman GC et al. 2007	3680	66.3 (10.8)	20.3m	13.4 (5)	MMSE
Cerebrovascular disease	Rowan EN et al. 2007	170	79.4*	27m	16 (5.1)	CAMCOG, CDR
	Teunissen CE et al. 2008	219	45.1 (11.1)	4y	11.2*	PASAT

Abbreviations: ADAS-cog: Alzheimer's Disease Assessment Scale-cognitive subscale, AVLT: Auditory Verbal Learning Test, BDAE: Boston Diagnostic Aphasia Examination, BGT: Bender Visual Motor Gestalt Test, BNT: Boston Naming Test, BSAT: Brixton Spatial Anticipation Test, BVRT: Benton Visual Retention Test, CAMCOG: Cambridge Cognitive Examination, CASI: The Cognitive Abilities Screening Instrument, CDR: Cognitive Drug Research, CDT: Clock Drawing Test, CERAD: Consortium to Establish a Registry for Alzheimer's Disease, COWAT: Controlled Word Association Test, DART: Dutch Version of the National Adult Reading Test, DM: Diabetes mellitus, DRST: Delayed Recognition Span Test, DS: Digit Span, DSST: Digit Symbol Substitution Test, EBMT: East Memory Boston Test, GPT: Grooved Pegboard Test, h: hours, Hcy: serum homocysteine, HHcy: hyperhomocysteinemia, LDCT: Letter Digit Coding Test, LDST: Letter Digit Substitution Test, m: months, MDRS: Mattis Dementia Rating Scale, MMSE: Mini-Mental State Examination, MSCST: Multi Domain Cognitive Screening Test, N: number, NART: National Adult Reading Test, PASAT: Paced Auditory Serial Addition Test, PPT: Purdue Pegboard Test, RAVLT: Rey Auditory Verbal Learning Test, RCPM: Raven's Coloured Progressive Matrices, RDT: Rosen Drawing Test, ROCF: Rey-Osterrieth Complex Figure Test, RSPM: Raven Standard Progressive Matrices, SD: standard deviation, SDLT: Serial Digit Learning Test, SDMT: Symbol Digit Modalities Test, SKT: Syndrome Short Test, SRT: Selective Reminding Test, TMT: Trail Making Test, VE: Visual Elevator Test, VFT: Verbal Fluency Test, VMI: Visual-Motor Integration, WAIS-R: Revised Wechsler Adult Intelligence Scale, WLT: Word Learning Test, y: years, *: median

Table 2. Homocysteine and Cognition. Randomize clinical studies.

Population	Article	N Treatment/Placebo	Age (years) mean (SD) Treatment/Placebo	Hcy (µmol/l) mean (SD)	Treatment	Follow-up week	Neurocognitive tests
General	Lewerin C et al 2005	126/69	75.7(4.7)/75.6(4)	17.8 (5.5)/16.1 (4.5)	0.5 mg cyanocob +0.8 mg folic acid+3mg B6	16	DS, block design, digit symbol and Visual Reproduction of Wechsler, Thurstone's Picture Memory Test,
	McMahon JA et al 2006	127/126	73.6 (5.8)/73.4 (5.7)	16.8 (5.4)/16.3 (4.4)	1mg folate+0.5mg B12+10mg B6	104	MMSE, RAVLT, WMS, COWAT, TMT, RSPM, NART, VFT
	Eussen SJPM et al 2007	130/65	82 (5)	19 15.6 (6.6); 2 ^o 14.5 (4.4)/15.8(5.6)	1 ^o 1mg cobalamin, 2 ^o 1mg cobalamin+0.4mg folate	24	MMSE, ROCF, TMT, DS, Stroop Test, VFT, RSPM, Similarities of Wais, AVLT, FTT
	Macpherson H et al 2012	28/28	71.9 (4.8)/70.3 (4.3)	14.1 (2.8)/15.1 (2.8)	Swisse Women's Ultravite 50+TM	16	NART, SUCCAB, CVLT-II
	Pinligas A et al 2014	56/60	[M: 28.9 (6.9); F: 32.8(7.7)]/[M: 30.9 (7.5); F: 31.2 (6.8)]	10.9 (1.6)/ 10.7 (1.6)	Supplementation of Multivitamin	16	SUCCAB
	Cheng D et al 2014	57/47	74.3 (9.6)/72.5 (7.0)	20.6 (6.5)/19.3 (4.2)	0.8mg folate+10mg B6+0.025mg B12	14	BCATs
	Van der Zwaluw et al 2014	2919	74.1 (6.5)	14.4*	0.4mg folate+0.5mg B12+0.015 D3	104	MMSE, RAVIT, DS, TMT, Stroop Test, SDMT,VFT
	Harris E. et al 2015	52/51	60.2(3.2)/M: 59.1(2.3); F: 60.1(3.4)	M: 15.1(3.4); F: 12.9(3)/ M: 15.5(3.9); F: 13(2.9)	Supplementation of Multivitamin	16	Stroop, Immediate and Delayed Recognition Memory, Spatial Working Memory and Contextual Memory, Reaction Time.
	Dangour A.D. et al 2015	99/102	79.9 (3.5)/80.1(3.7)	17.1(4.6)/17.2 (5.6)	1 mg Vitamin B12	52	CVLT, VFT, SLMT, Reaction Time.
	Alzheimer and cognitive decline	Aisen PS et al 2008	204/140	75.7 (8.0)/77.3 (7.9)	9.2 (3.4)/9.1 (2.8)	5mg folate+ 1mg B12+25mg B6	78
Kwok T et al 2011		70/70	79.1 (6.7)/72.2 (7.9)	14.3 (4.3)/13.9 (3.8)	5mg folate+ 1mg methylcobalamin	104	MMSE, MDRS
De Jager CA et al 2012		110/113	--	11.3*/11.6*	0.8mg folate+0.5mg B12+ 20mg B6	104	MMSE, HVLT-R, CERAD, CLOX, CDR, IQCODE
Parkinson's Disease	Köbe T. et al 2015	13/9	70(7.2)/70 (5.2)	15.9(7)/17.6(3.6)	Omega-3+aerobic exercise+CS.	26	MMSE, AVLT, TMT, DS, VFT, Stroop
	Barone P et al 2008	224/118	--	--	Rivastigmine	24	ADAS-cog
Schizophrenia	Litvinenko IV et al 2010	32/30	71.2 (5.8)/72.4 (3.7)	--	20mg memantine	52	MMSE, ADAS-cog, FAB, CDT, VFT
	Levine J et al 2006	20/22	40*/40*	>15	2mg folate+25mg pyridoxine+0.4mg B12	26	DS, RAVLT, WCST, ROCF or TCF
Eating Disorders	Loria V et al 2013	14/10	22.3 (7.6)/26.7 (10.0)	9.4 (2.4)/10(2)	5mg folate(Acfol)X2	26	TMT, Stroop Test
Kidney Disease	Brady CB et al 2009	339/320	--	--	40mg folate+100mg B6+2mg B12	52	TICSm, DS of WAIS, VFT

Abbreviations: ADAS-cog: Alzheimer's Disease Assessment Scale-cognitive subscale, AVLT: Auditory Verbal Learning Test, BCATs: Basic Cognitive Aptitude Tests, CDR: Cognitive Drug Research, CDT: Clock Drawing Test, CERAD: Consortium to Establish a Registry for Alzheimer's Disease, CLOX: executive clock drawing test, COWAT: Controlled Word Association Test, CS: Cognitive Stimulation; CVLT-II: California Verbal Learning Test, DS: Digit Span, FAB: Frontal Lobe Dysfunction Assessment Battery, FTT: Finger Tapping Test, HVLT-R: Hopkins Verbal Learning Test Revised, IQCODE: Informant Questionnaire on Cognitive Decline in the Elderly, MDRS: Mattis Dementia Rating Scale, MMSE: Mini-Mental State Examination, N: number, NART: National Adult Reading Test, RAVLT: Rey Auditory Verbal Learning Test, ROCF: Rey-Osterrieth Complex Figure Test, RSPM: Raven Standard Progressive Matrices, SD: standard deviation, SDMT: Symbol Digit Modalities Test, SLMT: Symbol Letter Modality Test, SUCCAB: Swinburne University Computerised Cognitive Assessment Battery, TCF: Taylor Complex Figure, TICSm: Telephone Interview for Cognitive Status-modified, TMT: Trail Making Test, VFT: Verbal Fluency Test, WAIS: Wechsler Adult Intelligence Scale, WCST: Wisconsin Card Sort Test, WMS: Wechsler Memory Scale, y: years, *: median

Table 3. Homocysteine and Cognition. Case-control studies.

Population	Article	N	Age (years) Mean (SD) Patients/Controls	Hcy (µmol/l) Mean (SD) Patients/Controls	Neurocognitive tests
Alzheimer and cognitive decline	Irizarry MC et al. 2005	377/88	72.5(70.3)(9.8)	9.1(8.7)(3.2)	BDS-1MC
	Hernanz A et al. 2007	51/44	AD 73.2(7.1); MCI 75.9(6.9)/ 73.5(3.2)	AD 14.8(7.4); MCI 13.2(6)/ 10.4(2.7)	MMSE
Parkinson's Disease	Gorgone G et al. 2009	63/64	75.7(9.2)/75.7(8.6)	19.1(4.4)/13.3(3.5)	MMSE, MDB
	Lepara O et al. 2009	30/30	80(0.9)/77.5(1)	16.1(1)/12.6(0.4)	MMSE
	Lidballe DL et al. 2011	80/83	83*/82*	15.6*/12.8*	MMSE
	Faux NG et al. 2011	101/172	AD 78.4(8.7); MCI 75.7(7.6)/ 70(7)	AD [M&F 9.7(0.3)]; MCI [M 10 (0.4), F 8.9 (0.3)] / M 9.6(0.2), F 8.4(0.1)	CVLT, RCFT, Stroop Test, VFT, Digit Symbol Coding Test, LM
Parkinson's Disease	Ford AH et al. 2012	431/1347	82.7(3.9)(81.2)(3.5)	13.3(4.5)/12.6(4.8)	TICS
	Kim G et al. 2013	200/121	74.8(7.2)	--	MMSE, BNT, Word List Memory Test, Constructional Recall Test
	Ozer F et al. 2006	39/28	67(9.3)/61.9(8.3)	15.2(9.6)/10.9(2.8)	STMS, CDT, SBST, WMS, BFR, JLO, Stroop Test, WCST, VFT
	Zoccollella S et al. 2009	121/154	--	17.5(10.2)/11(4.1)	--
Bipolar Disease	Camicicli R et al. 2009	51/50	71.5(4.7)/71.6(4.9)	13.6(3.8)/10.5(2.5)	MMSE, DHS
	Rodriguez MC et al. 2009	89/30	Group A 70(6.5), Group B 70.2(5.2), Group C 74.9(6.1)/ 68.5(3)	Group A 14.9(4.7), Group B 15.1(4.3), Group C 15.4(5.4)/ 8.5(1.9)	MMSE, BDS, FCSRT, CERAD Word List, BNT, APM, VFT, TMT, Stroop Test, DS
	Bialecka M et al. 2012	320/254	64.4(10.1)/64.8(9.6)	18(7.8)/14(9.6)	MMSE, WAIS-R, RCFT, DS, BVRT, TMT, AVLT, WCST, VFT
Depression	Dittmann S et al. 2008	74/42	42.5(12.2)/43(12.7)	10.2(5.2)/8.9(2.8)	TMT, German Versions of WAIS-R, LNST of WAIS-III, RBANS
	Dias V et al. 2009	65/49	37.8(10.5)/33.6(9.8)	1.4mg/l(0.4)/1.3mg/l(0.4)	WMS, SDMT, TMT, Stroop Test, Tower of Hanoi Test, COWAT, Bells Test, Comprehension and Similarities of WAIS-R
Schizophrenia	Alexopoulos P et al. 2010	25/25	69.8(5.8)/69.4(5.1)	13.3(5.5)/11.3(4.9)	Stroop test, D2-CP
	Ford AH et al. 2013	236/122	Hcy≥13: 62.1, Hcy<13: 62.4/ 63.8	--	MMSE, CERAD, BNT, VFT, Word List Memory Task
Cerebrovascular disease	Ayessa R et al. 2012	139/99	32.1(10.8)/27(6.1)	13.6(7.1)/11.1(3)	RAVLT, Digit Symbol of WAIS-III, GPT, Tol, RCFT, TMT, Stroop Test, CPT, LNST and Vocabulary of WAIS-III, FAS, Zoo Map Test, Eyes Task,
	Pavlovic AM et al. 2011	95/41	59.8(10.9)/57.7(10.6)	14.4(5)/8.9(3.9)	MMSE, TMT, WCST, RCFT, RAVLT, BNT, Animal Naming Test
Obstructive sleep apnea	Sales LV et al. 2013	14/13	37.2(6.9)/36(6.1)	16.7(8)/10.7(2.9)	WCST, DSST, DS, RCFT, LM and Verbal Paired Association Tests, Toulouse-Pieron Attention Test, Similarities Test
	Russo C et al. 2008	94/53	36.6(10.4)/37.1(12.1)	13.2(5.6)/9.8(2.5)	RCPM, TMT, RCFT, VFT, VSAT, PWAT, Immediate and Delayed Recall of a Short Story

Abbreviations: AD: Alzheimer's Disease, APM: Raven's Advanced Progressive Matrices, AVLT: Auditory Verbal Learning Test, BDS-1MC: Blessed Dementia Scale-Information Memory Concentration Scale, BFR: Benton's Face Recognition, BNT: Boston Naming Test, BVRT: Benton Visual Retention Test, CDT: Clock Drawing Test, CVLT: California Verbal Learning Test, CERAD: Consortium to Establish a Registry for Alzheimer's Disease, COWAT: Controlled Word Association Test, CPT: Continuous Performance Test, D2-CP: D2 Concentration Performance Test, DS: Digit Span, DSST: Digit Symbol Substitution Test, F: Female, FAS: FCSRT: Free and Cued Selective Reminding Test, GPT: Grooved Pegboard Test, Hcy: serum homocysteine, HHcy: hyperhomocysteinemia, JLO: Judgement Line Orientation, LM: Logical Memory, LNST: Letter-Number Sequencing Subtest, M: Male, MCI: Mild Cognitive Impairment, MDB: Mental Deterioration Battery, MMSE: Mini-Mental State Examination, PWAT: Paired Word Association Test, RAVLT: Rey Auditory Verbal Learning Test, RBANS: Repeatable Battery for the Assessment of Neuropsychological Status Form A, RCPM: Raven's Coloured Progressive Matrices, RCFT: Rey Complex Figure Test, SBST: Sozel Bellek Surecleri Testi, SD: standard deviation, SDMT: Symbol Digit Modalities Test, STMS: Short Test of Mental Status, TICS: Telephone Interview for Cognitive Status, TMT: Trail Making Test, Tol: Tower of London, VFT: Verbal Fluency Test, VSAT: Visual Search and Attention Test, WAIS-III: Wechsler Adult Intelligence Scale III, WAIS-R: Revised Wechsler Adult Intelligence Scale, WCST: Wisconsin Card Sorting Test, WMS: Wechsler Memory Scale, *: median.

Table 4. Homocysteine and Cognition. Cross-sectional case studies.

Population	Article	N	Age (years) mean (SD)	Hcy (µmol/l) mean (SD)	Neurocognitive tests	
General	Aleman A et al. 2005	400	60.2 (11.3)	13.2 (4.0)	MMSE, DS, DSST, RAVLT, TMT, DART, VFT, D&P	
	Elias MF et al. 2005	2096	GroupA: 45.3(2.7); GroupB: 54.6(3.2); GroupC: 65.2(4)	GroupA: 9.5(3.5); GroupB: 9.8(3.7); GroupC: 10.3(3.7)	Similarities of WAIS, WMS, LM, VR, HRNB, VOT, BNT	
	Ravaglia G et al. 2005	540	G1 77.4 (8.0); G2 73.1 (6.0); G3 72.1 (5.3)	G1 14.3; G2 12.7; G3 11.1	MMSE	
	Schafer JH et al. 2005	1047	59.3*	10 (4.1)	BNT, VFT, RCMP, ROCF, TMT, RAVLT, Stroop Test, PPT, FTT, Symbol Digit	
	Clark MS et al. 2005	200	59.9 (2.5)	10 (2.7)	CVLT, LNST, NART, Ten Unrelated Words	
	Wright CB et al. 2006	3298	66.7 (8.4)	9.2 (3.4)	MMSE	
	Feng L et al. 2006	451	64.2 (6.7)	13 (4.6)	RAVLT, TMT, SDMT, VFT, VR, DS, SSP, Block Desing Test	
	Fischer P et al. 2006	606	75.8 (0.4)	M:14.7 (4.9)/ F: 14 (6)	MMSE	
	De Lau L et al. 2007	1033	72.2 (7.4)	11.5 (4.1)	Stroop test, VFT, AVLT, MST	
	Elias MF et al. 2008	911	No-ApoE-ε1:62.6 (12.9)/ ApoE-ε1:60.7 (12.3)	No-ApoE-ε1:10 (3.6)/ ApoE-ε1:9.8 (3.2)	MMSE, Maine-Syracuse Neuropsychological Test Battery	
	Vidal JS et al. 2008	3914	74.2	15.2 (5.6)	MMSE, TMT, BVRT, IST, FWT	
	Koike T et al. 2008	99	75.4 (8.1)	11.0 (4.9)	MMSE	
	Chin AV et al. 2008	466	75.4 (6.1)	13.9 (6.1)	MMSE, NART-R, VFT, WMS, Digit Symbol Coding of WAIS	
	West RK et al. 2011	199	86.8 (4.0)	11.81 (4.3)	TMT, VFT, BNT, Shipley Vocabulary	
	Tangney CC et al. 2011	121	78.7 (5.7)	10.9 (3.1)	LM, EBMT, JLO, RSPM, SDMT, BNT, NART, DS	
	Xu LL et al. 2012	1412	>65	--	Short Portable Mental Status Questionnaire	
	Moorthy D et al. 2012	1955	66.8	10.7	MMSE	
	Feng L et al. 2013	228	65.4 (6.2)	13.4 (3.8)	DS, SSP, RAVLT, VFT, SDMT, TMT, VAT, DF, OANB	
	Allam M et al. 2013	45	68.1 (3.9)	18.6 (12.3)	MMSE, ACE, Clinical Dementia Rating Scale	
	Strain JJ et al. 2013	210	5.6 (0.3)	6.1 (1.4)	FTT, PLS, WI, CBCL, KBIT	
	Kong HG et al. 2013	662	71.6 (8.8)	15.9 (7.3)	MMSE, BCAT	
	Cheng Y et al. 2014	1889	73.4 (5.9)	17.2 (8.4)	CSID, CERAD, VFT, BNT, WLT, IUFT, Stick Desing	
	Aruna Agrawal et al 2015	639	--	M:12.7 (4.1) F:10.98 (3.82)	Medicaid Systems	
	Raszewski G. et al 2015	170	56.45 (3.54)	14.9 (7.5)	FTT, SDMT, Continuous Performance Test, Verbal Memory, Shifting Attention Test	
	Alzheimer and cognitive decline	Adunsky A et al. 2005	186	82.6 (7.1)	13.2 (2)	MMSE
		Manders M et al. 2006	157	83*	16.9 (1.5)	MMSE, ADAS-cog
		Kim J et al. 2007	1215	68.7 (5.4) (60-85)	--	Mayo Clinic Criteria
		388	69.2 (5.6)	17.6 (7.4)		
		827	68.4 (5.3)	15.7 (4.8)		
Sala I et al. 2008		325	74.4 (9.1)	12.8 (5.1)	MMSE, BDS, LM, BVRT, CERAD, BNT, Stroop Test, VFT, TMT, RSPM	
Li L et al. 2008		191	72.6 (9.2)	10.6 (3.6)	MMSE, WMS-R, CERAD, VFT, BNT, CDT, DS, WAIS-R, JLO, TMT, BVMT	
Nilsson K et al. 2010		448	75*	13.6*	MMSE	
Sachdev PS et al. 2012		757	78.5 (4.7)	11.5 (4.2)	MMSE, NART, TMT, RAVLT, LM, BNT, BVRT, VFT, BDT, Digit Symbol Coding of WAIS	
Bonetti F. et al 2015		318	G1: 76.5(5.6); G2: 76.8(4.9); G3: 79.5(4.3); G4: 78.2(4.9)	G1:11.2*; G2:12.2*; G3:24*; G4:19.2*	MMSE	

ADAS-cog	Annanaki T et al. 2008	40	60.8	9.3	MMSE, WAIS-R, WMS-R, WMS-III, BADS, Stroop Test, VFT
	Martin J et al. 2010	58	66 (8.3) (with levo: 66.8 (8.2); without levo: 62.8 (8.4))	13.06; with levo 14 (5.1) without levodopa 9.8 (2.5)	MMSE, CDT, VFT, Test Barcelona, NPI
Schizophrenia	Gonzalez C et al. 2014	41	32.3 (9.1)	16.5 (9.7)	WMS-III, RAVLT, WCST, TMT, JLO
	Frieling H et al. 2005	26	AN:27.8 (10.4); BN 24.5 (8.4)	12	TMT, Stroop Test, ROCF, HAWIER-R, D2, Wechsler Memory Screen Revised
Eating Disorders	Wilhelm J et al. 2010	20	AN 26.3 (8.6); BN 35.5 (5.9)	AN 12.6 (6.0); BN 10.9 (1.9)	D2, Stroop Test, SKT, MWT-B
	Osher Y et al. 2008	57	39.4 (12.7)	--	WCST, Stroop Test, BVRT, RAVLT, DS, DSST, Block Desing, Animal Naming
Cerebrovascular disease	Wong A et al. 2006	57	noHHcy 68.3 (12.2) HHcy 73.0 (6.8)	--	MMSE, ADAS-cog, MDRS Initiation/Perseveration Subtest
	Tay SY et al. 2006	169	65.5 (11.7)	TACI/PACI 14.5 (5.8); POCI 14.3 (4.2); LACI 13.7 (5.2)	MMSE
Cardiovascular disease	Gunstad J et al. 2006	128	69.1 (7.6)	11.7 (4.9)	MMSE, DS, GPT, COWAT, BVMT-R, BNT, VFT, CFT, VOT, Block Desing, CVLT, Stroop Test, TMT, DRS, Digit Symbol Coding, Similarities
	Kloppenborg RP et al. 2011	763	59 (10) HHcy 62.6 (10.7) NO HHcy 58.0 (9.9)	M: 14.13 (6.41); F:12.65 (4.71)	RAVLT, ROCF, VE, BSAT, VFT, DART
	Ford AH et al. 2012	155	Hcy <15: 65.5 (10.1) Hcy ≥15: 68.8 (10.4)	--	CAMCOG, CVLT, Digit Copy and Coding of WAIS
Kidney Disease	Tamura MK et al. 2011	3591	58.2 (11.0)	--	3MS
	Troen AM et al. 2012	183	54 (9.5)	15.4 (5.5)	NART, TMT, Digit Symbol Coding, Block Desing
Geriatric Patients	Hengstermann S et al. 2009	189	78.6 (7.3)	18.8*	MMSE, SKT
Multiple System Atrophy	Chen D. et al 2015	47	58.74 (10.18)	13.28 (4.13)	MMSE
Epilepsy	Di Rosa G et al. 2013	179	14.0 (4.2)	9.7 (3.1)	Total Intelligence quotient, Verbal Intelligence quotient, Performance Intelligence quotient

ADAS-cog: Alzheimer's Disease Assessment Scale-cognitive subscale; AN: Anorexia Nervosa; AVLT: Auditory Verbal Learning Test; BADS: Behavioral Assessment of the Dysexecutive Syndrome; BCAT: Basic Cognitive Aptitude Test; BDT: Block Desing Test; BDS: Blessed Dementia Scale; BN: Bulimia Nervosa; BNT: Boston Naming Test; BSAT: Brixton Spatial Anticipation Test; BVMT: Brief Visual Memory Test; BVRT: Benton Visual Retention Test; CAMCOG: Cambridge Cognitive Examination; CBCL: Child Behavior Checklist; CDT: Clock Drawing Test; CERAD: Consortium to Establish a Registry for Alzheimer's Disease; CFT: Complex Figure Test; CSID: Community Screening Instrument for Dementia; CVLT: California Verbal Learning Task; DART: Dutch Version of the National Adult Reading Test; DF: Design Fluency; DRS: Dementia Rating Scale; DS: Digit Span; DSST: Digit Symbol Substitution Test; D&P: Doors and People Memory Test; D2: Concentration Performance; EBMT: East Memory Boston Test; FTT: The finger-tapping test; FWT: The five word test; GPT: Grooved Pegboard Test; HAWIER-R: Hamburg Wechsler Intelligence; Hcy: serum homocysteine; HHcy: hyperhomocysteinemia; HRNB: Halstead-Ritan Neuropsychological Test Battery; IST: Isaacs Set Test; IUFT: IU Token Test; JLO: Judgement Line Orientation; KBIT: Kaufman Brief Intelligence Test; LM: Logical Memory; LNST: Letter-Number Sequencing Subtest; MMSE: Mini-Mental State Examination; N: number; MST: Memory Scanning Test; MWT-B: Mehrfachwahl-Wortschatz-Intelligenztest; NART: National Adult Reading Test; NPI: Neuropsychiatric Inventory; OANB: Object and Action Naming Battery; PPT: Perceptual Pattern Test; RAVLT: Rey-Osterrieth Complex Figure Test; RSPM: Raven Standard Progressive Matrices; SD: standard deviation; SDMT: Symbol Digit Modalties Test; SKT: Syndrom Kurztest; SSP: Spatial Span; TMT: Trail Making Test; VAT: Visual Association Test; VE: Visual Elevator Test; VFT: Verbal Fluency Test; VOT: The Hooper Visual Organization Test; VR: Visual Reproduction; WAIS-R: Revised Wechsler Adult Intelligence Scale; WCST: Wisconsin Card Sorting Test; WI: Woodcock-Johnson Test; WLT: Word Learning Test; WMS: Wechsler Memory Scale; 3MS: Modified Mini-Mental State Exam; *: median