

Far from the Hearth



*(Above) Martin Jones at West Stow, 1972 (with thanks to Ian Alister, Lucy Walker, Leonie Walker, and West Stow Environmental Archaeology Group); (Below) Martin Jones in a millet field, Inner Mongolia, 2010. (Photograph: X. Liu.)*





McDONALD INSTITUTE CONVERSATIONS

---

# Far from the Hearth

Essays in Honour of Martin K. Jones

Edited by Emma Lightfoot, Xinyi Liu & Dorian Q Fuller

*Published by:*

McDonald Institute for Archaeological Research  
University of Cambridge  
Downing Street  
Cambridge, UK  
CB2 3ER  
(0)(1223) 339327  
info@mcdonald.cam.ac.uk  
www.mcdonald.cam.ac.uk



McDonald Institute for Archaeological Research, 2018

© 2018 McDonald Institute for Archaeological Research  
*Far from the Hearth* is made available under a  
Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 (International) Licence:  
<https://creativecommons.org/licenses/by-nc-nd/4.0/>

ISBN: 978-1-902937-87-8

Cover image: *Foxtail millet field near Xinglonggou, Chifeng, China, photographed by Xinyi Liu, September 2014.*

Edited for the Institute by James Barrett (*Series Editor*) and Anne Chippindale.

---

---

# CONTENTS

Contributors	vii	
Figures	viii	
Tables	xvi	
Acknowledgements	xx	
<i>Foreword</i>	xxi	
JAMES H. BARRETT		
<b>Part I</b>	<b>Introduction</b>	<b>1</b>
	Introduction: Far from the Hearth	3
	XINYI LIU, EMMA LIGHTFOOT & DORIAN Q FULLER	
<b>Part II</b>	<b>A Botanical Battleground</b>	<b>7</b>
<i>Chapter 1</i>	The Making of the Botanical Battle Ground: Domestication and the Origins of the Worlds' Weed Floras	9
	DORIAN Q FULLER & CHRIS J. STEVENS	
<i>Chapter 2</i>	The Fighting Flora: An Examination of the Origins and Changing Composition of the Weed Flora of the British Isles	23
	CHRIS J. STEVENS & DORIAN Q FULLER	
<i>Chapter 3</i>	A System for Determining Plant Macro Archaeological Remains	37
	VICTOR PAZ	
<i>Chapter 4</i>	Phytoliths and the Human Past: Archaeology, Ethnoarchaeology and Paleoenvironmental Studies	51
	CARLA LANCELOTTI & MARCO MADELLA	
<i>Chapter 5</i>	Genetics and the Origins of European Agriculture	65
	TERRY BROWN	
<i>Chapter 6</i>	Martin Jones' Role in the Development of Biomolecular Archaeology	71
	TERRY BROWN, RICHARD P. EVERSLED & MATTHEW COLLINS	
<b>Part III</b>	<b>The Stomach and the Soul</b>	<b>75</b>
<i>Chapter 7</i>	'Rice Needs People to Grow it': Foraging/farming Transitions and Food Conceptualization in the Highlands of Borneo	77
	GRAEME BARKER, CHRISTOPHER O. HUNT, EVAN HILL, SAMANTHA JONES & SHAWN O'DONNELL	
<i>Chapter 8</i>	How did Foraging and the Sharing of Foraged Food Become Gendered?	95
	CYNTHIA LARBAY	
<i>Chapter 9</i>	Agriculture is a State of Mind: The Andean Potato's Social Domestication	109
	CHRISTINE A. HASTORF	
<i>Chapter 10</i>	Archaeobotanical and Geographical Perspectives on Subsistence and Sedentism: The Case of Hallan Çemi (Turkey)	117
	MANON SAVARD	
<i>Chapter 11</i>	Rice and the Formation of Complex Society in East Asia: Reconstruction of Cooking Through Pot Soot- and Carbon-deposit Pattern Analysis	127
	LEO AOI HOSOYA, MASASHI KOBAYASHI, SHINJI KUBOTA & GUOPING SUN	
<i>Chapter 12</i>	Food as Heritage	145
	GILLY CARR, MARIE LOUISE STIG SØRENSEN & DACIA VIEJO ROSE	

---

Contents

---

<b>Part IV</b>	<b><i>Between Fertile Crescents</i></b>	<b>153</b>
<i>Chapter 13</i>	From a Fertile Idea to a Fertile Arc: The Origins of Broomcorn Millet 15 Years On XINYI LIU, GIEDRE MOTUZAITE MATUZEVICIUTE & HARRIET V. HUNT	155
<i>Chapter 14</i>	A World of C <sub>4</sub> Pathways: On the Use of $\delta^{13}\text{C}$ Values to Identify the Consumption of C <sub>4</sub> Plants in the Archaeological Record EMMA LIGHTFOOT, XINYI LIU & PENELOPE J. JONES	165
<i>Chapter 15</i>	The Geography of Crop Origins and Domestication: Changing Paradigms from Evolutionary Genetics HARRIET V. HUNT, HUGO R. OLIVEIRA, DIANE L. LISTER, ANDREW C. CLARKE & NATALIA A.S. PRZELOMSKA	177
<i>Chapter 16</i>	The Adoption of Wheat and Barley as Major Staples in Northwest China During the Early Bronze Age HAIMING LI & GUANGHUI DONG	189
<i>Chapter 17</i>	When and How Did Wheat Come Into China? ZHIJUN ZHAO	199

---

## CONTRIBUTORS

GRAEME BARKER

Department of Archaeology, University of  
Cambridge, Cambridge CB2 3DZ, UK.  
*Email:* gb314@cam.ac.uk

JAMES H. BARRETT

McDonald Institute for Archaeological Research,  
University of Cambridge, Cambridge CB2 3ER, UK.  
*Email:* jhb41@cam.ac.uk

TERRY BROWN

Manchester Institute of Biotechnology, School of  
Earth and Environmental Sciences, University of  
Manchester, Manchester M1 7DN, UK.  
*Email:* Terry.Brown@manchester.ac.uk

GILLY CARR

Institute of Continuing Education, University of  
Cambridge, Cambridge CB23 8AQ, UK.  
*Email:* gcc20@hermes.cam.ac.uk

ANDREW C. CLARKE

McDonald Institute for Archaeological Research,  
University of Cambridge, Cambridge CB2 3ER, UK.  
*Email:* acc68@cam.ac.uk

MATTHEW J. COLLINS

Natural History Museum of Denmark, University of  
Copenhagen, Copenhagen DK-1123, Denmark.

&

Department of Archaeology, University of  
Cambridge, Cambridge CB2 3DZ, UK.  
*Email:* matthew.collins@snm.ku.dk

GUANGHUI DONG

MOE Key Laboratory of West China's  
Environmental System, Lanzhou University,  
Lanzhou 730000, China.  
*Email:* ghdong@lzu.edu.cn

RICHARD P. EVERSHERD

School of Chemistry, University of Bristol, Bristol  
BS8 1TS, UK.  
*Email:* R.P.Eversherd@bristol.ac.uk

DORIAN Q FULLER

Institute of Archaeology, University College  
London, London WC1H 0PY, UK.  
*Email:* d.fuller@ucl.ac.uk

CHRISTINE A. HASTORF

Department of Anthropology, University of  
California-Berkeley, Berkeley, CA 94720, USA.  
*Email:* hastorf@berkeley.edu

EVAN HILL

School of Natural and Built Environment,  
Queen's University Belfast, Belfast BT7 1NN, UK.  
*Email:* ehill08@qub.ac.uk

LEO AOI HOSOYA

Institute for Global Leadership,  
Ochanomizu University, Tokyo 112-8610, Japan.  
*Email:* hosoya.aoi@ocha.ac.jp

CHRISTOPHER O. HUNT

School of Natural Sciences and Psychology,  
University of Liverpool, Liverpool L3 5UX, UK.  
*Email:* c.o.hunt@ljmu.ac.uk

HARRIET V. HUNT

McDonald Institute for Archaeological Research,  
University of Cambridge, Cambridge, CB2 3ER, UK.  
*Email:* hvh22@cam.ac.uk

PENELOPE J. JONES

Menzies Institute for Medical Research, University  
of Tasmania, Sandy Bay, TAS 7050, Australia.  
*Email:* Penelope.Jones@utas.edu.au

SAMANTHA JONES

School of Geosciences, University of Aberdeen,  
Aberdeen AB24 3FX, UK.  
*Email:* samantha.jones@abdn.ac.uk

MASASHI KOBAYASHI

Hokuriku Gakuin University, I-11, Mitsukoji-machi,  
Kanazawa, Ishikawa Prefecture 920-1396, Japan.  
*Email:* masashi@hokurikugakuin.ac.jp

SHINJI KUBOTA

Kanazawa University, Kakuma-cho, Kanazawa,  
Ishikawa Prefecture 920-1192, Japan.  
*Email:* shinjikubota@hotmail.com

CARLA LANCELOTTI

CaSEs Research Group (Culture and Socio-  
Ecological Dynamics), Department of Humanities,  
Universitat Pompeu Fabra, Barcelona 08005, Spain.  
*Email:* carla.lancelotti@upf.edu

---

CYNTHIA LARBEBY

Department of Archaeology, University of  
Cambridge, Cambridge CB2 3DZ, UK.  
*Email:* cdal3@cam.ac.uk

HAIMING LI

MOE Key Laboratory of West China's  
Environmental System, Lanzhou University,  
Lanzhou 730000, China.  
*Email:* lihaimingboy@126.com

EMMA LIGHTFOOT

McDonald Institute for Archaeological Research,  
University of Cambridge, Cambridge CB2 3ER, UK.  
*Email:* elfl2@cam.ac.uk

DIANE L. LISTER

McDonald Institute for Archaeological Research,  
University of Cambridge, Cambridge CB2 3ER, UK.  
*Email:* dll1000@cam.ac.uk

XINYI LIU

Department of Anthropology, Washington  
University in St. Louis, St. Louis, MO 63130, USA.  
*Email:* liuxinyi@wustl.edu

MARCO MADELLA

CaSEs Research Group (Culture and Socio-  
Ecological Dynamics), Department of Humanities,  
Universitat Pompeu Fabra, Barcelona 08005, Spain.

&

ICREA (Institució Catalana de Recerca i Estudis  
Avançats), Barcelona 08010, Spain.

&

School of Geography, Archaeology and  
Environmental Studies, The University of  
Witwatersrand, Johannesburg 2000, South Africa.  
*Email:* marco.madella@upf.edu

GIEDRE MOTUZAITĖ MATUZEVIČIŪTĖ

Department of City Research, Lithuanian Institute  
of History, Vilnius 01108, Lithuania.  
*Email:* giedre.motuzaitė@gmail.com

DACIA VIEJO ROSE

Department of Archaeology, University of  
Cambridge, Cambridge CB2 3DZ, UK.  
*Email:* dv230@cam.ac.uk

SHAWN O'DONNELL

School of Natural and Built Environment, Queen's  
University Belfast, Belfast BT7 1NN, UK.  
*Email:* S.O'Donnell@qub.ac.uk

HUGO R. OLIVEIRA

Manchester Institute of Biotechnology, School of  
Earth and Environmental Sciences, University of  
Manchester, Manchester M1 7DN, UK.  
*Email:* hugo.oliveira@manchester.ac.uk

VICTOR PAZ

Archaeological Studies Program, University  
of the Philippines, Diliman, Quezon City 1101,  
Philippines.  
*Email:* vjpaz@up.edu.ph

NATALIA A.S. PRZELOMSKA

Department of Anthropology, National Museum  
of Natural History, Smithsonian Institution,  
Washington, DC 20560, USA.

&

Smithsonian's National Zoo & Conservation Biology  
Institute, National Zoological Park, Washington,  
DC 20008, USA.

*Email:* PrzelomskaN@si.edu

MANON SAVARD

Laboratoire d'archéologie et de patrimoine,  
département de biologie, chimie et géographie,  
Université du Québec à Rimouski, Québec G5L 3A1,  
Canada.

*Email:* Manon\_Savard@uqar.ca

MARIE LOUISE STIG SØRENSEN

Department of Archaeology, University of  
Cambridge, Cambridge CB2 3ER, UK.  
*Email:* mlss@cam.ac.uk

CHRIS J. STEVENS

Institute of Archaeology, University College of  
London, London WC1H 0PY, UK.  
*Email:* c.stevens@ucl.ac.uk

GUOPING SUN

Zhejiang Provincial Research Institute of Cultural  
Relics and Archaeology, Hangzhou 310014, China.  
*Email:* zjkgoffice@163.com

ZHIJUN ZHAO

Institute of Archaeology, Chinese Academy of Social  
Sciences, Beijing 100710, China.  
*Email:* zjzhao@cass.org.cn

## Figures

### Chapter 1

1.1	<i>Wild barley spikelets (Hordeum spontaneum).</i>	11
1.2	<i>Seed size increase over time standardized to percentage change, comparing Southwest Asia and China.</i>	12
1.3	<i>Charts showing founder weed taxa over time and proportion of cereals in the plant assemblage.</i>	16
1.4	<i>A field of wheat in which weedy oats and wild barley appear to be rather better than the crop.</i>	17

### Chapter 2

2.1	<i>Diagrammatic representation of seed-bank types—autumn sowing-tillage cycle.</i>	25
2.2	<i>Diagrammatic representation of seed-bank types—spring sowing-tillage cycle.</i>	25
2.3	<i>Relative presence and persistence of seed-banks types I–IV in the field after ard cultivation.</i>	26
2.4	<i>Relative presence and persistence of seed-banks types I–IV in the field after mouldboard plough cultivation.</i>	26
2.5	<i>Timeline of agricultural changes and number of introduced/reintroduced weed flora.</i>	30

### Chapter 3

3.1	<i>Identification and determination of plant macro remains.</i>	42
3.2	<i>Diagram of determination process.</i>	45

### Chapter 4

4.1	<i>Increase in phytolith studies in the last 15 years.</i>	52
-----	--	----

### Chapter 5

5.1	<i>The first ancient DNA sequences obtained from charred grain.</i>	66
-----	---	----

### Chapter 7

7.1	<i>Borneo, showing the location of the Kelabit Highlands and other locations.</i>	77
7.2	<i>Penan encampment in the Baram valley.</i>	78
7.3	<i>Kelabit longhouse at Pa'Daleh, southern Kelabit Highlands.</i>	79
7.4	<i>Map showing key sites and locations in the Kelabit Highlands.</i>	79
7.5	<i>Oxcal plots of summed probabilities from archaeological and landscape sites.</i>	82
7.6	<i>Stratigraphic summaries of the cores and geoarchaeological sites.</i>	83

### Chapter 11

11.1.	<i>Burn mark above the waterline after experimental cooking of liquid-rich food.</i>	130
11.2.	<i>The style of Jomon and Yayoi major cooking pots.</i>	131
11.3.	<i>Removing excess water after boiling rice (Central Thailand).</i>	134
11.4.	<i>Steaming stage of the yutori boil-and-steam rice-cooking method reconstructed with Yayoi pots.</i>	134
11.5.	<i>Cooking pot styles of the Tianluoshan site.</i>	136
11.6.	<i>Shift of proportions of cooking-pot styles in Hemudu culture.</i>	137
11.7.	<i>TLS round-body pots characteristic soot and burn mark.</i>	137
11.8.	<i>Layered burn deposits formed after experimental porridge cooking.</i>	138

### Chapter 12

12.1	<i>Photographs taken at the Refugee Camp in Idomeni, Greece, March/April 2016.</i>	146
12.2	<i>An example of a 'Mediterranean Diet' meal.</i>	147
12.3	<i>A Guernsey occupation-era kitchen, complete with food-related objects.</i>	149

### Chapter 13

13.1	<i>Locations of key millet sites across Eurasia.</i>	156
13.2	<i>Harriet Hunt visiting the Vavilov Herbarium, St Petersburg.</i>	158
13.3	<i>Martin Jones at a broomcorn millet field near Lanzhou, Gansu Province, western China.</i>	159
13.4	<i>Visiting millet sites in Gansu Province, western China.</i>	159

---

## Figures and Tables

---

<b>Chapter 15</b>		
15.1	<i>Martin Jones visiting the N.I. Vavilov Research Institute for Plant Industry, St Petersburg.</i>	178
15.2	<i>Barley exemplifies the complexity of inheritance of different segments of a domesticated crop's genome.</i>	183
<b>Chapter 16</b>		
16.1.	<i>Distribution of prehistoric sites in the NETP and Hexi Corridor.</i>	190
16.2.	<i>The actual yield percentage of the sites in the NETP and Hexi Corridor.</i>	192
16.3.	<i>Sum of the actual yield percentage of the sites in the NETP and Hexi Corridor.</i>	192
16.4.	<i>Carbonized plant seeds collected from Lijiaping Site.</i>	194
<b>Chapter 17</b>		
17.1.	<i>The potential routes for the spread of wheat into China.</i>	205
<b>Tables</b>		
<b>Chapter 1</b>		
1.1	<i>Presence/absence of a select roster of founder weeds.</i>	14–15
<b>Chapter 2</b>		
2.1	<i>Common weeds within British archaeobotanical assemblages.</i>	28–9
<b>Chapter 3</b>		
3.1	<i>Classifications of seeds based on preservation conditions.</i>	40
3.2	<i>Variables relevant in establishing the level of confidence of determination.</i>	40
<b>Chapter 4</b>		
4.1	<i>Phytolith production and taxonomic specificity for the world's major crops.</i>	54
<b>Chapter 6</b>		
6.1	<i>Projects and workshops funded by the NERC Ancient Biomolecules Initiative (1993–1998).</i>	73
<b>Chapter 7</b>		
7.1	<i>Radiocarbon dates from archaeological and palynological sites in the Kelabit Highlands.</i>	84–5
<b>Chapter 8</b>		
8.1	<i>Published studies on remains of starchy plants during prehistoric hunter-gatherer periods.</i>	100–101
<b>Chapter 10</b>		
10.1	<i>Archaeobotanical results from Hallan Çemi.</i>	118–19
<b>Chapter 14</b>		
14.1	<i>List of edible plants found in Haryana and their photosynthetic pathways.</i>	170–72
<b>Chapter 16</b>		
16.1.	<i>Calibrated radiocarbon data in the Hehuang Basin and Hexi Corridor.</i>	191
16.2.	<i>Charred seeds from the Lijiaping site, Linxia county, Gansu Province, China.</i>	193
<b>Chapter 17</b>		
17.1.	<i>Early wheat remains in last-century archaeological discoveries.</i>	201
17.2.	<i>Early wheat remains with only relative ages.</i>	202
17.3.	<i>Directly dated early wheat remains.</i>	203
17.4.	<i>List of archaeological cultures in the Central Asian Steppe.</i>	207

---

---

## *Acknowledgements*

The initial idea of editing this volume grew out of a conversation between Xinyi Liu and Graeme Barker at St John's College, Cambridge in June 2016. The editors subsequently discussed the provisional layout of the volume. By April of the following year, our list of agreed contributors was complete. Abstracts followed, and the chapters themselves soon after. First of all, the editors would like to pay tribute to our 36 authors, whose excellent work and timely contributions made it all possible.

For the last two-and-a-half years, the volume has been known as 'Fantastic Beasts' in order to keep it a secret from Martin. As we enter the final stage, we wish to extend our thanks to all who have ensured Martin remains blissfully unaware, including Lucy Walker, and we offer her our sincere thanks. We are extremely grateful for Harriet Hunt, Diane Lister, Cynthia Larbey and Tamsin O'Connell, who are kindly

organizing the gatherings to mark Martin's retirement and the publication of this volume.

With respect to the volume's production, we would like to thank the McDonald Institute for Archaeology Research for financial support. The McDonald Monograph Series Editor James Barrett oversaw and encouraged all aspects of this project, and we offer him sincere thanks. We would also like to acknowledge the support of Cyprian Broodbank, not least for allowing us to host the workshop at the institute, but also for his encouragement throughout all phases of the volume's implementation. Particular thanks must go to several key individuals: Anne Chippindale, Ben Plumridge, Emma Jarman, Simon Stoddart and Samantha Leggett. Finally, we are also grateful to the anonymous reviewers who recommended changes that have greatly enhanced the final version of this volume.

*Xinyi Liu, Emma Lightfoot and Dorian Fuller  
August 2018*

---

---

## Foreword

The 28-year term of Martin Jones as the first George Pitt-Rivers Professor of Archaeological Science witnessed, and in part created, a transformation in the fields of environmental and biomolecular archaeology. In this volume, Martin's colleagues and students explore the intellectual rewards of this transformation, in terms of methodological developments in archaeobotany, the efflorescence of biomolecular archaeology, the integration of biological and social perspectives, and the exploration of archaeobotanical themes on a global scale. These advances are worldwide, and Martin's contributions can be traced through citation trails, the scholarly diaspora of the Pitt-Rivers Laboratory and (not least) the foundations laid by the Ancient Biomolecules Initiative of the Natural Environment Research Council (1989–1993), which he chaired and helped create. As outlined in Chapter 6, Martin's subsequent role in the bioarchaeology programme of the Wellcome Trust (1996–2006) further consolidated what is now a central and increasingly rewarding component of archaeological inquiry. Subsequently, he has engaged with the European Research Council, as Principal Investigator of the Food Globalisation in Prehistory project and a Panel Chair for the Advanced Grant programme. As both practitioner and indefatigable campaigner, he has promoted the field in immeasurable ways, at critical junctures in the past and in on-going capacities as a research leader.

The accolades for Martin's achievements are many, most recently Fellowship of the British Academy. Yet it is as a congenial, supportive—and demanding—force within the Pitt-Rivers Laboratory that the foundations of his intellectual influence were laid. Here, each Friday morning, the archaeological science community would draw sticks to decide who would deliver an impromptu research report or explore a topical theme. Martin is among the most laid-back colleagues I have worked with, yet simultaneously the most incisive in his constructive criticism. As a provider of internal peer-review he was fearless without being unkind. The themed Pitt-Rivers Christmas parties were equally impactful—on one occasion Alice Cooper appeared, looking ever so slightly like our professor of archaeological science.

Martin's roles as a research leader extended to several stints as head of the Department of Archaeology, chairing the Faculty of Archaeology and Anthropology and serving as a long-term member of the Managing Committee of the McDonald Institute for Archaeological Research. Having started his professional career as an excavation-unit archaeobotanist in Oxford, he was a long-standing proponent of the highly successful Cambridge Archaeological Unit. In the wider collegiate community, he is a Fellow (and was Vice-Master) of Darwin College and was the staff treasurer of the Student Labour Club. In all roles he fought valiantly and often successfully for the interests of his constituency. His capacity to fight for deeply held priorities while recognizing the value of diverse perspectives was of utmost importance. His nostalgic enthusiasm for the debate with archaeological science that was engendered by the post-processual critique is one signal of an underlying appreciation of plurality. His active support for the recent merger of the Divisions of Archaeology and Biological Anthropology, within our new Department of Archaeology, is another. As a scientist (Martin's first degree, at Cambridge, was in Natural Sciences) he values the peer-reviewed journal article above all scholarly outputs, yet has authored as many highly regarded books as a scholar in the humanities. His *Feast: Why humans share food* has been translated into several languages and won Food Book of the Year from the Guild of Food Writers. He views academia and society as a continuum, campaigning for archaeobotanical contributions to global food security (e.g. by promoting millet as a drought-resistant crop) and working with world players such as Unilever to encourage archaeologically informed decisions regarding food products.

That Martin's achievements and influence merit celebration is clear. That his colleagues and students wish to honour him is equally so. Yet does the McDonald Conversations series publish *Festschriften*? This is a semantic question. As series editor I am delighted to introduce a collection of important papers regarding the past, present and future of archaeobotany, representing its methodological diversity and maturity. That this collection concurrently pays respect to a treasured colleague is a very pleasant serendipity.

Dr James H. Barrett

*Part II*  
**A Botanical Battleground**



---

## Chapter 1

# The Making of the Botanical Battleground: Domestication and the Origins of the World's Weed Floras

Dorian Q Fuller & Chris J. Stevens

'The development of plant communities on agricultural land can thus be seen in part as a battle between weed communities and human communities, in which stakes for both parties are high'

Martin Jones (1988, 86)

### Introduction

Martin Jones's work on the archaeology of British farming, pursued from the 1970s through the 1990s, combined big-picture evolutionary ecology with details of archaeobotanical evidence and individual weed ecologies. This approach considers the arable field as a habitat that is constantly evolving with changing human practice (M. Jones 1988). This was the 'botanical battleground' in which weed taxa competed with each other and the crop, and in which farmers competed with weeds. As such the arable ecosystem is defined in terms of cycles of human activity, rather than soil or climate conditions. Unlike biomes, in which shared characteristics of vegetation are determined largely by climatic constraints, the agricultural 'anthrome' (*sensu* Ellis 2011) represents something new to planet Earth from the start of the Holocene (or latest Pleistocene) created through the emergent mechanisms of culture. This has received attention in recent years as central to human niche construction or the emergence of an 'anthropocene' (e.g. Boivin *et al.* 2016; Ellis *et al.* 2013). Archaeobotany has a key role to play in documenting how these cultural ecosystems evolved and diverged. Research into the origins of agriculture has traditionally focused on social and economic transformations and the domesticated crops themselves; however, in this contribution we would like to explore the *botanical battlegrounds* that accompanied the earliest cultivation and domestication processes.

The origins of arable ecologies provide a context for the evolution of both weeds and domesticated crops from their respective wild ancestors. A weed is usually defined as a plant that grows where it is not wanted,

and as such is a human concept, as such rules do not apply in nature (Bunting 1960). Weed is a concept that arises within the history of human-plant relationships in which humans increasingly seek to control their environment. Prior to the start of cultivation weeds did not exist as such, but rather grew in their own 'natural' non-anthropogenic habitats. However, in some cases this natural habitat is a challenge to identify and Zohary (1950) classified such species as 'obligate' weeds. Nevertheless, as recognized by Harlan and de Wet (1965), there is a second definition of weed, which is a plant that thrives on disturbed ground, such as a cleared field. Such species then are pioneers and possess traits that allow for the rapid establishment of the plant and its acquisition of nutrients from the soil. This can be defined as an ecological strategy of fast resource acquisition (see Milla *et al.* 2015; Reich 2014). These ecological traits of weeds, or weediness, are shared with many domesticated cereals, suggesting parallel adaptations between crops and weeds.

As with domesticated species, some species growing in the cultivated field might be expected to evolve adaptations to this new arable ecology. Amongst such adaptations some of the key traits recognized as part of the domestication syndrome should then be considered, including changes in seed size, in germination patterns, or indeed the loss of germination. Ultimately a key distinction between weeds and crops is whether or not particular species within the cultivated field were volunteers or intentionally planted. The nature of this distinction plays an important role within the domestication syndrome; as crops evolved to be more readily harvested, so weeds utilized strategies in which they either became part of the harvest or avoided it.

### Activities of the arable and the origins of fields

One of the key distinctions that makes the archaeobotanical study of domestication processes feasible is the

distinction between evidence for human practice and evidence for evolutionary changes in plants, underwritten by genetic shifts in plant populations. This is the distinction between cultivation and domestication, a distinction perhaps best clarified in the work of Harris (1989; 2012) and Hillman and Davies (1990), but essentially a division between what people do, for example *cultivate*, and what happens to plants, *domestication* (Fuller 2007; Purugganan & Fuller 2009). This creates an evolutionary process that is inherently co-evolutionary, an entangled network of feedbacks between human practices (evolving through cultural transmission) and plant morphologies (evolving through genetic adaptations). Previously, we have explored this entanglement in terms of humans getting ‘trapped’ in ever-increasing labour investment in soil maintenance, and harvesting and crop-processing technologies, which in turn are rewarded by higher returns (Fuller *et al.* 2016). A notion that was inspired by conversations with Martin Jones is to see this as shifting interactions within the food web, with human activities influencing energy flows at many levels.

Nearly three decades ago, Martin Jones (1992, 213) highlighted the need to move beyond ‘oversimplified correlates of a “domestication event” to examining’ the wider influences of humans on the nutritional status and the species they consume, such as the soil conditions in which food plants grew. This view highlights the importance of the small details of the nature of cultivated fields, the species in them and how these competed and adapted over time. Rather than framing a singular shift from foraging to farming, we need to explore the evolving ecosystem of cultivated fields alongside the various ‘intermediate economies’ (*sensu* Harris 2012) through the two to three millennia of the protracted domestication processes (Fuller *et al.* 2014). By considering the arable system as a *botanical battleground* we can usefully frame the key variables in this transition process, in which plants favoured by people (crops) and those not (weeds) compete for resources, and in which humans strategically alter the conditions of soil, water and light resources; and through this framework we can perhaps see more clearly some of the commonalities and differences between crops and weeds in the making of agriculture.

Cultivation involved a number of transformations of the soil which established the parameters of competition. First, pre-existing vegetation was largely cleared from the small plots of cultivation. It is conceivable that small woody perennials were left in place. Seed-dispersal studies of recruitment in natural grasslands suggest that existing perennials can limit seed establishment, especially of species not already established, whereas in annual ecosystems there is

greater competition between seeds (Peart 1989). The act of cultivation creates a new type of habitat in which annual disturbance is both uniform and highly predictable, with the removal of the existing plant canopy providing repeated opportunities for seeds present in the soil seed-bank to participate. Field clearance tends to mean that sunlight is widely available for growth and germination, but faster-growing plants in the field may quickly shade out their neighbours. Tillage also creates deeper cracks, which may bury seeds more deeply than if they had fallen on natural soil surfaces. Certain human cultivation practices may counteract some of these factors. For example, planting in rows or well-spaced crops will reduce overshadowing and competition between the roots of different plants. People can also add both nutrients (manuring) and water (irrigating) to the soil, and one of the key questions asked of archaeobotanists is when such practices came about? And what methods, for example inferences from weed seed ecology or stable isotopes, can provide evidence for such practices (Bogaard *et al.* 2007; G. Jones *et al.* 2010; Madella *et al.* 2009)? Evidence from elevated  $\delta^{15}\text{N}$  in cereal grains from Greece suggests small intensively managed and manured fields (Vaiglova *et al.* 2014), something that may have been the norm for early arable systems (Bogaard 2005), with declining  $\delta^{15}\text{N}$  levels in cereal grains over the course of the Holocene suggesting a movement towards less intensive, more extensive systems (Araus *et al.* 2014; Styring *et al.* 2017). In China, early fields were also small-scale (<10 sq. m), allowing close management of water and soil, including manuring with household waste and drying out to increase rice yields and control weeds (Fuller & Qin 2009; Weisskopf *et al.* 2014; 2015).

From the point of view of plant competition, these fields appear generally nutrient-rich and therefore potentially favoured plant traits that fit a nutrient acquisitive strategy, as opposed to a conservation, or nutrient-allocation, strategy, as defined by Reich (2014) and Milla and colleagues (2015).

Many adaptations of cereal spikelets serve to facilitate the position of seeds for germination. For example, grass awns, as well as aiding animal and water dispersal, can move in daily cycles in response to ambient temperature. This action drives the spikelet along the soil surface until a suitable crack or depression is found, and in some cases enables the burial of spikelets (Kulić *et al.* 2009; Peart 1979). In wild wheats the two awns open and close on a daily cycle, serving to ratchet the spikelet into soil (Elbaum *et al.* 2007). In weedy species of oats (*Avena* sp.) the bent awn plays a key role in drilling spikelets into the soil, enabling survival through winter for spring germination (Somody



**Figure 1.1.** Wild barley spikelets (*Hordeum spontaneum*), hand-picked (left) and that have dehisced overnight and are projecting into cleared soil in the morning (right). (Photograph: D. Fuller, Iraqi Kurdistan, May 2012.)

*et al.* 1985). Experiments with the awned dicot *Erodium* indicated that seed burial was more effective in soils with plant litter than barren or compacted soil, a useful adaptation within grassland environments (Stamp 1989). While long awn morphology may be excellent for dispersal (Fig. 1.1), and it may play some role in deterring herbivory, the metabolic investment in creating awns will detract from the potential investment in seed nutrients that power the early seedling. Human harvesting and sowing, along with the development of non-shattering types, removes the need for dispersal mechanisms, hence reduced metabolic expenditure upon these structures is expected during domestication leading to a reduction in awns and barbs (Fuller 2007).


#### The evolution of seed size: automatic escalation

The new competition created by the tilled and sown field accounts for one of the key recurrent domestication traits, namely larger seed size. Increased seed size during domestication was attributed by Harlan and colleagues (1973) to selection relating to seedling vigour and competition, and to deeper planting; however, the latter explanation has often been emphasized at the expense of the former (cf. Zohary 2004). This seemingly forgotten explanation of Harlan and colleagues (1973) was that larger seed sizes in crops are expected to evolve in relation to the highly disturbed soils of early cultivation. Larger seeds have a series of competitive advantages, including being correlated with larger seedlings in many grasses and legumes

(Baskin & Baskin 2001, 214). Larger seedlings will have a head-start in competition for light and space in what, after competing vegetation is removed, is effectively a level playing field, as sown grain or grain from the seed-bank germinates. Hence larger grains have a selective advantage, while conversely, the competitive advantage of smaller grain sizes that might aid dispersal and burial though reduced seed mass is lost.

Fuller (2007) emphasized depth of burial as a possible cause of increased grain size, but while supported experimentally in some taxa it was not in others (Kluyver *et al.* 2013). Larger seeds had advantages in seedling emergence in lentil (*Lens culinaris*), mungbean (*Vigna radiata*), cowpea (*Vigna unguiculata*), lima bean (*Phaseolus lunatus*) and more weakly in pea, but no significant correlation was recorded for soybean, peanut or common bean. A further difference, among the pulses tested, was between species processing hypogeal germination, in which cotyledons remain in the soil providing food for the seedling, and those with epigeal germination, in which cotyledons are raised above the soil, where they become photosynthetic. As might be expected species processing hypogeal germination were better at emerging from depth generally, and it may be that selection for larger seeds in epigeal species might increase the photosynthetic area, providing more resources for initial growth (Kluyver *et al.* 2013).

That seed size increase predominantly correlates with domestication, not just in cereals and pulses grown for their seeds, but in numerous vegetables grown for their leaves and tubers, such as lettuce,



**Figure 1.2.** Seed size increase over time standardized to percentage change, comparing Southwest Asia (10,000–5000 BC) and China (6000–1000 BC) for selected crops. Linear regressions indicated for some taxa to illustrate trends. (Raw data from Fuller et al. 2014, except melon, from Fuller 2012.)

potato, beet, carrot and parsnip, indicates that seed-size increase was an evolutionary outcome arising from the cultivated environment (Kluyver *et al.* 2017). It is possible that this trait may be linked to other correlated traits, such as overall biomass of other organs that are linked in development to seed size, that is the effect of allometry or pleiotropy. Such changes, however, took millennia (Fuller *et al.* 2014; 2017), hence differences between generations within average seed size occurred on a minute scale that would be difficult to measure even with modern scientific techniques, let alone apparent to the naked eye. As such it is implausible that seed-size increase with initial domestication could be a target of conscious human manipulation. Instead, seed-size increase took place as part of the crops becoming incorporated into new arable ecologies, calling for more application of toolkits of comparative functional ecology to understanding domestication (Milla *et al.* 2015).

Archaeobotanical evidence allows us to put the timing and extent of changes in seed size into their cultural and geographical context and to explore comparisons across crops. Despite the effects of charring that may reduce seed sizes variably, charred archaeological seeds still document chronological trends during episodes of domestication (Fuller 2018; Fuller *et al.* 2014; 2017). Previous work has compiled

time series data for a range of annual crops, including Near Eastern cereals and pulses, North American composites, sumpweed (*Iva annua*) and sunflower (*Helianthus annuus*), Chinese (japonica) rice (*Oryza sativa*) and soybean (*Glycine max*) and Indian (indica) rice (*Oryza sativa*) and mungbean (*Vigna radiata*) (Fuller *et al.* 2012; 2014; Purugganan & Fuller 2011). One observation of Kluyver and colleagues (2017) is that the total size increase in cereals and pulses, grown for their seeds, is generally greater than that in vegetable crops. Indeed, when archaeobotanical data for size increase are plotted together, by standardizing these in terms of percentage change from the original (earliest/smallest size), some comparisons are striking (Fig. 1.2). First, it can be seen that in the Near Eastern cereals and representative pulses (lentil and pea, *Pisum sativum*) the trends of seed-size change are similar, with similar rates and total amount of change (average maximum being 45–65 per cent larger over 4000 years), with emmer wheat showing the slowest trend (although pea has a less clear trend). For China, rice showed a total increase towards the lower end of this spectrum at *c.* 50 per cent, while much more rapid and greater increase was evident in the soybean (>100 per cent increase) and in melons (*Cucumis melo*). Melon-seed size may be selected in part by simple allometry, as selection for larger fruits would developmentally

increase seed size, but selection from increased competitiveness in the botanical battleground of the cultivated field may have played a fundamental role, particularly in the early stages of domestication. Conscious selection of traits, such as fruit size or seed size, would be expected to increase the speed of change, and on this point it is worth noting that seed size in Chinese melons is relatively rapid in comparison with changes in cereal grain size (Fuller *et al.* 2014). Tree fruit seeds may also increase in size somewhat more rapidly (Fuller 2018).

Domestication of crops represents convergent evolution, involving similar adaptations. In this sense, crops emerged through domestication as tested warriors on the botanical battleground, with highly acquisitive ecological strategies. Indeed, crops appear to have been selected from wild ancestors that lay on the more acquisitive end of the annual herbs within a flora, processing characteristics that made them more adaptable to increased competition and disturbance (Cunniff *et al.* 2014). Sometimes, however, crops combine traits that are at odds with competitive adaptation within the ecological setting of their wild progenitors. For example, seed number and seed size can be regarded as trade-offs (e.g. Sadras 2007) in which plants may gain a competitive advantage through producing a greater number of seeds or by producing fewer, larger seeds (Harlan *et al.* 1973). However, both grain size and number have tended to increase with domestication. As crops come to lack the fall-back strategies of a seed-bank or perennating organs, this high investment and consumption habit can make them vulnerable to invaders that are less needy, the weeds, against which human cultural practices must evolve and adapt.

### The sources of weeds in early Western Asia

Archaeobotanical evidence tells us that weeds have been persistent within crops throughout the Old World for many millennia. So where did these weeds come from? And how did some come to be such strong actors in the arable theatre?

The list of plant species reported as weeds of cultivation worldwide is staggering, in the tens of thousands (Randall 2002), covering a diverse range of plant families and genera. However, it is unlikely they evolved *de novo* with the creation of the first arable fields, so in answer to where weeds came from, we might rather ask: what was the original geography and habitat of the 'wild progenitors' of weeds?

Just as crops have evolved from wild relatives, we should perhaps think of weeds as also deriving from wild weed progenitors. It may be the case that populations of the same taxonomic species can still

be found in less anthropogenic 'natural' habitats, the so-called 'facultative weeds' (Harlan & de Wet 1965; Hartmann-Shenkman *et al.* 2015; Zohary 1950). Other weeds, however, have been termed 'homeless' or 'obligatory' (Harlan & de Wet 1965; Hartmann-Shenkman *et al.* 2015; Willcox 2012; Zohary 1950), indicating taxa that are unknown outside their arable and highly anthropogenic habitats. In other words, the original habitat of their ancestors, pre-dating cultivation, either no longer exists in its original form, or the ancestral forms of these species have since become extinct.

The presence of these 'obligatory weeds' on early sites in the Levant, alongside early domesticated crops, or morphologically wild cereals, has emerged as a key argument for recognizing the beginnings of cultivation (Colledge 2002; Hartmann-Shenkman *et al.* 2015; Willcox 2012). At Epipaleolithic Abu Hureyra (11,200–10,100 BC), Hillman (2000) argued for emergence of an arable ecology based on increases in potential weed taxa alongside morphologically wild rye and einkorn wheat. While this is a suggestive pattern, its statistical robustness has been questioned and the data reinterpreted as broadening of plant diet and a shift in foraging across a wider range of environments; in other words, cultivation was not required as an explanation for the changes seen (Colledge & Conolly 2010).

The few large-sized grains of rye and einkorn from Abu Hureyra could suggest some cultivation, as their size falls near the upper end of the range in late Early PPNB sites (see Fuller 2012, fig 5.3), but occasional transient cultivation, alongside a predominant strategy of collecting from wild stands, is both more plausible and likely. Nevertheless, the taxa at Abu Hureyra, mainly rye and some einkorn wheat, were not the key founder crops of more widespread cereal agriculture, that is barley and emmer wheat. So the notion that there was a single centre of agricultural origins has passed into intellectual history.

In the early Holocene, evidence for a more extensive weed flora is found alongside morphologically wild and evolving cereals that were increasingly acquiring a domesticated character. Willcox (2012) compiled a list of 19 indicator weeds, drawn from obligatory and facultative weed lists, from which he excluded taxa with edible seeds and ruderals that might have grown upon human settlements. The facultative weeds mainly have their alternative habitat in the steppe through to the desert margins (Zohary 1950; 1962). In this regard many facultative weeds originate on the drier end of the spectrum from cereals that are regarded as native to the transition zone from steppe to open woodland (Hillman 2000). Only in some cases can these weeds be definitely identi-

**Table 1.1.** Presence/absence of a select roster of founder weeds, expanded from Willcox (2012) to include some taxa discussed by Hartmann-Shenkman et al. (2015), and other key weedy grasses. Note that not all wild seed taxa are included, as some hard-seeded taxa or minute taxa may survive from animal dung or be processed as food in their own right (e.g. *Chenopodiaceae*, *Cyperaceae*, *Polygonaceae*, *Juncaceae*). Primary archaeobotanical primary sources, cereal proportions and median ages are those reviewed in Maeda et al. (2016, supplementary materials), with additions from Arranz-Otaegui et al. (2016). These data are drawn from all types of contexts, but the presence of charred cereal grains suggests that crop/food processing is a major input to these assemblages.

		Northern sites																		
Age: 1000s BC		10.6	11	9.9	9.6	9.5	9.1	9.1	8.6	8.4	7.7	7.5	7.4	7.4	7.1	6.9	6.8	6.7	6.6	6.2
sites	Obligate/Facultative	Abu Hureyra 1	Dederiyeh	Qaramel	Mureybit-II	Hallan Cemi	Mureybit-III	Jarf al Ahmar	Djade	Çayönü (bp/ch)	Asikli Höyük	Halula	Abu Hureyra 2	Cafar Huyok IV-1	Ras Shamra (Vc)	Bouqras	Nebi Mend	El Kowm	Çatal Höyük	Sabi Abyad II
% Cereals		3.9	1.5	23	0.6	1.4	75	46.4	15.5	50.2	19.5	31.8	46.4	25.9	8.6	25	71.5	22.4	44	58.8
<i>Adonis</i>	O		X	X		X		X	X	X		X	X		X		X		X	X
<i>Bellevalia</i>	O	X		X	X	X	X	X	X		X	X			X			X	X	X
<i>Bupleurum</i>	O			X																
<i>Centaurea</i>	O		X		X	X	X	X	X			X	X			X			X	X
<i>Fumaria</i>	O		X		X	X		X	X	X		X	X							
<i>Galium</i>	O	X	X	X	X	X	X	X	X	X	X	X	X	X		X	X	X	X	X
<i>Glaucium</i>	O	X		X				X	X			X	X						X	X
<i>Heliotropium</i>	O	X	X	X	X	X	X	X	X		X	X	X	X		X		X	X	X
<i>Lolium temulentum</i>	O														X					
<i>Ornithogalum</i>	O	X		X				X	X				X		X					
<i>Papaver</i>	O			X				X				X	X				X	X		
<i>Phalaris</i>	O			X				X		X		X			X				X	X
<i>Silene/Gysposila</i>	O/F	X	X	X	X	X	X	X	X	X	X	X	X	X		X	X	X	X	X
<i>Teucrium</i>	O					X			X		X	X	X						X	
<i>Vaccaria</i>	O					X			X	X		X							X	
<i>Valerianella</i>	O								X		X	X			X				X	X
<i>Aegilops</i>	F			X				X	X					X	X	X		X		X
<i>Avena</i>	F	X										X			X					
<i>Coronilla</i>	F			X				X	X			X								
<i>Crucianella</i>	F		X	X				X			X	X			X			X	X	
<i>Erodium</i>	F	X	X					X					X						X	X
<i>Lolium cf. remotum</i>	F																			
<i>Lolium sp./ perenne/rigidum</i>	F	X		X				X	X	X		X		X	X	X	X	X	X	X
<i>Onobrychis</i>	F	X	X					X	X			X	X	X				X		
<i>Thymelaea</i>	F					X	X	X	X	X	X				X	X			X	
<i>Trifoliae</i>	F	X		X	X	X	X	X	X	X	X	X	X			X	X		X	X
<i>Trigonella astroites</i>	F				X		X	X	X		X	X				X				X
<b>Total weed taxa</b>		<b>11</b>	<b>9</b>	<b>15</b>	<b>8</b>	<b>11</b>	<b>8</b>	<b>20</b>	<b>19</b>	<b>9</b>	<b>10</b>	<b>20</b>	<b>13</b>	<b>6</b>	<b>11</b>	<b>9</b>	<b>6</b>	<b>9</b>	<b>15</b>	<b>14</b>

fied to species level in archaeological material. For example, Hartmann-Shenkman and colleagues (2015) were able to identify to species level 5 obligate weeds, as well as a longer list of 39 facultative weeds, from Atlit-Yam, dating to c. 6900 BC. Nevertheless, the long

list of these taxa and their recurrence across sites with both early domesticated and pre-domesticated (or intermediate) cereal finds suggests that the emergence of a weed flora was part and parcel of agricultural origins.


The Making of the Botanical Battleground

Table 1.1. (Continued).

		Southern sites															
Age: 1000s BC		9.9	9.4	9.3	9.1	9	8.6	8.6	8.5	7.9	7.7	7.6	7.5	7.4	7.2	7	6.9
sites	Obligate/Facultative	Geshet	Gilgal	Iraq ed-Dubb	Neiv Hagdud	Zahrat Adh-Dhra 2	el-Hemmeh	Aswad	Tell Qarassa	Beidha	Wadi Jilat 7	Aswad-2	Nahal Hemar	Ghoratifé	Basta	Ramad	Aitit Yam
% Cereals		0.5	?	16	28.8	36.3	52.6	71.2	47.2	0.5	8.3	74.9	4.9	51.9	29.6	82.3	20.5
<i>Adonis</i>	O				X			X	X		X	X		X		X	X
<i>Bellevalia</i>	O				X			X				X	X	X		X	
<i>Bupleurum</i>	O					X										X	X
<i>Centaurea</i>	O	X			X		X		X		X	X	X			X	X
<i>Fumaria</i>	O	X	X		X		X		X		X	X				X	X
<i>Galium</i>	O	X		X	X		X	X	X		X	X		X	X	X	
<i>Glaucium</i>	O			X												X	
<i>Heliotropium</i>	O				X	X			X							X	X
<i>Lolium temulentum</i>	O																X
<i>Ornithogalum</i>	O							X	X			X		X		X	
<i>Papaver</i>	O																
<i>Phalaris</i>	O	X			X		X		X				X	X		X	X
<i>Silene/Gysposila</i>	O/F			X	X		X	X	X			X		X	X	X	
<i>Teucrium</i>	O															X	
<i>Vaccaria</i>	O							X	X			X		X		X	
<i>Valerianella</i>	O								X								
<i>Aegilops</i>	F	X	X	X	X	X	X			X		X				X	
<i>Avena</i>	F		X		X	X		X	X		X	X	X	X	X	X	X
<i>Coronilla</i>	F															X	
<i>Crucianella</i>	F											X				X	
<i>Erodium</i>	F	X	X		X		X										
<i>Lolium cf. remotum</i>	F															X	
<i>Lolium sp./ perenne/ rigidum</i>	F							X	X	X		X		X		X	X
<i>Onobrychis</i>	F				X	X		X	X				X			X	
<i>Thymelaea</i>	F					X		X	X			X		X			X
<i>Trifoliae</i>	F	X					X	X	X		X	X	X	X	X	X	
<i>Trigonella astroites</i>	F				X		X	X	X			X		X		X	
<b>Total weed taxa</b>		<b>7</b>	<b>4</b>	<b>4</b>	<b>13</b>	<b>6</b>	<b>9</b>	<b>12</b>	<b>16</b>	<b>2</b>	<b>6</b>	<b>15</b>	<b>6</b>	<b>12</b>	<b>4</b>	<b>22</b>	<b>10</b>

A broader analysis of these data suggests the diversity of weed species increases during the pre-pottery Neolithic with greater cereal use (Table 1.1; Fig. 1.3). Thus, as cereal consumption increases, so does the evidence for a greater range of key weed taxa, implying that weed seeds were preserved through charring of crop-processing waste. For the southern Levant, the strength of this relationship is stronger

in the PPNB ( $r^2=0.769$  for the Late PPNB) than in the PPNA ( $r^2=0.169$  for PPNA), suggesting that over the era of domestication the arable ecological niche and its associated flora became increasingly entangled. Part of this can be attributed to the evolution and adaptation of key weed species shifting from their previous ecological strategies into emergent arable ecosystems. Additional factors, like the adoption of domesticated



**Figure 1.3.** (Left) Chart showing the number of founder weed taxa over time (from the select list in Table 1.1), plotted by site against median age; averages for northern and southern regions calculated for each millennium; (right) Chart showing correlation between number of founder weed taxa and total proportion of cereals in the plant assemblage, suggesting a positive correlation between weed presence and cereal dependency, especially in the southern Levant.

animals and the use of their dung as fuel, might also contribute to greater wild seed diversity, but taxa that are well known to survive dung, for example Chenopodiaceae, Cyperaceae and Polygonaceae (Filipović 2014; Spengler 2018), are not included amongst our founder-weed roster. A few minute-seeded grasses are also associated with surviving in dung (Filipović 2014), but are not included in our list. Small legumes (*Trifolia*, *Trigonella*) are ambiguous, however, and could be derived from dung; but we have left them on our weeds list following that of Willcox (2012). In addition, the predominance of cereal grains, alongside other larger grasses, which are normally digested and not included in dung (Wallace & Charles 2013), highlights major inputs from agriculture/food into the archaeobotanical record.

In a few cases we can point to potential morphological evolution in weeds that likely accompanied adaptation to cultivation, human harvesting and sowing. Large weed seeds accompanying cereal grains into storage are likely to get dispersed with sowing, thus creating selection for seed characteristics that mimic the crop, including potentially changes in size and the loss of dormancy mechanisms. An interesting case is provided by *Bupleurum*. Like most Apiaceae, *Bupleurum* spp. typically disperse as individual separated mericarps. In the obligate weed *B. subovatum*, however, mericarps remain fused in pairs, which make them closer in size to grains and spikelets, and this trait likely evolved as an adaptation to dispersal with seed-corn prior to 6900 BC (Hartmann-Shenkman *et al.* 2015).

*Lolium temulentum* (darnel) is another obligate weed, a large-grained grass close in length to barley. Available genetic data indicate that it is phylogenetically close to *L. remotum*, a flax weed primarily distributed across northern Eurasia, with which it

is interfertile, although both are predominantly self-fertilizing, much like wheat/barley (Charmet *et al.* 1996). Likewise it is also interfertile with *L. persicum*, which has a broadly Middle Eastern distribution from Baluchistan to Anatolia (Davis 1985). *L. temulentum* appeared only towards the end of the Pre-Pottery Neolithic, with examples from Atlit Yam (c. 6900 BC) and Ras Shamra VC (c. 7100 BC) (Hartmann-Shenkman *et al.* 2015; van Zeist & Bakker-Heeres 1986). The *Lolium* from Ramad (c. 7300 BC) was shorter, like *L. remotum* (see van Zeist & Bakker-Heeres 1985, 511), or perhaps *L. persicum*. *L. rigidum/perenne* types were widespread in the Neolithic Near East, making precise identifications a continuing challenge. It is plausible that once *L. persicum* invaded early cultivated fields, it differentiated into *L. remotum* and *L. temulentum*. *L. temulentum* evolving longer grains that mimic harvested barley or wheat grains that would be hard to remove during processing (Harlan & de Wet 1965). Subsequently *L. temulentum* was to spread as a frequent cereal weed through both Pakistan and Europe.

### Secondary domestications: weeds as sources of crops

In some cases, weeds became so well adapted to cultivation that they could even out-compete crops. Some of these ‘weeds’ themselves then became valued as resources that ultimately became domesticated. A farmer observing a weed-infested field (Fig. 1.4) might dismay at the reduced harvest of the favoured crop, but in times of need might decide that gathering the grains of these weeds would also provide an alternative source of calories, as recorded for *Bromus secalinus* (bromegrass) in Europe (M. Jones 1988), eventually cultivating the weed itself, turning it into



**Figure 1.4.** A field of wheat (*Triticum aestivum*) in which weedy oats (*Avena fatua*) and wild barley (*Hordeum spontaneum*) appear to be rather better than the crop. (Photograph: D. Fuller, Iraqi Kurdistan, May 2012.)

a crop. These are what botanists have referred to as secondary domesticates (e.g. Vavilov 1992). One way to explain these domestications is that they represent a case of conscious selection by farmers, who decided to transform a weed using the model of existing crops, thereby rapidly breeding it into a domesticate. But it is also possible that this began through inadvertent outcomes of the co-evolutionary battles of arable field, between weeds and farmers.

Europe's cultivated oat is a classic example of a secondary domesticate. *Avena sativa* (Oat) was itself domesticated from a weed (*Avena ludoviciana* or the *A. sterilis* complex) that in all likelihood was evolving for millennia as a weed of cultivation. Today *A. ludoviciana* is found on fallow fields and field edges, and river banks and oak scrub (Davis 1985), where one suspects it has been invasive from arable fields. Its ancestor has been shown to be *Avena sterilis* (Loskutov 2008), a native to Mediterranean and steppic habits of the Near East, growing upon limestone slopes and calcareous coastal soils, and is a recurrent weed on many early sites (Table 1.1). The widespread weedy oat today, *Avena fatua*, has no native habitat, and rep-

resents a probable parallel derivation from *A. sterilis* (Loskutov 2008). The genus *Avena* as a whole is largely circum-Mediterranean (Baum 1977), and while there is evidence for short-lived early cultivation of *A. sterilis* during the PPNA in Israel (Weiss *et al.* 2006), there is no evidence for a lasting tradition of cultivation or oat domestication in the Near East. Instead, the oat crops we know today appear to have been domesticated in central or eastern Europe around the Late Bronze Age to Early Iron Age and by the first millennium AD were widespread as a cultivated domesticate. They came into their own in the more marginal environments of northern Europe, Ireland and Scotland from around 2000 years ago, and possibly earlier in Scandinavia (Grabowski 2011). An unanswered question is whether or not the naked oat, widely cultivated in cooler and higher elevation parts of China, Tibet and the Himalayas, is derived from the same domestication. More likely, it represents a further secondary domestication of weedy *A. sterilis/ludoviciana* that dispersed eastwards with wheat and barley during the later Neolithic or Early Bronze Age (Stevens *et al.* 2016). The naked, free-threshing grains of east Asian

cultivated oats (*Avena nuda*) fit alongside other winter cereals in this zone, naked barley and free-threshing bread wheat; whereas European oat retained its hull, joining an agricultural milieu already dominated by hulled cereals, spelt wheat and hulled barley. This highlights how secondary domesticates were selected in each region for features paralleled within existing domesticates.

Oat domestication and secondary cereal domestications have been little studied. Like other cereals, oats have spikelets that do not dehisce from the panicle, and this can be diagnosed in preserved spikelet bases. It is conceivable this trait was unconsciously selected initially within weedy oats where it evolved as a mechanism by which they were more likely harvested and sown with seed corn. This appears the case for the semi-domesticated *A. abyssinica* in Ethiopia, probably derived from the wild shattering oats, *A. barbata*, and variant *A. vavilovii*, all weeds of highland wheat and barley (Baum 1977; Ladizinsky 1975). In contrast to *A. barbata* and *A. vavilovii*, *A. abyssinica* is shorter, blending into wheat and barley fields, has grains similar in size to barley and non-shattering spikelets. These spikelets are readily harvested by sickle, then threshed and processed and consumed with the main cereal crop, and in some cases it is cultivated on its own.

This example provides a model for the evolutionary trajectory for cultivated oats (*A. sativa*), in which domestication traits, probably greater grain size, then non-shattering, evolved through adaptations resulting from escalating co-evolutionary feedbacks through which weedy oats became an ever better mimic of the main crop, probably barley, in which at first it was tolerated as an edible weed, through to cultivation in its own right. In this scenario the evolution of secondary domesticates is just as unconscious as primary domestications (Fuller *et al.* 2010) and might be similarly protracted.

Mimicry of crops by weeds during their vegetative growth phase is a further common outcome of the botanical battleground. It is likely that all traditions of cultivation involve some degree of field weeding or roguing to remove competition to increase crop productivity, potentially selecting for weeds that look increasingly like the crop. The case of *A. abyssinica* is one case in point, being shorter in stature, whereas many wild oats stand tall above cereals. Others include *Camelina sativa* ssp. *linicola* N. Zing. that mimics flax in vegetative characters, has synchronous flowering with the crop and non-dehiscent capsules (Barrett 1983). Another form, *C. sativa* var. *crepitans* Sinskaya, has dehiscent capsules and co-occurs with rare dehiscent flax forms (*Linum usitatissimum* spp. *crepitans* Elladi).

Another well-documented mimic is *Echinochloa crus-galli* (barnyard millet: Barrett 1983). The wild form, barnyard grass, is widespread in wetlands across Eurasia, commonly occurring as a weed of rice. In Japan, a subspecies Japanese barnyard millet, *E. crus-galli* var. *utilis* (Ohwi & Yabuno) Kit. was cultivated and domesticated, during the Middle Jomon period long before the arrival of domesticated rice from China (Crawford 2011; Yabuno 1987). However, another weedy subspecies of this grass, *E. crus-galli* var. *oryzicola* (Vasinger) Ohwi, is well adapted to flooded paddy fields, mimicking rice in appearance from its seedling stage throughout its vegetative growth, making weeding near impossible (Barrett 1983), but usually flowering and setting seed before the rice harvest (de Wet *et al.* 1983a). In parts of the Caucasus in Russia a non-shattering form of *E. crus-galli* var. *oryzicola* has evolved in rice fields (also called *E. macrocarpa* Vasinger), in which spikelets remain on the panicle. These are reportedly cultivated sometimes in their own right and made into beer and flat breads (de Wet *et al.* 1983a), thus providing a parallel spectrum of adaptations to those of weedy and domesticated oats.

Rice fields have provided a potentially rich habitat for the evolution of other secondary domesticates. Kimata and colleagues (2000) proposed that all the native species of millets in India originated as weeds of rice, as their wild forms commonly occur in rice fields. However, this appears incorrect, as some native millets form primary staple foods within regional Neolithic traditions, for example *Panicum sumatrense* (little millet) in northwest India (Fuller 2006; Weber & Kashyap 2016) and *Brachiaria ramosa* (browntop millet) in southern India (Fuller 2006; Kingwell-Banham & Fuller 2014), before the arrival of rice. But it is likely true for Kodo millet (*Paspalum scrobiculatum*), the wild form being a widespread weed of rice, especially in dry (rainfed) fields (de Wet *et al.* 1983b; Moody 1989; Weisskopf *et al.* 2014). Early archaeobotanical finds comprise occasional grains associated with assemblages dominated by other millets or rice, but during the Iron Age on the Indian Peninsula it occurs with very high frequency and ubiquity, often out-numbering all other crops (Cooke & Fuller 2015), with plumper-grained, domesticated type forms occurring alongside narrower grain (wild types) (e.g. Kajale 1984). As the main form of early rice cultivation in India was likely rain fed (Fuller & Qin 2009; Weisskopf *et al.* 2014), the potential for poor yields due to low rainfall or drought and competition from weeds would have been high. In this context the more prolific grain-producing weeds, such as *Paspalum scrobiculatum*, could have been increasingly

attractive as fall-back foods, eventually evolving into domesticated crops.

### Losing the battle, winning the war?

In conceptualizing the arable field as a battleground between crops and their human allies and weeds, Martin Jones provided a framework that recognized a dynamic history in agriculture. Overall, secondary domesticates have received less attention than primary crops, but are key representatives of the botanical battleground, helping to adapt agriculture to a wider range of environmental conditions as humans took traditional crops beyond their native ranges, and hedging against crop failures through diversification.

Over the long history of agriculture, not only have weed assemblages changed, but the species that constitute weeds have evolved, and in this sense the arable ecosystems of the world represent a dynamic and changing anthropogenic ecology. Archaeobotanists have a unique vantage point, and a duty, to reveal more about this battleground. For one thing, agriculture has had and continues to have an unparalleled impact on global ecosystems, cultural stability and human population dynamics. Yet most scientific agricultural research draws on a shallow time depth of experiments and historical knowledge, whereas archaeobotany offers an approach to a holistic history of agricultural ecosystems.

### Acknowledgements

This paper was supported by a European Research Council grant 'Comparative Pathways to Agriculture' (ComPag, no. 323842).

### References

- Araus, J.L., J.P. Ferrio, J. Voltas, M. Aguilera & R. Buxóet, 2014. Agronomic conditions and crop evolution in ancient Near East agriculture. *Nature Communications* 5, 3953. DOI: 10.1038/ncomms4953
- Arranz-Otaegui, A., S. Colledge, L. Zapata, L.C. Teira-Mayolinid & J.J. Ibáñez, 2016. Regional diversity on the timing for the initial appearance of cereal cultivation and domestication in southwest Asia. *Proceedings of the National Academy of Sciences* 113(49), 14001–6.
- Barrett, S.H., 1983. Crop mimicry in weeds. *Economic Botany* 37(3), 255–82.
- Baskin, C.C. & J.M. Baskin, 2001. *Seeds. Ecology, biogeography, and evolution of dormancy and germination*. New York (NY): Academic Press.
- Baum, B.R., 1977. *Oats: Wild and Cultivated. A monograph of the genus Avena L. (Poaceae)*. Ottawa: Canada Department of Agriculture.
- Bogaard, A., 2005. 'Garden agriculture' and the nature of early farming in Europe and the Near East. *World Archaeology* 3(2), 177–96.
- Bogaard, A., T.H.E. Heaton, P. Poulton & I. Merbach, 2007. The impact of manuring on nitrogen isotope ratios in cereals: archaeological implications for reconstruction of diet and crop management practices. *Journal of Archaeological Science* 34, 335–43.
- Boivin, N.L., M.A. Zeder, D.Q. Fuller, *et al.*, 2016. Ecological consequences of human niche construction: examining long-term anthropogenic shaping of global species distributions. *Proceedings of the National Academy of Sciences* 113(23), 6388–96.
- Bunting, A.H., 1960. Some reflections on the ecology of weeds, in *The Biology of Weeds*, ed. J.L. Harper. (British Ecological Society Symposium 1.) Oxford: Blackwell Scientific, 11–26.
- Charmet, G., F. Balfourier & V. Chatard, 1996. Taxonomic relationships and interspecific hybridizations in the genus *Lolium* (grasses). *Genetics Resources and Crop Evolution* 43, 319–27.
- Colledge, S., 2002. Identifying pre-domestication cultivation in the archaeobotanical record using multivariate analysis presenting the case for quantification, in *The Dawn of Farming in the Near East*, eds. R.T.J. Cappers & S. Bottema. Berlin: ex oriente, 141–52.
- Colledge, S. & J. Conolly, 2010. Reassessing the evidence for the cultivation of wild crops during the Younger Dryas at Tell Abu Hureyra, Syria. *Environmental Archaeology* 15, 124–38.
- Cooke, M. & D.Q. Fuller, 2015. Agricultural continuity and change during the Megalithic and Early Historic Periods in South India, in *Megalithic traditions in India. Archaeology and ethnography, Volume 2*, eds. K.K. Basa, R.K. Mohanty & S.B. Ota. Delhi: Aryan Books International, 445–76.
- Crawford, G.W., 2011. Advances in understanding early agriculture in Japan. *Current Anthropology* 52(S4), S331–45.
- Cunniff, J., S. Wilkinson, M. Charles, G. Jones, M. Rees & C.P. Osborne, 2014. Functional traits differ between cereal crop progenitors and other wild grasses gathered in the Neolithic Fertile Crescent. *PLoS ONE* 9(1), e87586.
- Davis, P.H., 1985. *Flora of Turkey, Volume 9*. Edinburgh: Edinburgh University Press.
- de Wet, J.M.J., K.E. Prasada Rao, M.H. Mengesha & D.E. Brink, 1983a. Domestication of Sawa millet (*Echinochloa colona*). *Economic Botany* 37(3), 283–91.
- de Wet, J.M.J., K.E. Prasada Rao, M.H. Mengesha & D.E. Brink, 1983b. Diversity in Kodo millet, *Paspalum scrobiculatum*. *Economic Botany* 37(2), 159–63.
- Elbaum, R., L. Zaltzman, I. Burgert & P. Fratzl, 2007. The role of wheat awns in the seed dispersal unit. *Science* 316(5826), 884–6.
- Ellis, E.C., 2011. Anthropogenic transformation of the terrestrial biosphere. *Philosophical Transactions of the Royal Society of London A* 369(1938), 1010–35.
- Ellis, E.C., J.O. Kaplan, D.Q. Fuller, S. Vavrus, K.K. Goldewijk & P.H. Verburg, 2013. Used planet: a global history. *Proceedings of the National Academy of Sciences* 110(20), 7978–85.

- Filipović, D., 2014. Early Farming in Central Anatolia: An archaeobotanical study of crop husbandry, animal diet and land use at Neolithic Çatalhöyük. (BAR International series S2667.) Oxford: Archaeopress.
- Fuller, D.Q., 2006. Agricultural origins and frontiers in South Asia: a working synthesis. *Journal of World Prehistory* 20, 1–86.
- Fuller, D.Q., 2007. Contrasting patterns in crop domestication and domestication rates: recent archaeobotanical insights from the Old World. *Annals of Botany* 100(5), 903–24.
- Fuller, D.Q., 2012. New archaeobotanical information on plant domestication from macro-remains: tracking the evolution of domestication syndrome traits, in *Biodiversity in Agriculture: Domestication, evolution, and sustainability*, eds. P. Gepts, T.R. Famula & R.L. Bettinger. Cambridge: Cambridge University Press, 110–35.
- Fuller, D.Q., 2018. Long and attenuated: comparative trends in the domestication of tree fruits. *Vegetation History and Archaeobotany* 27(1), 165–76.
- Fuller, D.Q., R.G. Allaby & C.J. Stevens, 2010. Domestication as innovation: the entanglement of techniques, technology and chance in the domestication of cereal crops. *World Archaeology* 42(1), 13–28.
- Fuller, D.Q., S. Colledge, C. Murphy & C. J. Stevens, 2017. Sizing up cereal variation: patterns in grain evolution revealed in chronological and geographical comparisons, in *Miscelánea en homenaje a Lydia Zapata Peña (1965–2015)*, eds. J. Fernández Eraso, J.A. Mujika Alustiza, Á.A. Valbuena & M. García Díez. Bilbao: Servicio Editorial Universidad Del País Vasco, 131–49.
- Fuller, D.Q., T. Denham, M. Arroyo-Kalin, *et al.*, 2014. Convergent evolution and parallelism in plant domestication revealed by an expanding archaeological record. *Proceedings of the National Academy of Sciences* 111(17), 6147–52.
- Fuller, D.Q. & L. Qin, 2009. Water management and labour in the origins and dispersal of Asian rice. *World Archaeology* 41(1), 88–111.
- Fuller, D.Q., C.J. Stevens, L. Lucas, C. Murphy & L. Qin, 2016. Entanglements and entrapment on the pathway toward domestication, in *Archaeology of Entanglement*, eds. L. Der & F. Fernadini. Walnut Creek (CA): Left Coast Press, 151–72.
- Fuller, D.Q., G. Willcox & R. Allaby, 2012. Early agricultural pathways: moving outside the ‘core area’ hypothesis in Southwest Asia. *Journal of Experimental Botany* 63, 617–33.
- Grabowski, R., 2011. Changes in cereal cultivation during the Iron Age in southern Sweden: a compilation and interpretation of the archaeobotanical material. *Vegetation History and Archaeobotany* 20(5), 479–94.
- Harlan, J.R. & J.M.J. de Wet, 1965. Some thoughts about weeds. *Economic Botany* 19, 16–24.
- Harlan J.R., J.M.J. de Wet & E.G. Price, 1973. Comparative evolution of cereals. *Evolution* 27, 311–25.
- Harris, D.R., 1989. An evolutionary continuum of people-plant interaction, in *Foraging and Farming: The evolution of plant exploitation*, eds. D.R. Harris & G.C. Hillman. London: Routledge, 11–26.
- Harris, D.R., 2012. Evolution of agroecosystems: biodiversity, origins, and differential development, in *Biodiversity in Agriculture. Domestication, evolution, and sustainability*, eds. P. Gepts, T.R. Famula, L. Bettinger, S.B. Brush, A.B. Damania, P.E. McGuire & C.O. Qualset. Cambridge: Cambridge University Press, 21–56.
- Hartmann-Shenkman, A., M. Kislev, E. Galili, Y. Melamed & E. Weiss, 2015. Invading a new niche: obligatory weeds at Neolithic Atlit-Yam, Israel. *Vegetation History and Archaeobotany* 24, 9–18.
- Hillman, G.C. & M.S. Davies, 1990. Measured domestication rates in wild wheats and barley under primitive cultivation, and their archaeological implications. *Journal of World Prehistory* 4(2), 157–222.
- Hillman, G.C., 2000. Abu Hureyra 1: the Epipalaeolithic, in *Village on the Euphrates: The excavation of Abu Hureyra*, eds. A.M.T. Moore, G.C. Hillman & A.J. Legge. New York (NY): Oxford University Press, 327–98.
- Jones, G., M. Charles, A. Bogaard & J. Hodgson, 2010. Crops and weeds: the role of weed functional types in the identification of crop husbandry methods. *Journal of Archaeological Science* 37, 70–77.
- Jones, M.K., 1988. The arable field: a botanical battleground, in *Archaeology and the Flora of the British Isles: Human Influence on the Evolution of Plant Communities*, ed. M.K. Jones. Oxford: Oxford University Committee for Archaeology, 86–92.
- Jones, M.K., 1992. Food remains, food webs and ecosystems. *Proceedings of the British Academy* 77, 209–19.
- Kajale, M.D., 1984. New light on agricultural plant economy during 1st millennium B.C.: palaeobotanical study of plant remains from excavations at Veerapuram, District Kurnool, Andhra, Pradesh, in *Veerapuram: A type site for cultural study in the Krishna Valley*, eds. T.V.G. Sastri, M. Kasturibai & J.V. Prasada Rao. Hyderabad Birla: Archaeological and Cultural Research Institute, 1–15.
- Kimata, M., E.G. Ashok & A. Seetharam, 2000. Domestication, cultivation and utilization of two small millets, *Brachiaria ramosa* and *Setaria glauca* (Poaceae), in South India. *Economic Botany* 54(2), 217–27.
- Kingwell-Banham, E. & D.Q. Fuller, 2014. Brown top millet: origins and development, in *Encyclopaedia of Global Archaeology*, ed. C. Smith. New York (NY): Springer, 1021–4.
- Kluyver, T.A., M. Charles, G. Jones, M. Rees & C.P. Osborne, 2013. Did greater burial depth increase the seed size of domesticated legumes? *Journal of Experimental Botany* 64(13), 4101–8.
- Kluyver, T.A., G. Jones, B. Pujol, *et al.*, 2017. Unconscious selection drove seed enlargement in vegetable crops. *Evolution Letters* (1–2), 64–72.
- Kulić, I.M., M. Mani, H. Mohrbach, R. Thaokar & L. Mahadevan, 2009. Botanical ratchets. *Proceedings of the Royal Society of London B: Biological Sciences* 276(1665), 2243–7.
- Ladizinsky, G., 1975. Oats in Ethiopia. *Economic Botany* 29, 238–41.
- Loskutov, I.G., 2008. On evolutionary pathways of *Avena* species. *Genetic Resources and Crop Evolution* 55, 211–20.

- Madella, M., M.K. Jones, P. Echlin, A. Powers-Jones & M. Moore, 2009. Plant water availability and analytical microscopy of phytoliths: implications for ancient irrigation in arid zones. *Quaternary International* 193, 32–40.
- Maeda, O., L. Lucas, F. Silva, K-I. Tanno & D.Q Fuller, 2016. Narrowing the harvest: increasing sickle investment and the rise of domesticated cereal agriculture in the Fertile Crescent. *Quaternary Science Reviews* 145, 226–37.
- Milla, R., C.P. Osborne, M.M. Turcotte & C. Violle, 2015. Plant domestication through an ecological lens. *Trends in Ecology & Evolution* 30(8), 463–9.
- Moody, K., 1989. *Weeds Reported in Rice in South and Southeast Asia*. Laguna: IRRI.
- Peart, M.H., 1979. Experiments on the biological significance of the morphology of seed-dispersal units in grasses. *Journal of Ecology* 67(3), 843–63.
- Purugganan, M.D. & D.Q Fuller, 2009. The nature of selection during plant domestication. *Nature* 457, 843–8.
- Purugganan, M.D. & D. Q Fuller, 2011. Archaeological data reveal slow rates of evolution during plant domestication. *Evolution* 65(1), 171–83.
- Randall, R.P., 2002. *A Global Compendium of Weeds*. Melbourne: R.G. & F.J. Richardson.
- Reich, P.B., 2014. The world-wide ‘fast-slow’ plant economics spectrum: a traits manifesto. *Journal of Ecology* 102, 275–301.
- Sadras, V.O., 2007. Evolutionary aspects of the trade-off between seed size and number in crops. *Field Crops Research* 100(2), 125–38.
- Somody, C.N., J.D. Nalewaja & S.D. Miller, 1985. Self-burial of wild oat florets. *Agronomy Journal* 77(3), 359–62.
- Spengler, R.N., 2018. Dung burning in the archaeobotanical record of West Asia: where are we now? *Vegetation History and Archaeobotany*. DOI: 10.1007/s00334-018-0669-8
- Stamp, N.E., 1989. Efficacy of explosive vs. hygroscopic seed dispersal by an annual grassland species. *American Journal of Botany*, 555–61.
- Stevens, C.J., C. Murphy, R. Robert, L. Lucas, F. Silva & D.Q Fuller, 2016. Between China and South Asia: a Middle Asian corridor of crop dispersal and agricultural innovation in the Bronze Age. *Holocene* 26(10), 1541–55.
- Styring, A.K., M. Charles, F. Fantone, *et al.*, 2017. Isotope evidence for agricultural extensification reveals how the world’s first cities were fed. *Nature Plants* 3, 17076.
- Vaiglova, P., A. Bogaard, M. Collins, *et al.*, 2014. An integrated stable isotope study of plants and animals from Kouphovouno, southern Greece: a new look at Neolithic farming. *Journal of Archaeological Science* 42, 201–21.
- van Zeist, W. & J.H. Bakker-Heeres, 1985. Archaeobotanical studies in the Levant 1. Neolithic sites in the Damascus Basin: Aswad, Ghoraifé, Ramad. *Palaeohistoria* 24, 165–256.
- Vavilov, N.I., 1992. *Origin and Geography of Cultivated Plants* [trans. D. Love]. Cambridge: Cambridge University Press.
- Wallace, M. & M. Charles, 2013. What goes in does not always come out: the impact of the ruminant digestive system of sheep on plant material, and its importance for the interpretation of dung-derived archaeobotanical assemblages. *Environmental Archaeology* 18(1), 18–30.
- Weber, S. & A. Kashyap, 2014. The vanishing millets of the Indus civilization. *Archaeological and Anthropological Sciences* 8(1), 8–15.
- Weiss, E., M. Kislev & A. Hartman, 2006. Autonomous cultivation before domestication. *Science* 312, 1608–10.
- Weisskopf, A.R., E. Harvey, E. Kingwell-Banham, M. Kajale, R. Mohanty & D.Q Fuller, 2014. Archaeobotanical implications of phytolith assemblages from cultivated rice systems, wild rice stands and macro-regional patterns. *Journal of Archaeological Science* 51, 43–53.
- Weisskopf, A., L. Qin, J.L. Ding, P. Ding, G.P. Sun & D.Q Fuller, 2015. Phytoliths and rice: from wet to dry and back again in the Neolithic Lower Yangtze. *Antiquity* 89, 1051–63.
- Willcox, G., 2012. Searching for the origins of arable weeds in the Near East. *Vegetation History and Archaeobotany* 21, 163–7.
- Yabuno, T., 1987. Japanese barnyard millet (*Echinochloa utilis*, Poaceae) in Japan. *Economic Botany* 41(4), 484–93.
- Zohary, D., 2004. Unconscious selection and the evolution of domesticated plants. *Economic Botany* 58, 5–10.
- Zohary, M., 1950. The segetal plant communities of Palestine. *Plant Ecology* 2, 387–411.
- Zohary, M., 1962. *Plant Life of Palestine, Israel and Jordan*. New York (NY): Ronald.

