

Chapter/Review 12
Clinical Review for IDCNA IDC 33.4
Emerging and Re-Emerging Infectious Diseases

Title:

Human Monkeypox - Epidemiological and Clinical characteristics, Diagnosis and Prevention

Authors*:

Eskild Petersen, Anu Kantele, Marion Koopmans, Danny Asogun, Adesola Ogunlete, Chikwe Ihekweazu and Alimuddin Zumla

(* All authors contributed equally)

Institutional affiliations:

Eskild Petersen: Institute of Clinical Medicine, University of Aarhus, Denmark and The Royal Hospital, Muscat, Oman. European Society for Clinical Microbiology and Infectious Diseases Task Force for Emerging Infections, Basel, Switzerland. Email: eskind.petersen@gmail.com

Anu Kantele: Inflammation Center, Helsinki University Hospital and Helsinki University, Helsinki, Finland. Email: anu.kantele@helsinki.fi

Marion Koopmans: Viroscience Department, Erasmus Medical Centre, Rotterdam, The Netherlands. Email: m.koopmans@erasmusmc.nl

Danny Asogun: Department of Public Health, Faculty of Clinical Sciences, College of Medicine, Ambrose Alli University, Ekpoma, Nigeria. Email: asogun2001@yahoo.com

Adesola Ogunleye: Nigeria Centre for Disease Control, Jabi, Abuja, Nigeria.
Email: adesola.ogunleye@ncdc.gov.ng

Chikwe Ihekweazu: Nigeria Centre for Disease Control, Jabi, Abuja, Nigeria.
Email: chikwe.iheweazu@ncdc.gov.ng

Alimuddin Zumla: Division of Infection and Immunity, Center for Clinical Microbiology, University College London, and the National Institute of Health Research Biomedical Research Centre at UCL Hospitals, London, UK. Email: a.zumla@ucl.ac.uk

Mailing Addresses:

Institute of Clinical Medicine, University of Aarhus, Palle Juul-Jensens Boulevard 82, DK-8200 Aarhus N, Denmark

Inflammation Center, Helsinki University Hospital and Helsinki University, Stenbäckinkatu 9, PO BOX 100, FI-00029 HUS, Helsinki, Finland

Nigeria Centre for Disease Control, Plot 801, Ebitu Ukiwe Street, Jabi, Abuja, Nigeria.

Viroscience Department, Erasmus Medical Centre, Rotterdam, The Netherlands, Postbus 2040, 3000 CA Rotterdam, The Netherlands

Division of Infection and Immunity, Center for Clinical Microbiology, University College London, Gower Street, London, WC1E 6BT, United Kingdom

Keywords: Monkeypox, smallpox, West Africa, epidemic

Correspondence: Professor Eskild Petersen: Email: eskild.petersen@gmail.com

SYNOPSIS/ABSTRACT

Human monkeypox virus (MPXV) is a double-stranded DNA virus of the Orthopoxvirus genus of the family Poxviridae. Thought to be a rare and self-limiting disease, monkeypox has not attracted much attention since its first discovery in 1958. Recently, concern has been raised about its emergence and the occasionally severe clinical presentation bearing resemblance of its clinical presentation to smallpox, a deadly disease globally eradicated by vaccination 40 years ago. In September 2018, monkeypox drew the attention of global media, politicians and scientists, when three individual patients in the United Kingdom were diagnosed with monkeypox. The frequency and geographic distribution of human monkeypox cases across West and Central Africa have increased in recent years. On September 22, 2018, the Nigeria Centre for Disease Control (NCDC) commenced an outbreak investigation following the identification of a suspected case of monkeypox in an 11-year old child and as of 1st January 2019, there have been 311 suspected cases reported from 26 states (132 confirmed cases affecting children and adults of all ages) and 7 deaths. The majority of confirmed Monkeypox cases are under 40 years with a median age of 31 years, similar to the observed age range in DRC. This part of the population lack of cross-protective immunity since they were born only after the smallpox eradication campaigns had been discontinued. We review the epidemiology, clinical features and management of Monkeypox and discuss the growing public health threat of Monkeypox in light of discontinuation of the smallpox vaccination.

Key Points

- Human monkeypox is a zoonosis caused by the monkeypox virus (MPXV), a double-stranded DNA virus of the family Poxviridae.
- The frequency and geographic distribution of human monkeypox cases across West and Central Africa have increased in recent years
- The clinical presentation of monkeypox is similar to that of smallpox, in terms of symptoms onset, timing of rash occurrence, and rash distribution, but generally less severe than smallpox with lower fatality rate and scarification.
- The majority of confirmed Monkeypox cases are under 40 years with a median age of 31 years, a population born only after discontinuation of the smallpox vaccination campaign, and thus may reflect lack of cross-protective immunity.

INTRODUCTION

Human monkeypox (MPX) virus (MPXV) is a double-stranded DNA virus of the Orthopoxvirus genus of the family Poxviridae.^{1,2,3,4} Two genetic clades of the monkeypox virus have been characterized: West African and Central African. MPXV is one of the four orthopox virus species pathogenic for human, the other three are 1) variola major virus (VARV), the causative agent of smallpox, now eradicated, 2) variola minor virus (VARMV) and 3) cow pox virus (CPXV). There is a range of animal poxviruses several of which have zoonotic potential. Infections in humans have been described for vaccinia virus, cowpox virus, buffalopox virus and sporadic cases of camel pox.^{5,6} Monkeypox infects a wide range of mammalian species, but its natural host reservoir remains unknown.

PUBLIC HEALTH IMPORTANCE

Thought to be a rare and self-limiting disease,⁷ monkeypox has not attracted much attention since its discovery 70 years ago. The frequency and geographic distribution of human monkeypox cases have increased in recent years in a specific region of Africa (**Fig. 1**),⁸ and monkeypox has been recognized as increasing public health threat, particularly in regions in West Africa where there is close interaction between humans and wild animal reservoirs and in particular where there is evidence that the infection attack rate is increasing. The clinical presentation of monkeypox is similar to that of smallpox,² in terms of symptoms onset, timing of rash occurrence, and rash distribution,⁷ but generally less severe than smallpox in terms of complication rate, case fatality rate, and levels of scarification.

Recently, concern has been raised about its emergence as well as the resemblance of its clinical presentation to that of smallpox, a deadly disease globally eradicated by vaccination 40 years ago.⁹ During outbreaks, it has been challenging to clinically distinguish monkeypox from chickenpox, an unrelated herpesvirus infection. However, also the sporadic zoonotic infections with other orthopoxviruses calls for vigilance. Outbreaks of buffalopox have occurred with multiple human cases in India.¹⁰ Similarly, during outbreaks of vaccinia virus infection in cattle in Brazil, there is documented evidence of human infections.¹¹

Cross-immunity and protection

Various orthopox virus species share genetic and antigenic features,^{12,13,14} and an infection by any of these species may confer substantial protection against infection by the others.¹⁵ Vaccination with VACV protects against disease caused by VARV, MPXV, or CPXV.¹⁶ The immunological mechanisms underlying cross-protection by immunization with VACV appear to be diverse, with neutralizing antibodies among the principal components.¹⁷ Consistent with the ability of smallpox vaccine to provide cross-protection for humans against monkeypox, monkeys can be protected against monkeypox by immunization with the human smallpox vaccine.^{18,19}

Ever since smallpox vaccinations were discontinued in 1978, cross-protective immunity to various *Orthopox* viruses has waned, particularly in younger individuals lacking vaccinia induced immunity and the number of unvaccinated, susceptible individuals has grown worldwide. Indeed, these changes have been accompanied by an increased frequency and geographic distribution of human monkeypox cases in recent years.

EPIDEMIOLOGY

Discovery and animal reservoirs

The monkeypox virus was first detected in 1958 in an outbreak of a vesicular disease among captive monkeys transported to Copenhagen, Denmark from Africa for research purposes. Hence the name monkeypox.²⁰ The term is inappropriate since the largest animal reservoirs of the virus have been found in rodents, including squirrels and giant pouched rats both of which are hunted for food.²¹ Rodents are the largest group of mammals with over 1500 species. The extent of the wild animal reservoir, the natural history and pathogenesis of monkeypox in both animal and humans remains unknown, requiring characterization through ecological and epidemiological studies. Thus far, MPXV has been detected in diverse animal species: squirrels (rope and tree), rats, striped mice, dormice, and monkeys. In 1985, the virus was isolated from a rope squirrel in the Congo (DRC) and a dead infant mangabey monkey in Tai National Park, Cote d'Ivoire.²² During a large monkeypox outbreak following introduction of the virus through

animals imported into an animal trading company, at least 14 species of rodents were found to be infected.²³

Like humans, monkeys are considered disease hosts. Further studies are needed to understand how the virus persists in nature, explore pathogen-host associations and the effect of climatic and ecological factors influencing the shifts between geographic areas and the virus as a cause of disease in humans.²⁴

Transmission of MPXV to humans

Not only the specific animal host reservoir of monkeypox, but also mode of transmission of MPXV from animals to humans remain unknown. Aerosol transmission has been demonstrated in animals,^{25,26} and may explain a nosocomial outbreak in the Central African Republic.²⁷ However, indirect or direct contact with live or dead animals is assumed to be the driver of human monkeypox infections in humans^{28,29} Poverty and continued civil unrest force people to hunt small mammals (bushmeat) for obtaining protein-rich food, thus increasing exposure to wild rodents, which may carry monkeypox.³⁰

In August 1970 the first human case of monkeypox was identified in a 9-year-old child with smallpox-like vesicular skin lesions in the village of Bukenda in the Equatorial region of Zaire (now Democratic Republic of Congo - DRC).³¹ This patient was found during a period of intensified smallpox surveillance conducted 9 months after the eradication of smallpox in DRC had been certified by WHO.

Geographical endemicity and rise in number of cases

Ever since its discovery, the disease has been endemic to Central and West Africa with intermittent, sporadic cases of monkeypox transmitted from local wildlife reported among humans. Retrospective studies indicated that similar cases had occurred 1970–71 in the Ivory Coast, Liberia, Nigeria, and Sierra Leone.^{32,33,34,35} Subsequent enhanced surveillance observed a steady increase in the rate of human monkeypox cases. The number of human monkeypox cases reported has risen exponentially over the past 20 years, and has already exceeded that accumulated during the first 45 years since its first discovery.^{28,29,36,37,38,39,40,41,42,43,44,45}

A comprehensive enhanced surveillance study in the Democratic Republic of Congo (DRC) in 2004–05 showed a steep increase in incidence compared to data from a WHO enhanced

surveillance program carried out 1970–86 initiated by WHO reporting 404 cases.⁴⁶ The incidence was highest in forested regions, and in lower age groups not vaccinated as part of the smallpox eradication program.⁴² To date, human monkeypox cases have been reported from ten African countries – Democratic Republic of the Congo, Republic of the Congo, Cameroon, Central African Republic, Nigeria, Ivory Coast, Liberia, Sierra Leone, Gabon and South Sudan.^{1,36,47} The growing incidence of human monkeypox cases in Central and West Africa is considered a consequence of waning cross-protective immunity among the population after smallpox vaccination was discontinued in the early 1980s, following the eradication of smallpox.^{28,42} The deteriorating immunological status is not only related to waning vaccine-induced protection among those initially vaccinated, but probably – and even more – to the increasing proportion of those never given the vaccine, i.e. non-vaccinated younger age groups. Both mechanisms lead to a growing percentage of susceptible individuals in the endemic areas in Central and West Africa. Another central factor considered to contribute to the incidence of monkeypox is related to increasing contact between humans and small mammals potentially carrying MPXV. Humans invade jungles and forests, the natural environment of the reservoir species. Civil wars, refugee displacement, farming, deforestation, climate change, demographic changes and population movement may have led to a spread of monkeypox-infected animals and increased their interaction with humans across West and Central Africa.

Monkeypox cases in USA

Monkeypox remained an ignored global public health threat and was only given international attention, when the first cases outside Africa were detected in the United States in 2003.⁴⁸ After several Midwesterners developed fever, rash, respiratory symptoms, and lymphadenopathy, outbreak investigation linked the symptoms to exposure to pet prairie dogs (*Cynomys* species), and monkeypox virus was identified as a causative agent.⁴⁸ It spread rapidly. Monkeypox cases were reported from six states — Illinois, Indiana, Kansas, Missouri, Ohio, and Wisconsin – during the outbreak.⁴⁹ Molecular investigations identified a monkeypox virus of the West African genetic group (clade). Epidemiological studies concluded that the virus had been imported into the US, more specifically to Texas, from Ghana on April 9, 2003, with a shipment of small mammals of nine different species, including six genera of African rodents.⁵⁰ These comprised rope squirrels (*Funisciurus* sp.), tree squirrels (*Heliosciurus* sp.), African giant

pouched rats (*Cricetomys* sp.), brush-tailed porcupines (*Atherurus* sp.), dormice (*Graphiurus* sp.), and striped mice (*Lemniscomys* sp.). Some of the infected animals were housed in close proximity to prairie dogs later sold as pets.

Monkeypox cases in the United Kingdom and Israel

In September 2018, monkeypox again drew the attention of global media, politicians and scientists, when three individual patients in the United Kingdom were diagnosed with monkeypox.⁵¹ The first two had recently traveled in Nigeria, a country with an ongoing outbreak of the disease,⁵² and both were symptomatic during their flight home. The third case of monkeypox in the United Kingdom was diagnosed in a healthcare worker caring for one of the first two. As the clinical picture of the three patients' disease raised a concern over an exotic disease, special infection control measures were taken well before monkeypox was suspected. One of the primary cases reported contact with a person with suspected rash at a family gathering, and the consumption of bush meat.⁵² Secondary and tertiary human-to-human transmission of monkeypox do occur in endemic areas.^{53,54} Of note, definitive confirmation of human-to-human transmission in endemic areas is somewhat problematic, because even the secondary and tertiary cases may have been exposed to infected animals. The disease contracted by the British health care worker provides indisputable evidence of human-to-human transmission from an infected patient.

In October 2018, Israel reported a monkeypox case imported from Nigeria.⁵⁵ It is well known that travelers can act as sentinels of infectious diseases epidemics in the region visited. Not consistent with the reports of low levels of transmission in Nigeria, three cases imported to other countries from there within a couple of months should raise health authorities' concern.⁵⁶

Ongoing monkeypox outbreak in West Africa

On September 22, 2017, the Nigeria Centre for Disease Control (NCDC) commenced an outbreak investigation following the identification of a suspected case of monkeypox in an 11-year old child.⁵⁷ The data available indicate that the current outbreak is either a multisource outbreak or one stemming from previously undetected endemic transmission, since the cases were not epidemiologically linked.^{28,58,59,60} The exact zoonotic origin and role of environmental and ecological factors in the Nigerian outbreak are not yet known. New cases of monkeypox

continue to be detected in the country. Since the beginning of the outbreak on September 22nd 2018, as of 1st January 2019, there have been 311 suspected cases reported from 26 states (132 confirmed cases affecting children and adults of all ages) and 7 deaths reported.⁶⁰ The majority of confirmed monkeypox cases are between 21 and 40 years, with a median age of 31 years. The majority of confirmed Monkeypox cases are under 40 years with a median age of 31 years, similar to the observed age range in DRC.⁴² It is noteworthy that all were born after 1978, when the global vaccination programs for smallpox were discontinued.

MODES OF TRANSMISSION OF MPXV TO HUMANS

The exact mode of MPXV transmission to humans remains unknown. Primary animal-to-human infection is assumed to occur when handling monkeypox-infected animals, through direct (touch, bite or scratch) or indirect contact, although the exact mechanism(s) remains to be defined. The virus is assumed to enter the body through broken skin, respiratory tract, or the mucous membranes (eyes, nose, mouth). Secondary human-to-human transmission is considered common,^{37,53,61} presumably through large respiratory droplets or direct or indirect contact with body fluids, lesion material, and contaminated surfaces or other material, such as clothing or linens. Prolonged contact with patients render hospital staff and family members at greater risk of infection. Nosocomial transmission has been described.³⁸ There is no evidence, to date, that human-to-human transmission alone can sustain monkeypox infections in the human population.

There have only been a few genomic studies of the origins of monkeypox outbreaks. Human-to-human transmission has been described from primary human cases, secondary cases,^{53,62,63} and serial transmission across four cases has been observed.⁶⁴ In the current monkeypox outbreak in Nigeria, genomic studies of monkeypox virus isolates from humans,⁶⁰ indicate that the index case was not imported into Nigeria. Thus, the outbreak is considered to be a spillover from multiple sources of introduction into the human population. The zoonotic source(s) of the outbreak are being investigated at present, and it is unclear what, if any, environmental or ecologic changes might have facilitated the sudden reemergence of monkeypox in Nigeria. Case clustering has been identified within the various states, but no epidemiological linkages

between them have been detected thus far. Three family clusters have been identified, suggesting human-to-human transmission.^{58,59,60} In one family the secondary attack rate was 71%. However, most patients had no obvious epidemiologic linkage or person-to-person contact, indicating a probable multiple-source outbreak or, possibly, an endemic disease previously unrecognized.

CLINICAL FEATURES

CLINICAL FEATURES

The incubation period has been estimated at 5 to 21 days, and duration of symptoms and signs at 2 to 5 weeks. The illness begins with non-specific symptoms and signs that include fever, chills, headaches, lethargy, asthenia, lymph node swellings, back pain, and myalgia (muscle ache) and begins with a fever before skin rashes appear. Within 1 to 5 days after the onset of fever, skin rashes of varying sizes appear, first on the face (**Figs. 2A-D**), then across the body (**Figs. 3A-3D**), hands (**Fig. 4A**) and legs and feet (**Fig.4A-B**). The rash undergoes several stages of evolution from macules, papules, vesicles (fluid-filled blisters) (**Figs.2A-D**) and pustules (**Fig.3B, Fig 3D**), followed by resolution over time with crusts and scabs (**Fig.5**), which drop off on recovery. Various stages of the rash may show at the same time (**Fig.3B, Fig.5**). Areas of erythema (**Fig.2A**) and/or skin hyperpigmentation (**Fig.5**) are often seen around discrete lesions. Detached scabs may be considerably smaller than the original lesion. Inflammation of the pharyngeal, conjunctival, and genital mucosae may also be seen.

The clinical presentation of monkeypox includes symptoms and lesions that are difficult to distinguish from smallpox.^{37,60,65,66} Whilst the clinical manifestations of monkeypox are milder than smallpox, the disease can prove fatal, death rates ranging from 1% to 10%. Mortality is higher among children and young adults and the course more severe in immunocompromised individuals.⁶⁷ A range of complications have been reported, such as secondary bacterial

infections, respiratory distress, bronchopneumonia, encephalitis, and corneal infection with ensuing loss of vision, gastrointestinal involvement, vomiting and diarrhea with dehydration. Case fatality rates have varied between 1% and 10% in outbreaks, deaths occurring mostly among young adults and children. Particularly those with immunosuppression are at risk of severe disease. Lymphadenopathy is seen in up to 90% of patients that appears to be a clinical feature distinguishing human monkeypox from smallpox.

Previous smallpox vaccination confers some cross-protection against monkeypox and modifies the clinical picture towards a milder disease. Between 1980 and 1990, the clinical presentation of human monkeypox appears to have changed: primary human cases have increasingly been seen among those never vaccinated against smallpox. Compared to those vaccinated, the clinical picture described for the unvaccinated was more severe, with more vigorous and pleomorphic skin rashes, and higher mortality.^{62,66,68,69,70}

The primary differential diagnosis is severe chickenpox with lesions in palms and soles.^{7,65} The lesions in chickenpox are more superficial, occur in clusters of the same stage, with denser manifestations on the trunk than on face and extremities. Due to the non-specific nature of the symptoms and signs of monkeypox, a wide variety of differential diagnoses should be considered, ranging from chickenpox, molluscum contagiosum, measles, rickettsial infections, bacterial skin infections (such as those caused by *Staphylococcus aureus*), anthrax, scabies, syphilis, and drug reactions to other non-infectious causes of skin rash. A clinical sign differentiating monkeypox from smallpox and chickenpox is enlarged lymph nodes, particularly submental, submandibular, cervical, and inguinal nodes.⁷¹

SMALLPOX VACCINATION, MONKEYPOX PREVALENCE AND CHANGING CLINICAL PRESENTATIONS

In 1980, the Global Commission for the Certification of Smallpox Eradication (GCCSE) continued to designate monkeypox as a public health threat, recommending that the epidemiological, ecological and surveillance program on monkeypox be continued.^{35,72} In response, WHO supported an active surveillance program for human monkeypox from 1970 to 1986.⁴⁶ It was assumed to be endemic to DRC, but other Central and West African countries also reported cases of monkeypox in humans or circulation in wildlife. At the end of the smallpox eradication

campaign, the GCCSE stated that smallpox vaccination to prevent monkeypox was no longer justified, even if cross-protective immunity could not be relied on for long due to the vaccinations being discontinued. In retrospect, this resolution may have been an error.

Experimental studies of monkeys have shown immunization with smallpox vaccine to give cross-protection against monkeypox.¹⁹

Several reviews have summarized human monkeypox outbreaks over the past 38 years.^{28,29,42,73} Between November 2005 and November 2007, population-based surveillance studies conducted in nine health zones in central DRC identified 760 laboratory-confirmed human monkeypox cases. The average annual cumulative incidence across the zones was 5.53 per 10,000 (2.18–14.42). Factors associated with increased risk of infection included: living in forested areas, male gender, age < 15years, and absence of smallpox vaccination scar. Among those vaccinated, the risk of monkeypox was found to be 5.2-fold lower than among those unvaccinated (0.78 versus 4.05 per 10,000). Compared to surveillance data from the same region recorded in the 1980s, a 20-fold increase in human monkeypox incidence was observed. Between January 2001 and December 2004, the DRC Ministry of Health surveillance program reported 2,734 cases of suspected human monkeypox in 11 provinces, showing annual upward trends: 380 cases in 2001, 545 in 2002, 783 in 2003, and 1026 in 2004. The majority of cases (94%) were seen among children and young adults under 25 years of age.⁴¹ These patients had not been vaccinated against smallpox. Surveillance activities have been halted since 2005 because of the civil war.

DIAGNOSIS - LABORATORY, VIROLOGICAL, AND HISTOLOGICAL FEATURES

Optimal clinical specimens for laboratory analyses include specimens from skin lesions such as swabs of vesicular lesions, exudate or crusts stored in a dry, sterile tube (no viral transport media) and kept cold. A viral culture should be obtained by an oropharyngeal or nasopharyngeal swab. Skin biopsies of vesiculopustular rash or a sample of the roof of an intact skin vesicular lesion are valuable for analyzes. Reference laboratories with high containment facilities are required to make a definitive diagnosis using electron microscopy, culture and molecular analysis identification by PCR and sequencing. Serological testing requires paired

acute and convalescent sera for MPXV-specific IgM detection within 5 days of presentation, or IgG detection after 8 days.

Histology and immunohistochemistry of papular lesions may show acanthosis, individual keratinocyte necrosis, and basal vacuolization, along with a superficial and deep perivascular, lymphohistiocytic infiltrate in the dermis. Vesicular lesions show spongiosis with reticular and ballooning degeneration, multinucleated epithelial giant cells with epidermal necrosis with numerous eosinophils and neutrophils, and features of vasculitis and viral inclusions in keratinocytes. Intracytoplasmic, round-to-oval inclusions with sausage-shaped structures centrally, measuring 200-300 μm may be seen on electron microscopic observation.

TREATMENT

There is no specific treatment for monkeypox. Supportive care, symptomatic management, and treatment of secondary bacterial infections remain the main recommendations.

PREVENTION

Prevention of MPXV spread in endemic areas is highly challenging. It consists in avoiding any contact with rodents and primates as well as limiting direct exposure to blood and inadequately cooked meat. Efforts to halt bushmeat trade and consumption of wild animals are extremely difficult both culturally and economically, as this meat may be the only protein source available for the poorest people. Massive health education campaigns are needed to increase general awareness and to advise on proper handling of potential animal reservoir species (gloves, protective clothing, surgical mask) as well as avoiding close contact with anyone infected.

Infection control measures are vital to prevention of human-to-human transmission in health care. Improved nursing (gloves, protective clothing, surgical masks) and isolation practices require education as well as adequate facilities and staffing.

National health authorities should consider arranging immunization against smallpox for health care workers and those treating or exposed to patients with monkeypox or their samples.

Smallpox vaccination has been estimated to provide 85% cross-protection against monkeypox infection.³² The Centers for Disease Control and Prevention (CDC) recommended smallpox vaccination within 2 weeks, ideally before 4 days, after significant, unprotected exposure to diseased animal or a confirmed human case.

During an outbreak, the spread of monkeypox virus may be controlled by quarantining (at least for 6 weeks from the date of last exposure) the infected animals and tracing their contacts. Adherence to specific instructions from the local and global public health authorities be consulted. Increasing awareness and action (adequate decisions, medical staff, sampling, surveillance, education) both by local and international authorities are of central importance.

At hospitals in developed countries, when suspecting a case of monkeypox (e.g. patient with fever, skin lesions, and history of visiting endemic area or contact with patients) the patient should immediately be placed in a negative air pressure room, or a private room, if such facilities are unavailable. Standard, contact and droplet precautions should all be taken. Infection control personnel should be contacted without delay. In developed countries, likewise, increasing awareness among health care personnel about the disease and its endemic areas is an important precaution.

VACCINES AGAINST MONKEYPOX

Whilst new vaccines are being developed for monkeypox, there is a need for conducting controlled clinical trials to evaluate the impact of the use of smallpox vaccines for prevention of monkeypox or modifying disease severity. Studies should focus on the cost benefit of population level vaccination and investigation of alternative vaccination strategies such as targeting vaccination to affected areas, contacts and healthcare workers, and wider geographic areas. Currently, the CDC recommends pre-exposure smallpox vaccination for field investigators, veterinarians, animal control personnel, contacts of monkeypox patients, researchers and health-care workers caring for such patients and their contacts.³

Can the smallpox vaccine available be used to protect against monkeypox?

Percutaneous inoculation with VACV elicits a broad and heterogeneous serum antibody response targeting a large number of antigenic determinants of VACV.^{74,75} The viral inhibitory

activity of serum from immune subjects with cross-neutralizing activity to VACV, MPXV, and VARV is presumably composed of antibodies with diverse specificities.^{76,77,78}

Production of first generation live, attenuated vaccine vaccines has been reviewed by the WHO in 1988.⁹ A considerable proportion of the population may have contraindications for the vaccine candidates: 15.2–15.8% of the U.S. population have been estimated to have potential contraindications for taking the live, attenuated smallpox vaccine.⁷⁹

The rate of side effects associated with the live, attenuated vaccinia virus in the US in 1968 were 74 complications and one death per 1,000,000 primary vaccinations. Morbidity and mortality rates were highest for infants, with 112 complications and five deaths per 1,000,000 primary vaccinations.⁸⁰ In 2002, the U.S. Department of Defense resumed a program for widespread smallpox vaccinations due to a perceived threat of biological warfare. A total of 540,824 military personnel were vaccinated with a New York City Board of Health strain of vaccinia, “DryVax” from December 2002 through December 2003. Dryvax was produced by infection of skin of calves using the NYCBH strain as seed virus. Of these, 67 (1/8000) developed myopericarditis.^{81,82}

The highest rate of post-vaccine encephalitis, pvE, was found with the Bern strain (44.9 expected cases per million vaccines), followed by the Copenhagen strain (33.3 per million vaccines), the Lister strain (26.2 per million vaccines), and the NYCBH strain with the lowest rate (2.9 per million vaccinations).⁸³

ADDRESSING GAPS IN KNOWLEDGE AND STRENGTHENING PUBLIC HEALTH PREPAREDNESS

Most data available on monkeypox are obtained from individual case or outbreak reports, and from passive intermittent surveillance, none of which convey an accurate overall picture. The current major gaps in monkeypox knowledge, the changing epidemiological and clinical presentations, and the multifarious factors involved in monkeypox transmission argue the need to strengthen outbreak preparedness efforts. There remains an urgent need for developing public health and surveillance capacities in Central and West Africa to guide appropriate surveillance, data collection, prevention, preparedness and response activities to monkeypox and other emerging and re-emerging infections with epidemic potential. Advancing public

health preparedness and aligning proactive surveillance activities to priority research will require coordinated, locally led, multidisciplinary efforts adjusted closely to capacity development and training.

CONCLUSIONS

The spread of Monkeypox across West Africa over the past decade and the ongoing outbreak in Nigeria Monkeypox indicates it is no longer “a rare viral zoonotic disease that occurs primarily in remote parts of Central and West Africa, near tropical rainforests”. It’s potential for further spread both regionally and internationally remains a major concern.^{28,29} The ecological, zoonotic, epidemiological, clinical and public health aspects of monkeypox remain inadequately characterized.^{33,36,44,45} The first-generation live, attenuated vaccinia virus vaccines stored for emergency purposes in many countries cannot be used due to severe adverse reactions. Discontinuing the smallpox vaccination program has created an ecological gap where an increasing proportion of the population has either waning or non-existent immunity to MPXV. This development will further both the animal-to-human and human-to-human spread of the virus. Therefore, priority research and surveillance should urgently be conducted through a joint ‘One-Human-Animal-Environmental Health’ effort across Central and West Africa.^{84,85,86}

ACKNOWLEDGEMENTS

We thank the Nigeria CDC staff for providing the clinical photographs from the ongoing 2019 Monkeypox outbreak in Nigeria and patients who gave permission to have the photos taken.

AZ, DA and CI are members of the PANDORA-ID-NET Consortium. PANDORA-ID-NET, (EDCTP Reg/Grant RIA2016E-1609) funded by the European and Developing Countries Clinical Trials Partnership (EDCTP2) programme which is supported under Horizon 2020, the European Union's Framework Programme for Research and Innovation. AZ is in receipt of an National Institutes of Health Research (NIHR) senior investigator award.

CONFLICTS OF INTEREST

All authors have an interest in global public health and emerging and re-emerging infections. All authors have no other conflict of interest to declare.

REFERENCES

1. WHO.2019. Human Monkeypox. <https://www.who.int/emergencies/diseases/monkeypox/en/> (Accessed 22 Febr. 2019)
2. WHO 2019. Smallpox. <https://www.who.int/biologicals/vaccines/smallpox/en/> (accessed February 20th 2019)
3. CDC. Monkeypox. MMWR Morb Mortal Wkly Rep. 2018;67:306–10.
4. Shchelkunov SN, Totmenin AV, Babkin IV et al. Human monkeypox and smallpox viruses: genomic comparison. FEBS Lett 2001;509:66–70.
5. Pauli G, Blümel J, Burger R et al. Orthopox Viruses: Infections in Humans. Transfus Med Hemother. 2010;37: 351–64.
6. Bera BC, Shanmugasundaram K, Barua S et al. Zoonotic cases of camelpox infection in India. Vet Microbiol. 2011;152:29-38.
7. Cook GC, Zumla A. 2009. Chapter 47. Cutaneous viral diseases. Monkeypox. Pages 839-840. In Manson's Tropical Diseases. Editors: Cook GC, Zumla A. 22nd Edition.2009. Harcourt Brace Saunders, Publishing Group, London, UK.
8. Petersen E, Abubakar I, Ihekweazu C et al. Monkeypox - Enhancing public health preparedness for an emerging lethal human zoonotic epidemic threat in the wake of the smallpox post-eradication era. Int J Infect Dis. 2019;78:78-84.
9. Fenner F, Henderson DA, Arita I, Jezek Z, Ladnyi ID. Smallpox and its eradication. Geneva: World Health Organisation, Geneva 1988.
10. Singh RK, Hosamani M, Balamurugan V, Bhanuprakash V, Rasool TJ, Yadav MP. Buffalopox: an emerging and re-emerging zoonosis. Anim Health Res Rev. 2007;8:105-14.
11. Oliveira JS, Figueiredo PO, Costa GB, Assis FL, Drumond BP, da Fonseca FG, Nogueira ML, Kroon EG, Trindade GS. Vaccinia Virus Natural Infections in Brazil: The Good, the Bad, and the Ugly. Viruses. 2017 Nov 15;9(11). pii: E340.

12. Hughes AL, Irausquin S, Friedman R. The evolutionary biology of poxviruses. *Infect Genet Evol.* 2010;10:50–59.
13. Ichihashi Y, Oie M. Epitope mosaic on the surface proteins of orthopoxviruses. *Virology.* 1988;163:133–144.
14. Stanford MM, McFadden G, Karupiah G, Chaudhri G. Immunopathogenesis of poxvirus infections: forecasting the impending storm. *Immunol Cell Biol.* 2007;85:93-102.
15. McConnell S, Herman YF, Mattson DE, Huxsoll DL, Lang CM, Yager RH. Protection of Rhesus Monkeys against Monkeypox by Vaccinia Virus Immunization. *Am J Vet Res.* 1964; 25:192–5.
16. Hammarlund E, Lewis MW, Carter SV et al. Multiple diagnostic techniques identify previously vaccinated individuals with protective immunity against monkeypox. *Nat Med.* 2005;11:1005–11.
17. Moss B. Smallpox vaccines: targets of protective immunity. *Immunol Rev.* 2011;239:8–26.
18. Gispen R, Verlinde JD, Zwart P. Histopathological and virological studies on monkeypox. *Arch ges Virusforsch* 1967;21:205-216.
19. McConnell S, Hickman RL, Wooding WL, Jr., Huxsoll DL. Monkeypox: experimental infection in chimpanzee (*Pan satyrus*) and immunization with vaccinia virus. *Am J Vet Res* 1968;29:1675–80.
20. von Magnus P, Anderson EK, Petersen KB, Birch-Anderson A. A pox-like disease in cynomolgus monkeys. *Acta Pathol Microbiol Scand* 1959;46:156–76.
21. Doty JB, Malekani JM, Kalemba LN, Stanley WT, Monroe BP, Nakazawa YU, et al. Assessing Monkeypox Virus Prevalence in Small Mammals at the Human-Animal Interface in the Democratic Republic of the Congo. *Viruses.* 2017 Oct 3;9(10). pii: E283.
22. Radonić A, Metzger S, Dabrowski PW et al . Fatal monkeypox in wild-living sooty mangabey, Côte d'Ivoire, 2012. *Emerg Infect Dis.* 2014;20:1009-11.
23. Hutson CL, Lee KN, Abel J et al. Monkeypox zoonotic associations: insights from laboratory evaluation of animals associated with the multi-state US outbreak. *Am J Trop Med Hyg.* 2007;76:757-68.
24. Thomassen HA, Fuller T, Asefi-Najafabady S et al. Pathogen-host associations and predicted range shifts of human monkeypox in response to climate change in central Africa. *PLoS One.* 2013;8:e66071.
25. Prier JE, Sauer RM. A pox disease of monkeys. *Ann N Y Acad Sci* 1960;85:951-9.
26. Wenner HA, Macasaet D, Kamitsuka PS, Kidd P. Monkeypox. Clinical, virologic and immunologic studies. *Am J Epidemiol* 1968;87:551-66.

27. Nakoune E, Lampaert E, Ndjapou SG et al. A Nosocomial Outbreak of Human Monkeypox in the Central African Republic. *Open Forum Infect Dis.* 2017;4:ofx168.
28. Durski KN, McCollum AM, Nakazawa Y et al. Emergence of Monkeypox - West and Central Africa, 1970-2017. *MMWR Morb Mortal Wkly Rep.* 2018;67:306-10.
29. Sklenovská N, Van Ranst M. Emergence of Monkeypox as the Most Important Orthopoxvirus Infection in Humans. *Front Publ Hlth.* 2018;6:241.
30. Quiner CA, Moses C, Monroe BP et al. Presumptive risk factors for monkeypox in rural communities in the Democratic Republic of the Congo. *PLoS One.* 2017;12:e0168664.
31. Marennikova SS, Seluhina EM, Malceva NN, Cimiskjan KL, Macevic GR. Isolation and properties of the causal agent of a new variola-like disease (monkeypox) in man. *Bull World Hlth Organ* 1972;46:599–611.
32. Fine PE, Jezek Z, Grab B, Dixon H. The transmission potential of monkeypox virus in human populations. *Int J Epidemiol* 1988; 17: 643–50
33. Heymann DL, Szczeniowski M, Esteves K. Reemergence of monkeypox in Africa: a review of the past six years. *Br Med Bull* 1998; 54: 693–702.
34. Breman JG, Kalisa R, Steniowski MV, Zanotto E, Gromyko AI, Arita I. Human monkeypox, 1970–79. *Bull World Health Organ* 1980;58:165–82.
35. WHO.1980 The global eradication of smallpox: final report of the Global Commission for the Certification of Smallpox Eradication. World Health Organization, Geneva 1980.
36. WHO 2018. Human Monkeypox (MPX) <http://www.who.int/emergencies/diseases/monkeypox/en/> (Accessed 22nd February 2019).
37. Hutin YJ, Williams RJ, Malfait P et al. Outbreak of human monkeypox, Democratic Republic of Congo, 1996 to 1997. *Emerg Infect Dis* 2001;7:434.
38. Learned LA, Reynolds MG, Wassa DW, Li Y, Olson VA, Karem K, et al. Extended interhuman transmission of monkeypox in a hospital community in the Republic of the Congo, 2003. *Am J Trop Med Hyg.* 2005;73:428–34.
39. Reynolds MG, Emerson GL, Pukuta E et al. Detection of human monkeypox in the Republic of the Congo following intensive community education. *Am J Trop Med Hyg.* 2013;88:982–5.
40. Khodakevich L, Widy-Wirski R, Arita I et al. Monkey pox virus infection in humans in the Central African Republic. *Bull Soc Pathol Exot Filiales.*1985;78:311–20.

41. Rimoin AW, Kialu N, Kebela-Ilunga B, Mukaba T, Wright LL, Formenty P, et al. Endemic human monkeypox, Democratic Republic of Congo, 2001-2004. *Emerg Infect Dis.* 2007;13:934-7.
42. Rimoin AW, Mulembakani PM, Johnston SC, Lloyd Smith JO, Kialu NK, Kinkela TL, et al. Major increase in human monkeypox incidence 30 years after smallpox vaccination campaigns cease in the Democratic Republic of Congo. *Proc Natl Acad Sci U S A.* 2010;107:16262-7.
43. Kantele A, Chickering K, Vapalahti O, Rimoin AW. Emerging diseases-the monkeypox epidemic in the Democratic Republic of the Congo. *Clin Microbiol Infect.* 2016;22:658-9.
44. Hoff NA, Doshi RH, Colwell B et al. Evolution of a Disease Surveillance System: An Increase in Reporting of Human Monkeypox Disease in the Democratic Republic of the Congo, 2001-2013. *Int J Trop Dis Hlth.* 2017;25.pii: IJTDH.35885. doi:
45. Yinka-Ogunleye A, Aruna O, Ogoina D et al. Reemergence of Human Monkeypox in Nigeria, 2017. *Emerg Infect Dis.* 2018;24:1149-51.
46. Jezek Z, Fenner F. Human Monkeypox. *Monographs in Virology* vol. 17. Karger, Basel 1988. Tabel 7, pp. 49.
47. Formenty P, Muntasir MO, Damon I, Chowdhary V, Opoka ML, Monimart C et al . Human monkeypox outbreak cause by novel virus belonging to Congo Basin clade, Sudan, 2005. *Emerg Infect Dis.*2010;16:1539-45.
48. CDC, Centers for Disease Control and Prevention. Update: multistate outbreak of monkeypox--Illinois, Indiana, Kansas, Missouri, Ohio, and Wisconsin, 2003. *MMWR Morb Mortal Wkly Rep* 2003; 52:642.
49. CDC, Centers for Disease Control and Prevention, update: multistate outbreak of monkeypox--Illinois, Indiana, Kansas, Missouri, Ohio, and Wisconsin, 2003. *MMWR Morb Mortal Wkly Rep* 2003; 52: 589-90.
50. CDC, Centers for Disease Control and Prevention. Monkeypox in the United states. 2003 Outbreak. <https://www.cdc.gov/poxvirus/monkeypox/outbreak.html> (Accessed 22nd February 2019)
51. Public Health England 2018. Monkeypox -Latest update <https://www.gov.uk/government/news/monkeypox-case-in-england -accessed October 29th 2018> (Accessed 22nd February 2019).
52. Vaughan A, Aarons E, Astbury J, Balasegaram S, Beadsworth M, Beck CR, et al. Two cases of monkeypox imported to the United Kingdom, September 2018. *Euro Surveill.* 2018;23(38). doi: 10.2807/1560-7917.

53. Jezek Z, Arita I, Mutombo M, Dunn C, Nakano JH, Szczeniowski M. Four generations of probable person-to-person transmission of human monkeypox. *Am J Epidemiol* 1986; 123:1004-12.
54. Kalthan E, Tenguer J, Ndjapou SG, Koyazengbe TA, Mbomba J, Marada RM, et al. Investigation of an outbreak of monkeypox in an area occupied by armed groups, Central African Republic. *Med Mal Infect*. 2018;48:263-8.
55. Ministry of Health, State of Israel. Monkeypox Patient Diagnosed. https://www.health.gov.il/English/News_and_Events/Spokespersons_Messages/Pages/12102018_1.aspx (Accessed 19 February 2019).
56. WHO. Human Monkeypox in Nigeria – update 5 October 2018. <https://www.who.int/csr/don/05-october-2018-monkeypox-nigeria/en/> (Accessed 22nd February 2019).
57. Eteng WE, Mandra A, Doty J et al. Notes from the Field: Responding to an Outbreak of Monkeypox Using the One Health Approach - Nigeria, 2017-2018. *MMWR Morb Mortal Wkly Rep*. 2018;67:1040-1.
58. Nigeria CDC. 2018. An Update of Monkeypox Outbreak in Nigeria. <https://ncdc.gov.ng/diseases/sitreps/?cat=8&name=An%20Update%20of%20Monkeypox%20Outbreak%20in%20Nigeria> (Accessed 22nd February 2019)
59. Nigeria CDC 2019. Monkeypox. <https://ncdc.gov.ng/diseases/sitreps/?cat=8&name=An%20Update%20of%20Monkeypox%20Outbreak%20in%20Nigeria>. (Accessed 22nd February 2019)
60. Faye O, Pratt CB, Faye M et al. Genomic characterisation of human monkeypox virus in Nigeria. *Lancet Infect Dis*. 2018;18:246.
61. Jezek Z, Grab B, Szczeniowski MV, Paluku KM, Mutombo M. Human monkeypox: secondary attack rates. *Bull World Health Organ* 1988;66:465–70.
62. Jezek Z, Grab B, Dixon H. Stochastic model for interhuman spread of monkeypox. *Am J Epidemiol* 1987;126:1082-92.
63. Jezek Z, Szczeniowski M, Paluku KM, Mutombo M. Human monkeypox: clinical features of 282 patients. *J Infect Dis* 1987;156:293–98.
64. Nolen LD, Osadebe L, Katomba J, et al. Extended human-to-human transmission during a monkeypox outbreak in the Democratic Republic of the Congo. *Emerg Infect Dis*. 2016;22:1014–21.
65. Jezek Z, Szczeniowski M, Paluku KM, Mutombo M, Grab B. Human monkeypox: confusion with chickenpox. *Acta Trop* 1988;45:297–307.

66. Di Giulio DB, Eckburg PB. Human monkeypox: an emerging zoonosis. *Lancet* 2004;4:15-25.
67. Gordon SN, Cecchinato V, Andresen V et al. Smallpox vaccine safety is dependent on T cells and not B cells. *J Infect Dis.* 2011;203:1043-53.
68. Huhn GD, Bauer AM, Yorita K et al. Clinical characteristics of human monkeypox, and risk factors for severe disease. *Clin Infect Dis.* 2005;41:1742–51.
69. Damon IK. Status of human monkeypox: clinical disease, epidemiology and research. *Vaccine* 2011;29(Suppl 4):D54–9.
70. McCollum AM, Damon IK. Human monkeypox. *Clin Infect Dis.* 2014;58:260–7.
71. Osadebe L, Hughes CM, Shongo Lushima R et al. Enhancing case definitions for surveillance of human monkeypox in the Democratic Republic of Congo. *PLoS Negl Trop Dis.* 2017;11:e0005857.
72. WHO 1984. The current status of human monkeypox: Memorandum from a WHO Meeting. *Bull Wld Hlth Organ.* 1984;62:703–13.
73. Durski KN, McCollum AM, Nakazawa Y et al. Emergence of Monkeypox — West and Central Africa, 1970–2017. *MMWR Morb Mortal Wkly Rep.* 2018; 67:306–10.
74. Davies DH, Liang X, Hernandez JE et al. Profiling the humoral immune response to infection by using proteome microarrays: high-throughput vaccine and diagnostic antigen discovery. *Proc Natl Acad Sci U S A.* 2005;102:547–52.
75. Davies DH, Molina DM, Wrammert et al. Proteome-wide analysis of the serological response to vaccinia and smallpox. *Proteomics.* 2007;7:1678–86
76. Hughes CM, Newman FK, Davidson WB et al. Analysis of variola and vaccinia virus neutralization assays for smallpox vaccines. *Clin Vaccine Immunol.* 2012;19:1116-8.
77. Kennedy JS, Gurwith M, Dekker CL et al. Safety and immunogenicity of LC16m8, an attenuated smallpox vaccine in vaccinia-naive adults. *J Infect Dis.* 2011;204:1395-402.
78. Gilchuk I, Gilchuk P, Sapparapu G et al. Cross-Neutralizing and Protective Human Antibody Specificities to Poxvirus Infections. *Cell.* 2016;167:684-94.
79. Carlin EP, Giller N, Katz R. Estimating the Size of the U.S. Population at Risk of Severe Adverse Events from Replicating Smallpox Vaccine. *Public Health Nurs.* 2017;34:200-9.
80. Lane JM, Ruben FL, Neff JM, Millar JD. Complications of smallpox vaccination, 1968. *N Engl J Med* 1969;281:1201–8.
81. Eckart RE, Love SS, Atwood JE et al. Department of Defense Smallpox Vaccination Clinical Evaluation Team. Incidence and follow-up of inflammatory cardiac complications after smallpox vaccination. *J Am Coll Cardiol.* 2004;44:201-5

82. Neff J, Modlin J, Birkhead GS et al. and the Advisory Committee on Immunization. Practices; Armed Forces Epidemiological Board (2008). Monitoring the safety of a smallpox vaccination program in the United States: report of the joint Smallpox Vaccine Safety Working Group of the advisory committee on immunization practices and the Armed Forces Epidemiological Board. *Clin Infect Dis.* 46(Suppl 3):258-70.
83. Kretzschmar M, Walinga J, Teunis P, Xing S, Mikolajczyk R. Frequency of adverse events after vaccination with different vaccinia strains. *PLoS Med.* 2006 Aug;3(8):e272.
84. Bass J, Tack DM, McCollum AM et al. Enhancing health care worker ability to detect and care for patients with monkeypox in the Democratic Republic of the Congo. *Int Health.* 2013;5:237–43.
85. Zumla A, Dar O, Kock R et al. Taking forward a 'One Health' approach for turning the tide against the Middle East respiratory syndrome coronavirus and other zoonotic pathogens with epidemic potential. *Int J Infect Dis.* 2016;47:5-9.
86. Doshi RH, Guagliardo SAJ, Dzabatou-Babeaux A et al. Strengthening of Surveillance during Monkeypox Outbreak, Republic of the Congo, 2017. *Emerg Infect Dis.* 2018;24:1158-60.