

Modeling anxiety in healthy humans: A key intermediate bridge between basic and clinical sciences

Christian Grillon¹, Oliver J. Robinson², Brian Cornwell³, Monique Ernst¹

¹Section on the Neurobiology of Fear and Anxiety, National Institute of Mental Health, Bethesda, MD, USA.

² University College London, Institute of Cognitive Neuroscience, London, UK

³ Centre for Mental Health, Faculty of Health, Arts and Design, Swinburne University of Technology, Hawthorn, Australia

Corresponding author contact information:

Christian Grillon

Building 15K, Room 203

National Institute of Mental Health

Bethesda, MD 20814

phone: (310) 594 2894

christian.grillon@nih.gov

ABSTRACT (150-250 WORDS)

Animal models of anxiety disorders are important for elucidating neurobiological defense mechanisms. However, animal models are limited when it comes to understanding the more complex processes of anxiety that are unique to humans (e.g., worry) and to screen new treatments. In this review we outline how the *Experimental Psychopathology* approach, based on experimental models of anxiety in healthy subjects, can mitigate these limitations and complement research in animals. Experimental psychopathology can bridge basic research in animals and clinical studies, as well as guide and constrain hypotheses about the nature of psychopathology, treatment mechanisms and treatment targets. This review begins with a brief review of the strengths and limitation of animal models before discussing the need for human models of anxiety, which are especially necessary to probe higher-order cognitive processes. This can be accomplished by combining anxiety induction procedures with tasks that probe clinically-relevant processes to identify neurocircuits that are potentially altered by anxiety. The review then discusses the validity of experimental psychopathology and introduces a methodological approach consisting of five steps: 1) select anxiety-relevant cognitive or behavioral operations and associated tasks, 2) identify underlying neurocircuits supporting these operations in healthy controls, 3) examine the impact of experimental anxiety on the targeted operations in healthy control, 4) utilize findings from step 3 to generate hypotheses about neurocircuit dysfunction in anxious patients, and 5) evaluate treatment mechanisms and screen novel treatments. This is followed by two concrete illustrations of this approach and suggestions for future studies.

(now 7367 words; before 3505)

1. INTRODUCTION

Anxiety (1) disorders are a major public health problem. They are the most frequent classes of mental disorders in Western societies for which they are among the leading cause of disability (2) and come at huge individual and societal cost (3). Anxiety is also co-morbid with a variety of medical conditions, exacerbating symptoms, hampering recovery, and increasing risk for other mental disorders such as alcoholism and depression. Even when anxiety symptoms do not reach criteria for a disorder, they can cause misery and poor health (3).

Unfortunately, our understanding of anxiety symptoms remains limited and the treatment of anxiety disorders represents a significant challenge to mental health (4, 5). Current drug treatments often have sub-optimal efficacy as well as unwanted side-effects (5). This situation reflects the difficulty of translational research, which has struggled to capitalize on the array of new technologies to develop efficient treatments. It also points, to some extent, to the practical challenge of developing animal models of mental disorders (6). While animal models of anxiety provide critical insights into basic defense-survival mechanisms (4), promising novel treatments derived from animal models too often end up being ineffective in humans, and none have led to significant improvement to the current armamentarium of anxiolytic drugs, which has been stagnant for several decades (7). This lack of effectiveness has prompted the pharmaco-industry to close or downsize research on mental illnesses (8). Ultimately, this grim picture signals the need for a new approach to translational research to improve the synergy between basic science and clinical science.

In this paper, we argue that experimental models of anxiety in non-clinical, unmedicated humans can revitalize treatment discovery. The “experimental psychopathology” approach can bridge basic research in animals and clinical studies, as well as guide and constrain hypotheses about the nature of psychopathology, treatment mechanisms and treatment targets (9, 10).

This paper focuses on experimental models of anxiety as opposed to fear. Of prime consideration when developing experimental models of psychopathology is the nature of symptoms. Fear and anxiety are different features of anxiety disorders (11) with distinct underlying neurobiology (12). Fear is a phasic response to a well-defined and identifiable proximal threat, whereas anxiety is a more sustained aversive state generated by uncertain future threat (11, 13). Fear is above all a behavioral response that mobilizes the organism to act (fight or flight) in the face of a life-threatening situation. By contrast, anxiety is a feeling of apprehension, a perceived sense of unpredictability, and aversive anticipation (13, 14). While great progress has been made to elucidate the neurobiology of fear thanks to robust translational experimental models such as Pavlovian fear conditioning, comparatively much less is known about anxiety, partly because of the difficulty in modeling its symptoms in animals. More than fear, anxiety involves complex cognitive and executive processes mediated by high-order brain neurocircuits that are strikingly more developed in humans than animals, limiting the usefulness of animal models. Experimental models of anxiety in humans can characterize the normative interplay between anxiety and cognition, paving the way for identifying variability from these norms in pathological anxiety.

In this paper, we 1) briefly discuss the advantages and limitations of animal models and argue for the development of human models of anxiety, 2) present a rationale for the experimental psychopathology approach, 3) describe a neuroscience systems approach to experimental psychopathology in humans to investigate pathological anxiety and its treatments, 4) Provide an empirical illustration of this approach, and finally 5) review issues that need to be addressed in future studies.

2. ANIMAL AND HUMAN MODELS OF ANXIETY

Experimental models and translational research are necessary to improve our understanding of anxiety, anxiety disorders, and their treatments. Of all the mental disorders, anxiety disorders are arguably most amenable to translation because, unlike many psychiatric

symptoms, fear and anxiety are highly tractable in the laboratory. These defense-survival responses to threat can be readily induced and measured in animals and in humans, and their underlying neurocircuits are well-conserved across species (4, 12).

Why do we need to model anxiety in humans given that 1) basic research in animal models on neurobiological mechanisms can go far beyond research in humans and 2) clinical research comparing healthy controls to anxiety patients can identify variations from the norms? We argue that human models of anxiety are necessary for the same reasons we rely on animal models: to enhance basic knowledge and theories, define functional norms to help identify pathological mechanisms and treatment targets, and to screen new treatments. Animal and human models, however, have their own specific advantages, limitations, and can address distinct questions. They are complementary, but for progress to be made, their differing scopes, strengths and weaknesses must be acknowledged.

2.1 ANIMAL VS. HUMAN EXPERIMENTAL MODELS

The advantage of animal models over human models include the availability of more invasive experiments (but see (15)) and of more advanced technical tools (e.g., optogenetic techniques). Research in animal models have advanced the characterization of functional neurocircuits and underlying neurobiology, and the gene to phenotype relationship (16, 17). Animal models are also essential at the early stage of drug development to establish safety, pharmacokinetics, and early evidence of target effectiveness (18).

However, animal models have also important limitations. They have failed to deliver clinically effective psychopharmacological treatments for anxiety disorders (19, 20) and have limited application to test novel treatments (e.g., mindfulness, neurostimulation). This poor predictive validity of clinical efficacy has multiple causes, including poor construct validity (e.g., in modeling subjective experiences) and the difficulty of modeling the cognitive and behavioral symptoms listed under the major classification systems (e.g., DSM-5) (15, 21). As a result, the features of human anxiety modeled in animals can be fuzzy. In the defense of animal models, a key hurdle to their development is the current inadequate nosology of mental disorders based

on phenomenology (i.e., patients' reports, signs) rather than underlying biology, unknown pathogenic mechanisms, absence of biomarkers and recognized highly penetrant genetic variants, clinical heterogeneity, and sex differences in brain structures and functions (6, 22). In fact, animal models are expected to achieve greater validity and stronger clinical relevance when the critical features of anxiety and anxiety disorders (biological markers, genetic makeup, pathophysiology) become better defined, and the exploration of the mechanisms of behavior in animals more sophisticated, owing to the remarkable advances in basic neuroscience tools and methods.

However, the one irreducible issue concerns the cognitive, emotional and behavioral traits that are unique to humans (4). While animal models have been instrumental in elucidating defense-survival mechanisms, these defense-survival mechanisms are only one component of a complex system that contribute to human anxiety. Basic defense mechanisms interact with higher-order cognitive and executive functions mediated by neocortical circuits that are uniquely developed in humans (22). Above all, psychiatry is the field of medicine devoted to pathologies of the mind, which is the most complex and elusive faculty of the human being. The human mind harbors conscious and unconscious experiences and is the origin of conceptual abstraction and symbolic communication that cannot be assessed in animals (4, 15). In addition, the neural architecture of the brain is far more complex in humans than in animals and anxiety disorders often reflect disturbance of evolutionary recent circuits. The neurocircuitry of anxious feelings, mental representations, and emotion regulation in humans implicates prefrontal areas, which are uniquely developed in humans. Many psychiatric symptoms (e.g., worries) cannot be convincingly evaluated in animals, and while some domains of functioning (abnormal social behavior, working memory, attention control) can be modeled in animals, the analogy is only an approximation (4, 6).

Human models can address these limitations. Experimental models in humans provide insights into the core subjective, cognitive, and self-regulatory aspects of anxiety, which are responsible for functional impairment and bring patients to the clinic. Experimental psychopathology can characterize what is behaviorally and cognitively deviant from the norms

in anxiety disorders. Human models are valuable tools to identify potential pathophysiological mechanisms and treatment targets (23-25). Clearly, experimental psychopathology cannot achieve the detailed analysis of brain function that is possible in animal research. However, in coming years, improvement in neuroimaging techniques and computational psychiatry will enhance the yield of human research (26).

2.2 HUMAN EXPERIMENTAL MODELS VS. CLINICAL RESEARCH

Human models of anxiety in healthy individuals can inform and complement clinical research with patients. This is particularly important since research in patients is complex, costly and challenging with regards to recruitment and confounding factors, such as comorbidity and drug treatment (27). Fundamentally, experimental models in humans can test reciprocal causal and correlational effects of anxiety on cognition and executive functions (28). They can also help disentangle the role of anxiety in other conditions that are often comorbid with anxiety (see (29)).

Experimental models in healthy subjects also presents an advantage of scale in that they can leverage the benefit of relatively easy recruitment of participants (relative to psychiatric patients) to speed up research and widen its scope for an in-depth investigation of the multiple faceted aspects of anxiety.

In addition, a most promising application of human models relates to treatment, providing some understanding on how treatments work (23, 30). In addition, human models could be very useful for the screening of novel anxiolytics (24, 25), adding specificity to the high sensitivity of animal models (see Section 6.6). Experimental psychopathology may also be an instructive first step to test non-pharmacological treatments that cannot be assessed in animals such as cognitive and behavioral treatments (e.g., mindfulness) or neuromodulation (31).

Finally, because anxiety disorders are developmental brain disorders and because the manifestation of anxiety is different in children and in adults (32), experimental

psychopathology affords the opportunity to implement a developmental perspective for anxiety research.

3. EXPERIMENTAL PSYCHOPATHOLOGY: DEFINITION AND VALIDITY

Experimental psychopathology is conducted in healthy individuals, and employs experimental models of psychopathology (e.g., anxiety) to gain insights into normal and abnormal behaviors and their underlying neurobiological mechanisms (33, 34). The validity of experimental models of anxiety in healthy humans is consistent with the dimensional conceptualization of psychopathology (35, 36), which assumes *a continuum from normal to pathological anxiety*, and therefore a similar continuity for underlying mechanisms. While the conceptualization of anxiety disorders in terms of dimensions or categories has long been debated (37, 38), the dimensional view is espoused by pre-clinical (39) and clinical research (36). That anxiety disorders result from inappropriate activation or exaggeration of otherwise adaptive responses to threat provides the rationale behind the experimental psychopathology approach. The same anxiety system that is perturbed in anxiety disorders is also activated by anxiety challenges in normal individuals, albeit in less extreme, persistent or incapacitating forms (40-43) (see also section 4.3). Fear conditioning studies are a validation of this view. Indeed, many of the neural structures underlying normative fear mechanisms in healthy subjects have been implicated in pathological anxiety, especially symptoms that are fear-specific such as response to phobic stimuli (44). It follows that normative neuromechanisms of anxiety should provide a blueprint for the search of pathological mechanisms and drive hypotheses about neurocircuit dysfunction in pathological anxiety. However, while the experimental psychopathology approach is consistent with the dimensional view of anxiety, it remains a hypothesis and, as such, it is falsifiable. If it turns out that normative and pathological mechanisms differ, the dimensional view of psychopathology will have to be revisited.

4. A RECIPE FOR EXPERIMENTAL PSYCHOPATHOLOGY RESEARCH ON ANXIETY

Anxiety is multifaceted and cannot be expected to be recapitulated in a single model. Like animal models, the intent of human models is not to mimic the whole disease (6), but to develop specific tests of behavioral and psychological operations relevant to the symptoms and treatment of the target disorder. Findings informing how experimentally-induced anxiety affect these operations help *generate hypotheses* about biomarkers and underlying neurobiological dysfunction and *narrow the search* for treatment targets for the specific modelled symptoms.

While studies of fear-related defense mechanisms examine short-duration responses to threat cues (e.g., Pavlovian fear conditioning), studies focusing on anxiety examine sustained aversive states. Anxiety induction procedures include situations such as anxiogenic pharmacological challenges (e.g., 7.5% CO₂, CCK4) (45-52), darkness (53-55), or threat of unpredictable aversive stimuli (e.g., shock) (56-58). So far, most studies have investigated defensive mechanisms in *idle* subjects (i.e., subjects not involved in a complex task) and have enhanced our understanding of the clinical relevance of these anxiety states (59-66) and their underlying neurobiology (46, 67-77).

However, this type of investigation of sustained anxiety, which focuses uniquely on emotional expression, is limited in scope and does not fully capitalize on the advantages of experimental models in humans. Indeed, the cognitive aspects of anxiety state are for the most part ignored. This lapse is an issue since emotional and cognitive processes are not organized separately but contribute jointly to ongoing behavior (78) and subjective experience relies on cognitive processes such as attention and working memory (79). In addition, cognitive formulations of anxiety provide elaborated theoretical models of psychopathology (80-83) and define the processes targeted by successful cognitive therapies. These theories emphasize the role of information processing (i.e., cognition) in the etiology and maintenance of anxiety disorders. In overanxious individuals, faulty perception, encoding, storage, retrieval, interpretation, control, and action interfere with ongoing goals while deficits in executive function impairs self-regulation (emotion regulation) (81, 84-87). However, we currently have

little mechanistic understanding of the impact of anxiety on these processes and their dysregulation in anxiety disorders. It is to study these cognitive processes that experimental psychopathology can be most beneficial. A better characterization of the dynamic between anxiety and cognition with the objective of identifying core cognitive processes that are vulnerable to anxiety could significantly improve our understanding of pathological mechanisms.

Investigators have long used systems neuroscience to elucidate neurocircuits mediating behavioral plasticity and pharmacological effects (88). This approach begins with the characterization of the functional neural architecture that mediates the behavior under scrutiny to subsequently identify where and how plasticity occurs in response to experimental manipulations. The systems neuroscience approach has been used with great success to uncover underlying mechanisms of simple defense response to threat in animals (88, 89) and in humans (90). We argue that this approach can be applied to investigate how anxiety affects complex perceptual, cognitive and behavioral processes in experimental psychopathology studies in humans.

The overall strategy is to combine a primary cognitive or behavioral task with experimentally-induced anxiety. Its implementation follows 5 steps:

1) select anxiety-relevant cognitive or behavioral operations (e.g., attention bias for threat) and tasks (e.g., dot-probe) to probe these processes,

2) identify the underlying neurocircuits via connectivity/activation nodes in healthy volunteers,

3) examine the impact of experimental anxiety on the targeted operations and underlying neurocircuits in healthy volunteers,

4) utilize findings from step 3 to generate and constrain hypotheses about mechanisms of pathophysiology and neurocircuit dysfunction in anxious patients, and

5) finally, evaluate mechanisms of treatment efficacy and screen putative anxiolytic treatments in healthy individuals exposed to experimental anxiety.

4.1 STEP 1: FUNCTIONAL BEHAVIORAL AND COGNITIVE DOMAINS AND MARKERS OF ANXIETY

The first step is to select anxiety-relevant psychological and behavioral processes or constructs. One productive approach is to base this selection on findings from clinical research, which compares patients with anxiety disorders to individuals without psychiatric disorders. These studies have identified cognitive, executive, behavioral, neurobiological, and electrophysiological abnormalities in anxiety patients. Results of these studies have generated important theoretical models and psychological constructs that underscore key cognitive, executive and behavioral variables presumed to play a role in the origins and maintenance of anxiety disorders. These include negative bias in information processing (86, 91, 92), worry (93, 94), emotional dysregulation (95), behavioral inhibition (96), and avoidance (97). There are several established tasks and measures to investigate these constructs. For example, information processing biases can be evaluated using event-related potentials (ERPs) in response to sensory stimuli (98-100) or behavioral responses on cognitive tasks such as the dot-probe, visual search paradigms, and the Stroop test (86). These tasks have enhanced our understanding of how anxiety captures attention, but they have also generated conflicting theories regarding the nature and the stage of information processing deficits in anxiety (86). Resolution of these conflicts can be facilitated with studies of experimental anxiety in healthy subjects.

Another promising and more systematic approach relies on the Research Domain Criteria (RdoC) project, which seeks to relate dysfunction of specific neurobiological systems to symptoms (101). The RdoC is organized around fundamental functional constructs that are studied along the full range of variation, from normal to abnormal, across multiple units of analysis. These constructs are grouped into higher-level functional domains and are investigated using well-validated paradigms that have been selected by experts in the field.

Using a RDoC approach, experimental psychopathology seeks to better understand how anxiety changes behavior-brain relationships, and then relates these changes to pathological anxiety. One could argue that anxiety spans several RDoC constructs, the “potential threat” construct of the negative valence systems, and various constructs of the cognitive systems, including attention, perception, cognitive control, and working memory. For example, attentional bias for threat falls into the cognitive system and the negative valence system. From a RDoC perspective, experimental psychopathology explores the interaction between the domains of potential threat and various cognitive systems constructs. The RDoC represents a valuable framework for organizing experimental psychopathology research as it provides the scientific rationale to select constructs and validated tasks to probe these constructs. However, the list of constructs is not exhaustive, and it could be worthwhile to identify additional anxiety-relevant cognitive constructs (e.g., affective decision making) (102). This could be accomplished by following the example of research on the negative symptoms of schizophrenia. The MATRICS (Measurement and Treatment Research to Improve Cognition in Schizophrenia) initiative was designed to develop a consensus cognitive battery to stimulate the development of new drugs that target cognitive deficits in schizophrenia (103). A panel of experts was put together to identify relevant cognitive domains and constructs and select a battery of cognitive tests relevant to schizophrenia. A similar approach would greatly benefit the study of the core cognitive symptoms of anxiety.

4.2 STEP 2: NORMATIVE MECHANISMS IN HEALTHY SUBJECTS

This step belongs to the domain of cognitive neuroscience, which seeks to understand the neurocircuits of cognitive processes. Step 2 of the experimental psychopathology approach consists of identifying, in healthy individuals, the neural circuits underlying the behavioral/cognitive constructs selected in step 1. Neuroimaging techniques are the basic tools to examine these questions. They include most commonly magnetic resonance imaging (MRI), but also magnetoencephalography (MEG), and less frequently positron emission tomography (PET). Whereas information can be gleaned from structural brain data, functional data provide

more direct information about cognitive processes, since they record changes in neural activity that cooccur with changes in behavior. Two basic measures are supplied by fMRI: regional activation and functional connectivity. Most of the analyses, up till now, are correlational in nature. For example, working memory is a cognitive construct which has been “associated” with activations within the dorsolateral prefrontal cortex and parietal cortex. These regional activations are interpreted as forming the neural circuit that supports working memory. Naturally, the same is true for the functional connectivity measure.

Extended efforts have been made to formulate mathematical algorithms that could generate causal models. Two complementary methods, Granger causality (G-causality, (104)) and dynamic causal modeling (105) have been developed, but have not been extensively applied. Reasons for this include the many assumptions and requirements needed for their successful application, and their complexity. A more in-depth discussion of these techniques is outside the scope of this review but see (106). More recently, the introduction of machine learning tools, while large datasets become publicly available, may change the landscape of brain-behavior neuroscience research. For the perspective of this review, machine learning tools offer the possibility of exploiting a large array of behavioral and neuroimaging variables to not only refine understanding of neural circuits, but also discover novel mechanisms underlying specific behavioral processes.


4.3 STEP 3: STUDIES OF THE INTERPLAY OF ANXIETY AND COGNITION IN HEALTHY SUBJECTS

Once the behavioral/cognitive process (Step 1) and its underlying neurocircuits (Step 2) are delineated, Step 3 assesses the effect of experimentally-induced anxiety on these factors. Step 3 is the critical step that determines the relevance of the cognitive construct and associated neural circuit to anxiety. Promising results from this step are then followed up in steps 4 and 5. However, negative results signal the end of this specific investigation. Step 3 is the stage of experimental psychopathology that departs from but is the link to clinical research. In Step 3, cognitive tasks and their behavioral and neural signatures are explored in healthy individuals in a control condition and under a condition of experimentally-induced sustained

anxiety. Anxiety-related changes in cognitive performance and underlying neural substrates are then hypothesized to be markers of anxiety in general, and are expected to be relevant to clinical anxiety, which is what step 4 and 5 will test. Fig 1 illustrates a hypothetical scenario where a neurocircuit underlying response to a cognitive task is affected by induced-anxiety in healthy individuals. A network of three structures (ROI_1 , ROI_2 , ROI_3) is implicated functionally in the task. During an anxiety-induction procedure, another structure, ROI_x is activated and alters connectivity in the network, between ROI_1 and ROI_2 . These changes in neural activity and connectivity would be expected to be implicated in pathological anxiety.

Healthy subjects performing a cognitive task

No anxiety-induction



Anxiety-induction




Fig. 1. Diagram illustrating the 3rd step. The left side describes a hypothetical network of three structures (ROI_1 , ROI_2 , ROI_3) that is activated in response to a cognitive task in healthy individuals. The right side shows the activation of the same network when subjects are challenged with an anxiety induction procedure. When subjects are anxious, structure ROI_x comes on line and alters the connectivity in the network between ROI_1 and ROI_2 .

Ultimately, the aim of experimentally-induced anxiety is to uncover markers of anxiety that can lead to treatments. There has been a steady increase in studying the interplay of

anxiety and cognition in recent years. The great majority of studies use the threat of unpredictable shocks (28, 107-109, 110, 111-114) (also reviewed in (115)) or 7.5% CO₂ inhalation (29, 41, 42, 116). The trend that emerges is that anxiety induced by threat of shock can have detrimental but also beneficial effects depending on the nature of the task (reviewed in (115)). Generally, threat of shock facilitates early sensory processing, the detection of threat information, and interference from distractors. It impairs short-term memory but facilitates long-term memory. It does not have a uniform effect on inhibitory control, impairing cognitive inhibition in emotional Stroop tests, while facilitating response inhibition in go-nogo tasks. Breathing 7.5% CO₂ increases measures of hypervigilance such as orienting and alerting (41, 42), negative interpretation bias (117). Overall, many of these effects mirror the effect of clinical anxiety on cognitive processes.

Therefore, the next step (Step 4) is to evaluate the use of these findings in healthy subjects as markers of clinical anxiety

4.4 STEP 4: MECHANISMS OF PATHOPHYSIOLOGY

One objective of experimental psychopathology is to guide hypotheses about dysregulated cognitive processes and underlying neurocircuits in pathological anxiety. The dimensional perspective of psychopathology predicts that pathological anxiety is associated with perturbations in some of the processes and neural mechanisms engaged by experimental anxiety. In the example we provide above (Fig. 1), one would hypothesize that structure ROI_x and connectivity between ROI₁ and ROI₂ are potentially implicated in pathological anxiety.

What is the nature of the dysfunction that should be expected? From a dimensional perspective, the mechanisms that exacerbate responses to anxiety in healthy individuals should be amplified in anxiety patients (i.e., amplified excitatory mechanisms, and weakened inhibitory mechanisms). However, neuroadaptation following chronic anxiety may change the nature of these responses and underlying neurocircuits over time. Consistent with this view, while studies generally show increased defensive reactivity and increased amygdala activity in pathological

anxiety (66, 118), instances of hypo-reactivity in patients are not infrequent (119, 120). The current interpretation of these opposing effects is that they reflect distinct symptoms. For example, while pathological anxiety tends to be associated with exaggerated defensive responses as measured with startle (60, 61), increased illness chronicity and comorbidity is associated with blunted reactivity (120). A similar pattern of response has been reported at the neural level. In PTSD, for example, amygdala hyper-reactivity is associated with hypervigilance and intrusive thoughts, whereas amygdala hypo-reactivity leads to symptoms of dissociation and disengagement (119).

Another consideration concerns whether these maladaptive responses are triggered by specific events or whether they are chronically expressed. According to the diathesis-stress model of anxiety disorders, functional perturbation of these adaptive mechanisms should be greatest during periods of anxiety or stress (e.g., during experimental threat). Alternatively, these mechanisms could be chronically and maladaptively engaged, even in the absence of experimental threat.

4.5 STEP 5: TREATMENT IMPLICATIONS

Given the poor predictive validity of animal models, another promising clinical utilization of experimental psychopathology is treatment research. Although there is a dearth of research in this area, the two broad applications of experimental models in humans are to explore the mechanisms through which treatments exert their effect and to screen candidate anxiolytics in proof-of-concept studies.

Treatment mechanisms: The mechanism by which conventional pharmacological treatments, such as the SSRIs and the benzodiazepines, or cognitive behavioral therapy exert their effect is poorly understood. Experimental psychopathology is being used to clarify mechanisms responsible for the clinical efficacy of current pharmacological or psychological treatments. Results show that the effectiveness of SSRIs may be partly due to their down-regulating effect on attentional bias for threat (23, 30). Studies focused on defensive

mechanisms indicate the benzodiazepine alprazolam and the SSRIs citalopram, which are used to treat anxiety patients, reduce defense responses to unpredictable shock but not predictable shock in healthy subject (9, 10). Interestingly, alcohol, which is used for its anxiety-dampening effect, also reduces defense responses to unpredictable but not predictable shock (121). As research in humans and rodents point to distinct neurocircuits responsible for response to predictable and unpredictable threat, these results provide clues as to the structures that may be preferentially implicated in these anxiolytic effects (12).

As expected, given the role of cognitive dysfunction in anxiety disorders, there is also evidence that conventional anxiolytics act on cognitive processes. Studies in healthy volunteers have shown that 7 days of treatment with the benzodiazepine diazepam or with the SSRI citalopram reduces a pattern of attentional vigilance to threat (122, 123). Because cognitive models suggest that anxiety disorders are associated with attentional biases for threat, these results suggest that the therapeutic action of classical anxiolytics may partly be mediated by normalizing these biases.

Proof-of-concept: Anxiolytics that act preferentially through the γ -aminobutyric acid (GABA) or serotonergic systems have been the benchmark since the 50's and 80's respectively. To improve the efficacy of anxiolytics, research has sought to discover new compounds that either target these systems or have a novel mode of action, for example acting on the neuropeptide, glutamate, or endocannabinoid systems (20). However, despite important breakthroughs in basic science that have led to the development of a multitude of drug anxiolytic candidates, the ability to bring to the marketplace efficient new compounds has not improved. The lack of success of drug development can be traced to difficulties in selecting among the many candidate anxiolytic compounds because of costly and time-consuming clinical trials and the failure of animal models of anxiety to predict clinical efficacy. As long as candidate anxiolytics are safe, there is no reason why they could not be tested in models of anxiety using healthy individuals to establish whether they are actually anxiolytic.

The predictive validity of a model refers to the ability of the model to predict clinical efficacy. However, a model has predictive validity if it can successfully differentiate among *effective* and *ineffective* treatments. Animal models have a high level of sensitivity but poor specificity. The hope is that human models can add *specificity* and help drug sponsors make 'go-no-go' decisions about novel compounds and choose appropriate treatment doses for subsequent clinical trials. Given their high cost and long duration, clinical trials are limited in the number of novel compounds that can be tested. New drug development methods, such as experimental models in humans, may facilitate and speed up screening and improve the predictability and the efficacy of candidate anxiolytics.

Whether this approach will be successful remains to be seen. So far, the few studies that have tested putative anxiolytics in anxiety models in humans have had mixed success, as illustrated by translational research on antagonists of corticotropin-releasing factor (CRF) receptors. Based on strong preclinical evidence that CRF was a key mediator of stress-related responses, it has been suggested that drugs that target the CRF₁ systems could be developed to relieve anxiety (as well as depression and alcoholism) (124). CRF₁ antagonists have proved to be anxiolytic in a wide range of animal models (125), but results in human models have not been as consistent. The CRF₁ antagonist GSK876008 reduced startle potentiation in animal models of anxiety (126), but acute treatment with similar CRF₁ antagonist, Verucerfont (GSK561679), did not reduce startle potentiation or subjective feeling of anxiety to unpredictable shock in a double-blind, crossover design in healthy subjects (24). However, a 7-day treatment with another CRF antagonist (R317573), reduced subjective response but not physiological response to a 7.5% CO₂ inhalation challenge in a double-blind, randomized, placebo and active controlled study (25). These rather negative results should be considered in the light of clinical data that have shown CRF₁ antagonists to be clinically inefficient to treat generalized anxiety disorder, social anxiety disorders, depression, and comorbid anxiety-alcoholism (127, 128). It is noteworthy that the same CRF₁ antagonist that was tested in the fear-potentiated startle model in humans (24), was also found to have no anxiolytic effect when given chronically in individuals with PTSD (129). Although negative, these results have a positive side. The fear-

potentiated startle model in humans did not produce a false positive: a clinically ineffective drug was ineffective in the human model.

Beside screening putative anxiolytics, human models will be necessary to evaluate non-pharmacologic treatments, such as exercise (130), cognitive-behavioral behavioral (131), mindfulness, or neuromodulation (31). One advantage of experimental models in humans is that their sensitivity and specificity may be improved by targeting pathological mechanisms identified via the experimental psychopathology approach.

5. EXAMPLES OF EXPERIMENTAL PSYCHOPATHOLOGY APPROACH TO PRE-ATTENTIVE INFORMATION PROCESSING

Cognitive models emphasize the role of deficits in early information processing as a primary and persistent manifestation of pathological stress and anxiety, contributing to symptoms of hypervigilance and downstream malfunction of more complex cognitive operations (86, 132, 133). Various psychological constructs have been employed to study early information processing deficits in pathological anxiety, including attentional bias (86), sensory gating (134, 135), and pre-attentive perceptual processing (133, 136-138). This section illustrates how experimental psychopathology can provide clues about neurocircuit dysfunction using two specific examples: 1) perceptual sensitivity and 2) negative biases.

5.1 PERCEPTUAL SENSITIVITY

5.1.1 Step1: Functional domains and markers of anxiety: mismatch negativity

Effective early detection of environmental change is adaptive as it provides for rapid orienting to potential threats, driving the organism to adopt cautious behavioral strategies (139). Early perceptual responding to environmental changes can be examined with oddball stimulus paradigms. These paradigms assess how the organism responds to rare, deviant stimuli. A consistent electroencephalographic response to these unexpected stimuli has been

identified and labeled *mismatch negativity* (MMN). In other words, the MMN is a brain-evoked potential in response to stimuli that deviate (prediction errors) from an established familiar sequence of sensory stimuli (e.g., change in tone frequency) (140). It is considered a measure of preattentive detection because it can be elicited when attention is focused elsewhere (141).

As a measure of sensory perception and auditory discrimination, the MMN provides a window into the state of vigilance of the brain (142, 143). Consistent with this view, the MMN is abnormally elevated in a number of anxiety-related disorders, including PTSD (136, 137), panic disorder (138), and phobia (144). These findings support brain's heightened sensitivity to environmental changes in these patient populations (i.e., hypervigilance). Heightened MMN is also associated with behavioral inhibition, a temperamental vulnerability for later anxiety disorders (145). These results indicate that the MMN can be used as a proximal measure of anxious hypervigilance to gain insights into underlying neurocircuit dysfunction.

5.1.2 Step 2: Normative mechanisms in healthy subjects

While the neural regions contributing to MMN generation have long been elucidated (146), more recent works relying on Dynamic Causal Modeling (DCM) have provided a mechanistic understanding of how the magnetoencephalographic MMN (MMNm) is generated (147). The MMNm is generated within a well-established frontotemporal network composed of bilateral sources over the primary and secondary auditory cortex (superior temporal gyrus, STG), and inferior frontal gyri (IFG) (147) (Fig. 2, left). DCM shows that the MMN can be explained by changes in the strength of the connectivity between (extrinsic) and within (intrinsic) these cortical sources (Fig. 2, right) (147). Forward connections can be conceptualized as bottom-up processes transmitting sensory information from A1 to higher cortical levels and convey prediction errors (MMN). Backward connections represent top-down predictions based on prior sensory experience and explain away prediction errors (deviance detection).

5.1.3 Step 3: Studies of the interplay of anxiety and cognition

Consistent with the dimensional view of psychopathology, the MMNm is also increased by experimental anxiety in healthy subjects (98, 99). The underlying mechanisms responsible

for this increased has been recently elucidated via DCM and involves a rebalance of forward and backward connections. Threat of shock enhances post-synaptic gain in primary auditory cortex and modulation of feedforward pathway but attenuates the normal feedback signaling, which results in a failure to attenuate ascending prediction errors (98). These results suggest that anxiety-induced hypervigilance results from heightened sensitivity of bottom-up processes and failure of top-down modulation.




Fig. 2. Left. The MMNm arises from bilateral sources over primary, secondary auditory cortex (superior temporal gyrus, STG), and inferior frontal gyri (IFG) (146). Right. Dynamic causal modeling shows that the MMN results from changes within and among these cortical sources (147). Forward connections can be conceptualized as bottom-up processes transmitting sensory information from A1 to higher cortical levels and convey prediction errors (MMN). Backward connections represent top-down predictions based on prior sensory experience and explain away prediction errors (deviance detection). In the safe context, the MMN is mediated by changes in both extrinsic feedforward and feedback connectivity as well as intrinsic connectivity . Anxiety induced by threat of

shock suppresses feedback connectivity and the anxiolytic alprazolam reestablishes feedback connectivity (not shown).

Having established via DCM that the changes within the MMNm neurocircuit are responsible for the increased MMNm by anxiety, the next step will be to identify the structure(s) that cause(s) these changes. One potential structure is the amygdala. The amygdala plays a critical role in threat evaluation (148), boosts processing (149), is involved in novelty detection (150, 151), and, importantly, is activated by stimulus deviance in a time window that corresponds to IFG activation (99).

5.1.4 Step 4: mechanisms of pathophysiology

Results of experimental psychopathology constrain the search for the perturbation underlying the increased MMN in anxiety pathology. The identification of the normative mechanisms responsible for the increased MMN by anxiety leads to the hypotheses that comparable changes in the balance between feedforward and feedback signaling caused by threat of shock in healthy subjects are also responsible for the increased MMN in anxiety pathology. This hypothesis has not yet been tested.

5.1.5 Step 5: Treatment implications

Drugs acting on gamma-amino butyric acid-receptors can downregulate and upregulate hypervigilance (152, 153) and decrease or increase the MMN (154-156), respectively. In healthy subjects, the benzodiazepine alprazolam attenuates the threat-modulated MMNm and reestablishes the normal balance between feedforward and feedback signaling (98). This suggests that benzodiazepines may reduce hypervigilance, partly by dampening early deviance detection. These results suggest that treatment aimed at reducing hypervigilance symptoms in pathological anxiety should target the balance between feedforward and feedback implicated in deviance detection.

5.2 NEGATIVE BIASES

5.2.1 Step1: Functional domains and markers of anxiety: attention bias

Attention bias for threatening stimuli has long been associated with pathological anxiety (86). It reflects the propensity to rapidly detect and react to threat and can be investigated using a wide range of tests which reflect different aspects of attention, including the dot-probe, the emotional Stroop, and emotional face processing (86, 115, 157). A face processing task is the exemplar selected here.

5.2.2 Step 2: Normative mechanisms in healthy subjects

Neural models of emotion amplification and emotion regulation suggest several neural signals implicated in attention bias for threat. The amygdala automatically detects salient environmental stimuli (158, 159). Response flexibility is associated with structures that amplify the threat signals, such as the rodent's prelimbic cortex or its putative human equivalent, the dACC/dmPFC, and structures that protect goal-directed processing from threat distractors, including the ventromedial PFC (vmPFC), ventrolateral PFC (vlPFC), ACC, and IFG (160-162). It is likely that some of these structures work together for adaptive behavior. Experimental models in humans have helped refine our understanding of functional connectivity among these structures to support biases in attention in changing environments.

5.2.3 Step 3: Studies of the interplay of anxiety and cognition

Studies in healthy subjects during induced anxiety have highlighted the role of cortical-subcortical connectivity in attention bias for threat. More specifically, in a series of studies using an emotional face identification task, it was first reported that induced-anxiety drives attentional bias; subjects show quicker detection of fearful faces and greater defense response (i.e., as measured with startle) upon presentation of fearful faces during threat of shock compared to a safe condition (157, 163). It was then shown that this heightened attentional bias was associated with increased connectivity between the dACC/dmPFC and the amygdala, providing evidence of an "aversive amplification circuit" that strengthens amygdala response to

facilitate threat detection (Fig. 3) (164, 165). In addition, the strength of this connectivity correlated positively with the behavioral bias index, suggesting this circuit drives negative bias.

It has been argued that the dACC/dmPFC-amygdala neurocircuit serves to 'prime' the amygdala, maintaining the amygdala in a state of readiness. This provides a mechanism by which the amygdala can detect and react rapidly to alerting and potentially dangerous stimuli during sustained anxiety states without maintaining a sustained level of activation (166).




Fig. 3. Proposed model of attention bias for threat in anxiety. Increased bi-directional coupling between the dorsal anterior cingulate/dorsomedial prefrontal cortex (dACC/dmPFC) and amygdala (amg) promotes threat bias. Coupling is transiently activated during anxiety states in healthy control, facilitating the detection of threat. However, the coupling is chronically engaged in pathological anxiety, leading to maladaptive threat bias. Lowering serotonin has similar effect as induced-anxiety: it increases dACC/dmPFC-amygdala coupling. By increasing serotonin, SSRIs may reduce attention bias for threat via inhibition of dACC/dmPFC-amygdala coupling.

5.2.4 Step 4: mechanisms of pathophysiology

Critically, it was subsequently found that the same dACC/dmPFC-amygdala neurocircuit is overactive in anxiety patients, in the absence of experimental anxiety (i.e., without threat of shock) (92). In other words, the same neurocircuit responsible for threat bias, which is appropriately engaged and disengaged in healthy subjects by threat and safety, is chronically activated in anxious patients (i.e., in safe environments)

These studies validate the experimental psychopathology approach and further provide evidence that an otherwise self-protective neurocircuit is dysfunctional in anxiety disorders. The positive correlation in healthy individuals between trait anxiety and the strength of the dACC/dmPFC-amygdala coupling (164) suggest that this neurocircuit indexes a vulnerability to anxiety disorders.

5.2.5 Step 5: Treatment implications

The dACC/dmPFC-amygdala neurocircuit therefore represents a potential neural target for treatment. Neural markers are more proximal than behavioral markers, and as such, targeting the dACC/dmPFC-amygdala circuit may prove to be a powerful way of developing treatments .

Ideally, one would want to decrease dACC/dmPFC-amygdala coupling (92, 167). The mechanism by which activity in this neurocircuit can be reduced is not fully understood, but evidence shows that serotonin plays a key role. Serotonin affects affective processing (168, 169) and the first-line of psychopharmacological treatment for anxiety disorders, the SSRIs, alter serotonin neurotransmission (170). In addition, amygdala response to fearful faces is modulated by a serotonin transporter polymorphism (171). More direct evidence of serotonin involvement in dACC/dmPFC-amygdala coupling comes from findings that, in healthy subjects, reduced serotonin function following depletion of its precursor, tryptophan increases the strength of the connectivity in this circuit (172). These results suggest that SSRIs may reduce anxiety symptoms by increasing serotonin, which normalizes or reduces the excessive dACC/dmPFC-amygdala coupling responsible for promoting threat bias. This interpretation would be consistent with evidence that serotonin therapeutic effects is mediated by the serotonergic reduction in attention bias (169, 173). Fig. 3 illustrates this possibility. Given that increased dACC/dmPFC-amygdala coupling promotes threat biases and reducing serotonin increases dACC/dmPFC-amygdala, SSRIs may reduce threat biases by inhibiting dACC/dmPFC-amygdala coupling. Interestingly, psychological treatments (e.g., cognitive behavioral therapy)

of anxiety disorders, which can be as effective as pharmacological treatment (174), may also work by disengaging this circuit (167). .

6. FUTURE DIRECTIONS

Whether the experimental psychopathology approach will bring a new insight into psychopathology and will help screen novel treatments remain to be seen. Experimental psychopathology is still in its infancy, but recent progresses are encouraging, supporting further development of this approach to fully contribute to clinical research. Below, we briefly review of few issues that need to be addressed to take full advantage of experimental psychopathology.

1. *Sex and developmental changes*: Anxiety disorders are more prevalent in women (175) and girls experience more anxiety symptoms than boys. Experimental psychopathology can be applied to children and adolescents (176) and can contribute to elucidating the neural basis of this sex difference by characterizing developmental changes.
2. *Inter-individual differences*: Response to anxiety-inducing procedures (177-180) and cognitive performance (181) vary among individuals, reflecting the influence of temperamental, genetic, or environmental factors. It is likely that the interaction of cognition and experimental anxiety, as well as the treatment effects on these interactions, are similarly affected by inter-individual differences. It will be important to identify these factors since they may critically influence psychopathological mechanisms and treatment targets. By helping identify the inter-individual characteristics that determine treatment effects, experimental psychopathology carries implications for personalized medicine (182).
3. *Anxiety induction procedures*: Different classes of environmental threats activate different defense mechanisms. Defense responses to proximal vs. more distal or uncertain threats are distinguishable in terms of behavior, cognition, and neurobiological substrates (183) as well as psychopathology (11). Similarly, the response to bodily harm differs from that of social threat (184, 185). Aversive stimuli such as shocks have been successfully used in

experimental models in humans (60, 61, 186). However, threat of shock may not be appropriate to model all symptoms of anxiety. For example, low dose of CO₂ (e.g., 7.5%) challenge has been proposed as a model of general anxiety disorder (187). Behavioral avoidance tasks, where subjects anticipate a feared situation (e.g., anticipation of public speaking in social anxiety) is also a promising tool (188). For example, healthy controls and patients with social anxiety show distinct emotional responses and cognitive regulation to social threat compared to physical threat (185, 189) and avoidance performance in a virtual elevated plus-maze is associated with symptom of acrophobia but not social anxiety or trait anxiety (190). This approach can also be extended to more specific symptoms such as emotional distraction by worries and intrusive thoughts (191). Future works will need to examine the commonalities and differences in the effects of different types of anxiety induction procedures on cognitive and behavioral performance and underlying neurocircuits (192). It will be important to determine the extent to which different anxiety or stress induction procedures model symptoms that are similar or vary across disorders.

4. *Drug screening*: One exciting potential application of experimental psychopathology is to generate proof-of-concept evidence of the efficacy of novel psychopharmacological and psychological treatments (23, 24, 187). Currently, determining the efficacy of a drug or psychological intervention requires expensive and time-consuming clinical trials in patients. Experimental psychopathology could provide optimization of this process, leading to a rapid and affordable indication of efficacy at an early stage of treatment development (19). However, a question for future studies is the extent to which treatments that work on normative responses also work on pathological responses.

7. CONCLUSION

Animal models are essential to advance our understanding of anxiety, but their limitations are increasingly recognized. We have proposed a general framework centered on experimental psychopathology to improve research on pathological anxiety. Combining anxiety

challenges with tasks that probe clinically-relevant psychological or behavioral constructs to identify clinically-relevant neurocircuits provide a new approach to translational research on pathological anxiety.

Acknowledgements and Disclosures: This work was supported by the Intramural Research Program of the National Institute of Mental Health, project number ZIAMH002798 (clinical protocol 02-M-0321 (NCT00047853), 01-M-0185 (NCT00026559)) to CG, grant numbers MR/K024280/1 and MR/K0240817/1 to OJR.

OJR has completed consultancy work for IESO digital health and Brainbow and is running an Investigator Initiated Trial with Lundbeck. He holds an MRC Industrial Collaboration Award with Cambridge Cognition. The other authors declare no conflict of interest.

REFERENCES

1. Stein MB, Craske MG. Treating Anxiety in 2017: Optimizing Care to Improve Outcomes. *JAMA*. 2017:E1-E2.
2. Kessler RC, Chiu WT, Demler O, Merikangas KR, Walters EE. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2005;62:617-627.
3. Chisholm D, Sweeny K, Sheehan P, Rasmussen B, Smit F, Cuijpers P, Saxena S. Scaling-up treatment of depression and anxiety: a global return on investment analysis. *The Lancet Psychiatry*. 2016;3:415-424.
4. LeDoux JE, Pine DS. Using neuroscience to help understand fear and anxiety: a two-system framework. *Am J Psychiatry*. 2016;173:1083-1093.
5. Koen N, Stein DJ. Pharmacotherapy of anxiety disorders: a critical review. *Dialogues Clin Neurosci*. 2011;13:423-437.
6. Nestler EJ, Hyman SE. Animal models of neuropsychiatric disorders. *Nat Neurosci*. 2010;13:1161-1169.
7. Zoellner LA, Foa EB. Applying Research Domain Criteria (RDoC) to the study of fear and anxiety: A critical comment. *Psychophysiol*. 2016;53:332-335.
8. Nutt D, Goodwin G. ECNP Summit on the future of CNS drug research in Europe 2011: Report prepared for ECNP by David Nutt and Guy Goodwin. *Eur Neuropsychopharmacol*. 2011;21:495-499.
9. Grillon C, Chavis C, Covington MS, Pine DS. Two-week treatment with citalopram reduces contextual anxiety but not cued fear. *Neuropsychopharmacol*. 2009;34:964-971.
10. Grillon C, Baas JMP, Pine DS, Lissek S, Lawley M, Ellis V, Levine J. The benzodiazepine alprazolam dissociates contextual fear from cued fear in humans as assessed by fear-potentiated startle. *Biol Psychiatry*. 2006;60:760-766.
11. Association AP: Diagnostic and statistical manual of mental disorders: DSM-5. 5th ed. Washington, DC, American Psychiatric Association; 2013.
12. Davis M, Walker DL, Miles L, Grillon C. Phasic vs sustained fear in rats and humans: role of the extended amygdala in fear vs. anxiety. *Neuropsychopharmacol*. 2010;35:105-135.
13. Barlow DH: Anxiety and its disorders: the nature and treatment of anxiety and panic. New York, Guilford Press; 1988.
14. Mineka S, Zinbarg R: Conditioning and ethological models of anxiety disorders: stress-in-dynamic-context anxiety models. in *Perspective on Anxiety, Panic, & Fear*. Edited by Hope DA. Lincoln and London, University of Nebraska Press; 1996. pp. 135-210.
15. Rollin MDH, Rollin BE. Crazy like a fox: validity and ethics of animal models of human psychiatric disease. *Camb Q Healthc Ethics*. 2014;23:140-151.

16. Kaiser T, Feng G. Modeling psychiatric disorders for developing effective treatments. *Nature Medicine*. 2015;21:979.
17. Santos M, D'Amico D, Dierssen M. From neural to genetic substrates of panic disorder: Insights from human and mouse studies. *European Journal of Pharmacology*. 2015;759:127-141.
18. Bueters T, Ploeger BA, Visser SAG. The virtue of translational PKPD modeling in drug discovery: selecting the right clinical candidate while sparing animal lives. *Drug Discovery Today*. 2013;18:853-862.
19. Dawson GR, Goodwin G. Experimental medicine in psychiatry. *J Psychopharmacol*. 2005;19:565-566.
20. Griebel G, Holmes A. 50 years of hurdles and hope in anxiolytic drug discovery. *Nat Rev Drug Discov*. 2013;12:667-687.
21. Rodgers RJ. Animal models of 'anxiety': where next? *Behavioural Pharmacology*. 1997;8:477-496.
22. MacLean EL. Unraveling the evolution of uniquely human cognition. *Proceedings of the National Academy of Sciences of the United States of America*. 2016;113:6348-6354.
23. Pringle A, Harmer CJ. The effects of drugs on human models of emotional processing: an account of antidepressant drug treatment. *Dialogues in Clinical Neuroscience*. 2015;17:477-487.
24. Grillon C, Hale E, Lieberman L, Davis A, Pine DS, Ernst M. The CRH1 antagonist GSK561679 increases human fear but not anxiety as assessed by startle. *Neuropsychopharmacol*. 2015;40:1064-1071.
25. Bailey J, Papadopoulos A, Diaper A, Phillips S, Schmidt M, van der Ark P, Dourish C, Dawson G, Nutt D. Preliminary evidence of anxiolytic effects of the CRF(1) receptor antagonist R317573 in the 7.5% CO(2) proof-of-concept experimental model of human anxiety. *J Psychopharmacol*. 2011;25:1199-1206.
26. Pine DS. Clinical advances from a computational approach to anxiety. *Biol Psychiatry*. 2017;82:385-387.
27. Montoya A, Bruins R, Katzman MA, Blier P. The noradrenergic paradox: implications in the management of depression and anxiety. *Neuropsychiatric disease and treatment*. 2016;12:541-557.
28. Vytal K, Cornwell B, Arkin N, Grillon C. Describing the interplay between anxiety and cognition: From impaired performance under low cognitive load to reduced anxiety under high load. *Psychophysiology*. 2012;49:842-852.
29. Gillan CM, Vaghi MM, Hezemans FH, van Ghesel GS, Dafflon J, Brühl AB, Savulich G, Robbins TW. Experimentally-induced and real-world acute anxiety have no effect on goal-directed behaviour. *bioRxiv*. 2019:606145.
30. Harmer CJ, Cowen PJ. 'It's the way that you look at it'—a cognitive neuropsychological account of SSRI action in depression. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 2013;368.

31. Balderston NL, Roberts C, Masi E, Deng Z-D, Radman T, Luber B, Lisanby SH, Ernst M, Grillon C: A causal link between right dlPFC Activity and anxiety expression revealed using repetitive transcranial magnetic stimulation. . in 2019 Annual meeting of the Society of Biological Psychiatry. Chicago2019.
32. Pine DS, Helfinstein SM, Bar Haim Y, Nelson E, Fox NA. Challenges in developing novel treatments for childhood disorders: lessons from research on anxiety. *Neuropsychopharmacol.* 2008;34:213-228.
33. Forsyth JP, Zvolensky MJ. Experimental psychopathology, clinical science, and practice: An irrelevant or indispensable alliance? *Appl Prev Psychol.* 2001;10:243-264.
34. Vervliet B, Raes F. Criteria of validity in experimental psychopathology: application to models of anxiety and depression. *Psychol Med.* 2012;43:2241-2244.
35. Kotov R, Krueger RF, Watson D, Achenbach TM, Althoff RR, Bagby RM, ...Zimmerman M. The Hierarchical Taxonomy of Psychopathology (HiTOP): A dimensional alternative to traditional nosologies. *J Abnorm Psychol.* 2017;126:454–477.
36. Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, Sanislow C, Wang P. Research Domain Criteria (RDoC): Toward a New Classification Framework for Research on Mental Disorders. *Am J Psychiat.* 2010;167:748-751.
37. Ingvar B, A. LS, A. DA, Arnstein M, Eystein S, C. KH. A dimensional versus a categorical approach to diagnosis: Anxiety and depression in the HUNT 2 study. *Int J Methods Psychiatr Res.* 2009;18:128-137.
38. Watson D, Gamez W, Simms LJ. Basic dimensions of temperament and their relation to anxiety and depression: A symptom-based perspective. *J Res Pers.* 2005;39:46-66.
39. Calhoun GG, Tye KM. Resolving the neural circuits of anxiety. *Nat Neurosci.* 2015;18:1394-1404.
40. Robinson OJ, Krinsky M, Grillon C. The impact of induced anxiety on response inhibition. *Front Hum Neurosci.* 2013;7:69.
41. Garner M, Attwood A, Baldwin DS, James A, Munafò MR. Inhalation of 7.5% carbon dioxide increases threat processing in humans. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology.* 2011;36:1557-1562.
42. Garner M, Attwood A, Baldwin D, Munafò M. Inhalation of 7.5% carbon dioxide increases alerting and orienting attention network function. *Psychopharmacology.* 2012;223:67-73.
43. Costello CG. Two dimensional views of psychopathology. *Behaviour Research and Therapy.* 1994;32:391-402.
44. Greco JA, Liberzon I. Neuroimaging of Fear-Associated Learning. *Neuropsychopharmacology.* 2016;41:320-334.

45. Benkelfat C, Bradwejn J, Meyer E, Ellenbogen M, Milot S, Gjedde A, Evans A. Functional neuroanatomy of CCK4-induced anxiety in normal healthy volunteers. *Am J Psychiatry*. 1995;152:1180-1184.
46. Eser D, Leicht G, Lutz J, Wenninger S, Kirsch V, Schule C, Karch S, Baghai T, Pogarell O, Born C, Rupprecht R, Mulert C. Functional neuroanatomy of CCK-4-induced panic attacks in healthy volunteers. *Hum Brain Mapp*. 2009;511-522.
47. Bailey JE, Dawson GR, Dourish CT, Nutt DJ. Validating the inhalation of 7.5% CO₂ in healthy volunteers as a human experimental medicine: a model of generalized anxiety disorder (GAD). *Journal of Psychopharmacology*. 2011;25:1192-1198.
48. Roberson-Nay R, Gorlin EI, Beadel JR, Cash T, Vrana S, Teachman BA. Temporal stability of multiple response systems to 7.5% carbon dioxide challenge. *Biological Psychology*. 2017;124:111-118.
49. Papadopoulos A, Rich A, Nutt D, Bailey J. The effects of single dose anxiolytic medication on the CO₂ models of anxiety: differentiation of subjective and objective measures. *Journal of psychopharmacology*. 2010;24:649-656.
50. Bailey JE, Papadopoulos A, Seddon K, Nutt DJ. A comparison of the effects of a subtype selective and non-selective benzodiazepine receptor agonist in two CO₂ models of experimental human anxiety. *Journal of Psychopharmacology*. 2008;23:117-122.
51. Bailey JE, Argyropoulos SV, Kendrick AH, Nutt DJ. Behavioral and cardiovascular effects of 7.5% CO₂ in human volunteers. *Depress Anxiety*. 2005;21:18-25.
52. Bailey JE, Kendrick A, Diaper A, Potokar J, Nutt DJ. A validation of the 7.5% CO₂ model of GAD using paroxetine and lorazepam in healthy volunteers. *J Psychopharmacol*. 2007;21:42-49.
53. Grillon C, Pellowski M, Merikangas KR, Davis M. Darkness facilitates the acoustic startle in humans. *Biological Psychiatry*. 1997;42:453-460.
54. Grillon C, Morgan CA, Davis M, Southwick SM. Effect of darkness on acoustic startle in Vietnam veterans with PTSD. *Am J Psychiatry*. 1998;155:812-817.
55. Kamkwalala A, Norrholm SD, Poole JM, Brown A, Donley S, Duncan E, Bradley B, Ressler KJ, Jovanovic T. Dark-Enhanced Startle Responses and Heart Rate Variability in a Traumatized Civilian Sample: Putative Sex-Specific Correlates of Posttraumatic Stress Disorder. *Psychosomatic Medicine*. 2012;74:153-159.
56. Grillon C, Ameli R, Woods SW, Merikangas K, Davis M. Fear-potentiated startle in humans: effects of anticipatory anxiety on the acoustic blink reflex. *Psychophysiology*. 1991;28:588-595.
57. Schmitz A, Grillon C. Assessing fear and anxiety in humans using threat of predictable and unpredictable aversive events (the NPU-threat test). *Nature Protoc*. 2012;7:527-532.
58. Shankman SA, Robison-Andrew EJ, Nelson BD, Altman SE, Campbell ML. Effects of predictability of shock timing and intensity on aversive responses. *International Journal of Psychophysiology*. 2011;80:112-118.

59. Seddon K, Morris K, Bailey J, Potokar J, Rich A, Wilson S, Bettica P, Nutt DJ. Effects of 7.5% CO₂ challenge in generalized anxiety disorder. *Journal of Psychopharmacology*. 2011;25:43-51.
60. Grillon C, Pine DS, Lissek S, Rabin S, Bonne O, Vythilingam M. Increased anxiety during anticipation of unpredictable aversive stimuli in posttraumatic stress disorder but not in generalized anxiety disorder. *Biol Psychiatry*. 2009;66:47-53.
61. Grillon C, Lissek S, Rabin S, McDowell D, Dvir S, Pine DS. Increased anxiety during anticipation of unpredictable but not predictable aversive stimuli as a psychophysiological marker of panic disorder. *Am J Psychiatry*. 2008;165:898-904.
62. Gorka S, Liu H, Sarapas C, Shankman S. Time course of threat responding in panic disorder and depression. *International journal of psychophysiology*. 2015;98:87-94.
63. Gorka SM, Nelson BD, Shankman SA. Startle response to unpredictable threat in comorbid panic disorder and alcohol dependence. *Drug and Alcohol Dependence*. 2013;132:216-222.
64. Gorka SM, Hee D, Lieberman L, Mittal VA, Phan KL, Shankman SA. Reactivity to uncertain threat as a familial vulnerability factor for alcohol use disorder. *Psychological Medicine*. 2016:1-10.
65. Gorka SM, Lieberman L, Phan KL, Shankman SA. Startle Potentiation to Uncertain Threat as a Psychophysiological Indicator of Fear-Based Psychopathology: An Examination Across Multiple Internalizing Disorders. *J Abn Psychol*. 2017;126:8-18.
66. Moberg CA, Bradford DE, Kaye JT, Curtin JJ. Increased startle potentiation to unpredictable stressors in alcohol dependence: Possible stress neuroadaptation in humans. *Journal of Abnormal Psychology*. 2017;126:441-453.
67. Figel B, Brinkmann L, Buff C, Heitmann CY, Hofmann D, Bruchmann M, Becker MPI, Herrmann MJ, Straube T. Phasic amygdala and BNST activation during the anticipation of temporally unpredictable social observation in social anxiety disorder patients. *NeuroImage: Clinical*. 2019;22:101735.
68. Buff C, Brinkmann L, Bruchmann M, Becker MPI, Tupak S, Herrmann MJ, Straube T. Activity alterations in the bed nucleus of the stria terminalis and amygdala during threat anticipation in generalized anxiety disorder. *Social Cognitive and Affective Neuroscience*. 2017;12:1766-1774.
69. Brinkmann L, Buff C, Neumeister P, Tupak SV, Becker MPI, Herrmann MJ, Straube T. Dissociation between amygdala and bed nucleus of the stria terminalis during threat anticipation in female post-traumatic stress disorder patients. *Human Brain Mapping*. 2017;38:2190-2205.
70. Brinkmann L, Buff C, Feldker K, Tupak SV, Becker MPI, Herrmann MJ, Straube T. Distinct phasic and sustained brain responses and connectivity of amygdala and bed nucleus of the stria terminalis during threat anticipation in panic disorder. *Psychological Medicine*. 2017:1-14.
71. Herrmann MJ, Boehme S, Becker MPI, Tupak SV, Guhn A, Schmidt B, Brinkmann L, Straube T. Phasic and sustained brain responses in the amygdala and the bed nucleus of the stria terminalis during threat anticipation. *Human Brain Mapping*. 2016;37:1091-1102.

72. Alvarez R, Lissek S, Biggs A, Grillon C: Contextual fear conditioning in a virtual reality environment: A fMRI study. in *The Organization for Human Brain Mapping*. Chicago, IL2007.
73. Alvarez RP, Biggs A, Chen G, Pine DS, Grillon C. Contextual fear conditioning in humans: cortical-hippocampal and amygdala contributions. *J Neuroscience*. 2008;28:6211-6219.
74. Alvarez RP, Chen G, Bodurka J, Kaplan R, Grillon C. Phasic and sustained fear in humans elicits distinct patterns of brain activity. *Neuroimage*. 2011;1:389-400.
75. McMenamin BW, Langeslag SJE, Sirbu M, Padmala S, Pessoa L. Network organization unfolds over time during periods of anxious anticipation. *The journal of neuroscience*. 2014;34:11261-11273.
76. McMenamin BW, Pessoa L. Discovering networks altered by potential threat (“anxiety”) using quadratic discriminant analysis. *NeuroImage*. 2015;116:1-9.
77. Bijsterbosch J, Smith S, Bishop SJ. Functional connectivity under anticipation of shock: correlates of trait anxious affect versus induced anxiety. *Journal of Cognitive Neuroscience*. 2015:1-14.
78. Pessoa L. A Network Model of the Emotional Brain. *Trends in Cognitive Sciences*. 2017;21:357-371.
79. Baars BJ, Franklin S. How conscious experience and working memory interact. *Trends in Cognitive Sciences*. 2003;7:166-172.
80. Mathews A, Mackintosh BA. A cognitive model of selective processing in anxiety. *Cognitive therapy and research*. 1998;22:539-560.
81. Eysenck MW, Derakshan N, Santos R, Calvo MG. Anxiety and cognitive performance: attentional control theory. *Emotion*. 2007;7:336-353.
82. Kopp CB. Antecedents of self-regulation: A developmental perspective. *Developmental Psychology*. 1982;18:199-214.
83. Hofmann W, Schmeichel BJ, Baddeley AD. Executive functions and self-regulation. *Trends in cognitive sciences*. 2012;16:174-180.
84. Hartley CA, Phelps EA. Anxiety and decision-making. *Biological Psychiatry*. 2012;72:113-118.
85. Derryberry D, Reed MA. Anxiety-related attentional biases and their regulation by attentional control. *Journal of Abnormal Psychology*. 2002;111:225-236.
86. Bar-Haim Y, Lamy D, Pergamin L, Bakermans-Kranenburg MJ, van IMH. Threat-related attentional bias in anxious and nonanxious individuals: a meta-analytic study. *Psychol Bull*. 2007;133:1-24.
87. Bishop SJ. Trait anxiety and impoverished prefrontal control of attention. *Nature Neuroscience*. 2009;12:92-98.
88. Walters E, Carew T, Kandel E. Associative Learning in *Aplysia*: evidence for conditioned fear in an invertebrate. *Science*. 1981;211:504-506.

89. Davis M. The role of the amygdala in fear-potentiated startle: implication for animal models of anxiety. *Trends in Pharmacological Sciences*. 1992;13:35-41.
90. Kuhn M, Wendt J, Sjouwerman R, Büchel C, Hamm A, Lonsdorf TB. The neurofunctional basis of affective startle modulation in humans - evidence from combined facial EMG-fMRI. *bioRxiv*. 2019:567032.
91. Beck AT, Clark DA. An information processing model of anxiety: automatic and strategic processes. *Behav Res Ther*. 1997;35:49-58.
92. Robinson OJ, Lieberman L, Allen P, Vytal K, Grillon C. The dorsal medial prefrontal-amygdala 'aversive amplification' circuit in unmedicated generalized and social anxiety disorders. *The Lancet Psychiatry*. 2014;1:294-302.
93. Borkovec TD, Alcaine OM, Behar E: Avoidance theory of worry and generalized anxiety disorder. in *Generalized anxiety disorder: advances in research and practice*. Edited by Heimberg RG, Turk CL, Mennin DS. New York, Guilford; 2004. pp. 77–108.
94. Morris LW, Liebert RM. Effects of negative feedback, threat of shock, and level of trait anxiety on the arousal of two components of anxiety. *Journal of Counseling Psychology*. 1973;20:321-326.
95. Mennin DS, Heimberg RG, Turk CL, Fresco DM. Applying an Emotion Regulation Framework to Integrative Approaches to Generalized Anxiety Disorder. 2002;9:85-90.
96. Kagan J, Reznick JS, Snidman N. Biological basis of childhood shyness. *Science*. 1988;240:167-171.
97. Dymond S. Overcoming avoidance in anxiety disorders: The contributions of Pavlovian and operant avoidance extinction methods. *Neuroscience & Biobehavioral Reviews*. 2019;98:61-70.
98. Cornwell BR, Garrido MI, Overstreet C, Pine DS, Grillon C. The un-predictive brain under threat: a neuro-computational account of anxious hypervigilance. *Biol Psychiatry*. 2017;82:447-454.
99. Cornwell BR, Baas JMP, Johnson L, Holroyd T, Carver FW, Lissek S, Grillon C. Neural responses to auditory stimulus deviance under threat of electric shock revealed by spatially-filtered magnetoencephalography. *Neuroimage*. 2007;37:282-289.
100. Weinberg A, Klein DN, Hajcak G. Increased error-related brain activity distinguishes generalized anxiety disorder with and without comorbid major depressive disorder. *Journal of Abnormal Psychology*. 2012;121:885-896.
101. Insel T, Cuthbert B, Garvey M, Heinssen R, Kozak M, Pine DS, Quinn K, Sanislow C, Steinberg J, Wang P. Research Domain Criteria (RDoC): toward a new classification framework for research on mental disorders. *Am J Psychiat*. 2010;167.
102. Criteria NAMHCWoTaMfRD: Behavioral Assessment Methods for RDoC Constructs. in *A Report by the National Advisory Mental Health Council Workgroup on Tasks and Measures for Research Domain Criteria (RDoC)*. Bethesda, MD, Z:\PDF_3\Anxiety-disorders\RDoC; 2016. pp. pp. 163.

103. Green MF, Nuechterlein KH, Gold JM, Barch DM, Cohen J, Essock S, Fenton WS, Frese F, Goldberg TE, Heaton RK, Keefe RSE, Kern RS, Kraemer H, Stover E, Weinberger DR, Zalcman S, Marder SR. Approaching a consensus cognitive battery for clinical trials in schizophrenia: The NIMH-MATRICES conference to select cognitive domains and test criteria. *Biological Psychiatry*. 2004;56:301-307.
104. Brovelli A, Ding M, Ledberg A, Chen Y, Nakamura R, Bressler SL. Beta oscillations in a large-scale sensorimotor cortical network: directional influences revealed by Granger causality. *Proceedings of the National Academy of Sciences of the United States of America*. 2004;101:9849-9854.
105. Friston KJ, Harrison L, Penny W. Dynamic causal modelling. *NeuroImage*. 2003;19:1273-1302.
106. Friston K, Moran R, Seth AK. Analysing connectivity with Granger causality and dynamic causal modelling. *Current opinion in neurobiology*. 2013;23:172-178.
107. Robinson OJ, Bond RL, Roisier JP. The impact of threat of shock on the framing effect and temporal discounting: executive functions unperturbed by acute stress? *Frontiers in Psychology*. 2015;6.
108. Charpentier CJ, Hindocha C, Roiser JP, Robinson OJ. Anxiety promotes memory for mood-congruent faces but does not alter loss aversion. *Scientific Reports*. 2016;6:24746.
109. Sussman TJ, Szekely A, Hajcak G, Mohanty A. It's all in the anticipation: How perception of threat is enhanced in anxiety. *Emotion*. 2016;16:320-327.
110. Weymar M, Bradley MM, Hamm AO, Lang PJ. Encoding and reinstatement of threat: Recognition potentials. *Neurobiology of Learning and Memory*. 2014;107:87-92.
111. Yang Y, Miskovich TA, Larson CL. State anxiety impairs proactive but enhances reactive control. *Frontiers in Psychology*. 2018;9.
112. Clarke RJ, Johnstone T. Prefrontal inhibition of threat processing protects working memory from interference. *Frontiers in Human Neuroscience*. 2013;7.
113. Pessoa L, Padmala S, Kenzer A, Bauer A. Interactions between cognition and emotion during response inhibition. *Emotion*. 2012;12:192-197.
114. Choi J, Padmala S, Pessoa L. Impact of state anxiety on the interaction between threat monitoring and cognition. *Neuroimage*. 2012;59:1912-1923.
115. Robinson OJ, Vytal K, Cornwell BR, Grillon C. The impact of anxiety upon cognition: perspectives from human threat of shock studies. *Front Hum Neurosci*. 2013;7:203:203.
116. Bailey JE, Papadopoulos A, Lingford-Hughes A, Nutt DJ. D-Cycloserine and performance under different states of anxiety in healthy volunteers. *Psychopharmacology*. 2007;193:579-585.
117. Cooper R, Howard CJ, Attwood AS, Stirland R, Rostant V, Renton L, Goodwin C, Munafò MR. Acutely induced anxiety increases negative interpretations of events in a closed-circuit television monitoring task. *Cognition and Emotion*. 2013;27:273-282.

118. Shin LM, Liberzon I. The neurocircuitry of fear, stress, and anxiety disorders. *Neuropsychopharmacology*. 2009.
119. Forster GL, Simons RL, Baugh LA: Revisiting the role of the amygdala in posttraumatic stress disorder. in *The amygdala: Where emotions shape perception, learning and memories*. Edited by Ferry B. Rijeka, Croatia, InTech; 2017. pp. 113-135.
120. Lang PJ, McTeague LM, Bradley MM. RDoC, DSM, and the reflex physiology of fear: A biodimensional analysis of the anxiety disorders spectrum. *Psychophysiology*. 2016;53:336-347.
121. Bradford DE, Shapiro BL, Curtin JJ. How bad could It be? Alcohol dampens stress responses to threat of uncertain intensity. *Psychological science*. 2013;24:2541-2549.
122. Pringle A, Warren M, Gottwald J, Cowen PJ, Harmer CJ. Cognitive mechanisms of diazepam administration: a healthy volunteer model of emotional processing. *Psychopharmacology*. 2016;233:2221-2228.
123. Murphy SE, Yiend J, Lester KJ, Cowen PJ, Harmer CJ. Short-term serotonergic but not noradrenergic antidepressant administration reduces attentional vigilance to threat in healthy volunteers. *International Journal of Neuropsychopharmacology*. 2009;12:169-179.
124. Valdez GR. Development of CRF1 receptor antagonists as antidepressants and anxiolytics: progress to date. *CNS Drugs*. 2006;20:887-896.
125. Risbrough VB, Stein MB. Role of corticotropin releasing factor in anxiety disorders: a translational research perspective. *Horm Behav*. 2006;50:550-561.
126. Walker D, Yang Y, Ratti E, Corsi M, Trist D, Davis M. Differential Effects of the CRF-R1 Antagonist GSK876008 on Fear-Potentiated, Light- and CRF-Enhanced Startle Suggest Preferential Involvement in Sustained vs Phasic Threat Responses. *Neuropsychopharmacol*. 2009;34:1533-1542.
127. Griebel G, Holsboer F. Neuropeptide receptor ligands as drugs for psychiatric diseases: the end of the beginning? *Nat Rev Drug Discov*. 2012;11:462-478.
128. Schwandt ML, Cortes CR, Kwako LE, George DT, Momenan R, Sinha R, Grigoriadis DE, Pich EM, Leggio L, Heilig M. The CRF1 antagonist verucerfont in anxious alcohol-dependent women: translation of neuroendocrine, but not of anti-craving effects. *Neuropsychopharmacology*. 2016;41:2818-2829.
129. Dunlop BW, Binder EB, Iosifescu D, Mathew SJ, Neylan TC, Pape JC, Carrillo-Roa T, Green C, Kinkead B, Grigoriadis D, Rothbaum BO, Nemeroff CB, Mayberg HS. Corticotropin-Releasing Factor Type 1 Receptor Antagonism is Ineffective for Women with Posttraumatic Stress Disorder. *Biological Psychiatry*. 2017;82:866-874.
130. Lago TR, Hsiung A, Leutner BL, Duckworth CJ, Balderston NL, Chen KY, Grillon C, Ernst M. Exercise modulates the interaction between cognition and anxiety in humans. *Cognition & Emotion*. in press.
131. Gorka SM, Lieberman L, Klumpp H, Kinney KL, Kennedy AE, Ajilore O, Francis J, Duffecy J, Craske MG, Nathan J, Langenecker S, Shankman SA, Phan KL. Reactivity to

- unpredictable threat as a treatment target for fear-based anxiety disorders. *Psychological Medicine*. 2017;1-11.
132. Costello CG. Anxiety and the persisting novelty of input from the autonomic nervous system. *Behavior Therapy*. 1971;2:321-333.
133. Gray JA: *The psychology of fear and stress*. Cambridge, England, Cambridge University Press; 1987.
134. Ludewig S, Geyer MA, Ramseier M, Vollenweider FX, Rechsteiner E, Cattapan-Ludewig K. Information-processing deficits and cognitive dysfunction in panic disorder. *J Psychiatry Neurosci*. 2005;30:37-43.
135. Ludewig S, Ludewig K, Geyer MA, Hell D, Vollenweider FX. Prepulse inhibition deficits in patients with panic disorder. *Depress Anxiety*. 2002;15:55-60.
136. Morgan III CA, Grillon C. Abnormal mismatch negativity in women with sexual assault-related posttraumatic stress disorder. *Biol Psychiatry*. 1999;45:827-832.
137. Ge Y, Wu J, Sun X, Zhang K. Enhanced mismatch negativity in adolescents with posttraumatic stress disorder (PTSD). *Int J Psychophysiol*. 2011;79:231-235.
138. Chang Y, Xu J, Pang X, Sun Y, Zheng Y, Liu Y. Mismatch negativity indices of enhanced preattentive automatic processing in panic disorder as measured by a multi-feature paradigm. *Biol Psychol*. 2015;105:77-82.
139. Wessel JR, Aron AR. On the Globality of Motor Suppression: Unexpected Events and Their Influence on Behavior and Cognition. *Neuron*. 2017;93:259-280.
140. Näätänen R, Kujala T, Winkler I. Auditory processing that leads to conscious perception: A unique window to central auditory processing opened by the mismatch negativity and related responses. *Psychophysiology*. 2011;48:4-22.
141. Näätänen R, Paavilainen P, Titinen H, Jiang D, Alho K. Attention and mismatch negativity. *Psychophysiology*. 1993;30:436-450.
142. Fan L, Sun Y-B, Sun Z-K, Wang N, Luo F, Yu F, Wang J-Y. Modulation of auditory sensory memory by chronic clinical pain and acute experimental pain: a mismatch negativity study. *Scientific Reports*. 2018;8:15673.
143. Gené-Cos N, Ring HA, Pottinger RC, Barrett G. Possible Roles for Mismatch Negativity in Neuropsychiatry. *Cognitive and Behavioral Neurology*. 1999;12:17-27.
144. Mager R, Bullinger AH, Mueller-Spahn F, Kuntze MF, Stoermer R. Real-time monitoring of brain activity in patients with specific phobia during exposure therapy, employing a stereoscopic virtual environment. *Cyberpsychol Behav*. 2001;4:465-469.
145. Marshall PJ, Reeb BC, Fox NA. Electrophysiological responses to auditory novelty in temperamentally different 9-month-old infants. *Developmental science*. 2009;12:568-582.
146. Giard MH, Perrin F, Pernier J, Bouchet P. Brain generators implicated in the processing of auditory stimulus deviance: a topographic event-related potential study. *Psychophysiology*. 1990;27:6627-6640.

147. Garrido MI, Kilner JM, Stephan KE, Friston KJ. The mismatch negativity: A review of underlying mechanisms. *Clinical Neurophysiology*. 2009;120:453-463.
148. Davis M. The role of the amygdala in fear and anxiety. *Annual Review of Neuroscience*. 1992;15:353-375.
149. Vuilleumier P. How brains beware: neural mechanisms of emotional attention. *Trends Cogn Sci*. 2005;9:585-594.
150. Wessel JR, Danielmeier C, Morton JB, Ullsperger M. Surprise and error: common neuronal architecture for the processing of errors and novelty. *J Neuroscience*. 2012;32:7528-7537.
151. Ousdal OT, Andreassen OA, Server A, Jensen J. Increased Amygdala and Visual Cortex Activity and Functional Connectivity towards Stimulus Novelty Is Associated with State Anxiety. *PLOS ONE*. 2014;9:e96146.
152. D'Hulst C, Atack JR, Kooy RF. The complexity of the GABAA receptor shapes unique pharmacological profiles. *Drug Discovery Today*. 2009;14:866-875.
153. Pellow s, File SE. Multiple sites of action for anxiogenic drugs: behavioural, electrophysiological and biochemical correlations. *Psychopharmacol*. 1984;83:304-315.
154. Smolnik R, Pietrowsky R, Fehm HL, Born J. Enhanced Selective Attention After Low-Dose Administration of the Benzodiazepine Antagonist Flumazenil. *Journal of Clinical Psychopharmacology*. 1998;18:241-247.
155. Nakagome K, Ichikawa I, Kanno O, Akaho R, Suzuki M, Takazawa S, Watanabe H, Kazamatsuri H. Overnight Effects of Triazolam on Cognitive Function: An Event-Related Potentials Study. *Neuropsychobiology*. 1998;38:232-240.
156. Rosburg T, Marinou V, Haueisen J, Smesny S, Sauer H. Effects of lorazepam on the neuromagnetic mismatch negativity (MMNm) and auditory evoked field component N100m. *Neuropsychopharmacology*. 2004;29:1723-1733.
157. Grillon C, Charney DR. In the face of fear: anxiety sensitizes defensive responses to fearful faces. *Psychophysiology*. 2011;12:1745-1752.
158. Morris JS, DeGelder B, Weiskrantz L, Dolan RJ. Differential extrageniculostriate and amygdala responses to presentation of emotional faces in a cortically blind field. *Brain*. 2001;124:1241-1252.
159. Whalen PJ, Rauch SL, Etcoff NL, McInerney SC, Lee MB, Jenike MA. Masked presentations of emotional facial expressions modulate amygdala activity without explicit knowledge. *J Neurosci*. 1998;18:411-418.
160. Bishop S, Duncan J, Brett M, Lawrence AD. Prefrontal cortical function and anxiety: controlling attention to threat-related stimuli. *Nature neuroscience*. 2004;7:184-188.
161. Dolcos F, McCarthy G. Brain systems mediating cognitive interference by emotional distraction. *J Neurosci*. 2006;26:2072-2079.

162. MacDonald AW, Cohen JD, Stenger VA, Carter CS. Dissociating the role of the dorsolateral prefrontal and anterior cingulate cortex in cognitive control. *Science*. 2000;288:1835-1838.
163. Robinson OJ, Letkiewicz AM, Overstreet C, Ernst M, Grillon C. The effect of induced anxiety on cognition: threat of shock enhances aversive processing in healthy individuals. *Cogn Affect Behav Neurosci*. 2011;11:217–227.
164. Robinson OJ, Charney DR, Overstreet C, Vytal K, Grillon C. The adaptive threat bias in anxiety: amygdala-dorsomedial prefrontal cortex coupling and aversive amplification. *Neuroimage*. 2012;60:523-529.
165. Etkin A, Egner T, Kalisch R. Emotional processing in anterior cingulate and medial prefrontal cortex. *Trends in cognitive sciences*. 2011;15:85-93.
166. Vytal K, Overstreet C, Charney D, Robinson OJ, Grillon C. Sustained anxiety increases amygdala-dorsomedial prefrontal coupling: A mechanism for maintaining an anxious state. *J Psychiatry Neurosci*. 2014;39:321-329.
167. Robinson OJ, Krinsky M, Lieberman L, Vytal K, Ernst M, Grillon C. Anxiety-potentiated amygdala-medial frontal coupling and attentional control. *Transl Psychiatry*. 2016;6:e833.
168. Robinson OJ, Cools R, Crockett MJ, Sahakian BJ. Mood state moderates the role of serotonin in cognitive biases *Journal of Psychopharmacology*. 2010;24:573-583.
169. Harmer CJ. Serotonin and emotional processing: Does it help explain antidepressant drug action? *Neuropharmacology*. 2008;55:1023-1028.
170. Nutt DJ. The neuropharmacology of serotonin and noradrenaline in depression. *Int Clin Psychopharmacol*. 2002;17 (Suppl. 1):S1-S12.
171. Hariri AR, Mattay VS, Tessitore A, Kolachana B, Fera F, Goldman D, Egan MF, Weinberger DR. Serotonin transporter genetic variation and the response of the human amygdala. *Science*. 2002;297:400-403.
172. Robinson OJ, Overstreet C, Allen PS, Letkiewicz A, Vytal K, Pine DS, Grillon C. The role of serotonin in the neurocircuitry of negative affective bias: serotonergic modulation of the dorsal medial prefrontal-amygdala ‘aversive amplification’ circuit. *Neuroimage*. 2013;78:217-223.
173. Walsh AEL, Harmer CJ. The cognitive neuropsychological model of antidepressant response. *Current Opinion in Psychology*. 2015;4:124-130.
174. Butler AC, Chapman JE, Forman EM, Beck AT. The empirical status of cognitive-behavioral therapy: A review of meta-analyses. *Clinical Psychology Review*. 2006;26:17-31.
175. McLean CP, Asnaani A, Litz BT, Hofmann SG. Gender differences in anxiety disorders: Prevalence, course of illness, comorbidity and burden of illness. *Journal of Psychiatric Research*. 2011;45:1027-1035.
176. Patel N, Vytal K, Pavletic N, Stoodley C, Pine DS, Grillon C, Ernst M. Interaction of threat and verbal working-memory in adolescents. *Psychophysiology*. 2016;53:518-526.

177. Lieberman LC, Gorka SM, Sarapas C, Shankman SA. Cognitive flexibility mediates the relation between intolerance of uncertainty and safety signal responding in those with panic disorder. *Cogn Emot*. 2015.
178. Gorka SM, Lieberman L, Nelson BD, Sarapas C, Shankman SA. Aversive responding to safety signals in panic disorder: The moderating role of intolerance of uncertainty. *Journal of Anxiety Disorders*. 2014;28:731-736.
179. Roberson-Nay R, Beadel JR, Gorlin EI, Latendresse SJ, Teachman BA. Examining the latent class structure of CO2 hypersensitivity using time course trajectories of panic response systems. *Journal of Behavior Therapy and Experimental Psychiatry*. 2015;47:68-76.
180. Nay WT, Thorpe GL, Roberson-Nay R, Hecker JE, Sigmon ST. Attentional bias to threat and emotional response to biological challenge. *J Anxiety Disord*. 2004;18:609-627.
181. Kane MJ, Engle RW. The role of prefrontal cortex in working-memory capacity, executive attention, and general fluid intelligence: An individual-differences perspective. *Psychonomic Bulletin & Review*. 2002;9:637-671.
182. Insel TR. Translating scientific opportunity into public health impact: A strategic plan for research on mental illness. *Archives of General Psychiatry*. 2009;66:128-133.
183. Blanchard RJ, Blanchard DC: Anti-predator defense as models of fear and anxiety. in *Fear and Defense*. Edited by Brain PF, Blanchard RJ, Parmigiani S. London 1990. pp. 89-108.
184. Blanchard RJ, McKittrick CR, Blanchard DC. Animal models of social stress: effects on behavior and brain neurochemical systems. *Physiology & Behavior*. 2001;73:261-271.
185. Kogler L, Müller VI, Chang A, Eickhoff SB, Fox PT, Gur RC, Derntl B. Psychosocial versus physiological stress — Meta-analyses on deactivations and activations of the neural correlates of stress reactions. *NeuroImage*. 2015;119:235-251.
186. Kaye JT, Bradford DE, Curtin JJ. Psychometric properties of startle and corrugator response in NPU, affective picture viewing, and resting state tasks. *Psychophysiology*. 2016;53:1241-1255.
187. Baldwin DS, Hou R, Gordon R, Huneke NTM, Garner M. Pharmacotherapy in generalized anxiety disorder: novel experimental medicine models and emerging drug targets. *CNS Drugs*. 2017;31:307-317.
188. Price M, Anderson PL. Latent growth curve analysis of fear during a speech task before and after treatment for social phobia. *Behaviour research and therapy*. 2011;49:763-770.
189. Goldin PR, Manber T, Hakimi S, Canli T, Gross JJ. Neural bases of social anxiety disorder: Emotional reactivity and cognitive regulation during social and physical threat. *Archives of General Psychiatry*. 2009;66:170-180.
190. Biedermann SV, Biedermann DG, Wenzlaff F, Kurjak T, Nouri S, Auer MK, Wiedemann K, Briken P, Haaker J, Lonsdorf TB, Fuss J. An elevated plus-maze in mixed reality for studying human anxiety-related behavior. *BMC biology*. 2017;15:125-125.
191. Iordan AD, Dolcos S, Dolcos F. Brain activity and network interactions in the impact of internal emotional distraction. *Cerebral Cortex*. 2018:1-17.

192. Balderston NL, Liu L, Roberson-Nay R, Ernst M, Grillon C. The relationship between dlPFC activity during unpredictable threat and CO₂-induced panic symptoms. *Transl Psychiatry*. 2017;7:1266.

FIGURE LEGENDS

Fig. 1. Diagram illustrating the 5-step experimental approach. Step 1: Selection of a clinically-relevant construct (e.g., attention bias for threat) and tasks (e.g., dot-probe) that probe the construct; Step 2: characterization of the response (output) and underlying neurobiology associated with the task (input) probing the construct in healthy subjects. The response can be a behavioral, cognitive, a pharmacological, or electrophysiological measure. The region of interest ROI1, ROI2, and ROI3 represent 3 hypothetical structures that mediate the behavior tested by the task. The arrows represent hypothetical directional connectivity among these structures; Step 3: The task is combined with an anxiety induction procedure to determine how anxiety affects the response and underlying neurocircuits. In this example, a structure ROIx affects the connectivity between ROI1 and ROI2. In Step 4 the task is tested in patients to determine whether the same processes and neurocircuits affected by experimental anxiety in healthy subjects are also implicated in the patients. Patients can be tested with or without an anxiety induction procedure. Step 5: Healthy subjects are tested as in Step 3 (Task + induced-anxiety) but are also given a treatment to examine mechanisms of treatment responses or to screen putative anxiolytic treatments (do they affect the response and do they affect the same anxiety-neurocircuit identified in Step 2?).


Fig. 2. Left. The MMNm arises from bilateral sources over primary, secondary auditory cortex (superior temporal gyrus, STG), and inferior frontal gyri (IFG) (146). Right. Dynamic causal modeling shows that the MMN results from changes within and among these cortical sources (147). Forward connections can be conceptualized as bottom-up processes transmitting sensory information from A1 to higher cortical levels and convey prediction errors (MMN). Backward connections represent top-down predictions based on prior sensory experience and explain away prediction errors (deviance detection). In the safe context, the MMN is mediated by

changes in both extrinsic feedforward and feedback connectivity as well as intrinsic connectivity (not shown). Anxiety induced by threat of shock suppresses feedback connectivity. The anxiolytic alprazolam reestablishes feedback connectivity (not shown).

Fig. 3. Proposed model of attention bias for threat in anxiety. Increased bi-directional coupling between the dorsal anterior cingulate/dorsomedial prefrontal cortex (dACC/dmPFC) and amygdala (amg) promotes threat bias. Coupling is transiently activated during anxiety states in healthy control, facilitating the detection of threat. However, the coupling is chronically engaged in pathological anxiety, leading to maladaptive threat bias. Lowering serotonin has similar effect as induced-anxiety: it increases dACC/dmPFC-amygdala coupling. By increasing serotonin, SSRIs may reduce attention bias for threat via inhibition of dACC/dmPFC-amygdala coupling.

Healthy subjects
performing a cognitive task

No anxiety-induction



Anxiety-induction

