

Height and timing of growth spurt during puberty in young people living with vertically acquired HIV in Europe and Thailand.**Running title (40/40 characters):** Pubertal growth in young people with HIV

Group authorship: The European Pregnancy and Paediatric HIV Cohort Collaboration (EPPICC) study group *

*Please see Acknowledgements section for list of Writing Group members

*Authors

Siobhan CRICHTON¹Erik BELFRAGE²Intira Jeannie COLLINS¹Katja DOERHOLT³Ali JUDD¹Sophie LE COEUR⁴Vana SPOULOU⁵Henriette SCHERPBIER⁶Colette SMIT⁷Tessa GOETGHEBUER⁸Diana M. GIBB¹Antoni NOGUERA-JULIAN⁹Maria Luisa NAVARRO¹⁰Jose Tomas RAMOS¹¹

This is an open access article distributed under the Creative Commons Attribution License 4.0, (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Luisa GALLI¹²

Carlo GIAQUINTO¹³

Claire THORNE¹⁴

Santa ANSONE¹⁵

Magdalena MARCZYNSKA¹⁶

Liubov OKHONSKAIA¹⁷

Begoña MARTINEZ DE TEJADA¹⁸

Gonzague JOURDAIN¹⁹

Luc DECKER¹⁹

Luminita ENE²⁰

Ruth GOODALL¹

1 MRC Clinical Trials Unit at University College London, London, UK

2 Department of Pediatrics, Karolinska Institutet and University Hospital, Stockholm, Sweden

3 St George's University Hospitals NHS Foundation Trust, London, UK

4 French Institute for Demographic Studies (INED), Paris, France

5 Department of Infectious Diseases, University of Athens, Athens, Greece

6 Amsterdam University Medical Centers, Netherlands

7 HIV Monitoring Foundation, Amsterdam, Netherlands

8 Hopital St Pierre, Brussels, Belgium

9 Unitat d'Infectologia, Servei de Pediatria, Hospital Sant Joan de Deu, Universitat de Barcelona, Barcelona, Spain

10 Hospital General Universitario "Gregorio Marañón", Complutense University, Madrid, Spain

11 Paediatrics Department, Hospital Clínico Universitario San Carlos, Complutense University, Madrid, Spain

12 Department of Health Sciences, University of Florence, Meyer Children's University Hospital, Italy

13 Paediatric European Network for the Treatment of AIDS (PENTA), Padova, Italy

14 UCL Great Ormond Street Institute of Child Health, UCL, London, UK

15 Riga East University Hospital "Latvian Centre on Infectious Diseases", Latvia

16 Medical University of Warsaw, Hospital of Infectious Diseases, Warsaw, Poland

17 Republican Hospital of Infectious Diseases, St Petersburg, Russia;

18 Department of Obstetrics and Gynecology, and Faculty of Medicine, Geneva, Switzerland

19 Institut de Recherche pour le Developpement (IRD), UMI 174/PHPT, Chiang Mai,

Thailand;

20 Victor Babes Hospital, Bucharest, Romania

Corresponding Author: Siobhan Crichton

MRC Clinical Trials Unit at UCL, 90 High Holborn, London WC1V 6LJ.

Telephone no: +44 (0)20 7670-4913. E-mail: s.crichton@ucl.ac.uk

Funding: EPPICC receives funding from the PENTA Foundation (<http://penta-id.org>). The

MRC Clinical Trials Unit at UCL is supported by the Medical Research Council (programme number MC_UU_12023/26).

Conflicts of interest: None

ACCEPTED

Abstract

Objective: To describe growth during puberty in young people with vertically acquired HIV.

Design: Pooled data from 12 paediatric HIV cohorts in Europe and Thailand.

Methods: 1094 children initiating an NNRTI- or boosted PI-based regimen aged 1 to 10 years were included. Super Imposition by Translation And Rotation (SITAR) models described growth from age 8 years using 3 parameters (average height, timing and shape of the growth spurt), dependent on age and height-for-age z-score (HAZ) (World Health Organisation references) at ART initiation. Multivariate regression explored characteristics associated with these 3 parameters.

Results: At ART initiation median age and HAZ was 6.4[IQR:2.8,9.0] years and -1.2[-2.3,-0.2], respectively. Median follow-up was 9.1[6.9,11.4] years. In females, older age and lower HAZ at ART initiation were independently associated with a growth spurt which occurred 0.41(95% CI 0.20,0.62) years later in children starting ART age 6-<11 years compared to 1-<3 years and 1.50(1.21,1.78) years later in those starting with HAZ<-3 compared to HAZ \geq -1. Later growth spurts in females resulted in continued height growth into later adolescence. In males starting ART with HAZ<-1, growth spurts were later in children starting ART in the oldest age group, but for HAZ \geq -1 there was no association with age. Males and females who initiated ART with HAZ \geq -1 maintained a similar height to the WHO reference mean.

Conclusions: Stunting at ART initiation was associated with later growth spurts in females. Children with HAZ \geq -1 at ART initiation grew in height at the level expected in HIV negative children of a comparable age.

Key words: Growth, height, HIV, perinatal, puberty, Europe, Thailand

Introduction

Although young people living with HIV are at risk for poor height growth[1], treatment with antiretroviral therapy (ART) improves growth, with strongest gains in those treated at a young age[2]. While initial catch-up growth on ART has been well described[2] there are less data on long term growth, particularly during adolescence.

Delays in pubertal development have been reported in young people with HIV[3-7] with the onset of puberty[5] and sexual maturation[6] occurring 6 months later compared to HIV-exposed uninfected young people (HEU). Earlier puberty in the general population is associated with being taller and having higher BMI throughout childhood[8], and poor growth in children with PHIV has been shown to account for much of the delay in reaching sexual maturity[6]. There is also evidence that children starting ART with low height-for-age z-scores experience delays in the onset of puberty independently of age at ART initiation[3].

Poor growth during childhood can have implications for future health. Height velocity is associated with increased HIV replication[9] and progression to AIDS and death[10] with the association with death being independent of age, viral load and CD4 count[11]. The timing of puberty is also inversely associated with bone mass and density among HIV-negative adolescents[12] and delayed puberty may increase future risk of osteoporosis among young people with HIV, who themselves are at risk of poor bone health, either caused by HIV infection itself or prolonged exposure to ART[13]. Early growth failure has also been linked to poorer social and economic outcomes in later life in the general population[14].

In this study, statistical models which describe an individual's growth in terms of mean height throughout adolescence, and timing and shape of the adolescent growth spurt were

applied to longitudinal height measurements. The overall aim of this study was to explore the association between characteristics at ART initiation, in particular age and height-for-age z-score, and growth during adolescence.

Methods

17 paediatric HIV cohorts from 15 countries contributed individual level data to the European Pregnancy and Paediatric HIV Cohort Collaboration (EPPICC) between September 2016 and March 2017 using a modified HICDEP protocol (www.hicdep.org). Pseudo-anonymised data on all children at participating clinics were included. All cohorts received approval from local and/or national ethical committees. Five cohorts from three countries (Italy, Ukraine and three from Russia) where height data were not routinely collected (each with <20% of children having a height measurement at ART initiation) were excluded. Children from the remaining 12 cohorts were eligible provided they initiated ART with ≥ 2 nucleoside reverse transcriptase inhibitors (NRTI) + a non-nucleoside reverse transcriptase inhibitor (NNRTI) or boosted protease inhibitor (bPI); were 1-10 years old at ART initiation; not known to have horizontally-acquired HIV; and aged ≥ 8 years at the end of follow-up. We excluded children initiating ART after age 11 years. For those initiating ART at an older age it would be difficult to distinguish between changes in growth occurring as a result of a pubertal growth spurt and as a result of initiating ART. Children with no height recorded at ART initiation and/or after 8 years of age were excluded.

Height measurements were censored at the earliest of: 19th birthday, transfer to adult care, death or loss to follow-up. Height and BMI were converted to height-for-age z-scores (HAZ) and BMI-for-age z-scores (zBMI), using the World Health Organization (WHO) Growth Standard for measurements when children were aged <5 years old[15] and the WHO 2007

growth reference when aged 5-18 years [15, 16]. Data checks were carried out to detect implausible changes in height and/or HAZ. HAZ was categorised according to WHO definitions as: <-3 SD (severe stunting); -3 to <-2 SD (stunting); -2 to <-1 SD; and ≥-1 SD. zBMI was categorised as: <-2 SD (underweight); -2 to 1 SD (normal); >1 to 2 SD (overweight); and >2 SD (obese). HAZ and zBMI nearest to ART initiation (closest within 6 months before to 1 month after) were considered baseline measurements.

Other variables included were: sex, country (Thailand, UK/Ireland, Rest of Europe), age at ART initiation (1-2, 3-5 or 6 -10 years), initial ART regimen (PI, NNRTI), being born outside the country of the cohort (“born abroad”), and immunodeficiency at ART initiation classified using the WHO immunological classification: none (CD4% $>35\%$, $>30\%$, $>25\%$ or CD4 count >500 cells/mm³ in children <1 , 1-2, 3-4 and ≥ 5 years, respectively), mild (CD4% 30-35%, 25-30%, 20-25% or CD4 count 350-499cells/mm³), advanced (CD4% 25-29%, 20-24%, 15-19% or CD4 count 200-349 cells/mm³) or severe (CD4% $<25\%$, $<20\%$, 15-19% or in children >5 CD4 count <200 cells/mm³ OR $<15\%$)[17].

Statistical analysis

Characteristics at ART initiation were summarised by HAZ category. Mean height at age 16 years was summarised by age and HAZ at ART initiation and compared to the WHO reference height to quantify differences in height following the growth spurt. It was not possible to assess differences in final height as many adolescents transfer to adult care from age 16 years, ending follow-up in EPPICC.

Height was modelled using Super Imposition by Translation And Rotation (SITAR) models [18]. SITAR was developed to model growth during childhood and adolescence and

quantifies differences in growth via three parameters representing the timing and shape of the adolescent growth spurt, as well as average height. The models can explain up to 99% of the variation between individuals' growth [18] and can be summarised as:

$$y_{it} = a_i + h\left(\frac{t - b_i}{\exp^{-c_i}}\right)$$

where the outcome y_{it} is the height of individual i at age t and $h()$ is a natural cubic spline of height over age. The parameters a_i , b_i and c_i are participant specific random effects. a_i represents average height throughout adolescence; negative values indicate shorter height overall. b_i represents timing of the pubertal growth spurt; negative values indicate earlier puberty. c_i represents growth velocity, or the shape of the growth spurt; positive values indicate shorter growth spurts and a steeper growth velocity curve while negative values indicate the growth spurt occurs over a longer duration. Corresponding growth velocity curves can also be estimated as the first derivative of the modelled growth (height) curve.

Age at peak height velocity (APHV) is correlated with timing of puberty and often used as a proxy for timing of maturation. It commonly occurs in girls in Tanner stage 2 or 3 and in Tanner stage 3 or 4 for boys[19, 20], though there is variation in timing across Tanner stages [19]. Differences in the timing of the growth spurt estimated using SITAR models have been shown to be highly correlated with APHV[18].

All height measurements (in centimetres) from age 8 (or start of ART if after 8th birthday) to 18 years were included. Age and HAZ at ART initiation were added to the SITAR model as fixed effects which could influence the mean of a , b and c . Thus the estimated random effects a_i , b_i and c_i represent the individual differences in average height, timing and shape of the growth spurt not associated with differences in age or height at ART initiation. Models were fitted separately to males and females using a spline with 6 degrees of freedom. Log

transformations of both age and height [18] were considered but the untransformed data provided the best fit. Interactions between baseline height and age were added where appropriate (model comparison carried out using Bayes Information Criteria (BIC)).

To explore other factors (sex, country, initial ART regimen, WHO immunological classification, zBMI at ART initiation) associated with growth after allowing for differences in baseline age and height, the estimated a_i , b_i and c_i random effects from the SITAR model were analysed using multivariable linear regression. Interactions between each of the factors and sex and between immunological classification and HAZ and age at ART initiation were considered. A second model was fitted including zBMI at age 8 years instead of at ART initiation.

Modelling was repeated in countries where >5% of children were born abroad and >5% born in the country (UK & Ireland, Spain and Netherlands) to explore differences between those born abroad and those born in the cohort country. Sensitivity analyses were carried out in which i) models were fitted separately to children from Thailand and elsewhere ii) Thai specific growth reference data were used for Thai children[21], and iii) children starting ART after age 8 years were excluded.

Analyses were carried out using STATA/IC 15.1 and the SITAR package[22] in R v3.3.3.

Results

Patient characteristics

In total 1943 young people with HIV initiated ART on an eligible regimen age 1- 10 years and were ≥ 8 years at end of follow up (Figure 1). After excluding those with missing baseline height (n=721) and/or height after age 8 years (n=202), we included 1094 children in the

analysis. Children excluded due to missing height data were more likely to be from countries other than Thailand or UK/Ireland, be born abroad, and be younger at ART initiation than those who were included (Supplementary Table 1). The 1094 included children were followed-up for a median of 9.1[6.9,11.4] years after ART initiation. During this time 37,325 height measurements were recorded with a median of 32[19, 46] per child, of which 25,458 (median 21[11,32]) were from age 8 years onwards. The median time between height measurements was 2.8[1.4,3.9] months, with some variation by cohort ranging from every 1.8[1.0,2.8] months in Thailand to 8.3[4.7,11.3] months in Greece.

At ART initiation median HAZ was -1.2[-2.3,-0.2] and age was 6.4[2.8,9.0] years. Characteristics of children at ART initiation, stratified by baseline HAZ, are described in Table 1. More severe stunting was associated with residence in Thailand, not being born abroad, initiating on an NNRTI based regimen, earlier calendar year of ART initiation, higher viral load, more severe immunodeficiency and lower zBMI at ART initiation.

At the end of the study 493(45%) children had reached their 16th birthday while still in paediatric care (Figure 1); of whom 463(94%) had their height recorded within 6 months of their birthday. Children who survived to age 16 years but were no longer in follow up in paediatric care were more likely to reside in Thailand and start ART at a younger age. At age 16 years the mean (standard deviation) heights of males and females were 166(8.7)cm and 158(6.9)cm respectively, significantly shorter than the WHO reference mean height of 173(7.8)cm for males and 163(6.8)cm for females (both $p < 0.001$) (Supplementary Table 2).

Associations between age and HAZ at ART initiation and growth from age 8 years

Results from the SITAR models are available in Supplementary Table 3. Estimated mean height and corresponding growth velocity curves stratified by HAZ and by age are summarised in Figure 2A and 2B, respectively, for females and Figure 3A and 3B for males.

In females, across each of the baseline HAZ groups (Figure 2Ai-iv), children starting ART in the oldest age group had growth spurts on average 0.41(95% CI 0.20,0.62) years later than those starting ART in the youngest age group. Across the baseline age groups (Figure 2Bi-iii), females starting ART with low HAZ had later growth spurts; there was a 1.50(1.21,1.78) year delay in those with baseline $HAZ < -3$ compared to baseline $HAZ \geq -1$. The effect of this delay on overall height can be seen in Figures 2L-N; the differences in height are smaller from age 16 onwards (after the growth spurt) than at age 8.

In males, the association between baseline age and the timing of the growth spurt differed by baseline HAZ (Figure 3Ai-iv); there was no significant difference by age in males who started ART with $HAZ \geq -1$ (Figure 3Ai). In males with baseline HAZ of -2 to < -1 (Figure 3Aii), the growth spurt was 0.96(0.19,1.72) years later in those starting ART in the oldest compared to youngest age group. Similarly, for a baseline HAZ of -3 to < -2 (Figure 3Aiii) the corresponding delay in those starting ART in the oldest age group was 0.92(0.17,1.66) years, and for baseline $HAZ < -3$ it was 0.42(-0.32,1.16) years (Figure 3Aiv). The timing of the growth spurt in males did not differ significantly by baseline HAZ (Figure 3Bi-iii).

Females (Figure 2Bi) and males (Figure 3Bi), who started treatment with a baseline $HAZ \geq -1$ maintained a similar mean height to the WHO reference, regardless of baseline age.

Other factors associated with growth from age 8 years

Characteristics associated with variations in growth that remained after adjusting for differences in baseline HAZ and age are summarised in Table 2. Young people from Thailand were smaller throughout adolescence than those from other countries, but did not differ in the timing of the growth spurt. The shape of the growth spurt differed by country and was shorter in children from the UK and Ireland than elsewhere. Lower zBMI at ART initiation was significantly associated with a later growth spurt (a one SD decrease was associated with a 0.07(0.02,0.11) year delay in the growth spurt). In a second model (data not shown) a one SD decrease in zBMI at age 8 years was associated with a 0.16(0.09,0.22) year delay in the timing of the growth spurt, while other parameters did not change substantially. There was no evidence of any interactions.

In subgroup analysis (n=545) there was a significant interaction between gender and being born abroad on timing of the growth spurt (p=0.038). Females born abroad experienced a growth spurt 0.24(0.02,0.46) years earlier than those born in the cohort country; while there was no association in males. However, after adjusting for zBMI at age 8, the association was no longer significant (growth spurt for females born abroad was 0.18(-0.05,0.42) years earlier)

In the three sensitivity analyses where 1) models were fitted separately to children from Thailand and elsewhere, 2) Thai specific reference data were used for Thai children and 3) children starting ART age \geq 8 years were excluded, overall conclusions were unchanged (data not shown).

Discussion

In this study we described growth throughout adolescence in a large cohort of young people with vertically acquired HIV in Europe and Thailand. Although all adolescents in the study initiated ART before age 11 years, growth deficits remained throughout adolescence. Only children with $HAZ \geq -1$ when starting ART were able to achieve a similar height to the WHO reference at age 16 years, suggesting that for others, catch up growth associated with being on ART long term was not sufficient to restore height to what would be expected in an HIV-negative population.

We observed an association between older age at ART initiation and later growth spurts in males (with $HAZ < -1$ at ART initiation) and females, in line with findings from the Antiretroviral research for Watoto (ARROW) where attainment of each tanner stage and onset of menarche was delayed in those starting ART at older ages[3]. We also observed an association between stunting and later growth spurts, but only in females. The potential role of anthropometric parameters in early childhood on growth during puberty was highlighted in a study of 2539 young people with vertically acquired HIV and HIV exposed uninfected (HEU) young people from the USA[6]. Young people living with HIV reached sexual maturity on average 6 months later than the HEU group, but differences in HAZ prior to puberty accounted for up to 98% of the delay in males and (together with $zBMI$) 74% in females, suggesting much of the delay may be attributable to earlier poor growth [6]. Low HAZ at ART initiation was also associated with delayed attainment of all Tanner stages in males and females, and menarche in females, independently of age at ART initiation in the ARROW trial [3]. However, in males, the delay was reduced in those who had the greatest initial gains in CD4 after starting ART, but there was no similar association in females. Undernutrition early in life was also found to have a stronger association with adult height in

females than males in the Netherlands [23]. While this suggests females may be more sensitive to impairments early in life, and prior to ART, the mechanism underlying potential gender differences remains to be explained.

After accounting for HAZ and age at ART initiation we found no association between WHO immunological status or viral load at ART initiation and growth. Similarly, the ARROW trial found immune suppression prior to ART was not associated with delayed puberty or menarche[3]. Other studies have also reported a lack of association between clinical status at start of puberty and age at onset[4, 7]. However, in young people in the USA, low CD4 and high VL at first pubertal assessment were associated with later pubertal onset. Among males, prior CDC C, low nadir CD4% or high peak VL were also associated with later puberty[5]. However many of these young people initiated ART on mono or dual therapy and are likely to have substantially different treatment histories compared to our study.

We found zBMI at ART initiation and age 8 to be associated with the timing of the growth spurt, with no evidence of a difference between males and females. We also observed that females born abroad experienced an earlier pubertal growth spurt than females born in the cohort country but the differences in the timing of the growth spurt reduced after adjusting for zBMI at age 8. In females a relationship between low BMI and delayed puberty has been found in multiple studies[8] and rapid weight gain prior to puberty also linked to early onset[24]. Differences between young people born abroad and those born in the country may therefore be explained by periods of more rapid weight gain in children arriving from abroad, the majority from Africa, compared to those born in the country.

This study had several limitations; as with all observational studies our findings on the association between age and HAZ at ART initiation and growth should not be over

interpreted or assumed to be causative. At ART initiation stunting was strongly correlated with immunosuppression, viral load and zBMI and may be a marker for poor immunological status and other impairments. Children starting ART at older ages represent a group who have survived without treatment and possibly with limited access to care and so may be subject to a survivor bias. Had ART initiation been delayed in those who started at a young age, the observed delay in the growth spurt associated with starting ART at an older age may have been less in this group who would also have been more likely to have access to health care and regular monitoring. Nonetheless the findings provide insight in to growth patterns among children presenting to care and starting ART at different ages.

Inclusion criteria applied also leads to the potential for selection bias. We excluded children with missing height data. Multiple imputation was not possible since other data, such as immunological and virological status, at ART initiation, likely to be strong predictors of baseline height were missing in more than half of the children with missing heights. We excluded young people from Russia, Ukraine and Italy where height data were not routinely recorded. Further the cohorts included in EPPICC range from national coverage to city hospitals leading to potential for bias where children treated in large city hospitals are not representative of others in the country. Our analyses were restricted to children aged 1-10 years at ART initiation. The number of infants initiating ART under age 1 year was small, with high rates of missing baseline data.. A further limitation is the lack of quantitative measures of pubertal status such as Tanner stage and date of onset of menarche, which is not routinely collected by the majority of participating cohorts. However, differences in timing of the growth spurt are likely to be indicative of differences in timing of onset of puberty.

Finally, we used the WHO growth standard[15] and growth reference[16] to derive z-scores at ART initiation. While the WHO growth standards were developed to assess growth globally, children from Thailand were significantly shorter than those residing in Europe and WHO reference may overestimate stunting as compared to Thailand's own national growth reference[25] . However, in sensitivity analyses, using Thai reference data we did not find any difference in the associations between baseline HAZ and growth during adolescence..

Despite these limitations, the study has several strengths. The collaborative nature of the study provides a rich source of longitudinal height measurements from a large sample of young people living with HIV followed during childhood and adolescence and the use of SITAR models provides insight into growth during puberty in the absence of quantitative measures of pubertal status.

In summary, we have shown that children who initiate ART at younger ages are taller. Children who initiate ART with a 'normal' height for age z-score ($HAZ \geq -1$) remained with a "normal" height throughout adolescence. Those who initiated ART stunted or severely stunted were less likely to achieve 'normal' height. We also demonstrated that in females, regardless of age at ART initiation, stunting at time of initiation was associated with a later pubertal growth spurt, and this continued growth into later adolescents may allow those most severely stunted to catch-up somewhat. However, longer-term follow-up is required to understand the potential implications of delayed pubertal growth on outcomes in later life.

Acknowledgements

We thank all the patients for their participation in these cohorts, and the staff members who cared for them.

Author contributors: Writing Group (consisting of Project Team first (ordered alphabetically by name except for the first and last author), and also other Writing Group members (ordered alphabetically by cohort name)):

Project Team: Siobhan Crichton (EPPICC statistician), Eric Belfrage (Karolinska Institutet and University Hospital, Stockholm, Sweden), Intira Jeanne Collins (Collaborative HIV Paediatric Study (CHIPS), UK and Ireland), Katja Doerholt (Collaborative HIV Paediatric Study (CHIPS), UK and Ireland), Ali Judd (co-lead of EPPICC), Sophie Le Coeur (Thailand Program for HIV Prevention and Treatment (PHPT) Study Group, Thailand), Vana Spoulou (Greece Cohort, Greece), Ruth Goodall (EPPICC senior statistician).

Other Writing Group members: Henriette Scherpbier, Colette Smit (ATHENA paediatric cohort, Netherlands); Tessa Goetghebuer (Hospital St Pierre Cohort, Brussels, Belgium); Diana M. Gibb (Collaborative HIV Paediatric Study (CHIPS) and National Study of HIV in Pregnancy and Childhood (NSHPC), UK & Ireland); Antoni Noguera (CoRISPE-CAT, Catalonia cohort, Spain) Maria Luisa Navarro, Jose Tomas Ramos (CoRISPE-S, rest of Spain cohort, Spain); Luisa Galli (Italian Register for HIV infection in children, Italy); Carlo Giaquinto, Claire Thorne (Paediatric European Network for the Treatment of AIDS (PENTA), Italy); Santa Ansone (Latvian cohort, Latvia); Magdalena Marczyńska (Polish paediatric cohort, Poland); Liubov Okhonskaia (Republican Hospital of Infectious Diseases, St Petersburg, Russia); Begoña Martínez de Tejada (Swiss Mother and Child HIV Cohort Study, Switzerland); Gonzague Jourdain, Luc Decker (Thailand Program for HIV Prevention and Treatment (PHPT) Study Group, Thailand); Luminita Ene (“Victor Babes” Hospital Cohort, Romania).

APPENDIX: COLLABORATING COHORTS

Belgium: Hospital St Pierre Cohort, Brussels: Tessa Goetghebuer, MD, PhD; Marc Hainaut, MD PhD; Evelyne Van der Kelen, Research nurse; Marc Delforge, data manager.

Greece: Greek cohort: Vana Spoulou.

Italy: Italian Register for HIV infection in Children. Coordinators: Maurizio de Martino (Florence), Pier Angelo Tovo (Turin). Participants: Osimani Patrizia (Ancona), Domenico Larovere (Bari), Maurizio Ruggeri (Bergamo), Giacomo Faldella, Francesco Baldi (Bologna) Raffaele Badolato (Brescia), Carlotta Montagnani, Elisabetta Venturini, Catuscia Lisi (Florence), Antonio Di Biagio, Lucia Taramasso (Genoa), Vania Giacomet, Paola Erba, Susanna Esposito, Rita Lipreri, Filippo Salvini, Claudia Tagliabue (Milan), Monica Cellini (Modena), Eugenia Bruzzese, Andrea Lo Vecchio (Naples), Osvalda Rampon, Daniele Donà (Padua), Amelia Romano (Palermo), Icilio Dodi (Parma), Anna Maccabruni (Pavia), Rita Consolini (Pisa), Stefania Bernardi, Hyppolite Tchidjou Kuekou, Orazio Genovese (Rome), Paolina Olmeo (Sassari), Letizia Cristiano (Taranto), Antonio Mazza (Trento), Clara Gabiano, Silvia Garazzino (Turin), Antonio Pellegatta (Varese)

Latvia: Latvian Cohort (Santa Ansone)

Netherlands: The ATHENA database is maintained by Stichting HIV Monitoring and supported by a grant from the Dutch Ministry of Health, Welfare and Sport through the Centre for Infectious Disease Control of the National Institute for Public Health and the Environment.

CLINICAL CENTRES (PAEDIATRIC CARE)

Emma Kinderziekenhuis, Amsterdam, University Medical Centers: *HIV treating physicians:* D. Pajkrt, H.J. Scherpbier. *HIV nurse consultants:* A.M. Weijnsfeld, C.G de Boer. *HIV clinical virologists/chemists:* S. Jurriaans, N.K.T. Back, H.L. Zaaijer, B. Berkhout, M.T.E. Cornelissen, C.J. Schinkel, K.C.wolthers. **Erasmus MC–Sophia, Rotterdam:** *HIV*

treating physicians: P.L.A. Fraaij, A.M.C. van Rossum, C.L. Vermont. *HIV nurse consultants:* L.C. van der Knaap, E.G. Visser. *HIV clinical virologists/chemists:* C.A.B. Boucher, M.P.G Koopmans, J.J.A van Kampen, S.D. Pas. **Radboudumc, Nijmegen:** *HIV treating physicians:* S.S.V. Henriët, M. van de Flier, K. van Aerde. *HIV nurse consultants:* R. Strik-Albers. *HIV clinical virologists/chemists:* J. Rahamat-Langendoen, F.F. Stelma. **Universitair Medisch Centrum Groningen, Groningen:** *HIV treating physicians:* E.H. Schölvink. *HIV nurse consultants:* H. de Groot-de Jonge. *HIV clinical virologists/chemists:* H.G.M. Niesters, C.C. van Leer-Buter, M. Knoester. **Wilhelmina Kinderziekenhuis, UMC Utrecht, Utrecht:** *HIV treating physicians:* L.J. Bont, S.P.M. Geelen, T.F.W. Wolfs. *HIV nurse consultants:* N. Nauta. *HIV clinical virologists/chemists:* R. Schuurman, F. Verduyn-Lunel, A.M.J. Wensing.

COORDINATING CENTRE

Director: P. Reiss. *Deputy director:* S. Zaheri. *Data analysis:* D.O. Bezemer, A.I. van Sighem, C. Smit, F.W.M.N. Wit. *Data management and quality control:* M. Hillebregt, A. de Jong, T. Woudstra. *Data monitoring:* D. Bergsma, S. Grivell, R. Meijering, M. Raethke, T. Rutkens. *Data collection:* L. de Groot, M. van den Akker, Y. Bakker, M. Bezemer, A. El Berkaoui, J. Geerlinks, J. Koops, E. Kruijne, C. Lodewijk, E. Lucas, R. van der Meer, L. Munjishvili, F. Paling, B. Peeck, C. Ree, R. Regtop, Y. Ruijs, L. van de Sande, M. Schoorl, P. Schnörr, E. Tuijn, L. Veenenberg, S. van der Vliet, A. Wisse, E.C. Witte. *Patient registration:* B. Tuk.

Poland: Polish paediatric cohort: Head of the team: Prof Magdalena Marczyńska, MD, PhD. Members of the team: Jolanta Popielska, MD, PhD; Maria Pokorska-Śpiewak, MD, PhD; Agnieszka Ołdakowska, MD, PhD; Konrad Zawadka, MD, PhD; Urszula Coupland, MD, PhD. Administration assistant: Małgorzata Doroba. Affiliation: Medical University of

Warsaw, Poland, Department of Children's Infectious Diseases; Hospital of Infectious Diseases in Warsaw, Poland.

Romania: "Victor Babes" Hospital Cohort, Bucharest: Dr Luminita Ene.

Russia: Federal State-owned Institution "Republican Clinical Infectious Diseases Hospital" of the Ministry of Health of the Russian Federation, St Petersburg: Liubov Okhonskaia, Evgeny Voronin, Milana Miloenko, Svetlana Labutina

Spain: CoRISPE-cat, Catalonia: financial support for CoRISPE-cat was provided by the Instituto de Salud Carlos III through the Red Temática de Investigación Cooperativa en Sida. Members: Hospital Universitari Vall d'Hebron, Barcelona (Pere Soler-Palacín, Maria Antoinette Frick and Santiago Pérez-Hoyos (statistician)), Hospital Universitari del Mar, Barcelona (Antonio Mur, Núria López), Hospital Universitari Germans Trias i Pujol, Badalona (María Méndez), Hospital Universitari Josep Trueta, Girona (Lluís Mayol), Hospital Universitari Arnau de Vilanova, Lleida (Teresa Vallmanya), Hospital Universitari Joan XXIII, Tarragona (Olga Calavia), Consorci Sanitari del Maresme, Mataró (Lourdes García), Hospital General de Granollers (Maite Coll), Corporació Sanitària Parc Taulí, Sabadell (Valentí Pineda), Hospital Universitari Sant Joan, Reus (Neus Rius), Fundació Althaia, Manresa (Núria Rovira), Hospital Son Espases, Mallorca (Joaquín Dueñas) and Hospital Sant Joan de Déu, Esplugues (Clàudia Fortuny, Antoni Noguera-Julian).

Spain: CoRISPE-S and Madrid cohort: María José Mellado, Luis Escosa, Milagros García Hortelano, Talía Sainz (Hospital La Paz); María Isabel González- Tomé, Pablo Rojo, Daniel Blázquez (Hospital Doce de Octubre, Madrid); José Tomás Ramos (Hospital Clínico San Carlos, Madrid); Luis Prieto, Sara Guillén (Hospital de Getafe); María Luisa Navarro, Jesús Saavedra, Mar Santos, M^a Angeles Muñoz, Beatriz Ruiz, Carolina Fernandez Mc Phee, Santiago Jimenez de Ory, Susana Alvarez (Hospital Gregorio Marañón); Miguel Ángel Roa

(Hospital de Móstoles); José Beceiro (Hospital Príncipe de Asturias, Alcalá de Henares); Jorge Martínez (Hospital Niño Jesús, Madrid); Katie Badillo (Hospital de Torrejón); Miren Apilanez (Hospital de Donostia, San Sebastián); Itziar Pocheville (Hospital de Cruces, Bilbao); Elisa Garrote (Hospital de Basurto, Bilbao); Elena Colino (Hospital Insular Materno Infantil, Las Palmas de Gran Canaria); Jorge Gómez Sirvent (Hospital Virgen de la Candelaria, Santa Cruz de Tenerife); Mónica Garzón, Vicente Román (Hospital de Lanzarote); Abián Montesdeoca, Mercedes Mateo (Complejo Universitario de Canarias, La Laguna-Tenerife), María José Muñoz, Raquel Angulo (Hospital de Poniente, El Ejido); Olaf Neth, Lola Falcón (Hospital Virgen del Rocío, Sevilla); Pedro Terol (Hospital Virgen de la Macarena, Sevilla); Juan Luis Santos (Hospital Virgen de las Nieves, Granada); David Moreno (Hospital Carlos Haya, Málaga); Francisco Lendínez (Hospital de Torrecárdenas, Almería); Ana Grande (Complejo Hospitalario Universitario Infanta Cristina, Badajoz); Francisco José Romero (Complejo Hospitalario de Cáceres); Carlos Pérez (Hospital de Cabueñes, Gijón); Miguel Lillo (Hospital de Albacete); Begoña Losada (Hospital Virgen de la Salud, Toledo); Mercedes Herranz (Hospital Virgen del Camino, Pamplona); Matilde Bustillo, Carmelo Guerrero (Hospital Miguel Servet, Zaragoza); Pilar Collado (Hospital Clínico Lozano Blesa, Zaragoza); José Antonio Couceiro (Complejo Hospitalario de Pontevedra); Amparo Pérez, Ana Isabel Piqueras, Rafael Bretón, Inmaculada Segarra (Hospital La Fe, Valencia); César Gavilán (Hospital San Juan de Alicante); Enrique Jareño (Hospital Clínico de Valencia); Elena Montesinos (Hospital General de Valencia); Marta Dapena (Hospital de Castellón); Cristina Álvarez (Hospital Marqués de Valdecilla, Santander); Ana Gloria Andrés (Hospital de León); Víctor Marugán, Carlos Ochoa (Hospital de Zamora); Santiago Alfayate, Ana Isabel Menasalvas (Hospital Virgen de la Arrixaca, Murcia); Elisa de Miguel (Complejo Hospitalario San Millán-San Pedro, Logroño) and

Paediatric HIV-BioBank integrated in the Spanish AIDS Research Network and collaborating Centers.

Funding: This work has been partially funded by the Fundación para la Investigación y Prevención de SIDA en España (FIPSE) (FIPSE 3608229/09 , FIPSE 240800/09, FIPSE 361910/10), Red Temática de Investigación en SIDA (RED RIS) supported by Instituto de Salud Carlos III (ISCIII) (RD12/0017/0035 and RD12/0017/0037), project as part of the Plan R+D+I and cofinanced by ISCIII- Subdirección General de Evaluación and Fondo Europeo de Desarrollo Regional (FEDER), Mutua Madrileña 2012/0077, Gilead Fellowship 2013/0071, FIS PI15/00694 , CoRISpe (RED RIS RD06/0006/0035 y RD06/0006/0021).

Sweden: Karolinska University Hospital, Stockholm (Lars Naver, Sandra Soeria-Atmadja, Vendela Hagås).

Switzerland: *Members of the Swiss HIV Cohort Study (SHCS) and the Swiss Mother and Child HIV Cohort Study:* Aebi-Popp K, Anagnostopoulos A, Asner S, Battegay M, Baumann M, Bernasconi E, Böni J, Braun DL, Bucher HC, Calmy A, Cavassini M, Ciuffi A, Duppenhaler A, Dollenmaier G, Egger M, Elzi L, Fehr J, Fellay J, Francini K, Furrer H, Fux CA, Grawe C, Günthard HF (President of the SHCS), Haerry D (deputy of "Positive Council"), Hasse B, Hirsch HH, Hoffmann M, Hösli I, Huber M, Kahlert CR (Chairman of the Mother & Child Substudy), Kaiser L, Keiser O, Klimkait T, Kottanattu L, Kouyos RD, Kovari H, Ledergerber B, Martinetti G, Martinez de Tejada B, Marzolini C, Metzner KJ, Müller N, Nicca D, Paioni P, Pantaleo G, Perreau M, Polli Ch, Rauch A (Chairman of the Scientific Board), Rudin C, Scherrer AU (Head of Data Centre), Schmid P, Speck R, Stöckle M (Chairman of the Clinical and Laboratory Committee), Tarr P, Thanh Lecompte M, Trkola A, Vernazza P, Wagner N, Wandeler G, Weber R, Wyler CA, Yerly S. *Funding:* This study

has been financed within the framework of the Swiss HIV Cohort Study, supported by the Swiss National Science Foundation (grant #177499).

Thailand: Program for HIV Prevention & Treatment (PHPT). Participating hospitals: Lamphun: Pornpun Wannarit; Phayao Provincial Hospital: Pornchai Techakunakorn; Chiangrai Prachanukroh: Rawiwan Hansudewechakul; Chiang Kham: Vanichaya Wanchaitanawong; Phan: Sookchai Theansavetrakul; Mae Sai: Sirisak Nanta; Prapokklao: Chaiwat Ngampiyaskul; Banglamung: Siriluk Phanomcheong; Chonburi: Suchat Hongsiriwon; Rayong: Warit Karnchanamayul; Bhuddasothorn Chacheongsao: Ratchanee Kwanchaipanich; Nakornping: Suparat Kanjanavanit; Somdej Prapinklao: Nareerat Kamonpakorn, Maneeratn Nantarukchaikul; Bhumibol Adulyadej: Prapaisri Layangool, Jutarat Mekmullica; Pranangkla: Paiboon Luksanapisitkul, Sudarat Watanayothin; Buddhachinaraj: Narong Lertpienthum; Hat Yai: Boonyarat Warachit; Regional Health Promotion Center 6, Khon Kaen: Sansanee Hanpinitak; Nong Khai: Sathit Potchalongsin; Samutsakhon: Pimpraphai Thanasiri, Sawitree Krikajornkitti; Phaholpolphayuhasena: Pornsawan Attavinijtrakarn; Kalasin: Sakulrat Srirojana; Nakhonpathom: Suthunya Bunjongpak; Samutprakarn: Achara Puangsombat; Mahasarakam: Sathaporn Na-Rajsima; Roi-et: Pornchai Ananpatharachai; Sanpatong: Noppadon Akarathum; Vachira Phuket: Weerasak Lawtongkum; Chiangdao: Prapawan Kheunjan, Thitiporn Suriyaboon, Airada Saipanya.

Data management team: Kanchana Than-in-at, Nirattiya Jaisieng, Rapeepan Suaysod, Sanuphong Chailoet, Naritsara Naratee, and Suttipong Kawilapat.

UK & Ireland: Collaborative HIV Paediatric Study (CHIPS): CHIPS is funded by the NHS (London Specialised Commissioning Group) and has received additional support from Bristol-Myers Squibb, Boehringer Ingelheim, GlaxoSmithKline, Roche, Abbott, and Gilead

Sciences. The MRC Clinical Trials Unit at UCL is supported by the Medical Research Council (<https://www.mrc.ac.uk>) programme number MC_UU_12023/26.

CHIPS Steering Committee: Hermione Lyall (chair), Alasdair Bamford, Karina Butler, Katja Doerholt, Conor Doherty, Caroline Foster, Kate Francis, Ian Harrison, Julia Kenny, Nigel Klein, Gillian Letting, Paddy McMaster, Fungai Murau, Edith Nsangi, Helen Peters, Katia Prime, Andrew Riordan, Fiona Shackley, Delane Shingadia, Sharon Storey, Claire Thorne, Gareth Tudor-Williams, Anna Turkova, Steve Welch. **MRC Clinical Trials Unit:** Intira Jeannie Collins, Claire Cook, Siobhan Crichton, Donna Dobson, Keith Fairbrother, Diana M. Gibb, Lynda Harper, Ali Judd, Marthe Le Prevost, Nadine Van Looy. National Study of HIV in Pregnancy and Childhood, UCL: Helen Peters, Claire Thorne.

Participating hospitals: Republic of Ireland: Our Lady's Children's Hospital Crumlin, Dublin: K Butler, A Walsh. UK: University Hospitals Birmingham NHS Foundation Trust, Birmingham: L Thrasyvoulou, S Welch; University Hospitals Bristol NHS Foundation Trust, Bristol: J Bernatoniene, F Manyika; Calderdale and Huddersfield NHS Foundation Trust, Halifax: G Sharpe; Derby Teaching Hospitals NHS Foundation Trust: B Subramaniam; Middlesex: K Sloper; East Sussex Healthcare NHS Trust, Eastbourne: K Fidler, Glasgow Royal Hospital for Children, Glasgow: R Hague, V Price; Great Ormond Street Hospital for Children NHS Foundation Trust, London: M Clapson, J Flynn, A Cardoso M Abou – Rayyah, N Klein, D Shingadia; Homerton University Hospital NHS Foundation Trust, London: D Gurtin, Oxford University Hospitals NHS Foundation Trust, Oxford: S Yeadon, S Segal; King's College Hospital NHS Foundation Trust, London: C Ball, S Hawkins; Leeds Teaching Hospitals NHS Trust, Leeds: M Dowie; University Hospitals of Leicester NHS Trust, Leicester: S Bandi, E Percival; Luton and Dunstable Hospital NHS Foundation Trust, Luton: M Eisenhut; K Duncan, S Clough; Milton Keynes General University Hospital NHS Foundation Trust, Milton Keynes: Dr L Anguava, S Conway, Newcastle upon Tyne

Hospitals NHS Foundation Trust, Newcastle: T Flood, A Pickering; The Pennine Acute Hospitals NHS Trust, Manchester: P McMaster C Murphy; North Middlesex University Hospital NHS Trust, London: J Daniels, Y Lees; Northampton General Hospital NHS Trust, Northampton: F Thompson; London North West Healthcare NHS Trust, Middlesex; B Williams, S Pope; Nottingham University Hospitals NHS Trust, Nottingham: L Cliffe, A Smyth, S Southall; Portsmouth Hospitals NHS Trust, Portsmouth: A Freeman; Raigmore Hospital, Inverness: H Freeman; Royal Belfast Hospital for Sick Children, Belfast: S Christie; Royal Berkshire NHS Foundation Trust, Reading: A Gordon; Royal Children's Hospital, Aberdeen: D Rogahn L Clarke; Royal Edinburgh Hospital for Sick Children, Edinburgh: L Jones, B Offerman; Royal Free NHS Foundation Trust, London: M Greenberg ; Alder Hey Children's NHS Foundation Trust , Liverpool: C Benson, A Riordan; Sheffield Children's NHS Foundation Trust, Sheffield: L Ibberson, F Shackley; University Hospital Southampton NHS Foundation Trust, Southampton: SN Faust, J Hancock; St George's University Hospitals NHS Foundation Trust, London: K Doerholt, , K Prime, M Sharland, S Storey; Imperial College Healthcare NHS Trust, London: H Lyall, C Monrose, P Seery, G Tudor-Williams; Guy's and St Thomas' NHS Foundation Trust, London: , E Menson, A Callaghan; University Hospitals of North Midlands NHS Trust, Stoke On Trent: A Bridgwood, P McMaster; University Hospital of Wales, Cardiff: J Evans, E Blake; NHS Frimley Health Foundation Trust, Slough: A Yannoulis.

References

1. Williams PL, Jesson J. **Growth and pubertal development in HIV-infected adolescents.** *Current Opinion in HIV and AIDS* 2018; 13(3):179-186.
2. McGrath CJ, Diener L, Richardson BA, Peacock-Chambers E, John-Stewart GC. **Growth Reconstitution following Antiretroviral Therapy and Nutrition Supplementation: Systematic Review and Meta-Analysis.** *AIDS* 2015; 29(15):2009.
3. Szubert AJ, Musiime V, Bwakura-Dangarembizi M, Nahiry-Ntege P, Kekitiinwa A, Gibb DM, et al. **Pubertal development in HIV-infected African children on first-line antiretroviral therapy.** *AIDS* 2015; 29(5):609.
4. McHugh G, Rylance J, Mujuru H, Nathoo K, Chonzi P, Dauya E, et al. **Chronic morbidity among older children and adolescents at diagnosis of HIV infection.** *Journal of Acquired Immune Deficiency Syndromes* 2016; 73(3):275.
5. Williams PL, Abzug MJ, Jacobson DL, Wang J, Van Dyke RB, Hazra R, et al. **Pubertal onset in HIV-infected children in the era of combination Antiretroviral Treatment.** *AIDS* 2013; 27(12):1959.
6. Bellavia A, Williams PL, DiMeglio LA, Hazra R, Abzug MJ, Patel K, et al. **Delay in sexual maturation in perinatally HIV-infected youths is mediated by poor growth.** *AIDS* 2017; 31(9):1333-1341.
7. de Martino M, Tovo P-A, Galli L, Gabiano C, Chiarelli F, Zappa M, et al. **Puberty in perinatal HIV-1 infection: a multicentre longitudinal study of 212 children.** *AIDS* 2001; 15(12):1527-1534.

8. Willemsen RH, Dunger DB. **Normal variation in pubertal timing: genetic determinants in relation to growth and adiposity.** In: *Puberty from Bench to Clinic*: Karger Publishers; 2016. pp. 17-35.
9. Arpadi SM, Cuff PA, Kotler DP, Wang J, Bamji M, Lange M, et al. **Growth velocity, fat-free mass and energy intake are inversely related to viral load in HIV-infected children.** *The Journal of Nutrition* 2000; 130(10):2498-2502.
10. Benjamin Jr DK, Miller WC, Benjamin DK, Ryder RW, Weber DJ, Walter E, et al. **A comparison of height and weight velocity as a part of the composite endpoint in pediatric HIV.** *AIDS* 2003; 17(16):2331-2336.
11. Chantry CJ, Byrd RS, Englund JA, Baker CJ, MCKINNEY JR RE. **Growth, survival and viral load in symptomatic childhood human immunodeficiency virus infection.** *The Pediatric Infectious Disease Journal* 2003; 22(12):1033-1038.
12. Gilsanz V, Chalfant J, Kalkwarf H, Zemel B, Lappe J, Oberfield S, et al. **Age at onset of puberty predicts bone mass in young adulthood.** *The Journal of Pediatrics* 2011; 158(1):100-105. e102.
13. Eckard AR, Mora S. **Bone health in HIV-infected children and adolescents.** *Curr Opin HIV AIDS* 2016; 11(3):294-300.
14. Hodinott J, Behrman JR, Maluccio JA, Melgar P, Quisumbing AR, Ramirez-Zea M, et al. **Adult consequences of growth failure in early childhood--.** *The American Journal of Clinical Nutrition* 2013; 98(5):1170-1178.

15. WHO. **WHO child growth standards: length/height for age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age, methods and development.** World Health Organization; 2006.
16. WHO. **Growth reference data for 5-19 years. WHO Reference 2007.** World Health Organization; 2007.
17. WHO. **WHO case definitions of HIV for surveillance and revised clinical staging and immunological classification of HIV-related disease in adults and children.** World Health Organization; 2007.
18. Cole TJ, Donaldson MD, Ben-Shlomo Y. **SITAR—a useful instrument for growth curve analysis.** *International Journal of Epidemiology* 2010; 39(6):1558-1566.
19. Granados A, Gebremariam A, Lee JM. **Relationship between timing of peak height velocity and pubertal staging in boys and girls.** *Journal of Clinical Research in Pediatric Endocrinology* 2015; 7(3):235.
20. Tanner JM. **Growth of the human at the time of adolescence.** *Lectures on the scientific basis of medicine* 1953; 1:308.
21. Working Group on Using Weight and Height References in Evaluating Growth Status of Thai Children. **Manual on using weight and height references in evaluating the growth status of Thai children.** Bangkok: Department of Health, Ministry of Public Health; 2000.
22. Cole T, Cole MT. **sitar: Super Imposition by Translation and Rotation Growth Curve Analysis. R package version 1.0.10.** 2017.

23. Portrait FRM, van Winderden TF, Deeg DJH. **Early life undernutrition and adult height: The Dutch famine of 1944-45.** *Economics and Human Biology* 2017 Nov;27(Pt B):339-348.

24. Wagner IV, Sabin MA, Pfäffle RW, Hiemisch A, Sergeyev E, Körner A, et al. **Effects of obesity on human sexual development.** *Nature Reviews Endocrinology* 2012; 8(4):246.

25. Mo-Suwan L, Choprapawon C. **Comparison of Prevalence of Nutritional Status of Thai Children in the First 2 Years of Life Using National and International Growth Charts.** *J Med Assoc Thai* 2016; 99(1):58-64.

ACCEPTED

Tables and Figures

Figure 1: Flow chart of participants included in the study




Figure 2A: Mean height and growth velocity of females stratified by HAZ at ART initiation and 2B: mean height and growth velocity of females stratified by age at ART initiation





Figure 3A: Mean height and growth velocity of males stratified by HAZ at ART initiation and 3B: mean height and growth velocity of males stratified by age at ART initiation





Table 1: Characteristics of 1094 young people living with HIV at ART initiation stratified by height-for age z-scores

N(%) or median[IQR]	All	Height-for-age z-score at ART initiation				p-value
		<-3 SD (Severely stunted)	-3 to <-2 SD (Stunted)	-2 to <-1 SD	>= -1 SD	
All	1094	157(14)	187(17)	272(25)	478(44)	
Male	526(48)	83(53)	90(48)	117(43)	236(49)	0.207
Country						
UK & Ireland	517(47)	17(11)	56(30)	137(50)	307(64)	<0.001
Thailand	352(32)	121(77)	103(55)	84(31)	44(9)	
Other*	225(21)	19(12)	28(15)	51(19)	127(27)	
Ethnicity						
White	99(9)	10(6)	12(6)	26(10)	51(11)	<0.001
Black	484(44)	16(10)	49(26)	125(46)	294(62)	
Asian	365(33)	121(77)	106(57)	89(33)	49(10)	
Other	63(6)	6(4)	5(3)	19(7)	33(7)	
Unknown/Prohibited	83(8)	4(3)	15(8)	13(5)	51(11)	
Born abroad	370(35)	21(14)	51(28)	100(37)	198(43)	<0.001
Age						
Median[IQR] years	6.4[2.8,9.0]	6.4[2.7,9.0]	7.1[3.8,9.5]	6.4[3.2,8.7]	6.1[2.5,8.8]	0.236
1 to <3 years	280(26)	45(29)	35(19)	65(24)	135(28)	
3 to <6 years	237(22)	31(20)	43(23)	64(24)	99(21)	
6 to <11 years	577(53)	81(52)	109(58)	143(53)	244(51)	
Year started ART						
Median[IQR]	2004[2003,2007]	2003[2003,2008]	2004[200,2006]	2004[2003,2006]	2005[2003,2008]	<0.001
<2004	433(40)	79(50)	78(42)	116(43)	160(33)	
2004-2008	445(41)	69(44)	93(50)	106(39)	177(37)	
≥2008	216(20)	9(6)	16(9)	50(18)	141(30)	
NNRTI based regimen	880(80)	141(90)	167(89)	222(82)	350(73)	<0.001
Viral load						
Value present	980(90)	133(85)	167(89)	241(89)	439(92)	
Median log VL	5.0[4.5,5.5]	5.3[4.9,5.7]	5.0[4.7,5.5]	5.0[4.5,5.4]	5.0[4.2,5.5]	<0.001
≤400	36(4)	4(3)	4(2)	6(2)	22(5)	
>400 to 1,000	13(1)	1(1)	1(1)	1(0)	10(2)	
>1,000 to 10,000	91(9)	7(5)	11(7)	21(9)	52(12)	
>10,000 to 100,000	435(35)	39(29)	60(36)	89(27)	157(36)	
>100,000	495(51)	82(62)	91(54)	124(51)	198(45)	
WHO immunological classification						
Value present	1006(92)	145(92)	177(95)	258(95)	426(89)	
None or not significant	164(16)	7(5)	16(9)	40(16)	101(24)	<0.001
Mild	110(11)	5(3)	12(7)	27(10)	66(15)	

Advanced	129(13)	6(4)	12(7)	35(14)	76(18)	
Severe	603(60)	127(88)	137(77)	156(60)	183(43)	
zBMI						
Value present	1089(100)	157(100)	186(100)	271(100)	475(99)	
Median zBMI	-0.1[-1.1,0.8]	-1.1[-2.3,0.2]	-0.7[-1.6,0.3]	-0.1[-0.9,0.7]	0.3[-0.5,1.1]	<0.001
<-2 SD (Thinness)	133(12)	49(31)	34(18)	22(8)	28(6)	
-2 to <-1 SD	164(15)	32(20)	42(23)	42(16)	48(10)	
-1 to <1 SD	576(53)	59(38)	96(62)	158(58)	263(55)	
1 to <2 SD (Overweight)	160(15)	13(8)	12(6)	37(14)	98(21)	
>=2 SD (Obese)	56(5)	4(3)	2(1)	12(4)	38(8)	

*Other includes Belgium, Greece, Latvia, Netherlands, Poland, Romania, Spain, Sweden and Switzerland

ACCEPTED

Table 2: Association between characteristics at ART initiation and average height, timing and shape of growth spurt after adjustment for baseline age and HAZ in 918 young people living with HIV

	Average height			Timing of growth spurt			Shape of growth spurt		
	coef	95% CI	P-value	coef	95% CI	P-value	coef	95% CI	P-value
Female	0.58	-0.18, 1.34	0.132	0.04	-0.08, 0.16	0.485	0.007	-0.015, 0.029	0.555
Country (ref: Other Europe)									
Thailand	-3.06	-4.34, -1.78	<0.001	0.04	-0.16, 0.24	0.701	-0.030	-0.067, 0.007	0.116
UK & Ireland	-0.55	-1.65, 0.55	0.325	0.10	-0.07, 0.29	0.241	-0.033	-0.065, -0.001	0.044
Year (ref: <2004)									
2004 to <2008	-0.31	-1.17, 0.55	0.476	0.00	-0.14, 0.14	0.991	0.010	-0.015, 0.035	0.426
>=2008	-0.22	-1.33, 0.90	0.704	-0.08	-0.26, 0.10	0.369	0.011	-0.022, 0.043	0.512
NNRTI based regimen (ref: PI based regimen)	0.27	-0.78, 1.31	0.616	-0.04	-0.20, 0.13	0.673	-0.018	-0.048, 0.012	0.240
Viral load (per log increase)	-0.10	-0.51, 0.30	0.615	0.03	-0.03, 0.10	0.292	-0.002	-0.013, 0.010	0.787
WHO immunological classification (ref: None)									
Mild	-0.10	-1.56, 1.35	0.891	0.06	-0.17, 0.29	0.608	-0.003	-0.045, 0.039	0.900
Advanced	-0.56	-1.96, 0.84	0.433	-0.06	-0.29, 0.16	0.581	-0.016	-0.057, 0.024	0.426
Severe	0.28	-0.84, 1.39	0.628	-0.01	-0.18, 0.17	0.948	0.002	-0.034, 0.031	0.927
zBMI (per 1SD increase)	-0.32	-0.60, -0.05	0.021	-0.07	-0.11, -0.02	0.003	-0.007	-0.015, 0.001	0.083

Note: Individual size, tempo and velocity parameters were estimated using the SITAR model described in the results and table S1 and represent the differences in size, tempo and velocity unexplained by age and HAZ at ART initiation. Model included data from 918 of the 1094 children included in the SITAR model for which data on the explanatory variables were complete.