


Accepted Manuscript

Global glycosphingolipid analysis in urine and plasma of female Fabry disease patients

Wendy Heywood, Ivan Doykov, Justyna Spiewak, Jenny Hallqvist, Kevin Mills, Albina Nowak



PII: S0925-4439(19)30228-5

DOI: <https://doi.org/10.1016/j.bbadis.2019.07.005>

Reference: BBADIS 65511

To appear in: *BBA - Molecular Basis of Disease*

Received date: 19 March 2019

Revised date: 22 June 2019

Accepted date: 11 July 2019

Please cite this article as: W. Heywood, I. Doykov, J. Spiewak, et al., Global glycosphingolipid analysis in urine and plasma of female Fabry disease patients, *BBA - Molecular Basis of Disease*, <https://doi.org/10.1016/j.bbadis.2019.07.005>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Global Glycosphingolipid Analysis in Urine and Plasma of Female Fabry Disease Patients

Wendy Heywood<sup>1,2\*</sup>, Ivan Doykov<sup>1\*</sup> Justyna Spiewak<sup>1</sup>, Jenny Hallqvist<sup>1</sup>, Kevin Mills<sup>1,2§</sup>, Albina Nowak<sup>3,4§</sup>

\* Joint first authors, § joint last authors

<sup>1</sup> Inborn Errors of Metabolism Section, Genetics & Genomic Medicine Unit, Great Ormond Street Institute of Child Health, University College London, 30 Guilford Street, London, WC1N 1EH UK

<sup>2</sup> NIHR Great Ormond Street Biomedical Research Centre, Great Ormond Street Hospital and UCL Great Ormond Street Institute of Child Health

<sup>3</sup> Department of Endocrinology and Clinical Nutrition, University Hospital Zurich and University of Zurich, Zurich, Switzerland

<sup>4</sup> Department of Internal Medicine, Psychiatry University Clinic Zurich, Zurich, Switzerland

Keywords: Fabry disease; glycosphingolipid; biomarker, galabiosylceramide, lyso-Gb<sub>3</sub>; ceramide dihexoside

## Abstract

Fabry disease (FD) is an X-linked lysosomal storage disorder caused by deficiency of  $\alpha$ -galactosidase-A, which results in accumulation of the glycosphingolipid (GSL) globotriaosylceramide (Gb<sub>3</sub>). Gb<sub>3</sub> and globotriaosylsphingosine (lyso-Gb<sub>3</sub>) levels in plasma and urine are used routinely for diagnosis and treatment monitoring. FD female patients are problematic to diagnose and to predict when to begin treatment. Further biomarkers are needed to detect pre-symptomatic females that will develop the chronic symptoms associated with FD. A LC-MS/MS glycosphingolipidomic assay was developed to measure lyso-Gb<sub>3</sub> and GSLs from the lysosomal GSL degradation pathway, including globoside (Gb<sub>4</sub>), Gb<sub>3</sub>, ceramide dihexosides (CDH) and ceramide monohexosides (CMH). We analysed plasma and urine from a cohort of Fabry patients, grouped according to clinical symptoms and independent of treatment status (asymptomatic females n=18, symptomatic females n=18, males n= 27 and control urines n=16 and control plasmas n=58).

Multivariate and subsequent univariate analysis showed urine GSLs which had highest significance in identifying asymptomatic females were total levels of CDH, in particular the long chain isoforms C22:1,C22:0,C22:1-OH,C22:0-OH,C24:2,C24:0,C24:2-OH,C24:1-OH,C24:0-OH,C26:0 which likely represent Galabiosylceramide (Ga<sub>2</sub>) and not lactosylceramide. These long chain Ga<sub>2</sub> isoforms were found to be 5-fold elevated and more statistically significant ( $p<0.0001$ ) than plasma lyso-Gb<sub>3</sub> ( $p< 0.01$ ) in identifying asymptomatic Fabry female patients. Receiver operating characteristic curve analysis gave an area under the curve of 0.82 ( $p=0.001$ ) for lyso-Gb<sub>3</sub> and 0.88 ( $p=0.0006$ ) for long-chain CDH isoforms indicating the long chain CDH isoforms were as, if not more, a better biomarker for the identification of female FD patients.

## Introduction

Fabry disease (FD) is an X-linked, lysosomal storage disorder (LSD) that is caused by mutations of the *GLA*-gene that lead to markedly reduced or absent activity of the lysosomal enzyme alpha-galactosidase (GLA). This results in a progressive accumulation of the glycosphingolipid (GSL) globotriaosylsphingosine (Gb<sub>3</sub>) [1]. Traditionally, Gb<sub>3</sub> has been an accepted biomarker for FD but in recent years the deacylated version of Gb<sub>3</sub>, lyso-Gb<sub>3</sub>, has been observed to be more indicative of disease burden and monitoring the response to treatment [2]. The presence of lyso-Gb<sub>3</sub> in the circulation is not completely understood but is thought to be an inadvertent product of promiscuous acid ceramidase activity on Gb<sub>3</sub> [3]. Female FD patients who are heterozygous for the disease are typically harder to detect and monitor biochemically using Gb<sub>3</sub> [4]. Lyso-Gb<sub>3</sub> so far is proving to be more useful for identifying FD females [4-6]. However, not all FD females can be detected using plasma lyso-Gb<sub>3</sub> levels and usually require a mixture of glycosphingolipid, enzymology and genetics to diagnose. Therefore there is a need for better and additional biomarkers [5].

The GSL degradation pathway involves multiple steps, by highly specific enzymes, that result in the breakdown of the ganglioside GM<sub>1</sub> and globoside (Gb<sub>4</sub>), finally resulting in the production of sphingosine which is then recycled (figure 1). This results in various intermediate compounds including Gb<sub>3</sub>, lactosylceramide (LacCer), glucosylceramide (GlcCer) and ceramide. Galabiosylceramide (Ga<sub>2</sub> or Gb<sub>2</sub>) is also a substrate of alpha-galactosidase and is known to accumulate in FD but not to the extent as Gb<sub>3</sub>. To add to the complexity of GSLs, they also consist of multiple isoforms resulting from the different fatty acid chain lengths, double bonds and fatty acid modifications such as hydroxylation [7]. Modifications also can occur on the sphingosine backbone [8] which is pertinent to the discovery of the lyso-Gb<sub>3</sub> analogues that are relevant in FD [9]. The role and the function of the different isoforms for many GSLs are largely unknown [10] and with the effect on the overall GSL degradation pathway in FD patients has not been fully investigated. Recent mouse studies have indicated other GSL species could be affected [11, 12] and therefore, in this study we have looked at the GSL degradation pathway in more detail to see if any changes are also observed in humans. We have developed a multiplex UPLC LC-MS/MS glycosphingolipid screen to quantitate the main isoforms of the four GSL species globoside (Gb<sub>4</sub>), Gb<sub>3</sub>, ceramide dihexoside (CDH) and ceramide monohexoside (CMH) and lyso-Gb<sub>3</sub> (including analogues) [13]. Our primary objective was to use this assay to study the effect on the metabolic flux of other glycosphingolipids upstream and downstream of the enzymatic defect found in FD. The secondary aim was to see if any of these other lipids were of use for aiding in the diagnosis of FD, in particular in the

detection of FD females. Finally, we have compared those GSL species to the gold standard plasma lyso-Gb<sub>3</sub>, the currently used biomarker, to see if they can aid in the diagnosis of FD.

## Methods

### Patient samples

FD plasma and urine samples were obtained from University Hospital Zurich, Zurich, Switzerland, all with a *GLA*-mutation confirmed diagnosis, who presented for routine annual examinations at the specialized FD centre. All patients participated in this study were treated in the same hospital. All patients had a comprehensive work-up, including medical history, cardiac, renal, and neurological evaluations. ERT was initiated according to the written local guidelines. Accordingly, ERT was indicated in all males with pathogenic *GLA* mutation, independent from age, phenotype, and symptoms. In females, ERT was indicated if they had proteinuria of more than 300 mg per day, FD-typical kidney biopsy findings, signs of FD cardiomyopathy, such as left ventricular hypertrophy or arrhythmia, if the FD patients had stroke or transient ischemic attack (TIA), persistent FD-related neuropathic pains despite conventional analgetic therapy, and or gastrointestinal symptoms. In all patients, ERT was prescribed at the licensed dose of either 0.2 mg/kg body weight of recombinant agalsidase-a (Replagal) or 1 mg/kg body weight agalsidase-b (Fabrazyme) and given intravenously every 14 days. Each patients ERT status is indicated in supplementary data file 1 which includes all known clinical data and corresponding glycosphingolipid values. This study was conducted in accordance with the principles of the Helsinki Declaration and approved by the Zurich Ethical Committee; the approval number is 2017-00386. All patients signed a written informed consent.

Samples for analysis consisted of 27 males (19-65 yrs median age 47) all on ERT and symptomatic. Thirty-seven female patients were included and divided into 2 experimental groups FD female asymptomatic (age range 21-65 yrs, median 36.5 yrs) where patients were mostly not on ERT and a FD female group who were symptomatic (age range 19-78 yrs, median 51 yrs) where the majority of patients were on ERT. Control samples used in this analysis (plasma n= 64 M: F 32:32, urine n=12 M: F 4:8) were from healthy volunteers (age range 22-61 yrs, median 29 yrs).

**Sample preparation for LC-MS/MS analysis**

**Plasma lyso-Gb<sub>3</sub> analysis:** Lyso Gb<sub>3</sub> was extracted from 100 µl of plasma using 1 ml of acetone: methanol (1:1 v/v) containing 2 ng/ml di-methyl psychosine internal standard (Avanti Polar Lipids inc). Samples were shaken for 20 min, sonicated for 15 min in a sonicator bath and shaken again for 20 min at room temperature (RT). After centrifugation at 16000 g, supernatants was transferred and evaporated on a rotational evaporator at RT. Samples were reconstituted in 100 µl of methanol prior to LC-MS/MS analysis.

**Urine lyso-Gb<sub>3</sub> analysis:** Lyso-Gb<sub>3</sub> was extracted from 500 µl of urine using 600 µl 0.2% formic acid (FA), containing 50 µl of Internal standard solution at 0.1 µg /ml in methanol, followed by 50 µl of methanol. Samples were shaken for 20 min at RT and then cleaned using C18 SPE. 1 ml C18 cartridges were primed with 1 ml of methanol:chloroform (2:1 v/v), then equilibrated 2x 1ml 0.1 % FA. Samples are added and cartridges washed with 0.1 % FA. Lyso-Gb<sub>3</sub> was eluted in 500 µl of methanol:chloroform (2:1 v/v). Samples were dried using a rotational evaporator at RT and reconstituted 100 µl of methanol prior to LC-MS/MS analysis.

**Plasma GSL analysis:** GSLs were extracted from 50 µl of plasma using a 1 ml solution containing 0.5 µg/ µl C17-CTH (Matreya LLC, USA) and 0.5 µg / µl [D<sub>3</sub>]C16:0-Lactosylceramide (Matreya LLC, USA) internal standard in chloroform:methanol (2:1 v/v). Samples were shaken for 20 min at RT, sonicated for 15 min, shaken again for 20 min. 200 µl of ice-cold PBS was added and samples were shaken again for 20 min. Samples were spun at 16000 g for 10min. Using a glass pipette the bottom layer is carefully extracted and placed in a 5 ml glass vial. Samples were dried under nitrogen and reconstituted in 1 ml of chloroform. Samples were cleaned using C18 SPE cartridges (Biotage, UK). Cartridges were primed with 1 ml acetone:methanol (9:1 v/v) then equilibrated twice with chloroform before the addition of sample. Cartridges were washed 2 x 1 ml chloroform and then eluted with 700 µl of acetone:methanol (9:1 v/v). Samples were dried under nitrogen and reconstituted in 200 µl of methanol prior to MS analysis.

**Urine GSL analysis:** GSLs were extracted from 50 µl of urine using 200 µl solution containing 0.5 µg/µl C17-CTH (Matreya LLC, USA) and 0.5 µg/µl [D<sub>3</sub>]C16:0-Lactosylceramide (Matreya LLC, USA) internal standard in methanol:chloroform (2:1 v/v). Samples were shaken for 1 hr at RT. 150 µl of isopropanol was added and samples centrifuged for 10 min at 16000 g. Supernatants were transferred to a new vial and evaporated using a rotational evaporator at RT. Samples were reconstituted in 100 µl of methanol prior to MS analysis.

**UPLC-MS/MS analysis**

An ACQUITY UPLC system from Waters Corporation (Milford, MA) was used for the separation of GSLs before quantitative tandem mass spectrometry analysis. The reverse-phase UPLC methods used for plasma and urine GSL and lyso-Gb<sub>3</sub> analysis are described in supplementary data table S-1. The UPLC system was coupled to a Xevo TQ-S (Waters Ltd, UK) mass spectrometer operated in the multiple reaction monitoring (MRM) mode. MRM transitions corresponding to isoforms of CMH, CDH, Gb<sub>3</sub>, Gb<sub>4</sub>, and lyso-Gb<sub>3</sub> are given in supplementary data Table S-5. Internal standard transitions were included in all MRM files. Transition dwell times were set automatically with minimum of 0.03 sec and minimum of 8 data points per peak. The divert valve of the mass spectrometer was programmed to discard the UPLC effluent before (0 to 1 min) and after (2.8-3.5 GSL 5 min methods) and 8-10 min (10 min methods) to reduce system contamination. Galabiosylceramide (Gb<sub>2</sub>) is a structural isomer of lactosylceramide (LacCer) a product of alpha galactosidase from ganglioside degradation (figure 1 A). Being structural isomers these two GSLs have the same molecular weight and cannot be separated by conventional reverse phase chromatography. Similarly, it is not possible to separate structural isomers GlcCer and galactosylceramide (GalCer). Therefore, they are referred to as total CDH and total CMH in this study. We demonstrate that there is no requirement to separate these molecules using normal phase chromatography methods such as a HILIC chromatography.

The concentrations of lyso-Gb<sub>3</sub> and its GSLs were evaluated using the TargetLynx 4.1 software (Waters). C17-CTH was used as the internal standard for the quantification of Gb<sub>3</sub> and Gb<sub>4</sub> and [D<sub>3</sub>]C16:0-Lactosylceramide for quantitation of CDH and CMH. Dimethylpsychosine was used as IS for lyso-Gb<sub>3</sub>. Calibration curves were linear the origin was excluded. Urinary lipids were ratioed to urinary creatinine levels as described previously [14]. Further methods details are available in the supplementary information.

**Data analysis:** Excel and GraphPad Prism v 6 were used for all composition, comparative and ROC analysis. Both parametric and non-parametric analysis was used for comparisons where appropriate. Multivariate analysis including principal components and OPLS-DA analysis was performed using SIMCA v 14 (Umetrics, Sweden). Where there is no data a 0 value for that molecule is given for analysis.

Spearman Correlation matrix analysis was performed using an in-house Python v 3.6 script (URL <https://www.python.org/>).




Figure 1. A. Glycosphingolipid degradation pathway (Polo et al 2017[15]) with highlighted GSL species analysed including Lyso-Gb<sub>3</sub>. LacCer and Ga2 consist of CDH and GlcCer and GalCer consist of CMH.



## Results

### Glycosphingolipid Profile and Composition in Normal and Fabry Disease Patients

Figure 3 shows the total averaged abundance levels of each GSL in (i) plasma and (ii) urine. In control subjects, CDH and Gb<sub>3</sub> are the most abundant GSLs present in both plasma and in urine. Additionally, CDH makes up 47% and Gb<sub>3</sub> 38% of total plasma GSL in healthy controls. For total GSLs in urine, CDH consists of approximately 72% in females and 59% in males, with Gb<sub>3</sub> making up 22% of total glycosphingolipids in females and 35% in males, respectively. Plasma Gb<sub>4</sub> and CMH levels are much lower (Gb<sub>4</sub> = ~4%, CMH = ~11%) and in urine (Gb<sub>4</sub> ~0.31%, CMH = ~6%) in both males and females. Surprisingly, control females have a much higher level of total GSLs in urine than control males (figure 3 (ii)) of which the majority is comprised of CDH and which confirms previous reports [16]. Furthermore, the proportion of CDH in female urine was also observed to be higher than in men (CDH ~72% in females vs ~59% in males,  $p < 0.03$ ). However, males were observed to have a greater proportion of Gb<sub>3</sub>, 22% in females vs 35% in males ( $p < 0.02$ ).

Analysis of patient samples demonstrated the composition of the CDH and Gb<sub>3</sub> in FD does not appear to change due to the enzymatic defect observed in FD (figure 2iii), with FD females having on average 74% CDH vs 55% in FD males ( $p < 0.001$ ), and 24% Gb<sub>3</sub> vs 41% in males ( $p < 0.001$ ). In FD, plasma GSL levels do not alter greatly with only moderate total increases in all FD groups (figure 3 (i)). However, in urine, the increase is much more profound in both males and females which confirms why urine Gb<sub>3</sub> is typically used for diagnosis before lyso-Gb<sub>3</sub> was discovered [14].





Figure 2. Overview of the approximate average levels and composition of each GSL species in (i) plasma and (ii) urine for Control Female (Ctl), Fabry Female Asymptomatic, Fabry Female Symptomatic, Control Male and Fabry Male who are all symptomatic. (iii) Shows percentage composition comparison of each measured GSL species. CMH and Gb<sub>4</sub> are prevalent in plasma compared to urine. Changes in total levels of CDH and Gb<sub>3</sub> are more evident in the urine in Fabry disease. Females have both higher urinary CDH and Gb<sub>3</sub> levels which both increase in Fabry disease which also increase in males but Gb<sub>3</sub> levels and composition are higher in males.

### Glycosphingolipidomic changes observed in Fabry Disease Patients

Figure 4 shows that when comparing the total levels of each GSL in the FD groups, it was observed that there is a small but significant increase of approximately 1.4 fold of CMH in both FD female groups compared to controls ( $p < 0.001$ ). This increase is also observed in the FD male group but to a lesser extent of 1.3 fold ( $p < 0.05$ ). The only other significantly altered GSL in plasma was a slight increase of Gb<sub>3</sub> in the FD male group of appx 1.2 fold ( $p < 0.01$ ).

However greater changes in other GSLs were observed in urine. The lesser abundant GSLs CMH and Gb<sub>4</sub> were observed to be significantly affected in FD patient urine. CMH is slightly elevated (1.6 fold) in the male FD group ( $p < 0.05$ ) but Gb<sub>4</sub> was observed to be 3-fold elevated in all FD groups. Urinary Gb<sub>3</sub> has been previously shown to be significantly elevated by a much larger factor in all FD groups, and was observed in this study to increase by approximately 2.9-fold ( $p < 0.05$ ) in females and 25-fold in males ( $p < 0.01$ ). CDH was also observed to be elevated in all FD groups but by a greater fold change in the FD female groups by >2.7-fold in the asymptomatic group ( $p < 0.05$ ) and 17-fold in the male FD group ( $p < 0.01$ ). Statistical analyses of grouping female patients by ERT status as opposed to grouping according to symptomatic status, showed negligible differences (supplementary figure S4). Therefore, grouping by symptomatic status with an asymptomatic female group was used to identify GSLs that could potentially aid in biochemical identification of suspected female FD patients.



**Figure 3. Comparison of each GSL species in Fabry patients.** (i) Mean  $\pm$  SEM plasma values of CMH, CDH, Gb<sub>3</sub>, and Gb<sub>4</sub>. Significance in the Fabry female group for plasma is determined by one way ANOVA and for Fabry males by unpaired t-test (ii) Mean  $\pm$  SEM urine values for the lower abundant GSL species CMH and Gb<sub>4</sub> in urine. (iii) Mean  $\pm$  SEM urine values for the higher abundant GSL species CDH and Gb<sub>3</sub> in urine. Significance of urine GSLs was determined by use of non-parametric Kruskal Wallis for the Fabry female groups and Mann-Witney test for the Fabry male group comparison. \*  $p < 0.05$ , \*\*  $p < 0.01$ . AS Fabry Females = asymptomatic Fabry females, S Fabry Females = Symptomatic Fabry Females, Ctl Male = Control males, Ctl Female = Control females.

### Relationship between GSLs and their isoforms in Plasma and Urine

All various GSLs and their isoforms analysed in both FD patient plasma and urine were subjected to Pearson Correlation analysis in order to highlight the relationships between the various GSL species and isoforms. As control samples have much lower GSLs in urine, this analysis was only performed on the FD patient GSL values. The plasma correlation matrix demonstrated (Figure 2), as expected, a strong relationship of the GSL isoforms within their own GSL species group i.e. Gb<sub>3</sub> isoforms correlate with all other Gb<sub>3</sub> isoforms. However, a poor relationship was observed between the individual GSL group species. CDH appears to show a greater degree of correlation with CMH, Gb<sub>3</sub>, and Gb<sub>4</sub>. Lyso-Gb<sub>3</sub> and analogues [17] were assayed in plasma in order to see if there may be a particular Gb<sub>3</sub> isoform that is more susceptible to conversion to lyso-Gb<sub>3</sub>, therefore giving an indication of the mechanism of generation of lyso-Gb<sub>3</sub>. The area of strongest correlation with lyso-Gb<sub>3</sub> is indicated on the plasma correlation matrix and does demonstrate a strong correlation with Gb<sub>3</sub>. In particular, the shorter isoform chains of Gb<sub>3</sub> demonstrated the highest correlation with lyso-Gb<sub>3</sub> levels and particularly the Gb<sub>3</sub> isoforms of C16 and C18 (supplementary data figure S-1).

In the urine analyses, no significant relationship was observed between the GSL species. However, a weak relationship was observed between CDH and Gb<sub>3</sub> and particularly between the C24:1 and C24:2 hydroxylated isoforms of CDH and Gb<sub>3</sub> (urine correlation matrix plot figure 2ii). This may indicate this modification is independent to the expression of the GSLs.



Figure 4. Pearson correlation matrix of all the glycolipids quantitated in plasma and urine of Fabry patients (i). Red indicating a strong correlation and blue a negative correlation between each species. Plasma correlation shows no significant correlation between each of the individual glycolipids with each other. Lyso-Gb<sub>3</sub> is highlighted in the boxed area and demonstrates a stronger correlation with Gb<sub>3</sub> and more so with the short chain isoforms C16, C18 of Gb<sub>3</sub>. (ii) Urine analyses also demonstrated no relationship in expression between the GSL species. However, the hydroxylated long-chain species of CDH and Gb<sub>3</sub> appear to show a correlation between each other (highlighted in box area).


ACCEPTED MANUSCRIPT

## Multivariate analysis of Glycosphingolipidomic screen of Female Patients with Fabry Disease

In order to identify which GSLs, or specific GSL isoforms, could have potential in aiding the diagnosis of FD in females, a multivariate analysis of the data was also performed. Figure 4 shows Principal Component Analysis (PCA) of both urine (A) and plasma (B). All samples demonstrated a separate clustering of the control and FD patients; the first two components (t[1] and t[2]) explaining 55% and 12% of the variation for plasma and the first two components (t[1] and t[2]) explain 41% and 12% of the variation in urine. The FD groups demonstrated a far more varied distribution in both plasma and urine. The male group particularly shows a wide clustering in urine, more so than plasma and which is unsurprising as GSLs were observed to be affected more in the urine of FD patients.

In an attempt to identify female specific markers, a further statistical OPLS-DA analysis was performed on the FD female samples. A significant separation in both the urine and plasma GSL profiles was observed (OPLS-DA parameters are given in supplementary data table S-6). Interestingly, there appeared to be no clustering of the GSL profiles based on symptomatic status in FD females in urine or plasma. As shown by corresponding loading plots the GSLs in the plasma analyses that appear to drive the changes were increased levels of C18 CDH and Gb<sub>4</sub> and reduced levels of all CMH isoforms. In urine, there was a clear GSL profile for control and FD females which was driven by increased levels of lyso-Gb<sub>3</sub>, lyso-Gb<sub>3</sub>-analogues and longer chain CDH isoforms C22:1, C22:0, C22:1-OH, C22:0-OH, C24:2, C24:0, C24:2-OH, C24:1-OH, C24:0-OH, C26:0.





**Figure 5. Multivariate analysis of GSLs in Fabry female patients.** Scatter (top) and corresponding loading plots (bottom) of GSL PCA analysis of male and female control and Fabry patient urine (A) and plasma (B). Further sub-analysis of Fabry female patient samples using OPLS-DA is shown for urine (C) and plasma (D).

## **Glycosphingolipids biomarkers that could aid in the identification of female Fabry Disease Patients**

Multivariate analysis indicated that the urinary long chain CDH fatty acid isoforms C22:1, C22:0, C22:1-OH, C22:0-OH, C24:2, C24:0 C24:2-OH, C24:1-OH, C24:0-OH and C26:0 contributed the greatest to driving the differences between FD females and controls. In plasma, lyso-Gb<sub>3</sub> analogues and also total CMH levels contributed to the changes in FD female patients.

Analogues of lyso-Gb<sub>3</sub> are more predominant in urine than lyso-Gb<sub>3</sub> (supplementary figure S2C) therefore analogue levels were included in the total levels reported. Lyso-Gb<sub>3</sub> showed a significant difference in symptomatic FD females in plasma but less significantly for the asymptomatic female group in urine. In plasma, lyso-Gb<sub>3</sub> demonstrated only a small change in the asymptomatic female group. This indicated that lyso-Gb<sub>3</sub> is a marker of disease severity but offers little information in regards to confirming a diagnosis in asymptomatic heterozygous FD females. However, total urine CDH levels can be used to help identify FD females (figure 5(iii)) and in particular, the use of the 10 long chain isoforms of CDH C22:1, C22:0, C22:1-OH, C22:0-OH, C24:2, C24:0 C24:2-OH, C24:1-OH, C24:0-OH, C26:0 which demonstrate even better specificity ( $p < 0.001$ ) than total CDH ( $p < 0.05$ ) in its ability to discriminate FD female patients. Importantly, the asymptomatic female group who are the most difficult to diagnose, are more distinguishable from female controls when quantitating the longer chain CDH isoforms compared to using plasma lyso-Gb<sub>3</sub> (figure 6). A ROC analysis (figure 6B) of plasma CMH lyso-Gb<sub>3</sub> and urine total CDH and long chain CDH demonstrated the urine long chain CDH has the highest AUC 0.88 for detecting female FD patients.




Figure 6. Comparison of Female Fabry biomarkers (AS Fabry Females = asymptomatic Fabry females, S Fabry Females = Symptomatic Fabry Females, Ctl Male = Control males, Ctl Female = Control females). **A** Currently used biomarkers lyso-Gb<sub>3</sub> in urine and plasma show that plasma lyso-Gb<sub>3</sub> is more specific and has even greater significance in symptomatic patients. However long chain CD also have comparable ability to distinguish asymptomatic female patients from controls. **B** ROC analysis comparing sensitivity and specificity of all GSLs altered in Female Fabry patients. Significance was determined by use of non-parametric Kruskal Wallis for the Fabry female groups and Mann-Witney test for the Fabry male group comparison. \*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ , \*\*\*\*  $p < 0.0001$ .

## Discussion

This glycosphingolipidomic assay has provided significant amounts of useful data regarding GSL catabolism in humans and allowed us to obtain a global overview of GSL composition and abundance in both urine and plasma. To our knowledge, this is the first specific analyses in human samples targeting specifically all those GSLs directly upstream and downstream of the catabolic enzyme defect observed in FD. We did not include the gangliosides GM1, GM2, and GM3 as they are present in very low amounts. Whilst their degradation pathway is not directly related to Gb<sub>3</sub> catabolism (figure 1), their expression would be interesting to observe in FD to see if this branch of the GSL degradation pathway is affected.

The mechanism of generation of lyso-Gb<sub>3</sub> in FD by non-specific action of acid ceramidase has been proposed but the mechanism as yet not fully understood [3]. Multiple correlation analysis in figure 4 shows that lyso-Gb<sub>3</sub> and some of its analogues have the strongest correlation with the Gb<sub>3</sub> isoforms C16:1 and C18:0 ( $r^2 > 0.7$ ,  $p < 0.0001$ ). This relationship indicates these Gb<sub>3</sub> isoforms may be more likely to be converted to lyso-Gb<sub>3</sub> and/or its analogues. Interestingly, these isoforms are not the abundant isoforms of Gb<sub>3</sub> in plasma. Supplementary figure S2 shows the composition of the Gb<sub>3</sub> isoforms and lyso-Gb<sub>3</sub> analogues [17] and demonstrates that the C16:1 and C18:0 isoforms of Gb<sub>3</sub> are the 5<sup>th</sup> and 6<sup>th</sup> most abundant isoforms. In total they comprise approximately 7% and 5% of total Gb<sub>3</sub>. This observation needs to be verified by other independent cohorts but these isoforms may hint at a mechanism of lyso-Gb<sub>3</sub> formation which may or may not involve acid ceramidase [16, 18].

It was not possible to include lyso-Gb<sub>3</sub> in any GSL composition analyses because the levels are much lower (ng/ml range) compared to GSLs (µg/ml range). Figure 2 showed that there are significant differences in the urine GSL composition between males and females that are unchanged in FD. In addition, Figure 3 demonstrated that there are significant changes in the total levels of CDH and Gb<sub>3</sub> in both males and females. This observation that total levels increase but the composition does not, indicate that CDH is probably an equally an important substrate for alpha-galactosidase and is more relevant in Female FD than Gb<sub>3</sub>. This explains previous reports that have observed increased CDH in female FD patients [14, 16]. This study also indicated that Gb<sub>4</sub> and CMH, the upstream and downstream glycolipids to Gb<sub>3</sub> respectively, are more abundant in plasma and form a much lower proportion of the GSL profile in urine. Thus, this indicating that the kidney is probably the primary source of CDH and Gb<sub>3</sub>. In FD disease, plasma Gb<sub>4</sub> levels were unaffected but the downstream glycolipid, CMH, was increased. This observation was unexpected as the CDH to CMH step is downstream of the defect observed in the GSL degradation pathway caused by FD. Therefore, there may be another mechanism/pathway in FD tissues that causes this unpredicted increase of CMH. As

in urine, the predominant GSLs in plasma were Gb<sub>3</sub> and CDH which do not appear to change significantly in FD and confirms previous findings on the use of plasma Gb<sub>3</sub> as a poor biomarker for FD disease [19].

Ferraz *et al* used various mouse models of GSL degradation disorders and measured the main GSL and lyso compounds. In tissues from the FD model mouse, it was observed there were no other sphingolipid changes in the FD mouse model apart from Gb<sub>3</sub> and lyso-Gb<sub>3</sub>. CMH was not affected and indicating no feedback on GSL synthesis to avoid Gb<sub>3</sub> accumulation [12]. However, our data in humans demonstrated that CMH is increased in plasma and therefore indicates that there may be subtle different pathways between mice and human glycosphingolipid homeostasis. Kamani *et al* used a FD mouse model and also measured other products of the GSL degradation pathway including the mono and dihexacylceramides (CDH) [11] and also did not observe an elevation in CMH. However, significant elevation of CDH was observed in the kidney tissue of FD mice and confirming our observations in humans in this study.

GSL species comprise of multiple isoforms based on modification of the fatty acid chain. Little is understood about the biological relevance of each isoform. We took an unbiased analysis approach by using multivariate statistics to see if there were isoforms within the GSL profile with greater specificity for FD females. We confirmed plasma lyso-Gb<sub>3</sub> and also identified urinary long chain CDH isoforms as having greater changes in FD females, indicating they may have use clinically. Subsequent univariate analysis confirmed plasma lyso-Gb<sub>3</sub> as an excellent marker relevant to disease progression. However, the long chain urinary CDH isoforms were highly significant ( $p < 0.001$ ) and more so than lyso-Gb<sub>3</sub> ( $p < 0.01$ ) for the asymptomatic female group, indicating the long chain CDH isoforms could also be useful for identifying FD females. Gb<sub>3</sub> and CDH are the predominant GSLs in urine and this is demonstrated in figure 3. In control females, the percentage of CDH is ~79% and Gb<sub>3</sub> is ~22% whilst in males, these values are ~59% CDH and ~35% Gb<sub>3</sub>. In FD, this composition only changes for males with an increase of Gb<sub>3</sub> to 40%. This indicates the involvement of CDH could also be exploited for the diagnosis and as a feature in FD as both GSLs increase in FD to a similar degree in urine.

The analysis of CDH has been performed previously in FD heterozygotes [14] but is confounded by the fact that LacCer and Ga<sub>2</sub> are structural isomers with the same mass. Both LacCer and Ga<sub>2</sub> differ by configuration at a single chiral centre where the LacCer sugar moiety consists of a glucose and a galactose and Ga<sub>2</sub> consists of 2 galactose units (figure 1). These two isomers do not separate by conventional reverse phase chromatography. Boutin *et al* developed a method using normal phase chromatography to separate LacCer and Gb<sub>2</sub> and assessed the two isomers in FD disease urine [16]. They described that Gb<sub>2</sub> was not significantly different between males and females but LacCer was.

Therefore, it is likely LacCer is the driving dihexosylceramide GSL that contributes to our observation of differing CDH levels between females and males in this study. Boutin *et al* did not describe any significant changes in the levels of LacCer in FD but did for Ga<sub>2</sub>. The limitation of our study is that LacCer levels, although not altered in FD according to Boutin *et al*, could affect the sensitivity and specificity of Ga<sub>2</sub> detection. However, the Boutin *et al* study showed that hydroxylated and long chain LacCer could not be detected in urine. Therefore, the long chain CDH isoforms described in our study are likely to be comprised entirely of Ga<sub>2</sub> isoforms and this explains why the long chain isoforms are more specific at detecting the FD patients. The normal phase LC-MS/MS method developed by Boutin *et al* may overall be a more specific method for the analysis of Ga<sub>2</sub>, LacCer, and Gb<sub>3</sub> isoforms. However due to issues with complex matrices and throughput the use of normal phase methodology such as HILIC or amide column chromatography is not widespread compared to reverse phase chromatographic methods in clinical laboratories. Longer equilibration is required for HILIC analysis compared to reverse phase thereby increasing LC-MS/MS run times which makes it less attractive for high throughput analysis in many laboratories and exact buffer composition has to be tightly controlled which can be problematic for routine analysis [17, 20-22]. A clinical method to quantitatively measure lyso-compounds has been described previously using reverse phase C18 chromatography. The authors also describe that the use of a normal phase amide chromatography method was found only suitable as a qualitative second-tier method to separate the two hexosylsphingosines, glucosylsphingosine and galactosylsphingosine, that are relevant to Gaucher and Krabbe disease [15]. Our study highlights that by analysing the long chain CDH isoforms the well-established standard reverse phase methodology already employed in most diagnostic laboratories to diagnose Fabry patients could be easily augmented to include this additional biomarker in future FD studies. Our study however was unable to confirm that Gb<sub>3</sub> could discriminate the FD female patients unlike Boutin *et al* who showed that urinary Gb<sub>3</sub> was better than Ga<sub>2</sub>. Urinary Gb<sub>3</sub> has been shown previously to be useful as a biomarker in FD disease but less so for the identification of females and has largely been superseded by plasma lyso-Gb<sub>3</sub> as the biomarker of choice for diagnosing and monitoring treatment for heterozygotes [19]. To test the specificity of CDH, we compared its levels with plasma and urine lyso-Gb<sub>3</sub> (plus analogues) [23]. Urine lyso-Gb<sub>3</sub> appears to perform poorly compared to plasma lyso-Gb<sub>3</sub> and is likely due to the lower levels of lyso-Gb<sub>3</sub> in urine. In our study, plasma lyso-Gb<sub>3</sub> appears to be more specific for the symptomatic FD female group confirming its association with disease severity/progression [24]. It does, however, have limited use in detecting the asymptomatic heterozygote females. Suspected female FD patients all undergo biochemical confirmation by analysis of urinary Gb<sub>3</sub> and lyso-Gb<sub>3</sub>. Therefore, we propose that urinary Ga<sub>2</sub> (long chains) could be easily added into existing LC-MS/MS Gb<sub>3</sub> assays. One of the interesting questions

raised from these results is what the potential contribution to the disease pathology could come from raised Ga<sub>2</sub> levels? Does Ga<sub>2</sub> also have a lyso-compound that can also cause disease pathology like lyso-Gb<sub>3</sub> has been shown to do [25, 26]?

**Conclusions:** Urinary CDH is the more prominent GSL in females which means changes associated with FD are more likely to be detected. We demonstrate that the longer chain CDH isoforms C22:1, C22:0, C22:1-OH, C22:0-OH, C24:2, C24:0 C24:2-OH, C24:1-OH, C24:0-OH, C26:0 of Ga<sub>2</sub> in urine are elevated in FD and can discriminate asymptomatic heterozygotes better than Gb<sub>3</sub> and lyso-Gb<sub>3</sub>. We propose it may be more powerful to combine and analyse lyso-Gb<sub>3</sub> and Ga<sub>2</sub> together, particularly for females where there is an uncertainty of diagnosis. Furthermore, this could easily be incorporated into the conventionally used LC-MS analysis being employed in chemical pathology laboratories.

**Acknowledgments:** This study has been part funded by the UCL Biological Mass Spectrometry Centre. All research at Great Ormond Street Hospital NHS Foundation Trust and UCL Great Ormond Street Institute of Child Health is made possible by the NIHR Great Ormond Street Hospital Biomedical Research Centre. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. We would also like to thank the Peto foundation for their generous donations.

## References

- [1] I.Y. Desnick R, Eng C,  $\alpha$ -galactosidase A deficiency: Fabry disease, McGraw-Hill, New York, 2001.
- [2] A. Nowak, T.P. Mechtler, T. Hornemann, J. Gawinecka, E. Theswet, M.J. Hilz, D.C. Kasper, Genotype, phenotype and disease severity reflected by serum LysoGb3 levels in patients with Fabry disease, *Molecular genetics and metabolism*, 123 (2018) 148-153.
- [3] M.J. Ferraz, A.R. Marques, M.D. Appelman, M. Verhoek, A. Strijland, M. Mirzaian, S. Scheij, C.M. Ouairy, D. Lahav, P. Wisse, H.S. Overkleeft, R.G. Boot, J.M. Aerts, Lysosomal glycosphingolipid catabolism by acid ceramidase: formation of glycosphingoid bases during deficiency of glycosidases, *FEBS letters*, 590 (2016) 716-725.
- [4] B.E. Smid, L. van der Tol, M. Biegstraaten, G.E. Linthorst, C.E. Hollak, B.J. Poorthuis, Plasma globotriaosylsphingosine in relation to phenotypes of Fabry disease, *Journal of medical genetics*, 52 (2015) 262-268.
- [5] A. Nowak, T.P. Mechtler, R.J. Desnick, D.C. Kasper, Plasma LysoGb3: A useful biomarker for the diagnosis and treatment of Fabry disease heterozygotes, *Molecular genetics and metabolism*, 120 (2017) 57-61.
- [6] F.J. Alharbi, S. Baig, C. Auray-Blais, M. Boutin, D.G. Ward, N. Wheeldon, R. Steed, C. Dawson, D. Hughes, T. Geberhiwot, Globotriaosylsphingosine (Lyso-Gb3) as a biomarker for cardiac variant (N215S) Fabry disease, *J Inherit Metab Dis*, 41 (2018) 239-247.
- [7] D. Singh, H.C. Jarrell, E. Florio, D.B. Fenske, C.W. Grant, Effects of fatty acid alpha-hydroxylation on glycosphingolipid properties in phosphatidylcholine bilayers, *Biochimica et biophysica acta*, 1103 (1992) 268-274.
- [8] A.H. Merrill, Jr., Sphingolipid and glycosphingolipid metabolic pathways in the era of sphingolipidomics, *Chem Rev*, 111 (2011) 6387-6422.

- [9] C. Auray-Blais, M. Boutin, R. Gagnon, F.O. Dupont, P. Lavoie, J.T. Clarke, Urinary globotriaosylsphingosine-related biomarkers for Fabry disease targeted by metabolomics, *Analytical chemistry*, 84 (2012) 2745-2753.
- [10] C.A. Lingwood, Glycosphingolipid functions, *Cold Spring Harbor perspectives in biology*, 3 (2011).
- [11] M.A. Kamani, P. Provencal, M. Boutin, N. Pacienza, X. Fan, A. Novak, T.C. Huang, B. Binnington, B.C. Au, C. Auray-Blais, C.A. Lingwood, J.A. Medin, Glycosphingolipid storage in Fabry mice extends beyond globotriaosylceramide and is affected by ABCB1 depletion, *Future Sci OA*, 2 (2016) FSO147.
- [12] M.J. Ferraz, A.R. Marques, P. Gaspar, M. Mirzaian, C. van Roomen, R. Ottenhoff, P. Alfonso, P. Irun, P. Giraldo, P. Wisse, C. Sa Miranda, H.S. Overkleeft, J.M. Aerts, Lyso-glycosphingolipid abnormalities in different murine models of lysosomal storage disorders, *Molecular genetics and metabolism*, 117 (2016) 186-193.
- [13] M. Boutin, C. Auray-Blais, Multiplex tandem mass spectrometry analysis of novel plasma lyso-Gb(3)-related analogues in Fabry disease, *Analytical chemistry*, 86 (2014) 3476-3483.
- [14] K. Mills, P. Morris, P. Lee, A. Vellodi, S. Waldek, E. Young, B. Winchester, Measurement of urinary CDH and CTH by tandem mass spectrometry in patients hemizygous and heterozygous for Fabry disease, *J Inherit Metab Dis*, 28 (2005) 35-48.
- [15] G. Polo, A.P. Burlina, T.B. Kolamunnage, M. Zampieri, C. Dionisi-Vici, P. Strisciuglio, M. Zaninotto, M. Plebani, A.B. Burlina, Diagnosis of sphingolipidoses: a new simultaneous measurement of lysosphingolipids by LC-MS/MS, *Clinical chemistry and laboratory medicine : CCLM / FESCC*, 55 (2017) 403-414.
- [16] M. Boutin, I. Menkovic, T. Martineau, V. Vaillancourt-Lavigne, A. Toupin, C. Auray-Blais, Separation and Analysis of Lactosylceramide, Galabiosylceramide, and Globotriaosylceramide by LC-MS/MS in Urine of Fabry Disease Patients, *Analytical chemistry*, 89 (2017) 13382-13390.
- [17] M. Boutin, Y. Sun, J.J. Shacka, C. Auray-Blais, Tandem Mass Spectrometry Multiplex Analysis of Glucosylceramide and Galactosylceramide Isoforms in Brain Tissues at Different Stages of Parkinson Disease, *Analytical chemistry*, 88 (2016) 1856-1863.
- [18] V. Manwaring, M. Boutin, C. Auray-Blais, A metabolomic study to identify new globotriaosylceramide-related biomarkers in the plasma of Fabry disease patients, *Analytical chemistry*, 85 (2013) 9039-9048.
- [19] E. Young, K. Mills, P. Morris, A. Vellodi, P. Lee, S. Waldek, B. Winchester, Is globotriaosylceramide a useful biomarker in Fabry disease?, *Acta paediatrica*, 94 (2005) 51-54; discussion 37-58.
- [20] M. Sarbu, A.D. Zamfir, Modern separation techniques coupled to high performance mass spectrometry for glycolipid analysis, *Electrophoresis*, 39 (2018) 1155-1170.
- [21] P. Jandera, T. Hajek, Mobile phase effects on the retention on polar columns with special attention to the dual hydrophilic interaction-reversed-phase liquid chromatography mechanism, a review, *J Sep Sci*, 41 (2018) 145-162.
- [22] S. Wernisch, S. Pennathur, Evaluation of coverage, retention patterns, and selectivity of seven liquid chromatographic methods for metabolomics, *Anal Bioanal Chem*, 408 (2016) 6079-6091.
- [23] C. Auray-Blais, A. Ntwari, J.T. Clarke, D.G. Warnock, J.P. Oliveira, S.P. Young, D.S. Millington, D.G. Bichet, S. Sirrs, M.L. West, R. Casey, W.L. Hwu, J.M. Keutzer, X.K. Zhang, R. Gagnon, How well does urinary lyso-Gb3 function as a biomarker in Fabry disease?, *Clinica chimica acta; international journal of clinical chemistry*, 411 (2010) 1906-1914.
- [24] S.M. Rombach, N. Dekker, M.G. Bouwman, G.E. Linthorst, A.H. Zwinderman, F.A. Wijburg, S. Kuiper, M.A. Vd Bergh Weerman, J.E. Groener, B.J. Poorthuis, C.E. Hollak, J.M. Aerts, Plasma globotriaosylsphingosine: diagnostic value and relation to clinical manifestations of Fabry disease, *Biochimica et biophysica acta*, 1802 (2010) 741-748.
- [25] J.M. Aerts, J.E. Groener, S. Kuiper, W.E. Donker-Koopman, A. Strijland, R. Ottenhoff, C. van Roomen, M. Mirzaian, F.A. Wijburg, G.E. Linthorst, A.C. Vedder, S.M. Rombach, J. Cox-Brinkman, P.




Somerharju, R.G. Boot, C.E. Hollak, R.O. Brady, B.J. Poorthuis, Elevated globotriaosylsphingosine is a hallmark of Fabry disease, *Proc Natl Acad Sci U S A*, 105 (2008) 2812-2817.

[26] L. Choi, J. Vernon, O. Kopach, M.S. Minett, K. Mills, P.T. Clayton, T. Meert, J.N. Wood, The Fabry disease-associated lipid Lyso-Gb3 enhances voltage-gated calcium currents in sensory neurons and causes pain, *Neuroscience letters*, 594 (2015) 163-168.

ACCEPTED MANUSCRIPT

## Graphical abstract



**Highlights**

- Ceramide dihexoside (CDH) is the predominant glycosphingolipid in females
- Plasma ceramide monhexoside is increased in Fabry disease
- Urinary long-chain CDH isoforms are better than lyso-Gb3 to distinguish Fabry females

ACCEPTED MANUSCRIPT