

**Effectiveness of BCG vaccination against *Mycobacterium tuberculosis* infection in adults: a cross-sectional analysis of a UK-based cohort**

**Authors**

Anthea L Katelaris, Charlotte Jackson, Jo Southern, Rishi K Gupta, Francis Drobniowski, Ajit Lalvani, Marc Lipman, Punam Mangtani\*, Ibrahim Abubakar\*

\*Joint last authors

**Affiliations**

Anthea L Katelaris, Faculty of Public Health and Policy, London School of Hygiene and Tropical Medicine, London WC1E 7HT, UK

Charlotte Jackson, Institute for Global Health, University College London, London WC1E 6BT, UK

Jo Southern, National Infection Service, Public Health England, London NW9 5EQ, UK

Rishi K Gupta, Institute for Global Health, University College London, London WC1E 6BT, UK

Francis Drobniowski, Department of Medicine, Imperial College, London W12 0NN, UK

Ajit Lalvani, National Institute for Health Research Health Protection Research Unit in Respiratory Infections, Imperial College London W2 1PG, UK

Marc Lipman, UCL Respiratory, Division of Medicine, University College London, London, WC1E 6BT, UK

Punam Mangtani, Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, London WC1E 7HT, UK

Ibrahim Abubakar, Institute for Global Health, University College London, London, UK  
WC1E 6BT, UK

### **Corresponding author**

Dr Charlotte Jackson, Institute for Global Health, University College London, London; email: [c.r.jackson@ucl.ac.uk](mailto:c.r.jackson@ucl.ac.uk); phone: +44 (0)20 7670 4806.

### **Running head**

BCG effectiveness against *Mtb* infection

### **Summary statement**

This analysis found BCG was associated with a lower prevalence of LTBI (measured via IGRA) in adults with recent exposure to active tuberculosis. These results suggest BCG may provide durable protection against *Mtb* infection as well as disease.

### **Word count:**

Abstract: 200

Main text: 3507

## **ABSTRACT**

### **Background**

BCG appears to reduce acquisition of *Mycobacterium tuberculosis* (*Mtb*) infection in children, measured using interferon-gamma release assays (IGRAs). We explored whether BCG vaccination continues to be associated with decreased prevalence of *Mtb* infection in adults.

### **Methods**

We conducted a cross-sectional analysis of data from adult contacts of tuberculosis cases participating in a UK cohort study. Vaccine effectiveness (VE) of BCG, ascertained based on presence of a scar or vaccination history, against latent tuberculosis infection (LTBI), measured via IGRA, was assessed using multivariable logistic regression. The effects of age at BCG and time since vaccination were also explored.

### **Results**

Of 3453 recent tuberculosis contacts, 27.5% had LTBI. There was strong evidence of an association between BCG and LTBI (aOR=0.70, 95% CI 0.56-0.87, p=0.0017) yielding a VE of 30%. VE declined with time since vaccination, but there was evidence that LTBI prevalence was lower amongst vaccinated individuals even >20 years after vaccination, compared with non-vaccinated participants.

### **Conclusion**

BCG is associated with lower prevalence of LTBI in adult contacts of tuberculosis. These results contribute to growing evidence that suggests BCG may protect against *Mtb* infection

as well as disease. This has implications for immunisation programmes, vaccine development and tuberculosis control efforts worldwide.

**Key words**

BCG; Bacille Calmette-Guérin vaccine; tuberculosis; vaccine effectiveness

## INTRODUCTION

Tuberculosis (TB) is the leading cause of death from infectious disease worldwide.[1] The widely used Bacillus Calmette-Guérin (BCG) vaccine is the only licensed vaccine against *Mycobacterium tuberculosis* (*Mtb*).[2] Infant BCG vaccination has shown consistently high efficacy of 70-80% against childhood TB, namely meningitis and miliary TB.[3] The protective effect of BCG against adult pulmonary disease varies geographically,[4–6] which may in part be associated with varying exposure to *Mtb* or environmental mycobacteria [7,8], which may mask or block protection induced by BCG.[4,9–11]

Until recently it was not possible to determine if BCG vaccination prevents acquisition of *Mtb* infection or only limits progression from latent TB infection (LTBI) to active disease. This was due to limitations of the tuberculin skin test (TST), which can be positive following BCG vaccination and, to a lesser extent, exposure to environmental mycobacteria.[12,13] More recently developed interferon-gamma release assays (IGRAs) are more specific for *Mtb* infection,[14] as they measure interferon-gamma release from T-lymphocytes stimulated with antigens not present in BCG or most environmental mycobacteria.[15] Since the protective effect of BCG against *Mtb* infection was first demonstrated[16], several observational studies have investigated this phenomenon. A meta-analysis of 14 observational studies in children aged <16 years with recent exposure to active TB found a vaccine effectiveness (VE) of 19% against *Mtb* infection. In the six studies which followed up children who were IGRA-positive, BCG was associated with a 58% reduction in the risk of active TB.[17]

Data on whether this protection against *Mtb* infection continues into adulthood, as it appears to against disease,[18,19] are limited. Several observational studies of adults have found weak evidence of protection by BCG against infection measured via IGRA.[20–25] There has been no investigation of the influence of age at vaccination on VE of BCG against LTBI in

adults, or of changes in protection over time. These issues are important as, in order to assess the role of BCG (or similar vaccines in development) in TB control programmes, the effect of BCG on the full spectrum of TB should be understood.[26] In this cross-sectional study, we used baseline data from a large United Kingdom (UK) cohort study of adults at risk of *Mtb* infection to determine the presence and durability of BCG protection against *Mtb* infection, assessed by IGRA, and the potential factors that influence VE.

## **METHODS**

### **Study design**

A cross-sectional study was carried out amongst participants recruited to the UK PREDICT study who were recent contacts of patients with active TB ('contacts'). PREDICT was a prospective cohort study that aimed to assess the prognostic value of IGRAs in predicting the development of active TB among individuals with, or at risk of, LTBI, as previously described.[27] 10045 TB contacts, as well as recent migrants, aged  $\geq 16$  years were recruited between January 2011 and July 2015. Participants with evidence of active TB at baseline were excluded, and migrants were excluded from this analysis as it was unclear when their primary infection occurred. Contacts were recruited at contact tracing appointments for TB screening in TB clinics (part of the routine public health management of TB in the UK).

After obtaining informed consent, study nurses completed a questionnaire with participants, including on demographic, social, medical and TB exposure history, took blood samples for IGRA testing, and administered a Mantoux TST (which was read 48-72 hours later).

Most recruitment sites were in London, with one clinic each in Birmingham and Leicester. To increase the probability that any detected LTBI was due to recent infection, participants who reported previous contact with active TB (prior to that resulting in recruitment) were

excluded. Participants who reported a previous TB diagnosis were also excluded from this analysis, as a positive IGRA may reflect their prior TB disease rather than current LTBI.

### **Primary exposure, outcome and covariates**

The primary exposure of interest was previous BCG vaccination. Vaccination status was determined by inspection of both arms for a vaccination scar by a trained study or TB nurse, combined with a vaccination history and documentation from a personally-held vaccination record (e.g. the 'red book') if available. A positive response was recorded when a scar was observed or there was documented evidence of BCG administration, and a negative response recorded if no scar was seen. If participants thought they had been vaccinated but no scar was observed, the interviewer recorded 'unsure' and this was treated as missing. Participants who had received vaccination were asked the year of vaccination.

The outcome of interest was LTBI, measured via IGRA. LTBI was defined as a positive result on either or both of QuantiFERON-TB Gold In-Tube (QFT-GIT) and T-SPOT.TB. Those who were negative on both assays, or negative on one and indeterminate on the other, were considered not to have LTBI. Most participants had both IGRAs performed, however a small number were only tested with QFT-GIT, and LTBI was determined based on the single assay. If both assays were indeterminate, the outcome was considered missing.

Self-reported questionnaire data included details of the TB exposure, country of birth, ethnicity (based on the Enhanced Tuberculosis Surveillance system categories[28]), smoking status (never or ever), social risk factors (none, or any of homelessness, imprisonment or use of controlled drugs), and medical details, including whether the participant had diabetes, HIV or other immunosuppression.

Age at vaccination was calculated as the difference between vaccination and birth year, and

dichotomised based on the ages of vaccination recommended in typical BCG vaccination policies ( $\leq 2$  years to reflect infant vaccination,  $>2$  years for vaccination in childhood and at older ages). Time since vaccination was calculated as age at recruitment minus the stated age at vaccination, and grouped into three categories to avoid data sparsity ( $\leq 10$  years, 11-20 years,  $>20$  years).[5] TB incidence in country of birth was obtained from WHO estimates for the year 2000.[29] Absolute latitude of country of birth, found to affect VE in previous studies[9], was calculated using average country latitude data from Google Public Data Explorer[30], and collapsed into  $20^\circ$  groups.[9,17]

### **Statistical analysis**

To investigate the association between LTBI and BCG status (and all other covariates), cross-tabulation, unadjusted ORs and likelihood ratio tests (LRTs) were calculated. For variables with no natural reference group, the largest group was used as the baseline.

Multivariable analysis was performed using logistic regression and LRTs. Age group and sex were considered *a priori* confounders. Age group was treated as a categorical, rather than continuous, variable as this produced a better fit in bivariable analysis. Further confounding variables were included in the multivariable model based on the change-in-estimate method. All variables that were associated with BCG status or LTBI on bivariable analysis with  $p \leq 0.2$  (with none deemed to lie on the causal pathway) were added separately to the model. Any variable that resulted in a  $\geq 10\%$  relative change in the OR was kept in the model (with the variable that caused the largest change in the OR selected first if more than one variable caused a  $\geq 10\%$  change). All other variables were then individually re-added to the model until the addition of no further variables resulted in a 10% change in OR. All analyses used a 'complete-case' approach. Vaccine effectiveness for LTBI was calculated as  $VE = 1 - OR$ . This



OR is the prevalence odds ratio[31], which approximates the incidence rate ratio in cross-sectional studies.[32,33]

The roles of age at vaccination and time since vaccination on the association between BCG and LTBI were then explored. Two further multivariable logistic regression models were fitted separately (for age at BCG and time since BCG), using the same modelling strategy as above. The analysis of age at BCG was stratified by place of birth (UK or non-UK). Due to collinearity between age at vaccination and time since vaccination, analysis of time since BCG was stratified by age at BCG ( $\leq 2$  years or  $> 2$  years).

Six sensitivity analyses of the primary multivariable model were performed: 1) with additional adjustment for smoking status and any social risk factor; 2) including adjustment for TB incidence of country of birth; 3) including only participants with concordant IGRAs; 4) using binomial regression with a log-link to directly estimate the prevalence ratio (PR), which may not be well approximated by the OR as the outcome was common[34]; 5) restricting the analysis to household contacts of active TB cases, who likely had the most defined single TB exposure[35]; 6) including participants who reported earlier contact with people with active TB. We also performed additional analyses using TST results to measure LTBI. Two cut-off criteria were used to define positivity: firstly,  $TST \geq 5\text{mm}$ [36]; and secondly,  $TST \geq 5\text{mm}$  in BCG-naïve, or  $\geq 15\text{mm}$  in BCG-vaccinated participants (which better predicts disease progression[27]).

To examine bias from missing data, observations with missing values were investigated for association with the outcome and exposure using cross-tabulation, together with multivariable analysis of factors associated with missing BCG status.

Analysis was conducted using Stata v15.0.

## **RESULTS**

### **Participant inclusion**

Of 10045 participants enrolled in the PREDICT study, 9515 had no evidence of active TB and did not report a prior history of TB. 4310 participants were recent TB contacts, of whom 857 had missing data on LTBI and/or BCG status, leaving 3453 participants in this cross-sectional study (Figure 1).

### **Baseline characteristics and bivariable analysis**

Of 3453 participants, 86.9% (3000/3453) had received BCG vaccination. The median age was 32 years (interquartile range 25-43) and 50.0% were male. 2420/3444 (70.3%) were born outside the UK (Table 1).

The overall prevalence of LTBI was 27.5% (951/3453), 27.0% (809/3000) in those vaccinated and 31.4% (142/453) in the unvaccinated, yielding an unadjusted OR of 0.81 (95% CI 0.65-1.00,  $p=0.054$ ) (Table 1). Characteristics associated with increased LTBI prevalence on univariate analysis were male sex, older age, being born outside the UK, some ethnicities, lack of immunosuppression, having diabetes, lower latitudes of country of birth, TB incidence in country of birth, and household TB exposure.

### **Multivariable analysis**

3399 (98.4%) participants were included in the complete-case analysis. Apart from sex and age (included *a priori*), latitude of country of birth was the key confounding variable, shifting the OR away from the null, and the only covariate retained in the final model. After adjusting for sex, age group and latitude of country of birth, there was strong evidence of an association

between BCG and LTBI (OR 0.70, 95% CI 0.56-0.87,  $p=0.0017$ )(Table 2), thus the VE of BCG for LTBI in adult contacts of TB cases was 30% (95% CI 13-44%).

Sensitivity analyses produced largely similar results (Appendix 1). Additional analyses using TST results found that BCG was positively associated with LTBI using a 5mm TST threshold, and negatively associated using a stratified threshold based on BCG status (OR 0.50, 95% CI 0.40-0.63,  $p<0.001$ , Appendix 1).

There was no evidence of an association between missing LTBI data and BCG vaccination (OR 1.05, 95% CI 0.74-1.49,  $p=0.79$ ). However those without data on BCG status were more likely to have LTBI (OR 1.33, 95% CI 1.09-1.63)(Appendix 2).

### **Vaccine effectiveness by age at vaccination and time since vaccination**

Data on reported age at (and time since) vaccination were available for 2195 (73%) of 3000 BCG-vaccinated participants. Of the 641 people born in the UK, 497 (78%) were vaccinated aged  $>2$  years. Of the 1554 people born outside the UK, 1268 (82%) were vaccinated aged  $\leq 2$  years (Appendix 3). Protection against infection was observed following both infant and older age vaccination, among those born within and outside the UK, after adjusting for age group, sex and latitude of country of birth (Table 3). There was no clear pattern by age at vaccination, as the confidence intervals for the adjusted ORs overlapped.

The association between BCG and LTBI, adjusted for age group, sex, ethnicity and country of birth, appeared to vary with time since vaccination in both age at vaccination strata (Table 4). In those vaccinated at age 2 or younger, the protective association was greater in those vaccinated 11-20 years ago (OR 0.55 [95% CI 0.33-0.90]) than those vaccinated  $>20$  years ago (OR 0.78 [0.60-1.00]). As this was an adult cohort, there were no participants vaccinated  $\leq 10$  years ago in the group vaccinated in infancy. In participants vaccinated at age  $>2$  years,

the protective association was greatest in those vaccinated  $\leq 10$  years ago (OR 0.31 [95% CI 0.15-0.63]), versus those vaccinated 11-20 and  $>20$  years ago (OR 0.73 [0.48-1.10] and 0.67 [0.46-0.98] respectively). However there appeared to be evidence of protection more than 20 years later in both those vaccinated in infancy and those vaccinated in childhood and older ages, compared to those without BCG vaccination.

## **DISCUSSION**

### **Principal findings**

In this cross-sectional study, we found strong evidence of an association between BCG and LTBI in recent adult contacts of TB, with VE estimated as 30% (95% CI 13-44%). Protection against infection was seen following both infant and older age vaccination, among participants born within and outside the UK. The association appeared to differ by time since vaccination, in participants vaccinated at both  $\leq 2$  and  $>2$  years of age. In those vaccinated aged  $\leq 2$  years, protection was seen in those vaccinated 11-20 years ago and, though less so, vaccinated  $>20$  years ago. Among participants vaccinated at ages  $>2$  years, the protective association was greater for those vaccinated  $<10$  years ago than those vaccinated 10-20 years ago, where evidence of a protective effect was weak. In both groups, there remained evidence of some protection in participants vaccinated  $>20$  years ago.

### **Strengths and limitations**

To our knowledge, this is the largest adult study on the association between BCG and LTBI measured via IGRA in recent contacts of TB patients. The large sample size and rich dataset are key strengths. This enabled analysis by strata of age at BCG and time since vaccination and broad sensitivity analyses. Although the analysis of time since vaccination could not be adjusted for age at vaccination (and vice versa) due to collinearity, stratification helped

address this. Using both commercially-available IGRAs allowed a sensitivity analysis of those with concordant results, reducing the potential for outcome misclassification.

There are some limitations in a cross-sectional study. The observational design cannot prove causality, and there may be confounding by other factors such as health-seeking behaviour. Furthermore, as many participants were born in high-burden countries, some may have been exposed to *Mtb* before vaccination. However, to increase the probability that vaccination preceded *Mtb* exposure, we only included recent TB contacts without prior reported TB exposure. Also, most participants born outside the UK were vaccinated in infancy, limiting exposure to *Mtb* before vaccination.

#### *Measurement error*

Similar to prior studies, this study relied predominantly on BCG scar to measure vaccination. BCG does not always result in scar formation, but may still confer protection in these cases.[35,37,38] Therefore some vaccinated participants may have been misclassified as non-vaccinated, if there was no documented evidence otherwise. This would be non-differential with respect to IGRA, potentially biasing our VE estimates towards the null. Data on vaccination year was also likely subject to recall error, but again this is likely non-differential with regards to outcome.

Most data on covariates was self-reported. Prevalence of HIV and smoking were relatively low, possibly reflecting social desirability biases. Residual confounding is also possible. For example, latitude of country of birth was an important confounder and may be a proxy measure of potential exposure to nontuberculous mycobacteria, as well as *Mtb*, but does not take into account the time lived at that latitude. Similarly, TB incidence in country of birth is an incomplete measure of *Mtb* exposure.

There is currently no gold standard for diagnosing LTBI, and a positive IGRA cannot distinguish between true ongoing latent *Mtb* infection with dormant yet viable bacteria and an immunological memory response following exposure to *Mtb*. Long-term follow-up studies (or development of a gold standard test for diagnosing LTBI) would help assess if IGRA does measure true infection preceding risk of disease. We found increased prevalence of TST reaction  $\geq 5$ mm amongst vaccinated individuals, consistent with TST being affected by BCG. Stratified thresholds can improve the specificity of TST, as demonstrated by the protective association seen in our tiered 5/15mm analysis. However IGRAs are a more sensitive and specific surrogate marker for *Mtb* infection in BCG-vaccinated individuals.[14,15]

### *Missing data*

Twenty percent of otherwise eligible participants were excluded due to missing data on LTBI and/or BCG. Missing data on LTBI were predominantly due to logistic factors (e.g. failed blood collection), so likely missing completely at random. This is supported by the lack of association between missing outcome data and BCG vaccination, and is therefore unlikely to be an important source of bias.

BCG vaccination status was missing for 13% of participants, and these people were more likely to have LTBI. As it is harder to be certain of lack of vaccination, more unvaccinated participants may have been recorded as ‘unsure’ and therefore coded as missing (possibly leading to the association of missing BCG status with LTBI). However in these cases, when the missingness mechanisms are related to the exposure and covariates, but not the outcome, a logistic regression complete-case approach can still provide an unbiased estimate of the OR,[39] dependent on the model including the covariates with which missingness is related. While being non-UK born and of Pakistani ethnicity were associated with missing BCG status (and not included in the final model as they were not found to be important

confounders), these associations and numbers were small, so the overall bias of VE from missing BCG status is likely minimal.

### **Comparison with existing literature**

The estimated VE of 30% is broadly consistent with other studies in adults.[20–25] Three studies in TB contacts had higher VEs than this study (ORs 0.11-0.5), though these were small studies restricted to close contacts and had wide confidence intervals that crossed 1.[23–25] The lower VE in our analysis may reflect the limitations discussed above, which could have biased the estimate towards the null.

Although the confidence intervals overlapped, our results suggest a possible decline in protection with increased time since vaccination. This concurs with studies of the duration of protection against active TB.[5,18] Our finding that protection against infection was seen following both infant and older age vaccination is less consistent with other studies reporting that protection against active TB is greater when vaccination occurs at younger ages, in immunologically naïve subjects.[5,9,40] Our findings are likely influenced by several factors. We do not have data on why participants were vaccinated at particular ages. However, those born outside the UK and vaccinated aged >2 years are likely to have been vaccinated after moving to the UK, where they would have been offered school-aged BCG vaccination only after negative TST results under the UK's school-based BCG program, discontinued in 2005.[41] Usually only infant vaccination programs are offered in high TB burden settings. In addition, infant vaccination is less likely to result in scar formation compared with vaccination at older ages.[42] Therefore non-differential misclassification is more likely in those vaccinated earlier, leading to an underestimate of VE in the younger age group. Finally, measurement of these variables was susceptible to bias, due to missing data and recall errors

for age at vaccination. These exploratory results should therefore be interpreted cautiously and confirmed in other studies, including with longitudinal design.

### **Implications and future research**

This study suggests that BCG may protect against the acquisition of *Mtb* infection in adults and not only progression to disease.[17,26] This has implications for TB control strategies. The most recent WHO position paper on BCG reflects the growing evidence that BCG confers “a modest protective effect... against *Mtb* infection, representing a significant additional benefit of the BCG vaccine.”[43] Furthermore our data are consistent with evidence of durability of protection against disease,[18,19] which should be considered in cost-effectiveness and modelling studies by low-burden countries considering moving away from universal BCG vaccination to targeted vaccination of high-risk groups.[43]

The effect that BCG may have on wider transmission dynamics is unclear, though preliminary modelling studies suggest a vaccine that protects against *Mtb* infection would have a high population-level impact on TB incidence, across a range of exposure intensities.[44] Future modelling studies of TB transmission and burden should therefore include sensitivity analyses that account for some degree of protection against infection.[43]

Finally, these findings may contribute to understanding the mechanism of action of BCG. This is vital in the development and evaluation of new TB vaccines, since many vaccine candidates rely on a BCG-boosting strategy.[45] Novel vaccines must, and are starting to show they do, confer additional protection beyond the effect of BCG on both *Mtb* infection and disease progression.[46] Prevention of infection measured via IGRA is being examined as a shorter-term endpoint in efficacy trials of new TB vaccines, accelerating their assessment.[47] A recent trial of BCG revaccination versus vaccination with a novel subunit vaccine assessed vaccine efficacy using IGRA conversion as a measure of *Mtb* acquisition



and sustained IGRA positivity to reflect sustained infection.[48] Our study suggests BCG-like vaccines that prevent infection may also provide long-term durability of protection.

Our study contributes to the growing evidence that suggests BCG can act partially by providing protection against *Mtb* infection as well as disease. This has important implications for immunisation programmes, vaccine development and assessment, and TB control efforts worldwide.


## **ACKNOWLEDGEMENTS**

We are grateful to all members of the PREDICT Study Group: Ibrahim Abubakar, David Adeboyeku, Nabeela Bari, Jack Barker, Helen Booth, Graham Bothamley, Felix Chua, Dean Creer, Mathina Darmalingam, Robert N. Davidson, Martin Dediccoat, Jonathan J Deeks, Francis Drobniowski, Anne Dunleavy, Jose Figueroa, Chris Griffiths, Pranab Haldar, Mimi Haseldean, Andrew Hayward, Norman Johnson, Onn Min Kon, Heinke Kunst, Ajit Lalvani, Marc Lipman, Stefan Losewicz, Joanne Lord, William Lynn, Bobby Mann, Heather Milburn, John Moore-Gillon, Geoff Packe, Anton Pozniak, Frances Sanderson, Jo Southern.

We would also like to acknowledge Antonio Gasparini for his advice on the statistical methods.

## TABLES AND FIGURES

**Figure 1:** Recruitment of participants to the PREDICT study and inclusion in cross-sectional study.



TB = tuberculosis; LTBI = latent tuberculosis infection; BCG = Bacillus Calmette-Guérin.

\* Contacts were defined as people with a cumulative duration of exposure to the index case (pulmonary or extra-pulmonary TB) of >8 hours in a confined space during the period of infectiousness.

**Table 1:** Cross sectional study baseline characteristics and unadjusted ORs of their association with LTBI (n=3453).

Variable	Group	Number positive for LTBI	Percentage positive for LTBI	Crude OR	95% CI	p value*
BCG vaccination	No	142/453	31.4	1		
	Yes	809/3000	27.0	0.81	0.65-1.00	0.054
Sex (missing [m]=16)	Male	538/1726	31.2	1		
	Female	410/1711	24.0	0.70	0.60-0.81	<0.001
Age group, years (m=1)	16-25	213/965	22.1	1		
	26-35	328/1175	27.9	1.37	1.12-1.67	
	36-45	195/601	32.5	1.70	1.35-2.13	
	>45	215/711	30.2	1.53	0.24-0.33	<0.001
Country of birth (m=9)	UK	155/1024	15.1	1		
	Non-UK	790/2420	32.6	2.72	2.25-3.29	<0.001
Ethnicity (m=76)	Indian	292/967	30.2	1		
	White	137/709	19.3	0.55	0.44-0.70	
	Black African	190/536	35.5	1.27	1.01-1.59	
	Mixed	145/474	30.6	1.02	0.80-1.29	
	Pakistani	71/245	29.0	0.94	0.69-1.28	
	Bangladeshi	22/142	15.5	0.43	0.26-0.68	
	Black Caribbean	31/162	19.1	0.55	0.36-0.83	
	Black Other / Chinese/Other	35/142	27.3	0.76	0.50-1.13	<0.001
Any social risk factor	No	897/3227	27.8	1		
	Yes	54/226	23.9	0.82	0.59-1.12	0.20
HIV (m=111)	No	913/3317	27.5	1		
	Yes	7/25	28.0	1.02	0.43-2.50	0.96
Other immunosuppression (m=6)	No	928/3338	27.8	1		
	Yes	20/110	18.2	0.58	0.35-0.94	0.02
Diabetes (m=6)	No	869/3236	26.9	1		
	Yes	80/211	37.9	1.66	1.25-2.22	0.0007
Smoking status (m=16)	Ever	207/822	25.2	1		
	Never	741/2615	28.4	1.17	0.98-1.41	0.078
Latitude of country of birth (m=37)	<20°	265/779	34.0	1		
	20-40°	455/1378	33.0	0.96	0.79-1.15	
	>40°	221/1259	17.6	0.41	0.34-0.51	<0.001
TB incidence per 100 000 of country of birth (m=46)	≤10	26/115	22.6	1		
	11-40	200/1252	16.0	0.65	0.41-1.03	
	41-100	36/129	28.0	1.33	0.74-2.37	
	101-150	15/69	22.1	0.97	0.47-1.99	
	151-300	584/1621	36.0	1.93	1.23-3.02	
	≥300	77/222	34.7	1.82	1.08-3.05	<0.001
Setting of TB exposure (m=274)	Household	671/2149	31.2	1		
	Non-household	204/1030	19.8	0.54	0.46-0.65	<0.001

\*Derived from likelihood ratio test

OR = odds ratio; LTBI = latent tuberculosis infection; n = number; CI = confidence interval;

m = missing; UK = United Kingdom; HIV = human immunodeficiency virus

**Table 2:** Adjusted ORs for the association of LTBI and BCG vaccination from a multivariable logistic regression model (n=3399).

<b>Variable</b>	<b>Group</b>	<b>Adjusted OR*</b>	<b>95% CI</b>	<b>p value**</b>
BCG vaccination	No	1		
	Yes	0.70	0.56-0.87	0.0017
Sex	Male	1		
	Female	0.70	0.60-0.82	<0.001
Age group (years)	16-25	1		
	26-35	1.31	1.07-1.61	
	36-45	1.55	1.23-1.97	
	>45	1.38	1.10-1.74	0.0016
Latitude of country of birth	<20°	1		
	20-40°	0.92	0.76-1.12	
	>40°	0.42	0.34-0.52	<0.001

\*Adjusted for all other variables in the table

\*\*Likelihood ratio test p value

OR = odds ratio; LTBI = latent tuberculosis infection; BCG = Bacillus Calmette-Guérin; n = number; CI = confidence interval.

**Table 3:** Crude and adjusted ORs for the association of LTBI and age at BCG vaccination, by place of birth.

<b>Birthplace</b>	<b>Age at BCG</b>	<b>Proportion with LTBI (%)</b>	<b>Crude OR (95% CI)</b>	<b>p value*</b>	<b>Adjusted OR** (95% CI)</b>	<b>p value*</b>
UK (n=792)	No BCG	32/151 (21.2)	1		1	
	≤2 years	14/144 (9.7)	0.40 (0.20-0.79)		0.39 (0.19-0.77)	
	>2 years	72/497 (14.5)	0.63 (0.40-1.00)	0.021	0.62 (0.38-1.00)	0.018
Non-UK (n=1854)	No BCG	110/300 (36.7)	1		1	
	≤2 years	434/1268 (34.2)	0.90 (0.69-1.17)		0.83 (0.64-1.09)	
	>2 years	83/286 (29.0)	0.71 (0.50-1.00)	0.123	0.63 (0.44-0.90)	0.040

\*Likelihood ratio test p value

\*\*Adjusted for age group, sex and latitude of country of birth.

OR = odds ratio; LTBI = latent tuberculosis infection; UK = United Kingdom; BCG = Bacillus Calmette-Guérin; n = number; CI = confidence interval.

**Table 4:** Crude and adjusted ORs for the association of LTBI and years since BCG vaccination, by age of vaccination.

Age at BCG	Years since BCG	Proportion with LTBI (%)	Crude OR (95% CI)	p value*	Adjusted OR** (95% CI)	p value*
≤2 years (n=1867)	No BCG	142/453 (31.4)	1		1	
	≤10	-	-		-	
	11-20	27/160 (16.9)	0.44 (0.28-0.70)		0.55 (0.33-0.90)	
	>20	423/1254 (33.7)	1.11 (0.89-1.40)	<0.001	0.78 (0.60-1.00)	0.026
>2 years (n=1236)	No BCG	142/453 (31.4)	1		1	
	≤10	12/106 (11.3)	0.28 (0.15-0.53)		0.31 (0.15-0.63)	
	11-20	51/243 (21.1)	0.58 (0.40-0.84)		0.73 (0.48-1.10)	
	>20	92/434 (21.2)	0.59 (0.43-0.80)	<0.001	0.67 (0.46-0.98)	0.0014

\* Likelihood ratio test p value

\*\*Adjusted for age group, sex, ethnicity and country of birth (UK or non-UK)

OR = odds ratio; LTBI = latent tuberculosis infection; BCG = Bacillus Calmette-Guérin; n = number; CI = confidence interval; UK = United Kingdom.



## REFERENCES

1. World Health Organization. Global Health Estimates 2015: Deaths by Cause, Age, Sex, by Country and by Region, 2000-2015. Geneva: World Health Organization; 2016.
2. Casey RM. Global Routine Vaccination Coverage, 2015. *MMWR Morb Mortal Wkly Rep* [Internet]. **2016**; 65. Available from: <http://www.cdc.gov/mmwr/volumes/65/wr/mm6545a5.htm>
3. Trunz BB, Fine P, Dye C. Effect of BCG vaccination on childhood tuberculous meningitis and miliary tuberculosis worldwide: a meta-analysis and assessment of cost-effectiveness. *Lancet Lond Engl*. **2006**; 367(9517):1173–1180.
4. Narayanan PR. Influence of sex, age & nontuberculous infection at intake on the efficacy of BCG: re-analysis of 15-year data from a double-blind randomized control trial in South India. *Indian J Med Res*. **2006**; 123(2):119–124.
5. Abubakar I, Pimpin L, Ariti C, et al. Systematic review and meta-analysis of the current evidence on the duration of protection by bacillus Calmette-Guérin vaccination against tuberculosis. *Health Technol Assess Winch Engl*. **2013**; 17(37):1–372, v–vi.
6. Fine PE. Variation in protection by BCG: implications of and for heterologous immunity. *Lancet Lond Engl*. **1995**; 346(8986):1339–1345.
7. Reyn CF von. Correcting the record on BCG before we license new vaccines against tuberculosis. *J R Soc Med*. **2017**; 110(11):428–433.
8. Mangtani P, Rodrigues LC, Sterne JA, Abubakar I, Fine PEM. Reply to Kernodle and von Reyn. *Clin Infect Dis Off Publ Infect Dis Soc Am*. **2014**; 59(4):608–609.

9. Mangtani P, Abubakar I, Ariti C, et al. Protection by BCG vaccine against tuberculosis: a systematic review of randomized controlled trials. *Clin Infect Dis Off Publ Infect Dis Soc Am.* **2014**; 58(4):470–480.
10. Palmer CE, Long MW. Effects of infection with atypical mycobacteria on BCG vaccination and tuberculosis. *Am Rev Respir Dis.* **1966**; 94(4):553–568.
11. Andersen P, Doherty TM. The success and failure of BCG - implications for a novel tuberculosis vaccine. *Nat Rev Microbiol.* **2005**; 3(8):656–662.
12. Fine PE. BCG vaccination against tuberculosis and leprosy. *Br Med Bull.* **1988**; 44(3):691–703.
13. Wang L, Turner MO, Elwood RK, Schulzer M, FitzGerald JM. A meta-analysis of the effect of Bacille Calmette Guérin vaccination on tuberculin skin test measurements. *Thorax.* **2002**; 57(9):804–809.
14. Pai M, Zwerling A, Menzies D. Systematic review: T-cell-based assays for the diagnosis of latent tuberculosis infection: an update. *Ann Intern Med.* **2008**; 149(3):177–184.
15. Lalvani A, Pareek M. A 100 year update on diagnosis of tuberculosis infection. *Br Med Bull.* **2010**; 93:69–84.
16. Soysal A, Millington KA, Bakir M, et al. Effect of BCG vaccination on risk of *Mycobacterium tuberculosis* infection in children with household tuberculosis contact: a prospective community-based study. *Lancet Lond Engl.* **2005**; 366(9495):1443–1451.
17. Roy A, Eisenhut M, Harris RJ, et al. Effect of BCG vaccination against *Mycobacterium tuberculosis* infection in children: systematic review and meta-analysis. *BMJ.* **2014**; 349:g4643.

18. Mangtani P, Nguipdop-Djomo P, Keogh RH, et al. The duration of protection of school-aged BCG vaccination in England: a population-based case-control study. *Int J Epidemiol.* **2018**; 47(1):193–201.
19. Nguipdop-Djomo P, Heldal E, Rodrigues LC, Abubakar I, Mangtani P. Duration of BCG protection against tuberculosis and change in effectiveness with time since vaccination in Norway: a retrospective population-based cohort study. *Lancet Infect Dis.* **2016**; 16(2):219–226.
20. Chen C, Zhu T, Wang Z, et al. High Latent TB Infection Rate and Associated Risk Factors in the Eastern China of Low TB Incidence. *PloS One.* **2015**; 10(10):e0141511.
21. Chan P-C, Yang C-H, Chang L-Y, et al. Lower prevalence of tuberculosis infection in BCG vaccinees: a cross-sectional study in adult prison inmates. *Thorax.* **2013**; 68(3):263–268.
22. Pai M, Gokhale K, Joshi R, et al. Mycobacterium tuberculosis Infection in Health Care Workers in Rural India: Comparison of a Whole-Blood Interferon  $\gamma$  Assay With Tuberculin Skin Testing. *JAMA.* **2005**; 293(22):2746–2755.
23. Hill PC, Brookes RH, Fox A, et al. Longitudinal assessment of an ELISPOT test for Mycobacterium tuberculosis infection. *PLoS Med.* **2007**; 4(6):e192.
24. Lee SJ, Lee SH, Kim YE, et al. Risk factors for latent tuberculosis infection in close contacts of active tuberculosis patients in South Korea: a prospective cohort study. *BMC Infect Dis.* **2014**; 14:566.

25. Eriksen J, Chow JY, Mellis V, et al. Protective effect of BCG vaccination in a nursery outbreak in 2009: time to reconsider the vaccination threshold? *Thorax*. **2010**; 65(12):1067–1071.
26. Lalvani A, Sridhar S. BCG vaccination: 90 years on and still so much to learn.... *Thorax*. **2010**; 65(12):1036–1038.
27. Abubakar I, Drobniewski F, Southern J, et al. Prognostic Value of Interferon Gamma Release Assays and Tuberculin Skin Test in Predicting the Development of Active Tuberculosis: The UK PREDICT TB Cohort Study. *Lancet Infect Dis*. **2018**; .
28. Public Health England. Tuberculosis in England 2017 report. London: Public Health England; 2018.
29. WHO. WHO TB burden estimates. 2019. [Accessed 8 May 2019]. Available from: <https://www.who.int/tb/country/data/download/en/>
30. countries.csv. Dataset Publishing Language [Internet]. Google Dev. Public Data Explorer. 2012 [Accessed 16 May 2019]. Available from: [https://developers.google.com/public-data/docs/canonical/countries\\_csv](https://developers.google.com/public-data/docs/canonical/countries_csv)
31. Pearce N. Effect measures in prevalence studies. *Environ Health Perspect*. **2004**; 112(10):1047–1050.
32. Rodrigues L, Kirkwood BR. Case-control designs in the study of common diseases: updates on the demise of the rare disease assumption and the choice of sampling scheme for controls. *Int J Epidemiol*. **1990**; 19(1):205–213.

33. Reichenheim ME, Coutinho ES. Measures and models for causal inference in cross-sectional studies: arguments for the appropriateness of the prevalence odds ratio and related logistic regression. *BMC Med Res Methodol.* **2010**; 10:66.
34. Barros AJ, Hirakata VN. Alternatives for logistic regression in cross-sectional studies: an empirical comparison of models that directly estimate the prevalence ratio. *BMC Med Res Methodol.* **2003**; 3:21.
35. Guwatudde D, Nakakeeto M, Jones-Lopez EC, et al. Tuberculosis in Household Contacts of Infectious Cases in Kampala, Uganda. *Am J Epidemiol.* **2003**; 158(9):887–898.
36. NICE. National Institute for Health and Care Excellence (NICE) guideline—NG33 Tuberculosis [Internet]. United Kingdom: NICE; 2016. Available from: <https://www.nice.org.uk/guidance/NG33>
37. Young TK, Mirdad S. Determinants of tuberculin sensitivity in a child population covered by mass BCG vaccination. *Tuber Lung Dis Off J Int Union Tuberc Lung Dis.* **1992**; 73(2):94–100.
38. Rani SH, Vijayalakshmi V, Sunil K, Lakshmi KA, Suman LG, Murthy KJ. Cell mediated immunity in children with scar-failure following BCG vaccination. *Indian Pediatr.* **1998**; 35(2):123–127.
39. Bartlett JW, Harel O, Carpenter JR. Asymptotically Unbiased Estimation of Exposure Odds Ratios in Complete Records Logistic Regression. *Am J Epidemiol.* **2015**; 182(8):730–736.
40. Shapiro C, Cook N, Evans D, et al. A case-control study of BCG and childhood tuberculosis in Cali, Colombia. *Int J Epidemiol.* **1985**; 14(3):441–446.

41. Fine P. Stopping routine vaccination for tuberculosis in schools. *BMJ*. **2005**; 331(7518):647–648.
42. Fine PE, Ponnighaus JM, Maine N. The distribution and implications of BCG scars in northern Malawi. *Bull World Health Organ*. **1989**; 67(1):35–42.
43. World Health Organization. BCG vaccines: WHO position paper – February 2018. *Wkly Epidemiol Rec*. **2018**; 93(8):73–96.
44. Hawn TR, Day TA, Scriba TJ, et al. Tuberculosis Vaccines and Prevention of Infection. *Microbiol Mol Biol Rev MMBR*. **2014**; 78(4):650–671.
45. McShane H, Jacobs WR, Fine PE, et al. BCG: Myths, realities, and the need for alternative vaccine strategies. *Tuberculosis*. **2012**; 92(3):283–288.
46. Meeren OVD, Hatherill M, Nduba V, et al. Phase 2b Controlled Trial of M72/AS01E Vaccine to Prevent Tuberculosis. *N Engl J Med* [Internet]. **2018** [cited 2018 Dec 1]; . Available from: [https://www.nejm.org/doi/10.1056/NEJMoa1803484?url\\_ver=Z39.88-2003&rfr\\_id=ori%3Arid%3Acrossref.org&rfr\\_dat=cr\\_pub%3Dwww.ncbi.nlm.nih.gov](https://www.nejm.org/doi/10.1056/NEJMoa1803484?url_ver=Z39.88-2003&rfr_id=ori%3Arid%3Acrossref.org&rfr_dat=cr_pub%3Dwww.ncbi.nlm.nih.gov)
47. Ellis RD, Hatherill M, Tait D, et al. Innovative clinical trial designs to rationalize TB vaccine development. *Tuberc Edinb Scotl*. **2015**; 95(3):352–357.
48. Nemes E, Geldenhuys H, Rozot V, et al. Prevention of *M. tuberculosis* Infection with H4:IC31 Vaccine or BCG Revaccination. *N Engl J Med*. **2018**; 379(2):138–149.

## **FINANCIAL SUPPORT**

The data for this study is from the PREDICT study, which was supported by the National Institute for Health Research [grant numbers NIHR HTA 08/68/01, NIHR SRF-2011-04-001]. FD was supported by the NIHR Imperial Biomedical Research Center. The funders had no influence on the contents of this paper.

## **POTENTIAL CONFLICTS OF INTEREST**

All authors have completed the ICMJE Form for Disclosure of Potential Conflicts of Interest. CJ, JS and AJ report grants from the UK National Institute for Health Research during the conduct of the study. In addition, AJ reports grants from Wellcome Trust during the conduct of the study, and is a named inventor on patents pertaining to T cell-based diagnosis, including IGRA technologies. Some of these patents were assigned by the University of Oxford to Oxford Immunotec PLC resulting in royalty entitlements for the University of Oxford and AL. We declare no other relationships or activities that could appear to have influenced the submitted work.

## **AUTHOR CONTRIBUTIONS**

Drs Katelaris, Gupta, and Abubakar have full access to all data in the study and take responsibility for the integrity of the data and accuracy of the data analysis.

Study concept and design: Mangtani, Jackson, Katelaris, Abubakar

Acquisition, analysis, and interpretation of data: Katelaris, Jackson, Mangtani, Abubakar

Statistical analysis: Katelaris, Jackson, Gupta

Drafting of the manuscript: Katelaris, Jackson, Mangtani

Critical revision of the manuscript for important intellectual content: All authors

Obtained funding: Abubakar, Lalvani, Drobniowski, Lipman

Administrative, technical, or material support: Jackson, Gupta, Southern

Supervision: Jackson, Mangtani, Abubakar

## **ETHICS APPROVAL**

This study was approved by the LSHTM MSc Research Ethics Committee (reference #13431). Ethics approval for the PREDICT study was granted through the National Research Ethics Service (Brent Medical Ethics Committee, approval number 10/H0717/14) and is registered on clinicaltrials.gov (NCT01162265).

## **CORRESPONDING AUTHOR**

Dr Charlotte Jackson, Institute for Global Health, University College London, London; email: [c.r.jackson@ucl.ac.uk](mailto:c.r.jackson@ucl.ac.uk); phone: +44 (0)20 7670 4806.

## **CURRENT AFFILIATIONS**

Anthea L Katelaris, National Centre for Epidemiology and Population Health, Research School of Population Health, Australian National University, ACT 0200, Australia



## **SUPPLEMENTARY DATA**

### **Effectiveness of BCG vaccination against *Mycobacterium tuberculosis* infection in adults: a cross-sectional analysis of a UK-based cohort**

#### **AUTHORS**

Anthea Laura Katelaris, Charlotte Jackson, Jo Southern, Rishi K Gupta, Francis Drobniewski, Ajit Lalvani, Marc Lipman, Punam Mangtani\*, Ibrahim Abubakar\*

\*Joint last authors

#### **Appendix**

##### **Appendix 1: Sensitivity analyses**

*Table A1a:* Sensitivity analyses. Adjusted effect estimates for the association between LTBI and BCG vaccination.

*Table A1b:* Analysis of the association of BCG status with tuberculin skin test positivity

##### **Appendix 2: Missing data analysis**

*Table A2a:* Missingness analysis

*Table A2b:* Missingness analysis. Multivariable analysis of the association of covariates with missing BCG status (n=3657)

##### **Appendix 3: Bivariable analysis of age at BCG, by country of birth and ethnicity**

*Table A3:* The distribution of country of birth and ethnicity by age at BCG (column percentages)

## **Appendix 1: Sensitivity analyses**

Sensitivity analyses produced results largely similar to the main analysis (*Table A1a*).

Adjustment for smoking status and any social risk factor (as proxy measures of socioeconomic status) gave an OR of 0.67 (0.54-0.84). With adjustment for TB incidence of country of birth, the OR was 0.69 (0.55-0.87). When the analysis was restricted to participants with concordant IGRAs, the OR was 0.68 (95% CI 0.51-0.90). Using log binomial regression yielded a prevalence ratio (PR) of 0.80 (95% CI 0.69-0.92). When participants with prior TB contact were included, the OR increased slightly to 0.73 (95% CI 0.59-0.90) and restricting the analysis to household contacts produced a slightly lower OR of 0.65 (95% CI 0.49-0.87).

**Table A1a:** Sensitivity analyses. Adjusted effect estimates for the association between LTBI and BCG vaccination.

Sensitivity analysis	Number	Adjusted effect estimate*	95% CI	p value**
Original logistic regression model	3399	OR=0.70	0.56-0.87	0.0017
Additional adjustment for smoking and social risk factors	3384	OR=0.67	0.54-0.84	0.0007
Additional adjustment for TB incidence of country of birth	3390	OR=0.69	0.55-0.87	0.0017
Restricted to participants with concordant IGRAs	2760	OR=0.68	0.51-0.90	0.0088
Log binomial regression model	3399	PR=0.80	0.69-0.92	0.0031
Participants with prior contact with TB included	3927	OR=0.73	0.59-0.90	0.0033
Restricted to household contacts	2118	OR=0.65	0.49-0.87	0.0038

\*Adjusted for age group, sex and latitude of country of birth (unless otherwise stated)

\*\*Likelihood ratio test p value

CI = confidence interval; OR = odds ratio; PR = prevalence ratio; IGRAs = interferon-gamma release assay; TB = tuberculosis.

**Table A1b:** Analysis of the association of BCG status with tuberculin skin test positivity (n with TST results = 3119)

TST positive criteria	Vaccination	Proportion positive (%)	Crude OR (95% CI)	p value*	Adjusted OR** (95% CI)	p value *
All ≥5mm	No BCG	160/421 (38.0)	1		1	
	BCG	1371/2698 (50.8)	1.69 (1.37-2.08)	<0.001	1.59 (1.28-1.98)	<0.001
No BCG ≥5mm	No BCG	160/421 (38.0)	1		1	
	BCG	695/2698 (25.8)	0.57 (0.46-0.70)	<0.001	0.50 (0.40-0.63)	<0.001

\*Likelihood ratio test p value

\*\*Adjusted for age group, sex and latitude of country of birth; n = 3072 in the adjusted analyses.

OR = odds ratio; TST = tuberculin skin test; BCG = Bacillus Calmette-Guérin; CI = confidence interval; n= number.

## **Appendix 2: Missing data analysis**

We analysed missing data by outcome and primary exposure, which showed most variables had very low percentages of missing data (*Table A2a*). Data on the outcome (LTBI) was missing in 363 participants (8.4%). There was no association between missing outcome data and BCG vaccination (OR 1.05, 95% CI 0.74-1.49,  $p=0.79$ ). Data on BCG vaccination was missing in 555 participants (12.9%). Those with data missing on BCG status were more likely to have LTBI (OR 1.33, 95% CI 1.09-1.63).

Multivariable analysis of covariates associated with missing BCG status showed that age group, country of birth, latitude of country of birth, ethnicity and immunosuppression were independently associated with missing BCG status (*Table A2b*).

Although data were missing on the age of vaccination and years since vaccination for a quarter of subjects, there was no evidence that this was associated with the outcome (LTBI).

### *Missing data in the primary analysis*

There were little missing data on covariates, so the complete-case approach only resulted in a further 1.6% of participants being excluded from the primary analysis due to missing covariate data.

**Table A2a:** Missingness analysis. Odds ratios of LTBI positivity and BCG vaccination, in participants missing data on baseline variables versus those with data not missing (n=4310)

Variable	Number missing (%) <sup>*</sup>	OR of LTBI (n=3497)	95% CI	p value <sup>**</sup>	OR of BCG vaccination (n=3755)	95% CI	p value <sup>**</sup>
<i>LTBI</i>	363 (8.42)	-	-	-	1.05	0.74-1.49	0.79
<i>BCG vaccination</i>	555 (12.88)	1.33	1.09-1.63	0.005	-	-	-
<i>Sex</i>	23 (0.53)	0.90	0.32-2.52	0.85	0.80	0.23-2.76	0.73
<i>Age group, years</i>	7 (0.16)	1.27	0.11-13.99	0.85	0.15	0.009-2.42	0.12
<i>Country of birth</i>	21 (0.49)	2.91	1.05-8.05	0.03	0.60	0.13-2.84	0.52
<i>Ethnicity</i>	109 (2.53)	1.46	0.97-2.22	0.07	1.36	0.65-2.85	0.41
<i>Any social risk factor</i>	0 (0)	-	-	-	-	-	-
<i>HIV</i>	152 (3.53)	1.00	0.68-1.46	0.99	1.22	0.68-2.19	0.50
<i>Other</i>							
<i>immunosuppression</i>	16 (0.37)	3.18	0.85-11.86	0.07	0.38	0.07-1.94	0.22
<i>Diabetes</i>	16 (0.37)	1.44	0.42-4.96	0.55	0.75	0.09-6.45	0.79
<i>Smoking status</i>	30 (0.70)	0.70	0.26-1.90	0.48	0.30	0.11-0.80	0.01
<i>Latitude of country of birth</i>	58 (1.35)	0.94	0.50-1.78	0.85	0.46	0.22-0.95	0.03
<i>Age at vaccination (if vaccinated, n=3264)</i>	888 (27.21)	0.90	0.75-1.07	0.24	-	-	-
<i>Years since BCG (if vaccinated, n=3264)</i>	860 (26.35)	0.91	0.76-1.10	0.34	-	-	-
<i>Setting of TB exposure</i>	340 (7.89)	0.98	0.76-1.26	0.87	0.536	0.40-0.73	<0.001

\*Number missing out of total (n=4310)

\*\*Derived from chi-squared test

LTBI = latent tuberculosis infection; BCG = Bacillus Calmette-Guérin; n = number; CI = confidence interval; OR = odds ratio; HIV = human immunodeficiency virus.

**Table A2b:** Missingness analysis. Multivariable analysis of the association of covariates with missing BCG status (n=3657)

Variable	Group	Adjusted OR*	95% CI	p value**
<i>LTBI</i>	Negative	1		
	Positive	1.37	1.10-1.70	0.005
<i>Sex</i>	Male	1		
	Female	0.98	0.79-1.20	0.82
<i>Age group, years</i>	16-25	1		
	26-35	0.90	0.69-1.16	0.41
	36-45	0.62	0.44-0.87	0.006
	>45	1.15	0.85-1.56	0.36
<i>Country of birth</i>	UK	1		
	Non-UK	1.65	1.05-2.54	0.028
<i>Ethnicity</i>	Indian	1		
	White	0.79	0.54-1.16	0.23
	Black African	0.93	0.61-1.40	0.71
	Mixed	1.00	0.71-1.39	0.99
	Pakistani	1.44	0.48-1.00	0.048
	Bangladeshi	1.16	0.70-1.94	0.56
	Black Caribbean	1.09	0.64-1.85	0.74
	Black Other / Chinese / Other	0.66	0.35-1.27	0.22
<i>Any social risk factor</i>	No	1		
	Yes	0.97	0.62-1.52	0.90
<i>HIV</i>	No	1		
	Yes	0.86	0.25-2.91	0.81
<i>Other immunosuppression</i>	No	1		
	Yes	2.12	1.32-3.38	0.002
<i>Diabetes</i>	No	1		
	Yes	1.08	0.72-1.62	0.72
<i>Smoking status</i>	Ever	1		
	Never	1.09	0.84-1.42	0.50
<i>Latitude of country of birth</i>	<20°	1		
	20-40°	1.20	0.84-1.72	0.32
	>40°	1.67	0.98-2.86	0.06

\*Adjusted for all other variables in the table

\*\*Wald test p value

BCG = Bacillus Calmette-Guérin; n = number; OR = odds ratio; CI = confidence interval; LTBI = latent tuberculosis infection; UK = United Kingdom; HIV = human immunodeficiency virus.

### Appendix 3: Analysis of age at BCG by ethnicity and place of birth

**Table A3:** The distribution of age at BCG by ethnicity, stratified by place of birth (UK born or non-UK born)

Country of birth	Ethnicity	Age at BCG in years, n (%)			
		≤2		>2	
		n	(%)	n	(%)
<b>UK born</b>					
	Indian	41	(30)	95	(70)
	White	41	(16)	216	(84)
	Black African	11	(26)	31	(74)
	Mixed	13	(20)	51	(80)
	Pakistani	14	(36)	25	(64)
	Bangladeshi	10	(40)	15	(60)
	Black Caribbean	5	(10)	43	(90)
	Black other / Chinese / other	8	(35)	15	(65)
	Missing	1	(14)	6	(86)
	<b>Total</b>	144	(22)	497	(78)
<b>Non-UK born</b>					
	Indian	415	(84)	82	(17)
	White	136	(72)	53	(28)
	Black African	263	(86)	42	(14)
	Mixed	219	(84)	42	(16)
	Pakistani	92	(88)	13	(12)
	Bangladeshi	52	(84)	10	(16)
	Black Caribbean	32	(58)	23	(42)
	Black other / Chinese / other	45	(82)	10	(18)
	Missing	14	(56)	11	(44)
	<b>Total</b>	1,268	(82)	286	(18)

BCG = Bacillus Calmette-Guérin; n = number; UK = United Kingdom