

1 **Challenges and research priorities to understand interactions between climate, ice**
2 **sheets and global mean sea level during past interglacials**

3 Emilie Capron¹, Alessio Rovere², Jacqueline Austermann³, Yarrow Axford⁴, Natasha L. M.
4 Barlow⁵, Anders E. Carlson⁶, Anne de Vernal⁷, Andrea Dutton⁸, Robert E. Kopp⁹, Jerry F.
5 McManus³, Laurie Menviel¹⁰, Bette L. Otto-Bliesner¹¹, Alexander Robinson^{12,13,14}, Jeremy
6 D. Shakun¹⁵, Polychronis C. Tzedakis¹⁶, Eric W. Wolff¹⁷

7 ¹Physics of Ice, Climate and Earth, Niels Bohr Institute, University of Copenhagen, Tagensvej 16, DK-2200
8 Copenhagen, Denmark

9 ²MARUM – Center for Marine Environmental Sciences, University of Bremen, Leobener Strasse 8, 28359,
10 Bremen, Germany

11 ³Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY 10964-8000, USA

12 ⁴Department of Earth and Planetary Sciences, Northwestern University, Evanston, IL 60208, USA

13 ⁵School of Earth and Environment, University of Leeds, Leeds LS2 9JT, UK

14 ⁶College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, Corvallis, OR, 97331, USA

15 ⁷GEOTOP-UQAM, CP 8888 Montréal, H3C 3P8, Canada

16 ⁸Department of Geological Sciences, University of Florida, Gainesville, FL, 32611, USA; Department of
17 Geoscience, University of Wisconsin, Madison, WI, 53706, USA

18 ⁹Department of Earth and Planetary Sciences and Institute of Earth, Ocean and Atmospheric Sciences,
19 Rutgers University, New Brunswick, NJ, USA

20 ¹⁰Climate Change Research Centre, PANGEA, the University of New South Wales, Sydney, NSW, Australia

21 ¹¹Climate and Global Dynamics Laboratory, National Center for Atmospheric Research (NCAR) Boulder, CO
22 80305, USA

23 ¹²Potsdam Institute for Climate Impact Research, Potsdam, Germany

24 ¹³Complutense University of Madrid, 28040 Madrid, Spain

25 ¹⁴Geosciences Institute, UCM-CSIC, 28040 Madrid, Spain

26 ¹⁵Department of Earth and Environmental Sciences, Boston College, Chestnut Hill, MA 02467, USA

27 ¹⁶Environmental Change Research Centre, Department of Geography, University College London, England,
28 UK

29 ¹⁷Department of Earth Sciences, University of Cambridge, CB2 3EQ, Cambridge, UK

1 **Abstract:** Quaternary interglacials provide key observations of the Earth system's
2 responses to orbital and greenhouse gas forcing. They also inform on the capabilities of
3 Earth system models, used for projecting the polar ice-sheet and sea-level responses to
4 a regional warmth comparable to that expected by 2100 C.E. However, a number of
5 uncertainties remain regarding the processes and feedbacks linking climate, ice-sheet and
6 sea-level changes during past warm intervals. Here, we delineate the major research
7 questions that need to be resolved and future research directions that should be taken by
8 the paleoclimate, sea-level and ice-sheet research communities in order to increase
9 confidence in the use of past interglacial climate, ice-sheet and sea-level reconstructions
10 to constrain future predictions. These questions were formulated during a joint workshop
11 held by the PAGES-INQUA PALSEA (PALEo constraints on SEA level rise) and the
12 PAGES-PMIP QUIGS (QUaternary InterGlacialS) Working Groups in September 2018.

13 **Key-words:** Interglacials; Paleoclimatology; Polar ice sheets; Sea-level changes; Natural
14 archives; Earth System Modeling.

15 **Introduction**

16 Human-induced global warming has large environmental and societal implications,
17 including the potential for substantial and/or rapid polar ice-mass loss that leads to global
18 sea-level rise. To evaluate the risk of major current and future environmental changes, it
19 is essential to understand climate and cryosphere processes and feedbacks occurring
20 during past periods that were warmer than the pre-industrial. About 50 warm intervals,
21 referred to as interglacials, punctuated the Quaternary (0-2.6 Ma). Each lasted between
22 ~10 to ~30 thousand of years (ka) with the distribution of Northern Hemisphere ice
23 resembling the present, with minimal ice sheets outside of Greenland (PAGES working
24 group on Past Interglacials, 2016). Over the past ~450 ka, ice core records show that
25 interglacial Antarctic surface conditions were similar to, or even warmer than those of the
26 pre-industrial (e.g. Jouzel et al., 2007; Figure 1). Past interglacials are not perfect
27 analogues for future anthropogenic changes, as their warming was caused by different
28 forcing mechanisms. Interglacial polar warming resulted from changes in astronomical
29 forcing and an altered global heat distribution, while future warming is driven by
30 anthropogenic greenhouse gas forcing. Nonetheless, interglacials provide an opportunity
31 to utilize proxy data and Earth system models (ESM) to assess the effects of warmer-than-
32 pre-industrial polar temperatures on critical parts of the Earth system, such as polar ice
33 sheets and sea level.

1 In this context, the Last Interglacial (LIG, ~129-116 ka) has received particular
2 attention. Being the most recent interglacial prior to the Holocene, the LIG offers a great
3 amount of data, and as the warmest interglacial of the last 800 ka at many locations
4 (PAGES working group on Past Interglacials, 2016), it is especially informative despite
5 some major uncertainties that still prevail. Recent studies estimated that LIG high-latitude
6 surface ocean temperatures were warmer by at least 1°C and polar surface air
7 temperatures by >3-11°C relative to pre-industrial temperatures, and that peak warmth did
8 not occur synchronously across the globe (Capron et al., 2017 and references therein).
9 Nevertheless, there are too few continental surface temperature reconstructions and no
10 comprehensive global surface temperature estimate combining ice, marine, and terrestrial
11 paleo-temperature reconstructions.

12 While commonly reported figures suggest that the LIG was characterized by global
13 mean sea level (GMSL) 6-9 m higher than today (Dutton et al., 2015; Figure 1), there is
14 still considerable uncertainty attached to the amplitude and exact timing of this GMSL
15 peak. LIG sea-level reconstructions from coastal archives are hampered by uncertainties
16 in Glacial Isostatic Adjustment (GIA) related to Earth's internal viscoelastic structure and
17 the size of ice sheets during the preceding glaciation, which both remain poorly
18 constrained. For instance, while benthic $\delta^{18}\text{O}$ records suggest similar ice volumes
19 between the last two glacial maxima though the spatial distribution of ice might have been
20 different, a recent study proposes that the penultimate glacial maximum ice volume was
21 $21\pm 14\text{m}$ global sea-level equivalent smaller than during the most recent one (Rohling et
22 al., 2017). Additionally, mantle dynamic topography may bias LIG sea level
23 reconstructions, but it remains difficult to reliably quantify this effect (Austermann et al.
24 2017).

25 There are also large uncertainties regarding the timing and extent of mass loss from
26 the Greenland and Antarctic ice sheets and their respective contributions to the higher-
27 than-present GMSL throughout the LIG. In particular, the sea-level community still debates
28 the extent of variability in sea level within the LIG, with geomorphological studies inferring
29 global mean sea level to be relatively stable (Barlow et al., 2018) versus models with large
30 high-to-low swings (Kopp et al., 2009); and regional records do not agree on the number
31 or the timing of sea level fluctuations (Vyverberg et al., 2018 and references therein).

32 It would be valuable to assemble similar datasets for older interglacials to assess
33 the relationship between polar climate and ice sheets/sea level, and thus evaluate how

1 consistent this relationship has been. However, for these intervals, both climate and sea-
2 level data are more limited.

3 In September 2018, the PAGES-INQUA PALSEA (PALeo constraints on SEA level
4 rise) and the PAGES-PMIP QUIGS (QUaternary InterGlacialS) Working Groups held a
5 joint workshop gathering key expertise from the sea-level, ice-sheet and climate
6 paleocommunities, in order to formulate a series of priority research questions to motivate
7 further research and cross-cutting initiatives. Here, we report those key questions that may
8 provide future research foci for communities working on paleo climate, sea level and ice
9 sheets.

10 **Priority research questions**

11 To use past interglacial climate, ice-sheet and sea-level records to support our
12 predictions of future sea-level changes with greater confidence, eight research areas
13 require further investigation:

14 **1. Sea level and temperature.** How did interglacial sea-level highstands relate to regional
15 (e.g., high latitudes, tropics) and global temperatures? This can be subdivided into the
16 following questions:

- 17 • How did global and regional climates evolve during past interglacials?
- 18 • When and how high were sea level highstands during past interglacials?
- 19 • How does the timing of peak temperature and sea level compare within an
20 interglacial?

21 **2. Variability in sea level and climate.** Were there centennial- to millennial-scale
22 oscillations in GMSL or in climate during interglacials? This question is crucial for
23 estimating the maximum rates of sea-level changes when the configuration of the ice
24 sheets most closely resembled that of today (e.g., during the LIG and Marine Isotope
25 Stage 11; Figure 1).

26 **3. Contribution of ice sheets.** What determined the size of polar ice sheets in different
27 interglacial climates? Furthermore, what were the relative contributions and when were
28 peak contributions of the different ice sheets to GMSL rise during interglacials?

29 **4. Ice-sheet resilience.** Can large polar ice sheets persist under a large multi-millennial-
30 scale warming and how long must warmth persist to lead to their disappearance? For
31 example, a decade ago the Greenland ice sheet was considered to be largely deglaciated
32 during the LIG. However, more recent data and reconstructions suggest that this ice sheet
33 largely persisted through the LIG despite a regional warming of at least 4°C (e.g., Colville

1 et al., 2011; NEEM community members, 2013). Collectively, results from Greenland point
2 to important unanswered questions for the LIG. Meanwhile, similar research on other ice
3 sheets and during older interglacials is necessary too.

4 **5. Glacial maxima.** What was the size and extent of each ice sheet during past glacial
5 maxima? Was the partitioning of ice between ice sheets consistent between glacial
6 periods? These are crucial boundary conditions needed to correct interglacial sea-level
7 indicators for GIA effects.

8 **6. Interglacial oceanic circulation.** How did oceanic circulation evolve during
9 interglacials, and how did this affect polar climate and the Antarctic and Greenland ice
10 sheets? Also, could significant freshwater contributions from a Southern Hemisphere
11 source have impacted oceanic circulation and regional climate anomalies?

12 **7. Holocene versus older interglacials.** How can we use the Holocene, for which
13 paleodata are most abundant, to learn about older interglacials?

14 **8. Interglacial relevance to the future.** Which ice-sheet processes relevant to future
15 changes can be meaningfully constrained by interglacial reconstructions, and which are
16 fundamentally different from processes in past interglacials?

17 To provide answers to these research questions, several technical research issues
18 should first be addressed. There is still a strong need for highly-resolved climate and sea-
19 level records for the LIG, and even more so for prior interglacials. It is also necessary to
20 improve absolute and relative chronologies for paleoclimatic and sea-level records, along
21 with methodological developments to synchronize records from different types of archives
22 and fully integrate tracers and age uncertainties. Uncertainties remain on the rate of sea-
23 level rise prior to and during the LIG. Further, sea-level modeling requires a better
24 understanding of Earth's internal viscoelastic structure and dynamics. Paleoclimate
25 records provide insight into the spatio-temporal structure and amplitude of interglacial
26 climate change to inform ESM simulations. However, a better assessment of the
27 respective influence of the controlling factors and the seasonality of proxy records is
28 required to improve the interpretation of temperature reconstructions. Plans should be
29 implemented to fill geographical gaps, such as better spatial coverage of paleodata in the
30 Southern and Pacific Oceans and on continents, especially in the high latitudes. There is
31 an urgent need to find direct (near-field) evidence to detect and quantify ice-mass loss in
32 Antarctica, which is crucial to constrain the degree to which it may have contributed to
33 LIG, and other interglacial, GMSL. For Greenland, important inconsistencies exist
34 between ice-sheet model results, climate proxy reconstructions and geological evidence


1 for ice sheet size (Colville et al., 2011; NEEM community members, 2013; Yau et al.,
2 2016). Uncertainties in Greenland leave it unclear how much contribution from Antarctica
3 would be required to explain LIG far-field sea level estimates; and furthermore, maximum
4 contributions from the ice sheets may have been out of phase with each other (Dutton et
5 al., 2015). Finally, further coordinated ESM efforts should simulate the transient evolution
6 of interglacials, and explicitly include climate proxies (e.g., oxygen and carbon isotopes),
7 climate-ice-sheet interactions and dynamic vegetation. These modeling exercises should
8 be extended beyond the LIG when comprehensive datasets become available, to provide
9 improved quantitative understanding of warmth, sea level, ice sheets, and their
10 interactions through multiple interglacials.

11 **Concluding remarks**

12 Investigation of the community-identified research priorities described here will lead to an
13 improved understanding of climate and ice-sheet (and subsequently sea-level) responses
14 to astronomical and greenhouse gas forcing, and thereby the Earth system response to
15 global surface conditions similar to or warmer than the pre-industrial climate. Improved
16 data will be useful benchmarks to inform on model capabilities used for climate, ice-sheet
17 and sea-level projections in capturing millennial-scale patterns and amplitudes of the
18 responses in warm(er) worlds. Addressing these outstanding questions requires
19 organized interdisciplinary collaborations between scientists working on climate, ice
20 sheets, sea level and the solid Earth. Hence, cross-cutting activities such as the 2018
21 PALSEA-QUIGS workshop are necessary and should be repeated to facilitate knowledge
22 exchange between the different disciplines.

23 **Acknowledgments:** This paper is an outcome of the joint workshop organized by the
24 PAGES-INQUA PALSEA (PALeo constraints on SEA level rise) and the PAGES-PMIP
25 QUIGS (QUaternary InterGlacials) Working Groups in Galloway NJ (USA), 27-30
26 September 2018. We are very grateful to all the participants for the stimulating discussions
27 that contributed to this article. We also thank Past Global Changes (PAGES), the
28 International Union for Quaternary Research (INQUA), the United States National Science
29 Foundation (NSF) and Rutgers University, Institute of Earth, Ocean, and Atmospheric
30 Sciences for their financial support. E.C. acknowledges financial support from the
31 ChronoClimate project, funded by the Carlsberg Foundation.

1 **Figure 1. Key paleoclimatic records over the past 450 ka.**
 2 (A) Northern Hemisphere 21 June insolation, grey. (B) Benthic foraminifera $\delta^{18}\text{O}$
 3 composite, black (Lisiecki and Raymo, 2005), numbers designate the Marine Isotope
 4 Stages (MIS) corresponding to interglacial intervals. (C) Red Sea sea-level curve relative
 5 to present, green, and 95% probability interval, light grey (Grant et al. 2014). (D) Global
 6 mean sea level (GMSL) estimates relative to present for MIS 11 and the LIG (referred to
 7 MIS 5e), light grey shading indicates the uncertainty of GMSL maximum (Dutton et al.,
 8 2015). (E) Atmospheric CO_2 concentrations, red (Bereiter et al., 2015) and (F) Antarctic
 9 surface temperature relative to present, blue (Jouzel et al., 2007) from the EPICA Dome
 10 C ice core. The projected Antarctic warming by 2100 based on the Representative
 11 Concentration Pathways (RCP) 8.5 and RCP 2.6 scenarios is indicated with orange
 12 dashed lines.



13

1 References

- 2 Austermann, J., Mitrovica, J.X., Huybers, P., Rovere, A., 2017. Detection of a dynamic
3 topography signal in last interglacial sea-level records. *Science Advances* 3, e1700457,
4 doi: 10.1126/sciadv.1700457.
- 5 Barlow, N. L. M., McClymont, E. L., Whitehouse, P. L., Stokes, C. R., Jamieson, S. S.
6 R., Woodroffe, S. A., Bentley, M. J., Callard, S. L., Ó Cofaigh, C., Evans, D. J. A.,
7 Horrocks, J. R., Lloyd, J. M., Long, A. J., Margold, M., Roberts, D. H., Sanchez-Montes
8 M. L. (2018). Lack of evidence for a substantial sea-level fluctuation within the Last
9 Interglacial. *Nature Geoscience*, 11(9), 627-634.
- 10 Bereiter, B., S. Eggelston, J. Schmitt, C. Nehrbass-Ahles, T. F. Stocker, H. Fischer, S.
11 Kipfstuhl, and J. Chappellaz (2015), Revision of the EPICA Dome C CO₂ record from
12 800 to 600 kyr before present. *Geophys. Res. Lett.* 42, doi:10.1002/2014GL061957.
- 13 Capron, E., Govin, A., Feng, R., Otto-Bliesner, B. and E. W. Wolff. Critical evaluation of
14 climate syntheses to benchmark CMIP6/PMIP4 127 ka Last Interglacial simulations in
15 the high-latitude regions. *Quaternary Sci. Rev.* 168, 137-150.
- 16 Colville, E.J., Carlson, A.E., Beard, B.L., Hatfield, R.G., Stoner, J.S., Reyes, A.V.,
17 Ullman, D.J., 2011. Sr-Nd-Pb isotopic evidence for ice-sheet presence on southern
18 Greenland during the last interglacial. *Science*, 333, 620-623.
- 19 Dutton, A., Carlson, A.E., Long, A.J., Milne, G.A., Clark, P.U., DeConto, R., Horton, B.P.,
20 Rahmstorf, S., Raymo, M.E., 2015. Sea-level rise due to polar ice-sheet mass loss
21 during past warm periods. *Science* 349 (6244), doi: 10.1126/science.aaa4019.
- 22 Grant, K. M., Rohling, E. J., Bronk Ramsey, C., Cheng, H., Edwards, R. L., Florindo, F.,
23 Heslop, D., Marra, F., Roberts, A. P., Tamisiea, M. E., and F. Williams, 2014. Sea-level
24 variability over five glacial cycles. *Nature Communications* 5, doi: 10.1038/ncomms6076
- 25 Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G.,
26 Minster, B., Nouet, J., Barnola, J.-M., Fisher, H., Gallet, J.-C., Johnsen, S.,
27 Leuenberger, M., Loulergue, L., Luethi, D., Oerter, H., Parrenin, F., Raisbeck, G.,
28 Raynaud, D., Schilt, A., Schwander, J., Selmo, J., Souchez, R., Spahni, R., Stauffer, B.,
29 Steffensen, J.P., Stenni, B., Stocker, T.F., Tison, J.-L., Werner, M., Wolff, E.W., 2007.
30 Orbital and millennial Antarctic climate variability over the past 800,000 years. *Science*
31 317, 793-796.
- 32 Kopp, R.E., Simons, F.J., Mitrovica, J.X., Maloof, A.C. and Oppenheimer, M., 2009.
33 Probabilistic assessment of sea level during the last interglacial stage. *Nature* 462: 863-
34 867, doi:10.1038/nature08686.
- 35 Lisiecki, L.E., Raymo, M.E., 2005. A Pliocene-Pleistocene stack of 57 globally distributed
36 benthic $\delta^{18}\text{O}$ records. *Paleoceanography* 20, [http://dx.doi.org/](http://dx.doi.org/10.1029/2004pa001071)
37 10.1029/2004pa001071. PA1003.
- 38 NEEM community members, 2013. Eemian interglacial reconstructed from a Greenland
39 folded ice core. *Nature* 493, 489-493. <http://dx.doi.org/10.1038/>.
- 40 Past Interglacials Working Group of PAGES, 2016. Interglacials of the last 800,000
41 years. *Rev. Geophys.*, <http://dx.doi.org/10.1002/2015RG000482>.
- 42 Rohling, E.J., Hibbert, F. D., Williams, F. H., Grant, K. M., Foster, G. Hennekam, R., de
43 Lange G. J., Roberts, A. P., Yu, J., Webster, M., and Yokoyama, Y., 2017. Differences

- 1 between the last two glacial maxima and implications for ice sheet, $\delta^{18}\text{O}$ and sea level
2 reconstructions. *Quaternary Sci. Rev.* 176, 1-28.
- 3 Vyverberg, K., Dechnik, B., Dutton, A., Webster, J.M., Zwartz, D. and Portell, R.W., 2018.
4 Episodic reef growth in the granitic Seychelles during the Last Interglacial: Implications for
5 polar ice sheet dynamics. *Marine Geology*, 399, 170-187.
- 6 Yau, A. M., Bender, M. L., Robinson, A., and E. J. Brook, 2016. Reconstructing the last
7 interglacial at Summit, Greenland: Insights from GISP2. *PNAS*, 113, 9710-9715.