


# DNAJC6 Mutations Disrupt Dopamine Homeostasis in Juvenile Parkinsonism-Dystonia

Joanne Ng, MD, PhD,<sup>1,2</sup> Elisenda Cortès-Saladelafont, MD,<sup>1</sup> Lucia Abela, MD,<sup>1</sup> Pichet Termsarasab, MD,<sup>3,4</sup>  Kshitij Mankad, FRCR,<sup>5</sup> Snिया Sudhakar, FRCR,<sup>5</sup> Kathleen M. Gorman, MD,<sup>1,6</sup> Simon J.R. Heales, PhD,<sup>7</sup> Simon Pope, PhD,<sup>7</sup> Lorenzo Biassoni, MSc, FRCP, FEBNM,<sup>5</sup> Barbara Csányi, MD,<sup>1</sup> John Cain, FRCR, PhD,<sup>8</sup> Karl Rakshi, MBChB,<sup>9</sup> Helen Coutts, MD,<sup>9</sup> Sandeep Jayawant, MD, FRCPCH,<sup>10</sup> Rosalind Jefferson, MBBS, PhD,<sup>11</sup> Deborah Hughes, MSc,<sup>12</sup> Àngels García-Cazorla, MD, PhD,<sup>13</sup> Detelina Grozeva, PhD,<sup>14,15</sup> F. Lucy Raymond, MD, PhD,<sup>14,15</sup> Belén Pérez-Dueñas, MD, PhD,<sup>1,16</sup> Christian De Goede, MD,<sup>17</sup> Toni S. Pearson, MD,<sup>3,18</sup> Esther Meyer, PhD,<sup>1</sup> and Manju A. Kurian, MD, PhD<sup>1,6\*</sup>

<sup>1</sup>Molecular Neurosciences, Developmental Neurosciences Programme, UCL Great Ormond Street Institute of Child Health, London, United Kingdom

<sup>2</sup>Gene Transfer Technology Group, UCL Institute for Women's Health, London, United Kingdom

<sup>3</sup>Department of Neurology, Icahn School of Medicine at Mount Sinai, New York, USA

<sup>4</sup>Division of Neurology, Department of Medicine, Faculty of Medicine Ramathibodi Hospital, Mahidol University, Bangkok, Thailand

<sup>5</sup>Department of Radiology, Great Ormond Street Hospital for Children NHS Foundation Trust, London, United Kingdom

<sup>6</sup>Department of Neurology, Great Ormond Street Hospital for Children NHS Foundation Trust, London, United Kingdom

<sup>7</sup>Neurometabolic Unit, National Hospital for Neurology and Neurosurgery, London, United Kingdom

<sup>8</sup>Department of Nuclear Medicine and Imaging, Lancashire Teaching Hospitals, NHS Foundation Trust, Preston, United Kingdom

<sup>9</sup>Department of Paediatrics, East Lancashire Hospital NHS Trust, Lancashire, United Kingdom

<sup>10</sup>Department of Paediatric Neurology, John Radcliffe Hospital, Oxford University, NHS Foundation Trust, London, United Kingdom

<sup>11</sup>Department of Paediatrics, Royal Berkshire Hospital, NHS Foundation Trust, Reading, United Kingdom

<sup>12</sup>Molecular Neuroscience and Reta Lila Weston Laboratories, Institute of Neurology, Queen Square, London, United Kingdom

<sup>13</sup>Department of Neurology, Neurometabolic Unit and CIBERER Hospital Sant Joan de Déu, Esplugues de Llobregat, Barcelona, Spain

<sup>14</sup>Medical Genetics, Cambridge Institute for Medical Research, University of Cambridge, Cambridge, United Kingdom

<sup>15</sup>UK10K Project, Wellcome Trust Sanger Institute, Hinxton, Cambridge, United Kingdom

<sup>16</sup>Hospital Vall d'Hebron, Institut de Recerca (VHIR), Barcelona, Spain

<sup>17</sup>Department of Paediatric Neurology, Royal Preston Hospital, Lancashire Teaching Hospitals, NHS Foundation Trust, London, United Kingdom

<sup>18</sup>Department of Neurology, Washington University School of Medicine, St. Louis, Missouri, USA

**ABSTRACT: Background:** Juvenile forms of parkinsonism are rare conditions with onset of bradykinesia, tremor and rigidity before the age of 21 years. These atypical

presentations commonly have a genetic aetiology, highlighting important insights into underlying pathophysiology. Genetic defects may affect key proteins of the

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

\*Correspondence to: Prof. Manju A. Kurian, Neurogenetics Group, Developmental Neurosciences Programme, UCL Great Ormond Street Institute of Child Health, Room 111, Level 1, 30 Guilford Street, London, WC1N 1EH, United Kingdom; E-mail: manju.kurian@ucl.ac.uk

Dr. Ng, Dr. Cortès-Saladelafont, and Dr. Abela contributed equally to the manuscript.

**Funding agencies:** This work was funded by the UK Medical Research Council Doctoral Fellowship (to J.N.), a Wellcome Intermediate Fellowship and NIHR Professorship (to M.A.K.), Sir Jules Thorn Charitable Trust (to M.A.K.), Great Ormond Street Hospital Children's Charity (to J.N., E.M., and M.A.K.), Rosetrees Trust (to J.N., M.A.K.), Agustí Pedró i Pons Foundation, Universitat de Barcelona and Rio Hortega 2015-2017 Institute of Health Carlos III (to E.C.S.), Dystonia Medical Research Foundation (to P.T.), PS09/01132 ISCIII-FEDER, Institute of Health Carlos III (to A.G.C.), European Union Marie Curie training network (to S.J.R.H.), NIHR Biomedical Research Centres, Swiss National Science Foundation Advanced Postdoc. Mobility Fellowship (to L.A.), Cambridge Biomedical Research Centre and Wellcome Trust for UK10K (to F.L.R., D.G.). This study made use of data generated by the UK10K Project award WT091310. A full list of consortium members can be found at the UK10K Project website (<http://www.uk10k.org>). This research was supported by the National Institute for Health Research Biomedical Research Centre at Great Ormond Street Hospital for Children NHS Foundation Trust, University College London.

**Relevant conflicts of interest/financial disclosures:** Nothing to report.

Full financial disclosures and author roles may be found in the online version of this article.

**Received:** 22 October 2019; **Revised:** 15 February 2020; **Accepted:** 3 March 2020

Published online 00 Month 2020 in Wiley Online Library ([wileyonlinelibrary.com](http://wileyonlinelibrary.com)). DOI: 10.1002/mds.28063

endocytic pathway and clathrin-mediated endocytosis (CME), as in *DNAJC6*-related juvenile parkinsonism.

**Objective:** To report on a new patient cohort with juvenile-onset *DNAJC6* parkinsonism-dystonia and determine the functional consequences on auxilin and dopamine homeostasis.

**Methods:** Twenty-five children with juvenile parkinsonism were identified from a research cohort of patients with undiagnosed pediatric movement disorders. Molecular genetic investigations included autozygosity mapping studies and whole-exome sequencing. Patient fibroblasts and CSF were analyzed for auxilin, cyclin G-associated kinase and synaptic proteins.

**Results:** We identified 6 patients harboring previously unreported, homozygous nonsense *DNAJC6* mutations. All presented with neurodevelopmental delay in infancy, progressive parkinsonism, and neurological regression in childhood.  $^{123}\text{I}$ -FP-CIT SPECT (DaTScan) was performed in 3 patients and demonstrated reduced or absent tracer uptake in the basal ganglia. CSF neurotransmitter analysis revealed an isolated reduction of homovanillic acid. Auxilin levels were significantly reduced in both patient fibroblasts and CSF. Cyclin G-associated kinase levels in

CSF were significantly increased, whereas a number of presynaptic dopaminergic proteins were reduced.

**Conclusions:** *DNAJC6* is an emerging cause of recessive juvenile parkinsonism-dystonia. *DNAJC6* encodes the cochaperone protein auxilin, involved in CME of synaptic vesicles. The observed dopamine dyshomeostasis in patients is likely to be multifactorial, secondary to auxilin deficiency and/or neurodegeneration. Increased patient CSF cyclin G-associated kinase, in tandem with reduced auxilin levels, suggests a possible compensatory role of cyclin G-associated kinase, as observed in the auxilin knockout mouse. *DNAJC6* parkinsonism-dystonia should be considered as a differential diagnosis for pediatric neurotransmitter disorders associated with low homovanillic acid levels. Future research in elucidating disease pathogenesis will aid the development of better treatments for this pharmaco-resistant disorder. © 2020 The Authors. *Movement Disorders* published by Wiley Periodicals, Inc. on behalf of International Parkinson and Movement Disorder Society.

**Key Words:** auxilin; *DNAJC6*; dopamine; dystonia; parkinsonism

Classical Parkinson's disease (PD) is an age-related neurodegenerative disorder, mainly affecting adults aged >50 years. Patients typically present with resting tremor, bradykinesia, rigidity, and postural instability. To date, a number of early-onset genetic forms of PD (*Parkin*, *PINK1*, and *DJ-1*)<sup>1-3</sup> and complex parkinsonism syndromes (*ATP13A2*, *PLA2G6*, *FBXO7*, *SLC6A3*, *SLC39A14*, and *PANK2*) have been described.<sup>4-9</sup> Importantly, the study of such monogenic forms of disease have provided significant insight into the pathogenic mechanisms underlying sporadic PD.<sup>10</sup> More recently, two genes, namely *SYNJ1*<sup>11</sup> and *DNAJC6*,<sup>12-14</sup> encoding proteins involved in postendocytotic recycling of synaptic vesicles, have been identified in early-onset parkinsonism.

In this study, we report on 6 children from three families, presenting with juvenile parkinsonism-dystonia associated with novel, biallelic *DNAJC6* mutations. We delineate their clinical phenotype, neuroimaging features (including  $^{123}\text{I}$ -FP-CIT single-photon emission computed tomography [SPECT; DaTScan]) and pattern of cerebrospinal fluid (CSF) neurotransmitter metabolites. Furthermore, we utilized patient fibroblasts and CSF to investigate secondary effects on auxilin, cyclin G-associated kinase (GAK), and dopaminergic proteins.

## Materials and Methods

### Subject Recruitment:

A cohort of 232 children with undiagnosed movement disorders were recruited for research between 2012 and 2016 at UCL Great Ormond Street-Institute of Child

Health (London, UK). A subgroup of 25 patients were identified with juvenile parkinsonism, defined as onset of bradykinesia before 21 years of age, and at least one of the following signs: resting tremor, rigidity, and postural instability. All patients had detailed clinical assessment, undertaken by a movement disorder specialist. Review of (1) the clinical history, (2) features on neuroimaging, and (3) video recordings of the movement disorder at different time points was undertaken. Written informed consent was obtained from participating families, and the study was approved by the local ethics committees (Reference 13/LO/0168).

### Diagnostic CSF Neurotransmitter Analysis:

In order to rule out a primary neurotransmitter disorder, where possible, patients had a routine diagnostic lumbar puncture for CSF neurotransmitter analysis. Using standardized protocols,<sup>15</sup> CSF samples were collected, snap frozen in liquid nitrogen, and stored at  $-80^{\circ}\text{C}$ . Analysis was undertaken using high-pressure liquid chromatography (HPLC) with electrochemical detection and reversed-phase column.<sup>15</sup> Seven anonymized control pediatric CSF samples (with normal CSF neurotransmitter profiles) were obtained from the Neurometabolic Laboratory (National Hospital for Neurology and Neurosurgery, London, UK). All samples were processed and stored in accordance with the UK Royal College of Pathologists guidelines.

### Molecular Genetic Investigation:

From the subgroup of 25 patients with juvenile parkinsonism (16 singletons and 9 familial cases from 3 kindreds), we prioritized two consanguineous families

**TABLE 1.** Clinical characteristics of patients with DNAJC6 mutations from this cohort and literature

	Previously Described DNAJC6-Related Childhood Parkinsonism Dystonia Phenotypes in the Literature						Previously Described DNAJC6-Related Early Onset PD in the Literature											
	Pat. 1 A-III:1	Pat. 2 A-III:4	Pat. 3 A-III:5	Pat. 4 B-IV:2	Pat. 5 B-IV:4	Pat. 6 C	II-2 <sup>12</sup>	II-4	402 <sup>13</sup>	502	504	505	42029 <sup>21</sup>	GPS313 <sup>14</sup>	GPS314	PAL50	PAL54	BR-2662
Presenting age (years)/sex	20/F	12/M	10/M	28/F	19/F	18/F	18/M	13/M	44/F	24/F	31/F	17/M	10/F	48/M	44/F	62/M	46/F	57/M
Consanguinity	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	N	N	N	N	N
Country of origin	PK	PK	PK	PK	PK	PR	PL	PL	TR	TR	TR	TR	YM	NL	NL	BR	BR	BR
Onset of parkinsonism (y)	11	10	9	13	7	10	7	11	10	11	10	10	10	21	29	42	31	24
Parkinsonism at presentation	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	N	Y	N	Y	Y	N
Early development	D	D	D	D	D	D	N	N	N	D	D	D	N	N	N	N	N	N
Bradykinesia	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Tremor	+	+	-	+	-	+	+	+	+	+	+	+	+	+	+	+	+	+
Rigidity	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Hypomimia	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Postural instability	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Dystonic posturing	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Dysarthria	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
Eye movements abnormal	+	-	-	NR	NR	-	-	-	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
Loss of ambulation (y)	13	Unsteady gait	-	13	10	15	18	13	39	21-26	31	20-25	12	Wheelchair bound	-	-	-	-
Motor fluctuations	+	-	-	-	-	+	NR	NR	NR	NR	NR	NR	NR	-	-	+	+	+
Seizures	-	-	-	+	-	+	-	-	+	-	+	+	+	-	-	-	-	-
Cognition	CI	CI	CI	CI	CI	CI	N	N	CI	CI	CI	CI	CI	NR	NR	CI	NR	NR
Spasticity	-	-	-	-	-	-	-	-	+	+	+	+	+	-	-	-	-	-
Psychiatric features	Ax	Ax, PSB	-	Ax	-	AD, AgS	NR	NR	-	Psy after L-dopa, AgB, Hall	MCI	-	Psy, visual and auditory Hall	NR	Psy	NR	NR	NR
Other clinical features	MC, hypothyroid	MC, SD	MC	SD	SD	Mild BP	NR	NR	Scol	MCI	Negative MCI, Scol	-	-	NR	NR	NR	NR	NR
Brain imaging (MRI/CT)	Perisylvian, cerebellar	N	N	Mild, generalized	Mild, generalized, cerebellar	Mild, generalized, thin corpus callosum	N	N	Generalized	N	N	N	N	N	N	N	N	N
(Atrophy)	Absent	Reduced	Not performed	Not performed	Absent	Not performed	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
<sup>123</sup> I-FP-CIT SPECT (DatScan <sup>TM</sup> )	Absent	Reduced	Not performed	Not performed	Absent	Not performed	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
(Uptake in BG)	Absent	Reduced	Not performed	Not performed	Absent	Not performed	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR	NR
Genetics	c.766 C>T	c.766 C>T	c.766 C>T	c.766 C>T	c.766 C>T	c.2416 C>T	c.801-2A>G	c.801-2A>G	c.2200 C>T	c.2200C>T	c.2200 C>T	c.2200 C>T	c.2365 C>T	c.2779 A>G	c.2779 A>G	c.2223 A>T	c.2223 A>T	c.2038+3 A>G and c.1468+33del/-
DNAJC6 mutation	c.766 C>T	c.766 C>T	c.766 C>T	c.766 C>T	c.766 C>T	c.2416 C>T	c.801-2A>G	c.801-2A>G	c.2200 C>T	c.2200C>T	c.2200 C>T	c.2200 C>T	c.2365 C>T	c.2779 A>G	c.2779 A>G	c.2223 A>T	c.2223 A>T	c.2038+3 A>G and c.1468+33del/-
Protein change	R256* N	R256* N	R256* N	R256* N	R256* N	R806* N	NR	NR	0734X NR	0734X NR	0734X NR	0734X NR	Gln789* NR	R927G NR	R927G NR	Thr741* NR	Thr741* NR	NR
Response to L-dopa/carbidopa (maximum dose)	Unstained (150 mg/d)	Not tried	Not tried	Unstained (5.5 mg/kg/d)	No response (10 mg/kg/d)	Some response (200 mg/d)	NR	No response	Good	Good	Therapy refused	Good	Mild	Good, dosages limited by psychiatric side effects	Good	Good	Good	Good

<sup>a</sup> ATP13A2, ATP1A3, ATP7B, ATXN2 (SCA2), ATXN3 (SCA3), C19orf12, C9orf72, CSF1R, DCTN1, DDC, DJ1 (PARK7), EIF4G1, FBXO7, FMR1, FTL, GCH1, GIGYF2, GRN, HTRA2, HTT, LRRK2, MAPT, ND4, NPC1, PANK2, PARKIN (PARK2), PINK1, PPTX3, PLA2G6, POLG, PPP2R2B, PRKRA, PTS, QDPR, SLC6A3, SLC18A2, SLC30A10, SNCA, SPATACSIN (SPG11), SR (SNCG), SYNJ1, TAF1, TBP, TH, and UCHL1. Y, years of age; F, female; M, male; N, no; Y, yes; NR, not reported; PK, Pakistan; PL, Palestine; TR, Turkey; YM, Yemen; NL, Netherlands; BR, Brasil; CI, cognitive impairment; AD, attention deficit; AgB, aggressive behavior; Ax, anxiety; PSB, perseveration behavior; Psy, psychosis; Hall, hallucinations; MC, microcephaly; SD, sleep disturbance; BP, behavioral problems; MCI, myoclonus; Scol, scoliosis; BG, basal ganglia.

(Family A, 3 affected children; Family B, 2 affected children) for initial analysis. These families were investigated using an autozygosity mapping approach, given that the affected children were phenotypically similar and both families originated from the same region in Pakistan. Single-nucleotide polymorphism (SNP) genotyping was performed as previously described.<sup>16</sup> In addition, whole-exome sequencing (WES) was performed for 2 children (A:III-1 and B:IV-2) by UCL Genomics (average WES coverage as previously reported<sup>17</sup>), with an average *DNAJC6* coverage of 30×, with minimum coverage 10× for 82% of the gene. WES data were probed for putative disease-causing *DNAJC6* mutations in the remaining 20 cases (16 sporadic patients, 4 familial cases from a single kindred). This was undertaken through UCL Genomics (8 patients) and Wellcome Trust Sanger Institute (12 patients) within the Wellcome Trust UK10K Rare Diseases project, as previously reported.<sup>18</sup> For patients where *DNAJC6* mutations were identified, whole-exome data were also probed for other genes associated with early-onset dystonia-parkinsonism (Table 1).

### Direct Sanger Sequencing

Sanger sequencing was used to confirm variants identified on WES and to establish familial segregation. A genomic *DNAJC6* sequence (Ensembl transcript: ENST00000371069; NCBI reference sequence: NM\_001256864) was utilized to design primers, using Primer3 software (<http://bioinfo.ut.ee/primer3/>). Primers and polymerase chain reaction (PCR) amplification conditions are available on request. PCR products were cleaned up with MicroCLEAN (Web Scientific) and directly sequenced using Big Dye Terminator Cycle Sequencing System (Applied Biosystems Inc., Foster City, CA). Sequencing reactions were run on an ABI PRISM 3730 DNA Analyzer (Applied Biosystems Inc.) and analyzed with Chromas (<http://www.technelysium.com.au/chromas.html>).

### Fibroblast and CSF Immunoblotting:

Methods to assess protein expression in patient fibroblasts were as previously reported.<sup>17</sup> In brief, primary fibroblast lines were cultured from skin biopsies taken from Patients A-III:1 and B-IV:4 (c.766C>T; p.R256\*) and 2 age-matched healthy donor controls. Antibodies for auxilin and GAK (gift from Professor Green, National Institutes of Health, Washington, DC) and glyceraldehyde 3-phosphate dehydrogenase (GAPDH) horseradish peroxidase (HRP) conjugate (Cell Signaling Technology, Inc., Danvers, MA) were used.

Patients A-III:1, A-III:4, and B-IV:4 underwent lumbar puncture for diagnostic CSF neurotransmitter analysis, and three CSF aliquots were snap frozen and stored at -80°C. Seven age-matched control CSF samples were identified, from subjects without movement disorders on

no medication. Patient and control CSF samples were immunoblotted for auxilin, GAK, and dopaminergic proteins as described previously.<sup>19</sup> CSF protein was probed with the following antibodies: auxilin, GAK, tyrosine hydroxylase (MilliporeSigma, Burlington, MA), dopamine receptor 2 (MilliporeSigma), dopamine transporter (MilliporeSigma), vesicular monoamine transporter 2 (Santa Cruz Biotechnology, Inc., Santa Cruz, CA), and transferrin (Santa Cruz Biotechnology, Inc) as the loading control. Relative protein levels were quantified using ImageJ software (National Institutes of Health, Bethesda, MD) and normalized to the loading control and the mean percentage of optical densitometry of three replicates analyzed with standard error of the mean.

### Statistical Analysis

Statistical analysis was performed using Prism software (version 8; GraphPad Software Inc., La Jolla, CA), with data tested for Gaussian distribution and compared by the Student *t* test.

### Data Availability Statement:

All clinical and experimental data relevant to this study are contained within the article. For Families A, B, and C, there is no ethical approval in place for deposition of whole-exome sequencing genomic data into a public repository. Genomic data from UK10K are available at the EGA European Genomes Phenome Archive (<https://www.ebi.ac.uk/ega/home>), EGAS00001000128(UK10K RARE FIND). Details of statistical analysis can be shared upon request.

## Results

### Patient Cohort


A total of 232 children were referred with undiagnosed movement disorders for genetic research (Fig. 1A). Of these, 25 children (10.7%) had juvenile parkinsonism, 16 females and 9 males with a current median age of 14 years (range, 4–28). Fourteen of 25 had additional neurological features, including dystonia (14 patients), developmental delay/learning difficulties (14 patients), and seizures (4 patients; Fig. 1B).

### Molecular Genetic Investigations

Families A and B (Fig. 1A) were prioritized for autozygosity mapping studies. SNP genotyping revealed a 4.33-Mb region of common homozygosity in both families on chromosome 1, between rs640407 (64,267,606 base pairs [bp]) and rs2566784 (68,602,735 bp; Fig. 2A,B). This region showed a common haplotype in all affected children whereas unaffected siblings had a different haplotype. It was therefore considered to be the likely disease locus (Fig. 2B).

WES performed in Patients A-III:1 and B-IV:2 revealed 23,365 and 23,549 variants, respectively. Given familial

(A)



(B)

Family	Gender	Current age (years)	Clinical presentation	Parkinsonism	Dystonia	Dyskinesia	Tremor	Spasticity	Seizures	CSF Neurotransmitter analysis	MRI	Other clinical features Response to medication
1	M	4	Infancy	x	x					N	N	Deep Brain Stimulator inserted
2	F	6	NR	x	x				x	Abnormal	N	
3	F	23	Childhood	x	x					Abnormal	N	L-Dopa non-responsive
4	M	24	Childhood	x	x					Abnormal	N	Cognitive impairment, psychiatric symptoms
5	F	5	Early infancy	x						Abnormal	N	L-Dopa responsive
6	M	7	Infancy	x						N	Mild bilateral pallidal signal abnormalities	
7	M	4	Early infancy	x	x					N	N	Developmental delay
8	F	12	Infancy	x	x	x				Abnormal	N	Axial hypotonia
*9.1	F	16	Toddler	x	x	x		x		N	N	N cognition. Altered eye movements
*9.2	F	14	Toddler	x	x	x		x		N	N	N cognition. Altered eye movements
*9.3	F	12	Toddler	x	x	x		x		N	N	N cognition. Altered eye movements
*9.4	M	6	Toddler	x	x	x		x		N	N	N cognition. Altered eye movements
10	F	14	Childhood	x						N	N	L-Dopa responsive. Hypotonia. Developmental delay.
11	M	15	Toddler	x	x					Abnormal	Generalised atrophy	L-Dopa responsive
12	M	Died at 21 years	Childhood	x	x		x			NR	NR	Behavioural difficulties. Died at 21 years
13	F	8	Neonatal	x						Abnormal	N	Contractures. Eye movement abnormalities
14	F	Died at 20 years	Toddler	x	x			x		NP	Generalised atrophy	Behavioural abnormalities. Died at 21 years
15	F	Died at 23 years	Adolescence	x	x				x	Abnormal	Atrophy (caudate)	Myoclonus. Mental impairment. Died at 23 years
16	F	26	Adolescence	x				x		NR	NR	Mild cognitive impairment
17	F	18	Toddler	x	x				x	Abnormal	This CC, generalised atrophy	Early childhood motor delay, cognitive impairment. L-Dopa responsive.

**FIG. 1.** Juvenile parkinsonism cohort: clinical features and molecular genetic investigation. (A) Flowchart demonstrating the pathway of molecular genetic investigations in a subcohort of 25 children with juvenile parkinsonism. (B) Clinical characteristics of 20 children from 17 families. Early infancy <3 months; infancy 3 to 12 months; toddler 12 to 24 months; childhood 2 to 13 years; adolescence 13 to 18 years. \*Consanguineous family. M, male; F, female; N, normal; NP, not performed; NR, not reported. [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]


(A)

	Family A				Family B			
	Start of Homozygous Region		End of homozygous region		Start of Homozygous Region		End of Homozygous Region	
	Reference SNP Cluster ID (rs number)	Physical Position (bp)	rs number	Physical Position (bp)	rs number	Physical Position (bp)	rs number	Physical Position (bp)
<b>Chromosome 1</b>	rs2819130	64 231 541	rs2772304	68 636 580	rs41285364	63 786 518	rs6687262	79 509 109
<b>Chromosome 2</b>					rs300758	72 184	rs809672	12 688 452
<b>Chromosome 5</b>					rs1859295	135 327 491	rs13153997	151 080 003
<b>Chromosome 5</b>					rs13358102	171 650 965	rs6887234	173 616 645
<b>Chromosome 7</b>					rs12669653	128 746 301	rs3800707	134 472 116
<b>Chromosome 11</b>	rs10836509	36 177 369	rs3133269	67 804 156				
<b>Chromosome 12</b>					rs4469939	77 288 346	rs2114926	101 093 560
<b>Chromosome 17</b>					rs9911464	32 468 227	rs7216307	42 413 481
<b>Chromosome 18</b>					rs292324	5 167 441	rs4800629	22 901 551


(B)



(C)



(D)



**FIG. 2.** Molecular genetic investigations and DaTSCAN imaging. (A) Family A and B SNP array results showing homozygous regions detected. For each chromosome, the start and end point is specified using the Reference SNP Cluster ID (rs number) and physical position. (B) Homozygous SNPs are represented in light blue (AA) and dark blue (BB), heterozygous SNPs in red (AB), and “no calls” in white. (C) Sanger Sequencing confirms a homozygous *DNAJC6* mutation, c.766C > T (p.R256\*), in all affected children of Family A (A-III:1, A-III:4, and A-III:5) and Family B (B-IV:2, B-IV:4). Parents are heterozygous carriers. (D) I-123-DaTSCAN™ with SPECT imaging in a control subject, Patient A-III:1 (19 years 3 months), Patient A-III:4 (11 years 4 months), and Patient B-IV:4 (17 years). In Patients A-III:1 and B-IV:4, DaTSCAN findings indicate virtually complete absence of tracer uptake in the basal ganglia, with very high background activity, suggesting loss of presynaptic dopaminergic terminals, whereas Patient A-III:4 showed significantly reduced, albeit still visible, uptake in the head of caudate (left better than right, white arrows). [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

consanguinity and the autozygosity mapping results, targeted analysis for recessive pathogenic variants within the putative disease locus on chromosome 1 was undertaken. A single homozygous nonsense variant c.766C>T (p.R256\*) in *DNAJC6* (Chr1, 65,248,219–65,415,869) was identified both in A-III:1 and B-IV:2 on WES, located within the common region of homozygosity. No other pathogenic changes in previously reported genes causing juvenile parkinsonism-dystonia phenotypes were identified from the WES data. Direct Sanger sequencing confirmed the homozygous c.766C>T mutation in all 5 affected patients, and familial segregation studies revealed that all parents were obligate carriers in both kindreds, with unaffected siblings either wild type or heterozygous for the identified variant (Fig. 2C). WES/whole-genome sequencing data for the remainder of the parkinsonism-dystonia cohort (n = 20) was interrogated for *DNAJC6* mutations. This led to the identification of a homozygous nonsense variant (c.2416C>T) in a sixth unrelated patient (Patient C), which was confirmed on Sanger sequencing.

### Delineation of the Clinical Phenotype of *DNAJC6* Patients

#### Family A (3 Affected Patients)

Patients A-III:1, A-III:4, and A-III:5 are 3 affected children born to first-cousin parents, currently 20, 12, and 10 years old (Table 1). Two other brothers (A-III:2 and A-III:3), aged 17 and 15 years, have mild learning difficulties without evidence of a movement disorder. The paternal grandfather was diagnosed with PD in his fifties.

All 3 children were born at term after an uneventful antenatal period. Microcephaly was evident at birth (head circumference: <0.4th centile), but nonprogressive over time. All siblings had early neurodevelopmental delay and moderate learning difficulties.

A-III:1 is the eldest daughter, aged 20 years. She presented at 10 years, with a 6-week history of feeding difficulties, vomiting, and weight loss. Over time, she developed fever, unsteady gait, facial asymmetry, left-sided tremor, and generalized seizures and was diagnosed with an encephalitis of uncertain etiology. She recovered from this acute illness, but subsequently had progressive bradykinesia, with tremor and rigidity, and loss of independent ambulation at 13 years, associated with cognitive decline. She is now wheelchair dependent, with generalised cogwheel-rigidity, severe bradykinesia, multiple limb contractures and emotional lability (Video 1). She also has severe gut dysmotility, with recurrent vomiting, and required a gastrostomy for deteriorating bulbar dysfunction. CSF neurotransmitter analysis (age 11 years), while on levodopa therapy, revealed an isolated low 5-hydroxyindoleacetic acid (5-HIAA; Fig. 3A). At 12 years 11 months, when off L-dopa, CSF HVA, and HVA:5-HIAA ratio were low. Brain MRI showed evidence of right-sided atrophy

of the perisylvian region and right cerebellum by 19 years of age (Supporting Information Fig. S1). At 19 years, <sup>123</sup>I-FP-CIT SPECT (DaTScan) showed absent uptake in the basal ganglia when compared to normal subjects (Fig. 2D). At this stage, while on L-dopa treatment, her CSF HVA levels normalized (Fig. 3A). Her condition is refractory to medical treatment, with no clinical response to trihexyphenidyl, benzhexol, procyclidine, clobazam, rotigotine, and apomorphine. L-dopa has proven difficult to titrate because of marked drug sensitivity. She experiences an improvement in motor function and speech 30 minutes postdose, after which she returns to the *off* state. L-dopa dosages >150 mg/d have resulted in drug-related dyskinesias.

Her two brothers (A-III:4 and A-III:5) presented with fine motor difficulties at 8 years of age. They subsequently developed positional tremor, upper limb dystonic posturing, hypophonia, hypomimia, bradykinesia, cogwheel rigidity, and postural instability over 12 months (Videos 2 and 3). Like their sister, both have gastrointestinal complications with sialorrhea, recurrent vomiting, and feeding difficulties, necessitating gastrostomy insertion. A-III:4 is currently 12 years old and suffers from anxiety and perseveration. His CSF-HVA levels are at the lower limit of normal, with a low HVA:5-HIAA ratio <1.0 (Fig. 3A). MRI brain scan was normal. <sup>123</sup>I-FP-CIT SPECT (DaTScan) at 11 years showed profound reduction in tracer uptake in the basal ganglia (Fig. 2D). Both boys responded to treatment with transdermal rotigotine and oral trihexyphenidyl, but with increasing doses, both experienced dyskinesias, necessitating dose reduction.

#### Family B (2 Affected Patients)


Patients B-IV:2 and B-IV:4 are 2 affected girls, born to first-cousin parents, and currently 28 and 19 years old. Both were born uneventfully following a normal pregnancy, presenting with early feeding difficulties, hypotonia, and delayed milestones by 6 months old. Both made slow developmental progress, achieving independent ambulation and spoken language by 3 years of age.

B-IV:2 presented at 9 years with generalized seizures that stabilized with lamotrigine therapy. From 13 years of age, motor and cognitive deterioration ensued, with onset of parkinsonism and loss of speech and ambulation. She experienced anxiety and recurrence of seizures. She has severe antecollis, hypomimia, tremor, generalised cogwheel rigidity, bradykinesia, and positive glabellar tap (Video 4). MRI was normal until 18 years, after which there was radiological evidence of mild generalized atrophy. Several medications were tried without clinical benefit, including L-dopa (maximum, 10 mg/kg/d), selegiline, rotigotine, and trihexyphenidyl. The younger sibling, B-IV:4, presented at 7 years with gait deterioration, bradykinesia, and cogwheel rigidity. She lost


(A)

Patient	A-III:1	A-III:1	A-III:1	A-III:4	B-IV:4	B-IV:4	B-IV:4	C
Age at LP	11y 8m	12y 11m	19y 3m	9y 4m	4y 2m	4y 11m	14y 11m	14y
Levodopa treatment at time of LP	Y	N	Y	N	N	N	Y	N
HVA (nmol/L) (#) <sup>1,2</sup>	190 (71-565) <sup>1</sup>	60 (71-565) <sup>1</sup>	154 (71-565) <sup>1</sup>	115 (71-565) <sup>1</sup>	48 (71-565) <sup>1</sup>	40 (71-565) <sup>1</sup>	47 (71-565) <sup>1</sup>	32 (157-563) <sup>2</sup>
5-HIAA (nmol/L) (#) <sup>1,2</sup>	54 (58-220) <sup>1</sup>	147 (58-220) <sup>1</sup>	152 (58-220) <sup>1</sup>	173 (58-220) <sup>1</sup>	90 (58-220) <sup>1</sup>	56 (58-220) <sup>1</sup>	116 (58-220) <sup>1</sup>	87 (67-189) <sup>2</sup>
HVA:5-HIAA ratio (normal 1.0-3.7)	3.5	0.4	1.1	0.7	0.5	0.7	0.4	0.37
3-O-Methyldopa present (nmol/L if quantified) (#) <sup>2</sup>	Y	N	730	N	Not quantified	Not quantified	391	10 (<100) <sup>2</sup>
Neopterin (nmol/L) (#) <sup>1,2</sup>	11 (7-65) <sup>1</sup>	NP	13 (7-65) <sup>1</sup>	11 (7-65) <sup>1</sup>	Not quantified	12 (7-65) <sup>1</sup>	19 (7-65) <sup>1</sup>	15 (8-33) <sup>2</sup>
Tetrahydrobiopterin (nmol/L) (#) <sup>1,2</sup>	16 (9-39) <sup>1</sup>	NP	10 (9-39) <sup>1</sup>	14 (9-39) <sup>1</sup>	Not quantified	15 (9-39) <sup>1</sup>	17 (9-39) <sup>1</sup>	8 (9-32) <sup>2</sup>
Dihydrobiopterin (nmol/L) (#) <sup>1</sup>	4.8 (0.4-13.9) <sup>1</sup>	NP	5.6 (0.4-13.9) <sup>1</sup>	5.4 (0.4-13.9) <sup>1</sup>	Not quantified	0.5 (0.4-13.9) <sup>1</sup>	5.1 (0.4-13.9) <sup>1</sup>	Not quantified
5-MTHF (nmol/L) (#) <sup>3</sup>	66 (46-160) <sup>3</sup>	NP	71 (46-160) <sup>3</sup>	65 (46-160) <sup>3</sup>	Not quantified	Not quantified	82 (46-160) <sup>3</sup>	89 (40-160) <sup>3</sup>


(B)




(C)



(D)



(E)



**FIG. 3.** CSF neurotransmitter analysis and patient fibroblast and CSF immunoblotting. (A) CSF neurotransmitter analysis. Age-related reference ranges indicated in brackets after each value. Red: abnormal result. Gray: borderline result. Symbol (“#”) indicates reference range: <sup>1</sup>Keith Hyland, Robert A.H. Surtees, et al. *Pediatr Res* 1993;34:10–14; <sup>2</sup>Keith Hyland, *Future Neurol* 2006;1:593–603; <sup>3</sup>Surtees R, Hyland K. *Biochem Med Metab Biol* 1990;44:192–199. (B) Scatterplot of CSF HVA and 5-HIAA levels (nmol/L) measured by high performance liquid chromatography (patient = red shapes, control = black triangles). Medication at time of CSF sampling: A-III-1: co-careldopa, melatonin, glycopyronium; A-III-4: none; B-IV-4: L-dopa, pyridoxine; Control 1: none; Control 2: none. Immunoblot of auxilin and GAK in patient fibroblasts (C) and CSF (D) compared to controls. (E) Immunoblot of patient CSF for TH, DAT, VMAT, and D2R protein levels measured compared to controls. Graphs show mean protein percent optical density (OD) normalized to loading control in patients (red) and controls (black). LP, lumbar puncture; y, years; m, months; 5-MTHF, 5-methyltetrahydrofolate; NP, not performed. [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]



independent ambulation and speech by 10 years (Video 5). She has developed dystonic posturing, bulbar dysfunction (necessitating gastrostomy), and a disrupted sleep pattern. The MRI brain scan was initially normal, but by 16 years showed subtle global cerebral atrophy (particularly in the posterior regions) as well as cerebellar atrophy (Supporting Information Fig. S2).  $^{123}\text{I}$ -FP-CIT SPECT (DaTScan) at 17 years showed profound reduction in tracer uptake in the basal ganglia (Fig. 2D). CSF HVA and HVA:5-HIAA ratio were reduced at ages 4 and 14 years. CSF-HIAA levels were reduced at age 4 years (Fig. 3A). She showed an initial response to L-dopa, but developed emotional lability at 5.5 mg/kg/d, leading to drug withdrawal. There was no clinical improvement observed with trihexyphenidyl or chloral hydrate. She had a modest response to pramipexole, with improved facial expression, reduced tremor, and increase in voluntary movements.

### Family C

This 18-year-old girl is the third child of distantly related Latin American parents, with 2 healthy siblings. She initially presented with neonatal feeding difficulties and hypotonia. In infancy, she showed delay in attaining milestones and developed seizures characterized by staring episodes with loss of tone. She walked independently from 2 years, but by 10 years of age her gait deteriorated, leading to frequent falls, postural instability, and losing the ability to run. Over the next 4 years, she continued to deteriorate with worsening antecollis and bradykinesia (Video 6). She developed severe bulbar dysfunction with sialorrhea, dysarthria, and, dysphagia, leading to considerable weight loss. At 12 years, she developed generalized tonic-clonic seizures and atypical absences, responsive to lamotrigine and zonisamide therapy. MRI demonstrated subtle generalized cerebral atrophy and CSF HVA was low (Fig. 3A). Her movement disorder responded to L-dopa, with improved tremor, gait, and a reduction in drooling. A maximum of 200 mg/d was tolerated, but further increases led to intolerable drug-induced dyskinesias. After 4 months of treatment, she developed aggressive behavior and received treatment with quetiapine. By 16 years, she became increasingly sensitive to L-dopa, with peak-dose agitation, restlessness, and dyskinesia. Lowering the dose improved side effects, and continued to provide motor benefit, although the effects wore off 2 to 3 hours after administration. *On-off* phenomena were commonly reported, and in the *off* state, she was often akinetic and rigid. Introduction of trihexyphenidyl improved rigidity, but not immobility.

### Patient CSF and Fibroblast Analysis

CSF HPLC analysis of the *DNAJC6* patient cohort showed reduction in CSF-HVA levels ( $P = 0.002$ ) compared to controls (but not 5-HIAA levels) in 3 patients


(Fig. 3A,B). Patient fibroblasts showed reduced auxilin ( $P = 0.009$ ) and a trend for increased GAK protein ( $P = 0.11$ ; Fig. 3C). Patient CSF auxilin levels were even more significantly reduced ( $P = 0.0015$ ; Fig. 3D). Notably, CSF-GAK levels were significantly increased in patients ( $P = 0.0014$ ; Fig. 3D). CSF immunoblotting studies showed that several key components of the dopaminergic synapse were significantly reduced, including tyrosine hydroxylase (TH;  $P = 0.0001$ ), vesicular monoamine transporter (VMAT;  $P = 0.0002$ ), dopamine transporter (DAT;  $P = 0.0003$ ), and D2 receptor (D2R;  $P = 0.002$ ; Fig. 3E).

## Discussion

Juvenile parkinsonism attributed to *DNAJC6* mutations has only recently been reported. Here, we report on a further 6 patients from three families, with two previously unreported homozygous nonsense mutations in *DNAJC6*. Moreover, our findings on  $^{123}\text{I}$ -FP-CIT SPECT (DaTScan) imaging, CSF analysis, and immunoblotting suggest downstream dyshomeostasis of auxilin, GAK, and dopaminergic proteins in *DNAJC6*-related disease.

Our data confirms that all reported cases of juvenile-onset *DNAJC6*-parkinsonism have core clinical characteristics (Table 1), including (1) clinical presentation of progressive parkinsonism toward of the first decade (median, 10 years; range, 7–13), (2) significant neurological regression thereafter, and (3) loss of ambulation in mid-adolescence.<sup>12-14,21</sup> In contrast to adult-onset PD, childhood parkinsonian disorders rarely present with a “pure” parkinsonian phenotype, as illustrated by the classical primary pediatric monoamine neurotransmitter disorders.<sup>20</sup> Similarly, in early-onset *DNAJC6*-related disease, parkinsonism is commonly present in tandem with a multitude of other clinical features, including dystonia, moderate learning difficulties, epilepsy, and neuropsychiatric features<sup>12-14,21</sup> (Table 1). Furthermore, many of our patients had evidence of bulbar dysfunction, gut dysmotility, and sleep disturbance. The majority of our patients showed limited response to L-dopa and other standard therapies for parkinsonism-dystonia. They experienced severe, often intolerable, side effects with dopaminergic agents, including *on-off* phenomenon and severe dyskinesia, particularly at higher drug dosages.

$^{123}\text{I}$ -FP-CIT SPECT (DaTScan) was performed in 3 patients, demonstrating reduced tracer uptake in the basal ganglia, suggestive of impaired presynaptic dopamine uptake and striatonigral neurodegeneration. Postmortem studies have confirmed striatal dopamine deficiency in patients with parkinsonism.<sup>22,23</sup> Together, these observations suggest a neurodegenerative process in *DNAJC6* patients. MRI brain imaging further corroborates this



**FIG. 4.** Schematic representation of *DNAJC6*-encoded auxilin protein function. The role of auxilin in synaptic vesicle recycling and endocytic pathway. (A) A nascent clathrin-coated pit is formed at the presynaptic membrane, followed by membrane invagination; (B) the pit develops into a clathrin-coated vesicle, where the clathrin lattice consists of (C) clathrin triskelions formed by three crossed ankle regions. (D) Auxilin (pale green square) binds to an exposed domain in the heavy chain of the clathrin. (E) Auxilin facilitates a conformational change in clathrin that allows binding of Hsc70 and by ATPase-mediated activity, and the clathrin lattice is disrupted and distorted, leading to (F) clathrin disassembly allowing subsequent delivery of cargo neurotransmitters to the membrane or other vesicle in the endocytic pathway. Hsc70, heat shock cognate protein 70. [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

hypothesis; the mild generalized cerebral and/or cerebellar atrophy in 4 patients suggests that *DNAJC6*-related disorders may also be associated with neuronal loss in other regions of the central nervous system.

All 6 cases fit the juvenile phenotype associated with this gene, though more recently, *DNAJC6* mutations have been reported in early adult-onset PD.<sup>14</sup> Although there are a number of overlapping features (progressive parkinsonism, psychiatric features), affected patients presented later (range, 21–42 years) and seizures and cognitive decline are not reported.

Homozygous and compound heterozygous mutations in *DNAJC6* are predicted to result in loss of protein function. To date, splice-site variants,<sup>12</sup> large multiexonic deletions,<sup>24</sup> truncating mutations,<sup>13</sup> and missense mutations<sup>14</sup> have been reported. All 6 patients in our cohort had nonsense mutations, predicted to cause nonsense-mediated decay or premature protein truncation. Five of the 6 reported patients are from two consanguineous families originating from the same region in Pakistan, and all have the same nonsense mutation. SNP array confirmed a common haplotype at this disease locus for all affected children, suggesting a possible founder effect.

*DNAJC6* encodes for auxilin, a neuronally expressed J-chaperone protein involved in the uncoating of clathrin-coated vesicles<sup>25,26</sup> (Fig. 4). Auxilin modifies the

three-dimensional conformation of heavy-chain clathrin triskelions, leading to clathrin coat distortion, instability, and subsequent disassembly.<sup>25,26</sup> Neurotransmission involves rapid continuous recycling of synaptic vesicles through CME. Deficiency in auxilin ultimately results in impairment of synaptic vesicle recycling and impaired neurotransmission. Similarly, aberrant synaptic vesicular trafficking is also evident in other forms of early-onset parkinsonism, including *LRRK2*,<sup>27,28</sup> *VMAT2*,<sup>29</sup> and *SNCA*-related disease.<sup>30</sup> Clathrin-mediated endocytosis is crucial for the regulation of developmental signaling pathways through internalization of receptors or ligands and is required for axon and dendrite outgrowth.<sup>31</sup> Presence of developmental delay well before onset of parkinsonism in patients with *DNAJC6* mutations further corroborates the notion that auxilin is likely to have a central role in neurodevelopment, given its role in CME. In *Drosophila*, auxilin is crucial for Notch signaling, a developmental pathway that regulates neural stem-cell proliferation, survival, renewal, and differentiation, as well as neuronal specification of dopaminergic neurons.<sup>32–34</sup>

To investigate the downstream effects of *DNAJC6* mutations, we studied auxilin and GAK protein levels in 2 patients using patient fibroblasts and CSF. Auxilin is a neuron-specific protein, enriched in presynaptic terminals, whereas GAK is an ubiquitously expressed

protein.<sup>35,36</sup> Auxilin and GAK are highly homologous proteins that both have the ability to bind clathrin and clathrin adaptor protein 2 in order to initiate clathrin uncoating of endocytosed vesicles.<sup>37</sup> In the auxilin knockout mouse model, it is reported that upregulation of GAK can partially compensate for the loss of auxilin and decrease mortality.<sup>35</sup> We therefore wished to determine whether a similar compensatory mechanism was evident in our patients. In our study, we observed that patient fibroblast auxilin protein levels were significantly reduced when compared to controls, as previously reported.<sup>14</sup> We found that patient fibroblast GAK levels were slightly, but not significantly, increased, whereas patient CSF GAK protein levels were significantly increased. Our findings support upregulation of brain GAK levels in *DNAJC6* patients, partially compensating for auxilin reduction, as evident in the auxilin knockout mouse model.<sup>35</sup>

Diagnostic CSF neurotransmitter analysis revealed that levels of the stable dopamine metabolite (HVA) were either below the age-related reference ranges or close to the lower limit of normal in our patients, indicating impaired dopamine turnover. Indeed, CSF-HVA levels and HVA:5-HIAA ratios were comparable to those observed in TH deficiency, an inherited dopamine synthesis defect associated with central dopamine deficiency.<sup>20</sup> In order to determine how *DNAJC6* mutations may impact the dopaminergic system, we used patient CSF to analyze proteins involved in dopamine signaling and homeostasis. We observed that patient CSF had significantly reduced levels of VMAT, DAT, TH, and D2R when compared to controls. <sup>123</sup>I-FP-CIT SPECT (DaTScan) imaging additionally provides in vivo evidence of impaired DAT function in *DNAJC6* patients. VMAT and DAT are both synaptic transporters recycled through clathrin-mediated endocytosis.<sup>38,39</sup> The reduction in HVA associated with low VMAT/DAT protein levels may imply that the observed dopamine deficiency is associated with impaired clathrin-mediated neurotransmitter recycling. D2R is also postulated to be internalized through clathrin-mediated endocytosis.<sup>38,39</sup> Neurons internalize receptors to adjust excitability and degrade, resensitize, and recycle desensitized receptors.<sup>38,39</sup> *DNAJC6* mutations thus may affect D2R protein levels and normal postsynaptic function. It is likely that presynaptic D2R autoreceptor function will also be affected, leading to aberrant TH regulation.<sup>40</sup>

Overall, our findings suggest that the mechanisms governing *DNAJC6*-associated parkinsonism are likely to be multifactorial. Another plausible explanation for the reduction in synaptic protein levels may be as a result of neurodegeneration secondary to defective chaperone function. Auxilin and other J-chaperone proteins play a crucial role in regulating the folding and conformational change of proteins to maintain integrity in the neuron.<sup>41,42</sup> Indeed, in the auxilin knockout mouse model, there is

sequestration of clathrin cages in the cerebellum.<sup>35</sup> With impaired auxilin function, a cumulative effect of sequestered misfolded proteins and accumulation of clathrin coat components in assembled coats and cages may lead to apoptotic cascades and neurodegeneration. There is growing interest in the role of such chaperone proteins in human disease and mutations in eight distinct J proteins (*DNAJB2*, *DNAJB6*, *DNAJC5*, *DNAJC6*, *DNAJC12*, *DNAJC13*, *DNAJC19*, and *DNAJC29*) have been described.<sup>43,44</sup> Future research into such “chaperonopathies” may provide further insights into neurodegenerative disorders.

In conclusion, we report on a cohort of patients with previously unreported *DNAJC6* mutations associated with early neurodevelopmental delay, juvenile parkinsonism, and neurological regression in the second decade of life. We further demonstrate disturbance of dopamine homeostasis in patient-derived CSF and report on a possible GAK-mediated compensatory mechanism for auxilin deficiency. Mutations in *DNAJC6* are rare, but a likely under-recognized cause of parkinsonism-dystonia in infants and children. Elucidating the genetic diagnosis has important implications for families given that early diagnosis negates the need for extensive invasive investigations, facilitates treatment strategies, and aids genetic counseling for future pregnancies. The early clinical features and CSF neurotransmitter signature observed in our patients can mimic primary neurotransmitter disorders, and *DNAJC6* mutations should thus be considered as a differential diagnosis. We observed reduced auxilin and increased GAK protein levels, suggesting a possible compensatory role for GAK in this condition. Study of CSF synaptic proteins suggest downstream effects on dopamine synthesis, recycling, homeostasis, and signaling that may result from a combination of primary auxilin deficiency and neurodegeneration. Abnormal synaptic vesicle dynamics are increasingly recognized as a disease mechanism in neurodegenerative parkinsonian disorders, and future research into elucidating the pathogenesis of such conditions will no doubt assist the development of novel targeted treatments. ■

**Acknowledgments:** We thank the patients and families for their participation in this study. We acknowledge Kerra Pearce, Mark Kristiansen, and Alan Pittman (UCL Genomics) for assistance with the SNP arrays and whole-exome sequencing. We thank Professor Lois Greene for kindly providing the anti-auxilin and GAK antibodies.

## References

1. Kitada T, Asakawa S, Hattori N, et al. Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism. *Nature* 1998;392:605–608.
2. Valente EM, Abou-Sleiman PM, Caputo V, et al. Hereditary early-onset Parkinson's disease caused by mutations in PINK1. *Science* 2004;304:1158–1160.

3. Bonifati V, Rizzu P, van Baren MJ, et al. Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism. *Science* 2003;299:256–259.
4. Ramirez A, Heimbach A, Gründemann J, et al. Hereditary parkinsonism with dementia is caused by mutations in ATP13A2, encoding a lysosomal type 5 P-type ATPase. *Nat Genet* 2006;38:1184–1191.
5. Paisan-Ruiz C, Bhatia KP, Li A, et al. Characterization of PLA2G6 as a locus for dystonia-parkinsonism. *Ann Neurol* 2009;65:19–23.
6. Fonzo AD, Dekker MCJ, Montagna P, et al. FBXO7 mutations cause autosomal recessive, early-onset parkinsonian-pyramidal syndrome. *Neurology* 2009;72:240–245.
7. Ng J, Zhen J, Meyer E, et al. Dopamine transporter deficiency syndrome: phenotypic spectrum from infancy to adulthood. *Brain* 2014;137:1107–1119.
8. Tuschl K, Meyer E, Valdivia LE, et al. Mutations in SLC39A14 disrupt manganese homeostasis and cause childhood-onset parkinsonism-dystonia. *Nat Commun* 2016;7:11601.
9. Schneider SA, Bhatia KP, Hardy J. Complicated recessive dystonia parkinsonism syndromes. *Mov Disord* 2009;24:490–499.
10. Bonifati V. Genetics of Parkinson's disease—state of the art, 2013. *Parkinsonism Relat Disord* 2014;20(Suppl. 1):S23–S28.
11. Quadri M, Fang MY, Picillo M, et al. Mutation in the SYNJ1 gene associated with autosomal recessive, early-onset parkinsonism. *Hum Mutat* 2013;34:1208–1215.
12. Edvardson S, Cinnamon Y, Ta-Shma A, et al. A deleterious mutation in DNAJC6 encoding the neuronal-specific clathrin-uncoating co-chaperone auxilin, is associated with juvenile parkinsonism. *PLoS One* 2012;7:e36458.
13. Koroglu C, Baysal L, Cetinkaya M, Karasoy H, Tolun A. DNAJC6 is responsible for juvenile parkinsonism with phenotypic variability. *Parkinsonism Related Disord* 2013;19:320–324.
14. Olgiati S, Quadri M, Fang MY, et al. DNAJC6 mutations associated with early-onset Parkinson's disease. *Ann Neurol* 2016;79:244–256.
15. Hyland K, Surtees RA, Heales SJ, Bowron A, Howells DW, Smith I. Cerebrospinal fluid concentrations of pterins and metabolites of serotonin and dopamine in a pediatric reference population. *Pediatr Res* 1993;34:10–14.
16. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* 2010;26:841–842.
17. Meyer E, Carss KJ, Rankin J, et al. Mutations in the histone methyltransferase gene KMT2B cause complex early-onset dystonia. *Nat Genet* 2017;49:223–237.
18. Consortium UK, Walter K, Min JL, et al. The UK10K project identifies rare variants in health and disease. *Nature* 2015;526:82–90.
19. Ortez C, Duarte ST, Ormazábal A, et al. Cerebrospinal fluid synaptic proteins as useful biomarkers in tyrosine hydroxylase deficiency. *Mol Genet Metab* 2015;114:34–40.
20. Willemsen MA, Verbeek MM, Kamsteeg EJ, et al. Tyrosine hydroxylase deficiency: a treatable disorder of brain catecholamine biosynthesis. *Brain* 2010;133:1810–1822.
21. Elsayed LE, Drouet V, Usenko T, et al. A novel nonsense mutation in DNAJC6 expands the phenotype of autosomal-recessive juvenile-onset Parkinson's disease. *Ann Neurol* 2016;79:335–337.
22. Cheng HC, Ulane CM, Burke RE. Clinical progression in Parkinson disease and the neurobiology of axons. *Ann Neurol* 2010;67:715–725.
23. Burke RE, O'Malley K. Axon degeneration in Parkinson's disease. *Exp Neurol* 2013;246:72–83.
24. Vauthier V, Jaillard S, Journel H, Dubourg C, Jockers R, Dam J. Homozygous deletion of an 80 kb region comprising part of DNAJC6 and LEPR genes on chromosome 1P31.3 is associated with early onset obesity, mental retardation and epilepsy. *Mol Genet Metab* 2012;106:345–350.
25. Ungewickell E, Ungewickell H, Holstein SEH, et al. Role of auxilin in uncoating clathrin-coated vesicles. *Nature* 1995;378:632–635.
26. Eisenberg E, Greene LE. Multiple roles of auxilin and Hsc70 in clathrin-mediated endocytosis. *Traffic* 2007;8:640–646.
27. Migheli R, Del Giudice MG, Spissu Y, et al. LRRK2 affects vesicle trafficking, neurotransmitter extracellular level and membrane receptor localization. *PLoS One* 2013;8:e77198.
28. Xiong Y, Dawson Valina L, Dawson Ted M. LRRK2 GTPase dysfunction in the pathogenesis of Parkinson's disease. *Biochem Soc Trans* 2012;40:1074–1079.
29. Rilstone JJ, Alkhatir RA, Minassian BA. Brain dopamine-serotonin vesicular transport disease and its treatment. *N Engl J Med* 2013;368:543–550.
30. Alter SP, Lenzi GM, Bernstein AI, Miller GW. Vesicular integrity in Parkinson's disease. *Curr Neurol Neurosci Rep* 2013;13:362–362.
31. Yap CC, Winckler B. Adapting for endocytosis: roles for endocytic sorting adaptors in directing neural development. *Front Cell Neurosci* 2015;9:119.
32. Eun SH, Banks SML, Fischer JA. Auxilin is essential for Delta signaling. *Development* 2008;135:1089–1095.
33. Lathia JD, Mattson MP, Cheng A. Notch: from neural development to neurological disorders. *J Neurochem* 2008;107:1471–1481.
34. Tio M, Toh J, Fang W, Blanco J, Udolph G. Asymmetric cell division and Notch signaling specify dopaminergic neurons in *Drosophila*. *PLoS One* 2011;6:e26879.
35. Yim YI, Sun T, Wu LG, et al. Endocytosis and clathrin-uncoating defects at synapses of auxilin knockout mice. *Proc Natl Acad Sci U S A* 2010;107:4412–4417.
36. Ahle S, Ungewickell E. Auxilin, a newly identified clathrin-associated protein in coated vesicles from bovine brain. *J Cell Biol* 1990;111:19–29.
37. Scheele U, Kalthoff C, Ungewickell E. Multiple interactions of auxilin 1 with clathrin and the AP-2 adaptor complex. *J Biol Chem* 2001;276:36131–36138.
38. Ferguson SSG, Caron MG. G protein-coupled receptor adaptation mechanisms. *Semin Cell Dev Biol* 1998;9:119–127.
39. Sorkin A, Von Zastrow M. Signal transduction and endocytosis: close encounters of many kinds. *Nat Rev Mol Cell Biol* 2002;3:600–614.
40. Blackstone C. Infantile parkinsonism-dystonia: a dopamine “transportopathy”. *J Clin Invest* 2009;119:1455–1458.
41. Kakkar V, Prins LCB, Kampinga HH. DNAJ proteins and protein aggregation diseases. *Curr Top Med Chem* 2012;12:2479–2490.
42. Kampinga HH, Craig EA. The HSP70 chaperone machinery: J proteins as drivers of functional specificity. *Nat Rev Mol Cell Biol* 2010;11:579–592.
43. Koutras C, Braun JEA. J protein mutations and resulting proteostasis collapse. *Front Cell Neurosci* 2014;8:191.
44. Anikster Y, Haack TB, Vilboux T, et al. Biallelic mutations in DNAJC12 cause hyperphenylalaninemia, dystonia, and intellectual disability. *Am J Hum Genet* 2017;100:257–266.

## Supporting Data

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.

SGML and CITI Use Only  
DO NOT PRINT

Author Roles

(1) Research Project: A. Conception and Design; B. Acquisition of Data; C. Analysis and Interpretation of Data;  
(2) Manuscript: A. Writing of the First Draft, B. Review and Critique; (3) Other: A. Statistical Analysis (for Figure );  
B. Drafting of Figures; C. CSF Neurotransmitter Analysis and Interpretation.

J.N.: 1A, 1B, 1C, 2A, 3A

E.C.-S.: 1A, 1B, 1C, 2A, 3B

L.A.: 1B, 1C, 2A, 3B

P.T.: 1B, 1C

K.M.: 1B, 1C

S.S.: 1B, 1C

K.M.G.: 1B

S.J.R.H.: 3C

S.P.: 3C

L.B.: 1B, 1C

B.C.: 1B, 1C

J.C.: 1B, 1C

K.R.: 1B, 1C

H.C.: 1B, 1C

S.J.: 1B, 1C

R.J.: 1B, 1C

D.H.: 1B, 1C

À.G.-C.: 1B, 1C

D.G.: 1B, 1C

F.L.R.: 1B, 1C

B.P.-D.: 1B, 1C

C.D.G.: 1B, 1C

T.S.P.: 1B, 1C

E.M.: 1A, 1B, 1C, 2A

M.A.K.: 1A, 1B, 1C, 2A

Financial Disclosures

Nothing to report.