**Immunomonitoring of renal transplant
recipients in the early post-transplant
period by analysis of cytokine gene
expression in peripheral blood
mononuclear cells**

M. D. Thesis

Paul John Gibbs MBBS FRCS

ProQuest Number: U644224

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.

ProQuest U644224

Published by ProQuest LLC(2016). Copyright of the Dissertation is held by the Author.

All rights reserved.

This work is protected against unauthorized copying under Title 17, United States Code.
Microform Edition © ProQuest LLC.

ProQuest LLC
789 East Eisenhower Parkway
P.O. Box 1346
Ann Arbor, MI 48106-1346

Abstract

The study was performed to investigate whether monitoring of sequential changes in peripheral blood mononuclear cell (PBMC) cytokine gene expression is predictive of clinical events following renal transplantation, as suggested by earlier studies in our laboratory, using non-quantitative PCR.

RNA was extracted from PBMCs from 45 successive renal transplant recipients (including 15 rejectors) and 13 successive living donors pre-operatively and then daily for 14 and 7 days respectively. RNA was reverse transcribed to cDNA and interleukin (IL)-4, IL-10, transforming growth factor (TGF) β 1, tumour necrosis factor (TNF) α , β -actin, β 2-microglobulin and transferrin receptor gene expression measurement was performed using quantitative real time TaqMan PCR technology.

The results of earlier work in our laboratory using RT-PCR ELISA technology to monitor cytokine gene expression were, in part, confirmed and also extended. β -actin, β 2-microglobulin and transferrin receptor gene expression was found to be influenced by both surgery and acute rejection and so were inappropriate for use as housekeeping genes in the transplant setting. Statistically significant changes in gene expression of all four cytokines were seen in the renal transplant recipients, but only for IL-10 in the living donors. Increased gene expression of IL-10 and TGF- β 1 was observed in both rejecting and non-rejecting recipients. However, changes in gene expression patterns detected for IL-4 and TNF- α were markedly different in rejecting and

non-rejecting recipients. These changes in IL-4 and TNF- α gene expression preceded the clinical detection of acute rejection by more than 24 hours.

This study has demonstrated statistically significant changes in gene expression of IL-4 and TNF- α , which distinguish rejectors from non-rejectors following renal transplantation. While these changes may not reflect intragraft events, they were detectable prior to clinical and biochemical indications of acute rejection and therefore, may have a potential role in immunomonitoring of renal transplant recipients.

Contents

Title Page

Abstract

Table of Contents

List of Figures

List of Tables

Abbreviations

Acknowledgements

1.0 Introduction

1.1 The History of Renal Transplantation

1.1.1 Early Experience with Renal Transplantation

1.1.2 The Introduction of Cross-Matching

1.1.3 The Introduction of Tissue Typing

1.1.4 Advances in Immunosuppression

1.2 Major Histocompatibility Molecules and Antigen Presentation

1.2.1 The Distribution and Role of the Major Histocompatibility Molecules

1.2.2 Structure of the Major Histocompatibility Molecules

1.2.3 Activation of the Immune System Following Transplantation and Antigen Presentation

1.3 The T Cell Cascade and Acute Allograft Rejection

1.3.1 T Cell Activation

- 1.3.2 Costimulation
- 1.4 The Role of Cytokines in Acute Allograft Rejection**
 - 1.3.3 The Th1/Th2 Paradigm
 - 1.3.4 Th1 Cytokines and Acute Rejection
 - 1.3.5 Th1 and Th2 Cytokines and Acute Rejection
 - 1.3.6 Th2 Cytokines and Acute Rejection
 - 1.3.7 Th2 Cytokines and Acceptance
 - 1.3.8 Cytokine Networks and Redundancy
 - 1.3.9 Non-cytokine Influences on the Immune Response
 - 1.3.10 Why Might The Results Be Conflicting?
 - 1.3.11 Cytokine Gene Polymorphisms
 - 1.3.12 Influence of Cytokine Gene Polymorphisms on Rejection
 - 1.3.13 Why Might There Be This Disparity?
- 1.5 Immunological Monitoring**
 - 1.5.1 The Measurement of Constituents of the Urine for Immunological Monitoring
 - 1.5.2 The Measurement of Serum Proteins and Soluble Molecules for Immunological Monitoring
 - 1.5.3 The Measurement of Serum Cytokines for Immunological Monitoring
 - 1.5.4 The Measurement of Peripheral Blood Cell Surface Markers for Immunological Monitoring
- 1.6 Previous Work from Our Laboratory in the Field of Immunological Monitoring**
- 1.7 TaqMan Real Time PCR Detection**
- 1.8 Aims of the Study**

2 Patients, Method and Materials

2.1 Patient Recruitment and Rejection Diagnosis

2.2 Cell Separation

2.2.1 Blood Collection

2.2.2 Cell Separation

2.3 Total RNA Extraction, Quantitation and Standardisation

2.3.1 Total RNA Extraction

2.3.2 Total RNA Quantitation

2.3.3 Total RNA Standardisation

2.4 First-strand cDNA Synthesis

2.4.1 First-strand cDNA Synthesis

2.5 TaqMan Cytokine Gene Expression Analysis

2.5.1 TaqMan Cytokine Gene Expression Analysis

3 The Substantiation of the Results of the RT-PCR ELISA Gene Expression Work by Using the TaqMan System

3.1 Comparison of commercial primers/probes with designed primers/probes

3.2 Re-analysis of the Existing cDNA Samples from the RT-PCR ELISA Work Using the TaqMan System

3.3 Discussion

4 The Investigation of the Use of a House Keeping Gene for PCR Standardisation

4.1 Introduction

4.1.1 Why use a house keeping gene?

4.1.2 Which house keeping gene to use?

4.2 Investigation of Single PCR Tube Multiple Gene Expression Analysis Using Different TaqMan Reporter Dyes

4.3 Results

4.3.1 Individual variation in Expression of the Three House Keeping Genes

4.3.2 Correlation Between the Expression Data of the Three House Keeping Genes in both Living Donors and Transplant Recipients

4.3.3 Gene Expression Data for Each House Keeping Gene in Each of the Three Cohorts of Patients

4.4 Discussion

5 Results of the Cytokine Gene Expression Work

5.1 Patient Demographics

5.2 Results of Cytokine Gene Expression Measurement

6 Discussion

6.1 Interleukin-4

6.2 Interleukin-10

6.3 Transforming Growth Factor- β 1

6.3 Tumour Necrosis Factor- α

7 Summary and Conclusions

Appendices

Appendix I

Appendix II

Appendix III

Appendix IV

Appendix V

Appendix VI

Appendix VII

References

List of Figures

Fig. 1.1	Diagram illustrating the mechanism of drug actions
Fig. 1.2a	Diagram illustrating direct antigen presentation
Fig. 1.2b	Diagram illustrating indirect antigen presentation
Fig. 1.4	Diagram illustrating the Th1/Th2 Paradigm
Fig. 1.7	Scattergram showing the correlation between TaqMan duplicates
Fig. 2.5a	Diagram illustrating TaqMan initiation of polymerisation
Fig. 2.5b	Diagram illustrating strand displacement of the TaqMan
Fig. 2.5c	Diagram illustrating cleavage of the TaqMan probe
Fig. 2.5d	Diagram illustrating completion of polymerisation
Fig. 3.1a	Graph showing a TaqMan result of a non-rejector using in house designed IL-4 primers and probe
Fig. 3.1b	Graph showing a TaqMan result of a non-rejector using commercially designed IL-4 primers and probe
Fig. 3.1c	Graph showing a TaqMan result of a non-rejector using in house designed IL-10 primers and probe
Fig. 3.1d	Graph showing a TaqMan result of a non-rejector using commercially designed IL-10 primers and probe
Fig. 3.1e	Graph showing a TaqMan result of a rejector using in house designed IL-4 primers and probe
Fig. 3.1f	Graph showing a TaqMan result of a rejector using commercially designed IL-4 primers and probe
Fig. 3.1g	Graph showing a TaqMan result of a rejector using in house designed IL-10 primers and probe
Fig. 3.1h	Graph showing a TaqMan result of a rejector using commercially designed IL-10 primers and probe
Fig. 3.1i	Scattergram showing the correlation between in house and commercially designed IL-4 and IL-10 primers and probes
Fig. 3.2a	Graph showing the similarity between TaqMan results and RT- PCR ELISA results

- Fig. 3.2b Box and whisker plot showing IL-4 gene expression data from the non-rejector cohort in the RT-PCR ELISA study
- Fig. 3.2c Box and whisker plot showing IL-4 gene expression data from the non-rejector cohort in the RT-PCR ELISA study but analysed using TaqMan technology
- Fig. 3.2d Box and whisker plot showing IL-10 gene expression data from the non-rejector cohort in the RT-PCR ELISA study
- Fig. 3.2e Box and whisker plot showing IL-10 gene expression data from the non-rejector cohort in the RT-PCR ELISA study but analysed using TaqMan technology
- Fig. 3.2f Box and whisker plot showing IL-4 gene expression data from the rejector cohort in the RT-PCR ELISA study
- Fig. 3.2g Box and whisker plot showing IL-4 gene expression data from the rejector cohort in the RT-PCR ELISA study but analysed using TaqMan technology
- Fig. 3.2h Box and whisker plot showing IL-10 gene expression data from the rejector cohort in the RT-PCR ELISA study
- Fig. 3.2i Box and whisker plot showing IL-10 gene expression data from the rejector cohort in the RT-PCR ELISA study but analysed using TaqMan technology
- Fig. 4.2a Graph showing a TaqMan result of a non-rejector using in house designed IL-4 primers and probe
- Fig. 4.2b Graph showing a TaqMan result of a non-rejector using in house designed IL-4 primers and probe with the commercially produced β -actin primers and probe in the same tube
- Fig. 4.2c Graph showing a TaqMan result of a non-rejector using in house designed IL-10 primers and probe
- Fig. 4.2d Graph showing a TaqMan result of a non-rejector using in house designed IL-10 primers and probe with the commercially produced β -actin primers and probe in the same tube
- Fig. 4.2e Graph showing a TaqMan result of a non-rejector using the commercially produced β -actin primers and probe

- Fig. 4.2f Graph showing a TaqMan result of a non-rejector using the commercially produced β -actin primers and probe with the in house designed IL-10 primers and probe in the same tube
- Fig. 4.3.1a Graph showing the gene expression of all three potential house keeping genes in a non-rejecting transplant recipient
- Fig. 4.3.1b Graph showing the gene expression of all three potential house keeping genes in a rejecting transplant recipient
- Fig. 4.3.1c Graph showing the gene expression of all three potential house keeping genes in a living donor
- Fig. 4.3.2a Graph demonstrating the correlation between the gene expression data of β -2- and β -actin for the living donors
- Fig. 4.3.2b Graph demonstrating the correlation between the gene expression data of β -2- and transferrin receptor for the living donors
- Fig. 4.3.2c Graph demonstrating the correlation between the gene expression data of transferrin receptor and β -actin for the living donors
- Fig. 4.3.2d Graph demonstrating the correlation between the gene expression data of β -2- and β -actin for all the transplant recipients
- Fig. 4.3.2e Graph demonstrating the correlation between the gene expression data of β -2-microglobulin and transferrin receptor for all the transplant recipients
- Fig. 4.3.2f Graph demonstrating the correlation between the gene expression data of transferrin receptor and β -actin for all the transplant recipients
- Fig. 4.3.3a Box and whisker plot of the gene expression data of β -actin in the living donors
- Fig. 4.3.3b Box and whisker plot of the gene expression data of β -2-microglobulin in the living donors
- Fig. 4.3.3c Box and whisker plot of the gene expression data of transferrin receptor in the living donors

- Fig. 4.3.3d Box and whisker plot of the gene expression data of β -actin in the non-rejecting transplant recipients
- Fig.4.3.3e Box and whisker plot of the gene expression data of β 2-microglobulin in the non-rejecting transplant recipients
- Fig.4.3.3f Box and whisker plot of the gene expression data of transferrin receptor in the non-rejecting transplant recipients
- Fig. 4.3.3g Box and whisker plot of the gene expression data of β -actin in the rejecting transplant recipients
- Fig.4.3.3h Box and whisker plot of the gene expression data of β 2-microglobulin in the rejecting transplant recipients
- Fig. 4.3.3i Box and whisker plot of the gene expression data of transferrin receptor in the rejecting transplant recipients
- Fig. 4.3.3j Box and whisker plot of the gene expression data of β -actin in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis
- Fig.4.3.3k Box and whisker plot of the gene expression data of β 2-microglobulin in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis
- Fig. 4.3.3l Box and whisker plot of the gene expression data of transferrin receptor in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis
- Fig. 5.2a Box and whisker plot of the gene expression data of IL-4 in the living donors
- Fig. 5.2b Box and whisker plot of the gene expression data of IL-4 in the non-rejecting transplant recipients
- Fig. 5.2c Box and whisker plot of the gene expression data of IL-4 in the rejecting transplant recipients
- Fig. 5.2d Box and whisker plot of the gene expression data of IL-4 in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis
- Fig. 5.2e Box and whisker plot of the gene expression data of IL-10 in the living donors

- Fig. 5.2f Box and whisker plot of the gene expression data of IL-10 in the non-rejecting transplant recipients
- Fig. 5.2g Box and whisker plot of the gene expression data of IL-10 in the rejecting transplant recipients
- Fig. 5.2h Box and whisker plot of the gene expression data of IL-10 in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis
- Fig. 5.2i Box and whisker plot of the gene expression data of TGF- β 1 in the living donors
- Fig. 5.2j Box and whisker plot of the gene expression data of TGF- β 1 in the non-rejecting transplant recipients
- Fig. 5.2k Box and whisker plot of the gene expression data of TGF- β 1 in the rejecting transplant recipients
- Fig. 5.2l Box and whisker plot of the gene expression data of TGF- β 1 in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis
- Fig. 5.2m Box and whisker plot of the gene expression data of TNF- α in the living donors
- Fig. 5.2n Box and whisker plot of the gene expression data of TNF- α in the non-rejecting transplant recipients
- Fig. 5.2o Box and whisker plot of the gene expression data of TNF- α in the rejecting transplant recipients
- Fig. 5.2p Box and whisker plot of the gene expression data of TNF- α in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis
- Fig. VIIa Illustration showing a the typical appearance of a TaqMan output data graph

List of Tables

Table 1.4	Cytokine polymorphisms – position and phenotype
Table 2.5a	Standard TaqMan PCR conditions
Table 2.5b	Components of the TaqMan master mixes
Table 3.1a	Time points used in RT-PCR ELISA study
Table 3.1b	Correlation between in house and commercially designed IL-4 and IL-10 primers and probes
Table 3.2a	Results of the statistical analysis of IL-4 and IL-10 gene expression data from the non-rejector cohort from the RT-PCR ELISA study and after re-analysis with TaqMan technology
Table 3.2b	Results of the statistical analysis of IL-4 and IL-10 gene expression data from the rejector cohort from the RT-PCR ELISA study and after re-analysis with TaqMan technology
Table 4.3.2a	Results of the correlation between the expression of the three potential house keeping genes in the living donors
Table 4.3.2b	Results of the correlation between the expression of the three potential house keeping genes in the transplant recipients
Table 4.3.3a	Results of stepwise analysis of the gene expression data of the three potential house keeping genes in the living donors
Table 4.3.3b	Results of comparison to baseline analysis of the gene expression data of the three potential house keeping genes in the living donors
Table 4.3.3c	Results of stepwise analysis of the gene expression data of the three potential house keeping genes in the non-rejecting transplant recipients
Table 4.3.3d	Results of comparison to baseline analysis of the gene expression data of the three potential house keeping genes in the non-rejecting transplant recipients
Table 4.3.3e	Results of stepwise analysis of the gene expression data of the three potential house keeping genes in the rejecting transplant recipients

Table 4.3.3f	Results of comparison to baseline analysis of the gene expression data of the three potential house keeping genes in the rejecting transplant recipients
Table 4.3.3g	Results of stepwise analysis of the gene expression data of the three potential house keeping genes in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis
Table 4.3.3h	Results of comparison to baseline analysis of the gene expression data of the three potential house keeping genes in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis
Table 5.1a	Cause of the end-stage renal failure in all 30 non-rejecting and 15 rejecting renal transplant recipients
Table 5.1b	Demographics of the rejector and non-rejector recipient groups
Table 5.1c	Distribution of donor/recipient CMV status and donor/recipient sex for both sets of renal transplant recipients
Table 5.1d	Distribution of immunosuppressives regimes used in the two groups of renal transplant recipients
Table 5.1e	Type, mean age and sex of the living donors
Table 5.2a	Results of stepwise analysis of the cytokine gene expression data of the living donors
Table 5.2b	Results of the comparison to baseline analysis of the cytokine gene expression data of the living donors
Table 5.2c	Results of stepwise analysis of the cytokine gene expression data of the non-rejecting transplant recipients
Table 5.2d	Results of the comparison to baseline analysis of the cytokine gene expression data of the non-rejecting transplant recipients
Table 5.2e	Results of stepwise analysis of the cytokine gene expression data of the rejecting transplant recipients
Table 5.2f	Results of the comparison to baseline analysis of the cytokine gene expression data of the rejecting transplant recipients

Table 5.2g	Results of stepwise analysis of the cytokine gene expression data of the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis
Table 5.2h	Results of the comparison to baseline analysis of the cytokine gene expression data of the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis

Abbreviations

IgG	Immunoglobulin G
IgM	Immunoglobulin M
MHC	Major Histocompatibility Complex
HLA	Human Leucocyte Antigen
CyA	Cyclosporin A
FK506	Tacrolimus
TCR	T Cell Receptor
IL-	Interleukin-
IL-2R	Interleukin-2 Receptor
Ca ²⁺	Free Calcium
IFN	Interferon
CD	Cluster of Differentiation
APC	Antigen Presenting Cell
kB	kiloBase
kDa	kiloDalton
IP3	1,4,5-triphosphate
PKC	Protein Kinase C
CD40L	CD40 Ligand
Th	T Helper Cell
FNA	Fine Needle Aspiration
ELISA	Enzyme-Linked Immunoabsorption Assay
RT-PCR	Reverse Transcription Polymerase Chain Reaction
mRNA	messenger Ribonucleic Acid

TNF- α	Tumour Necrosis Factor- α
RNA	Ribonucleic Acid
CTL	Cytotoxic T Lymphocyte
PBMC	Peripheral Blood Mononuclear Cell
SNP	Single Nucleotide Polymorphism
TGF- β 1	Transforming Growth Factor- β 1
CMV	Cytomegalovirus
IL-1ra	Interleukin-1 receptor antagonist
PCR	Polymerase Chain Reaction
sIL-2R	soluble Interleukin-2 Receptor
ICAM-1	Intercellular Adhesion Molecule-1
VCAM-1	Vascular Adhesion Molecule-1
ELAM-1	Endothelial Adhesion Molecule-1
CRPu	urinary C-Reactive Protein
CRPs	serum C-Reactive Protein
β 2M	β 2-microglobulin
ATN	Acute Tubulonecrosis
SAA	Serum Amyloid A
sHLA	soluble Human Leucocyte Antigen
TNF- β	Tumour Necrosis Factor- β
CIM	Cytoimmunomonitoring
FCXM	Flow Cytometric Cross Match
cDNA	complimentary Deoxyribonucleic Acid
Taq	Thermus aquaticus DNA polymerase

C _T	Threshold Cycle (cycle at which PCR growth becomes exponential)
PBS	Phosphate Buffered Saline
DEPC	Diethyl Pyrocarbonate
UHQ	Ultra High Quality
ddH ₂ O	double distilled Water
FAM	6-carboxyfluorescein
TAMRA	6-carboxy-tetramethylrhodamine
CCD	Charge Coupled Device
VIC	reporter dye on commercial house keeping gene probes
T _m	Melting Temperature
PT	Pre-transplant
NS	Non-significant
GAPDH	glyceraldehyde-3-phosphate dehydrogenase
TfR	Transferrin Receptor
huBAc	human β -actin
B2MAV	β 2-microglobulin Average
BAAV	β -actin Average
TFRAV	Transferrin Receptor Average
BR	Before Rejection
PR	Post-rejection
PTBS	Pre-transplant Baseline
OKT3	Murine Anti-CD3 Monoclonal Antibody
MMF	Mycophenolate Mofetil

Acknowledgements

I would like to thank all the staff in the Histocompatibility and Immunology Laboratory at Southampton General Hospital for all their help and support.

I would also like to thank my two supervisors, Dr. W. M. Howell and Mr. S. A. Sadek, for their encouragement, constructive criticism and time.

I am very grateful to the patients and staff of the Wessex Renal and Transplant Unit. The patients for being so willing to take part in this study and the staff for helping out however and whenever they could.

My thanks also go out to the Wessex Renal Research Fund. They have very kindly provided the financial backing for this work.

Finally, I would like to thank Sarah and Mollie for making it all worth while.

1.0: Introduction

This introduction will give an overview of the cellular processes that occur after the introduction of a renal allograft into a new human host. In particular, antigen presentation, the T-cell cascade and acute allograft rejection will be considered. The role of cytokines in these events will be discussed, and a critical evaluation of the potential importance of cytokine gene promoter polymorphisms will be conducted.

The role of immunomonitoring in the management of transplant recipients will be reviewed and its potential benefits discussed. Previous work carried out by this laboratory on sequential T cell monitoring in renal allograft recipients will be examined, and the weaknesses considered.

Finally, the aims of the study will be presented.

1.1 The History of Renal Transplantation

Renal transplantation is now a routine surgical procedure in the United Kingdom. In 1997 nearly 1700 transplants were performed (The Report of the Working Party to Review Organ Transplantation, 1999) with a one year graft survival of more than 80% (Towards standards for organ and tissue transplantation in the United Kingdom, 1998). This is a tremendous advance on the early days of human renal transplantation in the 1930's and 1940's when patients did not survive more than a few days at best (Hume 1979).

1.1.1 Early Experience With Human Renal Transplantation

Voronoy performed the first human to human renal transplant in 1936, using a cadaveric kidney, but without immunosuppression. This, and others shortly afterwards, were unsuccessful and the patients died (Hume 1979).

In the early 1950's, work on renal transplantation was concentrated in Boston and Paris. The Boston group performed nine renal transplants between 1951 and 1953. Four of the nine showed measurable function, but this was only sufficient to maintain life in three. The French groups fared no better. They performed eight transplants in 1951. Only some produced urine, but in no case was this sufficient to improve the patient's uraemic state. Both groups were using immunosuppressive agents for the first time in an attempt to prevent rejection. This was either adrenocorticotrophic hormone or cortisone, but neither showed any definite effect (Hume 1979, Groth 1972).

In 1954, in Boston, a team headed by Dr. Joseph Murray performed the first successful human to human renal transplant. This was a living related transplant between identical twins. The graft functioned for eight years before the patient's death from a myocardial infarction (Murray 1955). This was a huge step forward, and one recognised by the joint award of the Nobel Prize for Medicine in 1990. Unfortunately, this progress would not benefit the vast majority of patients with end stage renal failure.

The next milestone was the introduction of first, azathioprine by Sir Roy Calne and then prednisolone by Zukoski during the 1960's. These drugs were able to suppress the immune system in a transplant recipient and, therefore, increase the chances of graft survival. With the two drugs used in combination the 1 year graft survival for kidney transplants improved to between 50 and 60% (Groth 1972).

1.1.2 The Introduction of Crossmatching

By the mid 1960's two forms of renal allograft rejection were recognised. Firstly, hyperacute rejection, which resulted in loss of the graft within the first few minutes after re-perfusion of the transplanted kidney. Secondly, rejection occurring after 7-10 days, termed acute rejection. Research into the former process identified the cause as pre-formed antibodies. These were detected when donor cells were incubated with serum from the recipient. From this discovery Paul Terasaki developed the lymphocytotoxic cross-match using small volumes of serum and a low number of cells (Talbot 1993). This assay

identified both IgG and IgM antibodies directed against recipient cells. As it became clear that IgM antibodies had little relevance to organ rejection, the addition of dithiothreitol, which deactivates IgM antibodies, became widely used. The separation of T and B cells led to the generally accepted principle in renal transplantation that a positive cross-match due to IgG to donor T cells was a contraindication to transplantation.

The disadvantages of cytotoxic cross-matching was that IgG had to be at a sufficient concentration to activate complement. In addition, the definition of a positive assay was operator dependent. These problems were overcome with the introduction of flow cytometry for the purposes of cross-matching. This allowed individual cell analysis rather than a system studying a whole cell population. Garavoy was the first to use flow cytometry for cross-matching purposes in renal transplant recipients (Talbot 1993).

1.1.3 The Introduction of Tissue Typing

Another advance that occurred in the 1960's was the commencement of tissue typing. It had long been recognised that it was important for donor and recipient to be ABO blood group compatible, as this resulted in better graft survival (Hume 1979). During the 1950's work was underway to identify similar serologically determined antigens on leucocytes. The first human major histocompatibility complex (MHC) antigen was identified in 1958 (Dausset 1958). These antigens are now referred to as the Human Leucocyte Antigens (HLA). Work continued during the 1960's to identify more of these

antigens and the polymorphic nature of the genes encoding them. By 1970 they had been codified into three series, namely A, B, and C and collectively known as the MHC class I antigens (Kissmeyer-nielson 1968, Sandberg 1970). In 1973 Van Leeuwen et al identified the first of the class II antigens (Van Leeuwen 1973). These are also divided into three series, designated as DR, DP, and DQ.

This knowledge gave rise to the clinical and laboratory practice of tissue typing. This affords a carefully designated level of HLA matching between recipient and donor. It was postulated that matching would reduce the severity and amount of rejection and therefore improve graft survival. This has been confirmed in renal transplantation with better matches being associated with lower rejection rates and increased graft survival (Opelz 1991). The possible reasons for this will be discussed later (section 1.2).

1.1.4 Advances in Immunosuppression

During the early 1970's, several commercial companies developed polyclonal antibody treatments to improve graft survival. Serum was extracted from horses and rabbits after the introduction of human antigens. The animal developed antibodies directed against human lymphocytes and thymocytes. When used with azathioprine and prednisolone as induction therapy, these drugs were shown to significantly reduce acute rejection episodes and improve both patient and graft survival by 10-15% at one year when compared to azathioprine and prednisolone alone. These preparations were

also shown to be better at treating acute allograft rejection, with fewer relapses and better survival (Cosimi 1981, Groth 1981).

The discovery by Borel et al of the drug cyclosporin A in 1976 was to prove the next major advance in transplantation. Multicentre trials in both Europe and Canada showed significantly better graft survival in renal transplant recipients receiving cyclosporin (with or without prednisolone) compared with dual therapy using azathioprine and prednisolone. The European study reported 1-year graft survival rates of 72% in the group receiving cyclosporin monotherapy compared with 52% in the control arm. While in the Canadian study, graft survival was 76% at 1-year for those receiving cyclosporin and prednisolone compared with 70% for those taking azathioprine and prednisolone (European Multicentre Trial Group 1983, Canadian Multicentre Transplant Study Group 1983).

In the last decade there have been many further advances with immunosuppressive therapy. This has been both in the number of agents available and in the formulation of existing products. Several of these have been shown to significantly reduce the rate of acute rejection and the need for biological agents to treat steroid resistant and severe acute rejection by targeting different steps in cell activation and proliferation (see figure 1.1). Unfortunately, none have had the profound affect on survival seen with the introduction of cyclosporin, but these drug therapies have increased the treatment options available (European Mycophenolate Mofetil Co-operation

Study Group 1995, Jordan 1997, Groth 1999, Kahan 1998, Nashan 1997, Pollard 1999).

The transplant community has come a very long way from the days of the early transplant pioneers. The ability exists to prevent acute rejection in the majority of renal transplant recipients. However, 30-40% of renal transplant recipients still experiences some degree of biopsy proven rejection (Pollard 1999, Groth 1999). This may lead to permanent damage to the transplanted organ, and decrease in long-term graft survival (Heaf 1998 and Ashraf 1998).

1.1.5 Future Challenges

With the demand for a renal transplant out stripping the supply (The Report of the Working Party to Review Organ Transplantation 1999), there is an ever-increasing need to optimise both the short and long-term graft survivals. Further advances in drug therapy maybe one answer. Others may include identification of genetic sub-types of recipients more or less likely to reject a transplanted organ. Monitoring of the immune system with appropriate and timely intervention should the evidence point to the initiation of a detrimental immune response may also be possible. These latter two points will be expanded upon in subsequent sections.

1.2 Major Histocompatibility Molecules & Antigen Presentation

The human immune system relies on the ability to differentiate between 'self' and both 'non-self' and 'modified-self' in order to survive. The molecular basis for this recognition resides with the major histocompatibility complex (MHC) molecules. There are two distinct types, namely, human leucocyte antigen (HLA) class I (A, B, and C) and class II (DR, DP, and DQ).

1.2.1 The Distribution and Role of the MHC Molecules

Class I antigens have an almost ubiquitous tissue distribution being present on almost all nucleated cells and platelets with the important exception of syncytiotrophoblast tissue and corneal endothelium (Daar 1984). Class II antigens, on the other hand, have a far more restricted distribution. They are expressed mainly on B lymphocytes, macrophages and monocytes, capillary endothelium and some other epithelia (Daar 1984), but are inducible by interferon (IFN)- γ (Arai 1990). The principal physiologic function of the cell surface histocompatibility molecules is to bind peptide fragments of 'foreign' proteins for presentation to appropriate antigen-specific T cells.

Class I MHC molecules bind to intracellularly synthesised peptides (e.g. viral antigens) and present them to CD8⁺ cytotoxic T lymphocytes. In this interaction, the T cell receptor recognises the MHC-peptide complex and the CD8 molecule binds to the non-polymorphic portion of the Class I molecule.

Since one of the important functions of CD8⁺ cytotoxic T lymphocytes is to eliminate virus-infected cells, and viral antigens can only be recognised when complexed to Class I MHC molecules, it is appropriate that expression of Class I HLA antigens is widespread (Neefjes 1990).

Class II molecules present antigenic peptides to CD4⁺ helper T cells. The nature of the antigens presented in association with Class II molecules differs. In general Class II molecules present exogenous antigens that are first internalised and processed by antigen-presenting cells (APCs) and then the Class II-peptide complex is transported to the cell surface (Neefjes 1990). APCs are highly specialised cells that are either derived from the macrophage/monocyte lineage, B lymphocytes and under certain conditions, endothelial cells (Sundstrom 1995).

1.2.2 Structure of the MHC Molecules

In humans, the genes encoding the MHC molecules are present on the short arm of chromosome 6 at position 6p21.3 encompassing a 3600kB region (Trowsdale 1991). The class I and II regions are well characterised (Koller 1989, Dunham 1989), and include the most polymorphic genes in the human genome. This is reflected in the variation of the amino acids lining the peptide-binding cleft of the expressed molecules. Functionally different HLA alleles can, therefore, present a different profile of processed antigenic peptides to T cells.

Class I antigens consist of a 45kDa α -chain polypeptide, with extracellular and transmembrane regions and a cytoplasmic tail. The extracellular component of the α -chain is non-covalently bound to a 12kDa polypeptide, β 2-microglobulin. The α -chain of each class I molecule is highly polymorphic, and encoded by closely linked genes on chromosome 6 within the MHC region, whereas the β 2-microglobulin is encoded by a non-polymorphic, non-MHC region of chromosome 15 (Bjorkman 1990). The class II molecule consists of two non-covalently bound α and β chain glycoproteins, again with extracellular, transmembrane and intracellular regions. Highly polymorphic A and B genes within the DR, DQ and DP sub-regions of the MHC again, encode each chain (Kappes 1988). Each molecule has been shown to have a groove in it's extracellular portion, which is the site of binding of small peptides for presentation to T cells via the cell surface T cell receptor (TCR) (Bjorkman 1987, Rudensly 1991).

1.2.3 Activation of the Immune System Following Transplantation and Antigen Presentation

The human immune system is activated by the detection of foreign antigens by T lymphocytes. These cells, however, do not have the ability to recognise free antigens, but react to the peptide/MHC complex, which is presented to the T cell receptor. In the case of MHC class I, the antigen is presented to CD8⁺ T cells. For CD4⁺ T cells this occurs when intracellularly processed antigen, in the form of small peptides, is associated with the MHC class II molecules on the surface of professional APCs. Therefore, the function of the MHC molecules is to present peptides to T cells as a surveillance mechanism so that

'foreign' peptides expressed by virally infected cells, tumour cells and, unfortunately, transplanted organs are rapidly identified. HLA class I molecules primarily present peptides from endogenous proteins to CD8⁺ (cytotoxic) T cells, while class II antigens bind peptides derived from exogenous proteins to CD4⁺ (helper) T cells (Neefjes 1990).

Organ transplantation results in the introduction of a large quantity of foreign antigen. These antigens can be presented to recipient CD4⁺ T cells in one of two ways. Firstly, the recipient T cells may interact directly with the APC's derived from the donor. This physiologically unusual stimulation is termed 'direct' presentation. In this pathway, the donor MHC alone or the donor MHC and peptide complex will appear as 'non-self' to the recipient CD4⁺ T cells, therefore activating these cells. In contrast, presentation of processed donor peptide by recipient APC's is called indirect presentation (see figures 1.2a and 1.2b). In other words, the donor MHC acts as the presenting molecule in the direct pathway whereas it may be the source of the peptide presented by self-MHC in the indirect pathway. Historically it has been thought to be the direct response that has been the more important in graft rejection (Lafferty 1976, Pescovitz 1984). However, there is mounting evidence that this is not the case and more emphasis should be placed on the indirect pathway (Gould 1999).

CD8⁺ cells do not require professional APCs as they interact with Class I antigens. In the transplantation setting, these cells may interact directly with donor cells of any type, as they will all express Class I antigens. Therefore,

any HLA class I disparity between the donor and recipient may lead to direct activation of recipient CD8⁺ T cells against donor cells of all types. For CD8⁺ T cells there is no equivalent to indirect antigen presentation.

Fig. 1.2a Direct Antigen Presentation

This is a diagrammatic representation of the process of direct antigen presentation.

Fig. 1.2b Indirect Antigen Presentation

This is a diagrammatic representation of the process of indirect antigen presentation.

1.3 The T Cell Cascade & Acute Allograft Rejection

Allograft rejection can be broadly divided into three categories. Firstly, hyperacute rejection, which is antibody mediated, and occurs within the first few hours following transplantation. These antibodies are pre-formed and usually directed against ABO blood group antigens or HLA antigens as a result of previous sensitisation. This form of rejection is now rarely seen due to modern cross-matching techniques. Secondly, there is acute rejection. This can have both cellular and humoral components. It usually occurs within the first two to three weeks following transplantation. This form of rejection is the focus of this work, and will be expanded on in this and other sections. Finally, there is chronic rejection, which can be detected in some grafts after only a few months, and is probably best termed chronic allograft nephropathy. The aetiology of this form of rejection is multifactorial, with chronic immune damage being one possible element contributing to the pathology.

Allograft rejection results from the co-ordinated activation of both alloreactive T cells and APCs. Following the release of cytokines and cell-cell interactions, several subsets of lymphocytes and other inflammatory leucocytes are recruited in the anti-allograft response. These include CD4⁺ T cells (primarily of helper phenotype), CD8⁺ T cells (primarily of cytotoxic phenotype) and antibody forming B cells. The transplanted organ sustains damage via the direct effect of graft-destructive T cells and via complement dependent and/or antibody dependent cell-mediated cytotoxicity (Strom 1975).

1.3.1 T Cell Activation

T cell activation commences with the immune system recognising 'non-self' or 'modified-self'. This involves the interaction of the TCR with a processed fragment of foreign peptide located within the groove of the MHC molecule (Unanue 1989, Germain 1994). This interaction can be between either a donor APC (direct recognition) or a recipient APC (indirect recognition) as discussed above. The relative roles of these two processes within the domain of renal transplantation has yet to be fully resolved (Gould 1999). Alternatively, CD8⁺ T cells may be activated by the direct interaction with donor MHC molecules. The TCR/CD3 complex, in association with either the CD4 or CD8 proteins, bind to Class I MHC molecules in the case of CD8 and Class II MHC for CD4 (Miceli 1991). Antigenic recognition stimulates a redistribution of cell-surface proteins and co-clustering of the TCR/CD3 complex with the CD4 or CD8 antigens. This multimeric complex includes other signalling molecules and functions as a unit in initiating T cell activation (Suthanthiran 1990, Brown 1989, Beyers 1992).

The TCR/CD3 complex is associated with two intracellular tyrosine kinases, while the cytoplasmic domains of CD4 and CD8 are linked with another (Klausner 1991). Following antigenic recognition of the TCR/CD3 complex tyrosine phosphorylation occurs. This triggers a series of events resulting in the release of the second messengers 1,4,5-triphosphate (IP₃) and diacylglycerol. IP₃ mobilises ionised calcium while diacylglycerol activates protein kinase C (PKC). The increase in concentration of intracellular free

Ca^{2+} and sustained PKC activation function synergistically in promoting the expression of several nuclear regulatory proteins and transcription of genes important for T cell growth and differentiation (Nishizuka 1992).

1.3.2 Costimulation

Stimulation of the T cell via the TCR/CD3 complex alone results in anergy. Full T cell activation requires co-stimulatory signals resulting from cell-cell interaction between the T cell and APC (Jenkins 1987, Schwartz 1996, Suthanthiran 1993). These include CD40L and CD28 on the T cell and CD40 and B7-1 and B7-2 respectively on the APC (Laman 1996, June 1990, Thompson 1989).

1.4 The Role of Cytokines in Acute Allograft Rejection

Cytokines are potent, pleiotrophic, non-antigen-binding polypeptides secreted by cells of the immune system. They are produced in response to a variety of stimuli including regulation of the immune response to foreign antigens and invading organisms. They are involved in cell recruitment, activation, differentiation and proliferation. They act on their target cell via specific surface receptors primarily through autocrine and paracrine stimulation (Paul 1989, Balkwill 1989, Arai 1990).

Acute allograft rejection is a T cell mediated response. It results in the specific recognition and destruction of an alloantigen-expressing donor organ. Secretion of cytokines by donor-specific, activated T cells is important in augmenting inflammation (Kirk 1995). Therefore, analysis of cytokine production during acute rejection episodes may enhance our understanding of the rejection process and the mechanisms associated with tolerance. The aim of such studies is to understand the interactions and role of cytokines in acute allograft rejection. This may lead to the ability to predict both graft acceptance and rejection early following allograft transplantation and, as a result, manipulate immunosuppressive therapy accordingly.

1.4.1 The Th1/Th2 Paradigm

In 1986 Mosmann et al (Mosmann 1986) described two functionally distinct CD4⁺ T helper (Th) cell subsets, characterised by divergent cytokine

elaboration patterns by in vitro analysis of murine Th cell clones. These findings were later corroborated in a human in vitro study (Del Prete 1991). These findings resulted in development of the so-called 'Th1/Th2 paradigm'.

This paradigm implies that naive Th cells, following activation, can differentiate into two distinct phenotypes producing different cytokine profiles. Th1 cells produce interleukin (IL)-2, interferon (IFN)- γ , and lymphotoxin, and promote cellular immune responses that induce potent cell-mediated immunity and allograft rejection. Conversely, Th 2 cells producing IL-4, IL-5, IL-6, and IL-10, are associated with humoral immunity and prolonged allograft survival and acquisition of donor specific tolerance (Mosmann 1989). As Th2 cells are antagonistic to Th1 development and function, a popular hypothesis states that preferential induction of allograft specific Th2 would down-regulate the Th1 driven rejection process, thereby promoting allograft acceptance (Piccotti 1997). See figure 1.4.

Figure 1.4. Diagram illustrating the Th1/Th2 Paradigm

This figure is a diagrammatic interpretation of the Th1/Th2 paradigm showing the differentiation of a naive CD4 T cell into either a Th1 cell or Th2 cell with their respective cytokine profiles and the negative inhibition these exhibit on the other Th subset.

Animal transplant models in the early 1990's seemed to confirm the Th1/Th2 paradigm, with Th1 cytokines being associated with rejection (Dallman 1991, Thai 1995, Wu 1992) and the Th2 cytokines with allograft acceptance (Bugeon 1992, Chen 1995, Sayegh 1995, Takuchi 1992). However, work published in 1995 (Steiger 1995) and in 1996 (Nickerson 1996, Konieczny 1996) suggested that the paradigm might be an over simplification of a complex process. This work showed the ability of IL-2 knockout mice to reject islet cell allografts (Steiger 1995) and IFN- γ knockout mice to reject skin and cardiac transplants (Konieczny 1996). Conversely, IL-4 knockout mice exhibit prolonged islet allograft acceptance (Nickerson 1996). It, therefore, became increasingly clear that the data derived from strictly controlled in vitro laboratory experiments differed from the findings of animal transplant models and in the clinical setting (Zhai 1999).

To further understand the role cytokines play in the process of acute allograft rejection, there are many studies looking at intragraft cytokine expression from both core biopsies and fine needle aspiration (FNA) samples. Most of these have studied the pattern of cytokine expression at the time of rejection and compared it to the pattern in quiescent grafts. Below is a summary of the work in this field, concentrating mainly on the work studying acute renal allograft rejection. The work published on cytokine expression outside of the allograft will be examined in the section on immunomonitoring (section 1.5).

1.4.2 Th1 Cytokines and Acute Rejection

The Th1/Th2 paradigm infers that acute allograft rejection is associated with the presence of Th1 and absence of Th2 cytokines. Unfortunately, despite early evidence to support this, there is increasing data that this is an oversimplification of what happens in clinical transplantation (Baan 1998, Piccotti 1997, Gimsa 1997).

Simpson and colleagues, in 1989, were one of the first groups to publish data supporting the role of IL-2 in acute renal allograft rejection. Using an ELISA system they demonstrated a rise in both urine and plasma levels of IL-2 preceded clinically apparent rejection. This rise was not seen in patients with stable renal function and fell following successful treatment (Simpson 1985). A group in Oxford was the first to show evidence for a predominance of Th1 cytokines in acute allograft rejection from within the graft. Using a reverse transcription polymerase chain reaction (RT-PCR) system, they reported IL-2 gene transcription, in sequential FNA samples taken from 7 human renal allografts, preceded clinical rejection, but was not detected in the absence of rejection (Dallman 1992).

Nast and colleagues in Los Angeles, who also looked at intragraft mRNA expression of FNAs in human renal transplant recipients, supported this finding. They detected the presence of another Th1 cytokine (IFN- γ) in 7/7 patients who either had or soon developed acute rejection compared with 0/8 patients with no rejection or treated rejection (Nast 1994). When Gorzynski et

al studied intragraft cytokine expression during acute allograft rejection of liver transplants they reported quantitative increases in both IL-2 and IFN- γ mRNA when compared with beta actin message. Interestingly they also reported a significant increase in IL-2 and IFN- γ mRNA transcripts in peripheral blood lymphocytes, of the same liver allograft rejectors, compared with non-rejectors (Gorzynski 1996). Iacono and colleagues reported similar findings in lung transplant recipients with refractory acute rejection. They identified elevated levels of IL-6 and IFN- γ mRNA transcripts in both bronchial alveolar lavage cells and peripheral blood lymphocytes when compared to expression of a house keeping gene, namely β -actin. What is interesting is that these levels fell significantly following treatment in 15/17 and all 15 had histological improvement on subsequent biopsies (Iacono 1997). However, Krams et al were unable to confirm these findings in renal allograft biopsies, using similar laboratory techniques. In their study, IL-2 and IFN- γ mRNA were detected infrequently in rejecting and rejected kidneys (Krams 1992). The Manchester group, also studying renal transplant core biopsies, found a similar lack of correlation between acute allograft rejection and both IL-2 and IFN- γ mRNA expression (Joseph 1995). Further conflicting data came from groups using in situ hybridisation techniques. Both groups examined core biopsies from kidneys without evidence of disease and those experiencing acute allograft rejection. Grimm et al found no cytokine staining in 8/14 rejectors. Two from the other six core biopsies from rejecting kidneys expressed IL-2 and 3 of them expressed IL-4 and IFN- γ (Grimm 1995). Vandenbroecke and colleagues studied IL-6, TNF- α and IFN- γ . No RNA transcripts were detected in the normal kidney samples for the three cytokines

examined. However in 6/8 samples from rejecting kidneys had elevated levels of IL-6, but not IFN- γ . TNF- α transcripts were detected in one patient exhibiting acute rejection, but in the absence of IL-6 (Vandenbroeke 1991). It would appear, therefore, that contrary to the Th1/Th2 paradigm, acute allograft rejection could occur in the absence of Th1 cytokines.

1.4.3 Th1 and Th2 Cytokines and Acute Rejection

Some groups have identified a correlation between expression of both Th1 and Th2 cytokines and acute rejection. Loong and colleagues detected, using both an in situ hybridisation technique and immunohistochemical staining, increased cytokine production for both Th1 cytokines (IL-2 and TNF- α) and Th2 cytokines (IL-4, IL-6 and IL-10) in core biopsies from 10 rejecting kidneys. They also identified increased levels of some adhesion molecules and IL-1 in the same specimens (Loong 1996). Similar findings were reported by Xu et al (Xu 1995). Using a RT-PCR system 98 human renal allograft biopsies were examined to identify intragraft expression of both Th1 and Th2 cytokines. They found a significant correlation between intragraft expression of both IL-2 mRNA and IL-10 mRNA with acute allograft rejection. However, neither IL-4 mRNA or IFN- γ mRNA expression were significant correlates with acute allograft rejection in their study.

1.4.4 Th2 Cytokines and Acute Rejection

Other studies have produced results that are even further away from the conventional Th1/Th2 paradigm. Krams et al used the PCR to examine intragraft expression of mRNA in rejecting and rejected human renal allografts. This study identified the presence of both IL-4 and IL-5 mRNA in both sample types, but IL-2 and IFN- γ message was rarely detected (Krams 1992). A group from Boston, USA reported similar findings. They used a competitive RT-PCR system to quantify intragraft gene expression in 60 human renal allograft core biopsies obtained for surveillance or at the time of graft dysfunction. They found the expression of IL-7, IL-10 and IL-15, but not that of IL-2 and IL-4 was significantly heightened during acute allograft rejection compared with non-rejection (Strehlau 1997).

1.4.5 Th2 Cytokines and Acceptance

One would expect, from the Th1/Th2 paradigm, that graft acceptance would be associated with increased expression of Th2 cytokines in the absence of Th1 cytokines. There is, however, little evidence to support this. Gorczynski et al have shown that, in a high proportion of liver biopsies obtained from patients without clinical evidence of rejection, IL-4 mRNA expression was present in the absence of IL-2 mRNA expression (Gorczynski 1996). In renal transplantation the only supporting evidence has been a case report. In this instance the peripheral blood cells of a patient who discontinued all immunosuppression produced high amounts of IL-4 protein when measured

using an ELISA system (Kusaka 1995). As discussed above, most published work that has identified increased levels of Th2 cytokines correlate with acute rejection rather than acceptance.

1.4.6 Cytokine Networks and Redundancy

Evidence for the Th1/Th2 paradigm in the clinical setting is conflicting. This might be explained, at least in part, by the existence of cytokine networks and redundancy. As previously discussed, there is considerable evidence that IL-2 is not essential for acute allograft rejection. This is further supported by work by Van Gelder et al (Van Gelder 1998). They found that blockade of the IL-2/IL-2R by a combination of CyA and an anti-IL-2R monoclonal antibody was insufficient to prevent acute allograft rejection in cardiac allograft recipients. It may be that other cytokines share certain functions in common with IL-2 in this situation. IL-7 and IL-15 are two possible candidates (Nickerson 1997). IL-7-triggered signals, not blocked by CyA, are important for lymphocyte differentiation and stimulation of CTL proliferation (Alderson 1990). IL-15 is produced by a wide variety of cell types, with the notable exception of mature T cells. IL-15 has similar biologic activities to IL-2. These may result from the fact that IL-15 binds to a trimolecular receptor which includes two members of the IL-2R complex (i.e. the IL-2R β chain and the common gamma [γ] chain) (Tagaya 1996).

There is some clinical evidence to support the hypothesis that these non-T cell derived T cell growth factors maybe important in acute allograft. Several

groups have detected the presence of IL-7 and IL-15 mRNA in graft biopsies at the time of allograft rejection. Two of these have been in human renal allografts (Pavlakis 1996, Strehlau 1997) while the other has been in human liver transplants (Baan 1998).

1.4.7 Non-cytokine Influences on the Immune Response

In addition to the role of non-T cell cytokines in clinical transplantation, there is another factor that may influence in vivo immune responses. Class II MHC is obviously crucial due to its part in antigen recognition and resulting T cell activation. However, there is now evidence that these molecules may actually be able to influence the type and size of response. Firstly, DiMolfetto and colleagues have demonstrated that the density of expression of MHC class II can influence Th cell differentiation (DiMolfetto 1998). Secondly, evidence that MHC class II polymorphisms may influence cytokine production has been provided by Petrovsky and Harrison (Petrovsky 1997). Their findings suggest HLA class II alleles, particularly at the DQ locus or genes in linkage with them, regulate IFN- γ expression by T cells.

Some work has been performed on peripheral blood mononuclear cells (PBMCs) extracted from the recipient of a renal allograft prior to transplantation. Kaminski et al stimulated PBMCs with an anti-CD3 antibody. Cells from those patients that went on to develop acute cellular rejection secreted higher mean levels of IL-2 and IFN- γ than those patients who had no rejection episodes, although this was only statistically significant for IFN- γ

(Kaminski 1995). Weimer et al also looked at in vitro cytokine secretion from PBMCs in 80 patients receiving a renal allograft. Using a cell stimulant and an ELISA to determine cytokine secretion, they reported lower pre-transplant IL-10 responses (14/80) were significantly associated with fewer acute rejection episodes and better 3 year graft survival (Weimer 1998).

1.4.8 Why Might The Results Be Conflicting?

It is apparent that the role of individual cytokines in acute allograft rejection is far from clear. There are numerous studies looking at both cytokine gene expression and protein levels at the time of rejection. There is evidence to support the Th1/Th2 paradigm described by Mosmann (Mosmann 1986), but it appears that in clinical human allograft transplantation the picture is far more complicated. Some of the disparity maybe explained by different material analysed (FNAs versus core biopsies), different techniques used in the analysis (antibody staining versus competitive RT-PCR versus non-competitive RT-PCR versus in situ hybridisation), different end points measured (gene expression versus protein levels), different organs and different immunosuppressive regimes. The most likely source of confusion probably results from the sampling. The vast majority of the work has been conducted on single time point biopsies. This may give misleading information on the presence or absence of a specific cytokine as there is experimental evidence that gene expression of IL-2, IL-6, IL-10, TNF- α and IFN- γ , following stimulation, peaks at 8 hours and is back at baseline levels by 24 hours (Fan 1998, Bill 1994). In a dynamic situation like the immune

system sequential measurement using a quantitative method would seem to be more informative. This would allow one to compare expression of a cytokine both to a pre-transplant level and to other cytokines detected.

1.4.9 Cytokine Gene Polymorphisms

Work by a number of groups has introduced another perspective into the potential role of cytokines in the process of acute allograft rejection. They have identified several cytokine gene polymorphisms, which may influence cytokine expression (see below). Most of these are single nucleotide polymorphisms (SNPs) in the upstream regulatory (promoter) sequence of the cytokine gene, for example, TNF- α at position -308, IL-4 at position -590 and IL-10 at position -1082, and may directly influence gene transcription. Other polymorphisms may affect expression by influencing secretion of the protein product. One such example is the signal sequence SNP at position +915 of the TGF- β 1 gene. Still others consist of a variable intron repeat (microsatellite) polymorphisms. For example the IFN- γ gene has a CA repeat in its first intron which varies from 11 to 15 repeats. While such microsatellites are unlikely to have a direct influence on cytokine expression, they may act as a surrogate marker for expression differences due to linkage disequilibrium with SNPs, which influence transcription (Pravica 1999).

Some polymorphisms have been shown to influence cytokine production in vitro and may affect outcome following organ transplantation, and include those associated with IFN- γ (Pravica 1999), TGF- β 1 (Awad 1998), TNF- α

(Wilson 1997), IL-4 (Rosenwasser 1995) and IL-10 (Turner 1997). Table 1.4 shows the polymorphism position, alleles and affect on cytokine production. These are not the only cytokine gene polymorphisms to be described. Many others exist, which potentially, may influence expression of their cytokines. These include IL-1 β at position -511, IL-2 at position -330 and IL-6 at position -174 (Bidwell 1999).

It is sometimes possible to identify a patient as either a high, intermediate or low producer of a particular cytokine. For example, for the IL-10 polymorphism at position -1082, GG homozygotes have been shown to be high producers whereas AG heterozygotes are intermediate and AA subtypes are low producers (Turner/Williams 1997). Usually, however, these genotypes are often combined into just two groups. For IL-10, for example, the genotypes IL-10 -1082 A/G and IL-10 -1082 A/A are defined as low producers (Sankaran 1999).

Table 1.4. Cytokine polymorphisms – position and phenotype

Cytokine	Polymorphism (position)	Alleles	Cytokine production
IFN- γ	First intron (CA) ⁿ repeat	12 repeats	High
		Other alleles	Low
TGF- β 1	Signal sequence (+915)	Arginine	High
		Proline	Low
TNF- α	Promoter (-308)	A	High
		G	Low
IL-4	Promoter (-590)	T	High
		C	Low
IL-10	Promoter (-1082)	G	High
		A	Low

This table shows the polymorphism position, the polymorphism type, the polymorphism alleles and affect the different polymorphisms have on cytokine production in vitro.

1.4.10 Influence of Cytokine Gene Polymorphisms on Rejection

Despite being a Th1 cytokine, the IFN- γ polymorphism in the first intron, has not been strongly associated with acute rejection in transplant recipients in isolation (Hutchinson/Pravica/Hajeer 1999). However, it is strongly associated with rejection when in combination with the high IL-10 genotype (Asderakis 2001).

The TGF- β 1 polymorphism at position +915 has not been shown to be associated with acute allograft rejection, but there is mounting evidence that the high producer genotype has a detrimental affect on the long-term survival of heart, lung, kidney and liver transplants (Hutchinson/Pravica/Hajeer 1999).

There is evidence that the TNF- α -308 promoter polymorphism is strongly associated with frequency and severity of rejection in heart, liver and kidney transplantation. The high TNF- α genotype (TNF -308A), in association with the low IL-10 genotype (-1082A/A) is more common in cardiac transplant recipients having multiple rejection episodes (Turner/Grant 1997), and is very strongly associated with death due to acute rejection in heart transplant patients (Hutchinson/Pravica/Perrey 1999). In HLA-DR mismatched kidney transplant recipients, the high TNF- α genotype (TNF -308A) has been shown to be significantly associated with both multiple rejection episodes and steroid resistant rejection episodes. When these results were combined with those for IL-10, the combination of TNF- α high (-308A)/IL-10 high (-1082G/G) had the worst prognosis (Sankaran 1999). However, work from our own centre

and by others has been unable to confirm any association between TNF- α genotype and acute allograft rejection in renal transplant recipients (Poole 2001, Kerman 2000, Young 2000, George 2000, Marshall/McLaren 2000). However, Bathgate et al have shown a link between the high TNF- α genotype –308 A/A and acute rejection in liver transplants. Interestingly, IL-10 had no significant association with acute rejection in this study (Bathgate 2000). When the high TNF- α genotype is from the donor, its effect appears to depend on the organ system. In kidney recipients it does not appear to alter outcome (Hutchinson/Pravica/Hajeer 1999, Poole 2001). Conversely, there is an effect in cardiac transplant recipients, but surprisingly the high TNF- α genotype is associated with a significantly better graft survival (Hutchinson/Pravica/Hajeer 1999).

The influence of the IL-4 promoter polymorphism on the clinical course in renal transplant recipients was studied in our centre by Poole et al in 120 donor/recipient pairs. This work showed that donor/recipient pairs both lacking the IL-4-T (high expresser) allele were significantly increased in the rejectors (Poole 2001).

The reported effect that the IL-10 genotype has on rejection episodes may vary. In heart transplant recipients the high IL-10 genotype appears to be protective (Turner/Williams 1997). This finding has been corroborated, by some, in renal transplant recipients (Kerma 2000, George 2000), and is supported in part by work from our laboratory. We have reported that the combination of recipient IL-10 –1082 G/G high producer/donor IL-10 –1082

A low producer was significantly decreased in renal transplant recipients experiencing multiple rejection episodes (Poole 2001). However, Sankaran et al have reported that the opposite is true in renal recipients, with the high genotype being associated with multiple (≥ 2) rejection episodes in HLA-DR mismatched kidneys and that this effect is exacerbated when combined with the high TNF α 308 producer genotype (Sankaran 1999). Finally, some groups have been unable to show any relationship between IL-10 genotype and rejection episodes in renal transplants (Young 2000, Marshall/McLaren 2000).

1.4.11 Why Might There Be This Disparity?

The first possibility, is that different cytokines, and therefore different cytokine gene polymorphisms, have different influences on different organ systems. From the sections above, it can be seen that cytokines may not always have the same effect and may not always be necessary for rejection or acceptance to occur. Secondly, different immunosuppressive regimes may influence cytokine production. This may either enhance or negate the importance of a cytokine gene polymorphism. Finally, the different studies discussed above use various criteria for classifying recipients as multiple rejectors, some involve donor and recipient status, while others group the genotypes differently, and finally, few mention the ethnic make-up of the studies. (Marshall/Welsh 2000)

Given the complexity of the cytokine network, it is perhaps not surprising that the impact of individual cytokine polymorphisms on outcome is proving

difficult to ascertain. This may be that the analysis being used is too crude to detect the influence of single cytokine polymorphisms. Recent work by Hutchinson and colleagues has used a novel approach to the problem. They have entered all the cytokine genotype data for IFN- γ , TNF- α , IL-10 and TGF- β 1 from a single transplant unit into a neural network computer program. This program apparently "learns" to predict rejection from the data supplied. The first observation was that the program was remarkably good at prediction of acute rejection. Secondly, the inclusion of both the donor and recipient cytokine genotype improved the predictive power even more (Hutchinson/Pravica/Hajeer 1999).

It would appear that cytokine gene polymorphisms have the ability, at least in part, to influence the outcome of clinical organ transplantation. Unfortunately, it is necessary to do further work before this information can be used prospectively to affect patient care and improve both graft and patient survival.

1.5 Immunological Monitoring

Regular monitoring of patients following surgical procedures is routine practice. This usually involves monitoring the patient's heart rate, blood pressure, temperature and oxygen saturation. Following renal transplantation it is no different, but the monitoring is more extensive and intensive. The latter also involves central venous pressure measurements, cardiac rhythm and, obviously, urine output.

For as long as the transplant is functioning, the renal transplant recipient will be monitored by both physical examination and various special investigations. Initially, this occurs at least daily, but as time progresses monitoring becomes less frequent. This is a reflection of the fact that most reversible causes of graft dysfunction occur within the first six months following renal transplantation.

Over the first two weeks or so the kidney recipient will have daily blood tests measuring renal function and monitoring the haemoglobin level and white blood cell count. Urine samples will be sent regularly to exclude infection. Also in the first few weeks, the transplant kidney will often undergo regular ultrasound and, more recently, duplex Doppler examination to exclude obstruction, monitor blood flow within the kidney and to confirm the absence of any perinephric collections. Kidneys exhibiting delayed graft function may need to be monitored with regular core biopsies or FNA samples to exclude acute allograft rejection as the clinician is unable to use renal function as an

indicator of immunological quiescence. Finally, some of the drugs used for immunosuppression, such as cyclosporin and tacrolimus, require monitoring of blood levels, as there is considerable individual variation in both absorbency and bioavailability of these compounds.

Monitoring, as described above, serves two purposes. Firstly, to monitor the health of the recipient and secondly, to monitor the 'health' of the transplanted kidney. Most of the investigations are to exclude renal dysfunction. This can be the result of several different processes, such as rejection, infection and obstruction. Early and accurate diagnosis of acute allograft rejection is essential for the administration of optimal immunosuppression, therefore optimising both the function and half-life of the transplant. The commonly used laboratory tests, such as creatinine and urea, are good indicators of a decrease in function of the transplanted kidney. Unfortunately, they lack specificity as to the cause of the dysfunction. More accurate diagnosis necessitates histological examination of biopsy material. Obtaining these specimens is invasive and there is a potential risk of damage to the organ. These samples are usually only taken when there is evidence of dysfunction, which may reflect organ damage.

Immunological monitoring of renal allograft recipients is based on the assumption that early recognition and treatment of acute allograft rejection may decrease the extent of irreversible damage in the graft and improves graft survival. Acute allograft rejection is the most ominous of several aetiologies that may underlie graft dysfunction. Serial monitoring, therefore, may be

helpful in the differential diagnosis of rejection, infectious complications and cyclosporin nephrotoxicity. It may also be possible to monitor treatment and, therefore, detect early failure to allow a change in anti-rejection therapy.

Over the past three decades, there has been a lot of work carried out in this field. Many different groups have studied various methods of immunological monitoring. This work will be reviewed in detail below. Much of this work has been carried out in the field of cardiac transplantation. This is because endomyocardial biopsy is invasive, expensive and potentially hazardous so the potential benefits of a monitoring tool are large. However, this review will focus on the use of immunological monitoring in the field of renal allograft transplantation.

1.5.1 The Measurement of Constituents of the Urine for Immunological Monitoring

Urine Cytology

Urine can contain cells, proteins and polypeptides. Therefore, urine can be used in several different ways for the purposes of immunological monitoring. Urine constituents have been used in isolation and in combination with other components of the urine and/or serum.

Several groups have shown that by monitoring the appearance of lymphocytes or an increase in the number of lymphocytes in the urine of renal transplant and pancreatic transplant (when there is urinary diversion of the exocrine pancreas) recipients correlates with immune events and precedes clinically

detectable rejection (Schumann 1977, Eggensperger 1988, Tyden 1987, Murphy 1973, Kauffman 1964). Sandoz et al went on to show that the accuracy of such tests seemed comparable to the diagnostic value of FNA from the graft itself (Sandoz 1986).

Simpson et al showed that the binding of goat antihuman globulin to renal tubular cells recovered from the urine of renal transplant recipients correlated well with impending or established acute allograft rejection, but not with other causes of renal dysfunction (Simpson 1985). This group then did further work on urine cytology and acute rejection. In this later study, they were able to repeat their earlier findings, and also show a good correlation between lymphocyturia and the presence of collecting duct cells in the urine when biopsy findings met acute rejection criteria in 48/51 patients. Two of the three patients in whom discrepant results were observed were later diagnosed as having an alternative cause for their biopsy findings (Simpson 1987).

Cytological examination is not only useful in renal transplant recipients, but has also been shown to correlate well with acute rejection in hepatic transplantation. In this clinical setting, the presence of inflammatory cells in the bile not only indicated the onset of rejection with a specificity of >94%, but also correlated with the persistence of rejection following treatment (Roberti 1992).

Urine cytology has the advantage that it is totally non-invasive and samples can easily be obtained on a daily basis. Not only can it be used for the

diagnosis of rejection episodes, but also ischaemic damage, drug toxicity and viral infections. However, causes of oliguria and anuria (acute tubular necrosis, obstruction and severe rejection) can make collection difficult. Also, cell disintegration in the urine due to proteolytic enzymes and bacterial cytotoxins, or hypertonic urine make evaluation difficult. Finally, additional cells from other parts of the urinary system and genital tract can disturb the differential (Hammer 1993).

Cytokines

Some groups have studied whether the measurement of cytokines (using bioassays and ELISA) in urine could be used for the purpose of immunological monitoring. Work published in this area has identified an association between elevated levels of urinary IL-2 (Simpson 1989), IL-6 (Van Oers 1988) and TNF- α (McLaughlin 1991) with acute rejection in renal transplant recipients when compared to controls and patients with stable renal function. For IL-2 there was also a significant rise in levels associated with CMV disease, but this is not seen with TNF- α measurement. However this rise in urinary TNF- α was only seen in 49% of rejectors. A rise in urinary IL-2 one to three days prior to conventional parameters in acute rejection in pancreatic transplants has also been demonstrated. However, elevated levels may also be seen in the presence of infection and acute pancreatitis (Georgi 1989).

Recently, Teppo and colleagues have studied the relationship between the IL-1 receptor antagonist (IL-1ra)/IL-1 β ratio in the urine of renal transplant

recipients. They measured the ratio in 69 twenty-four hour urine samples from 23 rejectors and 51 such samples from 17 patients with stable graft function. IL-1ra/IL-1 β ratios dipped at the time of rejection, and possibly before, compared with earlier samples, and rose again during treatment. However, the ratio remained unchanged in patients with stable graft function. These authors concluded that a high urinary IL-1ra/IL-1 β ratio might have a protective role after kidney transplantation (Teppo 1998). This study would have been improved if the pre-rejection monitoring had been done at multiple time points, rather than just one. This would have shown whether this ratio has a predictive value, as a sporadic value could not predict incipient rejection. They also failed to assess the influence of alternative causes of graft dysfunction on the urinary IL-1ra/IL-1 β ratio.

It is also possible to measure cytokine mRNA in the urine of renal transplant recipients. Jeyarajah and colleagues, using the PCR and Southern blotting, demonstrated the use of immunological monitoring for post-therapy surveillance following a rejection episode. They reported a consistent rise in IL-5 mRNA followed successful treatment and a fall in both IL-2 and IFN- γ , but due to small numbers only the former rise was statistically significant (Jeyarajah 1995).

Soluble IL-2 receptor

Simpson also studied levels of soluble IL-2 receptor (sIL-2R) in the urine of kidney recipients (Simpson 1989) and the bile of liver recipients (Simpson 1991). In the former group, she reported similar findings to those of urinary

IL-2. In the liver transplant recipients there was also a significant rise in mean bile sIL-2R levels which preceded clinical rejection, when compared with patients with stable function. There was overlap in ranges, but this was less of a problem if sequential measurements were analysed. Unfortunately in this paper there was no comment on the effect of CMV disease.

Adhesion Molecules

Some groups have examined the usefulness of detecting adhesion molecules in the urine to both predict rejection and monitor treatment. Chan et al studied the expression of intercellular adhesion molecule-1 (ICAM-1) on renal tubular cells excreted in the urine using immunocytology. They reported a significant increase in expression of ICAM-1 both before and during acute rejection (compared with patients with acute tubular necrosis) and demonstrated a significant drop in expression after successful treatment (Chan 1993). Bechtel and colleagues investigated the use of monitoring not only ICAM-1 in the urine of renal transplant recipients, but also vascular adhesion molecule-1 (VCAM-1) and endothelial adhesion molecule-1 (ELAM-1), as well as complement cleavage products C4d and C5b-9. They did not study expression, but measured the quantity of soluble molecules in the urine using an ELISA system. They reported significantly higher levels of both ICAM-1 and VCAM-1 in the urine of patients with acute steroid-resistant rejection compared with those with stable graft function and acute steroid-sensitive rejection. ELAM-1 was not detected in the urine of any patient. The results of C4d measurement were very similar to that of ICAM-1 and VCAM-1. C5b-9 measurement did not correlate with the severity of graft rejection. By using

immunohistochemistry on biopsies taken from the patients with steroid-resistant rejection, they were able to show that the degree of inflammation within the graft was reflected by the urinary concentration of soluble ICAM-1 and VCAM-1 (Bechtel 1994).

Interestingly, elevated excretion of soluble ICAM-1 into bile has also been noted during rejection of liver allografts (Adams 1993).

C-Reactive Protein

Chun first identified the association between urinary proteins and acute allograft rejection in 1972. This paper reported increased urinary C-reactive protein associated with rejection in renal transplant recipients (Chun 1972). Since then, Jürgen Steinhoff and colleagues have extensively studied the use of urinary proteins for their predictive value in diagnosing the cause of renal allograft dysfunction. By using the measurements (using immunoluminometric assays) of a combination of three proteins in the urine (C-reactive protein, α 2-macroglobulin and myeloperoxidase), they were able to confidently distinguish between stable graft function, acute allograft rejection, urinary tract infection, CMV infection and CyA nephrotoxicity. Admittedly, diuresis is a prerequisite, and more complex problems, especially the simultaneous occurrence of two or more pathological conditions, would need more elaborate investigation (Steinhoff 1997).

1.5.2 The Measurement of Serum Proteins and Soluble Molecules for Immunological Monitoring

β 2 microglobulin

As previously discussed in section 1.2, β 2 microglobulin (β 2M) is part of the HLA class I molecule. It is shed from nucleated cells in proportion to the rate of cellular turnover. It is readily filtered by the glomerular basement membrane and almost completely reabsorbed and catabolised by proximal renal tubules (Schweizer 1981).

Early reports suggested its usefulness in monitoring renal transplant recipients and predicting acute rejection episodes. Schweizer et al studied serum β 2M levels using radioimmunoassays in 52 renal transplant recipients. They reported that 43% of patients had a detectable elevation in serum β 2M prior to clinical detection of the rejection episode and a further 24% at the time of clinical detection. There were no rejection episodes where serum β 2M did not rise. What was disappointing was that there were 36 occasions where persistent elevations in serum β 2M were not due to rejection (Schweizer 1981). Very similar findings were reported in 8 cardiac transplant recipients experiencing 16 rejection episodes (Goldman 1983). In addition, Edwards and colleagues examined the usefulness of serum β 2M in monitoring renal transplant recipients. The advantage of the study over others was that it established the normal variation in serum β 2M in a population with stable renal function and a population on haemodialysis. They went on to study 93 recipients experiencing a total of 90 rejection episodes. Daily serum β 2M measurements using a solid-phase radioimmunoassay were taken in the early

post-transplant period. Forty-nine per cent of rejectors had a significant rise in their serum β 2M at least one day prior to the rise in serum creatinine and a further 42% had a simultaneous rise in both their serum β 2M and serum creatinine. Of the remaining 8 rejectors, all had a significant rise in their serum β 2M 24-48 hours after the rise seen in serum creatinine. Interestingly, they only reported 4 false positives. The weaknesses of this work are that there is no comment about immunosuppressive usage, the diagnosis of rejection was almost always made clinically, there is no mention of the effect of infection on serum levels and finally the effect of treatment on levels was not discussed (Edwards 1983). A later paper by Prischl et al attempted to answer some of these questions. Unfortunately, it was a retrospective review of renal transplant dysfunction in 49 recipients experiencing 100 episodes of graft deterioration. The diagnosis of the cause of the dysfunction was made retrospectively by studying the response to methyl prednisolone treatment and other parameters. In those that were diagnosed as experiencing acute rejection, serum β 2M rose in 46/52 and fell after treatment. In those with CyA toxicity, serum β 2M levels remained stable or even decreased. The urinary β 2M levels were significantly higher in those with CyA toxicity compared with rejectors and those with stable function (Prischl 1989). Not all workers, however, have concluded that the measurement of serum β 2M levels is a useful tool in immunological monitoring of renal transplant recipients. Vernon et al studied a similar sized cohort to Edwards, but reported dramatically different results. They found serum β 2M levels to be less specific and less sensitive than serum creatinine levels in monitoring renal transplant recipients for acute rejection episodes. They also were unable to confirm that elevations of serum β 2M

levels, due to acute rejection, preceded a rise in serum creatinine (Vernon 1985).

Adams and colleagues also found that the serum β 2M level was of no value in predicting acute rejection episodes in liver transplant recipients. However, they did report the level of biliary β 2M was significantly higher in patients experiencing acute rejection episodes compared with other complicated and uncomplicated recipients, and when the bile/serum ratio of β 2M was calculated these differences were even greater (Adams 1988).

C-Reactive Protein

As well as studying urine C-reactive protein (CRPu), Steinhoff has also used serum C-reactive protein (CRPs) to differentiate between different causes of renal transplant dysfunction. Using the ratio CRPs/CRPu it is possible to distinguish between urinary infection (quotient above 1.0) and acute rejection (quotient below 1.0) (Steinhoff 1992).

Other groups have also looked at CRPs. Lalla et al presented results from 31 renal transplant recipients whom had daily CRPs measurements. They reported elevated levels in 18/27 rejection episodes, when compared to baseline. Unfortunately, levels were also increased in infection, and with considerable overlap. There were also two unexplained elevations in CRPs levels (Lalla 1988). On the other hand, work in our own transplant unit has shown CRPs to be useful in monitoring renal transplant recipients. In uncomplicated renal transplant recipients with acute rejection there was a

significant rise in CRPs preceding acute rejection compared with ATN and CyA toxicity. This increase fell back towards baseline following successful treatment with methyl prednisolone, but remained elevated in those patients with steroid-resistant rejection (Harris 1995).

Serum Amyloid A

Maury et al have studied the value of serum amyloid A (SAA) protein in 31 renal transplant recipients and compared it to both CRPs and serum β 2M levels in the same patients. All three were predictive of acute rejection episodes, but the rise in SAA levels was the greatest. All three had false positives. The lowest number was seen with SAA, and the highest with serum β 2M. In the case of SAA, they were all caused by infective complications. The sensitivity was best for SAA and comparable for the other two. The positive predictive values were 80%, 64% and 55% for SAA, CRPs and serum β 2M respectively. The conclusion is that of the three, SAA is the most useful, but does not adequately differentiate between infection and acute rejection (Maury 1984).

It would appear that CRPs is of less use in liver transplant recipients. Only 2/7 patients developed elevated CRPs levels when acutely rejecting, whereas 5/7 patients with CMV disease and 7/9 with bacterial infections showed markedly elevated levels of CRPs (Tilg/Nordberg 1992).

Soluble HLA class I

Soluble HLA (sHLA) class I molecule levels have been measured in peripheral blood following transplantation. This has been done in 5 kidney and 8 simultaneous kidney and pancreas transplants with HLA-A2 negative recipients and HLA-A2 positive donors. Elevated levels of donor sHLA-A2 accompanied acute rejection, although not every elevation was associated with rejection. It was also noted that there was a less dramatic rise in total sHLA class I levels during rejection (DeVito-Haynes 1994). Drouet and colleagues studied the total amount of sHLA class I in renal transplant recipients. They reported elevated levels in 100% of CMV infections and 60% of acute rejection episodes. They also detected elevated levels on several occasions when no clinical events were diagnosed (Drouet 1995). In liver recipients it has been reported that it is also possible to detect a rise in total sHLA class I for 6 days preceding acute rejection. These levels appear to fall after successful treatment (Puppo 1994).

Soluble IL-2 Receptor

Other groups have examined the benefit of measuring serum levels of soluble IL-2 receptor (sIL-2R). Forsythe et al, using an ELISA system, demonstrated elevated levels of sIL-2R both preceding and at the time of rejection in 8 renal transplant recipients. In those who were successfully treated, the sIL-2R levels fell back to pre-rejection levels. Other causes of renal dysfunction were not associated with significant elevations of levels, but there were only 7 patient episodes in this category. Unfortunately, the control group of patients with stable graft function only consisted of 3 individuals (Forsythe 1989). Other

workers, using a sandwich enzyme immunoassay in 54 renal transplant recipients, also demonstrated sIL-2R successfully distinguished acute rejection from stable graft function and CyA toxicity, but not viral infection (Noronha 1990). A different group, this time using an ELISA system, were less convinced of the benefits of sIL-2R measurements in renal transplant recipients. They only detected significant elevations, from a pre-transplant baseline, in 5/24 acute rejection episodes (4/5 with elevated levels were successfully treated with a corresponding fall in sIL-2R levels). This was partly explained by the high levels in the pre-transplant levels compared with normal controls. What was interesting was that the sIL-2R levels in patients with stable renal function fell to normal control levels, significantly lower than pre-transplant levels (de Boccardo 1994).

Similar findings have been reported in liver transplant recipients. Significant rises in serum sIL-2R have been shown to precede acute rejection and CMV disease. The rises are larger in rejection, but there may be overlap with levels in those with stable graft function (Simpson 1991, Perkins 1989, Adams 1989).

Soluble CD23

Other groups have studied circulating serum levels of soluble CD23 (sCD23) after renal transplantation. This is a low affinity receptor for immunoglobulin E and is expressed on B lymphocytes, monocytes, macrophages and eosinophils. The results have been somewhat variable. One group found that 83% of rejectors showed elevation in sCD23 levels prior to clinical diagnosis,

but 73% of those with infective complications showed a similar increase (Traindl 1994). Another group found that serum levels of sCD23 were no different in rejectors compared with those with stable graft function and delayed graft function (Kutukuler 1995).

Soluble ICAM-1

Using an ELISA system, Stockenhuber et al studied serial serum measurements of soluble ICAM-1 in 49 renal transplant recipients. They reported significantly raised mean levels of soluble ICAM-1 in acute rejectors compared with patients with stable function, but levels were also raised in CMV disease and chronic rejection, but not CyA toxicity. In sequential measurements the rise in soluble ICAM-1 was seen 1-2 days before the clinical signs of rejection, and fell to baseline in patients with steroid-sensitive rejection (Stockenhuber 1993).

E-selectin

E-selectin is a glycoprotein expressed on the surface of vascular endothelial cells in response to TNF- α and IL-1. Using an ELISA system, Weston and colleagues measured serum levels of soluble E-selectin in renal transplant recipients. They were unable to show any significant difference in the levels of patients with stable renal function and those experiencing acute rejection, CyA toxicity and ATN. They, not surprisingly, drew the conclusion that monitoring of E-selectin was of no benefit in renal transplant recipients (Weston 1995).

1.5.3 The Measurement of Serum Cytokines for Immunological Monitoring

Tumour Necrosis Factor- α

Maury and Teppo were some of the first people to study the possible use of serum cytokines as tools for immunological monitoring. They studied sequential serum TNF- α , using a radioimmunoassay, in 10 renal transplant recipients treated with azathioprine and prednisolone and 40 healthy controls. They reported elevated levels associated with febrile rejection episodes, but not all rejection episodes, and also with bacterial and viral infection and pyrexia of unknown aetiology (Maury 1987). Other groups have subsequently studied this cytokine in both renal and non-renal transplant recipients. Noronha and colleagues reported similar results using an ELISA. They again identified elevated serum levels of TNF- α associated with acute renal allograft rejection. However, 3 patients in their series, who rejected, had undetectable levels. They also found that serum TNF- α measurements could differentiate between rejection and CyA toxicity, but not viral infection (Noronha 1990). McLaughlin's results were also disappointing, only identifying elevated levels in 65% of renal allograft rejectors (McLaughlin 1991). One study has reported results from studying both serum TNF- α and serum TNF- β in renal recipients, again using an ELISA. Both were noted to rise prior to rejection and infection, but there was a significantly larger mean increase of TNF- α in those that went on to reject compared with those that had an infection (Daniel 1995). Some, however, have not detected any association between TNF- α and rejection in renal transplant recipients (Tice 1993).

Studies of TNF- α in liver transplant recipients have revealed similar findings. Elevated TNF- α levels are associated with acute rejection, but also with infection, rendering this investigation unhelpful (Vogel 1990). One group has studied lymphotoxin (TNF- β) in a similar setting and found that relative increases in levels were associated with impending rejection, and that levels fell after successful treatment. Unfortunately the influence that infection may have on the level of lymphotoxin was not addressed (Sankary 1993).

In small bowel transplantation elevated TNF- α levels can be detected, but these may not always be associated with acute rejection episodes (Noguchi 1992).

Despite the evidence to support the role of the TNF- α -308 promoter polymorphism in the frequency of acute rejection episodes in cardiac transplant recipients, there is little evidence that measuring serum levels of this cytokine is of any predictive value in this group of patients (Grant 1996, Azzawi 1996, Torre-Amione 1998). However, sustained elevations in this cytokine may be a harbinger of profound morbidity. Two patients in one study, investigating the use of serum TNF- α levels in cardiac transplant recipients, had persistently high levels. One died within 30 days of the transplant due to a lymphoproliferative disorder and the other died within 3 months after repeated episodes of severe rejection, which were refractory to treatment (Kimball 1996).

Rondeau et al took a slightly different approach and measured TNF- α production (using an immunoradiometric assay) by cells of bronchioloalveolar lavage and peripheral blood mononuclear cells in cardiopulmonary transplant recipients. The results, however, were similar in that, rejection episodes were not associated with significant changes in TNF- α synthesis by either cell type (Rondeau 1990).

Interleukin-1

Maury and Teppo also studied IL-1 in renal transplant recipients. Daily blood samples were taken from twelve patients for the first 3-4 weeks following transplantation. Immunoreactive IL-1 was measured using a solid-phase radioimmunoassay. IL-1 levels were elevated 0-2 days prior to rejection, but there were also fluctuation of levels in 1/3 non-rejectors, reducing the usefulness of the measurements (Maury1987).

Interleukin-2

As IL-2 has such a key role in immune activation, it is hardly surprising that several different groups have studied measurement of serum levels of this cytokine in transplant recipients. Elevated serum IL-2 levels have been reported to precede signs of clinical rejection in renal transplant recipients and return to baseline following successful treatment. Elevated levels of serum IL-2 are not seen in CyA toxicity but there is some overlap with CMV disease. As well as these findings Simpson et al also found that 19/21 patients who had elevated pre-transplant serum levels of IL-2 (>15ng/ml as measured by an ELISA system) experienced acute allograft rejection compared with 15/70

recipients with low pre-transplant levels (Simpson 1989). Albert et al also noted an overlap between elevated serum IL-2 in acute rejection and viral infection in kidney recipients, but felt it was a useful investigation (Albert 1990). Johnson and colleagues found that elevated serum IL-2 levels correlated less well with the severity of the rejection. This was particularly well illustrated by 4 patients in their series who had severe rejection requiring monoclonal antibody therapy with extremely low or undetectable serum IL-2 levels. They were of the opinion that measuring several cytokines might be of more benefit than a single one (Johnson 1990). Not all groups studying renal transplant recipients have found the measurement of serum IL-2 to be of benefit, as they found alterations in levels appeared to occur randomly (Tice 1993), or were unable to differentiate from infective episodes (Daniel 1995).

In other transplant populations IL-2 has received mixed reviews. In pancreatic transplantation, elevations in serum levels have been shown to be associated with acute rejection, but also infection (Georgi 1989). In small bowel transplants acute rejection may be preceded by a serum rise in IL-2 (Noguchi 1992). In liver recipients there is a significant rise in IL-2 levels in both serum and bile associated with acute rejection and this rise usually precedes clinical diagnosis (Simpson 1991). In heart transplant recipients, however, IL-2 levels were of little help in predicting acute rejection episodes (Kimball 1996).

Interferon- γ

There is little evidence to suggest that serum level of IFN- γ is a reliable tool for immunological monitoring in renal transplantation. This is because its baseline levels fluctuate in stable graft function, infection and acute rejection (Steiner 1990, Daniel 1995).

Interleukin-4

Some work has been done on other cytokines. IL-4 has been shown to rise both before and after the clinical diagnosis of acute rejection in renal transplant recipients (Kutukculer 1995). In a different study of renal transplant recipients, serum IL-4 levels did not significantly change before either acute viral infections or acute rejection episodes (Daniel 1995). These studies suggest that the measurement of serum IL-4 levels is of limited value in immunological monitoring.

Interleukin-6

As previously mentioned (section 1.51), Van Oers reported elevated levels of IL-6 in the urine of patients experiencing acute rejection. This group also reported an elevation in serum IL-6 post-transplant and at the time of rejection, but these changes were less specific than those in urine, as IL-6 was also detected in the serum of healthy controls (Van Oers 1988). In a later study, using an ELISA technique, elevated serum IL-6 levels were detected in 6/7 renal transplant recipients who experienced an episode of acute rejection, as reported by Kutukculer et al, compared with those with stable function. The elevated IL-6 level preceded clinical rejection in 5/6 cases when it was

detected. The samples were collected during the first 14 days following transplantation. Unfortunately, there were no infective episodes in the sixteen patients studied and no mention of any experiencing CyA toxicity (Kutukculer 1995).

Yoshimura et al (using an IL-6-dependent cell line) also reported elevated levels of IL-6 in the serum of renal transplant recipients preceding clinical signs of acute rejection. They noted that these levels fell to baseline after successful treatment. These elevations differentiated dysfunction secondary to rejection from that of CyA toxicity as IL-6 levels fell in the case of toxicity. Unfortunately, a rise in IL-6 was also seen in infection and, as a result, failed to discriminate between rejection and infection (Yoshimura 1991). The Opelz group has reported similar findings when measuring serum IL-6 using an ELISA. An elevation of serum IL-6 preceded both clinical acute rejection and infection in a renal transplant population, but the mean elevation was significantly higher in the case of rejection. As in so many of these studies, there was considerable individual variation in absolute levels (Daniel 1995). Tice and colleagues were again unable to corroborate these findings in 40 kidney recipients using an ELISA (Tice 1993).

In other organ transplant settings elevated serum IL-6 levels have been associated with acute rejection in some studies, but not all. Noguchi detected elevated levels in small bowel transplants (Noguchi 1992) and Kimball had similar results in cardiac transplants (Kimball 1996). Interesting, this later report also indicates that IL-6 serum measurement gives a good and early

indication of the likelihood of response to treatment with elevated levels persisting in non-responders. Grant and colleagues were unable to corroborate these findings, detecting no association between serum IL-6 levels and acute rejection in cardiac transplant recipients (Grant 1996). In lung transplant recipients, acute rejection has also been associated with increased expression of IL-6 mRNA measured in bronchoalveolar lavage cells by RT-PCR, and that its degree of expression correlates with histological persistence of rejection (Iacono 1997). In liver transplant recipients, Tilg et al evaluated serum IL-6 activity by testing its capacity to induce proliferation of the IL-6-dependent hybridoma cell line B9. They were unable to show an association between elevated IL-6 levels and acute rejection, but did identify elevated levels with both bacterial and viral infection (Tilg/Nordberg 1992).

Interleukin-8

The results of studies examining serum IL-8 levels are interesting. In renal (Daniel 1995, Kutukculer 1995) and liver (Tilg/Ceska 1992) transplant recipients it has not been shown to be of any predictive value with regard to acute rejection, but does correlate with infection in the latter group. However, in cardiac recipients, IL-8 was an effective marker of significant rejection 2-4 days before the clinical diagnosis was made and also predicted resistance to monoclonal antibody therapy (Kimball 1996).

Interleukin-10

IL-10 has not been shown to be of predictive value in the diagnosis of acute rejection in renal transplant recipients, but does appear to be inducible following the additional treatment given for acute rejection. Daniel et al detected a significant rise in serum IL-10 levels within the first few days of the commencement of anti-rejection treatment (Daniel 1995). In cardiac transplant recipients no association was detected between IL-10 and acute rejection (Grant 1996).

1.5.4 The Measurement of Peripheral Blood Cell Surface Markers for Immunological Monitoring

As discussed earlier, when T cells are activated they release cytokines that lead to cell proliferation and differentiation. This activation is associated with increased expression of HLA molecules on the cell surface and the expression of other cell surface receptors and molecules e.g. IL-2 receptor and transferrin receptor (Stenzel 1978, Murphy 1986, Colvin 1987). These 'markers' could potentially be used to differentiate between active and quiescent cells and, therefore, have a role in monitoring transplant patients.

CD4/CD8 Ratio

Monitoring of total T cell levels has not been shown to be of value (Sengar 1977, Thomas 1977). As different subsets have different roles and actions in the immune response, groups started to study the relative increase or decrease in these respective groups. Using monoclonal antibodies and flow cytometry, Cosimi et al sequentially monitored 29 HLA-nonidentical renal transplant

recipients. They demonstrated that an elevated CD4/CD8 ratio was associated with acute rejection in the majority of patients experiencing rejection episodes (Cosimi 1981). In a comparable series of 30 patients, Ellis and colleagues, using similar techniques, reported high CD4/CD8 ratios were associated with rejection and a low ratio with stable graft function and quiescence in those who had previously experienced rejection (Ellis 1981). Kerman and colleagues also demonstrated elevated CD4/CD8 ratios with rejection in a series of 88 renal transplant recipients, but did think that it was a reliable index of rejection as increases in CD4/CD8 ratios were associated with both rejection and non-specific immune events. Also these elevated ratios only preceded the event on about 10% of occasions (Kerman/Van Buren 1983). Other groups have been unable to detect any relationship between CD4/CD8 ratio and acute renal allograft rejection, and have concluded that the CD4/CD8 ratio has no predictive value when correlating the results with rejection episodes (Von Willebrand 1983). One group detected a fall in the CD4/CD8 ratio associated with rejection in renal patients treated with azathioprine and prednisolone, but no change in the ratio in patients rejecting on CyA (Carter 1983).

Those groups that reported the ratio to be of benefit in predicting graft rejection have also indicated that it may differentiate between other causes of graft dysfunction. It has been proposed that the CD4/CD8 ratio might be of benefit in differentiating between those patients that are rejecting and those experiencing CyA toxicity. In the former group, rejection was always associated with an increased ratio, whereas, 91% of patients who were CyA

toxic had a ratio that remained low (Kerman/Flechner 1983). It has also been reported that viral complications are associated with lower ratios than in those rejecting, but this is not exclusive as rejection is seen in patients with low ratios, and may indicate a worse outcome (Colvin 1983).

Other groups have also reported conflicting results when studying CD4/CD8 ratios in combination with other surface markers in cardiac transplant recipients. Hoshinaga, using flow cytometry, studied the relationship between the percentage of transferrin receptor positive cells and the CD4/CD8 ratio in rejection and infection in 41 cardiac recipients. They reported increases in both parameters both preceding and during acute rejection episodes. There was a positive predictive rate of 72%. During infection the percentage of transferrin receptor positive cells increased to a level greater than that seen in rejection, but the CD4/CD8 ratio fell. Therefore, it may be possible to differentiate between infection and rejection. They also noted that a failure of the level of transferrin receptor positive cells to fall following treatment for rejection was highly predictive of a further acute rejection episode (Hoshinaga 1987). In a series of 10 cardiac transplant recipients, Coles et al studied the CD4/CD8 ratio and the percentage of cells expressing the IL-2 receptor. Neither measurement, in isolation or in combination, had any predictive value. The negative results may have resulted from the small numbers studied or the technique (smears of whole blood and an alkaline phosphatase anti- alkaline phosphatase technique) used (Coles 1987).

In renal transplantation CD4 and CD8 have been studied in combination with other surface antigens and molecules. In one study 83 patients had sequential monitoring of their CD4⁺ CD28⁻ T-helper/inducer lymphocytes during different clinical scenarios using flow cytometry. (CD28 is a costimulatory molecule found on the surface of T cells and interacts with B7.1 and B7.2 on the surface of B cells). A high level was not predictive of rejection, but a low level was significantly associated with viral and bacterial infection (Brando 1989). In another study, 60 renal recipients underwent sequential monitoring using flow cytometry, monoclonal antibodies and PBMCs. The reported findings associated with acute rejection were an increase in CD4/CD8 ratio, and an increase in both CD3CD25 and CD4 T cell subsets. (CD25 is the IL-2 receptor located on the surface of activated T cells). In the case of viral infection, the CD4/CD8 ratio fell and there were increases in CD8CD25, CD3DR and CD8 T cell subsets. Unfortunately there is no mention of the sensitivity and specificity of these observations (Valeri 1991).

Masri and colleagues attempted to overcome the problems of 'indirect cytokine measurement' described by other authors, by using flow cytometry, in vitro cell stimulation and a panel of monoclonal antibodies to study T cell subsets and the cytokines they produce. These workers examined 215 blood samples in 34 renal transplant recipients. Their results indicate that, following in vitro mitogen stimulation, patients with acute rejection expressed a combination of high intracellular CD4 and CD8 IL-2 levels, high CD4/CD69, and low intracellular levels for both the so called Th2 cytokines, IL-4 and IL-10 in CD4 and CD8 cells. (CD69 is a signal transduction molecule located on

the surface of T cells). Patients with good and stable function had low intracellular IL-2 levels in CD4 cells and 'normal' levels of both IL-4 and IL-10 in CD4 cells. Patients with viral infection had a pattern distinct from the other two. There is, however, no comment on whether the changes seen in the rejectors precede the clinical diagnosis. Also by using in vitro mitogen stimulation has the problem of 'indirect cytokine measurement' totally been avoided? (Masri 1998, Masri 1999).

IL-2 Receptor

As discussed above, IL-2 receptor (IL-2R) measurement in blood, urine and bile has been shown to predict rejection in some studies. IL-2R expression on the cell surface has also been discussed above and again its usefulness is debatable. Langley et al took a slightly different approach. These authors extracted lymphocytes from 57 renal transplant recipients and studied whether inducible levels of IL-2R correlated with clinical events. Increased levels of CD8 cells expressing inducible IL-2R correlated with acute rejection, the sensitivity and specificity of their results were 87% and 83% respectively (Langley 1993).

Position in the Cell Cycle

Using flow cytometry, Walle et al investigated the relationship of the cell cycle in PBMCs and immune status in renal allograft recipients. They reported that the 'activity index' (% of cells in G_1 x % of cells in SG_2M) was lowest in patients with stable function, intermediate in patients preceding a

rejection episode and high in those patients with infection (viral and bacterial). These differences were significant, and there was no overlap between the three groups, but the numbers of patients in each cohort were small (Walle 1984).

Cytoimmunomonitoring

Other groups have examined the usefulness of the relative frequency of activated lymphocytes, in peripheral blood, as a predictor of allograft rejection. Tanimura et al reported an 84% positive predictive value using flow cytometry in 30 renal transplant recipients. They measured spontaneous lymphocyte blastogenesis. Unfortunately there were no infective complications during the study, which has been reported to influence the results (Tanimura 1985). In a study in 54 heart transplant recipients, histological examination of a cytospin of PBMCs was compared to results from endomyocardial biopsies. This form of monitoring, termed cytoimmunomonitoring (CIM), gave a sensitivity and specificity of 88% and 86% respectively. Again, there are no results in patients with infective complications, but they do acknowledge that changes do occur during infection (Klanke 1987). Another study, in 16 cardiac transplant recipients using morphological inspection of PBMCs, concluded that this technique was not accurate enough to replace histological examination of biopsy material or that it could be used to guide a biopsy schedule (Wijngaard 1989). A later study, again studying spontaneous blastogenesis in renal transplant recipients, included patients with infective complications. This study shows a good correlation between increased spontaneous blastogenesis and acute rejection,

but also reported elevated counts in bacterial infection. The mean elevation was higher in the case of acute rejection, but there was considerable overlap of absolute values (de Boccardo 1994).

Caca and colleagues have compared flow cytometric analysis of the cell cycle in PBMCs with CIM. They reported an excellent correlation between the percentage of SG₂M cells and lymphoblasts, in 10 cardiac transplant recipients. They did not assess the technique's predictive values only the comparison between the two (Caca 1989). The same group has also shown that CIM may be of benefit in small bowel transplantation (Graeb 1992). In a more recent article, they claim good results from immunological monitoring using a combination of several techniques in both thoracic and liver transplant recipients. The tests they employed included CIM, CD4/CD8 ratios and the procalcitonin test. This later test allows for highly sensitive detection of bacterial and fungal infections and zoonoses (Hammer 1998).

Cross-Match

Some groups have used cross match techniques to monitor renal transplant recipients following transplantation. The rational of this approach is to determine whether the development of a positive cross match is associated with poorer graft outcome and survival. Piazza et al reported strong correlation between the subsequent development of a positive flow cytometric cross match (FCXM) and increased incidence of acute rejection, poor graft function and worse graft survival in 66 kidney recipients (Piazza 1999). Similar results, also using FCXM, were reported by Puig et al. They

performed FCXM on 78 cadaveric renal transplant recipients soon after receiving their kidneys. They reported that both acute rejection episodes and early graft loss correlated with a positive flow result (Puig 1995).

Panel Reactivity

Neylan et al examined the relationship of the percentage of panel reactive antibodies pre- and post-transplant in 64 renal transplant recipients. They reported a significant correlation between a post-transplant increase in the panel reactive antibodies of more than 10% and episodes of acute rejection. Patients with this increase had an 8x greater risk of rejection when compared to patients with no rise or a rise of less than 10% (Neylan 1997).

The problems with both the FCXM and panel reactivity studies relate to their retrospective nature and the use of infrequent monitoring. It would be interesting to know if the changes identified can be used to predict acute rejection or whether they result from the rejection process. If the later is true, they still may be of use in the management of these patients as they are potentially at risk of early graft loss.

Alloantigen Reactivity

DeBruyne et al used a limiting dilution analysis to monitor the frequency of alloantigen-reactive helper T lymphocytes in cardiac allograft recipients. Studying 12 patients, they identified that episodes of acute rejection are associated with an increase in the number of circulating donor-specific helper

T lymphocytes. This increase was frequently detected before the development of histologically defined rejection. Like so many methods of monitoring, absolute numbers were unhelpful, as there was considerable individual variation. Even when a percentage rise was examined, there was overlap between no rejection and mild rejection, and mild rejection and moderate rejection (DeBruyne 1993).

1.6 Previous Work from Our Laboratory in the Field of Immunological Monitoring

There is a considerable lack of agreement in the literature concerning the measurement of cytokines both at the time of rejection and preceding the diagnosis. (Discussed in section 1.4). This is true for both intragraft studies and serum studies in renal transplant recipients. In an attempt to clarify some of these issues, Tan et al monitored sequential changes in both Th1 and Th2 cytokine mRNA expression in T lymphocytes obtained from peripheral blood in a series of 43 renal transplant recipients (Tan 2001). Blood samples were obtained from all patients pre-transplant and then regularly within the first four weeks thereafter. If a patient experienced an episode of biopsy proven acute allograft rejection additional blood samples were obtained prior to the initiation of anti-rejection therapy and after completion of the treatment, up to 6 weeks post-transplant.

Mononuclear cells were extracted by density gradient centrifugation, the T cells were negatively selected using monoclonal antibodies and the RNA extracted using RNeasy. The extracted RNA was quantitated and standardised prior to reverse transcription to cDNA. An RT-PCR technique was combined with an ELISA method of detecting digoxigenin-labelled PCR products. This was used to semi-quantitatively detect sequential changes in expression of 6 cytokines (IL-2, IFN- γ , IL-4, IL-5, IL-10 and IL-13).

The two Th1 cytokines (IL-2 and IFN- γ) had similar patterns of expression. In the non-rejectors, expression of either cytokine failed to significantly change

from the baseline value. However, in the rejectors IL-2 expression decreased significantly on day 2 and this was maintained at a significantly lower level at the time the rejection was diagnosed and in the early post-treatment phase, but had reverted to a baseline level by a week post-treatment. IFN- γ also showed significantly lower expression at day 2 and in the early post-treatment phase, but not at the time of rejection diagnosis.

IL-5 expression was significantly lower than baseline at all time points following transplantation in the non-rejectors, but a relative rise in expression was detected by a return to baseline levels at the time of rejection, in the rejector group. Following treatment, the expression again significantly fell below baseline levels, reverting to a non-rejector pattern.

IL-13 expression did not significantly alter in expression following transplantation in the non-rejector group, but again rose at the time of rejection. This time significantly above baseline expression. This elevated levels returned to baseline following treatment and remained there.

IL-4 expression was significantly reduced in non-rejectors for the first week following transplantation, but had reverted to baseline levels by the week 2 sample. However, in rejectors the return to baseline was seen earlier, at the time of rejection. Levels fell again following treatment, but were at baseline levels at a week post-treatment.

IL-10 was the only cytokine to have significantly elevated levels of expression following transplantation. This was maintained at all time points in the non-rejector group, but at the time of rejection there was a relative fall in expression with levels comparable to baseline. Following treatment, expression levels significantly rose again, but this increase was only seen in the early post-treatment phase. Thereafter levels were not significantly different from baseline expression.

All of the significant changes described above were shown to be statistically significant at $p=0.05$ using the Wilcoxon rank signed test.

Sequential monitoring of peripheral T cell cytokine gene expression using a sensitive RT-PCR ELISA detection method revealed significant changes in cytokine gene expression. These results were intriguing, and constitute a first attempt to detect and analyse significant changes in peripheral cytokine gene expression following renal transplantation, which correlate with clinical events, including episodes of biopsy proven acute rejection. However, a number of important questions still need to be addressed. For example, The RT-PCR ELISA system used, while highly sensitive, is at best semi-quantitative. A quantitative methodology is therefore required to substantiate these findings. Also, the influence of major surgery (rather than transplantation per se) on cytokine gene expression was not investigated. This should be investigated in a non-transplant population. Finally, with a more sophisticated detection system it may be possible to use additional standardisation techniques in the form of measurement of the expression of

one or more house keeping genes, against which to compare cytokine gene expression.

1.7 TaqMan Real Time PCR Detection

The previous work from our laboratory on immunological monitoring has been described above (section 1.6). One of the main criticisms that may be directed towards this study was the method of measurement of the cytokine expression, namely RT-PCR ELISA. This is a highly specific and extremely sensitive technique, and at the time of the study was judged to be the optimum technique available. There are disadvantages in that it is technically demanding, it requires extensive post-PCR manipulations that add time to the analysis and may lead to laboratory contamination, and, most importantly, it measures product accumulation after a fixed number of cycles, selected to achieve maximal product levels.

The early cycles of PCR are characterised by an exponential increase in target amplification. As reaction components become limiting, the rate of target amplification decreases until a plateau is reached and there is little or no net increase in PCR product. Therefore, the amount of PCR product observed at the end of the reaction is very sensitive to slight variations in reaction components. This is because endpoint measurements are generally made when the reaction is beyond the exponential phase and a slight difference in a limiting component can have a drastic effect on the final amount of product, but will only affect the start of exponential product accumulation when the difference is large. More importantly, extensive experiments in many laboratories have now shown that widely differing amounts of template can generate equivalent amounts of PCR product when the plateau phase is

reached (Heid 1996). Therefore, there are potential advantages to measurement of PCR product accumulation during the exponential phase of the reaction, as the result will more accurately reflect the amount of starting material.

Higuchi et al pioneered the analysis of PCR kinetics by constructing a system that detected PCR product as it accumulated (Higuchi 1992, Higuchi 1993). 'Real-time' systems were later improved by probe-based, rather than intercalator-based, PCR product detection. The 5' nuclease assay provided a real-time method for detecting only specific amplification products. Holland et al were the first to demonstrate that cleavage of a target probe during a PCR could be used to detect amplification of the target-specific product. The technique used a probe labelled with ³²Phosphor on its 5' end and blocked on its 3' end so it could not act as a primer. During amplification, annealing of the probe to the target sequence generates a substrate that is cleaved by the 5' nuclease activity of Taq DNA polymerase when the enzyme extends from an upstream primer into the region of the probe. This dependence on polymerisation ensures that cleavage of the probe occurs only if the target sequence is being amplified. After PCR, cleavage of the probe was measured by using thin layer chromatography to separate cleavage products from intact probe (Holland 1991).

The development of fluorogenic probes made it possible to eliminate post-PCR processing for the analysis of probe degradation. The probe is an oligonucleotide with a reporter dye and a quencher dye attached. While the

probe is intact, the proximity of the quencher greatly reduces the fluorescence emitted by the reporter dye by Förster resonance energy transfer through space (Förster 1948). (See section 2.4 for more details). Several different reporter dyes are available allowing more than one target to be detected in a single PCR well.

The Applied Biosystems division of Perkin Elmer (Foster City, USA), developed the real-time PCR principle into the commercially available system called TaqMan.

The TaqMan PCR method offers several advantages over RT-PCR ELISA. Firstly, it is performed in a closed-tube system and requires no post-PCR manipulation of the sample. This not only reduces the risk of laboratory contamination, but also reduces the possibility for technical error and reduces the time required to perform an analysis. Secondly, the TaqMan system has universal PCR conditions. This enables multiple products to be amplified on the same 96-well plate, which increases productivity. Thirdly, the TaqMan PCR method is highly reproducible. Replicate amplifications can be analysed for each sample to identify a sub-optimal reaction. The sample can then be repeated if necessary. However, as can be seen from figure 1.7 there is superb correlation between the replicates and reaction failures are very uncommon. Fourthly, the system allows for a very large assay dynamic range (approaching 1 000 000-fold starting target). Finally, TaqMan detects PCR product accumulation during every cycle. The fluorescent threshold values (C_T), the point at which the reaction enters its exponential growth phase,

correlate linearly with the relative number of DNA or cDNA copies that were placed in the reaction tube (Heid 1996).

Figure 1.7. Scattergram showing the correlation between TaqMan duplicates

This is a scattergram demonstrating the correlation between replicate samples of cDNA analysed using commercially available TaqMan primers and probe for human β -actin. Pearson's correlation coefficient = 0.997. $p < 0.001$. (Data from this study).

BADP1 = β -actin duplicate 1

BADP2 = β -actin duplicate 2

1.8 Aims of the Study

These were three fold and are listed below.

1. To further study the use of intensive peripheral blood mononuclear cell cytokine gene expression measurement, using a fully quantitative technique, for the purposes of immunological monitoring in renal transplant recipients, concentrating on the cytokines IL-4, IL-10, TNF- α and TGF- β 1.
2. To design our own TaqMan primers and probes for the cytokines under investigation and confirm their efficacy by comparing the results with obtained using commercially available products. Then to use these in house designed primers and probes to substantiate the results of the earlier RT-PCR ELISA study using the fully quantitative TaqMan system.
3. To explore the possibility of using one or more house keeping genes in an attempt to allow further standardisation within the methodology.

Chapter 2: Patients, Methods and Materials

This chapter will describe, in detail, the patient recruitment, the laboratory techniques, equipment and consumables used during this study. However, as several of the steps used are routine laboratory procedures, the equipment and consumables for each of these stages are listed in an appendix. The methodology for the TaqMan cytokine gene expression analysis is novel, and so the consumables and equipment for this stage is given within the text.

2.1 Patient Recruitment and Rejection

Diagnosis

All patients admitted for a renal transplant or living donor nephrectomy to the Wessex Renal and Transplant Unit between June 1999 and July 2000 were approached about participating in this study. All patients were given a written information sheet to read at their leisure on admission to the ward (see Appendix I). If they were interested in taking part in the study, they were given the opportunity to ask any addition questions and have any queries answered by the author. If they agreed to participate in the study, written consent was obtained (see Appendix II) before a blood sample was taken. Data on all patients was collected prospectively on a data collection proforma (see Appendix III).

Daily blood samples were taken in order to measure the serum creatinine of all transplant recipients. If a patient had a rise in their serum creatinine of more than 20mmol/l on two successive days they underwent further investigation to identify the cause of their renal transplant dysfunction. This involved a urine culture to exclude infection and an ultrasound scan to exclude obstruction. Blood was also taken to monitor CyA levels. If no other explanation was found, then the patient underwent an ultrasound guided renal transplant core biopsy to detect acute allograft rejection. This was examined histologically and reported the same day.

Renal transplants experiencing acute tubular necrosis, diagnosed by the failure of the serum creatinine to fall at 24hours post-transplant in a well perfused kidney on duplex Doppler scanning, underwent an ultrasound guided renal transplant core biopsy to detect acute allograft rejection every five days until renal function recovered.

In this study, the diagnosis of acute allograft rejection was biopsy proven in all cases.

2.2 Cell Separation

This was the first stage of the laboratory work-up, and was performed in the Transplant Laboratory, St. Mary's Hospital, Portsmouth. The method used produced a high yield of peripheral blood mononuclear cells, without the need to obtain large quantities of blood from the patient (Böyum 1968). It was a quick process, only requiring an hour, which minimized the amount of cell manipulation prior to extraction of total RNA.

See appendix IV for consumables and equipment.

2.2.1 Blood Collection

The blood samples were collected prior to transplantation, usually at the time the samples were taken for cytotoxic cross-matching in the case of the cadaveric recipients or on the morning of surgery for the living donors and recipients, and then each morning between 8am and 9am. This was in order to exclude any diurnal variation in cytokine mRNA expression. The samples were taken each morning for the first 14 days post-transplant or until hospital discharge for the recipients and for the first 7 post-operative days or until hospital discharge for the donors, depending on which was the shorter. A total of 9ml of blood were taken on each occasion, in two 4.5ml pull sodium citrate 'vacutainer' bottles.

2.2.2 Cell Separation

Within two hours of the being obtained from the patients, the blood samples were placed in 50ml-skirted polypropylene test tube, and diluted up to a total volume of 22ml using single strength (1x) phosphate buffered saline (PBS). Each diluted sample was then divided into two aliquots of 11ml. Each aliquot was carefully layered on to 4ml of graduated density medium (Lymphoprep – density 1.077g/l, Nycomed, UK) in a 15ml conical polypropylene test tube. These were then spun in a tipping bucket centrifuge at 1100g for 20 minutes at room temperature (20°C), no additional acceleration or braking rates were added.

When the test tubes were removed from the centrifuge, the blood samples had clearly separated into four distinct layers. From top to bottom, these were the straw-coloured plasma, the white ‘buffy layer’, the clear density medium and finally the burgundy red cell and granulocyte layer. The buffy layer, which contains the mononuclear white cells, was extracted in approximately 2.5ml of serum using a Pasteur pipette and placed in another 50ml-skirted polypropylene test tube. The total sample of 5ml was washed with a further 12.5ml of 1x PBS. The suspension was spun again in order to pellet the mononuclear cells, at 400g for 10 minutes, again at 20°C. On that occasion maximum brake and acceleration rates were applied.

When the tube was removed the cell pellet could clearly be seen as a small yellow disc at the base. The serum/PBS was extracted using a 10ml

serological pipette, being careful to avoid the pellet, which remained in 200-300µl of fluid. The cells were re-suspended in this small volume by vortexing the test tubes.

The RNA extraction was performed using a commercially available product, namely RNazolB (Biogenesis, UK). This functions by promoting the formation of complexes of RNA with guanidinium and water molecules, and abolishes hydrophilic interactions of DNA and proteins. The DNA and proteins are efficiently removed from the aqueous phase, while RNA remains (Chomczynski 1987).

Homogenising the sample, by the addition of 1ml of RNazolB, commenced RNA extraction. This mixture was placed in a 1.5ml eppendorf tube and mixed thoroughly by vortexing. The homogenate was transferred to a -20°C freezer for temporary storage.

2.3 Total RNA Extraction, Quantitation and Standardisation.

The remainder of the laboratory work was performed at Southampton General Hospital, in the Wessex Immunology Laboratory.

Total RNA extraction was performed using a single step technique of guanidium thiocyanate-phenol-chloroform extraction. This method produced high yields of good quality undegraded RNA (Chomczynski 1987).

See appendix V for consumables and equipment.

2.3.1 Total RNA Extraction

The homogenised samples were transported from Portsmouth to Southampton at 4°C and placed on ice on arrival at the Immunology Laboratory to prevent RNA degradation.

While the centrifuge was cooled to 4°C, 130µl of chloroform was added to each eppendorf containing the RNeasy homogenate to deproteinize the solution. This solution was then shaken vigorously for approximately 30 seconds and left to stand in ice for an additional 5 minutes. The samples were then spun at 1200g for 15 minutes at 4°C.

Following centrifugation the samples were placed on ice. At this stage, the samples had separated into three distinct layers. The lower purple-coloured phase was the chloroform-phenol layer while the upper clear aqueous layer contained RNA. The third thin, white layer was at the inter-phase between the other two, and contained protein and DNA.

The upper phase (usually about 600µl), containing RNA, was carefully transferred to another sterile 1.5ml eppendorf containing 700µl of isopropanol cooled to approximately 4°C, in order to precipitate the RNA out of solution. Each sample was vortexed and left to stand on ice, in a refrigerator, overnight for a minimum of 18 hours. This step not only increased RNA yield. It also, for logistical reasons, allowed enough time for a TaqMan PCR run to be performed on both days RNA extraction was being performed.

The following day the centrifuge was again cooled to 4°C, and the samples spun at 1200g for 15 minutes at 4°C in order to pellet the RNA that had precipitated out of solution. The eppendorfs were placed on ice when removed from the centrifuge. The clear supernatant was extracted using a glass Pasteur pipette and discarded. Care was taken to avoid the RNA pellet that appeared as a white disc at the tip of the eppendorf.

The RNA pellet was then washed with 500µl of 75% ethanol, which had been stored at -20° C. This step, removes water-soluble ions. The solution was then re-centrifuged at 7500g for 8 minutes at 4°C in order to precipitate the RNA pellet again. The ethanol was then removed to dryness using another glass

Pasteur pipette, taking care not to disturb the pellet. Each eppendorf was then placed in a petri dish and transferred to a 45°C warming cabinet to completely dry the pellet. This usually took 10-15 minutes.

When completely dry the pellet becomes transparent, confirming complete evaporation of the ethanol. The pellet was then dissolved in 30µl of 0.1% diethyl pyrocarbonate (DEPC) treated double-distilled water. (DEPC is an inhibitor of ribonucleases). This was hastened by placing the eppendorfs in a 55°C water bath for 5 minutes. The eppendorf tubes were placed on ice on removal from the water bath.

To facilitate RNA quantitation, a 6µl aliquot of the RNA solution was added to a further 12µl of DEPC treated RNase free water in a 0.75ml eppendorf to give a one-third strength solution. This was again kept on ice prior to quantitation.

2.3.2 Total RNA Quantitation

The spectrophotometer used for quantitation was a GeneQuant (Pharmacia, UK). This was switched on and set up for RNA quantitation as follows:

Path length	=	5
Use 320nm?	=	Yes
RNA factor	=	40.0
Bases number	=	0
Oligo length	=	1
Mw calc	=	0.0
Ratio expected	=	2.000
Conc expected	=	0.000
Protein coeff.1	=	1.550
Protein coeff.2	=	0.760

Before the ultramicrovolume cell was used, it was sequentially washed with 0.1M solutions of sodium hydroxide and hydrochloric acid and then sequentially rinsed with ultra-high quality (UHQ) water (prepared by sterilising double distilled water), and DEPC-treated water. Excess liquid was flicked out of the cell before the next was used. Once the wash was completed the cell was dried with tissue paper and aspirated to dryness using a 10 μ l pipette.

A 'no RNA' reference was placed in the GeneQuant before use. This consisted of a 6 μ l aliquot of the same DEPC-treated water in which the RNA samples were dissolved. A similar volume of each one-third strength RNA sample was used for quantitation. The measurements at 260nm and 280nm were recorded, as was the 260/280 ratio. The 6 μ l aliquot measured and the

remaining third strength solution were discarded after use. Between each measurement the cell was thoroughly washed with both UHQ water and DEPC-treated water, dried with tissue and then aspirated to dryness. A 260/280 ratio of 1.9 was expected for good quality RNA.

If the 260 reading was '>3.000', then the concentration of the third strength solution was too high for the GeneQuant to measure. In this situation the 6 μ l aliquot measured was discarded, as above, and the cell rinsed. A further 12 μ l of DEPC water was added to the remaining 12 μ l of third strength solution therefore making it a one-sixth strength solution. A further 6 μ l aliquot of this latter RNA solution was again measured. Rarely this solution was still too concentrated. The measured 6 μ l was again discarded. This time a further 9 μ l of DEPC treated water was added to the one-sixth strength solution, which was then a one-ninth strength solution. Another 6 μ l aliquot was measured once the cell had been thoroughly rinsed again.

2.3.3 Total RNA Standardisation

Once the RNA quantitation had been performed, it was possible to calculate the total amount of total RNA present in the remaining stock solution. This was calculated by multiplying the 260nm absorbance by the 'RNA factor', the dilution factor of the measured solution and the volume of the remaining stock solution. The volume required to give a concentration of 125ng/ μ l was calculated. The volume added to the stock solution was the calculated volume minus the volume of the stock solution that remained.

A concentration of 125ng/μl of RNA produced aliquots of 1μg of total RNA in 8μl volumes. It was normally possible to obtain at least four such aliquots from each patient sample.

2.4 First-strand cDNA Synthesis

This was accomplished using a commercial kit (Pharmacia Biotech, UK), which utilises Moloney Murine Leukaemia Virus reverse transcriptase to catalyse the reaction. All reagents necessary to generate full-length first-strand cDNA from RNA templates 7kB or more in length were provided within the kit.

See appendix VI for consumables and equipment.

2.4.1 First-strand cDNA Synthesis

Following standardisation the four 8µl aliquots were combined in a single 0.75ml eppendorf and an additional 4µl of DEPC water was added to allow for evaporation losses at the time of denaturation. The sample was then denatured for 10min. at 65°C in a Hybaid thermal reactor before again being chilled on ice for another 5-10min.

While the sample was being heated, the cDNA reaction tubes were made up. Each previously labelled 0.75ml eppendorf tube (sterile and UV irradiated) contained 5µl of bulk first-strand reaction mix, 1µl of DTT and 1µl of dilute *Not-I*-d(T)₁₈ primer. To each tube was added 8µl of the chilled previously denatured standardised RNA solution. The 15µl mixture was thoroughly mixed before being placed in a 37°C water bath and incubated for 1 hour.

After this hour the 4 tubes of cDNA from each patient were combined and then were placed in a -70°C freezer to await subsequent molecular analysis.

2.5 TaqMan Cytokine Gene Expression

Analysis

PE Applied Biosystems developed the TaqMan system as a research tool for 'real time', *in vitro* quantitative evaluation of Polymerase Chain Reaction (PCR) products.

The PCR reaction exploits the 5' nuclease activity of the AmpliTaq Gold™ DNA Polymerase to cleave a TaqMan probe during PCR. A reporter dye [FAM (i.e. 6-carboxyfluorescein)] is attached at the 5' end of the TaqMan probe and a quencher dye [TAMRA (i.e. 6-carboxy-tetramethylrhodamine)] at the 3' end of the probe (See figure 2.5a).

When the probe is intact, the proximity of the reporter to the quencher dye results in suppression of the fluorescence of the reporter dye, primarily by Förster-type energy transfer (Förster 1948, Lakowicz, 1983).

During the PCR, cleavage of the probe separates the reporter dye and the quencher dye (see figure 2.5b), which results in increased fluorescence of the reporter (see figure 2.5c). The ABI Prism 7700 Sequence Detector (PE Applied Biosystems, UK) has a built in laser directed, via fibre optics, to each of the 96 wells. The fluorescence emission travels back through the cables to a charge-coupled device (CCD) camera detector. For each sample, the CCD camera collects the emission data between 520nm and 660nm once every few

seconds. Accumulation of PCR product is detected directly by monitoring the increase in fluorescence of the reporter dye throughout the amplification (see figure 2.5d).

During the PCR, if the target of interest is present, the probe specifically anneals between the forward and reverse primer sites. The 5'-3' nucleolytic activity of the AmpliTaq Gold DNA Polymerase cleaves the probe between the reporter and the quencher only if the probe hybridises to the target, and the polymerisation of the strand continues. The 3' end of the probe is blocked to prevent extension of the probe during the PCR. This process occurs in every cycle and does not interfere with the exponential accumulation of the product.

The increase in fluorescence signal is detected only if the target sequence is complementary to the probe and is amplified during the PCR. Therefore, any non-specific amplification is not detected.

There are several different reporter dyes that can be attached to the probes used for TaqMan PCR. The commercial primers/probe mixes produced by Perkin-Elmer for investigations of cytokine gene expression are labelled with the FAM dye. For this reason the probes that were designed for this study were also labelled with this dye. Perkin-Elmer also produced commercially available primers/probe mixes for several 'house-keeping' genes. These are labelled with a different dye, VIC, to enable 'in tube' standardisation if required.

Figure 2.5a.

Polymerisation

This figure is a diagrammatic representation of the structure of the probe with its reporter and quencher dyes, at the start of polymerisation. R = reporter dye.
Q = quencher dye.

Figure 2.5b.

Strand Displacement

This figure shows progression of the reaction with displacement of the probe.

Figure 2.5c.

Cleavage

This figure shows that cleavage of the probe separates the reporter and quencher dyes resulting in increased fluorescence of the reporter dye.

Figure 2.5d.

Polymerisation Complete

This figure shows that the reaction is complete with total separation of the probe.

Equipment: ABI Prism 7700 Sequence Detector [PE Applied Biosystems, UK] previously calibrated using Sequence Detection Systems Spectral Calibration Kit [PE Applied Biosystems, UK]
Pipettes of various ranges [Biohit, Alpha Laboratories, UK]

Consumables: MicroAmp Optical 96-Well Reaction Plate/Optical Caps [PE Applied Biosystems, UK]
Sterile pipette tips of various sizes [Greiner Labortechnik, UK]
Sterile 1.5 and 0.75ml eppendorfs [Greiner Labortechnik, UK]
TaqMan Universal PCR Master Mix, containing AmpliTaq Gold DNA Polymerase, AmpErase UNG, dNTPs with dUTP, Passive Reference and optimized buffer components. [PE Applied Biosystems, UK]
IL-4 primers/probe. [PE Applied Biosystems, UK]
IL-10 primers/probe. [PE Applied Biosystems, UK]
Human beta-actin primers/probe. [PE Applied Biosystems, UK]
Beta-2 microglobulin primers/probe. [PE Applied Biosystems, UK]
Transferrin receptor primers/probe. [PE Applied Biosystems, UK]

Biosystems, UK]

DEPC-treated RNase free water

Oligonucleotide Sequences

In addition to the commercially available primers and probes, the following primers and probes (see below) were designed, as part of the project, using PE Primer Express software [PE Applied Biosystems, UK]. This program allows the user to design TaqMan primers and probes for optimal PCR amplification and probe incorporation under a standard set of PCR conditions. The cDNA sequence of the target gene is imported into the PE Primer Express programme. The exon/exon boundaries are then highlighted as the software is designed to incorporate at least one boundary in either the primers or, ideally, the probe to minimise the risk of amplifying genomic DNA. The software produces a range of possible primers and probes that fulfil pre-defined rules. For both the probe and the primers these rules are that the percentage of Gs and Cs should be in the range 20-80% and there should be no runs of more than three consecutive Gs. For the probe only the rules are that firstly, the amplicon size is between 50-150 base pairs. Secondly, no G on the 5' end and thirdly, a T_m of between 65-67°C. For the primers only the rules are firstly, the two primers should be as close to the probe as possible without overlapping it and secondly, all primers have a T_m of 58-60°C. In the case of the probe, it is necessary to manually check that there are more Cs than Gs. In the case of the primers, it is desirable to have only 1-2 G+Cs in the five nucleotides at the 3' end.

For each set of primers and probe below, two bases either side of the exon/exon boundaries are highlighted.

IL-4	Sense Primer	GTGCACCGAGTTGACCGTAA
		[Oswell DNA Services, UK]
IL-4	Anti-sense Primer	TGTAGAACTGCCGGAGCACA
		[Oswell DNA Services, UK]
IL-4	Fam-labelled Probe	CTTTGCTGCCTCCAAGAACACAACTGAGA
		[PE Applied Biosystems, UK]
IL-10	Sense Primer	AGGCGCATGTGAACTCCCT
		[Oswell DNA Services, UK]
IL-10	Anti-sense Primer	CACGGCCTTGCTCTTGT
		[Oswell DNA Services, UK]
IL-10	Fam-labelled Probe	AGGCTACGGCGCTGTCATCGATTCT
		[PE Applied Biosystems, UK]
TNF α	Sense Primer	CCAGGGACCTCTCTCTAATCAGC
		[Oswell DNA Services, UK]
TNF α	Anti-sense Primer	GCTTGAGGGTTTGCTACAACATG
		[Oswell DNA Services, UK]
TNF α	Fam-labelled Probe	CAGTCAGATCATCTTCTCGAACCCCGAG
		[PE Applied Biosystems, UK]
TGF β 1	Sense Primer	CGCGTGCTAATGGTGGA
		[Oswell DNA Services, UK]
TGF β 1	Anti-sense Primer	GCTTCTCGGAGCTCTGATGTG
		[Oswell DNA Services, UK]
TGF β 1	Fam-labelled Probe	CCACACGAAATCTATGACAAGTTCAAGCAGAGTA
		[PE Applied Biosystems, UK]

2.5.1 TaqMan Cytokine Gene Expression Analysis

The conditions for every PCR reaction are universal (see table 2.5a). This allowed several different genes to be analysed, for each patient, on the same 96-well plate simultaneously. To reduce the error of a failed or sub-optimal PCR reaction, each sample was run in duplicate. The series was repeated, for that gene, if there was a discrepancy of more than one cycle between the two samples. The mean of the two was used for subsequent statistical analysis.

A TaqMan PCR master-mix was made up for each gene to be analysed. The individual components and volumes for a single sample are shown in table 2.5b. The concentrations of the primers and probe for the stock solution and the working solution are shown. The concentration of the stock, and hence, working solutions were not chosen randomly. Colleagues, within our institution, with previous experience of the TaqMan system recommended them. As these concentrations worked well with our primers and probes, we did not feel it necessary to experiment with other concentrations.

For speed and convenience, when more than one sample was being analysed, the volume of each component being added was multiplied by $n+1$, where n was the number of samples to be analysed and was less than 15, and $n+2$ when n was greater than 15. This allowed for some volume loss while pipetting.

1µl of cDNA from each sample was pipetted into a labelled sterile 0.75ml eppendorf. To this was carefully added 49µl of master-mix. This was thoroughly mixed by pipetting. The reaction mix was now complete. Two 25µl aliquots of each reaction mix were carefully transferred to adjacent wells of a MicroAmp Optical 96-Well Reaction Plate and cover with Optical Caps. The plate was then transferred to the ABI Prism 7700 Sequence Detector.

Before each PCR run, the thermal cycling conditions and reaction volume were checked, on the ABI Prism 7700 instrument, to confirm that the settings shown in table 2.5a had not been altered and the reaction volume was set for 25µl. The run could then be started.

When the run was completed, the data generated was saved and the output data exported into a Microsoft Excel file for later analysis. (The individual steps taken to initiate a run and to acquire the output data are described, in full, in appendix VII).

Table 2.5a. Standard TaqMan PCR conditions

Stage 1 (Hold)	Stage 2 (Hold)	Stage 3 (40 cycles)	
2min at 50°C	10min at 95°C	15s at 95°C	60s at 60°C

This table shows the standard TaqMan PCR. min = minute. °C = degrees

Celsius. s = seconds.

Table 2.5b. Components of the TaqMan master mixes

Components and Stock Concentration	Working Concentration	In House Designed	Commercial
TaqMan Universal PCR Master Mix (2x)	1x	25.0µl	25.0µl
RNAse Free Water	-	21.0µl	21.5µl
5' Primer (15µM)	300ηM	1.0µl	-
3' Primer (15µM)	300ηM	1.0µl	-
Probe (5µM)	100ηM	1.0µl	-
PE Commercial Primer/Probe Mix (20x)	1x	-	2.5µl
Total Volume	-	49.0µl	49.0µl

This table shows the components of the different master-mixes depending on whether the in house designed primers/probe or the commercial primers/probe were being used.

Chapter 3: The Substantiation of the Results of the RT-PCR ELISA Gene Expression Work by Using the TaqMan System

One of the limitations of the work previously undertaken in our laboratory was that the method of detection of cytokine gene expression was semi-quantitative, rather than fully quantitative. The results may be artefact, influenced by the RT-PCR ELISA system, rather than actual changes in expression even though the changes detected correlated with clinical events. It was, therefore, crucial to substantiate the results using the fully quantitative real time TaqMan gene detection system before progressing with this study.

This chapter will describe the initial experiments performed using TaqMan on stored cDNA samples collected in the previous RT-PCR ELISA study.

3.1 Comparison of commercial primers/probes with designed primers/probes

At the time this study commenced there were commercially available multiple cytokine gene expression plates and also pre-designed primers and probes for several individual cytokines. These reagents were extremely costly when used on the large scale required for our purposes. To overcome this problem, Primer Express software was used to design primers and probes for all four cytokines to be investigated (described in section 2.5). Having designed our own primers and probes, the first stage was to confirm their efficacy by comparing them with the commercial products. The primers and probes that were compared were those for IL-4 and IL-10 detection. These cytokines were selected as they showed the greatest difference in expression between the rejectors and non-rejectors in our previous work (see section 1.6). Stored cDNA from 18 patients from the earlier RT-PCR ELISA study (14 non-rejectors and 4 rejectors) were re-examined. All samples from the same patient series were analysed together on the same plate for a given cytokine. For each method (commercial and designed) standard TaqMan conditions were used as described in the previous chapter (section 2.4). The C_T value was used for data interpretation analysis.

The results will be presented with representative graphs, for visual comparison, from both rejectors and non-rejectors for both cytokines and both sets of primers and probes. The X-axis represents the time points. These are summarised in table 3.1a. The Y-axis represents the C_T value. Each sample was analysed in duplicate, hence two sets of results on each graph.

Figures 3.1a-h are a graphical representation of the results of 8 TaqMan analyses. They are the analysis of the commercial and 'in-house' designed primers of one non-rejector for both IL-4 and IL-10 and one rejector again for IL-4 and IL-10.

Table 3.1a. Time points used in RT-PCR ELISA study

Time point	Non-Rejectors	Rejectors
PT	Pre-transplant	Pre-transplant
2	2 days post-transplant	2 days post-transplant
3	1 week post-transplant	1 week post-transplant
4	2 weeks post-transplant	2 weeks post-transplant
5	4 weeks post-transplant	2 days after anti-rejection therapy
6	-	1 week after anti-rejection therapy
7	-	2 weeks after anti-rejection therapy

This table shows the time points and the code number they were given in the RT-PCR ELISA study.

Figure 3.1a. Graph showing a TaqMan result of a non-rejector using in house designed IL-4 primers and probe

This graph represents the results from the analysis of a non-rejector for the in house designed IL-4 primers and probe.

Figure 3.1b. Graph showing a TaqMan result of a non-rejector using commercially designed IL-4 primers and probe

This graph represents the results from the analysis of the same non-rejector for the commercially designed IL-4 primers and probe.

Figure 3.1c. Graph showing a TaqMan result of a non-rejector using in house designed IL-10 primers and probe

This graph represents the results from the analysis of a non-rejector for the in house designed IL-10 primers and probe.

Figure 3.1d. Graph showing a TaqMan result of a non-rejector using commercially designed IL-10 primers and probe

This graph represents the results from the analysis of the same non-rejector for the commercially designed IL-10 primers and probe.

Figure 3.1e. Graph showing a TaqMan result of a rejector using in house designed IL-4 primers and probe

This graph represents the results of the analysis of a rejector for the in house designed IL-4 primers and probe.

Figure 3.1f. Graph showing a TaqMan result of a rejector using commercially designed IL-4 primers and probe

This graph represents the results of the analysis of the same rejector for the commercially designed IL-4 primers and probes.

Figure 3.1g. Graph showing a TaqMan result of a rejector using in house designed IL-10 primers and probe

This graph represents the results of the analysis of a rejector for the in house designed IL-10 primers and probe.

Figure 3.1h. Graph showing a TaqMan result of a rejector using commercially designed IL-10 primers and probe

This graph represents the results of the analysis of the same rejector for the commercially designed IL-10 primers and probes.

In both rejectors and non-rejectors, demonstrated in the graphs above, there appeared to be an excellent correlation between the designed primers and probes and the commercially available primers and probes for both IL-4 and IL-10. This correlation is reiterated in figure 3.1i. This is a scattergram in which the mean value of each designed primers and probe value is plotted against the corresponding mean value of the commercial primers and probe.

Figure 3.1i. Scattergram showing the correlation between in house and commercially designed IL-4 and IL-10 primers and probes

This scattergram demonstrates the correlation between the mean C_T value for the in house designed primers and probe (plotted on the X-axis) and the mean C_T value for the commercially designed primers and probe (plotted on the Y-axis) for the detection of both IL-4 (green) and IL-10 (red). The numerical value on each axis corresponds to the C_T value.

IL4_AV = mean value for the in house designed IL-4 primers and probe.

IL4_C_AV = mean value for the commercially designed IL-4 primers and probe.

IL10_AV = mean value for in house designed IL-10 primers and probe.

IL10_C_AV = mean value for the commercially designed IL-10 primers and probe.

Table 3.1b. Correlation between in house and commercially designed IL-4 and IL-10 primers and probes

		Commercial IL-4	Commercial IL-10
Designed IL-4	Pearson's correlation coefficient	0.83	-
	Significance (2 tailed)	P<0.001	-
Designed IL-10	Pearson's correlation coefficient	-	0.76
	Significance (2 tailed)	-	P<0.001

The correlation that has been demonstrated graphically above between the in house designed primers and probes for IL-4 and IL-10 and those that had been commercially produced is confirmed statistically in this table. Using Pearson's correlation test, there is a highly significant correlation between the two sets of primers and probes.

The above graphics and statistical output data demonstrate the high degree of correlation between the in house designed primers and probes for IL-4 and IL-10 and those that were commercially produced for the same cytokines.

It can be seen that the commercial primers and probes were slightly more sensitive than the in house designed ones. The C_T value for the commercial primers and probe for both cytokines was 1-2 cycles lower than the designed primers and probes. This meant that it would not be possible to interchange the primers and probes, i.e. analyse some samples with the commercial primers and probe and others with the in house designed ones, as the C_T values would not be comparable. As the intention was to only use the in house designed primers and probes for all cytokines and as the C_T values for all time points for both cytokines were well within the measurable range, this was not a relevant observation for the purposes of this study. However, care would need to be taken when comparing data from different laboratories.

In the knowledge of these results, there was every justification to proceed to the next phase of the study using the in house designed primers and probes. Also, it was felt that the primers and probes for the cytokines TGF- β 1 and TNF- α , and other future cytokines could be safely designed using the Primer Express software without recourse to further confirmatory testing.

3.2 Re-analysis of the Existing cDNA Samples from the RT-PCR ELISA Work Using the TaqMan System

Having established the efficacy of our designed primers and probes, the next stage was to confirm the results of previous work from our laboratory investigating cytokine gene expression in 43 renal transplant recipients obtained with the RT-PCR ELISA. The stored cDNA samples were re-examined using TaqMan. Sufficient material was available for re-analysis of all but one subject. Therefore, 28 sample series of the non-rejectors and 14/15 of the rejectors were re-analysed using TaqMan and the in house designed primers and probes for IL-4 and IL-10. Each series was analysed in its entirety and each sample was analysed in duplicate. The mean of the C_T values was used for statistical analysis.

As TaqMan is fully quantitative and RT-PCR ELISA was thought to be semi-quantitative, it was thought inappropriate to directly compare one set of results to the other. However, as can be seen from the example in figure 3.2a, there was good visual comparability between the TaqMan results and those obtained using the RT-PCR ELISA system.

Figure 3.2a. Graph showing the similarity between TaqMan results and RT-PCR ELISA results

This is a graphical representation of the results obtained for IL-4 gene expression data from a non-rejector by both RT-PCR ELISA and TaqMan methods. The Y-axis on the left right is for the C_T value from the TaqMan analysis (both duplicates are shown). The Y-axis on the right is for the value from the RT-PCR ELISA plate reader results (only the mean is shown). The scale on the right is reversed as an increase in the C_T value represents a decrease in starting material. The X-axis represents the sequential time points 1 = pre-transplant: 2 = day 2 post-transplant: 3 = 1 week post-transplant: 4 = 2 weeks post-transplant: 5 = 4 weeks post-transplant.

IL4TM1 = IL-4 gene expression data of duplicate 1 obtained by TaqMan.

IL4TM2 = IL-4 gene expression data of duplicate 2 obtained by TaqMan.

IL4EL = mean IL-4 gene expression data obtained by RT-PCR ELISA.

In the previous study, the Wilcoxon Signed Ranks Test was used to compare the percentage change of each RT-PCR ELISA value of each time point with the value of the baseline reading, which were designated as 100% (Tan 2001). The percentage changes in the value was said to be significant if $p < 0.05$, and has been discussed earlier (see section 1.6).

As the RT-PCR ELISA system is thought to be semi-quantitative and the TaqMan system fully quantitative, it was inadvisable to directly compare the results obtained from the two different methods. It was, therefore necessary to compare the two sets of results indirectly. The data obtained from the TaqMan analysis of the cDNA from the previous study was also analysed using the Wilcoxon Signed Ranks Test. Each time point was, again, compared to the baseline (pre-transplant sample) value. If the same significant changes in cytokine gene expression were demonstrable using the TaqMan data, this would provide important validation of the findings of our earlier study with a quantitative methodology. In addition, such a finding would mean that results obtained from the use of TaqMan on additional patient series could be compared with findings obtained from our original investigation.

The results of the findings for both IL-4 and IL-10 for both techniques, and for rejectors and non-rejectors, are shown below (figures 3.2b-i) as box and whisker plots. In the case of the RT-PCR ELISA study, the Y-axis gives the absolute values from the spectrophotometer used to measure the colour change in each well and not the percentage change used for the statistical analysis. This value is proportional to the amount of starting material

measured. In the case of the TaqMan study, the Y-axis gives C_T value, which is inversely proportional to the amount of starting material measured. The X-axis represents the various time points on which the samples were collected. The results are also shown in tabular form (tables 3.2a and b).

Figure 3.2b. Box and whisker plot showing IL-4 gene expression data from the non-rejector cohort in the RT-PCR ELISA study

This box and whisker plot shows the IL-4 gene expression data for the non-rejectors using the original RT-PCR ELISA system. The Y-axis gives the values from the spectrophotometer readings, which are proportional to the amount of starting material. The sampling points are on the X-axis.

Figure 3.2c. Box and whisker plot showing IL-4 gene expression data from the non-rejector cohort in the RT-PCR ELISA study but analysed using TaqMan technology

This box and whisker plot shows the IL-4 gene expression data for the non-rejectors using the 'TaqMan' system. The Y-axis gives the C_T values, which are inversely proportional to the amount of starting material. The sampling points are on the X-axis.

Figure 3.2d. Box and whisker plot showing IL-10 gene expression data from the non-rejector cohort in the RT-PCR ELISA study

This box and whisker plot shows the IL-10 gene expression data for the non-rejectors using the original RT-PCR ELISA system. The Y-axis gives the values from the spectrophotometer readings, which are proportional to the amount of starting material. The sampling points are on the X-axis.

Figure 3.2e. Box and whisker plot showing IL-10 gene expression data from the non-rejector cohort in the RT-PCR ELISA study but analysed using TaqMan technology

This box and whisker plot shows the IL-10 gene expression data for the non-rejectors using the 'TaqMan' system. The Y-axis gives the C_T values, which are inversely proportional to the amount of starting material. The sampling points are on the X-axis.

Figure 3.2f. Box and whisker plot showing IL-4 gene expression data from the rejector cohort in the RT-PCR ELISA study

This box and whisker plot shows the IL-4 gene expression data for the rejectors using the original RT-PCR ELISA system. The Y-axis gives the values from the spectrophotometer readings, which are proportional to the amount of starting material. The sampling points are on the X-axis.

Figure 3.2g. Box and whisker plot showing IL-4 gene expression data from the rejector cohort in the RT-PCR ELISA study but analysed using TaqMan technology

This box and whisker plot shows the IL-4 gene expression data for the rejectors using the 'TaqMan' system. The Y-axis gives the C_T values, which are inversely proportional to the amount of starting material. The sampling points are on the X-axis.

Figure 3.2h. Box and whisker plot showing IL-10 gene expression data from the rejector cohort in the RT-PCR ELISA study

This box and whisker plot shows the IL-10 gene expression data for the rejectors using the original RT-PCR ELISA system. The Y-axis gives the values from the spectrophotometer readings, which are proportional to the amount of starting material. The sampling points are on the X-axis.

Figure 3.2i. Box and whisker plot showing IL-10 gene expression data from the rejector cohort in the RT-PCR ELISA study but analysed using TaqMan technology

This box and whisker plot shows the IL-10 gene expression data for the rejectors using the 'TaqMan' system. The Y-axis gives the C_T values, which are inversely proportional to the amount of starting material. The sampling points are on the X-axis.

Table 3.2a. Box and whisker plot showing IL-10 gene expression data from the rejector cohort in the RT-PCR ELISA study but analysed using TaqMan technology

Cytokines	Post-transplant time points			
	Day 2	1 week	2 weeks	4 weeks
IL-4 by RT-PCR ELISA	↓ expression p=0.001	↓ expression p=0.006	NS	NS
IL-4 by TaqMan	↓ expression p<0.001	↓ expression p<0.001	NS	NS
IL-10 by RT-PCR ELISA	↑ expression p=0.002	↑ expression p=0.008	↑ expression p=0.041	↑ expression p=0.001
IL-10 by TaqMan	↑ expression p<0.001	↑ expression p<0.001	↑ expression p=0.045	↑ expression p=0.008

This table shows the same results as the box and whiskers plots of the non-rejectors for both techniques and both cytokines. Arrows indicate the direction of the change in expression from the baseline value. The numbers given are the p values. NS = non-significant.

Table 3.2b. Results of the statistical analysis of IL-4 and IL-10 gene expression data from the rejector cohort from the RT-PCR ELISA study and after re-analysis with TaqMan technology

Cytokines	Post-transplant time points				
	Day 2	Pre-ART	Day 2 Post-ART	1 week Post-ART	4 weeks Post-ART
IL-4 by RT-PCR ELISA	↓ expression p=0.002	NS	↓ expression p=0.015	NS	NS
IL-4 by TaqMan	↓ expression p=0.003	NS	↓ expression p=0.016	NS	NS
IL-10 by RT-PCR ELISA	↑ expression p=0.004	NS	↑ expression p=0.005	NS	NS
IL-10 by TaqMan	↑ expression p=0.034	↑ expression p=0.019	↑ expression p=0.010	NS	NS

This table shows the same results as the box and whiskers plots of the rejectors for both techniques and both cytokines. Arrows indicate the direction of the change in expression from the baseline value. The numbers given are the p values. NS = non-significant.

3.3 Discussion

In section 1.6 the results of the previous immunological monitoring study from our laboratory were described. In this study, the gene expression of several cytokines was measured in peripheral T lymphocytes taken from renal transplant recipients prior to transplantation and then sequentially thereafter for up to four weeks. The measurement of the gene expression was performed using a semi-quantitative technique, namely a RT-PCR ELISA system. The results of the measurement of the gene expression for the cytokines IL-4 and IL-10 revealed distinct patterns for patients not experiencing an acute rejection episode compared with those that did. These cytokine gene expression profiles correlated with clinical events, i.e. the diagnosis of the rejection and subsequent treatment with anti-rejection therapy (0.5g IV methyl prednisolone given daily for three days).

One of the limitations of the study was that RT-PCR ELISA was perceived as a semi-quantitative system and, therefore, some of the changes in cytokine gene expression might have resulted from inaccuracies and inconsistencies within the post-PCR processing required in the RT-PCR ELISA.

Before embarking on a further study investigating the potential role of the measurement of cytokine gene expression from peripheral blood cells in renal transplant recipients, it was necessary to confirm the findings of the RT-PCR ELISA work. Therefore, the cDNA samples from the previous work were re-analysed using the fully quantitative real time TaqMan PCR system. This

system uses target specific fluorogenic labelled probes to measure the amount of PCR product during each cycle. The data collected make it possible to identify the point at which the PCR reaction in each well enters the exponential phase of amplification. A numerical value (between 0 and 40) can be attributed to this point which is directly related to the amount of starting material placed in the PCR reaction mix in each well. There are more detailed descriptions of this technique in sections 1.7 and 2.5.

Figure 3.2a is representative of the results of the IL-4 gene expression measurement of a non-rejecting patient using both the RT-PCR ELISA and the TaqMan system. There is an obvious visual similarity between the profiles of the gene expression detected by the two techniques. A similar picture was observed in the data obtained by both techniques for IL-4 gene expression in the rejectors and for IL-10 gene expression in both groups of patients.

As the output data of the two techniques is different (TaqMan output is linear while RT-PCR ELISA output is non-linear), it was not possible to directly compare the results of each method. Instead, the output data from the TaqMan system were analysed using the same statistical method applied to the data obtained from the RT-PCR ELISA work. Using the Wilcoxon Rank Signed Test, the mean value for each time point was compared to the baseline value for that particular patient. The results have been shown graphically in figures 3.2b-i in the form of box and whisker plots, and in tables 3.2a and b.

These results clearly show that the results of the RT-PCR ELISA study detected real changes in IL-4 and IL-10 gene expression in renal transplant recipients and were not in any way artefact. The re-analysis of the same stored cDNA samples, using the fully quantitative TaqMan system, has substantiated those findings. The results still closely correlated with clinical events and the expression profiles remain different for those experiencing a rejection episode compared with those that did not in the case of IL-4. However, for IL-10 gene expression, the differences between rejectors and non-rejectors may not be so marked as initially reported. The direction of the change of gene expression of both cytokines remains the same.

In the RT-PCR ELISA study it was not appropriate to quantitate the changes in gene expression detected. Using the TaqMan system this becomes possible. It can be seen from the box and whisker plots of the data obtained using the TaqMan system (figures 3.2c, e, g and i) that the mean gene expression of both cytokines in both groups of transplant recipients varies by several orders of magnitude. As each single point increment in C_T value equates to a doubling of starting material measured, these changes in gene expression are large as well as statistically significant.

The results of the TaqMan analysis of the gene expression of the cytokines IL-4 and IL-10 imply that the changes in gene expression of the other cytokines (IL-2, IFN- γ , IL-5 and IL-13) investigated in the RT-PCR ELISA study are also real changes. This gives further weight to those important and interesting observations.

These observations suggest that a RT-PCR ELISA, when used carefully with good laboratory technique, is a more reliable tool for detecting differences in gene expression than had been previously thought.

This phase of the study has confirmed that the results from the earlier immunological monitoring work from our laboratory had identified real changes in IL-4 and IL-10 gene expression following renal transplantation and both before and after the initiation of anti-rejection therapy. These changes were detected in mRNA (later transcribed into cDNA) extracted from T lymphocytes obtained sequentially from the peripheral blood mononuclear cells of renal transplant recipients in the first four weeks following transplantation. These changes produced profiles that were different in recipients experiencing acute allograft rejection compared with those that had an uncomplicated early post-transplant period. These significant findings justified the plan to elaborate on the earlier work by continuing to measure cytokine gene expression in peripheral blood mononuclear cells of further renal transplant recipients in an attempt to further clarify the role of cytokines in acute allograft rejection. It would also aim to increase our knowledge of the use of peripheral blood mononuclear cell cytokine gene expression as a method of immunological monitoring. This study would use peripheral blood mononuclear cells rather than T lymphocytes, in the knowledge that any changes identified using the Taqman system could be confidently compared with the earlier results.

Chapter 4: The Investigation of the Use of a House Keeping Gene for PCR Standardisation

The next phase of the study was to investigate the potential benefits of using one or more house keeping genes for the purposes of standardisation within the PCR, i.e. to control for any potential fluctuations in cDNA concentrations. Specifically, the experiments were designed to determine whether human β -actin, β 2-microglobulin and transferrin receptor were suitable house keeping genes in our study cohort of renal transplant donors and recipients. For one or both of these genes to be suitable, their expression would need to be independent of clinical events.

This chapter will introduce the potential benefits of using a reliable house keeping gene in this study and review the usage of β -actin as a house keeping gene in the fields of cytokine gene expression and organ transplantation. It will then describe the results of the experiments performed in the search of a suitable house keeping gene for this study. Finally, the results of the experiments will be discussed.

4.1 Introduction

It remains a possibility that the differences seen in cytokine gene expression demonstrated in the RT-PCR ELISA work from our laboratory and repeated using TaqMan quantitative PCR (described in Chapter 3) might be influenced by variations in the amount of cDNA being analysed. In the RT-PCR ELISA study, and in this present study, every attempt was made to minimise this potential problem. When the RNA was extracted, the concentration of each individual solution was measured using a spectrophotometer. Each solution was then standardised to a concentration of 125µg/µl. This meant that the 8µl of RNA solution that was added to the cDNA transcription reagents contained 1µg of RNA. Of course, this standardisation was for total RNA and not for messenger RNA (mRNA). It was the mRNA that was measured, after transcription, to establish the level of gene expression. Despite the precautions taken, there were still two possible reasons for fluctuations in measurements that were not the result of variations in expression. Firstly, was the amount of mRNA constant in 1µg of total RNA? Secondly, was the cDNA transcription reaction 100% efficient, or at least constant? If it were possible to control for such variations, this would improve the quality of the results by negating the fluctuations in apparent expression being caused by variations in the amount of cDNA measured.

4.1.1 Why use a house keeping gene?

The rationale behind using a house keeping gene is two-fold. Firstly, it can be simply used as a marker of the presence or absence of good quality cDNA or DNA. For example, in the event of a low or absent result from the gene under investigation, if the house keeping gene has been successfully amplified, then there is probably adequate material to amplify, but the gene under investigation is either not expressed or expressed at an undetectable level. On the other hand, if neither the house keeping gene, or the gene under investigation, amplify then the reaction has either failed or there is inadequate quality cDNA or DNA to amplify. Secondly, a house keeping gene can be used to standardise the results obtained from the gene or genes under investigation by controlling for the amount of material added to the PCR. This is possible as the expression of a house keeping gene is presumed to be constant over time, and not affected by the influences that are under investigation. If expression of a house keeping gene was indeed constant, then it would be possible to use it to control for possible fluctuations in the amount of the cDNA in the reaction as is often practiced in many laboratories (Lipman 1998, Suzuki 2000). Therefore, one would expect its measurement to be constant over time, in the same individual, if the same amount of substrate were added. If there were a variation in the measurement of the house keeping gene, then this could reflect a variation in the amount or concentration of the mRNA and/or cDNA added. Using a house keeping gene to standardise for the quantity of cDNA added should allow for a more accurate assessment of the changes in expression of the gene or genes under investigation. In this

particular study, the expression of the house keeping gene should not have been influenced by surgery, solid organ transplantation, acute allograft rejection and anti-rejection treatment.

4.1.2 Which house keeping gene to use?

At the time the study commenced, Applied Biosystems (Foster City, Ca) produced commercially available TaqMan primers and probes that were marketed as specific for several potential human house keeping genes. It was not feasible to test them all on our cohort of patients. Three genes were chosen, namely β -actin, β 2-microglobulin and transferrin receptor. The reasons for this choice are explained below.

Glyceraldehyde-3-phosphate dehydrogenase

One of the house keeping genes not selected, but available, was glyceraldehyde-3-phosphate dehydrogenase (GAPDH). This gene has previously been used in studies measuring cytokine gene expression. One particular study used it as a house keeping gene when studying cytokine gene expression in renal transplant core biopsies (Lipman 1998). In Lipman's study, gene expression was measured using a competitive template PCR technique. In our study we were using the TaqMan PCR system. Early experience, in our institution, with TaqMan and the use of GAPDH as a house keeping gene had been disappointing. This particular gene had been found to be unreliable as a house keeping gene on this particular system due to inconsistent and unexplainable results, possibly as the result of the presence of

pseudogenes (Dr M. Pickett, Department of Molecular Microbiology, personal communication). For this reason it was not considered further. Others have also found that the expression of this gene, despite being a popular choice as a house keeping gene, is unreliable and should be thoroughly tested before use (Suzuki 2000).

β -actin

β -actin is a cytoskeletal protein found in all cells, and is encoded by a gene located on chromosome 7 position p22-p12. It is a popular house keeping gene with many references to its use in the scientific literature (Suzuki 2000). It has been used in a wide range of human tissues, in a non-transplant setting, including vascular smooth muscle cells (Mao 1999), normal and malignant bronchial epithelial cells (Willey 1998), normal and malignant hepatocytes (Chang 1998) and prostatic tissue (Galvan 1997). Of more relevance to this study, it has been used as a house keeping gene in studies investigating cytokine gene expression in both animal models (Overbergh 1999, El-Naggar 1998) and human cell lines (Halminen 1999). β -actin has also been used as a house keeping gene in animal models of both liver transplantation (Rokahr 1998) and renal transplantation (Naar 1998). Groups studying human solid organ transplantation have also used β -actin as a house keeping gene in the investigation of both liver transplant recipients (Schiano 1999, Flach 1998) and renal transplant recipients (Zhao 1998). This study is investigating cytokine gene expression in peripheral blood mononuclear cells (PBMCs). β -actin has been as a house keeping gene in other studies using PBMCs in humans (Dassi 1998). It has also been used when investigating cytokine gene

expression in the field of human transplantation. Iacono and co-workers used β -actin for the purposes of standardisation when measuring cytokine gene expression in cells recovered from bronchoalveolar lavage procedures and peripheral blood lymphocytes of lung transplant recipients (Iacono 1997). Sharma et al also used β -actin as an internal control when measuring cytokine gene expression in 127 renal transplant core biopsies (Sharma 1996). Gorczynski and colleagues also used β -actin as an internal control in their study measuring cytokine gene expression in peripheral lymphocytes and core biopsies of liver transplant recipients. They also measured β -actin gene expression in peripheral lymphocytes of 12 non-transplant controls (Gorczynski 1996).

There was, therefore, ample evidence in the literature to support our choice of β -actin as a potential house keeping gene. This was given additional support by several publications from groups already using TaqMan. These groups had reported the successful usage of human β -actin as a standardisation tool when using this system system (Heid 1996, Kruse 1997, Härtel 1999, Kreuzer 1999).

To add credence to the investigation of the use of β -actin as a house keeping gene, it was decided that we would also investigate two other potential house keeping genes, namely β -2-microglobulin and transferrin receptor.

β-2-microglobulin

The β-2-microglobulin (β2M) polypeptide, as discussed earlier (see section 1.2), is part of the expressed Class I HLA molecule and is encoded on a non-MHC region of chromosome 15. Like β-actin, β2M had been used as a house keeping gene in various human studies. It has been used as a house keeping gene in tissues from attenuated human cell lines (Morehouse 2000), leukaemia patients (Palligaard 1999), prostatic biopsies (Moriyama 1999), the female reproductive tract (Gipson 1999), colonic biopsies (Leccia 1998) and brain tissue (Waha 1998). β2M has also been used as a house keeping gene in a quantitative RT-PCR system studying cytokine gene expression in a porcine model (Reddy 2000).

We were aware that there is substantial evidence (see section 1.5.2) that soluble β2M levels increased before and during acute rejection episodes. These results were derived from measurements of soluble β2M measured in the serum. These elevations are probably the result of cellular damage to donor cells with subsequent shedding of the Class I molecule. Measurement of β2M gene expression in recipient peripheral blood mononuclear cells should not be influenced by this process and still had the potential to be useful as a house keeping gene.

Transferrin receptor

Transferrin receptor (TfR) was the third gene chosen as a potential house keeping gene. This was a less strong candidate that was chosen, in part, as a comparison with the other two markers. Though TfR has the potential to act

as a house keeping gene in some cohorts of patients, i.e. the living donors, it was not thought to be appropriate as an ideal house keeping gene in renal transplant recipients (see below). Therefore, if TfR gene expression varied while β -actin and β 2M gene expression did not, then this would support the value of either or both genes as good house keeping genes.

The gene is located on chromosome 3 at position q29. The gene product of TfR is a disulfide-linked transmembrane glycoprotein. It is a widely distributed cell surface receptor present on rapidly proliferating and specialised quiescent cells that have particular needs for high levels of iron (Huebers 1987, May 1985). During T-cell activation, TfR expression follows the expression of IL-2R in a sequence that is necessary to initiate cell proliferation in quiescent T lymphocytes (May 1985, Pattanapanyasat 1991). In cardiac transplant recipients it has been shown that the percentage of TfR expressing CD4/CD8 cells increases both before and during allograft rejection and diminishes after successful treatment (Hoshinaga 1987). More recently, in a murine cardiac transplant model, the combination of IL-2R blockade and TfR blockade, with monoclonal antibodies, significantly prolonged graft survival when compared to controls or either agent alone (Woodward 1999).

4.2 Investigation of Single PCR Tube Multiple Gene Expression Analysis Using Different TaqMan Reporter Dyes

The probes designed for the purpose of detecting cytokine gene expression were labelled with the fluorescent reporter dye 6-carboxyfluorescein (FAM), whereas those used for detection of the house keeping genes had a alternative reporter dye, namely VIC. This potentially facilitates the measurement of the expression of both the gene under investigation and the house keeping gene in the same PCR tube. This has the obvious advantages of enabling direct measurement of the house keeping gene in the same sample as the gene under investigation. It also allows the analysis of more samples on each TaqMan run. Before we adopted this method, it was felt best to compare the measurement of both the house keeping gene and the cytokine gene under investigation under two different circumstances, i.e. separately or in a single tube. Figures 4.2a-f clearly demonstrate that if both the house keeping gene and the cytokine gene (on these occasions IL-4 and IL-10) under investigation are both analysed in a single PCR tube there was a greater variation in the C_T values of some of the duplicates at some time points. Given our earlier results, when analysing the stored cDNA from the RT-PCR ELISA study (see chapter 3), these results were disappointing. The variation in the C_T value from identical duplicates should be less than half of one cycle. In these experiments the variation was far larger in some instances. This was thought to be the result of a possible interaction between the commercially designed house keeping primers and probe and the designed primers and probes for both IL-4 and IL-10. This variation was unacceptable. In a more extensive study, with

additional time, it would be possible to manipulate and optimise the system to facilitate single tube multiple gene expression measurement. However, for the purposes of this study, it was decided to analyse both genes in separate PCR tubes.

The format for all six graphs is the same. The Y-axis represents the C_T value and the X-axis represents time. Time point 1 is the day of transplantation, time point 2 being day 1 post- transplant etc. Each reaction was performed in duplicate and both series are shown.

Figure 4.2a. Graph showing a TaqMan result of a non-rejector using in house designed IL-4 primers and probe

Figure 4.2b. Graph showing a TaqMan result of a non-rejector using in house designed IL-4 primers and probe with the commercially produced β -actin primers and probe in the same tube

These graphs demonstrate the variation of IL-4 expression depending on whether IL-4 was being analysed alone (figure 4.2a) or whether both IL-4 and human β -actin (huBAC) were analysed in the same PCR tube (figure 4.2b).

Figure 4.2c. Graph showing a TaqMan result of a non-rejector using in house designed IL-10 primers and probe

Figure 4.2d. Graph showing a TaqMan result of a non-rejector using in house designed IL-10 primers and probe with the commercially produced β -actin primers and probe in the same tube

These graphs demonstrate the variation of IL-10 expression depending on whether IL-10 was being analysed alone (figure 4.2c) or whether both IL-10 and huBAC were analysed in the same PCR tube (figure 4.2d).

Figure 4.2e. Graph showing a TaqMan result of a non-rejector using the commercially produced β -actin primers and probe

Figure 4.2f. Graph showing a TaqMan result of a non-rejector using the commercially produced β -actin primers and probe with the in house designed IL-10 primers and probe in the same tube

These graphs demonstrate the variation of huBac expression depending on whether huBac was being analysed alone (figure 4.2e) or whether both IL-10 and huBac were analysed in the same PCR tube (figure 4.2f).

4.3 Results

This section will present the results of the gene expression data of the 3 house keeping genes studied. The results will be presented both graphically and in tabular form. The results from the three different groups of patients, living donors, non-rejectors and rejectors, will be presented separately for comparison.

4.3.1 Individual variation in Expression of the Three House Keeping Genes

Using the standard TaqMan methodology described in section 2.4, all patient samples were analysed for all three house keeping genes. All three house keeping genes were normally analysed in separate PCR runs, but each patient series was always analysed completely in one plate for each house keeping gene. Figure 4.3.1a shows a typical example of the expression patterns of all three genes in a non-rejector. Figure 4.3.1b shows a typical example of the expression pattern of all three genes in a rejector. Figure 4.3.1c shows a typical example of the expression pattern of all three genes in a living-donor.

Figure 4.3.1a. Graph showing the gene expression of all three potential house keeping genes in a non-rejecting transplant recipient

This graph represents the expression data for all three house keeping genes for a non-rejecting transplant recipient. The Y-axis is the CT value. The X-axis is time in days where 1 is the day of transplantation.

B2MAV = average value of the duplicates obtained from the measurement of β -2-microglobulin gene expression at each time point.

BAAV = average value of the duplicates obtained from the measurement of β -actin gene expression at each time point.

TFRAV = average value of the duplicates obtained from the measurement of transferrin receptor gene expression at each time point.

Figure 4.3.1b. Graph showing the gene expression of all three potential house keeping genes in a rejecting transplant recipient

This graph represents the expression data for all three house keeping genes for a rejecting transplant recipient. The Y-axis is the CT value. The X-axis is time in days where 1 is the day of transplantation.

B2MAV = average value of the duplicates obtained from the measurement of β -2-microglobulin gene expression at each time point.

BAAV = average value of the duplicates obtained from the measurement of β -actin gene expression at each time point.

TFRAV = average value of the duplicates obtained from the measurement of transferrin receptor gene expression at each time point.

Figure 4.3.1c. Graph showing the gene expression of all three potential house keeping genes in a living donor

This graph represents the expression data for all three house keeping genes for a living-donor. The Y-axis is the CT value. The X-axis is time in days where 1 is the day of transplantation.

B2MAV = average value of the duplicates obtained from the measurement of β -2-microglobulin gene expression at each time point.

BAAV = average value of the duplicates obtained from the measurement of β -actin gene expression at each time point.

TFRAV = average value of the duplicates obtained from the measurement of transferrin receptor gene expression at each time point.

4.3.2 Correlation Between the Expression Data of the Three House Keeping Genes in both Living Donors and Transplant Recipients

It can be seen from the three representative graphs in the previous section, that there appeared to be a similarity in the variation of expression of all three house keeping genes in all three cohorts of patients. β -actin and β 2-microglobulin were both expressed at a comparable level, whereas transferrin receptor tended to be expressed at a lower level. However the magnitude of change of expression for all three genes from one time point to the next was the same.

The next six graphs are scattergrams comparing the expression data of each of the house keeping genes with each other for both living donors and transplant recipients. The gene expression data for each individual house keeping gene from each time point of each patient in the group were plotted against the corresponding value for a different house keeping gene.

Figure 4.3.2a. Graph demonstrating the correlation between the gene expression data of β -2- and β -actin for the living donors

This graph demonstrates the correlation between the gene expression data of β -2-microglobulin (B2MAV plotted on the X-axis) and β -actin (BAAV plotted on the Y-axis) for the living donors. The scale on both axes is the C_T value.

Figure 4.3.2b. Graph demonstrating the correlation between the gene expression data of β -2- and transferrin receptor for the living donors

This graph demonstrates the correlation between the gene expression data of β -2-microglobulin (B2MAV plotted on the X-axis) and transferrin receptor (TFRAV plotted on the Y-axis) for the living donors. The scale on both axes is the C_T value.

Figure 4.3.2c. Graph demonstrating the correlation between the gene expression data of transferrin receptor and β -actin for the living donors

This graph demonstrates the correlation between the gene expression data of transferrin receptor (plotted on the X-axis) and β -actin (BAAV plotted on the Y-axis) for the living donors. The scale on both axes is the C_T value.

Figure 4.3.2d. Graph demonstrating the correlation between the gene expression data of β -2- and β -actin for all the transplant recipients

This graph demonstrates the correlation between the gene expression data of β -2-microglobulin (B2MAV plotted on the X-axis) and β -actin (BAAV plotted on the Y-axis) for all the transplant recipients. The scale on both axes is the C_T value.

Figure 4.3.2e. Graph demonstrating the correlation between the gene expression data of β -2-microglobulin and transferrin receptor for all the transplant recipients

This graph demonstrates the correlation between the gene expression data of β -2-microglobulin (B2MAV plotted on the X -axis) and transferrin receptor (TFRAV plotted on the Y-axis) for all the transplant recipients. The scale on both axes is the C_T value.

Figure 4.3.2f. Graph demonstrating the correlation between the gene expression data of transferrin receptor and β -actin for all the transplant recipients

This graph demonstrates the correlation between the gene expression data of transferrin receptor (TFRAV plotted on the X-axis) and β -actin (BAAV plotted on the Y-axis) for all the transplant recipients. The scale on both axes is the C_T value.

The visual correlation between the gene expression data of the three house keeping genes of both living donors and transplant recipients can be clearly seen in the above graphs 4.32a-f.

Using Pearson's correlation test it was possible to show that these visual similarities were highly significant, not just in the donors but also in the recipients. Tables 4.3.2a and 4.3.2b give the results of the correlation test with correlation coefficients and p values for living donors and transplant recipients respectively.

Table 4.3.2a. Results of the correlation between the expression of the three potential house keeping genes in the living donors

		β -2-microglobulin	β -actin	transferrin receptor
β -2-microglobulin	Pearson's correlation	-	0.803	0.807
	Significance (2 tailed)	-	<0.001	<0.001
β -actin	Pearson's correlation	0.803	-	0.836
	Significance (2 tailed)	<0.001	-	<0.001
transferrin receptor	Pearson's correlation	0.807	0.836	-
	Significance (2 tailed)	<0.001	<0.001	-

This table presents the results of the correlations demonstrated graphically in figures 4.3.2 a, b and c. The table shows that the correlation between β -2-microglobulin and β -actin, β -2-microglobulin and transferrin receptor, and β -actin and transferrin receptor is highly significant in all three cases for the living donors.

Table 4.3.2b. Results of the correlation between the expression of the three potential house keeping genes in the transplant recipients

		β -2-microglobulin	β -actin	transferrin receptor
β -2-microglobulin	Pearson's correlation	-	0.897	0.848
	Significance (2 tailed)	-	<0.001	<0.001
β -actin	Pearson's correlation	0.897	-	0.904
	Significance (2 tailed)	<0.001	-	<0.001
transferrin receptor	Pearson's correlation	0.848	0.904	-
	Significance (2 tailed)	<0.001	<0.001	-

This table presents the results of the correlations demonstrated graphically in figures 4.3.2 d, e and f. The table shows that the correlation between β -2-microglobulin and β -actin, β -2-microglobulin and transferrin receptor, and β -actin and transferrin receptor is highly significant in all three cases for the transplant recipients.

4.3.3 Gene Expression Data for Each House Keeping Gene in Each of the Three Cohorts of Patients

To use the expression of a house keeping gene for PCR standardisation its expression must be constant and not influenced by the processes under investigation.

The box and whisker graphs (figures 4.3.3a-l) in this section demonstrate the mean gene expression data for each of the house keeping genes (β -actin, β 2-microglobulin and transferrin receptor) in each of the three cohorts of patients i.e. the living donors, the non-rejecting transplant recipients and the rejecting transplant recipients. For the latter group there are six graphs rather than the three for the other two groups. This is because the first set of three shows the house keeping gene expression data sequentially, starting with the pre-transplant sample. The second set of three box and whisker plots shows the same data but starting 6 days before the diagnosis and subsequent treatment of the acute rejection and continuing for 6 days after.

Figure 4.3.3a. Box and whisker plot of the gene expression data of β -actin in the living donors

This graph is a box and whisker plot and shows the results of the measurement of β -actin gene expression in the living donor cohort. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-nephrectomy sample.

Figure 4.3.3b. Box and whisker plot of the gene expression data of $\beta 2$ -microglobulin in the living donors

This graph is a box and whisker plot and shows the results of the measurement of $\beta 2$ -microglobulin gene expression in the living donor cohort. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-nephrectomy sample.

Figure 4.3.3c. Box and whisker plot of the gene expression data of transferrin receptor in the living donors

This graph is a box and whisker plot and shows the results of the measurement of transferrin receptor gene expression in the living donor cohort. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-nephrectomy sample.

Figure 4.3.3d. Box and whisker plot of the gene expression data of β -actin in the non-rejecting transplant recipients

This graph is a box and whisker plot and shows the results of the measurement of β -actin gene expression in the non-rejecting transplant recipients. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-transplant sample.

Figure 4.3.3e. Box and whisker plot of the gene expression data of β 2-microglobulin in the non-rejecting transplant recipients

This graph is a box and whisker plot and shows the results of the measurement of β 2-microglobulin gene expression in the non-rejecting transplant recipients. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-transplant sample.

Figure 4.3.3f. Box and whisker plot of the gene expression data of transferrin receptor in the non-rejecting transplant recipients

This graph is a box and whisker plot and shows the results of the measurement of transferring receptor gene expression in the non-rejecting transplant recipients. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-transplant sample.

Figure 4.3.3g. Box and whisker plot of the gene expression data of β -actin in the rejecting transplant recipients

This graph is a box and whisker plot and shows the results of the measurement of β -actin gene expression in the rejecting transplant recipients. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-transplant sample.

Figure 4.3.3h. Box and whisker plot of the gene expression data of $\beta 2$ -microglobulin in the rejecting transplant recipients

This graph is a box and whisker plot and shows the results of the measurement of $\beta 2$ -microglobulin gene expression in the rejecting transplant recipients. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-transplant sample.

Figure 4.3.3i. Box and whisker plot of the gene expression data of transferrin receptor in the rejecting transplant recipients

This graph is a box and whisker plot and shows the results of the measurement of transferring receptor gene expression in the rejecting transplant recipients. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-transplant sample.

Figure 4.3.3j. Box and whisker plot of the gene expression data of β -actin in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis

This graph is a box and whisker plot and shows the results of the measurement of β -actin gene expression in the rejecting transplant recipients re-aligned to the day of rejection diagnosis. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted from left to right starting with the pre-transplant sample and then the five samples before rejection (BABR5 –BABR1) and the five days post-rejection (BAPR1-BAPR5).

Figure 4.3.3k. Box and whisker plot of the gene expression data of β 2-microglobulin in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis

This graph is a box and whisker plot and shows the results of the measurement of β 2-microglobulin gene expression in the rejecting transplant recipients re-aligned to the day of rejection diagnosis. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted from left to right starting with the pre-transplant sample and then the five samples before rejection (B2MBR5 –B2MBR1) and the five days post-rejection (B2MPR1-B2MPR5).

Figure 4.3.31. Box and whisker plot of the gene expression data of transferrin receptor in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis

This graph is a box and whisker plot and shows the results of the measurement of transferrin receptor gene expression in the rejecting transplant recipients re-aligned to the day of rejection diagnosis. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted from left to right starting with the pre-transplant sample and then the five samples before rejection (TFRBR5 –TFRBR1) and the five days post-rejection (TFRPR1-TFRPR5).

The following eight tables (tables 4.3.3a-h) are a tabular representation of the previous twelve graphs (figures 4.3.3a-l). The first table of each pair shows the direction of change of the mean of the gene expression data for each house keeping gene in a sequential stepwise fashion. The analysis compared the mean pre-transplant value with the mean value from the first post-transplant day, the mean value of the first post-transplant day with the mean value of the second post-transplant day etc. The statistical analysis was performed using Wilcoxon's Rank Signed Test. The second table of each pair shows the direction of change of the mean of the gene expression data for each house keeping gene for each time-point compared to the mean baseline pre-transplant value. The statistical analysis was again performed using Wilcoxon's Rank Signed Test.

There are separate tables for the living donors (tables 4.3.3a and b), the non-rejecting transplant recipients (tables 4.3.3c and d), and the rejecting transplant recipients (tables 4.3.3e and f).

The seventh and eighth tables (tables 4.3.3g and h) are slightly different from the previous ones. They again present the data from the rejecting cohort of transplant recipients. However, on this occasion the data are re-aligned to the day on which the diagnosis of rejection was made (in table 4.3.3g and h this is time point BR1). Treatment was commenced on day BR1; therefore, day PR1 in these tables corresponds to the first day following the initiation of anti-rejection treatment. The analysis again performed in the two ways described above. Firstly, comparing the mean expression data from each house keeping

in a sequential stepwise manner. The mean expression data from day five prior to the diagnosis of rejection was compared with that from day four prior to rejection etc. Secondly, comparing the mean value of each time-point with the mean pre-transplant value. The statistical analysis was again performed using Wilcoxon's Rank Signed Test in both cases.

The graphs above (figures 4.3.3a-l) and the tables below (tables 4.3.3a-h) show that there were statistically significant changes in mean gene expression of all three potential house keeping genes in both the rejecting and non-rejecting transplant recipients. These variations in expression seen in the recipients and the results of the analysis of β -actin, β 2-microglobulin and transferrin receptor gene expression obtained from the living donors are discussed below in section 4.4.

Table 4.3.3a. Results of stepwise analysis of the gene expression data of the three potential house keeping genes in the living donors

Time points in days in sequential steps (1 = pre-nephrectomy, 2 = day 1 post-nephrectomy etc.)		1-2	2-3	3-4	4-5	5-6	6-7	7-8	8-9
β -actin	Direction of change of gene expression	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS
β 2-microglobulin	Direction of change of gene expression	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS
Transferrin receptor	Direction of change of gene expression	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS

This table shows the stepwise direction of change of the mean gene expression data for the three house keeping genes in the living donor cohort and the level of significance of these changes. Statistical analysis was performed using Wilcoxon's rank signed test. NS = non-significant.

Table 4.3.3b. Results of comparison to baseline analysis of the gene expression data of the three potential house keeping genes in the living donors

Time points in days comparing the pre-nephrectomy value with each subsequent value (1 = pre-nephrectomy, 2 = day 1 post-nephrectomy etc.)		1-2	1-3	1-4	1-5	1-6	1-7	1-8	1-9
β -actin	Direction of change of gene expression	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS
β 2-microglobulin	Direction of change of gene expression	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS
Transferrin receptor	Direction of change of gene expression	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS

This table shows the direction of change of the mean gene expression data of each time point compared to the baseline (pre-nephrectomy) value for the three house keeping genes in the living donor cohort. Statistical analysis was performed using Wilcoxon's rank signed test. NS = non-significant.

Table 4.3.3c. Results of stepwise analysis of the gene expression data of the three potential house keeping genes in the non-rejecting transplant recipients

Time points in days in sequential steps (1 = pre-transplant, 2 = day 1 post-transplant etc.)		1-2	2-3	3-4	4-5	5-6	6-7	7-8	8-9	9-10	10-11	11-12	12-13	13-14	14-15
β -actin	Direction of change of gene expression	↑	-	-	-	-	-	-	-	-	-	-	-	-	-
	p value	<0.001	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
β 2-microglobulin	Direction of change of gene expression	↑	-	-	-	-	-	↓	-	↑	-	-	-	-	-
	p value	0.012	NS	NS	NS	NS	NS	0.026	NS	0.033	NS	NS	NS	NS	NS
Transferrin receptor	-	↑	-	-	-	-	-	↓	-	-	-	-	-	-	-
	p value	<0.001	NS	NS	NS	NS	NS	0.036	NS	NS	NS	NS	NS	NS	NS

This table shows the stepwise direction of change of the mean gene expression data for the three house keeping genes in the non-rejecting transplant recipients and the level of significance of these changes. Statistical analysis was performed using Wilcoxon's rank signed test. NS = non-significant.

Table 4.3.3d. Results of comparison to baseline analysis of the gene expression data of the three potential house keeping genes in the non-rejecting transplant recipients

Time points in days comparing the pre-transplant value with each subsequent value (1 = pre-transplant, 2 = day 1 post-transplant etc.)		1-2	1-3	1-4	1-5	1-6	1-7	1-8	1-9	1-10	1-11	1-12	1-13	1-14	1-15
β -actin	Direction of change of gene expression	↑	-	↑	-	-	↑	-	-	↑	↑	-	-	-	-
	p value	<0.001	NS	0.048	NS	NS	0.015	NS	NS	0.016	0.016	NS	NS	NS	NS
β 2-microglobulin	Direction of change of gene expression	↑	-	↑	-	↑	↑	-	↑	↑	↑	-	-	-	-
	p value	0.012	NS	0.042	NS	0.033	0.009	NS	0.046	0.009	0.004	NS	NS	NS	NS
Transferrin receptor	Direction of change of gene expression	↑	↑	↑	-	↑	↑	↑	↑	↑	↑	-	-	-	-
	p value	<0.001	0.009	0.006	NS	0.005	0.002	0.049	0.008	0.003	0.026	NS	NS	NS	NS

This table shows the direction of change of the mean gene expression data of each time point compared to the baseline (pre-transplant) for the three house keeping genes in the non-rejecting transplant recipients and the level of significance of these changes. Statistical analysis was performed using Wilcoxon's rank signed test. NS =non-significant.

Table 4.3.3e. Results of stepwise analysis of the gene expression data of the three potential house keeping genes in the rejecting transplant recipients

Time points in days in sequential steps (1 = pre-transplant, 2 = day 1 post-transplant etc.)		1-2	2-3	3-4	4-5	5-6	6-7	7-8	8-9	9-10	10-11	11-12	12-13	13-14	14-15
β -actin	Direction of change of gene expression	-	-	-	↓	-	-	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	0.018	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
β 2-microglobulin	Direction of change of gene expression	-	-	-	↓	-	-	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	0.016	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Transferrin receptor	Direction of change of gene expression	-	-	-	↓	-	-	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	0.037	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

This table shows the stepwise direction of change of the mean gene expression data for the three house keeping genes in the rejecting transplant recipients and the level of significance of these changes. Statistical analysis was performed using Wilcoxon's rank signed test. NS = non-significant.

Table 4.3.3f. Results of comparison to baseline analysis of the gene expression data of the three potential house keeping genes in the rejecting transplant recipients

Time points in days comparing the pre-transplant value with each subsequent value (1 = pre-transplant, 2 = day 1 post-transplant etc.)		1-2	1-3	1-4	1-5	1-6	1-7	1-8	1-9	1-10	1-11	1-12	1-13	1-14	1-15
β -actin	Direction of change of gene expression	-	-	↑	-	-	↑	↑	-	-	↑	↑	-	↑	-
	p value	NS	NS	0.002	NS	NS	0.023	0.028	NS	NS	0.05	0.017	NS	0.021	NS
β 2-microglobulin	Direction of change of gene expression	-	-	↑	-	-	-	-	-	-	-	↑	-	-	-
	p value	NS	NS	0.005	NS	NS	NS	NS	NS	NS	NS	0.041	NS	NS	NS
Transferrin receptor	Direction of change of gene expression	-	-	↑	-	↑	↑	↑	-	-	↑	↑	-	-	↑
	p value	NS	NS	0.016	NS	0.041	0.023	0.008	NS	NS	0.038	0.05	NS	NS	0.046

This table shows the direction of change of the mean gene expression data of each time point compared to the baseline (pre-transplant) for the three house keeping genes in the rejecting transplant recipients and the level of significance of these changes. Statistical analysis was performed using Wilcoxon's rank signed test. NS =non-significant.

Table 4.3.3g. Results of stepwise analysis of the gene expression data of the three potential house keeping genes in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis

Time points in days in sequential steps (BR5 = 5 days before diagnosis of rejection, BR4 = 4 days before diagnosis of rejection etc.)		PTBS- BR5	BR5- BR4	BR4- BR3	BR3- BR2	BR2- BR1	BR1- PR1	PR1- PR2	PR2- PR3	PR3- PR4	PR4- PR5
β -actin	Direction of change of gene expression	-	-	-	↑	↓	-	↓	-	-	-
	p value	NS	NS	NS	0.004	0.009	NS	0.021	NS	NS	NS
β 2-microglobulin	Direction of change of gene expression	-	-	-	↑	↓	-	↓	-	-	-
	p value	NS	NS	NS	0.006	0.007	NS	0.028	NS	NS	NS
Transferrin receptor	Direction of change of gene expression	-	-	-	↑	-	↓	-	-	-	-
	p value	NS	NS	NS	0.028	NS	0.025	NS	NS	NS	NS

This table shows the stepwise direction of change of the mean gene expression data for the three house keeping genes in the rejecting transplant recipients when the data were re-aligned to the day when the diagnosis of rejection was made. The level of significance of these changes is also shown. Statistical analysis was performed using Wilcoxon's rank signed test. PTBS = pre-transplant baseline. BR = before rejection. PR = post-rejection. NS = non-significant.

Table 4.3.3h. Results of comparison to baseline analysis of the gene expression data of the three potential house keeping genes in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis

Time points in days comparing the pre-transplant value (PTBS) with each value before (BR) and post (PR) the diagnosis of rejection.		PTBS-BR5	PTBS-BR4	PTBS-BR3	PTBS-BR2	PTBS-BR1	PTBS-PR1	PTBS-PR2	PTBS-PR3	PTBS-PR4	PTBS-PR5
β -actin	Direction of change of gene expression	-	↑	↑	↑	↑	↑	-	-	-	-
	p value	NS	0.041	0.056	0.001	0.001	0.033	NS	NS	NS	NS
β 2-microglobulin	Direction of change of gene expression	-	↑	↑	↑	↑	-	-	-	-	-
	p value	NS	0.013	0.09	0.01	0.007	NS	NS	NS	NS	NS
Transferrin receptor	Direction of change of gene expression	-	↑	↑	↑	↑	-	-	-	↑	-
	p value	NS	0.017	0.033	0.003	0.021	NS	NS	NS	0.018	NS

This table shows the direction of change from the baseline of the mean gene expression data for the three house keeping genes in the rejecting transplant recipients when the data were re-aligned to the day when the diagnosis of rejection was made. The level of significance of these changes is also shown. Statistical analysis was performed using Wilcoxon's rank signed test. PTBS = pre-transplant baseline. BR = before rejection. PR = post-rejection. NS = non-significant.

4.4 Discussion

The inherent expression of a house keeping gene should be constant. If this is the case, then variations in expression of such a gene can be interpreted as variations in the amount of cDNA being measured. This enables the variation in expression of the gene or genes under investigation to be standardised. This means that variations in expression of the gene or genes under investigation are genuine changes and not the result of varying amounts of cDNA being measured.

In the previous study on cytokine gene expression performed in our laboratory using RT-PCR ELISA technology, no house keeping gene was used for the purposes of standardisation. Even though the changes in cytokine gene expression detected corresponded to clinical events, there remained the possibility that at least some of these changes may have been artefact resulting from changes in the amount of cDNA measured. To address this issue, as part of this study, an attempt was made to find a suitable house keeping gene to use for standardisation in renal transplant recipients.

As discussed at the start of this chapter, various house keeping genes have been used for standardisation when measuring cytokine gene expression in both the transplant and non-transplant settings. The most frequently used were β -actin and GAPDH. As mentioned earlier, β -actin was chosen along with $\beta 2$ -microglobulin and transferrin receptor to undergo investigation in this study.

Using commercially available primers and probes for all three potential house keeping genes, and standard TaqMan PCR conditions (described in section 2.5), the gene expression of all patient samples (donors, non-rejectors and rejectors) was measured.

The initial representative graphs shown in section 4.3.1 (figures 4.3.1a-c) show that there was an obvious visual similarity between the expression of β -actin, β 2-microglobulin and transferrin receptor genes. This was despite the difference in the function of the proteins of the three different genes, and the potential role of transferrin receptor in antigen recognition and acute rejection. This similarity was seen in the living donors, the non-rejecting recipients and the rejecting recipients.

The visual similarities in the expression patterns of all three genes seen in the representative graphs were real. The correlations between the expression of β -actin, β 2-microglobulin and transferrin receptor genes were highly significant in both the living donors and the transplant recipients (figures 4.3.2a-f and tables 4.3.2a and b). This suggested that any changes seen in one of the potential house keeping genes would be seen in the other two genes as well.

In the living donor cohort the mean expression data results for all three genes were encouraging (figures 4.3.3a-c and tables 4.3.3a and b). There were no statistically significant changes in the mean gene expression of any of the three genes, at any of the time points. This implied that the surgery of the donor nephrectomy did not appear to significantly influence the expression of

the three genes. Therefore, it could be concluded that, in this group of patients it was possible to use β -actin, β 2-microglobulin or transferrin receptor as house keeping genes and use them for the purpose of standardisation.

The results of the gene expression analyses were less encouraging in the non-rejecting transplant recipients (figures 4.3.3d-f and tables 4.3.3c and d). In this group of patients, there was a statistically significant increase in mean expression of all three genes between the first and second time points, i.e. between the pre-transplant sample and the first post-transplant sample. This statistically significant difference between the mean expression of all three house keeping genes seen in the transplant recipients but not the donors must result from the effect of transplantation. There were also significant changes seen at other time points. All three genes were expressed at levels significantly higher than pre-transplant baseline values at several points in both the first and second week following transplantation.

There was no obvious explanation for the significant rise detected in the mean expression of β -actin, β 2-microglobulin and transferrin receptor genes at about a week post-transplant, but may reflect the initiation of immune processes that do not result in clinical rejection as described by Lipman et al (Lipman 1998).

The significant changes in gene expression of all three genes, particularly at the time of transplantation, seen in the non-rejecting recipients make all three genes poor candidates for house keeping genes in this group of patients.

In the group of patients experiencing an acute rejection episode the mean expression data of the three house keeping genes was different again (figures 4.3.3g-i and tables 4.3.3e and f). When the results were aligned to the day of transplantation there were no significant rises in expression of any of the three genes immediately following transplantation as seen in the non-rejecting recipients. However, there were again several significant increases in gene expression seen in all three genes when compared to pre-transplant levels. With rejection occurring at different times in different patients this data was difficult to analyse. The data was, therefore, re-aligned to the time when the diagnosis of the acute rejection episode was made (figures 4.3.3j-l and tables 4.3.3g and h). When this was performed a striking pattern emerged. There were significant increases in the mean expression of all three genes before the diagnosis of an acute rejection episode was made. These increases in expression levels were detectable several days prior to the clinical and biochemical changes associated with acute allograft rejection. Following initiation of the anti rejection therapy (0.5g of methyl prednisolone on three successive days in all but two of the patients who received a ten day course of OKT3) there was a significant reduction in expression levels. This resulted in expression levels returning to pre-transplant values.

These results cast serious doubt on the use of β -actin, β 2-microglobulin and transferrin receptor genes as house keeping genes in renal transplant recipients, and possibly in the transplant setting as a whole. The placement of a transplant, the process of acute allograft rejection and the commencement of anti-rejection therapy appear to have significant influences on the expression

of β -actin, β 2-microglobulin and transferrin receptor. In the case of transplantation there may be a different affect in the different groups. In the non-rejectors it appears to increase the expression of all three genes at the time of transplantation. This effect is not seen in the rejectors and probably reflects the more complex events occurring in the early stages of acute rejection and it's treatment, rather than a different effect in the two different groups.

The changes in transferrin receptor expression during acute rejection were not unexpected. Up-regulation of this gene is associated with up-regulation of the IL-2 receptor on activated T cells (May 1985, Pattanapanyasat 1991) and blockade of this receptor with monoclonal antibodies prolongs graft survival in animal transplant models (Woodward 1999). Also in cardiac transplant recipients, it has been reported that there is a significant increase in the number of cells expressing transferrin receptor at the time of an acute rejection episode (Hoshinaga 1987). The decreased expression detected following treatment of the rejection episode probably reflects the immunosuppressive effect of methyl prednisolone and/or OKT3.

The significant increase in expression of the transferrin receptor gene immediately following renal transplantation in the non-rejectors, but not the rejector group is more difficult to explain. This difference may, at least in part, be the result of the relatively small numbers in the other two groups. A more likely explanation is the results reflect a mixed picture in a group of patients at

various stages of immunological activity as some of the recipients showed clinical evidence of rejection within the first 48 – 72 hours.

Increased β 2-microglobulin expression in the non-rejecting recipients immediately following transplantation is unlikely to be the result of the surgery as there is no significant rise seen in the living donor group. It probably reflects immune activation following transplantation. Increased expression in the recipients at the time of an acute rejection episode probably results from an up-regulation of class I antigens in activated mononuclear cells in the peripheral blood. The decreased expression detected following treatment of the rejection episode, as with transferrin receptor, probably reflects the immunosuppressive effect of methyl prednisolone and/or OKT3.

Changes in β -actin gene expression in the non-rejectors at the time of transplantation are again unlikely to be related to the surgery. The significant changes seen in the rejectors at the time of the acute rejection episode may well result from the cell proliferation associated with the cytokine cascade seen in acute allograft rejection. The decrease following the treatment again probably reflects the immunosuppressive effect of the anti-rejection therapy.

These observations and results suggest that it would be inadvisable to use β -actin, β 2-microglobulin and transferrin receptor genes for the purpose of standardisation within the renal transplant population, and probably any transplant scenario. Transplantation, acute allograft rejection and anti-

rejection therapy all influence expression of the three genes under investigation.

The inconsistency of expression and response to acute allograft rejection and anti-rejection therapy of the three genes chosen as potential house keeping genes in this study limits their usefulness for the purposes of standardisation. Scientists in fields of research other than human renal transplantation have also identified limitations. Some of these groups have also found that expression of β -actin is not constant. In a rat brain ischaemia model, expression of this gene was found to increase 24hours after a surgical insult. This work was also performed using TaqMan technology. They concluded that there was a need for careful choice of control house keeping genes used for RNA/cDNA comparison (Harrison 2000). In another rat model, both β -actin and GAPDH gene expression were shown to vary significantly between tissues of the same animal and following surgery (Weisinger 1999). β -actin gene expression has also been shown to be inconsistent in stressed biologically diverse cell lines. Interestingly, GAPDH gene expression varied considerably as well (Zhong 1999). The expression of β -actin and GAPDH genes, in response to various different stimulations, in porcine macrophage and spleen cell lines have also been shown to be variable (Foss 1998).

There is supporting evidence that immunosuppressives may affect expression of house keeping genes. Tacrolimus and rapamycin have been shown to alter β -actin synthesis in stimulated human primary T lymphocytes. Tacrolimus strongly enhanced β -actin synthesis while rapamycin reduced its synthesis

(Miyamoto 1999). This observed effect might, in part, explain the significant rise in β -actin gene expression seen immediately following transplantation in the non-rejectors. The loading dose of Neoral given prior to transplantation in the studied transplant recipients may have a similar affect to the Tacrolimus in the study described above. This effect may not just be limited to β -actin gene expression and may be similar for the other two genes. The reason a significant rise may not be seen in the rejecting group, may still be the relatively small numbers in this group, or, as mentioned above, it may reflect varying degrees of immunological activation in recipients that went on to show clinical signs of acute allograft rejection and receive anti-rejection therapy.

In a different study, β 2-microglobulin gene expression, in human hepatocytes, was reduced by the addition of glucocorticoids. Another observation from this study was that intrahepatic β 2-microglobulin gene expression was higher in acute rejection than in resolving rejection (Garcia-Monzon 1998).

There is, therefore, mounting evidence that some of the genes (β -actin, β 2-microglobulin and GAPDH) previously used for the purposes of RNA and cDNA standardisation, when measuring gene expression, may not be as reliable as once thought. The results presented in this chapter cast further doubt on the use of house keeping genes in the field of transplantation as it would appear that the expression of β -actin, β 2-microglobulin and transferrin receptor genes are influenced by placement of the transplant, acute allograft rejection and anti-rejection therapy.

The aim of this part of the study was to identify a house keeping gene that could be used to control for fluctuations in the amounts of cDNA being analysed when measuring cytokine gene expression in a renal transplant population. It has not been possible to do that, as none of the three genes tested are suitable.

There were, however, no significant changes in the mean expression of any of the three chosen genes in the living donor group. This suggests that our method of RNA standardisation and cDNA transcription (described at the beginning of this chapter and in chapter 2) did give constant cDNA concentrations. The laboratory methods were the same for both the living donors and the transplant recipients, and the recruitment period for both groups of patients was the same. This means that any subsequent changes in cytokine gene expression detected in the next phase of the study were almost certainly genuine and not an artefact (discussed in more detail in chapter 6).

Chapter 5: Results of the Cytokine Gene Expression Work

This chapter will first describe the demographics of the three groups of patients within the study, namely the living donors, the non-rejecting and rejecting renal transplant recipients. Following this, the results of the cytokine gene expression work will be presented for all four cytokines studied (IL-4, IL-10, TGF- β 1 and TNF- α) in all three groups of patients. The results will be presented in both graphic and tabular forms. Included in the tables will be the results of the statistical analyses performed on the data obtained.

The results presented below will be discussed in the next chapter.

5.1 Patient Demographics

The living donor group contained 13 patients. The group experiencing an episode of acute allograft rejection contained 15 patients. The final group of non-rejecting transplant recipients was the largest group consisting of 30 patients. All rejection episodes were biopsy proven. Two patients experienced 2 episodes of acute rejection within the study period. All rejection episodes were initially treated with three doses of 0.5g of methyl prednisolone. Two patients with rejection went on to have a 14 day course of the monoclonal antibody OKT3. A single non-rejector experienced an episode of CyA toxicity during the study period. Three transplant recipients in each group experienced a period of biopsy proven acute tubular necrosis, all of which recovered within the study period of 14 days. In both cases the numbers were too small to be able to draw any useful conclusions from the effect of CyA toxicity and acute tubular necrosis on cytokine gene expression in the peripheral blood lymphocytes of the renal transplant recipients in this study.

Table 5.1a gives a summary of the causes of the end-stage renal failure in the 45 renal transplant recipients categorised into rejectors and non-rejectors. Table 5.1b gives a summary of the demographics of both the non-rejector and the rejector groups. Table 5.1c shows the distribution of the CMV status of the donor/recipient pairs and sex of the donor/recipient pairs in both groups of transplant recipients. Table 5.1d shows the distribution of immunosuppression regimes given to the two sets of recipients. Table 5.1e shows the demographics of the living donors.

Table 5.1a Cause of the end-stage renal failure in all 30 non-rejecting and 15 rejecting renal transplant recipients

Cause of Renal Failure	Non-Rejectors	Rejectors
Glomerulonephritides	8	2
Diabetes	5	3
Reflux Nepohropathy	4	2
Hypertension	3	-
Drug Toxicity	2	-
Unknown	2	-
Adult Polycystic Kidneys	1	3
IgA Nephropathy	2	-
Renal Dysplasia	1	1
Focal Segmental Glomerulosclerosis	1	2
Interstitial Nephritis	1	-
Henoch-Scholein Purpura	-	1
Posterior Urethral Valves	-	1

This table shows the cause of the end-stage renal failure in all 30 non-rejecting and 15 rejecting renal transplant recipients.

Table 5.1c. Demographics of the rejector and non-rejector recipient groups

Recipient Group	Mean Age	Sex of Recipient	Transplant Number	Form of Dialysis	Mean Mismatch (A: B: DR)	Mean Donor Age	Donor Sex	Type of Donor	Mean Cold Ischaemic Time
Non-Rejectors	40.7yrs	12F 18M	20 1 st (67%) 10 2 nd (33%)	5 PD 8 CAPD 17 HD	1.0: 1.2: 0.6	42.3yrs	12M 18F	4 LR 4 LUR 22 C	12.5Hr
Rejectors	37.6yrs	7F 8M	12 1 st (80%) 3 2 nd (20%)	4 PD 3 CAPD 8 HD	0.8: 1.2: 0.7	40.5yrs	8M 7F	3 LR 12 C	13.5Hr

This table shows the demographics of the rejector and non-rejector recipient groups. F = female. M = male. PD = pre-dialysis. CAPD = continuous ambulatory peritoneal dialysis. HD = haemodialysis. LR = living related. LUR = living unrelated. C = cadaveric. Hr = hours.

Table 5.1c. Distribution of donor/recipient CMV status and donor/recipient sex for both sets of renal transplant recipients

Donor to Recipient CMV Status	Donor to Recipient Sex	Non- Rejectors	Rejectors
Positive to positive	-	10 (33%)	5 (33%)
Positive to negative	-	8 (27%)	6 (40%)
Negative to positive	-	5 (17%)	1 (7%)
Negative to negative	-	7 (23%)	3 (20%)
-	Male to Male	5 (33%)	4 (13%)
-	Male to Female	3 (20%)	8 (27%)
-	Female to Male	3 (20%)	14 (47%)
-	Female to Female	4 (27%)	4 (13%)

This table shows the distribution of donor/recipient CMV status and donor/recipient sex for both sets of renal transplant recipients.

Table 5.1d. Distribution of immunosuppressives regimes used in the two groups of renal transplant recipients

Immunosuppressive Regime	Non-rejectors	Rejectors
Cyclosporin (Neoral) Prednisolone Azathioprine	12 (40%)	4 (27%)
Cyclosporin (Neoral) Prednisolone Rapamune	8 (27%)	3 (20%)
Cyclosporin (Neoral) Prednisolone Simulect	5 (17%)	1 (7%)
Cyclosporin (Neoral) Prednisolone MMF	3 (10%)	3 (20%)
Cyclosporin (Neoral) Prednisolone MMF & Simulect	2 (7%)	1 (7%)
Cyclosporin (Neoral) Prednisolone RAD Trial	-	3 (20%)

This table shows the distribution of immunosuppressives regimes used in the two groups of renal transplant recipients. MMF = mycophenolate mofetil

Table 5.1e. Type, mean age and sex of the living donors

Donor Type	Number	Mean Age	Sex
Living Related	9	43	M 0 F 9
Living Unrelated	4	50	M 1 F 3

This table shows the type, mean age and sex of the living donors.

5.2 Results of Cytokine Gene Expression Measurement

In this section the results of the cytokine gene expression measurement using Taqman real time quantitative PCR technology will be presented. This will be done initially in a graphical format using box and whisker plots and then in summary tables.

There will be four graphs for each cytokine. The first will show the results of the cytokine gene expression measurement from the living donors, the second will show the results from the non-rejectors and the final two for each cytokine will show the results from the rejectors. The first three graphs for each cytokine will be in the same format with the C_T value on the Y-axis and time on the X-axis. Time points 1 to 9 will be shown for the donors and time points 1 to 15 will be shown for the recipients. Time point 1 corresponds to the baseline value, either pre-nephrectomy, in the case of the living donors, or pre-transplant, in the case of the recipients. Time point 2 corresponds to the first post-operative day, time point 3 to the second post-operative day, and so on, in both groups of patients.

The fourth graph for each cytokine is different. The axes are the same with the C_T value on the Y-axis and time on the X-axis. However, the time points are different. These graphs again show the results of the cytokine gene expression measurement in the rejectors, but this time they are re-aligned around the day that the rejection was diagnosed and anti-rejection treatment was started. The

baseline (pre-transplant) value is still shown, then the value for each of the five days preceding the diagnosis of rejection (BR) are shown followed by the value for each of the five days after the initiation of anti-rejection treatment (PR). The order of the graphs interleukin-4, interleukin-10, transforming growth factor- β 1 and tumour necrosis factor- α .

The tables 5.2a – 5.2h show a summary of the results of the statistical analysis of the cytokine gene expression measurements for each cohort of patients. There are two tables each for the living donors and the non-rejectors, and four tables for the rejectors. The first tables, for all three groups, show the results of the stepwise statistical analysis. Using the Wilcoxon rank signed test, the mean value of each time point was compared with the mean value of the next time point. In other words, the mean baseline value was compared with the mean value from the first post-operative day, which was compared to the mean value from the second post-operative day etc. The second tables, for all three groups, show the results of the statistical analysis comparing the mean value of each successive time point with the mean baseline value for that particular cytokine. This analysis was again performed using the Wilcoxon rank signed test.

The third and fourth tables for the rejector cohort use the same basic format. The third table is a summary of the stepwise analysis and the fourth again compares each value with the baseline. However, like the graphs, these tables show the analysis when the results are re-aligned to the day the diagnosis of rejection was made.

Figure 5.2a. Box and whisker plot of the gene expression data of IL-4 in the living donors

This graph is a box and whisker plot and shows the results of the measurement of interleukin-4 gene expression in the living donor cohort. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-nephrectomy sample.

Figure 5.2b. Box and whisker plot of the gene expression data of IL-4 in the non-rejecting transplant recipients

This graph is a box and whisker plot and shows the results of the measurement of interleukin-4 gene expression in the non-rejecting transplant recipients. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-transplant sample.

Figure 5.2c. Box and whisker plot of the gene expression data of IL-4 in the rejecting transplant recipients

This graph is a box and whisker plot and shows the results of the measurement of interleukin-4 gene expression in the rejecting transplant recipients. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-transplant sample.

Figure 5.2d. Box and whisker plot of the gene expression data of IL-4 in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis

This graph is a box and whisker plot and shows the results of the measurement of interleukin-4 gene expression in the rejecting transplant recipients re-aligned to the day of rejection diagnosis. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted from left to right starting with the pre-transplant sample and then the five samples before rejection (BR5 –BR1) and the five days post-rejection (PR1-PR5).

Figure 5.2e. Box and whisker plot of the gene expression data of IL-10 in the living donors

This graph is a box and whisker plot and shows the results of the measurement of interleukin-10 gene expression in the living donor cohort. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-nephrectomy sample.

Figure 5.2f. Box and whisker plot of the gene expression data of IL-10 in the non-rejecting transplant recipients

This graph is a box and whisker plot and shows the results of the measurement of interleukin-10 gene expression in the non-rejecting transplant recipients. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-transplant sample.

Figure 5.2g. Box and whisker plot of the gene expression data of IL-10 in the rejecting transplant recipients

This graph is a box and whisker plot and shows the results of the measurement of interleukin-10 gene expression in the rejecting transplant recipients. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-transplant sample.

Figure 5.2h. Box and whisker plot of the gene expression data of IL-10 in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis

This graph is a box and whisker plot and shows the results of the measurement of interleukin-10 gene expression in the rejecting transplant recipients re-aligned to the day of rejection diagnosis. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted from left to right starting with the pre-transplant sample and then the five samples before rejection (BR5 –BR1) and the five days post-rejection (PR1-PR5).

Figure 5.2i. Box and whisker plot of the gene expression data of TGF- β 1 in the living donors

This graph is a box and whisker plot and shows the results of the measurement of transforming growth factor- β 1 gene expression in the living donor cohort. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-nephrectomy sample.

Figure 5.2j. Box and whisker plot of the gene expression data of TGF- β 1 in the non-rejecting transplant recipients

This graph is a box and whisker plot and shows the results of the measurement of transforming growth factor- β 1 gene expression in the non-rejecting transplant recipients. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-transplant sample.

Figure 5.2k. Box and whisker plot of the gene expression data of TGF- β 1 in the rejecting transplant recipients

This graph is a box and whisker plot and shows the results of the measurement of transforming growth factor- β 1 gene expression in the rejecting transplant recipients. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-transplant sample.

Figure 5.2l. Box and whisker plot of the gene expression data of TGF- β 1 in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis

This graph is a box and whisker plot and shows the results of the measurement of transforming growth factor- β 1 gene expression in the rejecting transplant recipients re-aligned to the day of rejection diagnosis. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted from left to right starting with the pre-transplant sample and then the five samples before rejection (BR5 – BR1) and the five days post-rejection (PR1-PR5).

Figure 5.2m. Box and whisker plot of the gene expression data of TNF- α in the living donors

This graph is a box and whisker plot and shows the results of the measurement of tumour necrosis factor- α gene expression in the living donor cohort. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-nephrectomy sample.

Figure 5.2n. Box and whisker plot of the gene expression data of TNF- α in the non-rejecting transplant recipients

This graph is a box and whisker plot and shows the results of the measurement of tumour necrosis factor- α gene expression in the non-rejecting transplant recipients. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-transplant sample.

Figure 5.2o. Box and whisker plot of the gene expression data of TNF- α in the rejecting transplant recipients

This graph is a box and whisker plot and shows the results of the measurement of tumour necrosis factor- α gene expression in the rejecting transplant recipients. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted sequentially from left to right starting with the pre-transplant sample.

Figure 5.2p. Box and whisker plot of the gene expression data of TNF- α in the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis

This graph is a box and whisker plot and shows the results of the measurement of tumour necrosis factor- α gene expression in the rejecting transplant recipients re-aligned to the day of rejection diagnosis. The C_T value is plotted on the Y-axis. Time is plotted on the X-axis. The top set of numbers along the X-axis is the number of samples measured at each time point. The lower set is the time points plotted from left to right starting with the pre-transplant sample and then the five samples before rejection (BR5 –BR1) and the five days post-rejection (PR1-PR5).

Table 5.2a. Results of stepwise analysis of the cytokine gene expression data of the living donors

Time points in days in sequential steps (1 = pre-nephrectomy, 2 = day 1 post-nephrectomy etc.)		1-2	2-3	3-4	4-5	5-6	6-7	7-8	8-9
Interleukin-4	Direction of change of gene expression	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS
Interleukin-10	Direction of change of gene expression	↑	-	-	-	-	-	-	-
	p value	0.016	NS	NS	NS	NS	NS	NS	NS
Transforming growth factor- β 1	Direction of change of gene expression	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS
Tumour necrosis factor- α	Direction of change of gene expression	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS

This table shows the stepwise direction of change of the mean gene expression data for the four cytokine genes in the living donor cohort and the level of significance of these changes. Statistical analysis performed using Wilcoxon's rank signed test. NS = non-significant.

Table 5.2b. Results of the comparison to baseline analysis of the cytokine gene expression data of the living donors

Time points in days comparing the pre-nephrectomy value with each subsequent value (1 = pre-nephrectomy, 2 = day 1 post-nephrectomy etc.)		1-2	1-3	1-4	1-5	1-6	1-7	1-8	1-9
Interleukin-4	Direction of change of gene expression	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS
Interleukin-10	Direction of change of gene expression	↑	↑	-	-	-	-	-	-
	p value	0.016	0.003	NS	NS	NS	NS	NS	NS
Transforming growth factor-β1	Direction of change of gene expression	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS
Tumour necrosis factor-α	Direction of change of gene expression	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS

This table shows the direction of change of the mean gene expression data of each time point compared to the baseline (pre-nephrectomy) value for the four cytokine genes in the living donor cohort and the level of significance of these changes. Statistical analysis performed using Wilcoxon's rank signed test. NS = non-significant.

Table 5.2c. Results of stepwise analysis of the cytokine gene expression data of the non-rejecting transplant recipients

Time points in days in sequential steps (1 = pre-transplant, 2 = day 1 post-transplant etc.)		1-2	2-3	3-4	4-5	5-6	6-7	7-8	8-9	9-10	10-11	11-12	12-13	13-14	14-15
Interleukin-4	Direction of change of gene expression	↓	-	-	-	-	-	-	-	↑	-	-	-	-	-
	p value	<0.001	NS	NS	NS	NS	NS	NS	NS	0.017	NS	NS	NS	NS	NS
Interleukin-10	Direction of change of gene expression	↑	-	-	-	-	-	-	-	-	-	-	-	-	-
	p value	<0.001	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Transforming growth factor-β1	Direction of change of gene expression	↑	-	-	-	-	-	↓	-	-	-	-	-	-	-
	p value	<0.001	NS	NS	NS	NS	NS	0.05	NS	NS	NS	NS	NS	NS	NS
Tumour necrosis factor-α	Direction of change of gene expression	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

This table shows the stepwise direction of change of the mean gene expression data for the four cytokine genes in the non-rejecting transplant recipients and the level of significance of these changes. Statistical analysis performed using Wilcoxon's rank signed test. NS = non-significant.

Table 5.2d. Results of the comparison to baseline analysis of the cytokine gene expression data of the non-rejecting transplant recipients

Time points in days comparing the pre-transplant value with each subsequent value		1-2	1-3	1-4	1-5	1-6	1-7	1-8	1-9	1-10	1-11	1-12	1-13	1-14	1-15
Interleukin-4	Direction of change of gene expression	↓	↓	↓	↓	-	-	-	-	-	-	-	-	-	-
	p value	<0.001	0.003	0.015	0.008	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Interleukin-10	Direction of change of gene expression	↑	↑	↑	↑	-	↑	-	-	-	-	-	-	-	-
	p value	<0.001	0.005	0.029	0.05	NS	0.025	NS	NS	NS	NS	NS	NS	NS	NS
Transforming growth factor- β 1	Direction of change of gene expression	↑	-	↑	-	-	↑	↑	↑	↑	↑	-	↑	-	-
	p value	<0.001	NS	0.031	NS	NS	0.005	0.032	0.013	0.019	0.036	NS	0.039	NS	NS
Tumour necrosis factor- α	Direction of change of gene expression	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

This table shows the direction of change of the mean gene expression data of each time point compared to the baseline (pre-transplant) value for the four cytokine genes in the non-rejecting transplant recipients and the level of significance of these changes. Statistical analysis performed using Wilcoxon's rank signed test. NS = non-significant.

Table 5.2e. Results of stepwise analysis of the cytokine gene expression data of the rejecting transplant recipients

Time points in days in sequential steps (1 = pre-transplant, 2 = day 1 post-transplant etc.)		1-2	2-3	3-4	4-5	5-6	6-7	7-8	8-9	9-10	10-11	11-12	12-13	13-14	14-15
Interleukin-4	Direction of change of gene expression	↓	↑	-	-	-	↓	↑	-	-	-	↑	-	-	-
	p value	0.036	0.034	NS	NS	NS	0.043	0.019	NS	NS	NS	0.018	NS	NS	NS
Interleukin-10	Direction of change of gene expression	↑	-	-	-	-	-	-	-	-	-	-	-	-	-
	p value	0.002	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Transforming growth factor- β 1	Direction of change of gene expression	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Tumour necrosis factor- α	Direction of change of gene expression	-	-	-	↓	-	↑	-	-	-	-	-	-	-	-
	p value	NS	NS	NS	0.019	NS	0.016	NS	NS	NS	NS	NS	NS	NS	NS

This table shows the stepwise direction of change of the mean gene expression data for the four cytokine genes in the rejecting transplant recipients and the level of significance of these changes. Statistical analysis performed using Wilcoxon's rank signed test. NS = non-significant.

Table 5.2f. Results of the comparison to baseline analysis of the cytokine gene expression data of the rejecting transplant recipients

Time points in days comparing the pre-transplant value with each subsequent value		1-2	1-3	1-4	1-5	1-6	1-7	1-8	1-9	1-10	1-11	1-12	1-13	1-14	1-15
Interleukin-4	Direction of change of gene expression	↓	-	-	-	-	-	-	-	-	-	-	-	-	-
	p value	0.036	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Interleukin-10	Direction of change of gene expression	↑	↑	↑	-	↑	↑	↑	↑	-	↑	↑	-	↑	↑
	p value	0.002	0.016	0.001	NS	0.013	0.005	0.002	0.023	NS	0.05	0.017	NS	0.005	0.022
Transforming growth factor- β	Direction of change of gene expression	-	-	↑	-	-	↑	↑	-	-	↑	↑	-	↑	↑
	p value	NS	NS	0.001	NS	NS	0.016	0.016	NS	NS	0.033	0.013	NS	0.004	0.041
Tumour necrosis factor- α	Direction of change of gene expression	-	-	↑	-	-	↑	↑	↑	-	↑	↑	-	↑	↑
	p value	NS	NS	0.001	NS	NS	0.013	0.005	0.033	NS	0.05	0.022	NS	0.008	0.005

This table shows the direction of change of the mean gene expression data of each time point compared to the baseline (pre-transplant) value for the four cytokine genes in the rejecting transplant recipients and the level of significance of these changes. Statistical analysis performed using Wilcoxon's rank signed test. NS = non-significant.

Table 5.2g. Results of stepwise analysis of the cytokine gene expression data of the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis

Time points in days in sequential steps (BR5 = 5 days before diagnosis of rejection, BR4 = 4 days before diagnosis of rejection etc.)		PTBS- BR5	BR5- BR4	BR4- BR3	BR3- BR2	BR2- BR1	BR1- PR1	PR1- PR2	PR2- PR3	PR3- PR4	PR4- PR5
Interleukin-4	Direction of change of gene expression	-	-	-	↑	-	↓	-	-	↑	-
	p value	NS	NS	NS	0.001	NS	0.011	NS	NS	0.028	NS
Interleukin-10	Direction of change of gene expression	↑	-	↑	-	↓	-	-	-	-	-
	p value	0.050	NS	0.041	NS	0.046	NS	NS	NS	NS	NS
Transforming growth factor- β	Direction of change of gene expression	-	-	-	↑	↓	-	↓	-	-	-
	p value	NS	NS	NS	0.008	0.016	NS	0.028	NS	NS	NS
Tumour necrosis factor- α	Direction of change of gene expression	-	-	-	↑	↓	-	-	-	-	-
	p value	NS	NS	NS	0.006	0.004	NS	NS	NS	NS	NS

This table shows the stepwise direction of change of the mean gene expression data, when re-aligned to the time of rejection, for the four cytokine genes in the rejecting transplant recipients and the level of significance of these changes. Statistical analysis performed using Wilcoxon's rank signed test. NS = non-significant. PTBS = pre-transplant baseline. BR = before rejection. PR = post-rejection.

Table 5.2h. Results of the comparison to baseline analysis of the cytokine gene expression data of the rejecting transplant recipients with the data re-aligned to the day of rejection diagnosis

Time points in days comparing the pre-transplant value (PTBS) with each value before (BR) and post (PR) the diagnosis of rejection.		PTBS-BR5	PTBS-BR4	PTBS-BR3	PTBS-BR2	PTBS-BR1	PTBS-PR1	PTBS-PR2	PTBS-PR3	PTBS-PR4	PTBS-PR5
Interleukin-4	Direction of change of gene expression	-	-	-	↑	↑	-	↓	-	-	-
	p value	NS	NS	NS	0.009	0.033	NS	0.025	NS	NS	NS
Interleukin-10	Direction of change of gene expression	↑	↑	↑	↑	↑	↑	↑	↑	↑	-
	p value	0.050	0.008	0.002	0.001	0.006	0.013	0.021	0.036	0.037	NS
Transforming growth factor-β	Direction of change of gene expression	-	↑	↑	↑	↑	↑	-	-	-	↑
	p value	NS	0.016	0.030	0.001	0.001	0.010	NS	NS	NS	0.021
Tumour necrosis factor-α	Direction of change of gene expression	-	↑	↑	↑	↑	-	-	-	-	↑
	p value	NS	0.016	0.008	0.001	0.001	NS	NS	NS	NS	0.021

This table shows the direction of change of the mean gene expression data, when re-aligned to the time of rejection, of each time point compared to the pre-transplant baseline value for the four cytokine genes in the rejecting transplant recipients and the level of significance of these changes. Statistical analysis performed using Wilcoxon's rank signed test. NS = non-significant. PTBS = pre-transplant baseline. BR = before rejection. PR = post-rejection.

The results of the cytokine gene expression measurement from the living donors and transplant recipients shown in figures 5.2a to 5.2p and tables 5.2a to 5.2h demonstrate significant changes in the expression of all four cytokines. Statistically significant changes are not seen in all cohorts. Changes that are significant vary in both magnitude and direction between different cytokines and different cohorts.

These results and there possible implications are discussed in the next chapter.

Chapter 6: Discussion

In this chapter the results presented in the previous chapter will be discussed, dealing with each individual cytokine in turn. The findings of this study will be compared with the results from the earlier RT-PCR ELISA work from our laboratory and similar work from other centres. Finally, conclusions will be drawn as to the utility of monitoring expression of individual cytokines as a marker of immunological events, including acute allograft rejection in renal transplant recipients.

6.1 Interleukin-4

The IL-4 gene is located on chromosome 5. It encodes a multifunctional anti-inflammatory cytokine that is mainly produced by activated T lymphocytes, but also mast cells, and is central to the generation and regulation of immune responses (Nelms 1999, Chomarat 1998). IL-4 promotes type 2 T helper (Th2) cell differentiation and stability and antagonises the effects of IFN- γ on type 1 T helper (Th1) lymphocytes and macrophages (Paludan 1998). Biopsy specimens from transplanted kidneys have been analysed to detect the presence of IL-4 gene transcripts. These have been reported in both stable and rejecting grafts (see below), therefore, the role of this cytokine in the rejection process is not clear. Work from our own unit studying the IL-4 -590 gene promoter polymorphism showed that donor/recipient pairs both lacking the IL-4-T (high expresser) allele were significantly increased in the rejectors suggesting that an ability to produce increased quantities of IL-4 may be important in controlling the processes leading to acute rejection or tolerance (Poole 2001), i.e. low expression = rejection and high expression = tolerance. This is in keeping with IL-4 being an anti-inflammatory and Th 2 type cytokine. These latter results, combined with those from the previous RT-PCR ELISA, encouraged us to study this cytokine further.

The results of the RT-PCR ELISA study measuring IL-4 gene expression in T lymphocytes obtained from the peripheral blood of 43 renal transplant recipients, and confirmed using quantitative TaqMan technology, were described in chapter 3. This study demonstrated differences in IL-4 gene

expression between renal transplant recipients experiencing an episode of acute allograft rejection when compared with those who had an uneventful early post-transplant recovery. The statistically significant fall in IL-4 gene expression detected for a week following renal transplantation in non-rejectors was not maintained in the rejectors, but was re-established following anti-rejection therapy.

In this study, IL-4 gene expression was again measured in peripheral blood cells obtained from renal transplant recipients in the early post-operative period. This time, however, the extraction process was simplified, and gene expression was measured in the whole mononuclear cell subset. In addition to the renal transplant recipients, blood samples were also obtained from living donors both before and following the nephrectomy. This enabled us to study the potential influence of surgery alone on IL-4 gene expression.

The results of the measurement of IL-4 gene expression in the living donors were shown graphically in figure 5.2a and the results of the statistical analysis summarised in tables 5.2a and b. Expression did not appear to be influenced by surgery. There were no significant changes detected in IL-4 gene expression in the living donors between both the pre-operative baseline value and any of the post-operative values, or sequential day to day values. This confirms that the IL-4 findings of the RT-PCR ELISA study are real changes influenced by non-surgical immunological processes. It also suggests that any significant findings identified in the IL-4 gene expression of the transplant

recipients in this study are also the result of immunosuppressive treatment or immunological events.

The results of the measurement of IL-4 gene expression in the non-rejecting and rejecting renal transplant recipients were shown graphically in figure 5.2b and the results of the statistical analysis summarised in tables 5.2c and d. IL-4 gene expression did change significantly following renal transplantation in this cohort. As in the RT-PCR ELISA study, there was a significant reduction in IL-4 gene expression seen immediately following transplantation. This reduction was detected for the first five days in this study, after which the levels returned to baseline values. There was a significant increase in IL-4 gene expression detected between the eighth and ninth post-operative days, but neither of the individual day values was significantly different from the pre-transplant baseline value.

The decrease in IL-4 gene expression detected was large, corresponding to more than a 3-cycle rise in C_T value. This equates to at least an 8-fold reduction in expression of the IL-4 gene by the peripheral blood mononuclear cells within the first 24 hours following renal transplantation. This reduction is unlikely to be the result of the surgical insult, as there was no such change seen in the donors. However, it may be due to direct immunosuppressive action on these cells of the induction therapy given immediately prior to transplantation (Neoral 8mg/kg and 0.5g methyl prednisolone). The other possibility is that this apparent reduction in IL-4 gene expression may be the result of sequestration of cells with increased IL-4 gene expression into either

lymph nodes or the graft itself, leaving the quiescent or low expresser cells in the peripheral blood.

The results of the measurement of IL-4 gene expression in the rejecting renal transplant recipients were shown graphically in figures 5.2c and d and the results of the statistical analysis summarised in tables 5.2e, f, g and h. In this group there was also a significant fall in IL-4 gene expression detected on the first day post-transplant. This reduction in expression was at least as great as that seen in the non-rejectors equating to a 4-cycle or 16-fold decrease in IL-4 gene expression. However, this reduction in expression was followed by a significant increase in expression returning the value from the second post-transplant day back to a level that was not significantly different from the pre-transplant baseline value for IL-4.

Other significant changes were detected in IL-4 gene expression in the rejector group between days 5 and 6, 6 and 7, and 10 and 11. As the day on which the diagnosis of rejection was made and anti-rejection therapy varied from patient to patient, these results were difficult to interpret. To attempt to clarify the influence that acute allograft rejection and subsequent anti-rejection therapy may have on IL-4 gene expression the results were re-aligned with respect to the day acute allograft rejection was diagnosed.

When the expression data were re-analysed from the five days prior to the diagnosis and the five days following the initiation of anti-rejection therapy a coherent pattern emerged. There was a significant increase in IL-4 gene

expression, compared with the pre-transplant baseline value, detected for the two days immediately prior to the diagnosis of acute allograft rejection. Following the start of anti-rejection therapy, there was a significant sequential reduction in IL-4 gene expression when compared with both the pre-treatment and the pre-transplant values. This 6-cycle reduction approximately equates to 64-fold decrease in IL-4 gene expression over a two-day period. The cessation of anti-rejection treatment was followed by a significant increase in IL-4 gene expression returning the value to a pre-transplant baseline value.

The changes detected in IL-4 gene expression correlate well with clinical events in patients experiencing acute allograft rejection following renal transplantation. IL-4 gene expression initially falls following transplantation, in rejectors, in a similar fashion to non-rejectors, but returns to a baseline value by the second post-transplant day. This change and the changes detected following anti-rejection treatment are very similar to the findings of the RT-PCR ELISA work. However, the RT-PCR ELISA work failed to show a significant rise in IL-4 gene expression at the time of rejection.

High dose immunosuppression appears to result in a substantial reduction in IL-4 gene expression detectable in peripheral blood mononuclear cells, as seen by the fall in expression following induction therapy immediately post-transplant and following anti-rejection therapy.

Groups other than our own have studied IL-4 levels in renal transplant recipients experiencing acute allograft rejection. Some have been single time

point studies comparing intragraft IL-4 gene expression with clinical rejection or tolerance. Loong and colleagues detected increased levels of IL-4 in specimens from kidneys experiencing acute allograft rejection (Loong 1996). Krams et al detected increased levels of IL-4 mRNA in biopsy specimens from both rejecting and rejected kidneys, but did not compare these results with samples from stable grafts (Krams 1992). However, other groups looking for intragraft IL-4 expression in rejecting renal transplants have been unable to detect elevated levels (Strehlau 1997, Xu 1995). Joseph and co-workers were only able to detect IL-4 gene expression in kidneys not experiencing acute rejection, but not all stable grafts expressed IL-4 gene transcripts (Joseph 1995).

One group has studied the gene expression of IL-2, IL-4, IL-6, IFN- γ and TGF- β 1 in peripheral blood lymphocytes and core biopsies from liver transplant recipients. Unfortunately, as with so many of these studies, the sampling was at single time points. However, the peripheral blood lymphocyte samples were taken at the same time as the liver biopsies were performed. Using semi-quantitative RT-PCR technology, they were unable to detect a significant difference in IL-4 message expression between the rejectors and the non-rejectors for both liver biopsies and peripheral blood lymphocytes. IL-4 transcripts were detected in the vast majority of core biopsies irrespective of the rejection status. However, in the absence of rejection IL-4 transcription was noted with significantly more frequency than IL-2 transcription. In addition to this observation, they were able to report a good correlation between intragraft and peripheral blood lymphocyte cytokine

profiles, suggesting that 'lymphocyte populations regulating graft rejection were predominant both locally (intragraft) and peripherally (peripheral blood lymphocytes)' (Gorczynski 1996).

Other studies have measured serum IL-4 levels before and after renal transplantation. In one study daily serum samples were collected from 16 renal transplant recipients for the first 14 post-operative days. Seven of these experienced an episode of acute allograft rejection. Five of the seven had elevated levels of serum IL-4 prior to the diagnosis of rejection, while the remaining two had elevated levels detected after the diagnosis was made. Levels fell after treatment (Kutukculer 1995). In a different study, serum IL-4 levels were measured in patients experiencing either acute viral infections or acute allograft rejection. Neither group had a significant elevation in serum IL-4 levels prior to their diagnosis (Daniel 1995).

Other groups have measured serum IL-4 levels in non-transplant patients before and after surgery. In a series of 32 patients undergoing coronary artery bypass surgery, there was no change in IL-4 levels either during surgery or in the immediate post-operative period (Nathan 2000). Another study measured serum IL-4 levels before and after bowel resections for Crohn's disease in 22 patients. No rise in IL-4 levels was detected in any of the patients (Hildebrandt 1999).

Steroids have been shown to down regulate IL-4 mRNA in patients suffering from allergic rhinitis. In a study using an in situ hybridisation technique there

was a significant reduction in IL-4 mRNA levels in the group receiving topical steroid treatment compared to controls (Kondo 1999).

By intensively monitoring renal transplant recipients during the first two weeks following transplantation, and by using quantitative TaqMan PCR technology, we have confirmed that IL-4 gene expression in peripheral blood cells correlates well with clinical events in renal transplant recipients. There was a large and significant reduction in IL-4 gene expression immediately following renal transplantation that was maintained for 5 days in patients who do not experience a rejection episode. In those experiencing an acute rejection episode there was a significant rise in IL-4 gene expression prior to the diagnosis of the rejection and subsequently a large decrease in expression following anti-rejection therapy. These changes were not the result of surgery, as IL-4 gene expression levels did not vary in the living donors who underwent nephrectomies in our study. Other groups have reported similar results. Measuring serum IL-4 levels in non-transplant patients, Nathan and Hildebrandt were also unable to detect changes in levels of this cytokine following surgery (Nathan 2000, Hildebrandt 1999). The fall in IL-4 gene expression following transplantation and acute rejection may be due to the administration of steroids as this has been shown to reduce IL-4 expression in other studies (Kondo 1999).

Work from our own laboratory was performed in an effort to identify a link between the -590 IL-4 promoter polymorphism and the incidence and severity of acute rejection episodes in renal transplant recipients. The results of this

work suggested that the high expresser genotype might be protective against rejection as a lack of this genotype in both the donor and recipients of a renal transplant was more common in rejectors. This might suggest that one would expect to see increased IL-4 gene expression in the non-rejectors with an opposite pattern in the rejectors. However, in this study IL-4 gene expression is seen to fall, and then return to baseline values, but not to increase, in non-rejectors. One explanation for these observations is the sequestration of IL-4 producing cells into the graft in the early transplant period, resulting in a relative decrease in the IL-4 gene expression in the remaining peripheral blood mononuclear cells. The presence of the increased numbers of anti-inflammatory IL-4 producing cells in the graft preventing rejection. In the rejectors the role of IL-4 producing cells may be completely the opposite. In this situation the increased IL-4 gene expression is in response to the inflammatory acute rejection process, hence the rise detected prior to the initiation of anti-rejection treatment. This may explain why some studies have reported the presence of IL-4 gene transcripts in both rejectors and non-rejectors. IL-4 producing cells may be present in both situations, but performing different tasks.

These findings suggest that IL-4 is an important cytokine in acute allograft rejection in renal transplant recipients. The detrimental effect of a low expresser genotype, the observation that IL-4 gene expression increased preceding rejection and fell following anti-rejection therapy, and finally that the IL-4 gene expression pattern was different in non-rejectors compared with

rejectors implies that IL-4 might be involved in the immune processes involved in the cascade of events that result in acute allograft rejection.

In summary, IL-4 gene expression in peripheral blood mononuclear cells correlates well with intragraft events and anti-rejection therapy in the first two weeks following renal transplantation. Distinct expression patterns for rejectors and non-rejectors were demonstrable confirming the findings of the RT-PCR ELISA work. These differences in expression pattern precede the clinical and biochemical detection of acute rejection. Also, after successful anti-rejection therapy IL-4 gene expression levels initially rise above and then return to pre-transplant levels in a similar fashion to the early expression pattern seen in non-rejectors. Therefore, measurement of IL-4 gene expression in peripheral blood of renal transplant recipients may be of benefit in monitoring these patients both before rejection is confirmed and then after to monitor the effectiveness of treatment.

6.2 Interleukin-10

The IL-10 gene is located on chromosome 1. It encodes a ubiquitous cytokine with strong anti-inflammatory effects produced by many different cell types including lymphocytes, monocytes and macrophages. It has a diverse array of actions, which differ depending on cell type, nature of stimulus and the cellular microenvironment. However, it does play a major regulatory role in the course of the inflammatory responses by inhibiting the expression of genes encoding pro-inflammatory cytokines. It is a stress cytokine produced in response to cellular injury (Stordeur 1998, Pretlani 1999, Baud 1998, Donnelly 1999). IL-10 has been detected in core biopsy specimens from grafts experiencing acute rejection (see below), but it has not been established whether the presence of this cytokine indicates a role in initiating the process or a response to it. These observations have been compounded by work investigating the effect of IL-10 gene promoter polymorphisms. In cardiac transplant recipients the low IL-10 expresser genotype (-1082A/A) is more common in patients having multiple rejection episodes (Turner/Grant 1997), whereas in kidney transplant recipients the high IL-10 expresser genotype (-1082G/G) was associated with multiple rejection episodes (Sankaran 1999). In liver transplants the IL-10 genotype did not appear to influence outcome (Bathgate 2000). These findings, and those from the RT-PCR ELISA work from our laboratory, justified further study of this cytokine.

In the RT-PCR ELISA study described in chapter 3, differences in IL-10 gene expression were detected between the rejectors and non-rejectors. Both groups

showed a significant increase in IL-10 gene expression, compared to the pre-transplant baseline value, on the second post-operative day. In the non-rejectors this difference was maintained throughout the study period. In the rejectors there was a relative fall in IL-10 gene expression at the time of rejection as the levels returned to baseline value. Following anti-rejection treatment, the levels again increased, re-establishing the difference in expression seen after two days. However, when the samples from the RT-PCR ELISA study were re-analysed using quantitative Taqman technology the pattern was different. The statistically significant rise seen in IL-10 gene expression was maintained at the time of rejection. This meant that there was no difference between the expression patterns detected in the rejectors and non-rejectors. This finding suggested that the differences originally detected in IL-10 gene expression between the rejector and non-rejector groups, in the RT-PCR ELISA study, may have resulted from difficulties with a more technically demanding semi-quantitative laboratory method rather than true changes in gene expression.

As for IL-4, the IL-10 gene expression in this study was measured in the mononuclear cells obtained from the peripheral blood of both transplant recipients and living donors undergoing native nephrectomies. The results of the measurement of IL-10 gene expression in the living donors were shown graphically in figure 5.2e and the results of the statistical analysis summarised in tables 5.2a and b.

Unlike IL-4, IL-10 gene expression did appear to be influenced by surgery. There was a significant rise in IL-10 gene expression detected for the first two days following surgery in the samples obtained from the living donors. The 2 cycle increase was equivalent to an approximate 4-fold rise in gene expression, and was not seen to start to fall until the fourth post-operative day. The observation that IL-10 gene expression was influenced by surgery alone may alter the significance of the findings from the RT-PCR ELISA study and others where the measurement of IL-10 levels has been undertaken in transplant recipients experiencing early rejection episodes.

A similar picture was seen for IL-10 gene expression in the group of transplant recipients experiencing no rejection. (The results of the measurement of IL-10 gene expression in the non-rejecting were shown graphically in figure 5.2f and the results of the statistical analysis summarised in tables 5.2c and d.) There was a significant increase in IL-10 gene expression for most of the first week following transplantation. This initial 4 cycle increase in expression was larger than that seen in the living donors. It would appear that the initial rise in IL-10 gene expression is the result of surgical trauma and unlikely to be the result of an independent immunological process. However, the induction immunosuppression or immunological processes may influence the duration and magnitude of the significant elevation seen.

In the rejector group the IL-10 gene expression pattern was repeated. (The results of the measurement of IL-10 gene expression in rejecting renal

transplant recipients were shown graphically in figures 5.2g and h and the results of the statistical analysis summarised in tables 5.2e, f, g and h.) There was a significant increase in expression from the pre-transplant value to the value from the first post-transplant day. This was comparable with the rise in expression detected in the non-rejectors. There were no further significant changes from day to day, but the expression level stayed significantly higher than the pre-transplant baseline level for the majority of the study period. On the three days the value was not significantly higher than baseline (fourth, ninth and twelfth) the sample numbers on these particular days were lower, which may account for the lack of significance.

The similarity in the results from the three groups suggests that the initial increase in expression seen is the result of surgery and is unlikely to be the result of any transplant related immunological process. However, the duration of this significant increase in IL-10 gene expression increased from 2 days in the donors, to 6 days in the non-rejecting recipients and finally through all 14 days in the rejectors. Prolongation of the increased IL-10 gene expression in the recipients, particularly those experiencing acute rejection could either be due to the acute rejection process or the anti-rejection treatment. To study this further, the IL-10 gene expression data were re-aligned to the day of rejection diagnosis and the subsequent commencement of anti-rejection therapy (figure 5.2h and tables 5.2g and h). When the mean values were compared to the pre-transplant baseline value, all but the fifth post-rejection value were significantly higher, which is very similar to the findings of the sequential analysis. However, when the day-to-day variation was compared, there was a

significant increase in IL-10 gene expression between the fourth and third days before rejection is diagnosed. Also, there was a significant fall in expression between the second day before rejection and the day of diagnosis. However, both these changes occurred with all expression levels being significantly greater than the pre-transplant baseline value. No significant changes occurred after the initiation of anti-rejection treatment.

Unlike IL-4, high dose steroids did not appear to influence level of IL-10 gene expression to any significant extent, but may influence the duration of the increased expression detected. The most significant change in expression appears to be the result of the surgery. However, the immunological processes that result in acute allograft rejection may have some influence on IL-10 gene expression. An increase in expression was seen preceding rejection, which may result reflect a response to increasing expression of pro-inflammatory mediators. The subsequent fall in expression detected before the diagnosis of rejection was made may be the result of either sequestration of IL-10 secreting cells into the graft or a result of negative feed back from pro-inflammatory cytokines. However, these changes in expression are probably less significant than the overall pattern of increased IL-10 gene expression post-transplant compared with pre-transplant values.

Several groups have identified a link between IL-10 gene expression and acute allograft rejection in kidney transplant recipients. As well as detecting increased IL-4 gene levels, Loong et al also detected raised intragraft IL-10 gene expression levels in core biopsies from 10 kidneys experiencing acute

rejection episodes (Loong 1996). Xu and co-workers reported similar findings when studying 98 core biopsies from renal transplants. Intragraft IL-10 gene expression significantly correlated with acute allograft rejection (Xu 1995). Strehlau and colleagues also studied intragraft IL-10 gene expression, this time, in 60 renal transplant core biopsies. They reported similar findings with heightened intragraft IL-10 gene expression significantly correlating with acute rejection (Strehlau 1997). Not all groups, however, have been able to identify a difference in IL-10 gene expression when comparing rejecting with non-rejecting renal transplant biopsy specimens. Joseph et al studied the intragraft gene expression of several cytokines, including IL-10, in core biopsies from 50 renal transplant recipients. They reported that intragraft IL-10 gene expression showed no variation with graft rejection and it was not a predictor of stable function (Joseph 1995).

When used to monitor transplant recipients, serum IL-10 has not been shown to be of predictive value in the diagnosis of acute rejection, but has been detected following the additional treatment given for acute rejection. Daniel et al reported a significant rise in serum IL-10 levels within the first few days of the commencement of anti-rejection treatment in renal transplant recipients experiencing an acute rejection episode (Daniel 1995). In cardiac transplant recipients no association was detected between serum IL-10 levels and acute rejection (Grant 1996).

Steroid pre-treatment has been shown to enhance IL-10 production in patients undergoing cardiac surgery, including heart and heart and lung transplantation

(Wan 1997). In another study in patients having cardiac surgery, high dose methyl prednisolone was shown to reduce the production of IL-6 and IL-8 but not IL-10 (Kawamura 1999).

Several studies have detected elevated levels of IL-10 in the serum of patients following surgery. In a series of 34 patients undergoing either an open or laparoscopic Nissen's fundoplication, elevated levels of IL-10 were detected in the serum within a few hours of the procedure and remained elevated for 48 hours (Zieren 2000). Another study measured serum IL-10 levels before and after bowel resections for Crohn's disease in 22 patients. Elevated levels of IL-10 were also detected in this group following the surgery (Hildebrandt 1999). Pre-operative serum levels of IL-10 were compared with post-operative levels in a series of 23 patients operated on for abdominal aortic aneurysms. IL-10 levels were significantly elevated immediately following the surgery. A week after surgery the levels were lower, but still significantly higher than baseline measurements (Holmberg 1999). In a study of 12 patients having major upper gastrointestinal surgery, elevated levels of systemic IL-10 were detected up to 66 hours after the surgery (van Berge Henegouwen 1998).

Not all studies, however, have shown that the increase in serum IL-10 following surgery is maintained beyond 24 hours. In one series of 32 patients undergoing coronary artery bypass surgery, there was a 117-fold increase in serum IL-10 levels in the immediate post-operative period, but levels had returned to pre-operative levels by the following day (Nathan 2000). In another study, comparing cytokine responses in 14 patients having

laparoscopic cholecystectomies with 14 patients undergoing open cholecystectomies, no elevation in IL-10 levels were detected at either 24 hours or 7 days after surgery (Bellon 1997).

In this study we have detected a significant rise in IL-10 gene expression following both living donor nephrectomy and renal transplantation. Results confirm the observations of the repeat TaqMan analysis of the samples from the RT-PCR ELISA study. These findings suggest that the differences originally identified between the gene expression levels of the non-rejectors and rejectors may have been due to the semi-quantitative method used rather than actual changes in IL-10 gene expression and that IL-10 gene expression may not be a good candidate for monitoring transplant patients.

The findings in this study are in agreement with other studies that have identified elevated serum IL-10 levels following various surgical procedures. The duration of these elevated levels were greatest in the patients experiencing acute rejection episodes. Those transplant recipients without rejection also had elevated levels for longer than the donors did. This difference may be related to the methyl prednisolone given as a single dose to all recipients at the time of surgery. It is also given, in multiple doses (3 x 0.5g on successive days), to those experiencing acute allograft rejection. Steroids have been reported not to reduce IL-10 serum levels following surgery, while other cytokine levels have fallen, and steroids may actually enhance IL-10 production.

Studies investigating the affect of the IL-10 promoter polymorphism on rejection following organ transplantation have produced conflicting results. One study suggested that the high IL-10 producer genotypes were at increased risk of multiple acute rejection episodes following renal transplantation (Sankaran 1999). Other workers report the opposite to be true. In heart transplantation the high IL-10 producer genotype appeared to be protective (Turner/Grant 1997). This finding has been corroborated, by some, in renal transplant recipients (George 2000, Kerman 2000), and is supported, at least in part, by work from our own laboratory. We have reported that when the recipient had the high IL-10 producer genotype (and the donor had the low IL-10 producer genotype) they experienced significantly fewer multiple rejection episodes (Poole 2001). It has also been reported that the -1082 IL-10 promoter polymorphism genotype does not appear to have any affect on rejection episodes in renal transplant recipients (Young 2000, Marshall/McLaren 2000). Therefore, the influence of IL-10 promoter polymorphisms may differ according to the organ transplanted, although there are also other possible confounding variables between the published studies such as immunosuppression regimes.

In summary, this study detected significant changes in IL-10 gene expression in peripheral blood mononuclear cells in the first week following native nephrectomy and in the first 14 days following renal transplantation. The most striking of these was the initial rise in IL-10 gene expression following transplantation. This increase in expression appeared to be prolonged by the administration of high dose, intravenous methyl prednisolone. IL-10 gene

expression does not correlate well with intragraft immunological events. Although there were significant changes detected prior to the diagnosis of acute rejection they are of limited use for the purposes of immunological monitoring. The reason being that the gene expression levels before and after the changes were all statistically significantly elevated compared to pre-transplant levels. Also, the initiation of anti-rejection treatment did not influence expression levels. They continued to be elevated both before treatment, during treatment and after treatment had ceased.

In contrast to the RT-PCR ELISA study, in the mononuclear cell population in this study and in the re-analysis of the RT-PCR ELISA samples, there was no obvious differences in the IL-10 gene expression between the rejectors and non-rejectors that could be used for rejection monitoring purposes.

The findings of this study and those examining the role of the IL-10 promoter polymorphism in acute rejection suggest that this cytokine may not have a central role in the processes involved with the initiation and/or regulation of the acute rejection process. Its expression appears to be influenced by steroid administration, but this may be a surrogate marker for increasing levels of inflammation, and it may reflect the prolonged need for the anti-inflammatory influences of this cytokine. However, because of the less marked differences in expression patterns between rejectors and non-rejectors, IL-10 gene expression in peripheral blood mononuclear cells is a poor candidate for immunological monitoring.

6.3 Transforming Growth Factor- β 1

The TGF- β 1 gene is located on chromosome 19. It encodes for a multifunctional cytokine with anti-inflammatory, immunosuppressive and profibrogenic properties. TGF- β 1 is produced by many cell lines and is a prominent member of the cytokine cascade involved in tissue repair and wound healing (Suthanthiran 1998, O’Kane 1997). It is becoming well recognised that this cytokine is responsible, at least in part, for fibrotic changes identified in kidneys, and other organs, associated with chronic allograft nephropathy (Jain 2000, Hutchinson 1999). There is now evidence that it may also be involved in the acute rejection process. This evidence comes from studies in both animal models and in humans. Elevated levels of TGF- β 1 message and receptor have been detected in transplanted kidneys experiencing acute cell-mediated rejection (Cohen 1998). There is also evidence that the TGF- β 1 high producer genotype is associated with multiple rejection episodes and steroid resistant rejection in kidney transplant recipients (Hutchinson 1999). The level of TGF- β 1 gene expression in peripheral blood mononuclear cells before and after renal transplantation was not investigated in the RT-PCR ELISA study. With the increasing evidence that it might be involved in the rejection process it was felt that this cytokine warranted further investigation, to determine whether sequential monitoring could be used for immunological monitoring in renal transplant recipients.

As for IL-4 and IL-10, TGF- β 1 gene expression in this study was measured in the mononuclear cells obtained from the peripheral blood of both transplant recipients and living donors undergoing native nephrectomies, both before and up to 14 days (recipients) or 7 days (donors) after surgery.

The results of the measurement of TGF- β 1 gene expression in the living donors were shown graphically in figures 5.2i and the results of the statistical analysis summarised in tables 5.2a and b. Like IL-4, TGF- β 1 gene expression was not significantly altered following surgery in the living donors. There were no significant changes detected in TGF- β 1 gene expression in the living donors between the pre-operative baseline value and either any of the post-operative values, or sequential day to day values. This means that any significant findings identified in the TGF- β 1 gene expression of the transplant recipients in this study were also the result of immunosuppression therapy or immunological events.

In the renal transplant recipients the pattern of the results was different. The results of the measurement of TGF- β 1 gene expression in the non-rejecting recipients were shown graphically in figures 5.2j and the results of the statistical analysis summarised in tables 5.2c and d. In non-rejecting recipients there were significant changes in TGF- β 1 gene expression following transplantation. There was an approximate 3-cycle increase in expression from the pre-transplant baseline value to the value on the first post-operative day. This significant increase in expression was maintained for the majority of the 14 days monitored following transplantation. However, there was a

statistically significant decrease in TGF- β 1 gene expression between the sixth and seventh post-operative days, but on both days the expression remained significantly higher than the pre-transplant baseline value.

The results of the measurement of TGF- β 1 gene expression in the rejecting renal transplant recipients were shown graphically in figures 5.2k and l. The results of the statistical analysis were summarised in tables 5.2e, f, g and h. In the group of recipients experiencing an episode of acute rejection, there were both differences and similarities compared with the non-rejectors. There was no significant initial rise in TGF- β 1 gene expression following transplantation, as seen in the non-rejectors. There were also no significant day-to-day changes in expression throughout the study period. However, there were several days, starting with the third post-operative day, on which the TGF- β 1 gene expression was significantly higher than the pre-transplant baseline value.

When the data were re-aligned around the day of rejection diagnosis and initiation of anti-rejection treatment, a distinct pattern became apparent. There was significantly elevated TGF- β 1 gene expression, compared to the pre-transplant baseline value, for the four days immediately prior to the initiation of anti-rejection medication and for the first day following treatment. There was also a significant increase in expression from the third to the second day preceding the confirmation of the diagnosis. This was followed by a significant fall in expression between the second day before diagnosis and the day the diagnosis was made. There was a further significant fall in TGF- β 1

gene expression between the first and second post-treatment days, which reduced expression levels to pre-transplant values.

The changes detected in TGF- β 1 gene expression following renal transplantation were probably the result of immunological and/or immunosuppressive mechanisms. The increase in TGF- β 1 gene expression in the non-rejecting recipients was unlikely to be the result of increased activation secondary to surgical tissue damage and/or wound healing resulting from the operation itself, as surgery did not affect expression in the living donors. The initial increase in expression seen in the transplant recipients may be the result of monocyte activation following the re-perfusion injury and subsequent tissue damage ensuing within the transplanted kidney. If this were the case then one would expect a similar rise to be detected in the rejectors. The fact that this rise is not seen suggested that the processes leading to TGF- β 1 gene expression in the rejectors and non-rejectors might be different. As TGF- β 1 has immunosuppressive properties it might be that the increased gene expression in the non-rejectors was the result of cell activation following donor recipient cell interaction leading to tolerance. Another possible explanation for the increase in TGF- β 1 gene expression following renal transplantation was the direct effect of the immunosuppressive treatment. There is work that proposes that Cyclosporin (the main immunosuppressive used in our series) leads to an increase in TGF- β 1 expression (Khanna 1994, Hojo 1999).

When the data in the rejector group were re-aligned there was an increase in gene expression (compared to pre-transplant baseline values) prior to the diagnosis of acute rejection. This rise in gene expression may reflect increased activation due to the further damage occurring within the graft as a result of the rejection process. It may also reflect increased stimulation resulting directly from the T cell activation responsible for the ongoing rejection process. The significant fall in TGF- β 1 gene expression in peripheral blood mononuclear cells detected prior to the initiation of anti-rejection treatment may be the result of cell sequestration in the damaged graft. The continuing fall after initiation of anti-rejection treatment may be the result of continued sequestration of peripheral blood mononuclear cells into the graft or the immunosuppressive effects of methyl prednisolone.

It is well recognised that TGF- β 1 is an important component of the process that results in tissue repair and wound healing (O'Kane 1997, Martin 1997, Grande 1997). However, the expression levels of this gene in peripheral blood mononuclear cells do not appear to be altered by surgery, as there were no significant changes in expression of this gene in the living donor group in this study. Local levels at the site of injury may well increase, but this was not reflected in the expression levels in the peripheral blood mononuclear cells in our cohort of donors.

Other groups have studied TGF- β 1 gene expression in core biopsies from renal transplants. As with other cytokines, the results are contradictory. Some have reported elevated levels of message at the time of acute rejection in both

animal models and clinical renal transplantation settings (Cohen 1998). Some have reported elevated TGF- β 1 gene expression in core biopsies from human renal transplants at the time of acute rejection episodes compared with biopsies from non-rejecting kidneys (Loong 1996). Others have not been able to correlate the presence and extent of TGF- β 1 message with acute rejection in kidney transplant recipients, because it has been detected in both rejectors and non-rejectors (Xu 1995, Sharma 1996). The inconsistency of these results, as with many from intragraft biopsy studies, may reflect the limitations of single time point measurement of what is a highly dynamic and changeable system. It may also reflect different roles for TGF- β 1 depending upon the balance of rejection versus tolerance within the graft.

One group has studied the gene expression of IL-2, IL-4, IL-6, IFN- γ and TGF- β 1 in peripheral blood lymphocytes and core biopsies from liver transplant recipients. Unfortunately the sampling was again at single time points. However, the peripheral blood lymphocyte samples were taken at the same time as the liver biopsies were performed. Using semi-quantitative RT-PCR technology, this group was unable to detect a significant difference in TGF- β 1 message expression between the rejectors and the non-rejectors for both liver biopsies and peripheral blood lymphocytes. However, they were able to report a good correlation between intragraft and peripheral blood lymphocyte cytokine profiles, suggesting that 'lymphocyte populations regulating graft rejection were predominant both locally (intragraft) and peripherally (peripheral blood lymphocytes)' (Gorczynski 1996). This latter comment, at first glance, goes against the hypothesis of cell sequestration

accounting for some of the changes in TGF- β 1 gene expression that were detected. However, as the changes detected were between values that were both increased compared with baseline values, the hypothesis remains possible. The findings of Gorczynski's study are supported by this study. Significantly increased TGF- β 1 gene expression levels were detected in both non-rejectors and rejectors.

Little work has been published on the potential influence of the TGF- β 1 signal sequence polymorphism at position +915 on acute rejection. Work in our own laboratory has suggested that donor low expresser genotypes may be protective against acute rejection episodes, but this remains speculative (Poole 2001). These findings are supported by the observation that renal transplant recipients who are genetically high TNF- α expressers and genetically high TGF- β 1 expressers have a higher relative risk of multiple rejection episodes than patients who are genetically high TNF- α expressers and genetically low TGF- β 1 expressers (Hutchinson 1999). However, the Oxford group have reported that this polymorphism does not appear to influence either the incidence or severity of acute rejection episodes in renal transplant recipients (Marshall/McLaren 2000).

The results from our study have identified elevated TGF- β 1 gene expression levels, when compared to pre-transplant values, in both non-rejectors and rejectors. Therefore, it is probably not surprising that the results from polymorphism studies have not been able to link episodes of acute rejection with genotype. This is not the case for chronic rejection. In this situation there

is a strong link between high expresser genotype and chronic renal transplant nephropathy (Hutchinson/Pravica/Hajeer 1999).

In this study we have detected significant increases in expression of the TGF- β 1 gene in peripheral blood mononuclear cells following renal transplantation. Compared with pre-transplant baseline values these increases were seen in both rejectors and non-rejectors. These changes cannot be explained simply by the surgical insult as no change in expression was detected in the living donors following native nephrectomy. In the rejectors, a significant increase in expression was seen prior to the diagnosis of rejection, similar to that seen for IL-4 gene expression. However, prior to the commencement of anti-rejection therapy there was a significant decrease in expression as was seen for IL-10. This decrease in expression levels was more marked than the decrease seen for IL-10 and also expression levels continued to fall following the start of treatment.

The increased production of TGF- β 1 message in all recipients may be the result of activation of peripheral cells by cell damage, secondary to the re-perfusion injury, and immunological activation following transplantation. This may be why some studies have detected TGF- β 1 message in core biopsies from stable and rejecting kidneys. If such studies had used a fully quantitative assay, then it might have been possible to detect higher levels in rejecting grafts compared with non-rejecting ones. The significant rise in TGF- β 1 gene expression seen in the rejectors prior to the clinical diagnosis may reflect further cell activation. The fall in expression that precedes anti-rejection

treatment is possibly the result of subsequent recruitment of activated cells into a transplanted organ experiencing significant cellular damage and increased immunological activation.

The results obtained from the measurement of TGF- β 1 gene expression in this study are very similar to the results from the measurement of the gene expression for the three house keeping genes described in chapter 4. It is likely that the processes that led to the increased TGF- β 1 gene expression result in cell activation and proliferation. This would probably result in increased expression of β 2-microglobulin, β -actin and transferrin receptor genes (see chapter 4). This may explain the parallel changes that were detected.

In the patients experiencing a rejection episode there was no initial rise in gene expression following transplantation, as seen in the non-rejecting recipients. Also there was a significant rise followed by a significant reduction in TGF- β 1 gene expression prior to the clinical (seen in the realigned data), and subsequently biopsy proven, diagnosis of rejection, but this change occurred on a background of elevated levels. However, following anti-rejection treatment, these levels returned to pre-transplant values. These observations are novel and may be of potential benefit in monitoring renal transplant recipients in the future. However, the elevated levels detected in the non-rejectors would potentially make data interpretation difficult. Also the most discriminating change detected around the time of rejection was that of

the expression levels in the rejectors returning to pre-transplant levels, but this occurred after the diagnosis of acute rejection is made.

6.4 Tumour Necrosis Factor- α

The TNF- α gene is located on chromosome 6 with the MHC. The gene product is a pro-inflammatory cytokine with multiple actions, produced by both lymphocytes and macrophages. Its pro-inflammatory functions suggest that it may play an important role in initiating and orchestrating the rejection response (Azzawi 1999). Other evidence, from gene polymorphism studies, also supports a prominent role for TNF- α in acute transplant rejection. The TNF- α -308 high expresser promoter polymorphism is associated with increased frequency and severity of rejection in heart, liver and kidney transplantation (Turner/Grant 1997, Hutchinson/Pravica/Perrey 1999, Bathgate 2000, Sankaran 1999).

The level of TNF- α gene expression in peripheral blood mononuclear cells before and after renal transplantation was not investigated in the RT-PCR ELISA study. It was chosen in this study because there have been many studies, both renal and non-renal, that had implicated this cytokine as a important component of the rejection process. As for the other three cytokines, TNF- α expression in this study was measured in the mononuclear cells obtained from the peripheral blood of both transplant recipients and living donors undergoing native nephrectomies.

The results of the measurement of TNF- α gene expression in the living donors were shown graphically in figure 5.2m. The results of the statistical analysis

were summarised in tables 5.2a and b. As for IL-4 and TGF- β 1, TNF- α gene expression was not significantly altered by surgery in the living donors. There were no significant changes detected in TNF- α gene expression in the living donors between the pre-operative baseline value and any of the post-operative values. There were also no significant changes detected in TNF- α gene expression in the living donors between sequential day to day values. Therefore, any significant findings identified in the TNF- α gene expression of the transplant recipients in this study were likely to be the result of immunosuppression therapy or immunological events.

The results of the measurement of TNF- α gene expression in the non-rejecting renal transplant recipients were shown graphically in figure 5.2n. The results of the statistical analysis are summarised in tables 5.2c and d. In the non-rejecting transplant recipients the results of the TNF- α gene expression measurement were different from the other three cytokines. As for the donors, there were no significant changes detected in the expression levels of the TNF- α gene. This was both when comparing the result from each day with the baseline value, and when comparing the day to day variation. This was a very important observation as IL-4, IL-10 and TGF- β 1 gene expression had varied significantly following transplantation in the non-rejecting recipients. Therefore, any significant changes in expression levels from the recipients experiencing a rejection episode were most likely to be the result of the rejection process and subsequent anti-rejection treatment. Potentially, this would make TNF- α gene expression in peripheral blood lymphocytes of renal transplant recipients a very useful parameter for immunological monitoring.

The results of the measurement of TNF- α gene expression in the rejecting renal transplant recipients were shown graphically in figures 5.2o and p and the results of the statistical analysis summarised in tables 5.2g and h. In the rejectors there were, indeed, significant changes in expression detected. By the third post-operative day, TNF- α gene expression had increased by 4-cycles compared to the pre-transplant baseline value. After some significant day-to-day changes in expression between the third and sixth post-operative days, which were difficult to interpret without re-aligning the data, TNF- α gene expression increased further. On the eighth, ninth, twelfth and fourteenth post-transplant days TNF- α gene expression had increased to a level 5-cycles higher than the pre-transplant baseline value. This equated to an approximate 32-fold increase in gene expression. These large increases in expression were probably related to the rejection process. To aid interpretation, the data were re-aligned to the day of rejection diagnosis, as was done for the other three cytokines.

When the data from the rejectors were re-aligned to the day of diagnosis of rejection the pattern of expression was similar to that seen for TGF- β 1. When comparing each day with the pre-transplant baseline value there was a significant increase in TNF- α gene expression for the 4 days immediately prior to the day of rejection diagnosis. In the stepwise analysis comparing the day-to-day changes in expression there was a significant increase TNF- α gene expression between days 3 and 2 before rejection diagnosis followed by a significant decrease in expression between days 2 and 1 before rejection diagnosis. There was another fall in TNF- α gene expression between the day

before and the day after rejection diagnosis (commencement of anti-rejection therapy). This fall in expression was not significant in itself, but it did result in the expression levels for that, and subsequent days, returning to pre-transplant baseline values.

The increase in TNF- α gene expression detected in the rejectors in the re-aligned data most probably reflects increased cell activation due to the rejection process. From this study it was not possible to conclude whether this rise initiated the resulting rejection process or was part of it. The significant fall in TNF- α gene expression in peripheral blood mononuclear cells detected prior to the initiation of anti-rejection treatment might be the result of recruitment and sequestration of activated cells into the graft. The continuing fall after initiation of anti-rejection treatment may result from either the continued sequestration of peripheral blood mononuclear cells into the graft or the immunosuppressive effects of methyl prednisolone.

In our study we were unable to detect any changes in TNF- α levels following nephrectomy in the living donors. Work performed in non-transplant centres has also reported a lack of response of TNF- α levels to surgery. Serum levels of TNF- α did not increase in 21 patients following major intestinal surgery (Sasada 1999). In another series of 12 patients undergoing upper gastrointestinal resections, TNF- α fell following surgery, but had reached steady state before 18 hours post-operatively (van Berge Henegouwen 1998). However, not all studies have failed to detect a rise in TNF- α following surgery. The same group that reported no change in IL-10 levels following

cholecystectomy did detect a rise in serum TNF- α levels at 24 hours with a return to normal levels by 7 days (Bellon 1997).

Intragraft TNF- α gene expression levels have been studied on several occasions in most forms of solid organ transplantation. Again, most have been single time point studies, but do suggest that TNF- α is an important cytokine in the process that results in acute allograft rejection. Loong and colleagues detected enhanced TNF- α mRNA in biopsy specimens from kidneys with acute rejection compared with biopsies from kidneys with chronic rejection or no rejection (Loong 1996). Joseph et al were only able to detect TNF- α gene expression in kidneys with acute rejection, but not all rejecting kidneys expressed TNF- α (Joseph 1995). Krams and co-workers were able to detect TNF- α message in rejecting and rejected renal transplant biopsies, but they did not compare their results with samples from non-rejecting, stable transplants (Krams 1992). In contrast, Vandembroecke et al were only able to detect TNF- α mRNA in 1/8 biopsies from acutely rejecting kidney transplants and none of the stable renal grafts (Vandembroecke 1991).

Many groups have measured serum TNF- α levels in transplant recipients, again in most forms of solid organ transplantation. Maury and Teppo were some of the first to study sequential serum TNF- α in 10 renal transplant recipients and 40 healthy controls. They reported elevated levels associated with febrile rejection episodes, but not all rejection episodes (Maury 1987). Noronha and colleagues reported similar results identifying elevated serum levels of TNF- α associated with acute renal allograft rejection. However, 3

patients in their series, who rejected, had undetectable levels. (Noronha 1990). McLaughlin's results were also equivocal, only identifying elevated levels in 65% of renal allograft rejectors (McLaughlin 1991). One study has reported results from studying both serum TNF- α and serum TNF- β in renal recipients. Both were noted to rise prior to rejection and infection, but there was a significantly larger mean increase of TNF- α in those that went on to reject compared with those that had an infection (Daniel 1995). Some, however, have not detected any association between TNF- α and rejection in renal transplant recipients (Tice 1993).

Studies of TNF- α in liver transplant recipients have revealed similar findings in that elevated TNF- α levels are associated with acute rejection. Vogel and colleagues collected sequential serum samples from 20 liver transplant recipients. They detected elevated levels of TNF- α in the samples of those patients experiencing acute rejection episodes. They also reported that the peak level measured preceded the clinical rejection episode (Vogel 1990).

Several groups have measured serum TNF- α levels following cardiac transplantation. None have reported that the results obtained were of any benefit as a predictor of acute rejection episodes in this group of patients (Grant 1996, Azzawi 1996, Torre-Amione 1998).

Rondeau et al took a different approach and measured TNF- α production by different cell subsets following heart-lung transplants. Peripheral blood mononuclear cells and cells obtained from bronchioloalveolar lavage were

studied. They reported that rejection episodes were not associated with significant changes in the ability of either cell type to synthesise TNF- α . As with many of these studies, the cells were collected at single time points, at the time of rejection. The rejection episode in all patients was also beyond three months following transplantation (Rondeau 1990).

The results obtained from the measurement of TNF- α gene expression are the most clinically significant of the four cytokines examined in this study. This was the only cytokine in which the expression levels did not vary significantly in the non-rejectors. The statistically significant rise in expression detected in the rejectors was most likely to be the result of the rejection process, although we have not proven this. These results agree with the majority of the work, previously reported, implicating TNF- α as a very important cytokine in the process of acute allograft rejection. This study is not able to elucidate the position or role that TNF- α has in the initiation of acute rejection, but can only say that the gene expression of this pro-inflammatory cytokine, in peripheral blood mononuclear cells, is strongly influenced by the mechanisms involved with the acute rejection process. At least this is suggested by the strong association between the expression change and clinical and intragraft events.

These results of TNF- α gene expression measurement in this study complement other work published examining the potential influence of the –308 (high expresser) promoter polymorphism on transplant outcome. There is strong evidence that the high in vitro expresser allele of the –308 promoter

polymorphism is associated with increased rejection episodes and steroid resistant rejection in cardiac, liver and renal transplantation (Turner/Grant 1997, Hutchinson/Pravica/Perrey 1999, Bathgate 2000, Sankaran 1999).

In this study we have been able to detect significant differences in TNF- α gene expression in renal transplant recipients experiencing an acute rejection episodes compared with both the donors and, more significantly, the non-rejecting recipients. This observation, in combination with the significant rise and subsequent fall in expression levels prior to the clinical diagnosis, makes the measurement of TNF- α gene expression in the peripheral blood mononuclear cells of renal transplant recipients a very promising parameter for potential immunological monitoring.

Chapter 7: Summary and Conclusions

This study sought to answer some of the questions created by work previously performed in our laboratory and to expand our knowledge and understanding of cytokine gene expression in the early post-operative period following renal transplantation. The previous work had identified differences in the gene expression pattern of several cytokines when comparing rejectors and non-rejectors. This present study aimed to confirm the findings of the semi-quantitative methodology used previously by re-analysing the same cDNA samples using fully quantitative TaqMan PCR technology. The study then set out to investigate other areas of interest by recruiting a new cohort of 45 renal transplant recipients and 13 living renal donors. The methodology was simplified by using peripheral blood mononuclear cells rather than T lymphocytes, rendered quantitative by TaqMan technology and the sampling frequency increased to daily. It was then possible, using these new cohorts of patients and their cDNA samples, to identify whether the use of a house keeping gene would improve the quality and reliability of the results by controlling for fluctuations in the quantity of cDNA analysed. It was also possible to investigate the influence of surgery on cytokine gene expression by comparing the gene expression patterns of the same cytokines in patients undergoing a similar procedure to a renal transplant, but without the immunosuppression, i.e. the living donors, to those of the transplant recipients. The main aim, however, was to investigate whether intensive monitoring of cytokine gene expression in renal transplant recipients, in the

first two weeks following surgery, had the potential to be used as a method of immunological monitoring. Finally, results from this study have generated potential new avenues of enquiry and investigation. These points will be discussed in depth below.

Peripheral blood samples were collected pre-operatively and then daily for up to 14 days in 13 living kidney donors and 45 renal transplant recipients, 15 of which experienced an episode of biopsy proven acute allograft rejection. All patients were transplanted and managed in the same unit with well-established treatment protocols. Peripheral blood mononuclear cells were extracted from the samples on the day of collection. Total RNA was extracted from these cells and quantitated before being reverse transcribed into cDNA. This enabled the samples to be stored. The gene expression of three potential house keeping genes (β -actin, β 2-microglobulin and transferrin receptor) and four cytokine genes (IL-4, IL-10, TGF- β 1 and TNF- α) were measured using fully quantitative real time TaqMan PCR technology. The results were presented in chapters 4 and 5 and discussed in chapter 6.

A potential weakness of the cytokine expression monitoring performed in our laboratory prior to this study was that the measurement of the gene expression was performed using a semi-quantitative technique, namely RT-PCR ELISA, for end point PCR product detection. In the experiments described in chapter 3, using the same cDNA samples, it was possible to repeat the analysis for the cytokines IL-4 and IL-10, but this time using fully quantitative exponential phase PCR TaqMan technology. For IL-4, the two sets of data showed very

close correlation, although it was not possible to directly compare the results statistically. There remained distinct patterns for rejectors and non-rejectors, which corresponded very well with intragraft events, confirming the results of the earlier work. However, the results from the re-analysis of the samples for IL-10 gene expression were unable to confirm the earlier findings. There were still significant changes in IL-10 gene expression following renal transplantation, but there was no obvious difference detected in expression patterns between rejectors and non-rejectors in the first two weeks following surgery.

These results were encouraging. They confirmed that the significant changes in gene expression of these two cytokines following renal transplantation were likely to be the result of immunological processes and not methodological artefact. The confirmation that IL-4 gene expression patterns appeared to vary with rejection status and the confirmation of alterations in IL-10 gene expression following transplantation justified further investigation of peripheral cytokine mRNA monitoring following allotransplantation.

In this study, and in the RT-PCR ELISA study, great care was taken to quantify the amount of RNA extracted in order to standardise the concentrations of the RNA solutions prior to reverse transcription. The object of this was to reduce the potential error that apparent variations in cytokine gene expression were in fact the result of variations in substrate amount. However, this step could not absolutely guarantee standard concentrations of cDNA in PCR reactions for two reasons. Firstly, the RNA extracted and

measured was total RNA, while it is only mRNA that is reverse transcribed. Methods exist for extracting mRNA alone. However, these techniques had not been used in our laboratory hitherto and insufficient time was available in this project to compare results obtainable from total RNA and mRNA extraction methods. The methodology applied in this study also afforded comparability with our previous RT-PCR ELISA study and has been used in many other gene expression studies of this sort. Secondly, it was not possible to calculate the efficiency of the reverse transcription process. One way to overcome this problem was to use a house keeping gene or genes. The expression of this gene(s), if constant, would allow for standardisation of the TaqMan results of the cytokine genes under investigation, therefore controlling for any potential fluctuations in cDNA concentration.

For a house keeping gene to be effective it must be expressed at a constant level and not be influenced by the biological processes under investigation. In this study these were surgery and renal transplantation. Before choosing a definitive house keeping gene for standardisation, expression levels were measured in three different potential house keeping genes in order to determine the most appropriate for use in our study group. Three genes were chosen. These were β -actin, β 2-microglobulin and transferrin receptor. Expression levels of each of the three genes were measured in the three different groups of patients in this study.

In the living donors the results were encouraging. There were no significant variations in expression levels of any of the three genes following surgery.

This was an important finding, since it strongly suggests that the process of RNA extraction and cDNA transcription used in this study did indeed produce standard solutions of cDNA.

The results were different when expression of the three genes was measured in the transplant recipients. In both the rejectors and non-rejectors there were significant variations in gene expression. These were associated with the initiation of immunosuppression, the detection of acute allograft rejection and the commencement of anti-rejection treatment. These changes were not in line with some of the changes detected in IL-4 and IL-10 gene expression, but similar to the patterns detected in the other two cytokines studied, particularly in the rejector cohort of transplant recipients.

The conclusions to be drawn from this part of the study were that none of the three genes studied were good candidates for house keeping genes, and, therefore, great care must be taken when choosing a suitable house keeping gene for standardisation. This is particularly true in the field of transplantation. These observations were supported by work performed by other groups that have identified inconsistent expression levels in well-recognised 'house keeping' genes in non-transplant settings.

So is it necessary to use a house keeping gene for the purpose of standardisation in this setting? The results of the gene expression analysis from the living donor data suggest that our methodology for RNA standardisation and cDNA transcription was indeed reliable. As the samples

and analysis from this cohort of patients were managed synchronously and in an identical manner to those from the transplant recipients, the implication is that the changes in cytokine gene expression detected are the result of the transplantation process. Therefore, it could be argued that standardisation is unnecessary. However, when using quantitative techniques, it would be ideal to be able to reinforce findings with the data from a reliable house keeping gene or genes. This would require extensive testing to ensure that the processes under investigation in the cell type or types being studied do not influence the chosen gene(s).

It was an important objective of this study to separate the influences of surgery on cytokine gene expression from those of the immunosuppression and acute rejection and its subsequent treatment. This would enable us to be confident that any changes we detected in the cytokine gene expression of the transplant recipients were due to immunological processes alone. It is obviously unethical to perform transplantation without immunosuppression. Use of a cohort of patients undergoing living donor nephrectomy as a surgical control was, therefore, a logical substitute.

Cytokine gene expression was measured pre-operatively and daily for the first week post-operatively. Expression levels of the cytokines IL-4, TGF- β 1 and TNF- α were not influenced by surgery. There were no significant variations in expression of any of the three cytokines, either when comparing pre-operative values to post-operative values, or when comparing day to day values. These findings agreed with the vast majority of published data.

Expression of the cytokine IL-10, however, did appear to be influenced by surgery. Following the donor nephrectomy there was a significant rise in the expression levels of this cytokine. This increased expression was detected for the first two days following surgery. Levels were back to pre-operative levels by the third post-operative day. This finding confirmed work performed by other groups.

These findings meant that care must be taken when interpreting the IL-10 expression data in the transplant recipients, but any significant changes in expression levels detected in the other three cytokines were most likely to the result of processes related to the transplant itself.

When comparing changes in cytokine gene expression in rejectors and non-rejectors, the expression pattern for IL-4 was different and correlated well with intragraft events. In the non-rejectors, following transplantation, there was a large decrease in expression levels that lasted for several days before the levels returned to pre-operative values. These changes are probably the result of the transplant and not surgery as no change in expression was detected in the control group. The changes may be due to either the commencement of immunosuppression immediately prior to surgery or possibly the recruitment of IL-4 expressing cells from the peripheral blood and into the graft. In the rejectors the pattern of expression was very different. The initial decrease in gene expression seen in the non-rejectors was only detected for a day before levels returned to baseline values on the second post-operative day. There was then a significant increase in expression prior to the

diagnosis of acute rejection followed by a significant fall in expression after commencement of anti-rejection therapy. Levels fell below pre-transplant levels following anti-rejection treatment before returning to baseline levels after cessation of the additional immunosuppression.

Explanation of these changes is speculative. The rise in IL-4 gene expression prior to the detection of acute rejection may be causative in the rejection process or result from it. The fall in expression detected immediately following transplantation in the non-rejectors and following anti-rejection therapy in the rejectors may be due to either simple decreased expression following treatment with methyl prednisolone or the sequestration of the more active cells from the peripheral blood into the graft. Either way, this cytokine may be of value in immunomonitoring renal transplant recipients and has the potential to predict the onset of acute allograft rejection prior to clinical and biochemical signs become apparent.

The pattern of IL-10 gene expression detected in rejectors was very similar to the pattern detected in the non-rejectors. This was in stark contrast with the IL-4 data. For IL-10, both groups showed a similar initial rise in expression following the transplant operation, as was seen following living donor nephrectomy. This rise in expression was maintained for most of the first post-operative week in the non-rejectors and almost the entire study period in the rejectors. There was a significant fall in IL-10 expression immediately prior to the diagnosis of acute rejection, but this was on a background of

increased expression levels both prior to and following the commencement of anti-rejection therapy.

It would appear that surgery and steroids have more influence on the expression of the IL-10 gene than any rejection process. Other groups working in non-transplant settings indirectly support these results. These findings, together with the poor discrimination between the rejector and non-rejector expression patterns make this a poor candidate for immunomonitoring.

There were significant changes in expression of TGF- β 1 in relation to transplantation and at the time of acute rejection. In the non-rejectors there was an initial rise in expression levels following transplantation when compared to pre-transplant values. These elevated levels were maintained for most of the study period, with a sustained period of elevation between the sixth and tenth post-operative days. Although there was no initial statistically significant rise in TGF- β 1 seen in the rejectors, there were elevated levels detected on day 3 and several subsequent days. When the data were re-aligned there was a more discernible pattern. Expression levels were elevated for the five days prior and one day after the rejection was diagnosed when compared to pre-transplant values. When examining the day to day variation there was a significant rise followed by a significant decrease in expression levels prior to the detection of acute rejection. A further significant fall following anti-rejection treatment returned levels to pre-transplant values.

Changes in TGF- β 1 gene expression, therefore, correlated well with intragraft events, but the elevated levels detected in both rejectors and non-rejectors compromise its role in immunomonitoring. The significance of the lack of initial rise in TGF- β 1 gene expression seen in the rejectors is as yet undetermined, but may be a very important observation.

Results obtained for TNF- α were the most interesting with regard to its immunomonitoring potential. There were no significant changes in expression levels of this cytokine in the non-rejecting cohort when comparing day to day values and post-transplant values to the pre-transplant baseline value. Therefore changes detected in expression of this gene in the rejector group must be related to the rejection process. When the data were re-aligned there were significant elevations in expression levels for the four days prior to the diagnosis of rejection. When examining the day to day variation, there was a significant rise followed by a significant fall in gene expression prior to the diagnosis of acute rejection. These observations make TNF- α a very good candidate for immunomonitoring.

Study Conclusions

Using a simpler and quicker cell extraction technique and a fully quantitative method of measuring cytokine gene expression this study has confirmed that changes in cytokine gene expression of peripheral blood mononuclear cells correlate well with intragraft events in renal transplant recipients in the early post-transplant period. It has also been able to show that most of the changes

in cytokine gene expression detected are not related to the surgery involved with the renal transplant as cytokine gene expression has also been studied in a group of living donors. The study has also demonstrated that great care must be taken when choosing a house keeping gene for the purposes of standardisation. Both immunosuppression and acute rejection influenced gene expression of the three genes selected as potential house keeping genes in this project. Interestingly, this means they may have a potential role in immunomonitoring, an observation first made many years ago. Finally, this study has demonstrated that immunomonitoring of renal transplant patients has potential in the clinical setting. The changes in expression of both IL-4 and TNF- α differ significantly between rejectors and non-rejectors, and that these changes are detectable prior to the clinical signs of rejection.

The patients in this study received a number of different immunosuppressive drugs in several different combinations. As the two transplant cohorts were not on the same treatment regimes either within the groups or across the groups, this adds additional importance to these observed variations in expression patterns detected. The variations in expression are significant enough to have been detected despite the variability of the treatment regimes.

This study has confirmed the work previously performed by our group identifying the potential use of cytokine gene expression in PBMCs as a method of immunological monitoring of renal transplant recipients. The method and results described in this study have several benefits over the majority of work previously published in this field. Unlike most studies that have simply compared rejectors with non-rejectors we have also been able to

use the patients as their own controls. This was possible because of the intensive sequential monitoring undertaken. This enabled us to study cytokine expression patterns in rejectors and non-rejectors, therefore, enabling us to study changes in expression levels as well as absolute values. This study has been able to detect changes in the level of cytokine gene expression in relation to clinical events and not just whether the cytokine is detectable or not. The comparison of absolute values in single time point studies comparing rejectors with non-rejectors may have contributed to some of the confusion in the literature.

Another advantage of our method is that potentially it is fully quantitative and highly reproducible. This allows comparisons that are not possible with other non-quantitative methods such as PCR-ELISA. Our method also has the advantage that it is not dependent upon renal function, as the volume of urine output does not affect the measurement of gene expression in PBMCs. Monitoring cytokine levels in urine - by whatever technique - might be compromised by this. This method does not require prolonged manipulation of the extracted cells in vitro, reducing the potential of cell activation by other routes than those being studied.

This study has not been able to draw any conclusions about the effect of either infection or CyA toxicity on cytokine gene expression. This was because there was no episodes of infective complications within the first two weeks in any of the recipients and so few patients had biopsy proven CyA nephrotoxicity. This was unfortunate, as other studies have shown that both these

complications can affect the results of immunological monitored parameters. The effect of CyA toxicity on cytokine gene expression needs to be examined since this may enhance our understanding of the relative importance of the role each cytokine plays in acute rejection. Hopefully, as our group continues to collect samples it will be possible to explore this avenue at a later date. In the case of infection this is less relevant in the first two weeks following renal transplantation. CMV disease is very unlikely to be the cause of renal dysfunction this soon after transplantation and bacterial infection of the urine is relatively simple to diagnose with modern dipstick tests. Differentiating between CMV disease and acute rejection as causes of graft dysfunction is more important in settings where acute rejection episodes are not concentrated in the first few weeks following surgery, such as cardiac transplantation.

The method of cytokine gene expression measurement used in this study is relatively simple to perform with the appropriate equipment and a degree of scientific training. It is possible to obtain a cytokine gene expression result the same day as the blood is taken. This is of great potential benefit as changes in gene expression appear to occur prior to changes detected either clinically or biochemically. This would allow the clinician to perform a core biopsy early and potentially initiate anti-rejection treatment prior immunologically mediated damage occurring within the transplanted organ. Also, high throughput TaqMan technology is now available, permitting 384 samples, as opposed to 96 samples, to be analysed in a single run. This improved methodology is also more rapid and can be performed using smaller reaction

volumes, reducing reagent costs and conserving cDNA samples for further studies.

This study has identified that it is possible to detect significant changes in cytokine gene expression prior to clinical and biochemical signs of acute allograft reaction and also following the commencement of anti-rejection therapy. Positive and negative predictive rates for each cytokine were not calculated, as the number in the rejection group was small. However, it is unlikely that the measurement of the gene expression of a single cytokine would be enough to predict acute rejection or to monitor the success of anti-rejection treatment. It is therefore important that as many potential molecules involved or potentially involved in acute rejection are tested. This does not need to be limited to cytokines. Ongoing work in our laboratory is studying the expression of more genes including additional cytokines, adhesion molecules and markers of cellular activation. As the method in this latest study is identical to the one in this study, a large cDNA sample bank is being built up. This will allow a larger number of samples to be tested, therefore, improving the quality of the results still further. Advances in quantitative PCR technology, involving more rapid, high throughput PCR thermal cycling will facilitate this.

The findings of this study suggest that prior to the clinical detection of acute rejection, there is sequestration of 'active' cells into the transplanted kidney. The ultimate aim of the approach developed in this study should be simultaneous peripheral blood sampling and renal transplant fine needle

aspiration to test this hypothesis. This would allow an excellent understanding of the relationship between cytokine gene expression measured in peripheral blood mononuclear cells expression of mononuclear cells residing within the transplanted kidney both at the time of active rejection, but also at times of tolerance.

The technique developed in this study may also have other applications. It may be of potential use in monitoring the effectiveness of drug therapies, either in the short-term or, with potentially more impact on graft survival, in the long-term. This may be of particular relevance to the association between TGF- β 1 and chronic allograft nephropathy.

This study has confirmed the findings of earlier work from our laboratory identifying the potential role of the measurement of cytokine gene expression in cells from peripheral blood of renal transplant recipients in the early post-operative period. It has also shown that there may be several potential markers of rejection activity and that changes may precede cellular damage. There is also the potential for the method to be used for long term monitoring of patients and assessing the effectiveness of new treatment modalities.

Appendices

Appendix I

Patient Information Sheet: Cytokine Study

As I am sure you are aware, acute rejection is the result of your white blood cells recognising the transplanted kidney as foreign to you. These white cells release chemicals, called cytokines, into the blood. These cytokines excite other white cells and may propagate a rejection episode. During a rejection episode there may be some damage to your transplanted kidney.

We know from previous studies that there are changes in the levels of certain cytokines during a rejection episode.

Recent work that has been carried out in Manchester and other centres, suggests that some people may produce more of these cytokines than others do. This is a phenomenon known as gene polymorphism.

The objective of this study is twofold. Firstly, to see if by measuring the level of cytokines in your blood we are able to predict and therefore prevent the rejection before it occurs, therefore reducing the risk of damage to your kidney. Secondly, to identify those people who are theoretically more likely to produce higher levels of cytokines and see if they do in real life. This may allow us to individualise the immunosuppression each patient receives.

Mr. P. Gibbs FRCS

June 1999

Appendix II

Consent Form

Hospital
Sticker

I agree to be a patient in the MD project being carried out by Mr. Paul Gibbs
at St. Mary's Hospital, Portsmouth.

I understand that this will involve me having extra blood samples taken, on
top of the routine ones, both before my renal transplant and most days during
my subsequent stay in hospital.

I have agreed to participate in the study of my own free will and have had all
my questions answered to my satisfaction.

I understand that I can withdraw at any stage without my subsequent
treatment and care being affected in any way.

Signed.....

Date.....

Witness.....

Appendix III

A Cytokine Study Patient Proforma

Name	DOB	Hosp.No.
DOT	Tx No.	Original Disease
Recipient Tissue Type	Donor Tissue Type	Mismatch
HD / CAPD / PD	LRTx / CRTx	C.I.T.
Donor Age	Donor Sex	Cause of Death
Immunosuppression	1° Graft Function	Y / N
Biopsy Performed Y / N	Date of Biopsy	
Biopsy Result		
Anti-rejection Treatment Commenced On		
Type of Treatment & Duration		
Discharge Date		
Additional Infomation		

Appendix IV

Cell Separation

Equipment: MSE Mistral 3000i centrifuge [Sanyo, UK]
Microflow pathfinder laminar flow cabinet [MDH, UK]

Consumables: Lymphoprep, specific gravity 1.077g/l [Nycomed, UK]
Phosphate buffered saline without calcium and
magnesium (10x) [Life Technologies, UK]
Sterile 50ml skirted conical-based polypropylene tubes
[Greiner Labortechnik, UK]
Sterile 15ml conical-based polypropylene tubes
[Greiner Labortechnik, UK]
Sterile 3.5ml Pasteur pipettes [Greiner Labortechnik,
UK]
Sterile 10ml serological pipettes [Greiner Labortechnik,
UK]

Appendix V

Total RNA Extraction, Quantitation and Standardisation

Equipment:	Heraeus Sepatec Contifuge 17RS centrifuge [Heraeus, UK] Windsor incubator, set at 45°C [Sandrest, UK] Whirlimixer [Jencons Scientific, UK] Water bath, set at 55°C [Grant Instruments, UK] GeneQuant RNA/DNA Calculator [Pharmacia, UK] Ultramicrovolume cell [Pharmacia, UK] Pipettes of various ranges [Gilson, Anachem, UK; Eppendorf, Merck, UK; Biohit, Alpha Laboratories, UK] Series 2100 clinical autoclave [Prestige Medical, UK] Elgastat 4 Optional Water Purifier [Elga, UK]
Consumables:	RNAzolB [Biogenesis, UK] 0.1% Diethyl Pyrocarbonate [Sigma Chemical Company, UK] treated double-distilled water is prepared by adding 1µl to every 100ml of double-distilled water, leaving the solution to stand overnight before sterilising it in an autoclave. Chloroform [Merck, UK] Isopropanol [Merck, UK]

Absolute ethanol [James Burroughs, UK]

Sterile pipette tips of various sizes [Greiner
Labortechnik, UK]

Sterile 1.5 and 0.75ml eppendorfs [Greiner
Labortechnik, UK]

Appendix VI

First-strand cDNA Synthesis

Equipment: Amplirad UV cabinet [Gene Research Instrumentation,
UK]

Laminar airflow cabinet [MDH, UK]

Pipettes of various ranges [Gilson, Anachem, UK;
Eppendorf, Merck, UK; Biohit, Alpha
Laboratories, UK]

Hybaid thermal reactor [Hybaid, UK]

Whirlimixer [Jencons Scientific, UK]

Water bath, set at 37°C [Grant Instruments, UK]

Consumables: First-strand cDNA Synthesis [Pharmacia Biotech, UK]

DEPC treated RNase free water

Sterile 0.75ml eppendorfs [Greiner Labortechnik, UK]

Appendix VII

TaqMan Cytokine Gene Expression Analysis

As was mentioned in section 2.4.1, the first step was to ensure that the thermal cycling conditions and reaction volume were checked, on the ABI Prism 7700 instrument. This confirmed that the cycle conditions were set correctly and the reaction volume was set for 25 μ l. Selecting the ‘thermal cycler conditions’ window enabled this to be done. The final step before commencing the run was to select the ‘show set up’ window, and label all wells as ‘unknown’. Selecting ‘analysis’ and then clicking on ‘run’ started the run, which would take approximately one hour and fifty-six minutes.

When the run was complete those samples containing the FAM labelled probe (those under investigation) were left selected as ‘unknown’ in the ‘show set up’ window. Samples containing the VIC labelled probe must be labelled differently. This was done by first clicking on ‘sample type’ and selecting the positive control as VIC. Those wells that contained the VIC probe were then selected as ‘sample positive control’. Any unused wells, on either screen, were selected as ‘not in use’.

To analyse the data, the ‘analysis’ window was chosen and scrolled down to the ‘analyze’ icon, which was selected. The computer output data is graphical and a typical example is shown below (Figure VIIa).

The computer automatically selected the threshold at which the PCR reaction became exponential. Where quantitation of the PCR product, in a given reaction, reached this threshold was termed the C_T value for that reaction. This equated to the cycle number at which the reaction became exponential and was directly proportional to the amount of cDNA in the starting reaction mix.

To export the C_T value for each well into a Microsoft Excel file, 'OK' was selected on the graphics window, followed by 'File', and to 'export results', the destination was selected (3½ Floppy) followed by 'export'.

Fig. VIIa. Illustration showing a the typical appearance of a TaqMan output data graph

This graphic is a copy of a typical output data graft generated by cytokine gene expression analysis using TaqMan RT-PCR technology. The X-axis represents cycle number. The Y-axis is a log scale measuring product. Each individual plot represents an individual well.

References

Adams DH, Burnett D, Stockley RA, Hubscher SG, McMaster P, Elias E. Biliary β 2-microglobulin in liver allograft rejection. *Hepatology* 1988; 8: 1565-70.

Adams DH, Mainolfi E, Elias E, Neuberger JM, Rothlein R. Detection of circulating intercellular adhesion molecule-1 after liver transplantation – evidence of local release within the liver during graft rejection. *Transplantation* 1993; 55: 83-7.

Adams DH, Wang L, Hubscher SG, Elias E, Neuberger JM. Soluble interleukin-2 receptors in serum and bile of liver transplant recipients. *Lancet* 1989; 1(8636): 469-71.

Albert FW, Schmidt U, Seyfert UT, Würmell W, Gumbel W. Interleukin-2 measurements in peripheral blood after kidney transplantation. *Transplant Proc* 1990; 22: 1852-3.

Alderson MR, Sassenfeld HM, Widmer MB. IL-7 enhances cytolytic T lymphocyte generation and induces lymphokine activated killer cells from human peripheral blood. *J Exp Med* 1990; 172: 577-87.

Arai KI, Lee F, Miyajima A, Miyatake S, Arai N, Yokota T. Cytokines: coordinators of immune and inflammatory responses. *Annu Rev Biochem* 1990; 59: 783-836.

Asderakis A, Sanakaran D, Dyer P, Johnson RW, Pravica V, Sinnott PJ, Roberts I, Hutchinson IV. Association of polymorphisms in the human interferon-gamma and interleukin-10 genes and acute and chronic kidney transplant outcome: the cytokine effect on transplantation. *Transplantation* 2001; 71: 674-7.

Ashraf S, Parrott NR, Dyer P, Roberts I, Johnson RWG. Clinical response and temporal patterns of acute cellular rejection: relationship to chronic transplant nephropathy. *Transpl Int* 1998; 11 (suppl 1): S5-S9.

Awad MR, El-Gamel A, Hasleton P, Turner DM, Sinnott PJ, Hutchinson IV. Genotypic variation in the transforming growth factor-beta 1 gene association with TGF- β 1 production, fibrotic lung disease and graft fibrosis after lung transplantation. *Transplantation* 1998; 67: 1014-1020.

Azzawi M, Grant SCD, Hasleton PS, Yonan N, Campbell CS, Deiraniya A-K, Rahman A, Hutchinson IV. TNF α mRNA and protein in cardiac transplant biopsies: comparison with serum TNF α levels. *Cardiovas Res* 1996; 32: 551-6.

Azzawi M, Hasleton PS, Hutchinson IV. TNF-alpha in acute cardiac transplant rejection. *Cytokine Cell Mol Ther* 1999; 5: 41-9.

Baan CC, Niesters HGM, Metselaar HJ, Mol WM, Loonen EH, Zondervan PE, Tilanus HW, Ijzermans JMN, Schalm SW, Weimar W. Increased intra-graft IL-15 mRNA expression after liver transplantation. *Clin Transplant* 1998; 12: 212-8.

Baan CC, Weimar W. Intra-graft cytokine gene expression: implications for clinical transplantation. *Transpl Int* 1998; 11: 169-180.

Balkwill FR, Burke F. The cytokine network. *Immunol Today* 1989; 10: 299-304.

Bathgate AJ, Pravica V, Perrey C, Therapondos G, Plevris JN, Hayes PC, Hutchinson IV. The effect of polymorphisms in tumour necrosis factor- α , interleukin-10, and transforming growth factor- β 1 genes in acute hepatic allograft rejection. *Transplantation* 2000; 69: 1514-7.

Baud L, Fouqueray B, Suberville S, Doublier S. Interleukin-10 a logical candidate for suppressing glomerular filtration? *Exp Nephrol* 1998; 6: 22-7.

Bechtel U, Scheuer R, Landgraf R, König A, Feucht HE. Assessment of soluble adhesion molecules (sICAM-1, sVCAM-1, sELAM-1) and

complement cleavage products (sC4d, sC5b-9) in urine. Clinical monitoring of renal allograft recipients. *Transplantation* 1994; 58: 905-11.

Bellon JM, Manzano L, Bernrdos L, Ga-Honduvilla N, Larrad A, Bujan J, Alvarez-Mon M. Cytokine levels after open and laparoscopic cholecystectomy. *Eur Surg Res* 1997; 29: 27-34.

Beyers AD, Spruyt LL, Williams AF. Molecular associations between the T-lymphocyte antigen receptor complex and the surface antigens CD α , CD4, or CD8 and CD5. *Proc Natl Acad* 1992; 89: 2945-49.

Bidwell J, Keen L, Gallagher G, Kimberly R, Huizinga T, McDermott MF, Oksenberg J, McNicholl J, Pociot F, Hardt C, D'Alfonso S. Cytokine gene polymorphism in human disease: on-line databases. *Genes Immunity* 1999; 1: 3-19.

Bill O, Garlisi CG, Grove DS, Holt GE, Mastro AM. IL-2 mRNA levels and degradation rates change with mode of stimulation and phorbol ester treatment of lymphocytes. *Cytokine* 1994; 6: 102-10.

Bjorkman PJ, Parham P. Structure, function and diversity of Class I major histocompatibility complex molecules. *Annu Rev Biochem* 1990; 59: 253-88.

Bjorkman PJ, Saper MA, Samraoui B, Bennett WS, Strominger JL, Wiley DC. Structure of the human class I histocompatibility antigen HLA-A2. *Nature* 1987; 329: 506-12.

Bjorkman PJ, Saper MA, Samraoui B, Bennett WS, Strominger JL, Wiley DC. The foreign antigen binding site and T cell recognition regions of class I histocompatibility antigens. *Nature* 1987; 329: 512-8.

Borel JF, Feurer C, Gubler HU, Stähelin A. Biological effects of cyclosporin A: a new anti-lymphocytic agent. *Agents Actions* 1976; 6: 468-75.

Böyum A. Separation of leukocytes from blood and bone marrow. *Scand J Clin Invest* 1968; Suppl 97: 7.

Brando B, Sommaruga E, Civati G, Busnach G, Broggi ML, Seveso M, Minetti L. Monitoring of CD4+CD8- T-helper cells in renal transplant recipients. *Transplant Proc* 1989; 21: 1192-3.

British Transplant Society. Towards standards for organ and tissue transplantation in the United Kingdom. 1998.

Brown MH, Cantrell DA, Brattsand G, Crumpton MJ, Gullberg M. The CD2 antigen associates with the T-cell antigen receptor/CD3/antigen complex on the surface of human T lymphocytes. *Nature* 1989; 339: 551-3.

Bugeon L, Cuturi MC, Hallet MM, Paneau J, Chabannes D, Souillou JP. Peripheral tolerance of an allograft in adult rats. Characterisation by low interleukin-2 and interferon- γ mRNA levels by strong accumulation of major histocompatibility complex transcripts in the graft. *Transplantation* 1992; 54: 219-25.

Caca K, Schübel C, Krombach F, Kemkes BM, Hammer C. DNA analysis of circulating blood mononuclear cells for diagnosis of rejection in heart transplant patients. *Transplant Proc* 1989; 21: 2523-4.

Canadian Multicentre Transplant Study Group. A randomised clinical trial of Cyclosporine in cadaveric renal transplantation. *New Engl J Med* 1983; 309: 809-13.

Carter NP, Cullen PR, Thompson JF, Bewick ALT, Wood RFM, Morris PJ. Monitoring lymphocyte subpopulations in renal allograft recipients. *Transplant Proc* 1983; 15: 1157-9.

Chan RD, Greenstein SM, Tellis V, Corey HE. Renal tubular cells express ICAM-1 during allograft rejection. *Transplant Proc* 1993; 25: 915-6.

Chang TJ, Juan CC, Yin PH, Chi CW, Tsay HJ. Up-regulation of β -actin, cyclophilin and GAPDH in N1S1 rat hepatoma. *Oncol Rep* 1998; 5: 469-71.

Chen N, Field EH. Enhanced type 2 and diminished type 1 cytokines in neonatal tolerance. *Transplantation* 1995; 59: 933-41.

Chomarat P, Banchereau J. Interleukin-4 and interleukin-13: their similarities and discrepancies. *Int Rev Immunol* 1998; 17: 1-52.

Chomczynski P, Sacchi N. Single-step method of RNA extraction by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 1987; 162: 156-9.

Chun PK, Wong L, Paik YK, Siemsen AW, Hokama Y. Urinary C-reactive protein and lysozyme in renal transplantation. *Hawaii Med J* 1972; 31: 262-5.

Cohen AH, Nast CC. TGF-beta in renal allograft rejection. *Miner Electrolyte Metab* 1998. 24: 197-201.

Coles M, Rose M, Yacoub M. Appearance of cells bearing the interleukin-2 receptor in peripheral blood of cardiac transplant patients and their correlation with rejection episodes. *Transplant Proc* 1987; 19: 2546-7.

Colvin RB, Cosimi AB, Burton RC, Delmonico FL, Jaffers G, Rubin RH, Tolkoff-Rubin NE, Giorgi JV, McCluskey RT, Russell PS. Circulating T-cell subsets in 72 human renal allograft recipients: The OKT4⁺/OKT8⁺ cell ratio correlates with reversibility of graft injury and glomerulopathy. *Transplant Proc* 1983; 15: 1166-9.

Colvin RB, Fuller TC, MacKeen L, Kung PC, Ip SH, Cosimi RB. Plasma interleukin-2 receptor levels in renal allograft recipients. Clin Immunol Immunopath 1987; 43: 273-6.

Cosimi AB. The clinical value of antilymphocyte antibodies. Transplant Proc 1981; 13: 462-8.

Cosimi AB, Colvin RB, Burton RC, Rubin RH, Goldstein G, Kung PC, Hansen WP, Delmonico FL, Russell PS. Use of monoclonal antibodies to T-cell subsets for immunologic monitoring and treatment in recipients of renal allografts. New Eng J Med 1981; 305: 308-14.

Daar AS, Fuggle SV, Fabre JW, Ting A, Morris PJ. The detailed distribution of HLA A, B, C antigens in normal human organs. Transplantation 1984; 38: 287-94.

Daar AS, Fuggle SV, Fabre JW, Ting A, Morris PJ. The detailed distribution of major histocompatibility complex Class II antigens in normal human organs. Transplantation 1984; 38: 294-8.

Dallman MJ, Larsen CP, Morris PJ. Cytokine gene transcription in vascularised organ graft: analysis using semi-quantitative polymerase chain reaction. J Exp Med 1991; 174: 493-6.

Dallman MJ, Roake J, Hughes D, Toogood G, Morris PJ. Sequential analysis of IL-2 Gene transcription in renal transplantation. *Transplantation* 1992; 53: 683-5.

Daniel V, Pasker S, Wiesel M, Carl S, Pomer S, Staehler G, Schnobel R, Weimer R, Opelz G. Cytokine monitoring of infection and rejection in renal transplant recipients. *Transplant Proc* 1995; 27: 884-6.

Dassi C, Signorini S, Gerthoux P, Cazzaniga M, Brambilla P. Cytochrome p450 1B1 mRNA measured in blood mononuclear cells by quantitative reverse transcription PCR. *Clin Chem* 1998; 44: 2416-21.

Dausset J. Iso-leuco-anticorps (Iso-leuko-antibodies). *Acta Haematol* 1958; 20: 156-66.

De Boccardo G, Albano S, Alvarellos T, Massari P, Giraudo C. Soluble IL-2R assay and spontaneous blastogenesis in the immunological monitoring of kidney transplant patients. *Transplant Proc* 1994; 26: 2854-6.

DeBruyne LA, Ensley RD, Olsen SL, Taylor DO, Carpenter BM, Holland C, Swanson S, Jones KW, Karwande SV, Renlund DG, Bishop DK. Increased frequency of alloantigen-reactive helper T lymphocytes is associated with humancardiac allograft rejection. *Transplantation* 1993; 56: 722-7.

Del Prete GF, De Carli M, Mastromauro C, Biagiotti R, Macchia D, Falagiani P, Ricci M, Romagnani S. Purified protein derivative of *Mycobacterium tuberculosis* and excretory-secretory antigens of *Toxocara canis* expand in vitro human T cells with stable and opposite type 1 T helper or type 2 T helper profile of cytokine production. *J Clin Invest* 1991; 88: 346-50.

DeVito-Haynes LD, Jankowska-Gan E, Sollinger HW, Knechtle SJ, Burlingham WJ. Monitoring of kidney and simultaneous pancreas-kidney transplantation rejection by release of donor-specific, soluble HLA class I. *Hum Immunol* 1994; 40: 191-201.

DiMolfetto L, Neal HA, Wu A, Reilly C, Lo D. The density of the class II MHC cell receptor ligand influences IFN- γ /IL-4 ratios in immune response in vivo. *Cell Immunol* 1998; 183: 70-9.

Donnelly RP, Dickensheets H, Finbloom DS. The interleukin-10 signal transduction pathway and regulation of gene expression in mononuclear phagocytes. *J Interferon Cytokine Res* 1999; 19: 563-73.

Drouet M, Petit B, Peyronnet P, Aussel L, Venot J, Leroux-Robert C, Fauchet R. Soluble HLA class I antigens in kidney allograft recipients. *Transplant Proc* 1995; 27: 1671.

Dunham I, Sargent CA, Dawkins RL, Campbell RD. An analysis of variation in the long-range genomic organisation of the human major histocompatibility

complex class II region by pulsed field gel electrophoresis. *Genomics* 1989; 5: 787-96.

Edwards LC, Helderman JH, Hamm LL, Ludwin D, Gailiunas Jr P, Hull AR. Noninvasive monitoring of renal transplant function by analysis of beta2-microglobulin. *Kidney Int* 1983; 23: 767-70.

Eggensperger D, Schweitzer S, Ferriol E, O'Dowd G, Light JA. The utility of cytodiagnostic urinalysis for monitoring renal allograft injury. *Am J Nephrol* 1988; 8: 27-34.

Ellis TM, Lee HM, Mohanakumar T. Alterations in human regulatory T lymphocyte populations after renal allografting. *J Immunol* 1981; 127: 2199-2203.

El-Naggar MM, Ukai K, Takeuchi K, Sakakura Y. Expression of interferon-gamma, interleukin-4 and interleukin-5 mRNA in the nasal mucosal membrane of rats with allergic rhinitis. *Scand J Immunol* 1998; 47: 554-60.

European Multicentre Trial Group. Cyclosporin in cadaveric renal transplantation: one year follow-up of a multicentre trial. *Lancet* 1983; 2(8357): 968-9.

European Mycophenolate Mofetil Cooperation Study Group. Placebo-controlled study of mycophenolate mofetil combined with cyclosporin and corticosteroids for prevention of acute rejection. *Lancet* 1995; 345: 1321-5.

Fan J, Nishanian P, Breen EC, McDonald M, Fahey JL. Cytokine gene expression in normal human lymphocytes in response to stimulation. *Clin Diagn Lab Immunol* 1998; 5: 335-340.

Flach R, Speidel N, Flohe S, Borgermann J, Dresen IG, Erhard J, Schade FU. Analysis of intragraft cytokine expression during early reperfusion after liver transplantation using semi-quantitative RT-PCR. *Cytokine* 1998; 10: 445-51.

Förster VT. Zwischenmolekulare energiewanderung und fluoreszenz. *Ann Physics* 1948; 2: 55-75.

Forsythe JLR, Shenton BK, Parrott NR, Taylor RMR, Proud G. Plasma interleukin 2 receptor levels in renal allograft dysfunction. *Transplantation* 1989; 48: 155-7.

Foss DL, Baarsch MJ, Murtaugh MP. Regulation of HPRT, GAPDH and β -actin mRNA expression in porcine immune cells and tissues. *Animal Biotech* 1998; 9: 67-78.

Galvan B, Christopoulos TK. Quantitative reverse transcriptase-polymerase chain reaction for prostate-specific antigen. *Clin Biochem* 1997; 30: 391-7.

Garcia-Monzon C, Majano PL, Solis JA, Rodriguez S, Colina F, Lopez-Botet M, Moreno-Gonzalez E, Moreno-Otero R. Intrahepatic enhanced expression of beta2-microglobulin conformational epitope in acute liver allograft rejection: evidence of modulation by glucocorticoids. *Dig Dis Sci* 1998; 43: 1755-62.

George S, Turner D, Reynard M, Turakhia G, Navarrete C, Sweny P, Fernando ON, Powis SH, Varghese Z. Cytokine gene polymorphism may predict rejection following renal transplantation. *Transplantation* 2000; 69 Suppl: S220.

Georgi BA, Dempsey RA, Corry RJ. Interleukin-2 assay in serum and urine as a means of monitoring pancreatic allograft rejection. *Transplant Proc* 1989; 21: 2784-5.

Germain RN. MHC-dependent antigen processing and peptide presentation: providing ligands for T lymphocyte activation. *Cell* 1994; 76: 287-99.

Gimsa U, Mitchison A. New perspectives on the Th₁/Th₂ paradigm. *Curr Opin Org Transpl* 1997; 2: 18-22.

Gipson IK, Spurr-Michaud S, Moccia R, Zhan Q, Toribara N, Ho SB, Gargiulo AR, Hill 3rd JA. MUC4 and MUC5B transcripts are prevalent mucin messenger ribonucleic acids of the human endocervix. *Biol Reprod* 1999; 60: 58-64.

Goldman MR, Lippman R, Landwehr D, Szentpetery S, Wolfgang T, Lee HM, Hess M, Mendez-Pico G, Hastillo A, Lower RR. Beta 2 microglobulin and the diagnosis of cardiac transplant rejection. *Transplantation* 1983; 36: 209-11.

Gorzynski RM, Adams RB, Levy GA, Chung SW. Correlation of peripheral blood lymphocyte and intragraft cytokine mRNA expression with rejection in orthotopic liver transplantation. *Surgery* 1996; 120: 496-502.

Gould DS, Auchincloss Jr H. Direct and indirect recognition: the role of the MHC antigens in graft rejection. *Immunol Today* 1999; 20: 77-82.

Graeb C, Stangl MJ, Mebert H, Fischer T, Hammer C. Cytoimmunologic monitoring of small bowel allograft recipients. *Transplant Proc* 1992; 24: 1153-4.

Grande JP. Role of transforming growth factor-beta in tissue injury and repair. *Proc Soc Exp Bio Med* 1997; 214: 27-40.

Grant SCD, Lamb WR, Brooks NH, Brenchley PEC, Hutchinson IV. Serum cytokines in human heart transplant recipients. Is there a relationship to rejection? *Transplantation* 1996; 62: 480-91.

Grimm PC, McKenna RM, Gospodarek EM, Jeffery JR, Rush DN. Low frequency of infiltrating cells intensely expressing T cell cytokine mRNA in human renal allograft rejection. *Transplantation* 1995; 59: 579-84.

Groth CG. Antilymphocyte globulin (ALG). *Transplant Proc* 1981; 13: 460-1.

Groth GC. Landmarks in clinical renal transplantation. *Surg Gynaecol Obstet* 1972; 134: 323-8.

Groth CG, Bäckman L, Morles J-M et al for the Sirolimus European Renal Transplant Study Group. Sirolimus (rapamycin)-based therapy in human renal transplantation. *Transplantation* 1999; 67: 1036-42.

Halminen M, Sjöroos M, Makela MJ, Waris M, Terho E, Lovgren T, Ilonen J. Simultaneous detection of IFN-gamma and IL-4 mRNAs using RT-PCR and time-resolved fluorometry. *Cytokine* 1999; 11: 87-93.

Hammer C, Reichenspurner H, Klima G. Immunologic parameters for the diagnosis of graft dysfunction. *Transplant Proc* 1993; 25: 26-9.

Hammer C, Reichenspurner R, Meiser B, Reichart B. Cytoimmunology in monitoring: the Munich experience. *Transplant Proc* 1998; 30: 873-4.

Harris KR, Digard NJ, Slapak M, Lee HA. Serum C-reactive protein: A useful, economical marker of immune activation in cadaveric renal transplantation. *Transplant Proc* 1995; 27: 1046-7.

Harrison DC, Medhurst HD, Bond BC, Campbell CA, Davis RP, Philpott KL. The use of quantitative RT-PCR to measure mRNA expression in a rat model of focal ischaemia. *Molecular Brain Research* 2000; 75: 143-9.

Härtel C, Bein G, Kirchner H, Klüher H. A human whole blood assay for analysis of T-cell function by quantification of cytokine mRNA. *Scand J Immunol* 1999; 49: 649-54.

Heaf JG, Ladefoged J. The effect of acute rejection on long-term renal graft survival is mainly related to the initial renal damage. *Transpl Int* 1998; 11(suppl 1): S26-S31.

Heid CA, Stevens J, Livak KJ, Williams PM. Real time quantitative PCR. *Genome Res* 1996; 6: 986-94.

Higuchi R, Dollinger G, Walsh PS, Griffith R. Simultaneous amplification and detection of specific DNA sequences. *Biotechnology* 1992; 10: 413-7..

Higuchi R, Fockler C, Dollinger G, Watson R. Kinetic PCR: Real monitoring of DNA amplification reactions. *Biotechnology* 1993; 11: 1026-30.

Hildebrandt U, Kessler K, Pistorius G, Lindemann W, Ecker KW, Feifel G, Menger MD. Granulocyte elastase and systemic cytokine response after laparoscopic-assisted and open resections in Crohn's disease. *Dis Colon Rectum* 1999; 42: 1480-6.

Hojo M, Morimoto T, Maluccio M, Asano T, Morimoto K, Lagman M, Shimbo T et al. Cyclosporine induces cancer progression by a cell-autonomous mechanism. *Nature* 1997; 397: 530-4.

Holland PM, Abramson RD, Watson R, Gelfand DH. Detection of specific polymerase chain reaction product by utilizing the 5' and 3' exonuclease activity of *Thermus aquaticus* DNA polymerase. *Proc Natl Acad Sci USA* 1991;88: 7276-80.

Holmberg A, Berqvist D, Westman B, Siegbahn A. Cytokine and fibrinogen response in patients undergoing open abdominal aortic aneurysm surgery. *Eur J Vasc Endovasc Surg* 1999; 17: 294-300.

Hoshinaga K, Mohanakumar T, Wood NL, Szentpetery S, Lower RR. Increase in transferrin receptors as a marker for rejection and infection in human cardiac transplantation. *Transplant Proc* 1987; 19: 2534-6.

Huebers HA, Finch CA. The physiology of Transferrin and Transferrin Receptors. *Physiol Rev* 1987; 67: 520- 82

Hume DM. Early experiences in organ homotransplantation in man and the unexpected sequelae thereof. *Am J Surg* 1979; 137: 152-61.

Hutchinson IV. The role of transforming growth factor-beta in transplant rejection. *Transplant Proc* 1999; 31(7A): 9S-13S.

Hutchinson IV, Pravica V, Hajeer A, Sinnott PJ. Identification of high and low responders to allografts. *Rev Immunogenetics* 1999; 1: 323-333.

Hutchinson IV, Pravica V, Perrey C, Sinnott P. Cytokine gene polymorphisms and relevance to forms of rejection. *Transpalnt Proc* 1999; 31: 734-6.

Iacono A, Dauber J, Keenan R, Spichty K, Cai J, Grgurich W, Burckart G, Smaldone G, Pham S, Otori NP, Yousem S, Williams P, Griffith B, Zeevi A. Interleukin 6 and interferon- γ gene expression in lung transplant recipients with refractory acute cellular rejection. Implications for monitoring and inhibition by treatment with aerosolized cyclosporine. *Transplantation* 1997; 64: 263-9.

Jain S, Furness PN, Nicholson ML. The role of transforming growth factor beta in chronic renal allograft nephropathy. *Transplantation* 2000; 69: 1759-66.

Jenkins MK, Schwartz RH. Antigen presentation by chemically modified splenocytes induces antigen specific T cell unresponsiveness in vitro and in vivo. *J Exp Med* 1987; 165: 302-19.

Jeyarajah DR, Kadakia RA, O'Toole K, Newell KA, Josephson MA, Spargo BH, Woodle ES, Thislethwaite jr JR. Changes in urinary cytokine mRNA profile after successful therapy for acute cellular renal allograft rejection. *Transplant Proc* 1995; 27: 887-9.

Johnson CP, Chaharmahal A, Buchmann E, Roza AM, Adams MB. Plasma IL-2 levels and diagnosis of renal transplant rejection. *Transplant Proc* 1990; 22: 1849-51.

Jordan ML, Naraghi R, Shapiro R, Smith D, Vivas CA, Scantlebury VP, Gritsch HA, McCauley J, Randhawa P, Demetris AJ, McMichael J, Fung JJ, Starzl TE. Tacrolimus rescue therapy for renal allograft rejection – five-year experience. *Transplantation* 1997; 63: 223-8.

Joseph JV, Guy SP, Brenchley PE, Parrott NR, Short CD, Johnson RW, Hutchinson IV. Th1 and Th2 cytokine gene expression in human renal transplants. *Transplant Proc* 1995; 27: 915-6.

June CH, Ledbetter JA, Linsley PS, Thompson CB. Role of the CD28 receptor in T-cell activation. *Immunol Today* 1990; 11: 211-6.

Kahan BD, Podbielski J, Napoli KL, Katz SM, Meier-Kriesche H-U, Van Buren CT. Immunosuppressive effects and safety of a sirolimus/cyclosporine combination regimen for renal transplantation. *Transplantation* 1998; 66:1040-6.

Kaminski ER, Kaminski A, Bending MR, Chan R, Heads, A, Szydlo RM, Pereira RS. In vitro cytokine profiles and their relevance to rejection following renal transplantation. *Transplantation* 1995; 60: 703-6.

Kappes D, Strominger JL. Human Class II major histocompatibility complex genes and proteins. *Annu Rev Biochem* 1988; 57: 991-1028.

Kauffman jnr HM, Clark RF, Magee JH. Lymphocytes in urine as an aid in the early detection of renal homograft rejection. *Surg Gynecol Obst* 1964; 119: 25.

Kawamura T, Inada K, Nara N, Wakusawa R, Shigeatsu E. Influence of methyl prednisolone on cytokine balance during cardiac surgery. *Crit Care Med* 1999; 27: 545-8.

Kerman RH, Buyse IM, Warnell S, McKissick E, Garcia C, Kahan BD. Cytokine genotyping in caucasian and African-American renal allograft recipients. *Transplantation* 2000; 69 Suppl: S168.

Kerman RH, Flechner SM, Van Buren CT, Payne W, Kahan BD. Immunologic monitoring of renal allograft recipients treated with cyclosporine. *Transplant Proc* 1983; 15 (Suppl 1): 2302-5.

Kerman RH, Van Buren CT, Payne W, Flechner S, Kahan BD. Monitoring of T-cell subsets and immune events in renal allograft recipients. *Transplant Proc* 1983; 15: 1170-2.

Khanna A, Li B, Stenzel KH, Suthanthiran M. Regulation of new synthesis in mammalian cells by cyclosporine. Demonstration of transforming growth factor beta-dependant mechanism of inhibition of cell growth. *Transplantation* 1994; 57: 577-82.

Kimball PM, Radovancevic B, Isom T, Spickard A, Frazier OH. The paradox of cytokine monitoring – predictor of immunologic activity as well as immunologic silence following cardiac transplantation. *Transplantation* 1996; 61: 909-15.

Kirk AD, Bollinger RR, Finn OJ. Rapid, Comprehensive analysis of human cytokine mRNA and its application to the study of acute renal allograft rejection. *Hum Immunol* 1995; 43: 113-28.

Kissmeyer-Nielson F, Svejgaard A, Hauge M. Genetics of the human HL-A transplantation system. *Nature* 1968; 219: 1116-9.

Klanke D, Hammer C, Dirchedl P, Kemkes BM, Reichart B, Gokel M, Krombach F. Sensitivity and specificity of cyto-immunological monitoring in correlation with endomyocardial biopsies in heart transplant patients. Transplant Proc 1987; 19: 3781-3.

Klausener RD, Samelson LE. T-cell antigen receptor activation pathways: the tyrosine kinase connection. Cell 1991; 64: 875-8.

Koller BH, Geraghty DE, DeMars R, Duvick L, Rich SS, Orr HT. Chromosomal organisation of the human major histocompatibility complex Class I gene family. J Exp Med 1989; 169: 469-80.

Kondo H, Nachtigal D, Frenkiel S, Schotman E, Hamid Q. Effects of steroids on nasal inflammatory cells and cytokine profile. Laryngoscope 1999; 109: 91-7.

Konieczny BT, Saleem SS, Lowry RP, Lakkis FG. Vigorous cardiac and skin allograft rejection in the absence of IFN- γ . J Am Soc Nephrol 1996; 7: 1887-95.

Krams SM, Falco DA, Villaneuva JC, Rabkin J, Tomlanovich SJ, Vincenti F, Amend WJ, Melzer J, Garovoy MR, Roberts JP et al. Cytokine and T cell receptor gene expression at the site of allograft. Transplantation 1992; 53: 151-6.

Kreuzer KA, Lass U, Landt O, Nitsche A, Laser J, Ellerbrok H, Pauli G, Huhn D, Schmidt CA. Highly selective and specific fluorescence reverse transcription-PCR assay for pseudogene-free detection of β -actin transcripts as quantitative reference. Clin Chem 1999; 45: 297-300.

Kruse N, Pette M, Toyka K, Rieckmann P. Quantification of cytokine mRNA expression by RT PCR in samples of previously frozen blood. J Immunol Methods 1997; 210: 195-203.

Kusaka S, Grailer AP, Fechner JH, Burlington WJ. Evidence for a possible Th2 bias in human renal transplant tolerance. Transplant Proc 1995; 27: 225-6.

Kutukculer N, Clark K, Rigg KM, Forsythe JLR, Proud G, Taylor RMR, Shenton BK. The value of posttransplant monitoring of interleukin (IL)-2, IL-3, IL-4, IL-6, IL-8 and soluble CD23 in the plasma of renal allograft recipients. Transplantation 1995; 59: 333-40.

Lafferty KJ, Bootes A, Dart G, Talmage DW. Effect of organ culture on the survival of thyroid allografts in mice. Transplantation 1976; 22: 138-49.

Lakowicz JR. Energy transfer. In: Principles of Fluorescent Spectroscopy, Plenum Press, N.Y. 1983: pp. 303-39.

Lalla MLT, Eklund B, Ahonen. The clinical significance of serum C-reactive protein after renal transplantation. Transplant Proc 1988; 20: 402-4.

Laman JD, Claassen E, Noelle RJ. Functions of CD40 and its ligand, gp39 (CD40L). *Crit Rev Immunol* 1996; 16: 59-108.

Langley J, Dickerman R, Vergne-Marini P, Williams J, Jensik S, Kenyon N, Anderson J. Inducible IL-2 on CD8 lymphocytes as a postoperative predictor of rejection in renal allograft recipients. *Transplant Proc* 1993; 25: 939-43.

Leccia JR, Milano G, Formento P, Francoual M, Formento JL, Pierrefite V. A clinically applicable assay for tumoral thymidylate synthase combining reverse transcriptase-PCR and high performance liquid chromatography. *Eur J Cancer* 1998; 34: 182-7.

Lipman ML, Shen Y, Jeffery JR, Gough J, McKenna RM, Grimm PC, Rush DN. Immune-activation gene expression in clinically stable renal allograft biopsies: molecular evidence for subclinical rejection. *Transplantation* 1998; 66: 1673-81.

Loong CC, Chen A, Lui WY, King KL, Lin CY. Expression of cytokines, growth factors and adhesion molecules in rejecting human renal allograft. *Transplant Proc* 1996; 28: 1445-6.

Mao D, Lee JK, VanVickle SJ, Thompson RW. Expression of collagenase-3 (MMP-13) in human abdominal aortic aneurysms and vascular smooth muscle cells in culture. *Biochem Biophys Res Commun* 1999; 261: 904-10.

Marshall SE, McLaren AJ, Haldar NA, Morris PJ, Welsh KI. The impact of recipient cytokine genotype on acute rejection after renal transplantation. Presented at the BTS, Cardiff 2000.

Marshall SE, Welsh KI. Tumour necrosis factor polymorphisms and acute rejection: proven link or provisional association? *Transplantation* 2000; 69: 1237-8.

Martin P. Wound healing – aiming for perfect skin regeneration. *Science* 1997; 276: 75-81.

Masri MA, Stephan A, Barbari A, Rizk S, Kamel G. Future trends in immunological monitoring of transplant patients. *Transplant Proc* 1998; 30: 2979-81.

Masri MA, Stephan A, Barbari A, Rizk S, Kamel G. Logical approach to transplant patient monitoring. *Transplant Proc* 1999; 31: 3353-5.

Maury CPJ, Teppo A-M. Comparative study of serum amyloid-related protein SAA, C-reactive protein, and β 2-microglobulin as markers of renal allograft rejection. *Clin Nephrol* 1984; 22: 284-92.

Maury CPJ, Teppo A-M. Raised serum levels of cachectin/tumor necrosis factor α in renal allograft rejection. *J Exp Med* 1987; 166: 1132-7.

Maury CPJ, Teppo A-M. Serum immunoreactive interleukin-1 in renal transplant recipients. *Transplant Proc* 1987; 19: 4349-50.

May WS Jr, Cuatrecasas P. Transferrin receptor: it's biological significance. *J Membr Biol* 1985; 88: 205-15.

McLaughlin PJ, Aikawa AA, Davies HM, Bakran A, Sells RA, Johnson PM. Tumour necrosis factor in renal transplantation. *Transplant Proc* 1991; 23: 1289-90.

Miceli MC, Parnes JR. The role of CD4 and CD8 in T-cell activation. *Semin Immunol* 1991; 3: 133-141.

Miyamoto S, Safer B. Immunosuppressants FK506 and rapamycin have different effects on the biosynthesis of cytoplasmic β -actin during the early period of T cell activation. *Biochem J* 1999; 344: 803-12.

Morehouse CA, Owen RD. Exposure of Daudi cells to low-frequency magnetic fields does not elevate MYC steady-state mRNA levels. *Radiat Res* 2000; 153: 663-9.

Moriyama N, Yamaguchi T, Takeuchi T, Sakamoto E, Ueki T, Tsujimoto G, Kawabe K. Semiquantitative evaluation of α 1A-adrenoceptor subtype mRNA in human hypertrophied and non-hypertrophied prostates: regional comparison. *Life Sci* 1999; 64: 201-10.

Mosmann TR, Cherwinski H, Bond MW, Giedlin MA, Coffman RL. Two types of murine helper T cell clone. I. Definition according to profiles of lymphokine activities and secreted proteins. *J Immunol* 1986; 136: 2348-57.

Mosmann TR, Coffman RL. TH1 and TH2 cells: Different patterns of lymphokine secretion lead to different functional properties. *Annu Rev Immunol* 1989; 7: 145-73.

Murphy PA. The Interleukins. In: *Kidney Transplant Rejection: Diagnosis and Treatment*. Ed. Williams GM, Solez K. New York 1986: Maul Dekker, p55.

Murphy GP, Williams PD, Merrin CE. Diagnostic value of lymphocyturia in renal allograft rejection in man. *Urology* 1973; 2: 227-30.

Murray JE, Merrill JP, Harrison JH. Renal homotransplantation in identical twins. *Surg Forum* 1955; 6: 565-73.

Naar JD, Fisher RA, Saggi BH, Obias V, Goggins WC, Tawes JW, Wakely jnr PE, Posner MP. The immunological role of IFN-gamma in ACI to Lewis kidney transplantation. *J Surg Res* 1998; 78: 103-7.

Nashan B, Moore R, Amlot P, Schmidt A-G, Abeywickrama K, Souillou J-P for the CHIB 201 International Study Group. Randomised trial of basiliximab versus placebo for control of acute rejection in renal allograft recipients. *Lancet* 1997; 350: 1193-8.

Nast CC, Zuo XJ, Prehn J, Danovitch GM, Wilkinsin A, Jordan SC. Gamma-interferon gene expression in human renal allograft fine-needle aspirates. *Transplantation* 1994; 57: 498-502.

Nathan N, Preux PM, Feiss P, Denziot Y. Plasma interleukin-4, interleukin-10, and interleukin-13 concentrations and complications after coronary artery bypass graft surgery. *J Cardiothorac Vasc Anesth* 2000; 14: 156-60.

•

Neefjes JJ, Stollory V, Peters PJ, Geuze HJ, Ploegh HL. The biosynthetic pathway of MHC Class II but not Class I molecules intersects with the endocytic route. *Cell* 1990; 61: 171-83.

Nelms K, Keegan AD, Zamorano J, Ryan JJ, Paul WE. The IL-4 receptor: signaling mechanisms and biological functions. *Ann Rev Immunol* 1999; 17: 701-38.

Neylan J, de Smet W, Stijlemans B, Brinker K, Gonwa T, Moran J, Segers A on behalf of the anti-CD45 study group. Detection of clinically relevant antibodies pretransplant and posttransplant with PRA-STAT. *Transplant Proc* 1997; 29: 330-2.

Nickerson P, Steiger J, Zheng XX, Steele AW, Steurer W, Roy-Chaudhury P, Strom TB. Manipulation of cytokine networks in transplantation. *Transplantation* 1997. 63: 489-94.

Nickerson PW, Zheng XX, Steiger J, Steele AW, Steurer W, Roy Chaudhury P, Muller W, Strom TB. Prolonged islet allograft acceptance in the absence of interleukin 4 expression. *Transpl Immunol* 1996; 4: 81-5.

Nishizuka Y. Intracellular signalling by hydrolysis of phospholipids and activation of protein kinase C. *Science* 1992; 258: 607-14.

Noguchi K, Yoshida Y, Yagihashi A, Kita Y, Takeaka T, Terasawa K, Hayashi S, Konno A, Kobayashi M, Nakamura K, Reyes J, Abu-Elmagd K, Demetris AJ, Tzakis AG, Todo S, Iwaki Y, Starzl TE. Serum levels of interleukin-6, tumor necrosis factor- α , and interleukin-2 in rejecting human small bowel allografts. *Transplant Proc* 1992; 24: 1152.

Noronha IL, Daniel V, Rambausek M, Waldherr R, Opelz G. Soluble Interleukin-2 receptor (sIL-2R) and Tumour necrosis factor plasma levels in renal allograft recipients. *Transplant Proc* 1990; 22: 1859-60.

O'Kane S, Ferguson MW. Transforming growth factor beta and wound healing. *Int J Biochem Cell Bio* 1997; 29: 63-78.

Opelz G. HLA matching should be utilised for improving kidney transplant success rate. *Transplant Proc* 1991; 23: 46-50.

Overbergh L, Valckx D, Waer M, Mathieu C. Quantification of murine cytokine mRNAs using real time quantitative reverse transcriptase PCR. *Cytokine* 1999; 11: 305-12.

Pallisgaard N, Clausen N, Schroder H, Hokland P. Rapid and sensitive minimal residual disease detection in acute leukaemia by quantitative real-time RT-PCR exemplified by t(12;21) TEL-AML1 fusion transcript. *Genes Chromosomes Cancer* 1999; 26: 355-65.

Paludan SR. Interleukin-4 and interferon-gamma: the quintessence of a mutual antagonistic relationship. *Scand J Immunol* 1998; 48: 459-68.

Pattanapanyasat K, Hoy TG. Expression of cell surface Transferrin Receptor and intracellular ferritin after in vitro stimulation of peripheral blood T lymphocytes. *Eur J Haematol* 1991; 47: 140-5.

Paul WE. Pleiotrophy and redundancy: T cell-derived lymphokines in the immune response. *Cell* 1989; 57: 521-4.

Pavlakakis M, Strelau J, Lipman M, Shapiro M, Maslinski W, Strom TB. Intragraft IL-15 Transcripts are increased in human renal allograft rejection. *Transplantation* 1996; 62: 543-5.

Perkins JD, Nelson DL, Rakela J, Grambsch PM, Krom RAF. Soluble interleukin-2 receptor level as an indicator of liver allograft rejection. *Transplantation* 1989; 47: 77-81.

Pescovitz MD, Thistlethwaite JR jr, Auchinson H jr. Effect of class II antigen matching on renal allograft survival in miniature swine. *J Exp Med* 1984; 160: 1495-1508.

Petrovsky N, Harrison LC. HLA class II-associated polymorphism of interferon- γ production. *Human Immunol* 1997; 53: 12-16.

Piazza A, Borrelli L, Buonomo O, Pisani F, Valeri M, Torlone N, Felici A, Monaco PI, Adorno D, Casciani CU. Flow cytometry crossmatch and kidney allograft outcome. *Transplant Proc* 1999; 31: 314-6.

Piccotti JR, Chan SY, Vanbushkirk AM, Eichwald EJ, Bishop DK. Are Th2 helper lymphocytes beneficial, deleterious, or irrelevant in promoting allograft survival? *Transplantation* 1997; 63: 619-24.

Pollard SG, Lear PA, Ready AR, Moore RH, Johnson RWG on behalf of the U.K. Neoral Renal Study Group. Comparison of microemulsion and conventional formulations of cyclosporine A in preventing acute rejection in de novo kidney transplant patients. *Transplantation* 1999; 68: 1325-1331.

Poole KL, Gibbs PJ, Evans PR, Sadek SA, Howell WM. Influence of patient and donor cytokine genotype on renal allograft rejection: evidence from a single centre. *Transplant Immunol* 2001; 8: 259-65.

Pravica V, Asderakis A, Perrey C, Hajeer A, Sinnott PJ, Hutchinson IV. In vitro production correlates with CA repeat polymorphism in the human IFN- γ gene. *Eur J Immunogenet* 1999; 26: 1-3.

Pretolani M. Interleukin-10: an anti-inflammatory cytokine with therapeutic potential. *Clin Exp Allergy* 1999 29: 1164-71.

Prischl F, Gremmel F, Schwabe M, Schindler J, Balcke P, Kopsa H, Pinter G, Schwarzmeier J, Zazgornik J. Beta-2-microglobulin for differentiation between ciclosporin A nephrotoxicity and graft rejection in renal transplant recipients. *Nephron* 1989; 51: 330-7.

Puig N, Pallardó LM, Villalba JV, Sánchez J, Crespo J, Rodriguez R, Montoro JA. Donor-specific flow cytometric cross-match after kidney transplantation. *Transplant Proc* 1995; 27: 2369-70.

Puppo F, Pellicci R, Brenci S, Nocera A, Morelli N, Dardano G, Bertocchi M, Antonucci A, Ghio M, Scudeletti M et al. HLA class-I-soluble antigen serum levels in liver transplantation. A predictor marker of acute rejection. *Hum Immunol* 1994; 40: 166-70.

Reddy NR, Wilkie BN. Quantitation of porcine cytokine and β 2-microglobulin mRNA expression by reverse transcription polymerase chain reaction. *J Immunol Methods* 2000; 233: 83-93.

Roberti I, Lieberman KV, Manzarbeitia C, Schwartz M, Reisman L, Miller C. Evidence that the systematic analysis of bile cytology permits monitoring of hepatic allograft rejection. *Transplantation* 1992; 54: 471-4.

Rokahr KL, Sharland AF, Sun J, Wang C, Sheil AG, Yan Y, McCaughan GW, Bishop GA. Paradoxical early immune activation during acceptance of liver allografts compared with rejection of skin grafts in a rat model of transplantation. *Immunology* 1998; 95: 257-63.

Rondeau E, Cerrina J, Delarue F, Ladurie FL, Herve Ph, Chapelier A, Dartevielle Ph, Sraer JD. Tumour necrosis factor alpha (TNF-alpha) production by cells of broncioloalveolar lavage (BAL) and peripheral blood mononuclear cells (PBMC) in cardiopulmonary transplant recipients. *Transplant Proc* 1990; 22: 1855-6.

Rosenwasser LJ, Klemm DJ, Dreback JK, Inamura H, Mascali JJ, Klinnert M, Borish L. Promoter polymorphisms in the chromosome 5 gene cluster in asthma and atopy. *Clin Exp Allergy* 1995; 25 Suppl 2: 74-8.

Rudensly AY, Preston Hurlburt P, Hong SC, Barlow A, Janeway CA. Sequence analysis of peptides bound to MHC Class II molecules. *Nature* 1991; 353: 622-7.

Sandberg L, Thorsby E, Kissmeyer-Nielson F, Lindholm A. In: Terasaki P (ed) *Histocompatibility testing*. Copenhagen: Munkgaard, 1970, pp 165-9.

Sandoz PF, Bielman D, Mihatsch M, Thiel G. Value of urinary sediment in the diagnosis of interstitial rejection in renal transplants. *Transplantation* 1986; 41: 343-8.

Sankaran D, Asderakis A, Ashraf S, Roberts ISD, Short CD, Dyer PA, Sinnott PJ, Hutchinson IV. Cytokine gene polymorphisms predict acute graft rejection following renal transplantation. *Kid Int* 1999; 56: 281-8.

Sankary H, Chong A, Frye E, Scuderi P, Foster P, McChesney L, Williams J. Daily determinations of serum lymphotoxin allows for accurate early diagnosis of hepatic allograft rejection. *Transplant Proc* 1993; 25: 928-30.

Sasada K, Miyazaki M, Nakagawa K, Ito H, Ambiru S, Shimizu H, Nakajima N. Augmented enhancement of in vitro production of cytokines in peripheral blood mononuclear cells in patients undergoing simultaneous resection of the liver and gastrointestinal tract. *Crit Care Med* 1999; 27: 929-36.

Sayegh MH, Akalin E, Hancock WW, Russell ME, Carpenter CB, Linsley PS, Turka LA. CD28-B7 blockade after alloantigenic challenge in vivo inhibits Th1 cytokines but spares Th2. *J Exp Med* 1995;181: 1869-74.

Schiano TD, Fiel MI, Abe K, Thung SN, Bodenheimer HC. Absence of hepatitis G virus within liver tissue of patients undergoing transplantation for cryptogenic cirrhosis. *Transplantation* 1999; 67: 1193-7.

Schumann GB, Burleson RL, Henry JB, Jones DB. Urinary cytodiagnosis of acute renal allograft rejection using cytocentrifuge. *Am J Clin Pathol* 1977; 67: 134-40.

Schwartz RH. Models of T cell anergy: Is there a common molecular mechanism? *J Exp Med* 1996; 184: 1-8.

Schweizer RT, Moore R, Bartus SA, Bow L, Hayden. Beta 2-microglobulin monitoring after renal transplantation. *Transplant Proc* 1981; 13: 1620-3.

Sengar DPS, Rashid A, Harris JE. Post-transplant monitoring of renal allograft recipients for T, B and Null lymphocyte subpopulations. *Clin Exp Immunol* 1977; 28: 123-9.

Sharma VK, Bologa RM, Xu GP, Li B, Mouradian J, Wang J, Serur D, Rao V, Suthanthiran M. Intragraft TGF-beta1 mRNA: a correlate of interstitial fibrosis and chronic allograft nephropathy. *Kidney Int* 1996; 49: 1297-1303.

Simpson MA, Madras PN, Cornaby AJ, Etienne T, Dempsey RA, Clowes GHA, Monaco AP. Sequential determinations of urinary cytology and plasma and urinary lymphokines in the management of renal allograft recipients. *Transplantation* 1989; 47: 218-23.

Simpson MA, Madras PN, Crosson AW, Georgi BA, Monaco AP. Correlation of binding of human immunoglobulin, urinary cytology, and needle biopsy in the early post-transplant period. *Transplant Proc* 1987; 19: 1662-3.

Simpson M, Madras PN, Monaco AP. Binding of antihuman globulin by exfoliated renal tubular cells following renal transplantation. *Transplantation* 1985; 39: 400-5.

Simpson MA, Young-Fadok TM, Madras PN, Freeman RB, Dempsey RA, Shaffer D, Lewis D, Jenkins RL, Monaco AP. Sequential interleukin 2 and interleukin 2 receptor levels distinguish rejection from cyclosporine toxicity in liver allograft recipients. *Arch Surg* 1991; 126: 717-20.

Steiger J, Nickerson PW, Steurer W, Moscovitch Lopatin M, Strom TB. IL-2 knockout recipient mice reject islet cell allografts. *J Immunol* 1995; 155:489-98.

Steiner G, Zlabinger G, Karamelic J, Pohanka E, Kovarik J, Woloszczuk W. Interferon-gamma, neopterin and interleukin-2 receptor levels in the serum of kidney transplant recipients. *Transplant Proc* 1990; 22: 1857-8.

Steinhoff J, Bühner U, Feddersen A, Wood WG, Wiedermann G, Preuss R, Fricke L, Färber P, Hoyer J, Sack K. Analysis of C-reactive protein in urine as an aid in the clinical diagnosis of disturbed renal transplant function. *Transplant Proc* 1992; 24: 2735-7.

Steinhoff J, Einecke G, Niederstadt C, De Groot K, Fricke L, Machnik H, Sack K. Renal graft rejection or urinary tract infection? The value of myeloperoxidase, C-reactive protein and α 2-macroglobulin in the urine. *Transplantation* 1997; 64: 443-7.

Stenzel KH, Rubin AL, Nogovgradsky A. Blastogenic response to mitogens: an approach to the study of cell-surface topography in lymphocytes from transplant recipients. *Transplant Proc* 1978; 10: 589-91.

Stockenhuber F, Kramer G, Schenn G, Pec M, Keil M, Gnant M, Steininger R, Balcke P, Mühlbacher. Circulating ICAM-1: Novel parameter of renal graft rejection. *Transplant Proc* 1993; 25: 919-20.

Stordeur P, Goldman M. Interleukin-10 as regulatory cytokine induced by cellular stress: molecular aspects. *Int Rev Immunol* 1998; 16: 501-22.

Strehlau J, Pavlakis M, Lipman M, Shapiro M, Vasconcellos L, Harmon W, Strom T. Quantitative detection of immune activation transcripts as a diagnostic tool in kidney transplantation. *Proc Natl Acad Sci USA* 1997; 94: 695-700.

Strom TB, Carpenter Cb, Busch GJ. Identity and cytotoxic capacity of cells infiltrating renal allografts. *N Engl J Med* 1975; 292: 1257-63.

Sundstrom JB, Ansari AA. Comparative study of the role of professional versus semiprofessional or non-professional antigen presenting cells in the rejection of vascularised organ allografts. *Trans Immunol* 1995; 3: 273-89.

Suthanthiran M. Acute rejection of renal allografts: Mechanistic insights and therapeutic options. *Kidney Int* 1997; 51: 1289-1304.

Suthanthiran M. A novel model for antigen-dependent activation of normal human T-cells. Transmembrane signalling by crosslinkage of the CD3/T cell receptor- α/β complex with the cluster determinant 2 antigen. *J Exp Med* 1990; 171: 1965-79.

Suthanthiran M. Signalling features of T-cells: Implications for the regulation of the anti-allograft response. *Kidney Int* 1993; 44 (suppl 43): S3-S11.

Suthanthiran M, Strom T. Mechanisms and management of acute renal allograft rejection. *Surg Clinics North Am* 1998; 78: 77-94.

Suzuki T, Higgins PJ, Crawford DR. Control selection for RNA quantitation. *BioTechniques* 2000; 29: 332-7.

Tagaya Y, Bamford RN, DeFilippis AP, Waldman TA. IL-15: a pleiotrophic cytokine with diverse receptor/signalling pathways whose expression is controlled at multiple levels. *Immunity* 1996; 4: 329-36.

Takeuchi T, Lowry RP, Konieczny B. Heart allografts in murine systems. The differential activation of Th2-like effector cells in peripheral tolerance. *Transplantation* 1992; 53: 1281-94.

Talbot D. The flow cytometric crossmatch in perspective. *Transplant Immunol* 1993; 1: 155-62.

Tan L, Howell WM, Smith JL, Sadek SA. Sequential monitoring of peripheral T-lymphocyte cytokine gene expression in the early post renal allograft period. *Transplantation* 2001; 71: 751-9.

Tanimura S, Taguchi Y, Kunori T, Satomi S, Kaneda I, Kasai M. Monitoring of in vivo blastogenic response in kidney transplant patients by flow cytometry. *Transplant Proc* 1985; 17: 652-5.

Teppo A-M, Honkanen E, Ahonen J, Gronhagen-Riska C. Does increased urinary interleukin-1 receptor antagonist/interleukin-1 β ratio indicate good prognosis in renal transplant recipients? *Transplantation* 1998; 66: 1009-14.

Thai NL, Fu S, Qian S, Sun H, Gao L, Wang SC, Demetris AJ, Woo J, Thomson AW, Duquesnoy RJ, Fung JJ. Cytokine mRNA profiles in mouse

orthotopic liver transplantation. Graft rejection is associated with augmented TH1 function. Transplantation 1995; 59: 274-81.

The Royal College of Surgeons of England. The report of the working party to review organ transplantation. January 1999.

Thomas F, Thomas J, Mendez G, Lower R, Lee HM. Differential effects of immunosuppressive drugs on human T, B and K lymphocytes. Transplantation 1977; 24: 435-41.

Thompson CB, Lindsten T, Ledbetter JA, Kunkel SL, Young HA, Emerson SG, Leiden JM, June CH. CD 28 activation pathway regulates the production of multiple T-cell derived lymphokines/cytokines. Proc Natl Acad Sci USA 1989; 86: 1333-7.

Tice DG, Heintz S, Scruggs B, Hubbell C, Szmalc FS. Immune monitoring: Does the reality fulfill the Promise? Transplant Proc 1993; 25: 2654.

Tilg H, Ceska M, Vogel W, Herold M, Margeiter R, Huber C. Interleukin-8 serum concentrations after liver transplantation. Transplantation 1992; 53: 800-3.

Tilg H, Nordberg J, Vogel W, Luger TA, Herold M, Aulitzky WE, Margreiter R, Huber C. Circulating serum levels of interleukin 6 and C-reactive protein after liver transplantation. Transplantation 1992; 54: 142-6.

Torre-Amione G, MacLellan W, Kapadia S, Weilbaecher D, Farmer J, Young J, Mann D. Tumor necrosis factor- α is persistently expressed in cardiac allografts in the absence of histological or clinical evidence of rejection. *Transplant Proc* 1998; 30: 875-7.

Traindl O, Gisinger C, Reading S, Rosenkranz AR, Zlabinger GJ. Circulating serum levels of soluble CD23 (sCD23) after renal transplantation. *Clin Transplant* 1994; 8: 87-92.

Trowsdale J, Ragoussis J, Campbell RD. Map of the human MHC. *Immunol Today* 1991; 12: 433-6.

Turner DM, Grant SC, Yonan N, Sheldon S, Dyer PA, Sinnott PJ, Hutchinson IV. Cytokine gene polymorphism and heart transplant rejection. *Transplantation* 1997; 64: 776-9.

Turner DM, Williams DM, Sankaran D, Lazarus M, Sinnott PJ, Hutchinson IV. An investigation of polymorphism in the interleukin-10 gene promoter. *Eur J Immunogenet* 1997; 24: 1-8.

Tyden G, Reinholt FP, Brattstrom C, Lundgren G, Wilczek H, Bolinder J, Ostman J, Groth CG. Diagnosis of rejection in recipients of pancreatic grafts with enteric exocrine diversion monitoring pancreatic juice cytology and amylase excretion. *Transplant Proc* 1987; 19: 3892-4.

Unanue ER, Cerotoni J-C. Antigen presentation. FASEB J 1989; 3: 2496-2502.

Valeri M, Piazza A, Torlone N, Poggi E, Adorno D, Casciani CU. Immunologic monitoring in kidney transplantation. Transplant Proc 1991; 23: 2275-6.

Van Berge Henegouwen MI, van der Poll T, van Deventer SJ, Gouma DJ. Peritoneal cytokine release after elective gastrointestinal surgery and postoperative complications. Am J Surg 1998; 175: 311-6.

Vandenbroecke C, Caillat-Zucman S, Legendre C, Noel LH, Kreis H, Woodrow D, Bach JF, Tovey MG. Differential in situ expression of cytokines in renal allograft rejection. Transplantation 1991; 51: 602-9.

Van Gelder T, Baan CC, Balk AHMM, Knoop CJ, Holweg, CTJ, Van der Meer P, Mochtar B, Zondervan PE, Niesters HGM, Weimar W. Blockade of the IL-2/IL-2 receptor pathway with a monoclonal anti Interleukin-2 receptor antibody (BT563) does not prevent the development of acute heart allograft rejection in man. Transplantation 1998; 65:405-410.

Van Leeuwen A, Schuit HRE, Van Rood JJ. Typing for MLC (LDII) the selection of non-stimulator cells by MLC inhibition tests using ST identical stimulator cells (MISIS) and fluorescent antibody studies. Transplant Proc 1973; 5: 1539-42.

Van Oers MH, Van der Heyden AA, Aarden LA. Interleukin-6 in serum and urine of renal transplant recipients. Clin Exp Immunol 1988; 71: 314-9.

Vernon WB, DeLong ER, Briner WH, Bollinger RR. Renal transplant monitoring with serum β -2-microglobulin and creatinine. Transplant Proc 1985; 17: 646-51.

Vogel W, Tilg H, Herold O, Dietze O, Aulitzky WE, Margreiter R, Huber C. Cytokine production after liver transplantation. Transplant Proc 1990; 22: 1854.

Von Willebrand E. OKT4/8 ratio in the blood and in the graft during episodes of human renal allograft rejection. Cell Immunol 1983;77: 196-201.

Waha A, Watzka M, Koch A, Pietsch T, Przkora R, Peters N, Wiestler OD, von Deimling A. A rapid and sensitive protocol for competitive reverse transcriptase PCR analysis of cellular genes. Brain Pathol 1998; 8: 13-8.

Walle A, Suthanthiran M, Taufield PA, Yankiver B, Rubin AL, Stenzel KH, Darzynkiewicz Z. Multiparameter analysis of cell activation in renal allograft recipients. Transplant Proc 1984; 16: 1586-7.

Wan S, LeClerc J-L, Vincent J-L. Cytokine responses to cardiopulmonary bypass: lessons learned from cardiac transplantation. Ann Thorac Surg 1997; 63: 269-76.

Weimer R, Zipperle S, Daniel V, Carl S, Staehler G, Opelz G. Superior 3-year kidney graft function in patients with impaired pretransplant Th2 responses. *Transpl Int* 1998; 11 Suppl 1: S350-6.

Weisinger G, Gavish M, Mazurika C, Zinder O. Transcription of β -actin, cyclophilin and GAPDH genes: tissue- and treatment-specificity. *Biochim Biophys Acta* 1999; 1446: 225-32.

Weston SD, Lycett AE, Edwards C, Morgan JDT, Horsburgh T, Veitch PS, Bell PRF. Lack of correlation of soluble E-selectin level with renal transplant rejection. *Transpl Immunol* 1995; 3: 50-4.

Wijngaard PLJ, van der Meulen A, Schuurman H-J, Gmelig Meyling FHJ, Heyn A, Borleffs JCC, Jambroes G. Cytoimmunologic monitoring for the diagnosis of acute rejection after heart transplantation. *Transplant Proc* 1989; 21: 2521-22.

Wiley JC, Crawford EL, Jackson CM, Weaver DA, Hoban JC, Khuder SA, DeMuth JP. Expression measurement of many genes simultaneously by quantitative RT-PCR using standardized mixtures of competitive templates. *Am J Respir Cell Mol Biol* 1998; 19: 3-5.

Wilson AG, Symons JA, McDowell TL, McDevitt HO, Duff GW. Effects of a polymorphism in the human tumor necrosis factor α promoter on transcription activation. *Proc Natl Acad Sci USA* 1997; 94: 3195-9.

Woodward JE, Bayer AL, Baliga P. Enhanced allograft survival via simultaneous blockade of transferrin receptor and interleukin-2 receptor. *Transplantation* 1999; 68: 1369-76.

Wu CJ, Lovett M, Wong-Lee J, Moeller MF, Kikaura M, Goralski TJ, Billingham ME, Starnes VA, Clayberger C. Cytokine gene expression in rejection cardiac allografts. *Transplantation* 1992; 54:326-32.

Xu GP, Sharma VK, LI B, Bologa R, Li Y, Mouradian J, Wang J, Serur D, Rao V, Stenzel KH et al. Intragraft expression of IL-10 Messenger RNA: a novel correlate of renal allograft rejection. *Kid Int* 1995; 48: 1504-7.

Yoshimura N, Oka T, Kahan BD. Sequential determinations of serum interleukin 6 levels as an immunodiagnostic tool to differentiate rejection from nephrotoxicity in renal allograft recipients. *Transplantation* 1991; 51: 172-6.

Young CJ, Senkbeil R, Hudson S, Thomas J. Cytokine polymorphism and renal allograft rejection in the mycophenolate mofetil era. *Transplantation* 2000; 69 Suppl: S168.

Zhai Y, Ghobrial RM, Busuttill RW, Kupiec-Weglinski JW. Th1 and Th2 cytokines in organ transplantation: Paradigm lost? *Crit Rev Immunol* 1999; 19: 155-72.