

RESEARCH ARTICLE

# Amyloid beta, tau, synaptic, neurodegeneration, and glial biomarkers in the preclinical stage of the Alzheimer's continuum

Marta Milà-Alomà<sup>1,2,3,4</sup> | Gemma Salvadó<sup>1,2</sup> | Juan Domingo Gispert<sup>1,2,3,5</sup> |  
Natalia Vilor-Tejedor<sup>1,3,6,7</sup> | Oriol Grau-Rivera<sup>1,2,4,8</sup> | Aleix Sala-Vila<sup>1,2</sup> |  
Gonzalo Sánchez-Benavides<sup>1,2,4</sup> | Eider M. Arenaza-Urquijo<sup>1,2,4</sup> | Marta Crous-Bou<sup>1,2,4,9</sup> |  
José Maria González-de-Echávarri<sup>1,2</sup> | Carolina Minguillon<sup>1,2,4</sup> | Karine Fauria<sup>1,4</sup> |  
Maryline Simon<sup>10</sup> | Gwendlyn Kollmorgen<sup>11</sup> | Henrik Zetterberg<sup>12,13,14,15</sup> |  
Kaj Blennow<sup>12,13</sup> | Marc Suárez-Calvet<sup>1,2,4,8</sup> | José Luis Molinuevo<sup>1,2,3,4</sup> | for the ALFA  
study<sup>†</sup>

<sup>1</sup> Barcelonaβeta Brain Research Center (BBRC), Pasqual Maragall Foundation, Barcelona, Spain

<sup>2</sup> IMIM (Hospital del Mar Medical Research Institute), Barcelona, Spain

<sup>3</sup> Universitat Pompeu Fabra, Barcelona, Spain

<sup>4</sup> Centro de Investigación Biomédica en Red de Fragilidad y Envejecimiento Saludable (CIBERFES), Madrid, Spain

<sup>5</sup> Centro de Investigación Biomédica en Red Bioingeniería, Biomateriales y Nanomedicina, Madrid, Spain

<sup>6</sup> Centre for Genomic Regulation (CRG), The Barcelona Institute for Science and Technology, Barcelona, Spain

<sup>7</sup> Department of Clinical Genetics, ERASMUS MC, Rotterdam, the Netherlands

<sup>8</sup> Servei de Neurologia, Hospital del Mar, Barcelona, Spain

<sup>9</sup> Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts, USA

<sup>10</sup> Roche Diagnostics International Ltd, Rotkreuz, Switzerland

<sup>11</sup> Roche Diagnostics GmbH, Penzberg, Germany

<sup>12</sup> Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, University of Gothenburg, Mölndal, Sweden

<sup>13</sup> Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden

<sup>14</sup> Department of Neurodegenerative Disease, UCL Institute of Neurology, Queen Square, London, United Kingdom

<sup>15</sup> UK Dementia Research Institute at UCL, London, United Kingdom

## Correspondence

José Luis Molinuevo and Marc Suárez-Calvet,  
Alzheimer Prevention Program -Barcelonaβeta  
Brain Research Center, Wellington 30, 08005,  
Barcelona, Spain.

E-mail: [jlmolinuevo@barcelonabeta.org](mailto:jlmolinuevo@barcelonabeta.org)  
(J.L.M.), [msuarez@barcelonabeta.org](mailto:msuarez@barcelonabeta.org)  
(M.S.-C.)

<sup>†</sup>The complete list of collaborators of the ALFA  
Study can be found in the Acknowledgments  
section.

## Abstract

**Introduction:** The biological pathways involved in the preclinical stage of the Alzheimer's *continuum* are not well understood.

**Methods:** We used NeuroToolKit and Elecsys® immunoassays to measure cerebrospinal fluid (CSF) amyloid-β (Aβ)42, Aβ40, phosphorylated tau (p-tau), total tau (t-tau), neurofilament light (NfL), neurogranin, sTREM2, YKL40, GFAP, IL6, S100, and α-synuclein in cognitively unimpaired participants of the ALFA+ study, many within the Alzheimer's *continuum*.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2020 The Authors. *Alzheimer's & Dementia* published by Wiley Periodicals, Inc. on behalf of Alzheimer's Association

**Funding information**

"la Caixa" Foundation, Grant/Award Number: ID 100010434; Alzheimer's Association and an international anonymous charity foundation

**Results:** CSF t-tau, p-tau, and neurogranin increase throughout aging only in A $\beta$ -positive individuals, whereas NfL and glial biomarkers increase with aging regardless of A $\beta$  status. We modelled biomarker changes as a function of CSF A $\beta$ 42/40, p-tau and p-tau/A $\beta$ 42 as proxies of disease progression. The first change observed in the Alzheimer's *continuum* was a decrease in the CSF A $\beta$ 42/40 ratio. This is followed by a steep increase in CSF p-tau; t-tau; neurogranin; and, to a lesser extent, in NfL and glial biomarkers.

**Discussion:** Multiple biological pathways are altered and could be targeted very early in the Alzheimer's *continuum*.

**KEYWORDS**

Alzheimer's disease, biomarker, neurodegeneration, neuroinflammation, preclinical

**1 | BACKGROUND**

The natural history of Alzheimer's disease (AD) comprises a long asymptomatic or preclinical stage characterized by pathophysiological changes that start decades before symptoms arise.<sup>1-3</sup> In the new 2018 research framework, AD is defined based on biomarker evidence of amyloid- $\beta$  (A $\beta$ ) and tau pathology, while clinical manifestations are used for grading severity.<sup>4</sup> According to this framework, the term "Alzheimer's disease" is applied whenever there is evidence of A $\beta$  and tau pathology, regardless of the clinical manifestations. When there is evidence of A $\beta$  pathology but not tau, the term "Alzheimer's pathologic change" is used. Together, individuals with either "Alzheimer's pathologic change" or "Alzheimer's disease" belong to the so-called "Alzheimer's *continuum*."

AD cerebrospinal fluid (CSF) core biomarkers allow an accurate diagnostic and early identification of AD pathology.<sup>5</sup> AD CSF core biomarkers comprise A $\beta$ 42 and the A $\beta$ 42/40 ratio, phosphorylated tau (p-tau), and total tau (t-tau), which reflect A $\beta$  pathology, tau pathology, and neurodegeneration, respectively. However, multiple additional pathophysiological processes occur in these early stages of the Alzheimer's *continuum* such as neuronal and axonal damage,<sup>6-9</sup> synaptic dysfunction,<sup>10-12</sup> neuroinflammation and glial response,<sup>13-15</sup> and  $\alpha$ -synuclein or TDP-43 co-pathology.<sup>16-18</sup> In this context, the development of drugs targeting these processes may potentially modify the evolution of the disease.

The pathophysiological events that occur in this early stage remain to be fully elucidated. This can be explained by several reasons. (1) It is particularly challenging to recruit individuals in the earliest extreme of the Alzheimer's *continuum* (A $\beta$ -positive but still tau-negative, ie, preclinical Alzheimer's pathologic change). (2) Most cohorts include elders but not middle-age adults, when AD pathology most likely starts. (3) Most studies include a single or very low number of biomarkers and therefore it is difficult to assess the relationships between them. Recently, a very interesting study in the BioFINDER cohort modelled the changes in CSF and plasma biomarkers in cognitively unimpaired, subjective cognitive decline, and mild cognitively impaired

individuals.<sup>19</sup> They proposed a model with a sequence of events starting with changes in A $\beta$ ; followed by tau biomarkers; and, only after A $\beta$  positron emission tomography (PET) became abnormal, changes in neuronal injury, and synaptic and glial biomarkers.

Herein, we focused in cognitively unimpaired individuals and, very particularly, at the earliest stage of the Alzheimer's *continuum*. The main aim of our study is to define the pathophysiological events that occur in the preclinical stage of the *continuum*. We addressed the challenges described above by studying the well-characterized ALFA+ cohort, which includes cognitively unimpaired individuals, mainly in their middle age, and with a high prevalence of individuals that are A $\beta$  positive but still tau negative.<sup>20</sup> Moreover, we measured several CSF biomarkers that mark the main pathogenic events described in AD: A $\beta$  pathology (A $\beta$ 42, A $\beta$ 42/40 ratio), tau pathophysiology (p-tau), neurodegeneration (t-tau), axonal damage (neurofilament light [NfL]), synaptic dysfunction (neurogranin), microglial (sTREM2) and astroglial-related response (GFAP, YKL40, S100), other neuroinflammatory biomarkers (interleukin 6 [IL6]), and  $\alpha$ -synuclein. We investigated how these CSF biomarkers change with age, sex, and A $\beta$  pathology. Importantly, we modelled the sequence of biomarker changes during the preclinical stage of the Alzheimer's *continuum* to provide a model of the main pathophysiological changes in the earliest stage of the disease *continuum*.

**2 | METHODS****2.1 | ALFA participants and study design**

The ALFA+ cohort is a nested longitudinal study of the ALFA (for Alzheimer's and Families) study.<sup>20</sup> The ALFA cohort was established as a research platform to characterize preclinical AD in 2743 cognitively unimpaired individuals, aged between 45 and 75 years old, and enriched for family history of AD (excluding autosomal-dominant AD). In the nested ALFA+ study, participants are longitudinally followed up and undergo a more comprehensive evaluation. CSF samples in

these participants were obtained by lumbar puncture following standard procedures (see supporting information). Herein, we included the first consecutive 381 participants of ALFA+.

## 2.2 | CSF biomarker measurements

CSF t-tau and p-tau were measured using the electrochemiluminescence immunoassays Elecsys® Total-tau CSF and phosphor-tau(181P) CSF on a fully automated cobas e601 instrument (Roche Diagnostics International Ltd.). The rest of the biomarkers were measured with the prototype NeuroToolKit (Roche Diagnostics International Ltd.) on a cobas e411 or e601 instrument (supporting information). All measurements were performed at the Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden.

## 2.3 | CSF amyloid- $\beta$ and p-tau cutoffs derivation

A $\beta$  pathology positivity (A+) and tau pathology positivity (T+) were defined by CSF A $\beta$ 42/40 ratio and CSF p-tau, respectively.<sup>21</sup> We derived the cutoffs for each of these biomarkers using a two-Gaussian mixture modelling (GMM; supporting information). The cutoff was defined as the mean plus two standard deviations of the non-pathologic Gaussian distribution (ie, the Gaussian with the higher mean for CSF A $\beta$ 42/40 ratio and the Gaussian with the lower mean for p-tau and p-tau/A $\beta$ 42 ratio). The resulting cutoffs were 0.071 for the A $\beta$ 42/40 ratio, 24 pg/mL for p-tau, and 0.013 for the p-tau/A $\beta$ 42 (Figure S1 in supporting information).

## 2.4 | Statistical analyses

For each of the CSF biomarkers, we excluded the extreme values defined as either those that fell outside of three times the interquartile range below the first quartile (Q1) or those above the third quartile (Q3). In the main text, all the analyses were performed excluding extreme values, but including them rendered similar results (see Tables S2–S4 in supporting information). We tested for normality of the distribution for each biomarker using the Kolmogorov-Smirnov test and visual inspection of histograms. CSF A $\beta$ 42, p-tau, t-tau, NfL, neurogranin, YKL-40, GFAP, IL6, S100,  $\alpha$ -synuclein, and the p-tau/A $\beta$ 42 ratio did not follow a normal distribution and were thus log<sub>10</sub>-transformed. CSF A $\beta$ 40, A $\beta$ 42/40 ratio, and sTREM2 followed a normal distribution and were not transformed.

We conducted a one-way analysis of variance (ANOVA) to test statistically significant differences on age and education among AT groups. The mean levels of CSF biomarkers among AT (A $\beta$ /tau pathology) groups were assessed by a one-way analysis of covariance (ANCOVA) adjusting for age and sex. Cognitive performance (Mini-Mental State Examination [MMSE] and Free and Cued Selective Reminding Test [FCSRT]) was assessed by an ANCOVA adjusting for age, sex, and education. These comparisons were followed by Tukey corrected *post*

### HIGHLIGHTS

- Multiple cerebrospinal fluid (CSF) biomarkers are altered very early in the Alzheimer's *continuum*.
- CSF A $\beta$ 42/40 ratio changes first in the preclinical stage of the Alzheimer's *continuum*.
- CSF p-tau, t-tau, and neurogranin increase after the decrease in CSF A $\beta$ 42/40 ratio.
- CSF p-tau, t-tau, and neurogranin change specifically in A $\beta$ -positive individuals.
- CSF neurofilament light (NfL) and glial biomarkers increase with age regardless of A $\beta$  status.

### RESEARCH IN CONTEXT

1. Systematic review: The authors reviewed the literature using traditional sources. Publications describing the natural history of Alzheimer's disease (AD) and its associated cerebrospinal fluid amyloid- $\beta$ , tau, synaptic, neurodegeneration, and glial biomarkers are cited throughout the manuscript.
2. Interpretation: Our findings show that multiple biological pathways are altered in the earliest stages of the preclinical Alzheimer's *continuum* and thus could be targeted in asymptomatic individuals.
3. Future directions: This article proposes a framework for the generation of new hypotheses and the conduct of further studies. Some of these include, for example, the gathering of longitudinal data to assess biomarker profile progression, the addition of biomarkers related to other pathophysiological processes as well as neuroimaging biomarkers.

*hoc* pairwise comparisons. Differences in the frequencies of sex and apolipoprotein E (APOE)- $\epsilon$ 4 categories were assessed by the Pearson's  $\chi^2$  test. Correlations between CSF biomarkers were tested by partial correlations adjusted by age.

To study the association of each CSF biomarker with demographic characteristics and APOE- $\epsilon$ 4 status, we computed a linear regression model with age, sex, and APOE- $\epsilon$ 4 status as predictor variables. We conducted these analyses stratifying by A $\beta$  status and, additionally, including in the linear regression an "A $\beta$ 42/40 ratio x age" interaction term.

We plotted CSF biomarkers levels as a function of CSF A $\beta$ 42/40, p-tau, and p-tau/A $\beta$ 42. We corrected each of the CSF biomarker values by age and sex and computed the mean and standard deviation of each biomarker in the A–T– group (reference group). We next

converted CSF biomarker values to z-scores by subtracting the mean and dividing by the standard deviation of the reference group. The relationship between each CSF biomarker and the proxies of disease progression (ie, CSF A $\beta$ 42/40, p-tau, and p-tau/A $\beta$ 42) were modelled using a robust local weighted regression method (rlowess; "smooth" function in Matlab and a span of 300) and we plotted the resulting model.<sup>22,23</sup> Moreover, we calculated the linear regression slopes for each of the biomarkers in each of the negative or positive groups, using the previously described cutoffs. We computed the *P*-values testing the null hypothesis of whether the regression slope being equal to zero. In addition, the probability of the two regression slopes being equal was also calculated.

For all the analyses, we applied a false discovery rate (FDR) multiple comparison correction following the Benjamini-Hochberg procedure.<sup>24</sup> All tests were two-tailed, with a significance level of  $\alpha = 0.05$ . Statistical analyses were performed in SPSS IBM 20.0 and R software (<http://www.r-project.org/>). Figures were built using R and Matlab (v2018b).

### 3 | RESULTS

#### 3.1 | Participants' characteristics

Table 1 summarizes the demographic characteristics and CSF biomarkers measurements of ALFA+ study participants. Of note, 13 (3.4%) participants fell into the A-T+ group (non-AD pathologic change). Because our aim was to study the Alzheimer's *continuum*, we excluded these 13 participants from all analyses, and they are only depicted in Table 1 and Figure 1 for descriptive purposes. AT groups differed in years of age and education and prevalence of APOE- $\epsilon$ 4 status, but not in sex distribution or MMSE and FCSRT cognitive scores (Table 1).

#### 3.2 | CSF biomarkers and AT groups. Correlations between CSF biomarkers

We observed significant differences in the mean values of CSF NfL, neurogranin, sTREM2, YKL40, GFAP, S100, and  $\alpha$ -synuclein between AT groups (Figure 1). Specifically, CSF NfL, neurogranin, sTREM2, YKL-40, GFAP, and  $\alpha$ -synuclein were significantly higher in the A+T+ group compared to the A-T- and the A+T- groups (Table 1 and Figure 1). Adding years of education in the analyses as a covariate did not modify the results. Of note, none of the studied CSF biomarkers was increased in the A+T- group compared to the A-T- group.

We also tested the correlations between the CSF biomarkers. In a partial correlation adjusting by age, all CSF tau-related, synaptic dysfunction, neuronal injury, and glial markers significantly and positively correlate with each other. There was a negative correlation between CSF A $\beta$ 42/40 and CSF p-tau, t-tau, NfL, neurogranin, GFAP, and S100, but not with the rest of the biomarkers (Figure S2 in supporting information).

#### 3.3 | Associations of CSF biomarkers with age, sex, and APOE- $\epsilon$ 4 status

In the whole sample, CSF A $\beta$ 42, A $\beta$ 42/40, p-tau, t-tau, NfL, neurogranin, sTREM2, YKL-40, GFAP, and S100 were significantly associated with age (Table 2). We stratified the sample into those participants with normal AD biomarkers (A-T-; *n* = 237) and those within the Alzheimer's *continuum* biomarkers group (A+T\*; *n* = 131). In the A-T- group, there was a significant association with a positive direction between age and CSF NfL, YKL-40, and GFAP (Figure 2). In contrast, in the A+T\* group, there was a significant association with a positive direction between age and CSF p-tau, t-tau, NfL, neurogranin, sTREM2, YKL-40, GFAP, and  $\alpha$ -synuclein (Figure 2). Importantly, we found a significant interaction between age and CSF A $\beta$ 42/40 only in the models with CSF p-tau, t-tau, and neurogranin as outcomes (Figure 2).

Interestingly, there were sex differences in several CSF biomarkers, adjusting by the effect of age and APOE- $\epsilon$ 4 status (Table 2). CSF NfL was higher in men, while CSF neurogranin was higher in women (Table 2). Minor (< 10%) but significant differences were observed in CSF GFAP and IL6 (higher in men) and CSF A $\beta$ 40 and YKL-40 (higher in women). Including the CSF A $\beta$ 42/40 ratio as a covariable did not change the results, indicating that the observed differences in CSF biomarkers between sexes are not driven by A $\beta$  pathology. In the studied sample, men had a higher education compared to women (Table S1 in supporting information). Adding education as a covariable in the aforementioned analyses only changed the observed sex differences to non-significant for A $\beta$ 40 (*P* = .075). Unlike age and sex, APOE- $\epsilon$ 4 status was only significantly associated with CSF A $\beta$ 42 and A $\beta$ 42/40 ratio (Table 2).

#### 3.4 | Pathophysiological model of changes in the preclinical Alzheimer's *continuum*

Finally, we modelled the trajectories of the standardized (z-scores) CSF biomarkers in the preclinical Alzheimer's *continuum* applying a robust local weighted regression method. Cognizant that this is a cross-sectional analysis and to understand the changes against A $\beta$  and tau, we anchored the model to: (1) A $\beta$ 42/40 ratio, (2) p-tau, and (3) p-tau/A $\beta$ 42 ratio, as proxies of disease progression. Figure 3 shows the resulting plots. Moreover, for each CSF biomarker and each model, we computed the slopes of a given CSF biomarker before and after the cutoff for the CSF A $\beta$ 42/40 ratio, p-tau, or p-tau/A $\beta$ 42 ratio, respectively, and tested whether these slopes were statistically significantly different.

Anchoring to CSF A $\beta$ 42/40 as a proxy of disease progression, we observed that CSF p-tau, t-tau, and neurogranin start to significantly increase as soon as the CSF A $\beta$ 42/40 ratio becomes positive and continue to increase across the preclinical Alzheimer's *continuum*, eventually reaching the highest z-scores of all CSF biomarkers (3 z-scores for CSF p-tau and t-tau and 2.5 z-scores for CSF neurogranin, compared to their basal levels, Figure 3A). In the CSF

**TABLE 1** Participants' characteristics and CSF biomarkers by AT group

	Total (n = 381)	A-T- (n = 237, 62.2%)	A+T- (n = 100, 26.2%)	A+T+ (n = 31, 8.1%)	A-T+ (n = 13, 3.4%)	P-Value
Age, years	61.2 (4.68)	60.6 (4.44)	61.9 (5.01)	63.8 (4.41) <sup>†</sup>	61.2 (4.51)	.0004 <sup>a</sup>
Female, n (%)	232 (60.9)	146 (61.6)	56 (56.0)	21 (67.7)	9 (69.2)	.56
Education, years	13.4 (3.51)	13.5 (3.46)	13.7 (3.50)	11.9 (3.52) <sup>†</sup>	14.8 (3.59)	.044 <sup>a</sup>
APOE-ε4 carriers, n (%)	201 (52.8)	97 (40.9)	81 (81.0) <sup>‡</sup>	18 (58.1)	5 (38.5)	<.0001 <sup>a</sup>
MMSE	29.1 (0.95)	29.1 (0.92)	29.2 (0.93)	28.8 (1.13)	29.2 (1.17)	.32
FCSRT	15.2 (1.15)	15.2 (1.16)	15.3 (1.11)	15.0 (1.37)	15.7 (0.63)	.45
<b>CSF biomarkers</b>						
Aβ42 (pg/mL)	1302 (564)	1469 (514)	858 (277) <sup>‡</sup>	1016 (370) <sup>‡</sup>	2454 (383)	<.0001 <sup>a</sup>
Aβ42/40	0.075 (0.020)	0.086 (0.009)	0.054 (0.010) <sup>‡</sup>	0.044 (0.012) <sup>‡,§</sup>	0.097 (0.019)	<.0001 <sup>a</sup>
Aβ40 (ng/mL)	17.4 (5.03)	16.8 (4.76)	15.9 (3.63)	22.9 (3.59) <sup>‡,§</sup>	27.0 (3.23)	<.0001 <sup>a</sup>
p-tau (pg/mL)	15.9 (6.28)	13.8 (4.23)	15.6 (4.18) <sup>†</sup>	29.8 (4.86) <sup>‡,§</sup>	27.6 (4.58)	<.0001 <sup>a</sup>
t-tau (pg/mL)	196 (68.1)	175 (48.6)	191 (44.9) <sup>‡</sup>	338 (52.8) <sup>‡,§</sup>	317 (51.6)	<.0001 <sup>a</sup>
p-tau/Aβ42	0.014 (0.008)	0.010 (0.002)	0.020 (0.007) <sup>‡</sup>	0.032 (0.011) <sup>‡,§</sup>	0.011 (0.002)	<.0001 <sup>a</sup>
NfL (pg/mL)	81.5 (26.3)	75.7 (23.4)	82.1 (22.6)	115 (33.8) <sup>‡,§</sup>	103 (28.1)	<.0001 <sup>a</sup>
Neurogranin (pg/mL)	796 (326)	712 (250)	743 (224)	1366 (293) <sup>‡,§</sup>	1423 (328)	<.0001 <sup>a</sup>
sTREM2 (ng/mL)	7.91 (7.57)	7.65 (1.95)	7.50 (2.06)	9.85 (2.69) <sup>‡,§</sup>	11.24 (2.84)	<.0001 <sup>a</sup>
YKL-40 (ng/mL)	147 (52.9)	138 (44.0)	142 (46.4)	212 (64.9) <sup>‡,§</sup>	206 (74.3)	<.0001 <sup>a</sup>
GFAP (ng/mL)	7.54 (2.29)	7.20 (2.10)	7.44 (2.25)	10.3 (2.17) <sup>‡,§</sup>	7.93 (2.01)	<.0001 <sup>a</sup>
IL6 (pg/mL)	3.89 (1.46)	3.84 (1.34)	3.94 (1.65)	4.17 (1.76)	3.62 (1.31)	.54
S100 (ng/mL)	1.02 (0.23)	0.99 (0.20)	1.03 (0.24)	1.14 (0.27) <sup>‡</sup>	1.13 (0.33)	.010 <sup>a</sup>
α-synuclein (pg/mL)	198 (80.8)	187 (79.2)	181 (53.9)	298 (66.9) <sup>‡,§</sup>	322 (57.9)	<.0001 <sup>a</sup>

Notes: Data are expressed as mean (M) and standard deviation (SD) or percentage (%), as appropriate. One-way ANOVA followed by Tukey corrected *post hoc* comparisons was used to compare age and education and Pearson's  $\chi^2$  test to compare sex and APOE-ε4 status between AT groups. MMSE and FCSRT scores were compared with an ANCOVA adjusted by age, sex, and education. CSF biomarkers were compared with an ANCOVA adjusted by age and sex followed by Tukey corrected *post hoc* comparisons. The *P*-values indicated in the last column refer to the AT group effect. We did not include the A-T+ group in the analyses, but this group is included in the table for the sake of completeness. *P*-values are corrected for multiple comparisons using FDR approach.

Abbreviations: Aβ40, amyloid-β 40; Aβ42, amyloid-β 42; ANCOVA, analysis of covariance; ANOVA, analysis of variance; APOE, apolipoprotein E; CSF, cerebrospinal fluid; FCSRT, Free and Cued Selective Reminding Test (total recall); GFAP, glial fibrillary acidic protein; IL6, interleukin 6; MMSE, Mini-Mental State Examination; NfL, neurofilament light; p-tau, phosphorylated tau; sTREM2, soluble triggering receptor expressed on myeloid cells 2 (TREM2); t-tau, total tau.

<sup>a</sup>Significant values.

Pairwise *post hoc* comparisons:

\**P* < .001 versus A-T-.

<sup>†</sup>*P* < .05 versus A+T-.

<sup>‡</sup>*P* < .0001 versus A-T-.

<sup>§</sup>*P* < .0001 versus A+T-.


<sup>||</sup>*P* < .05 versus A-T-.

<sup>¶</sup>*P* < .01 versus A-T-.

Aβ42/40-positive participants, the higher absolute slopes are those of CSF p-tau and t-tau, followed by CSF neurogranin (Table 3). These slopes in the CSF Aβ42/40-positive participants were significantly different from the slopes of the CSF Aβ42/40-negative participants (Figure 3A; Table 3). The rest of the CSF biomarkers, except for CSF IL6 and S100, also increased in the CSF Aβ42/40-positive participants, but that increase was considerably less pronounced (as shown by lower absolute slopes). The slopes of CSF sTREM2 and YKL40 in Aβ-positive participants were also significantly different from those that were Aβ-negative. Remarkably, CSF NfL significantly increased as a function of CSF Aβ42/40 in Aβ-positive individuals, although the slope

was not significantly different than that of Aβ-negative individuals (Table 3).

When we used CSF p-tau as a proxy of disease progression, we observed that CSF Aβ42/40 dramatically decreases before CSF p-tau becomes positive. Importantly, CSF Aβ42/40 values already reach a decrease of four z-scores from the basal levels at the point when CSF p-tau becomes positive, and they plateau after (Figure 3B, Table 3). This result strongly suggests that changes in soluble Aβ precede those of tau pathophysiology. Remarkably, CSF neurogranin also started to increase before the CSF p-tau cutoff, reaching almost two z-scores from the basal levels at the point when CSF p-tau becomes positive,



**FIGURE 1** Comparison of cerebrospinal fluid (CSF) biomarkers between AT groups. Dot and box plots depicting the levels of each CSF biomarker in each of the AT groups. The box plots depict the median (horizontal bar), interquartile range (IQR, hinges), and  $1.5 \times$  IQR (whiskers). Because our goal is to assess CSF biomarkers in the Alzheimer's *continuum*, we did not include the A-T+ group (ie, non-AD pathologic change) in the analyses, but this group is included in the figure for the sake of completeness. P-values were assessed by a one-way analysis of covariance adjusted by age and sex, followed by Tukey corrected pair-wise *post hoc* comparisons

**TABLE 2** Effect of age, sex, and APOE-ε4 status on CSF biomarkers

Biomarker	Age		Sex			APOE-ε4		
	β (SE)	P-value	Male, M (SD)	Female, M (SD)	P-value	Non-carriers, M (SD)	Carriers, M (SD)	P-value
Aβ42	-0.12 (0.050)	.023 <sup>a</sup>	1203 (529)	1306 (526)	.26	1437 (563)	1114 (446)	<.0001 <sup>a</sup>
Aβ42/40	-0.24 (0.047)	<.0001 <sup>a</sup>	0.073 (0.019)	0.074 (0.019)	.47	0.081 (0.018)	0.067 (0.019)	<.0001 <sup>a</sup>
Aβ40	0.090 (0.052)	.092	16.3 (4.82)	17.6 (4.63)	.043 <sup>a</sup>	17.6 (5.03)	16.6 (4.42)	.33
p-tau	0.24 (0.051)	<.0001 <sup>a</sup>	15.1 (5.79)	15.7 (6.02)	.20	15.4 (6.26)	15.5 (5.64)	.81
t-tau	0.23 (0.052)	<.0001 <sup>a</sup>	188 (65.1)	195 (64.4)	.20	193 (69.5)	191 (60.2)	.93
NfL	0.41 (0.046)	<.0001 <sup>a</sup>	88.7 (24.5)	75.5 (25.4)	<.0001 <sup>a</sup>	79.5 (25.8)	81.7 (25.9)	.81
Neurogranin	0.15 (0.052)	.008 <sup>a</sup>	723 (287)	808 (310)	.013 <sup>a</sup>	782 (314)	767 (294)	.94
sTREM2	0.21 (0.051)	<.0001 <sup>a</sup>	7.81 (2.22)	7.78 (2.09)	.94	8.03 (2.17)	7.59 (2.09)	.27
YKL-40	0.41 (0.048)	<.0001 <sup>a</sup>	140 (50.3)	148 (51.1)	.043 <sup>a</sup>	147 (50.9)	144 (51.0)	.94
GFAP	0.33 (0.049)	<.0001 <sup>a</sup>	7.93 (2.29)	7.27 (2.28)	.043 <sup>a</sup>	7.50 (2.43)	7.55 (2.19)	.91
IL6	-0.068 (0.053)	.20	4.09 (1.51)	3.77 (1.42)	.043 <sup>a</sup>	3.94 (1.52)	3.86 (1.42)	.76
S100	0.12 (0.052)	.029 <sup>a</sup>	1.04 (0.24)	1.00 (0.21)	.20	1.00 (0.22)	1.03 (0.23)	.76
α-synuclein	0.092 (0.053)	.092	194 (83.3)	193 (74.2)	.86	196 (81.4)	192 (60.2)	.91

Notes: Each CSF biomarker was assessed by a linear model with age, sex, and APOE-ε4 status as main effects. The standardized regression coefficients (β) and standard errors (SE) are depicted for age and mean (M) and standard deviation (SD) for sex and APOE-ε4. P-values are corrected for multiple comparisons using FDR approach.

Abbreviations: Aβ40, amyloid-β 40; Aβ42, amyloid-β 42; APOE, apolipoprotein E; CSF, cerebrospinal fluid; GFAP, glial fibrillary acidic protein; IL6, interleukin 6; NfL, neurofilament light; p-tau, phosphorylated tau; sTREM2, soluble TREM2; t-tau, total tau.

<sup>a</sup>Significant values.

and plateaus in individuals that are already tau positive (Figure 3B, Table 3). The rest of CSF biomarkers (except CSF IL6) also increase as a function of CSF p-tau but their increase is less pronounced (< 0.5 z-scores). As expected, CSF t-tau levels parallel those of CSF p-tau. When we used CSF p-tau/Aβ42 as a proxy of disease progression, the results were very similar to those of CSF p-tau. Before CSF p-tau/Aβ42 becomes positive (Figure 3C), CSF Aβ42/40 significantly decreases to later plateau, and eventually reach a decrease of six z-scores. In contrast, the rest of CSF biomarkers do not significantly change before the CSF p-tau/Aβ42 cutoff, but they significantly increase after surpassing this cutoff, except for CSF IL6 and S100 (Table 3). CSF p-tau and t-tau have the steeper slopes and reach >4 z-scores of its basal levels. CSF neurogranin reaches two z-scores from its basal levels and the rest of CSF biomarkers showed a more moderate increase (< 2 z-scores).


## 4 | DISCUSSION

In this cross-sectional study we aimed at determining CSF biomarker changes in the preclinical Alzheimer's *continuum* in a well-characterized cohort of cognitively unimpaired individuals. Our main results are the following. (1) Changes in soluble Aβ occur earlier than in any other biomarker studied. (2) After soluble Aβ biomarkers become positive, there is a steep increase in tau-related (p-tau and t-tau) and synaptic dysfunction (neurogranin) CSF biomarkers and, to a lesser extent, in axonal injury (NfL) and glial (sTREM2, YKL40, GFAP) biomarkers. (3) Tau-related and synaptic dysfunction biomarkers increase across age only in Aβ-positive individuals, whereas axonal damage and glial biomarkers increase during aging in both Aβ-positive

and -negative individuals. Altogether, our results show that the first observable changes in the Alzheimer's *continuum* are those in soluble Aβ, and as soon as individuals become Aβ positive, changes in tau pathology and synaptic dysfunction occur. Changes in CSF NfL and glial biomarkers also occur during disease progression, but these are less pronounced.

We initially observed that all CSF biomarkers (except IL6) increased in the A+T+ group, but not in the A+T- group. However, this approach has the limitation that it simplifies the preclinical stage of the Alzheimer's *continuum* in only two stages, that is A+T- and A+T+. Therefore, we applied two additional approaches in order to define more precisely the CSF biomarker changes in the Alzheimer's *continuum*. We first assessed the changes of CSF biomarkers as a function of age and, second, as a function of Aβ (as defined by the CSF Aβ42/40 ratio) and tau (as defined by CSF p-tau) pathophysiology.

We observed that all biomarkers (except CSF Aβ40, IL6, and α-synuclein) increase with age. Interestingly, for CSF p-tau, t-tau, and neurogranin, this increase is specifically linked to Aβ pathology. When we used the CSF Aβ42/40 ratio as a proxy of disease progression, CSF p-tau, t-tau, and neurogranin are the biomarkers that change more markedly, and before the Aβ-positivity cutoff. This finding is consistent with the notion that CSF p-tau and neurogranin are biomarkers mainly related to AD, and less linked to other forms of neurodegeneration.<sup>6,25,26</sup> In contrast, CSF NfL and glial biomarkers CSF YKL40, GFAP, and sTREM2 (the latter approaching significance) change through aging in both Aβ-positive and -negative individuals. Although this increase is more pronounced in Aβ-positive than in negative individuals (as shown by the β slopes), the fact that there is no interaction between Aβ and age indicates that Aβ does not




**FIGURE 2** Association of cerebrospinal fluid (CSF) biomarkers with age. Scatter plots representing the associations of each of the CSF biomarkers with age in the A-T- (ie, normal AD biomarkers) and the A+T\* (ie, Alzheimer's *continuum*) groups. Each point depicts the value of the CSF biomarker of an individual and the solid lines indicate the regression line for each of the groups. The standardized regression coefficients ( $\beta$ ) and the *P*-values are shown and were computed using a linear model adjusting for age, sex, and apolipoprotein E (APOE)- $\epsilon 4$ . Additionally, we also computed the "A $\beta$ 42/40 x age" interaction term. All *P*-values are corrected for multiple comparisons using the FDR approach

significantly modify the association between age and the specific CSF biomarker. These results are in line with previous studies that show that glial biomarkers increase with aging and also in other neurological diseases besides AD.<sup>27-36</sup> Of note, our sample comprises a higher prevalence of A+T- than A+T+ individuals. We speculate that the increase of CSF NfL and glial markers throughout age would be more pronounced in the latter stages of the preclinical Alzheimer's *continuum* (ie, A+T+) and in early symptomatic stages. When we modeled the changes of CSF biomarkers as a function of the CSF A $\beta$ 42/40 ratio, we observe similar results, that is CSF NfL and glial biomarker changes are less pronounced than those of tau-related and synaptic

biomarkers. It is remarkable that CSF NfL behaves differently from CSF t-tau, despite the fact that both are biomarkers of neurodegeneration. Unlike CSF t-tau, CSF NfL may reflect a degree of age-related neuronal and axonal injury independent from A $\beta$  pathology. Moreover, CSF NfL less marked increase as a function of CSF A $\beta$ 42/40 may reflect that neuronal and axonal injury may be more prominent in later stages of preclinical AD, where A $\beta$  and tau pathology are already present. These findings are also consistent with previous studies showing that CSF t-tau and NfL might provide different information regarding neurodegeneration. While CSF t-tau mainly reflects an A $\beta$ -related change in tau metabolism and secretion that eventually



**FIGURE 3** Cerebrospinal fluid (CSF) biomarker trajectories. The graphs represent the z-scores changes of each CSF biomarker using the mean and the standard deviation of that CSF biomarker in the A-T- group as a reference. The resulting z-scores are shown as a function of CSF A $\beta$ 42/40 (A) p-tau (B) or p-tau/A $\beta$ 42 (C) using a robust local weighted regression method. The solid lines depict the trajectory of each CSF biomarker. The dashed lines depict the cutoff for CSF A $\beta$ 42/40, p-tau, and p-tau/A $\beta$ 42, respectively. The horizontal axis direction of CSF A $\beta$ 42/40 (A) was inverted.



**TABLE 3** Association between each CSF biomarker and A $\beta$ 42/40, p-tau, and p-tau/A $\beta$ 42

Model 1: CSF A $\beta$ 42/40 as proxy of disease progression					
	A $\beta$ 42/40 negative		A $\beta$ 42/40 positive		Slopes difference
	B (SE)	P-value	B (SE)	P-value	P-value*
p-tau	13.8 (8.30)	.46	-66.2 (11.2)	.002 <sup>a</sup>	<.0001 <sup>a</sup>
t-tau	15.0 (8.24)	.24	-52.3 (9.94)	<.0001 <sup>a</sup>	<.0001 <sup>a</sup>
NfL	-0.82 (11.8)	.9	-29.1 (9.24)	.004 <sup>a</sup>	.085
Neurogranin	21.9 (7.91)	.064	-39.2 (8.94)	.0001 <sup>a</sup>	<.0001 <sup>a</sup>
sTREM2	14.5 (11.2)	.46	-20.9 (7.03)	.005 <sup>a</sup>	.015 <sup>a</sup>
YKL-40	9.83 (9.80)	.46	-31.2 (9.09)	.002 <sup>a</sup>	.005 <sup>a</sup>
GFAP	-6.22 (11.7)	.66	-24.7 (8.74)	.007 <sup>a</sup>	.23
IL6	8.95 (11.2)	.53	0.064 (10.7)	.10	.57
S100	-23.0 (11.2)	.21	-2.75 (10.1)	.87	.22
$\alpha$ -synuclein	11.0 (11.1)	.46	-21.7 (6.56)	.003 <sup>a</sup>	.019 <sup>a</sup>
Model 2: CSF p-tau as proxy of disease progression					
	p-tau negative		p-tau positive		Slopes difference
	B (SE)	P-value	B (SE)	P-value	P-value*
A $\beta$ 42	0.12 (0.013)	<.0001 <sup>a</sup>	0.023 (0.041)	.71	.066
A $\beta$ 42/40	-0.24 (0.032)	<.0001 <sup>a</sup>	-0.041 (0.056)	.63	.028 <sup>a</sup>
A $\beta$ 40	0.13 (0.008)	<.0001 <sup>a</sup>	0.076 (0.024)	.022 <sup>a</sup>	.066
t-tau	0.19 (0.006)	<.0001 <sup>a</sup>	0.14 (0.022)	<.0001 <sup>a</sup>	.070
NfL	0.076 (0.016)	<.0001 <sup>a</sup>	0.12 (0.053)	.083	.58
Neurogranin	0.17 (0.007)	<.0001 <sup>a</sup>	0.057 (0.041)	.31	.055
sTREM2	0.12 (0.013)	<.0001 <sup>a</sup>	0.061 (0.022)	.029 <sup>a</sup>	.066
YKL-40	0.11 (0.014)	<.0001 <sup>a</sup>	0.12 (0.061)	.13	.96
GFAP	0.074 (0.015)	<.0001 <sup>a</sup>	-0.012 (0.055)	.90	.19
IL6	-0.014 (0.018)	.43	0.006 (0.071)	.93	.94
S100	0.073 (0.018)	<.0001 <sup>a</sup>	-0.064 (0.061)	.47	.066
$\alpha$ -synuclein	0.085 (0.013)	<.0001 <sup>a</sup>	0.089 (0.029)	.022 <sup>a</sup>	.96
Model 3: CSF p-tau/A $\beta$ 42 as proxy of disease progression					
	p-tau/A $\beta$ 42 negative		p-tau/A $\beta$ 42 positive		Slopes difference
	B (SE)	P-value	B (SE)	P-value	P-value*
A $\beta$ 42	-48.2 (44.1)	.51	-55.7 (9.21)	<.0001 <sup>a</sup>	.87
A $\beta$ 42/40	-387 (59.3)	<.0001 <sup>a</sup>	-152 (13.5)	<.0001 <sup>a</sup>	.001 <sup>a</sup>
A $\beta$ 40	-71.6 (31.0)	.22	32.0 (9.21)	.001 <sup>a</sup>	.033 <sup>a</sup>
p-tau	84.9 (36.2)	.13	109 (11.2)	<.0001 <sup>a</sup>	.75
t-tau	54.4 (35.1)	.40	86.2 (10.8)	<.0001 <sup>a</sup>	.65
NfL	63.0 (54.5)	.51	46.3 (10.9)	.0001 <sup>a</sup>	.83
Neurogranin	12.3 (35.2)	.86	59.8 (10.2)	<.0001 <sup>a</sup>	.58
sTREM2	-31.9 (51.9)	.86	30.9 (8.7)	.0008 <sup>a</sup>	.58
YKL-40	-12.6 (44.9)	.86	43.1 (10.8)	.0002 <sup>a</sup>	.58
GFAP	6.12 (53.6)	.91	36.8 (10.5)	.0008 <sup>a</sup>	.75
IL6	68.6 (51.0)	.47	10.6 (12.7)	.44	.58
S100	24.3 (56.0)	.86	3.28 (11.6)	.78	.83
$\alpha$ -synuclein	-13.6 (52.6)	.86	31.2 (7.83)	.0002 <sup>a</sup>	.65

Notes: For each CSF biomarker we computed the linear regression unstandardized coefficients (B) of the z-scores and standard errors (SE) in each of the negative or positive groups. P-values are corrected for multiple comparisons using FDR approach.

Abbreviations: A $\beta$ 40, amyloid- $\beta$  40; A $\beta$ 42, amyloid- $\beta$  42; CSF, cerebrospinal fluid; GFAP, glial fibrillary acidic protein; IL6, interleukin 6; NfL, neurofilament light; p-tau, phosphorylated tau; sTREM2, soluble TREM2; t-tau, total tau.

<sup>a</sup>Significant values.

\*P-value tests whether the regression slopes of the two groups are equal.

may translate into AD-type neurodegeneration, CSF NfL also reflects neurodegeneration not due to A $\beta$  pathology.<sup>37</sup> The underlying cause of that increase on CSF NfL in A $\beta$ -negative cognitively unimpaired individuals still needs to be clarified, but we speculate that age-related factors such as vascular and/or other neurodegeneration-related factors (eg, co-pathology with Lewy body disease, TDP-43, or hippocampal sclerosis) may at least partially explain those differences. Together, these findings also favor the use of CSF NfL as marker of neurodegeneration ("N") in the amyloid/tau/neurodegeneration (ATN) framework, instead of CSF t-tau. The CSF glial markers parallel the increase of CSF NfL in the model using CSF A $\beta$ 42/40 as a proxy of disease progression. Both CSF NfL and CSF glial biomarkers increase as a function of A $\beta$  pathology (as shown as decreased CSF A $\beta$ 42/40 ratio) in the A $\beta$ -positive but not the A $\beta$ -negative group. This may indicate that A $\beta$  pathology underlies the neuronal injury and glial response in these individuals. Yet, this idea does not exclude that, in other individuals, other mechanisms different from A $\beta$  pathology may also trigger neuronal injury and glial response.

Interestingly, we found sex differences in some of the studied CSF biomarkers. Men showed higher levels of NfL in CSF, which is consistent with previous studies,<sup>38,39</sup> and with the fact that the prevalence of A-T-N+ individuals (ie, neurodegeneration positive, as measured by hippocampal volume or cortical thickness, but A $\beta$ - and tau-negative) is higher in men than in women.<sup>40</sup> An unexpected finding was that CSF neurogranin is overall higher in women than in men. Whether this indicates that women may have a greater susceptibility to synaptic dysfunction remains to be clarified in further studies. The effect of sex, often overlooked, should be further studied to better understand AD pathogenesis and design preventive strategies.

The main limitation of cross-sectional biomarker studies in the sporadic preclinical Alzheimer's *continuum* is the lack of a proxy for the temporal evolution of the disease, such as the concept of estimated years from symptom onset in autosomal-dominant AD. We chose the CSF A $\beta$ 42/40 ratio to understand how biomarkers evolve during the *continuum* because changes in soluble A $\beta$  are those that occur earlier in the Alzheimer's *continuum*, both in autosomal-dominant and sporadic AD, and occur before than A $\beta$  PET becomes positive.<sup>41,42</sup> Moreover, CSF A $\beta$ 42/40 ratio is less affected by pre-analytical factors and inter-individual differences than CSF A $\beta$ 42. Remarkably, the CSF A $\beta$ 42/40 ratio cutoff used here is higher (and thus expected to be more sensitive) than the one commonly used for diagnostic purposes, given that our goal is to sensitively detect very early changes in A $\beta$  pathology. This is the reason why we used a two-Gaussian mixture modelling to determine the cutoff instead of conducting a comparison with a gold-standard (such as A $\beta$  PET), which would have rendered a more specific cutoff for deposited A $\beta$  but less sensitive for early soluble A $\beta$ -related pathophysiological changes. Being aware that using the CSF A $\beta$ 42/40 ratio as a proxy of disease progression may seem that we are assuming a sequence of events in which soluble A $\beta$  comes first, we also explored CSF biomarker changes using CSF p-tau and the p-tau/A $\beta$ 42 ratio as proxies, the latter being highly correlated with A $\beta$  PET load.<sup>43</sup> We similarly computed the CSF p-tau and p-tau/A $\beta$ 42 cutoffs using a two-Gaussian mixture modelling and these resulted lower (thus expected

to be more sensitive) than those usually used in the clinical setting. Even with such sensitive cutoffs, changes in CSF A $\beta$ 42/40 always come first, before CSF p-tau or the p-tau/A $\beta$ 42 become positive or before any other biomarker changes. Using these models, we also observed the initial increase in CSF neurogranin, earlier than changes in neuronal and axonal injury and glial-related biomarkers. Overall, these results are similar to those that have been reported in autosomal-dominant AD in both cross-sectional studies and longitudinal studies, in which initial changes in A $\beta$  are followed by tau-related biomarkers and, afterward, neurodegeneration and glial markers.<sup>1,2,15,44-47</sup> Recently, a very interesting study in the BIOFINDER cohort modelled the changes in several CSF and plasma biomarkers in predementia individuals.<sup>19</sup> The authors used A $\beta$  PET as a proxy of the disease progression, instead of CSF A $\beta$ 42/40, but their results were similar. The main difference with our study is that we observe earlier changes in CSF neurogranin, while they observed an early inflection of CSF NfL. We may argue that changes in soluble A $\beta$  are more linked to early synaptic dysfunction, while A $\beta$  deposition (as measured by A $\beta$  PET) captures a slightly later event that may be more associated to neuronal injury. Moreover, our study includes a high number of cognitively unimpaired individuals that are A $\beta$  positive but still tau negative, which may have allowed us to observe these early changes in CSF neurogranin.

It is worth noting the local regression methods we applied to model the association between CSF biomarker changes as a function of CSF A $\beta$ 42/40, p-tau, or p-tau/A $\beta$ 42. These methods are particularly suited to model non-linear associations without the need to specify any function or fit a model a priori. Still, the smoothing parameter ("span") needs to be specified. To this end, we chose a smoothing parameter that produced curves with a maximum of two inflection (change) points to avoid overfitting. Similar methods have been applied in previous reports displaying the cross-sectional variation of CSF and other biomarkers against proxies of AD progression.<sup>2,41,45</sup>

The main limitations of our study are the following. (1) It is a cross-sectional analysis and longitudinal studies are needed to confirm the results. (2) A $\beta$ - and tau-pathology cutoffs derived herein might not be applied to clinical cohorts because ALFA+ is a very specific cohort aimed at studying preclinical AD and with a high percentage of APOE- $\epsilon$ 4 carriers and A $\beta$ -positive individuals. Moreover, the diagnostic and prognostic value of these cutoffs in clinical population has not been assessed. (3) Although it includes CSF biomarkers related to different pathophysiological processes, we do not include biomarkers related to vascular function or TDP-43 pathology. (4) We did not include neuroimaging biomarkers such as structural magnetic resonance imaging or A $\beta$  PET. (5) We measured total levels of  $\alpha$ -synuclein, which probably does not reflect Lewy body disease or  $\alpha$ -synuclein deposition as phosphorylated or oligomeric forms do,<sup>48-51</sup> but most likely reflects neuronal injury.

In conclusion, our study shows that biomarkers reflecting multiple pathophysiological pathways change very early in the Alzheimer's *continuum*. tau-related and synaptic biomarkers are those that change earlier and more markedly, as soon as there is evidence of incipient A $\beta$  pathology. In order to develop therapeutic strategies targeting this early stage, it is fundamental to understand "which" are the

biological pathways involved and “when” in the long preclinical Alzheimer's *continuum* they are involved. Our results favor the idea of targeting tau and synaptic dysfunction in the earliest stages of the preclinical Alzheimer's *continuum*, as soon as alterations in A $\beta$  occur.

#### ACKNOWLEDGMENTS

This publication is part of the ALFA study (ALzheimer and FAMilies). The authors would like to express their most sincere gratitude to the ALFA project participants and relatives without whom this research would have not been possible. We also thank Tania Menchón, Carolina Herrero, Noemí Carranza, Charlotte Frick, Linnéa Lagerstedt, and Irina Nilsson for technical assistance. Collaborators of the ALFA study are: Annabella Beteta, Raffaele Cacciaglia, Alba Cañas, Carme Deulofeu, Irene Cumpido, Ruth Dominguez, Carles Falcon, Sherezade Fuentes, Laura Hernandez, Gema Huesa, Jordi Huguet, Paula Marne, Tania Menchón, Grégory Operto, Maria Pascual, Albina Polo, Sandra Pradas, Mahnaz Shekari, Anna Soteras, and Marc Vilanova.

The authors thank Roche Diagnostics International Ltd. for providing the kits to measure CSF biomarkers. ELECSYS, COBAS, and COBAS E are registered trademarks of Roche. The Elecsys  $\beta$ -Amyloid (1-42) CSF immunoassay in use is not a commercially available IVD assay. It is an assay that is currently under development and for investigational use only. The measuring range of the assay is 200 (lower technical limit) – 1700 pg/mL (upper technical limit). The performance of the assay beyond the upper technical limit has not been formally established. Therefore, use of values above the upper technical limit, which are provided based on an extrapolation of the calibration curve, is restricted to exploratory research purposes and is excluded for clinical decision making or for the derivation of medical decision points.

#### CONFLICTS OF INTEREST

JLM has served/serves as a consultant or at advisory boards for the following for-profit companies, or has given lectures in symposia sponsored by the following for-profit companies: Roche Diagnostics, Genentech, Novartis, Lundbeck, Oryzon, Biogen, Lilly, Janssen, Green Valley, MSD, Eisai, Alector, BioCross, GE Healthcare, ProMIS Neurosciences. KB has served as a consultant or at advisory boards for Abcam, Axon, Biogen, Lilly, MagQu, Novartis, and Roche Diagnostics, and is a co-founder of Brain Biomarker Solutions in Gothenburg AB, a GU Ventures-based platform company at the University of Gothenburg. HZ has served at scientific advisory boards for Roche Diagnostics, CogRx, Samumed, and Wave, and has given lectures in symposia sponsored by Alzecure and Biogen, and is a co-founder of Brain Biomarker Solutions in Gothenburg AB, a GU Ventures-based platform company at the University of Gothenburg. GK is a full-time employee of Roche Diagnostics GmbH. MS is a full-time employee of Roche Diagnostics International Ltd. The remaining authors declare that they have no conflicts of interest.

#### ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The ALFA+ study (ALFA-FPM-0311) was approved by the Independent Ethics Committee “Parc de Salut Mar,” Barcelona, and registered at Clinicaltrials.gov (Identifier: NCT02485730). All participating sub-

jects and signed the study's informed consent form that had also been approved by the Independent Ethics Committee “Parc de Salut Mar,” Barcelona.

#### AUTHOR CONTRIBUTIONS

Marta Milà-Alomà, Gemma Salvadó, Juan Domingo Gispert, Natalia Vilor-Tejedor, José Maria González-de-Echávarri, Marc Suárez-Calvet, and José Luis Molinuevo analyzed and interpreted the data. Kaj Blennow, Henrik Zetterberg, and Marc Suárez-Calvet analyzed the CSF samples. Maryline Simon and Gwendlyn Kollmorgen developed and provided the NeuroToolKit (Roche). Oriol Grau-Rivera, Aleix Sala-Vila, Gonzalo Sánchez-Benavides, Eider M. Arenaza-Urquijo, Marta Crous-Bou, José Maria González-de-Echávarri, Carolina Minguillon, Karine Fauria, Marc Suárez-Calvet, and José Luis Molinuevo contributed with ALFA+ participants' data. Marta Milà-Alomà, Juan Domingo Gispert, Marc Suárez-Calvet, and José Luis Molinuevo designed the study and wrote the manuscript. All authors critically reviewed and approved the final manuscript.

#### FUNDING

The project leading to these results has received funding from “la Caixa” Foundation (ID 100010434), under agreement LCF/PR/GN17/50300004 and the Alzheimer's Association and an international anonymous charity foundation through the TriBEKa Imaging Platform project (TriBEKa-17-519007). Additional support has been received from the Universities and Research Secretariat, Ministry of Business and Knowledge of the Catalan Government under the grant no. 2017-SGR-892. MSC received funding from the European Union's Horizon 2020 Research and Innovation Program under the Marie Skłodowska-Curie action grant agreement No 752310, and currently receives funding from Instituto de Salud Carlos III (PI19/00155) and from the Spanish Ministry of Science, Innovation and Universities (Juan de la Cierva Programme grant IJC2018-037478-I). JDG is supported by the Spanish Ministry of Science and Innovation (RYC-2013-13054). NVT is supported by the Spanish Ministry of Science, Innovation and Universities—Spanish State Research Agency (FJC2018-038085-I). OGR is supported by the Spanish Ministry of Science, Innovation and Universities (FJCI-2017-33437). ASV is the recipient of an Instituto de Salud Carlos III Miguel Servet II fellowship (CP II 17/00029). EMAU is supported by the Spanish Ministry of Science, Innovation and Universities—Spanish State Research Agency (RYC2018-026053-I). CM was supported by the Spanish Ministry of Economy and Competitiveness (grant n° IEDI-2016-00690). KB holds the Torsten Söderberg Professorship in Medicine at the Royal Swedish Academy of Sciences, and is supported by the Swedish Research Council (#2017-00915); the Swedish Alzheimer Foundation (#AF-742881), Hjärnfonden, Sweden (#FO2017-0243); and a grant (#ALFGBG-715986) from the Swedish state under the agreement between the Swedish government and the County Councils, the ALF-agreement. HZ is a Wallenberg Academy Fellow supported by grants from the Swedish Research Council (#2018-02532), the European Research Council (#681712), and a grant (#ALFGBG-720931) from the Swedish state under the agreement between the Swedish government and the County Councils.

## REFERENCES

- Fagan AM, Xiong C, Jasielec MS, et al. Longitudinal change in CSF biomarkers in autosomal-dominant Alzheimer's disease. *Sci Transl Med*. 2014;6:226ra30-226ra30.
- Bateman RJ, Xiong C, Benzinger TL, et al. Clinical and biomarker changes in dominantly inherited Alzheimer's disease. *N Engl J Med*. 2012;367:795-804.
- Jack CR, Holtzman DM. Biomarker modeling of Alzheimer's disease. *Neuron*. 2013;80:1347-1358.
- Jack CR, Bennett DA, Blennow K, et al. NIA-AA research framework: toward a biological definition of Alzheimer's disease. *Alzheimer's Dement*. 2018;14:535-562.
- Olsson B, Lautner R, Andreasson U, et al. CSF and blood biomarkers for the diagnosis of Alzheimer's disease: a systematic review and meta-analysis. *Lancet Neurol*. 2016;15:673-684.
- Bos I, Vos S, Verhey F, et al. Cerebrospinal fluid biomarkers of neurodegeneration, synaptic integrity, and astroglial activation across the clinical Alzheimer's disease spectrum. *Alzheimer's Dement*. 2019;15:644-654.
- Sjögren M, Blomberg M, Jonsson M, et al. Neurofilament protein in cerebrospinal fluid: a marker of white matter changes. *J Neurosci Res*. 2001;66:510-516.
- Skillback T, Farahmand B, Bartlett JW, et al. CSF neurofilament light differs in neurodegenerative diseases and predicts severity and survival. *Neurology*. 2014;83:1945-1953.
- Price JL, Ko AI, Wade MJ, Tsou SK, McKeel DW, Morris JC. Neuron number in the entorhinal cortex and CA1 in preclinical Alzheimer disease. *Arch Neurol*. 2001;58:1395-1402.
- Arendt T. Synaptic degeneration in Alzheimer's disease. *Acta Neuropathol*. 2009;118:167-179.
- Maslah E, Mallory M, Alford M, et al. Altered expression of synaptic proteins occurs early during progression of Alzheimer's disease. *Neurology*. 2001;56:127-129.
- DeKosky ST, Scheff SW. Synapse loss in frontal cortex biopsies in Alzheimer's disease: correlation with cognitive severity. *Ann Neurol*. 1990;27:457-464.
- Heneka MT, Carson MJ, El Khoury J, et al. Neuroinflammation in Alzheimer's disease. *Lancet Neurol*. 2015;14:388-405.
- Schöll M, Carter SF, Westman E, et al. Early astrocytosis in autosomal dominant Alzheimer's disease measured in vivo by multi-tracer positron emission tomography. *Sci Rep*. 2015;5:16404.
- Suárez-Calvet M, Araque Caballero MÁ, Kleinberger G, et al. Early changes in CSF sTREM2 in dominantly inherited Alzheimer's disease occur after amyloid deposition and neuronal injury. *Sci Transl Med*. 2016;8(369):369ra178.
- Hamilton RL. Lewy bodies in Alzheimer's disease: a neuropathological review of 145 cases using alpha-synuclein immunohistochemistry. *Brain Pathol*. 2000;10:378-384.
- Lippa CF, Fujiwara H, Mann DM, et al. Lewy bodies contain altered alpha-synuclein in brains of many familial Alzheimer's disease patients with mutations in presenilin and amyloid precursor protein genes. *Am J Pathol*. 1998;153:1365-1370.
- Amador-Ortiz C, Lin WL, Ahmed Z, et al. TDP-43 immunoreactivity in hippocampal sclerosis and Alzheimer's disease. *Ann Neurol*. 2007;61:435-445.
- Palmqvist S, Insel PS, Stomrud E, et al. Cerebrospinal fluid and plasma biomarker trajectories with increasing amyloid deposition in Alzheimer's disease. *EMBO Mol Med*. 2019; 11(12):e11170.
- Molinero JL, Gramunt N, Gisbert JD, et al. The ALFA project: a research platform to identify early pathophysiological features of Alzheimer's disease. *Alzheimer's Dement Transl Res Clin Interv*. 2016;2:82-92.
- Jack CR, Bennett DA, Blennow K, et al. A/T/N: an unbiased descriptive classification scheme for Alzheimer disease biomarkers. *Neurology*. 2016;87:539-547.
- Orfanidis SJ. *Introduction to Signal Processing*. Englewood Cliffs, New Jersey: Prentice-Hall; 1996.
- Cleveland WS. Robust locally weighted regression and smoothing scatterplots. *J Am Stat Assoc*. 1979;74:829-836.
- Benjamini Y, Hochberg Y. Controlling the false discovery rate - a practical and powerful approach to multiple testing. *J R Stat Soc Ser B*. 1995;57:289-300.
- Wellington H, Paterson RW, Portelius E, et al. Increased CSF neurogranin concentration is specific to Alzheimer disease. *Neurology*. 2016;86:829-835.
- Tarawneh R, D'Angelo G, Crimmins D, et al. Diagnostic and prognostic utility of the synaptic marker neurogranin in Alzheimer disease. *JAMA Neurol*. 2016;73:561.
- Heslegrave A, Heywood W, Paterson R, et al. Increased cerebrospinal fluid soluble TREM2 concentration in Alzheimer's disease. *Mol Neurodegener*. 2016;11:3.
- Suárez-Calvet M, Kleinberger G, Araque Caballero MÁ, et al. sTREM2 cerebrospinal fluid levels are a potential biomarker for microglia activity in early-stage Alzheimer's disease and associate with neuronal injury markers. *EMBO Mol Med*. 2016;8:466-476.
- Suárez-Calvet M, Morenas-Rodríguez E, Kleinberger G, et al. Early increase of CSF sTREM2 in Alzheimer's disease is associated with tau related neurodegeneration but not with amyloid- $\beta$  pathology. *Mol Neurodegener*. 2019;14:1-14.
- Piccio L, Deming Y, Del-Águila JL, et al. Cerebrospinal fluid soluble TREM2 is higher in Alzheimer disease and associated with mutation status. *Acta Neuropathol*. 2016;131:925-933.
- Henjum K, Almdahl IS, Årskog V, et al. Cerebrospinal fluid soluble TREM2 in aging and Alzheimer's disease. *Alzheimers Res Ther*. 2016;8:17.
- Alcolea D, Carmona-Iragui M, Suárez-Calvet M, et al. Relationship between  $\beta$ -secretase, inflammation and core cerebrospinal fluid biomarkers for Alzheimer's disease. *J Alzheimers Dis*. 2014;42:157-167.
- Alcolea D, Martínez-Lage P, Sánchez-Juan P, et al. Amyloid precursor protein metabolism and inflammation markers in preclinical Alzheimer disease. *Neurology*. 2015;85:626-633.
- Morenas-Rodríguez E, Alcolea D, Suárez-Calvet M, et al. Different pattern of CSF glial markers between dementia with Lewy bodies and Alzheimer's disease. *Sci Rep*. 2019;9:7803.
- Illán-Gala I, Alcolea D, Montal V, et al. CSF sAPP $\beta$ , YKL-40, and NfL along the ALS-FTD spectrum. *Neurology*. 2018;91:e1619-e1628.
- Craig-Schapiro R, Perrin RJ, Roe CM, et al. YKL-40: a novel prognostic fluid biomarker for preclinical Alzheimer's disease. *Biol Psychiatry*. 2010;68:903-912.
- Mattsson N, Insel PS, Palmqvist S, et al. Cerebrospinal fluid tau, neurogranin, and neurofilament light in Alzheimer's disease. *EMBO Mol Med*. 2016;8:1184-1196.
- Zetterberg H, Skillbäck T, Mattsson N, et al. Association of cerebrospinal fluid neurofilament light concentration with Alzheimer disease progression. *JAMA Neurol*. 2016;73:60.
- Lleó A, Alcolea D, Martínez-Lage P, et al. Longitudinal cerebrospinal fluid biomarker trajectories along the Alzheimer's disease continuum in the BIOMARKAPD study. *Alzheimer's Dement*. 2019;15: 742-753.
- Jack CR, Wiste HJ, Weigand SD, et al. Age-specific and sex-specific prevalence of cerebral  $\beta$ -amyloidosis, tauopathy, and neurodegeneration in cognitively unimpaired individuals aged 50-95 years: a cross-sectional study. *Lancet Neurol*. 2017;16:435-444.
- Palmqvist S, Mattsson N, Hansson O. Cerebrospinal fluid analysis detects cerebral amyloid- $\beta$  accumulation earlier than positron emission tomography. *Brain*. 2016;139:1226-1236.
- Vllassenko AG, McCue L, Jasielec MS, et al. Imaging and cerebrospinal fluid biomarkers in early preclinical Alzheimer disease. *Ann Neurol*. 2016;80:379-387.

43. Hansson O, Seibyl J, Stomrud E, et al. CSF biomarkers of Alzheimer's disease concord with amyloid- $\beta$  PET and predict clinical progression: a study of fully automated immunoassays in BioFINDER and ADNI cohorts. *Alzheimer's Dement*. 2018;14:1470-1481.
44. Benzinger TLS, Blazey T, Jack CR, et al. Regional variability of imaging biomarkers in autosomal dominant Alzheimer's disease. *Proc Natl Acad Sci U S A*. 2013;110:E4502-E4509.
45. Schindler SE, Li Y, Todd KW, et al. Emerging cerebrospinal fluid biomarkers in autosomal dominant Alzheimer's disease. *Alzheimer's Dement*. 2019;15:655-665.
46. McDade E, Wang G, Gordon BA, et al. Longitudinal cognitive and biomarker changes in dominantly inherited Alzheimer disease. *Neurology*. 2018;91:e1295-e1306.
47. Fleisher AS, Chen K, Quiroz YT, et al. Associations between biomarkers and age in the presenilin 1 E280A autosomal dominant Alzheimer disease kindred: a cross-sectional study. *JAMA Neurol*. 2015;72:316-324.
48. Wang Y, Shi M, Chung KA, et al. Phosphorylated  $\alpha$ -synuclein in Parkinson's disease. *Sci Transl Med*. 2012;4(121):121ra20.
49. Hansson O, Hall S, Öhrfelt A, et al. Levels of cerebrospinal fluid  $\alpha$ -synuclein oligomers are increased in Parkinson's disease with dementia and dementia with Lewy bodies compared to Alzheimer's disease. *Alzheimers Res Ther*. 2014;6:25.
50. Majbour NK, Vaikath NN, van Dijk KD, et al. Oligomeric and phosphorylated alpha-synuclein as potential CSF biomarkers for Parkinson's disease. *Mol Neurodegener*. 2016;11:7.
51. Tokuda T, Qureshi MM, Ardah MT, et al. Detection of elevated levels of  $\alpha$ -synuclein oligomers in CSF from patients with Parkinson disease. *Neurology*. 2010;75:1766-1772.

#### SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

**How to cite this article:** Milà-Alomà M, Salvadó G, Gispert JD, et al. Amyloid beta, tau, synaptic, neurodegeneration and glial biomarkers in the preclinical stage of the Alzheimer's disease continuum. *Alzheimer's Dement*. 2020;1-14.  
<https://doi.org/10.1002/alz.12131>