

Title: Knowledge of potential harms and benefits of tamoxifen among women considering uptake of breast cancer preventive therapy

Authors: R.J. Thorneloe¹, L. Hall¹, F.M. Walter², L. Side³, K.E. Lloyd¹, S.G. Smith¹, on behalf of the ENGAGE investigators.

Affiliations:

1. Leeds Institute of Health Sciences, University of Leeds, Leeds, UK
2. Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK
3. Wessex Clinical Genetics Service, University Hospitals Southampton, Southampton, UK

Corresponding author: Dr Samuel G. Smith, University of Leeds, Leeds Institute of Health Sciences, Worsley Building, Clarendon Way, Leeds, LS2 9NL, T: 0113 343 0892, E: s.smith1@leeds.ac.uk

Abstract

Background: Tamoxifen reduces breast cancer incidence in women at increased risk, but may cause side-effects. We examined women's knowledge of tamoxifen's potential harms and benefits, and the extent to which knowledge reflects subjective judgements of awareness and decision quality.

Methods: After a hospital appointment, 408 (55.7%) women at increased risk of breast cancer completed a survey assessing objective knowledge about the potential benefit (risk reduction) and harms (endometrial cancer, thromboembolic events, and menopausal side-effects) of tamoxifen, and subjective tamoxifen knowledge and decisional quality. 258 (63.2%) completed the 3-month follow-up survey.

Results: 15.7% of participants recognised the potential benefit and three major harms of using tamoxifen. These women were more likely to have degree level education (vs. below degree level) (OR=2.24, 95%CI: 1.11-4.55) and good numeracy (vs. poor numeracy) (OR=5.91, 95%CI: 1.33-26.19). Tamoxifen uptake was higher in women who recognised all harms and benefits (vs. not recognising) (OR=2.47, 95%CI: 0.94-6.54). 65.8% of tamoxifen users were unaware of its potential benefit and harms. Most (87.1%) women reported feeling informed about tamoxifen, and subjective decisional quality was high (Mean [SD], 17.03 [1.87], out of 18).

Conclusions: Women considering preventive therapy may need additional support to make informed decisions about tamoxifen preventive therapy.

Background

Breast cancer affects over 54,000 UK women each year (1). Women with first- and second-degree relatives with breast cancer are at increased risk of developing the disease (2). Breast cancer risk can be calculated using established models (3). The National Institute of Health and Care Excellence (NICE) considers women with a lifetime risk of breast cancer of 17-30% to be at moderate risk, and those exceeding this level to be at high risk (4). Women within these categories are eligible for early mammographic screening, and have the option of using tamoxifen or anastrozole for preventive therapy.

Tamoxifen can reduce breast cancer risk among this population by at least 30% (5,6). Risk reduction of breast cancer among women using anastrozole is estimated to be 53% (7), and was the recommended option for postmenopausal women in the 2017 updated NICE guidance CG164 (4).

Both medications have the potential for adverse effects. Tamoxifen increases the risk of gynaecological and vasomotor symptoms, thromboembolic events and endometrial cancer (5,8-10). Women using anastrozole are more likely to experience musculoskeletal problems (arthralgia, joint stiffness and carpal tunnel syndrome), vasomotor symptoms and hypertension (7,11). The majority of side-effects start within 12-months of initiation, but can occur throughout treatment (8,9,11).

There is widespread reluctance to use preventive therapy among patients, with fewer than one in 7 eligible women initiating therapy after it has been offered (12-14). Lack of information on chemoprevention has been reported by patients as one barrier to initiation (15,16). Healthcare professionals in primary and secondary care have expressed concern about their ability to discuss the harms and benefits with patients (17). A national survey of United Kingdom (UK) general practitioners (GPs) indicated just over half were aware tamoxifen could reduce breast cancer risk in healthy women, and only 42% felt comfortable discussing its harms and benefits (18). Women may feel dissatisfied with the support they receive from healthcare professionals, and could leave

appointments without a clear understanding of the potential harms and benefits of chemoprevention.

Assessing objective patient knowledge is one approach to evaluating the quality of decisions occurring in this setting. However, it is also important to consider subjective methods, such as the extent to which people feel informed and satisfied with their involvement in the decision-making process (19,20). Subjective knowledge assessments may reflect perceived adequacy of the information provided (20), but are often only weakly associated with objective knowledge (21). To date, no studies have included subjective assessments of decision quality in women considering preventive therapy, indicating that aspects of informed decision-making have not been fully considered. In women making breast cancer treatment decisions, clinician communication style (22,23) and receipt of a decision support tool (24) were related to subjective decision quality, and ethnic minority groups were more likely to report lower quality decision-making experiences (25).

In this study we assessed objective knowledge about the potential harms and benefits of tamoxifen in women at increased risk of breast cancer, and also women's subjective assessments of their knowledge and the quality of their decision about chemoprevention. We examined the socio-demographic, health-service and psychological factors associated with objective knowledge about tamoxifen, and the extent to which knowledge was related to tamoxifen uptake and subjective decisional quality.

Material and methods

Participants

Recruitment took place between September 2015 and December 2016 in hospital outpatient clinic settings. In the UK, women who present to primary care due to concerns about their breast cancer

risk are referred to secondary care if their GP believes they are likely to meet NICE criteria for moderate or high breast cancer risk (4). Four types of clinics in secondary and tertiary care were used to identify women at increased risk of breast cancer: family history clinics (n=12), breast clinics (n=4), clinical genetics centres (n=3), and a family history clinic with genetics support (n=1). Most of the clinics were located in major cities across England. Following their appointment, women were approached by a research nurse or a healthcare professional to discuss the study. Women were eligible if they were aged 18 years or older, spoke English, had discussed preventive therapy with a healthcare professional, were assessed as having a 'moderately high' or 'high' risk of breast cancer according to NICE guidelines (4); and had no known contraindications for tamoxifen use. Women were excluded if they were unable to consent, read English, or had a previous diagnosis of breast cancer.

Measures

Women were invited to complete a baseline survey containing measures of knowledge, healthcare professional satisfaction, and information provision. Women returning baseline questionnaires were sent a follow-up questionnaire at 3-months containing a measure of subjective decisional quality and an item assessing their decision about uptake of tamoxifen for preventive therapy. Length of follow-up was decided on the basis that this was a reasonable amount of time to consider the harms and benefits of tamoxifen, and speak with a GP about obtaining a prescription. The full baseline survey and follow up survey are available here: <https://osf.io/ud67j>.

Objective knowledge about tamoxifen

The items assessing objective knowledge were adapted from a 6-item questionnaire used previously (26). Knowledge was assessed by listing three potential harms of using tamoxifen for preventive therapy (endometrial cancer, menopausal symptoms, blood clotting) and one potential benefit

(breast cancer). For each potential harm and benefit, women were asked 'Who is more likely to experience the following...?' Responses were: 'Women who take tamoxifen', 'Women who do not take tamoxifen', 'Both groups are equally likely', and 'Unsure'. The correct response for the three potential harms was 'Women who take tamoxifen', and the correct response for the potential benefit was 'Women who do not take tamoxifen'. All other responses were marked as incorrect. Women who correctly answered all three potential harms and the potential benefit of using tamoxifen were classified as having good knowledge in the main analysis. Women who had missing data on any of the four items were excluded from the analysis ($n = 25$). The Kuder-Richardson reliability coefficient (KR-20) for this 4-item scale was 0.65.

Subjective knowledge about tamoxifen

A single item assessed perceived knowledge about tamoxifen: 'How informed do you feel about tamoxifen and its use by women at increased risk of breast cancer?' Responses were: 'Not very informed at all' (=1), 'Quite uninformed' (=2), 'Quite well informed' (=3), and 'Very well informed' (=4). We wanted to examine the group of women who reported feeling strongly uninformed, therefore participants who responded 'Not very informed at all' were coded as being uninformed. All other responses were coded as informed in the analysis.

Healthcare professional satisfaction

The validated genetic counselling satisfaction scale was used (27). This 6-item scale assesses patient satisfaction with the appointment and healthcare professional. Example items include: 'The clinician considered any stress I was facing', 'The clinician was concerned about my wellbeing', and 'The appointment was helpful to me'. Each item was scored on a 4-point scale ('strongly disagree' [=1] to 'strongly agree' [=4]). Items were summed to create a scale score ranging from 6 – 24, with higher scores indicating stronger healthcare professional satisfaction. Cronbach's alpha for the total scale was 0.93.

Information provision

A single item assessed whether women reported receiving information about preventive therapy during their appointment: 'During your hospital appointment, did the clinician give you a leaflet about tamoxifen? ('Yes', 'No', and 'Unsure'). No and unsure responses were combined, with missing data for this item also included in this category.

Subjective decision quality

The brief measure of subjective decision quality for breast cancer treatment was adapted for the preventive therapy setting and used in the 3-month follow-up questionnaire (20). This scale assesses six dimensions of subjective decision quality: 'How "right for you" was your decision about tamoxifen' (fit) ('not at all right for me' [=1]; 'neither right or wrong' [=2]; 'completely right for me' [=3]); 'How much information did you have for your decision about tamoxifen' (adequacy of information) ('not enough information' [=1]; 'enough information' [=2]; 'too much information [=3]); 'How much time did you have for your decision about tamoxifen' (adequacy of time) ('not enough time' [=1]; 'just right' [=2]; 'too much time [=3]); 'How much involvement did you have in your decision about tamoxifen' (adequacy of involvement) ('not enough involvement' [=1]; 'just right [=2]; 'too much involvement' [=3]); 'How much regret do you have with regard to your decision about tamoxifen?' (regret) ('no regret' [=1]; 'some regret' [=2]; a lot of regret [=3]); 'How satisfied are you with the decision you made about tamoxifen' (satisfaction) ('not at all satisfied [=1]; 'somewhat satisfied' [=2]; 'totally satisfied' [=3]). Each item included a 'still deciding' response (=4) and women who stated they were 'still deciding' were excluded from the analysis involving this outcome (43.2%; 111/257). Responses were re-coded in line with the original guidelines (20). A single scale score was calculated (ranging from 6 to 18), with higher scores indicating higher subjective decision quality. Cronbach's alpha for the total scale was 0.75.

Preventive therapy decision

Women were asked to indicate which of seven statements applied to them with regard to their decision about using tamoxifen for preventive therapy. The options were: 'I decided immediately that I did not want to take tamoxifen', 'After some thought, I decided that I did not want to take tamoxifen', 'I am still deciding if I want to take tamoxifen', 'I met with my GP to talk about tamoxifen, and I decided against it', 'I met with my GP to talk about tamoxifen, but they would not prescribe it', 'I have a prescription for tamoxifen from my GP', and 'I am currently taking tamoxifen'. Women were classified as initiating tamoxifen if they reported having a prescription for tamoxifen from their GP or were currently taking tamoxifen (13).

Analysis

The analysis plan was pre-registered (DOI 10.17605/OSF.IO/YE6D2). To establish potential bias among the sample retained at follow-up, women responding to the baseline questionnaire only were compared with respondents to the 3-month questionnaire with regard to objective and perceived knowledge about tamoxifen, healthcare professional satisfaction, and information provision. T-tests and Pearson's Chi-square test were used where appropriate. Differences in socio-demographic and clinical factors between responders and non-responders to the baseline questionnaire have previously been reported, and no differences were observed (13).

Measures were described using percentages (with 95% Confidence Intervals [CI]) and means (with Standard Deviations [SD]). A multivariable logistic regression model was used to identify factors associated with good objective knowledge of tamoxifen, defined as recognising all three major potential harms and the potential benefit. This model was adjusted for participant characteristics, information provision, healthcare professional satisfaction and subjective knowledge about tamoxifen. In a sensitivity analysis, the proportion of women who correctly identified the benefit and

at least 2 out of the 3 potential harms associated with tamoxifen were classified as having good knowledge about tamoxifen.

To identify the factors associated with higher subjective decisional quality, we used a multivariable linear regression model adjusted for the same factors previously described as well as objective knowledge about tamoxifen. Multivariable logistic regression was used to examine factors associated with preventive therapy uptake (yes vs. no). Missing data were deleted listwise when examining descriptive data and univariable associations, and pairwise when multivariable analyses were used. A value of $P \leq 0.05$ was considered to be statistically significant. The analysis was completed in SPSS version 21.0.

RESULTS

Sample

In total, 732 women were invited to complete a survey; 408 women (55.7%, 95% CI: 52.1-59.4) returned the baseline survey and 258 (63.2%, 95% CI: 58.4-67.9) women provided uptake data at least 3 months after their appointment.

Participant characteristics, clinical information and descriptions of the measures are presented in Table 1. The mean age of baseline respondents was 45.3 years ($SD \pm 7.82$). The majority of women had children, were married or cohabiting, had attained less than a degree level of education, correctly answered the numeracy item, were white, had good or excellent health, and were employed. The sample included more women who were classified as having a 'moderately high' risk of breast cancer than a 'high' risk of breast cancer. Patient satisfaction with healthcare professionals was high (mean=19.30, $SD \pm 3.77$ out of 24).

There were no differences between women who provided baseline data only and those who were retained at 3-month follow-up with regards to objective knowledge, healthcare professional satisfaction, and information provision (Table S1). A higher proportion of women who provided baseline and 3-month data felt informed about tamoxifen (91.0%, 95% CI: 86.8-94.2) compared with women who provided baseline data only (80.3%, 95% CI: 72.9-86.4).

Knowledge about tamoxifen harms and benefits and information provision

The majority of women reported feeling informed about tamoxifen and its use by women at increased risk of breast cancer (87.1%; 95% CI: 83.4-90.2). However, in the objective assessment, only 15.7% (95% CI: 12.2-19.7) of women correctly identified the potential benefit (breast cancer risk reduction) and all three potential harms (endometrial cancer, menopausal symptoms, blood clotting) of using tamoxifen. Overall, 60.9% of women (95% CI: 55.8-65.7) were aware tamoxifen could reduce breast cancer risk. Half of the sample were aware women taking tamoxifen were more likely to experience menopausal symptoms (50.1%; 95% CI: 45.1-55.2) and blood clotting (49.7%; 95% CI: 44.7-54.8), and 27.3% (95% CI: 22.9-32.0) had knowledge about the increased risk of endometrial cancer associated with tamoxifen. A fifth (18.2%; 95% CI: 14.5-22.4) of women recognised all three potential harms associated with taking tamoxifen.

In a multivariable analysis, women with a higher level of education (odds ratio (OR)=2.24, 95% CI: 1.11-4.55, $p = 0.025$) and with higher numeracy levels (OR=5.91, 95% CI: 1.33-26.19, $p = 0.019$) were more likely to have good knowledge about the potential benefits and harms of tamoxifen (Table 2). Sensitivity analyses yielded similar results (Table S2).

In total, 71.1% (95% CI: 66.4-75.4) of women stated their healthcare professional gave them a leaflet about tamoxifen during their appointment, with 28.9% (95% CI: 24.6-33.6) of women reporting that they did not receive or were unsure whether they received a leaflet. A higher proportion of women

who received a leaflet about tamoxifen had good knowledge on its harms and benefits (17.9%), compared with those who did not receive a leaflet or were unsure (10.1%). In the multivariable analysis (Table 2), this difference was not statistically significant (OR=1.54, 95% CI: 0.66 – 3.59, p = 0.313).

Subjective decisional quality

Among women who were followed-up at 3 months (n=258), 257 completed the decisional quality scale. Of this group, 111 (43.2%; 95% CI: 37.1-49.5) stated they were still deciding about tamoxifen (on ≥ 1 item) and 146 (56.8%; 95% CI: 50.5-62.9) had made their decision regarding tamoxifen.

Overall, women reported a high level of decisional quality about tamoxifen, with a mean \pm SD score of 17.03 ± 1.87 out of a possible 18. Most women reported having no regret about their decision (92.5%; 95% CI: 88.2, 96.8; 135/146), having the right amount of involvement in their decision (93.8%; 95% CI: 89.9, 97.7; 137/146), felt enough information was provided (90.4%; 95% CI: 85.6, 95.2; 132/146), and had the right amount of time to make a decision (95.9%; 95% CI: 92.7, 99.1; 140/146). Endorsement was lower for whether the decision was 'completely right for me' (78.1%; 95% CI: 71.4, 84.8; 114/146), and for feeling totally satisfied with their decision (77.4%; 95% CI: 70.6, 84.2; 113/146).

In a multivariable model including women who had reached a decision about tamoxifen initiation (n=128), those who felt more informed about tamoxifen (vs. less informed) ($\beta = 0.22$, p = 0.018) and who were from more disadvantaged socioeconomic backgrounds (vs. least deprived) ($\beta = 0.27$, p = 0.015) reported higher decisional quality (Table 3) (n=18 missing data).

Factors associated with uptake of tamoxifen

Uptake of tamoxifen at 3 month follow-up was 14.7% (95% CI: 10.6-19.7, 38/258). A higher proportion of women with good objective knowledge about the potential benefits and harms of

using tamoxifen initiated chemoprevention (27.9%; 95% CI: 15.3-43.7, 12/43), compared with those with poor knowledge (12.2%; 95% CI: 8.00-17.5, 25/205) (Table 4). There were no statistically significant associations in the multivariable model (Table 4). Despite greater knowledge among women using tamoxifen compared with non-users, levels were not optimal: 65.8%, 95% CI: 48.6-80.4 (25/38) of this group failed to recognise the major potential benefit and three major potential harms of the drug.

Discussion

In this multi-centre survey of healthy women considering breast cancer primary prevention, only 1 in 6 women recognised the potential benefits and harms of tamoxifen, i.e. that it can reduce breast cancer risk, but may cause thromboembolic events, endometrial cancer and menopausal side-effects. A third of women in this sample did not recognise any of these harms. Knowledge was particularly poor among women with lower levels of education and numeracy. Knowledge was higher among women who reported tamoxifen use at follow-up, but even among this group two thirds failed to recognise the key benefit and all three harms. Women's decisions regarding breast cancer preventive therapy appear to be based on incomplete information, and this may be a particular problem for those with low literacy and numeracy skills.

The finding of poor knowledge among users of tamoxifen may be problematic from a safety perspective. A lower level of awareness of these potential harms may hinder recognition of an adverse event and delay or prevent help-seeking behaviour. Healthcare professionals counselling women who are initiating tamoxifen for primary prevention should promote awareness of these signs and symptoms during the counselling process. This should be presented using absolute risk information so patients are aware these harms are possible, but can accurately gauge the likelihood of experiencing them. If GPs are to take on this role, they are likely to require information and support. A recent national survey of GPs indicated nearly 60% felt uncomfortable discussing the

harms and benefits of tamoxifen for primary prevention (18), although this study was conducted prior to its being licensed for this indication, in November 2018. Low knowledge among women initiating preventive therapy may also have detrimental effects on long-term adherence to the medication (28), which has been shown to be problematic in this context (9,11).

As part of their discussion about tamoxifen, a substantial minority of women did not report receiving written information during their appointment. Clinical centres should make efforts to improve dissemination to ensure all women considering preventive therapy are provided with written informational support. However, improvements may be needed to the information currently being provided, as receipt of information may not substantially affect women's knowledge of tamoxifen's harms and benefits. This may be because the information was overly complex, which is supported by the observation that women with lower levels of education and numeracy had poorer knowledge. NICE decision tools are freely available (4). While these aids frame numerical information using evidence-based approaches (29), they are lengthy documents that contain complex terminology. These aids should be user-tested with the appropriate groups to ensure comprehension and usability (30,31). Alternative risk communication tools, such as interactive websites and 'gist-based information', have been shown to be acceptable to patients (31-33), and can improve breast cancer risk perceptions, knowledge and interest in using chemoprevention (34,35). Developing similar tools for use in the context of breast cancer prevention the UK could be a useful next step.

The data presented here highlight a gap between participants' perceived and actual levels of knowledge about tamoxifen. The low levels of objective knowledge reported were not reflected in women's judgements about their own awareness, with nearly all respondents indicating that they felt quite or very informed about tamoxifen. Weak associations between subjective and objective knowledge measures have previously been reported (21). Furthermore, objective knowledge did not affect subjective decisional quality, with average scores for this construct being notably high. The

high levels of subjective decisional quality suggest that most women felt supported and involved in their decision about chemoprevention, and were satisfied with their choice. While there are no 'correct' responses to subjective decision quality scales (20), our data indicate women may be unaware of gaps in their knowledge, and as a result believe their decisions to be of high quality. Healthcare professionals should be aware that subjective assessments of knowledge and decisional quality may not reflect their own interpretations of a high quality decision in the preventive therapy setting. Clinicians may also need to go beyond simple checks of comprehension, and instead use techniques such as the 'teach back' method to verify understanding in verbal consultations (36,37). This involves the patient reporting back to the communicator their interpretation of the message conveyed, with misunderstandings addressed and clarified. Such measures could enhance the quality of informed decision making about tamoxifen chemoprevention for women at increased risk of developing breast cancer, even if uptake did not increase.

This study had limitations. While response rates were high, and there were few differences among responders and non-responders, those who did not complete the questionnaires may have scored differently on the assessments. The proportion of women using tamoxifen was small, and therefore the confidence intervals for estimates involving this outcome were wide. These data were collected prior to the NICE 2017 updated guidelines recommending anastrozole for postmenopausal women, and therefore our findings may not be generalisable to women considering this drug. The reliability of the subjective decision scale was below levels previously reported (20), and may have affected the associations reported. The study relied on self-reported uptake data, and therefore estimates may not be as reliable as more objective assessments. In this study, women were followed up at 3 months to investigate their decision about tamoxifen as this was decided to be a reasonable amount of time to consider the harms and benefits, and speak with a GP. However, data from the subjective decisional quality scale indicated approximately half of the sample were still considering whether to use tamoxifen. A longer follow-up may have affected the proportion of women currently using

tamoxifen, and their responses to the decision quality scale. Furthermore, their decision to use breast cancer chemoprevention may have been influenced by discussions in primary care.

Tamoxifen's off-licence status has previously been identified as a barrier to GP prescribing for breast cancer prevention (16), and the data presented here were collected before this became a licensed indication in November 2018.

In conclusion, the majority of women at increased risk of breast cancer considering the use of tamoxifen for primary prevention reported low levels of knowledge in relation to its major potential benefit and harms. Knowledge was particularly poor among women with lower levels of education and numeracy, which has the potential to exacerbate socioeconomic inequalities. While information leaflets and support tools are available and appear to be provided to the majority of women faced with this decision, they do not appear to be effective at supporting adequate understanding of the potential harms and benefits. Objective knowledge was not associated with women's subjective assessments of tamoxifen knowledge or decisional quality. Healthcare professionals should therefore be cautious about assuming an informed decision about breast cancer prevention has been made, and techniques such as the 'teach back' method could be usefully employed. There is also scope for developing and testing decision aids about primary prevention of breast cancer using different formats and "gist" versus "verbatim" detail.

Acknowledgements The authors acknowledge the contribution of the ENGAGE collaborators (in alphabetical order): Vanessa Adamson, Sarah Ainsworth, Malin Akerlund, Ivanna Baker, Julian Barwell, Jayne Beesley, Lisa Brock, Chrissie Butcher, Janice Carpenter, Martyn Clark, Shirley Cocks, Veronica Conteh, Martina Coulding, Sue Darby, Angela Duckworth, Gareth Evans, Catherine Fensom, Julie Fletcher, Kate Foster, Sara Grieg, Elaine Gullaksen, Jana Gurasashvili, Lisa Hardstaff, Rachel Hart, Kathryn Hoare, Jonathan Hoffman, Christopher Holcombe, Lynne Horton, Antony Howell, Farah Islam, Emma Jenkinson, Karen Jewers, Manisha Joshi, Amy Kirkby, Peter Kneeshaw, Natalie Knife, Jalal Kokan, Jin Li, Nicola Lunt, Douglas Macmillan, Karen Makinson, Evangelos Mallidis, Sarah Manyangadze, Charity Masvaure, Raksha Mistry, Alice Ngumo, Jane Ooi, Ashraf Patel, Vanessa Pope, Laura Price, Fiona Rabson, Lisa Richardson, Stephanie Ridgway, Karen Riley, Lorraine Roberts, Janet Ryan-Smith, Vian Salih, Nicky Scott, Mike Shere, Andrew Sloan, Nita Solanky, Amanda Taylor, Dinesh Thekkinkattil, Heather Thomas, Mangesh Thorat, Barbara Townley, Jayant S. Vaidya, Lynda Wagstaff, Shane Walsh, Lynsey Waring, Donna Watterson, Charlotte Westley, Lesley Wilkinson, Nicola Willis, and Julia Wiseman.

We also thank the women participating in the study.

Availability of data and material: Participants did not provide explicit consent for their data to be shared in public repositories. Therefore, data may not be made publicly available due to ethical restrictions. We can share the anonymised version of the data to individual qualified researchers upon request. Data requests may be sent to the corresponding author of this paper.

Funding: SGS was supported by a Cancer Research UK postdoctoral fellowship (C42785/A17965) during the collection of these data. He also acknowledges funding support from a Yorkshire Cancer Research University Academic Fellowship. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Ethical approval and consent to participate: Ethical approval was awarded by the National Research Ethics Service Committee North West—Preston (14/ NW/1408). All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from all individual participants included in this study.

Conflict of interest: RJT has received honorarium from Novartis. SS has received consulting fees from Luto Research. All other authors have declared no conflicts of interest.

Authorship: S.G.S and L.S designed the study. R.J.T. and K.E.L analysed and interpreted the results. R.J.T, L.H, F.M.W, L.S, K.E.L, and S.G.S helped to write the manuscript.

References

1. Cancer Research UK. Breast cancer statistics [Internet]. 2017 [cited 2018 June]. Available from: <http://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/breast-cancer>
2. Nelson HD, Zakher B, Cantor A, Fu R, Griffin J, O'Meara ES, et al. Risk factors for breast cancer for women aged 40 to 49 years: a systematic review and meta-analysis. *Ann Intern Med*. 2012 May 1;156(9):635–48.
3. Brentnall AR, Cuzick J, Buist DSM, Bowles EJA. Long-term accuracy of breast cancer risk assessment combining classic risk factors and breast density. *JAMA Oncol*. 2018 Sep 1;4(9):e180174.
4. National Institute for Health and Care Excellence. Familial breast cancer: classification, care and managing breast cancer and related risks in people with a family history of breast cancer [Internet]. 2017 [cited 2018 June]. Available from: <https://www.nice.org.uk/guidance/CG164>
5. Cuzick J, Sestak I, Bonanni B, Costantino JP, Cummings S, DeCensi A, et al. Selective oestrogen receptor modulators in prevention of breast cancer: an updated meta-analysis of individual participant data. *Lancet*. 2013 May 25;381(9880):1827–34.
6. Cuzick J, Sestak I, Cawthorn S, Hamed H, Holli K, Howell A, et al. Tamoxifen for prevention of breast cancer: extended long-term follow-up of the IBIS-I breast cancer prevention trial. *Lancet Oncol*. 2015 Jan;16(1):67–75.
7. Cuzick J, Sestak I, Forbes JF, Dowsett M, Knox J, Cawthorn S, et al. Anastrozole for prevention of breast cancer in high-risk postmenopausal women (IBIS-II): an international, double-blind, randomised placebo-controlled trial. *The Lancet*. 2014 Mar 22;383(9922):1041-8.
8. Cuzick J, Forbes JF, Sestak I, Cawthorn S, Hamed H, Holli K, et al. Long-term results of tamoxifen prophylaxis for breast cancer—96-month follow-up of the randomized IBIS-I trial. *J Natl Cancer Inst*. 2007 Feb 21;99(4):272–82.
9. Smith SG, Sestak I, Howell A, Forbes J, Cuzick J. Participant-reported symptoms and their effect on long-term adherence in the international breast cancer intervention study I (IBIS I). *J Clin Oncol*. 2017 Jun 29;35(23):2666-73.
10. Cuzick J, Forbes J, Edwards R, Baum M, Cawthorn S, Coates A, et al. First results from the International Breast Cancer Intervention Study (IBIS-I): a randomised prevention trial. *The Lancet*. 2002 Sep 14;360(9336):817–24.
11. Sestak I, Smith SG, Howell A, Forbes J, Cuzick J. Early participant-reported symptoms as predictors of adherence to anastrozole in the International Breast Cancer Intervention Studies II. *Ann Oncol*. 2017 Nov 6;29(2):504-9.
12. Smith SG, Sestak I, Forster A, Partridge A, Side L, Wolf MS, et al. Factors affecting uptake and adherence to breast cancer chemoprevention: a systematic review and meta-analysis. *Ann Oncol*. 2016 Apr;27(4):575–90.

13. Hackett J, Thorneloe R, Side L, Wolf M, Horne R, Cuzick J, et al. Uptake of breast cancer preventive therapy in the UK: results from a multicentre prospective survey and qualitative interviews. *Breast Cancer Res Treat.* 2018 Aug 1;170(3):633–40.
14. Thorneloe RJ, Horne R, Side L, Wolf MS, Smith SG, Adamson V, et al. Beliefs about medication and uptake of preventive therapy in women at increased risk of breast cancer: results from a multicenter prospective study. *Clin Breast Cancer.* 2019 Feb;19(1):e116–26.
15. Heisey R, Pimlott N, Clemons M, Cummings S, Drummond N. Women's views on chemoprevention of breast cancer. *Can Fam Physician.* 2006 May 10;52(5):624–5.
16. Port ER, Montgomery LL, Heerdt AS, Borgen PI. Patient reluctance toward tamoxifen use for breast cancer primary prevention. *Ann Surg Oncol.* 2001 Aug;8(7):580–5.
17. Smith SG, Side L, Meisel SF, Horne R, Cuzick J, Wardle J. Clinician-reported barriers to implementing breast cancer chemoprevention in the UK: a qualitative investigation. *Public Health Genomics.* 2016 Aug 23;19(4):239–49.
18. Smith SG, Foy R, McGowan JA, Kobayashi LC, Censi A de, Brown K, et al. Prescribing tamoxifen in primary care for the prevention of breast cancer: a national online survey of GPs' attitudes. *Br J Gen Pract.* 2017 Jun;67(659):e414–27.
19. Sepucha KR, Belkora JK, Chang Y, Cosenza C, Levin CA, Moy B, et al. Measuring decision quality: psychometric evaluation of a new instrument for breast cancer surgery. *BMC Med Inform Decis Mak.* 2012 Jun 8;12(1):51–61.
20. Resnicow K, Abrahamse P, Tocco RS, Hawley S, Griggs J, Janz N, et al. Development and psychometric properties of a brief measure of subjective decision quality for breast cancer treatment. *BMC Med Inform Decis Mak.* 2014 Dec 5;14(1):110–9.
21. Sepucha KR, Fagerlin A, Couper MP, Levin CA, Singer E, Zikmund-Fisher BJ. How does feeling informed relate to being informed? The DECISIONS survey. *Med Decis Making.* 2010 Oct;30(5 Suppl):77S–84S.
22. Martinez KA, Resnicow K, Williams GC, Silva M, Abrahamse P, Shumway DA, et al. Does physician communication style impact patient report of decision quality for breast cancer treatment? *Patient Educ Couns.* 2016 Dec 1;99(12):1947–54.
23. Katz SJ, Janz NK, Abrahamse P, Wallner LP, Hawley ST, An LC, et al. Patient reactions to surgeon recommendations about contralateral prophylactic mastectomy for treatment of breast cancer. *JAMA Surg.* 2017 Jul 1;152(7):658–64.
24. Hawley ST, Li Y, An LC, Resnicow K, Janz NK, Sabel MS, et al. Improving breast cancer surgical treatment decision making: the iCanDecide randomized clinical trial. *J Clin Oncol.* 2018 Jan 24;36(7):659–66.
25. Katz SJ, Wallner LP, Abrahamse PH, Janz NK, Martinez KA, Shumway DA, et al. Treatment experiences of Latinas after diagnosis of breast cancer. *Cancer.* 2017;123(16):3022–30.
26. Fagerlin A, Zikmund-Fisher BJ, Smith DM, Nair V, Derry HA, McClure JB, et al. Women's decisions regarding tamoxifen for breast cancer prevention: responses to a tailored decision aid. *Breast Cancer Res Treat.* 2010 Feb 1;119(3):613–20.

27. DeMarco TA, Peshkin BN, Mars BD, Tercyak KP. Patient satisfaction with cancer genetic counseling: a psychometric analysis of the genetic counseling satisfaction scale. *J Genet Couns.* 2004 Aug 1;13(4):293–304.
28. Rottman BM, Marcum ZA, Thorpe CT, Gellad WF. Medication adherence as a learning process: insights from cognitive psychology. *Health Psychol Rev.* 2017;11(1):17–32.
29. Spiegelhalter D. Risk and uncertainty communication. *Annu Rev Stat Its Appl.* 2017;4(1):31–60.
30. Raynor DK, Knapp P, Silcock J, Parkinson B, Feeney K. “User-testing” as a method for testing the fitness-for-purpose of written medicine information. *Patient Educ Couns.* 2011 Jun;83(3):404–10.
31. Smith SG, Wolf MS, Obichere A, Raine R, Wardle J, von Wagner C. The development and testing of a brief ('gist-based') supplementary colorectal cancer screening information leaflet. *Patient Educ Couns.* 2013 Dec;93(3):619–25.
32. Smith SG, Raine R, Obichere A, Wolf MS, Wardle J, von Wagner C. The effect of a supplementary ('gist-based') information leaflet on colorectal cancer knowledge and screening intention: a randomized controlled trial. *J Behav Med.* 2015;38(2):261–72.
33. Meisel SF, Freeman M, Waller J, Fraser L, Gessler S, Jacobs I, et al. Impact of a decision aid about stratified ovarian cancer risk-management on women's knowledge and intentions: a randomised online experimental survey study. *BMC Public Health.* 2017 Nov 16;17(1):882-94.
34. Kukafka R, Fang J, Vanegas A, Silverman, Crew KD. Pilot study of decision support tools on breast cancer chemoprevention for high-risk women and healthcare providers in the primary care setting. *BMC Med Inform Decis Mak.* 2018;18(1):134-45
35. Kukafka R, Yi H, Xiao T, Thomas P, Aguirre A, Smalletz C, et al. Why breast cancer risk by the numbers is not enough: evaluation of a decision aid in multi-ethnic, low-numerate women. *J Med Internet Res.* 2015 Jul 14;17(7):e165.
36. Griffey RT, Shin N, Jones S, Aginam N, Gross M, Kinsella Y, et al. The impact of teach-back on comprehension of discharge instructions and satisfaction among emergency patients with limited health literacy: a randomized, controlled study. *J Commun Healthc.* 2015 Mar 1;8(1):10–21.
37. Kornburger C, Gibson C, Sadowski S, Maletta K, Klingbeil C. Using “Teach-Back” to promote a safe transition from hospital to home: an evidence-based approach to improving the discharge process. *J Pediatr Nurs.* 2013 May 1;28(3):282–91.

TABLES AND FIGURES

Table 1. Description of demographic, clinical and psychological data

Mean (\pm SD) for continuous variables; % (95% CI) for categorical variables		
Baseline (n= 408)		<i>n</i>
Age, mean \pm SD	45.30 \pm 7.82	408
Children		
Yes	77.0 (72.6, 81.0)	314
No	23.0 (19.0, 27.4)	94
White ethnic group		
White	95.5 (93.0, 97.3)	384
Other	4.5 (2.7, 7.0)	18
Missing, n	6	
Education level		
Degree or above	44.2 (39.3, 49.3)	176
Below degree level	55.8 (50.7, 60.7)	222
Missing, n	10	
Health status		
Poor	4.0 (2.3, 6.4)	16
Fair	19.5 (15.7, 23.7)	78
Good	60.0 (55.0, 64.8)	240
Excellent	16.5 (13.0, 20.5)	66
Missing, n	8	
Risk level		
Moderate	59.6 (54.6, 64.4)	243
High	39.0 (34.2, 43.9)	159
Unclear	1.5 (0.5, 3.2)	6
SES		
Low (most deprived)	29.9 (25.5, 34.7)	120
Middle	32.7 (28.1, 37.5)	131
High (least deprived)	37.4 (32.7, 42.3)	150
Missing, n	7	
Employment		
Full-time	85.3 (81.5, 88.6)	348
All other employments	14.7 (11.4, 18.5)	60
Marital status		
Married or cohabiting	74.3 (69.7, 78.5)	298
Unmarried	25.7 (21.5, 30.3)	103
Missing, n	7	
Numeracy		
Good numeracy	81.8 (77.6, 85.5)	323
Poor numeracy	18.2 (14.5, 22.4)	72
Missing, n	13	
Received a leaflet about tamoxifen		
Yes	71.1 (66.4, 75.4)	290
No	25.7 (21.6, 30.3)	105
Unsure	3.2 (1.7, 5.4)	13
Knowledge		
Potential benefit and harms of using		

tamoxifen⁺		
Yes	15.7 (12.2, 19.7)	60
No	84.3 (80.3, 87.8)	323
Missing, n	25	
Tamoxifen can reduce breast cancer risk		
Yes	60.9 (55.8, 65.7)	238
No	39.1 (34.3, 44.2)	153
Missing, n	17	
Tamoxifen can increase risk of menopausal symptoms		
Yes	50.1 (45.1, 55.2)	197
No	49.9 (44.8, 54.9)	196
Missing, n	15	
Tamoxifen can increase risk of blood clotting		
Yes	49.7 (44.7, 54.8)	193
No	50.3 (45.2, 55.3)	195
Missing, n	20	
Tamoxifen can increase risk of endometrial cancer		
Yes	27.3 (22.9, 32.0)	107
No	72.7 (68.0, 77.1)	285
Missing, n	16	
Satisfaction with healthcare professional, mean ± SD	19.30 (3.77)	375
Missing, n	33	
Felt informed about tamoxifen		
Yes	87.1 (83.4, 90.2)	351
No	12.9 (9.8, 16.6)	52
Missing, n	5	
Three month follow-up (n = 258)		
Subjective decisional quality, Mean ± SD	17.03 ± 1.87	146
'Still deciding', n	111	
Missing, n	151	

Key: CI, Confidence Interval; SES, Socioeconomic status; SD, Standard deviation.

*= This variable included the potential benefit of tamoxifen (a reduction in breast cancer risk) and three potential harms of tamoxifen (increased risk of menopausal symptoms, blood clotting and endometrial cancer).

Table 2. Knowledge about the potential harms and benefits of tamoxifen by participant characteristics and univariable and multivariable logistic regression model (n=317)

Good knowledge ⁺ (%; N)		Univariable		Multivariable	
		OR (95% CI)	P-value	OR (95% CI)	P-value
Age					
≤ 35 years	10.3 (4)	0.66 (0.20 - 2.11)	0.479	0.54 (0.11 - 2.61)	0.443
36 - 49 years	16.9 (41)	1.16 (0.61 - 2.21)	0.644	0.71 (0.31 - 1.62)	0.417
≥ 50 years	14.9 (15)	Ref		Ref	
Children					
Yes	16.2 (48)	1.21 (0.61 - 2.40)	0.585	1.38 (0.54 - 3.56)	0.504
No	13.8 (12)	Ref		Ref	
Ethnic group*					
White	15.1 (55)	-	-	-	-
Other	28.6 (4)	-	-	-	-
Education level					
Degree or above	22.2 (38)	2.40 (1.36 - 4.25)	0.003	2.24 (1.11 - 4.55)	0.025
Below degree level	10.6 (22)	Ref		Ref	
Health status					
Poor*	20.0 (3)	-		-	
Fair	6.8 (5)	0.36 (0.12 - 1.13)	0.079	0.39 (0.11 - 1.41)	0.149
Good	18.3 (42)	1.12 (0.53 - 2.39)	0.764	1.13 (0.48 - 2.65)	0.777
Excellent	16.7 (10)	Ref		Ref	
Risk level					
Moderate	18.9 (43)	1.82 (0.99 - 3.33)	0.052	1.96 (0.95 - 4.04)	0.069
High	11.3 (17)	Ref		Ref	
Unclear*	0.0 (0)	-		-	
SES					
Low (most deprived)	14.3 (16)	0.83 (0.42 - 1.67)	0.606	0.97 (0.43 - 2.22)	0.945
Middle	15.0 (19)	0.88 (0.45 - 1.71)	0.704	0.87 (0.39 - 1.93)	0.734
High (least deprived)	16.7 (23)	Ref		Ref	
Employment					
Full-time	16.1 (53)	Ref		Ref	
All other employments	13.0 (7)	0.78 (0.33 - 1.81)	0.556	1.07 (0.42 - 2.74)	0.894
Marital status					
Married or cohabiting	16.3 (46)	1.16 (0.60 - 2.21)	0.662	0.99 (0.43 - 2.29)	0.985
Unmarried	14.4 (14)	Ref		Ref	
Numeracy					
Good numeracy	19.0 (58)	7.72 (1.84 - 32.43)	0.005	5.91 (1.33 - 26.19)	0.019
Poor numeracy	2.9 (2)	Ref		Ref	
Received a leaflet about tamoxifen					
Yes	17.9 (49)	1.94 (0.97 - 3.89)	0.062	1.54 (0.66 - 3.59)	0.313
No/unsure	10.1 (11)	Ref		Ref	
Satisfaction with healthcare professional					
Satisfaction score, mean ± SD	19.34 ± 3.55	1.00 (0.93 - 1.09)	0.922	0.98 (0.90 - 1.07)	0.664
Felt informed about tamoxifen					
Yes	17.4 (58)	4.85 (1.15 - 20.55)	0.032	5.36 (0.67 - 42.66)	0.113

No	4.2 (2)	Ref		Ref	
----	---------	-----	--	-----	--

*category not included in univariable and multivariable analyses due to insufficient cases.

*= This variable included the potential benefit of tamoxifen (a reduction in breast cancer risk) and three potential harms of tamoxifen (increased risk of menopausal symptoms, blood clotting and endometrial cancer).

Table 3. Subjective decisional quality by participant characteristics and univariable and multivariable linear regression model (n = 128)

Decisional quality (mean ± SD)		Univariable			Multivariable		
		B (SE)	β	P-value	B (SE)	β	P-value
Age							
≤ 35 years	16.41 ± 2.15	-0.26 (0.56)	-0.04	0.652	0.17 (0.74)	0.03	0.816
36 – 49 years	17.24 ± 1.51	0.58 (0.39)	0.14	0.139	0.50 (0.48)	0.12	0.294
≥ 50 years	16.67 ± 2.59	Ref	-	-	Ref	-	-
Children							
Yes	17.13 ± 1.58	0.48 (0.37)	0.11	0.205	0.24 (0.52)	0.05	0.644
No	16.66 ± 2.66	Ref	-	-	Ref	-	-
Ethnic group							
White	17.04 ± 1.89	0.20 (0.78)	0.02	0.796	0.73 (0.97)	0.07	0.451
Other	16.83 ± 1.33	Ref	-	-	Ref	-	-
Education level							
Degree or above	17.08 ± 1.86	0.11 (0.31)	0.03	0.728	-0.21 (0.41)	-0.05	0.604
Below degree level	16.97 ± 1.90	Ref	-	-	Ref	-	-
Health status							
Poor	17.17 ± 0.98	0.13 (0.83)	0.01	0.871	0.18 (1.12)	0.02	0.876
Fair	16.19 ± 2.65	-0.84 (0.49)	-0.17	0.091	-1.13 (0.58)	-0.23	0.054
Good	17.28 ± 1.65	0.25 (0.39)	0.07	0.526	0.04 (0.46)	0.01	0.936
Excellent	17.03 ± 1.66	Ref	-	-	Ref	-	-
Risk level							
Moderate	17.15 ± 1.41	0.38 (0.33)	0.10	0.248	0.25 (0.38)	0.06	0.515
High	16.77 ± 2.57	Ref	-	-	Ref	-	-
Unclear†	-	-	-	-	-	-	-
SES							
Low (most deprived)	17.36 ± 1.11	0.40 (0.41)	0.09	0.333	1.25 (0.51)	0.27	0.015
Middle	16.84 ± 2.28	-0.12 (0.36)	-0.03	0.731	0.08 (0.42)	0.02	0.855
High (least deprived)	16.97 ± 1.84	Ref	-	-	Ref	-	-
Employment							
Full-time	17.10 ± 1.90	0.48 (0.44)	0.09	0.281	0.60 (0.53)	0.11	0.258
All other	16.62 ± 1.66	Ref	-	-	Ref	-	-
Marital status							
Married or cohabiting	17.13 ± 1.57	0.53 (0.38)	0.12	0.169	0.41 (0.51)	0.09	0.418
Unmarried	16.60 ± 2.75	Ref	-	-	Ref	-	-
Numeracy							
Good numeracy	17.07 ± 1.80	0.37 (0.43)	0.07	0.389	0.42 (0.49)	0.08	0.393
Poor numeracy	16.70 ± 2.29	Ref	-	-	Ref	-	-
Received a leaflet about tamoxifen							
Yes	17.13 ± 1.37	0.34 (0.34)	0.08	0.323	0.21 (0.42)	0.05	0.620
No/unsure	16.79 ± 2.75	Ref	-	-	Ref	-	-
Satisfaction with healthcare professional							
Satisfaction score, mean ± SD	-	0.02 (0.04)	0.03	0.729	-0.02 (0.05)	-0.04	0.642
Felt informed about							

tamoxifen							
Yes	17.13 ± 1.77	1.88 (0.67)	0.23	0.005	1.78 (0.74)	0.22	0.018
No	15.25 ± 2.76	Ref	-	-	Ref		-
Knowledge about harms and benefits							
Yes	17.33 ± 1.41	0.37 (0.41)	0.08	0.365	0.23 (0.48)	0.04	0.640
No	16.97 ± 1.99	Ref	-	-	Ref		

†category not included in univariable and multivariable analyses due to insufficient cases.

Table 4. Uptake of tamoxifen by participant characteristics and univariable and multivariable logistic regression model (n = 187)

Uptake of tamoxifen (N; %)		Univariable		Multivariable	
		OR (95% CI)	P-value	OR (95% CI)	P-value
Age					
≤ 35 years†	1 (3.8)	-		-	-
36 - 49 years	29 (17.3)	1.46 (0.63 - 3.39)	0.378	1.31 (0.46 - 3.76)	0.617
≥ 50 years	8 (12.5)	Ref		Ref	
Children					
Yes	36 (17.6)	5.43 (1.26 - 23.34)	0.023	3.71 (0.73 - 18.80)	0.113
No	2 (3.8)	Ref		Ref	
Ethnic group†					
White	37 (15)	-	-	-	-
Other	1 (11.1)	Ref		Ref	
Education level					
Degree or above	20 (17.2)	1.41 (0.71 - 2.82)	0.327	1.54 (0.61 - 3.89)	0.360
Below degree level	18 (12.9)	Ref		Ref	
Health status					
Poor†	0	-	-	-	-
Fair	5 (10.6)	0.68 (0.20 - 2.32)	0.538	0.81 (0.19 - 3.56)	0.783
Good	25 (16.6)	1.13 (0.46 - 2.82)	0.787	1.28 (0.43 - 3.84)	0.661
Excellent	7 (14.9)	Ref		Ref	
Risk level					
Moderate	24 (15.1)	1.05 (0.52 - 2.15)	0.885	0.75 (0.32 - 1.77)	0.510
High	14 (14.4)	Ref		Ref	
Unclear†	0	-	-	-	-
SES					
Low (most deprived)	7 (11.9)	0.78 (0.30 - 2.03)	0.613	1.57 (0.54 - 4.58)	0.412
Middle	14 (16.3)	1.13 (0.52 - 2.47)	0.759	1.66 (0.64 - 4.27)	0.296
High (least deprived)	16 (14.7)	Ref		Ref	
Employment					
Full-time	32 (14.5)	Ref		Ref	
All other employments	6 (16.2)	1.14 (0.44 - 2.96)	0.783	1.88 (0.62 - 5.68)	0.265
Marital status					
Married or cohabiting	33 (16.7)	2.16 (0.80 - 5.81)	0.127	1.64 (0.47 - 5.70)	0.435
Unmarried	5 (8.5)	Ref		Ref	
Numeracy					
Good numeracy	32 (15.3)	1.37 (0.50 - 3.76)	0.536	1.21 (0.35 - 4.23)	0.765
Poor numeracy	5 (11.6)	Ref		Ref	
Received a leaflet about tamoxifen					
Yes	28 (14.9)	1.05 (0.48 - 2.29)	0.903	0.62 (0.24 - 1.58)	0.314
No/unsure	10 (14.3)	Ref		Ref	
Satisfaction with healthcare professional					
Satisfaction score, mean ± SD	19.54 ± 3.34	1.02 (0.93 - 1.12)	0.689	1.06 (0.94 - 1.20)	0.349
Felt informed about tamoxifen†					
Yes	38 (16.3)	-	-	-	-
No	0 (0.0)	Ref		Ref	

Knowledge about harms and benefits					
Yes	12 (27.9)	2.79 (1.27 - 6.12)	0.011	2.47 (0.94 - 6.54)	0.067
No	25 (12.2)	Ref		Ref	

†category not included in univariable and multivariable analyses due to insufficient cases.

SUPPLEMENTARY FILES

Table S1. Univariable comparison of retention (n = 408)

Mean (\pm SD) for continuous variables; % (n) for categorical variables	Baseline only (n = 150)	Baseline and 3 months (n = 258)	P-value
Knowledge about harms and benefits			
Yes	12.6 (17)	17.3 (43)	0.242
No	87.4 (118)	82.7 (205)	
Satisfaction with healthcare professional			
Satisfaction score, mean \pm SD	19.30 \pm 3.68	19.30 \pm 3.83	0.987
Received a leaflet about tamoxifen			
Yes	68.0 (102)	72.9 (188)	0.310
No/unsure	32.0 (48)	27.1 (70)	
Felt informed about tamoxifen			
Yes	80.3 (118)	91.0 (233)	0.003
No	19.7 (29)	9.0 (23)	

P-value tests for significant differences between the 'baseline' and 'baseline and 3-month' cohorts using χ^2 tests and t-tests.

Table S2. Sensitivity analysis: Knowledge about benefits and ≥ 2 harms of tamoxifen by participant characteristics and univariable and multivariable logistic regression model (n =317)

Knowledge about benefits and ≥ 2 harms of tamoxifen (%; N)		Univariable		Multivariable		
		OR (95% CI)	P-value	OR (95% CI)	P-value	
Age						
	≤ 35 years	25.6 (10)	0.78 (0.34 - 1.79)	0.556	0.73 (0.23 - 2.26)	0.579
	36 - 49 years	32.9 (80)	1.11 (0.67 - 1.83)	0.687	0.71 (0.37 - 1.36)	0.303
	≥ 50 years	30.7 (31)	Ref		Ref	
Children						
	Yes	33.1 (98)	1.38 (0.81 - 2.35)	0.240	1.61 (0.76 - 3.41)	0.219
	No	26.4 (23)	Ref		Ref	
Ethnic group*						
	White	31.2 (114)	-	-	-	-
	Other	42.9 (6)	-	-	-	-
Education level						
	Degree or above	43.3 (74)	2.67 (1.71 - 4.17)	≤ 0.001	2.95 (1.68 - 5.18)	≤ 0.001
	Below degree level	22.2 (46)	Ref		Ref	
Health status						
	Poor*	33.3 (5)	-	-	-	-
	Fair	16.2 (12)	0.39 (0.17 - 0.88)	0.023	0.47 (0.18 - 1.18)	0.108
	Good	36.2 (83)	1.14 (0.62 - 2.07)	0.675	1.14 (0.57 - 2.29)	0.713
	Excellent	33.3 (20)	Ref		Ref	
Risk level						
	Moderate	35.5 (81)	1.57 (1.00 - 2.47)	0.052	1.79 (1.03 - 3.14)	0.041
	High	26.0 (39)	Ref		Ref	
	Unclear*	20.0 (1)	-	-	-	-
SES						
	Low (most deprived)	28.6 (32)	0.77 (0.45 - 1.33)	0.354	1.06 (0.55 - 2.04)	0.860
	Middle	31.5 (40)	0.89 (0.53 - 1.49)	0.657	0.900 (0.48 - 1.70)	0.742
	High (least deprived)	34.1 (47)	Ref		Ref	
Employment						
	Full-time	31.0 (102)	Ref		Ref	
	All other employments	35.2 (19)	1.21 (0.66 - 2.21)	0.540	1.88 (0.91 - 3.90)	0.089
Marital status						
	Married or cohabiting	32.3 (91)	1.06 (0.65 - 1.75)	0.807	1.15 (0.59 - 2.26)	0.686
	Unmarried	30.9 (30)	Ref		Ref	
Numeracy						
	Good numeracy	34.0 (104)	1.99 (1.05 - 3.74)	0.034	2.43 (1.09 - 5.40)	0.030
	Poor numeracy	20.6 (14)	Ref		Ref	
Received a leaflet about tamoxifen						
	Yes	34.7 (95)	1.69 (1.02 - 2.81)	0.041	1.14 (0.61 - 2.12)	0.684
	No/unsure	23.9 (26)	Ref		Ref	
Satisfaction with healthcare professional						
	Satisfaction score, mean \pm SD	19.44 (3.44)	1.02 (0.96 - 1.08)	0.629	0.99 (0.93 - 1.07)	0.836
Felt informed about tamoxifen						
	Yes	34.5 (115)	3.69 (1.52 - 8.95)	0.004	3.73 (1.18 - 11.79)	0.025
	No	12.5 (6)	Ref		Ref	

*category not included in univariable and multivariable analyses due to insufficient cases.

⁺= This variable included the potential benefit of tamoxifen (a reduction in breast cancer risk) and at least two of the three potential harms of tamoxifen (increased risk of menopausal symptoms, blood clotting and endometrial cancer). In total, 31.6% (121) of women identified the benefit and ≥ 2 harms of tamoxifen.