

## Accepted Manuscript

Menopausal hormone therapy prior to the diagnosis of ovarian cancer is associated with improved survival

Katharine K. Brieger, Siri Peterson, Alice Lee, Bhramar Mukherjee, Kelly M. Bakulski, Aliya Alimujiang, Hoda Anton-Culver, Michael S. Anglesio, Elisa V. Bandera, Andrew Berchuck, David D.L. Bowtell, Georgia Chenevix-Trench, Kathleen R. Cho, Daniel W. Cramer, Anna DeFazio, Jennifer A. Doherty, Renée T. Fortner, Dale W. Garsed, Simon Gayther, Aleksandra Gentry-Maharaj, Ellen L. Goode, Marc T. Goodman, Holly R. Harris, Estrid Høgdall, David G. Huntsman, Hui Shen, Allan Jensen, Sharon E. Johnatty, Susan J. Jordan, Susanne K. Kjaer, Jolanta Kupryjanczyk, Diether Lambrechts, Karen McLean, Usha Menon, Francesmary Modugno, Kirsten Moysich, Roberta Ness, Susan J. Ramus, Jean Richardson, Harvey Risch, Mary Anne Rossing, Britton Trabert, Nicolas Wentzensen, Argyrios Ziogas, Kathryn L. Terry, Anna H. Wu, Gillian E. Hanley, Paul Pharoah, Penelope M. Webb, Malcolm C. Pike, Celeste Leigh Pearce

PII: S0090-8258(20)32296-4  
DOI: doi:[10.1016/j.ygyno.2020.06.481](https://doi.org/10.1016/j.ygyno.2020.06.481)  
Reference: YGYNO 978001

Published in: *Gynecologic Oncology*

Received date: 2 April 2020  
Accepted date: 7 June 2020

Cite this article as: Brieger KK, Peterson S, Lee A, Mukherjee B, Bakulski KM, Alimujiang A, Anton-Culver H, Anglesio MS, Bandera EV, Berchuck A, Bowtell DDL, Chenevix-Trench G, Cho KR, Cramer DW, DeFazio A, Doherty JA, Fortner RT, Garsed DW, Gayther S, Gentry-Maharaj A, Goode EL, Goodman MT, Harris HR, Hgdall E, Huntsman DG, Shen H, Jensen A, Johnatty SE, Jordan SJ, Kjaer SK, Kupryjanczyk J, Lambrechts D, McLean K, Menon U, Modugno F, Moysich K, Ness R, Ramus SJ, Richardson J, Risch H, Rossing MA, Trabert B, Wentzensen N, Ziogas A, Terry KL, Wu AH, Hanley GE, Pharoah P, Webb PM, Pike MC, Pearce CL, Menopausal hormone therapy prior to the diagnosis of ovarian cancer is associated with improved survival, *Gynecologic Oncology*, doi:[10.1016/j.ygyno.2020.06.481](https://doi.org/10.1016/j.ygyno.2020.06.481)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

## Menopausal hormone therapy prior to the diagnosis of ovarian cancer is associated with improved survival.

Katharine K. Brieger<sup>1</sup>, Siri Peterson<sup>1</sup>, Alice Lee<sup>2</sup>, Bhramar Mukherjee<sup>1,3</sup>, Kelly M. Bakulski<sup>1</sup>, Aliya Alimujiang<sup>1</sup>, Hoda Anton-Culver<sup>4</sup>, Michael S Anglesio<sup>5</sup>, Elisa V. Bandera<sup>6</sup>, Andrew Berchuck<sup>7</sup>, David D. L. Bowtell<sup>8,9</sup>, Georgia Chenevix-Trench<sup>10</sup>, Kathleen R. Cho<sup>11</sup>, Daniel W. Cramer<sup>12,13</sup>, Anna DeFazio<sup>14,15</sup>, Jennifer A. Doherty<sup>16</sup>, Renée T. Fortner<sup>17</sup>, Dale W. Garsed<sup>8,9</sup>, Simon Gayther<sup>18</sup>, Aleksandra Gentry-Maharaj<sup>19</sup>, Ellen L. Goode<sup>20</sup>, Marc T. Goodman<sup>21,22</sup>, Holly R. Harris<sup>23,24</sup>, Estrid Høgdal<sup>25,26</sup>, David G. Huntsman<sup>5,27</sup>, Hui Shen<sup>28</sup>, Allan Jensen<sup>25</sup>, Sharon E. Johnatty<sup>10</sup>, Susan J. Jordan<sup>29,30</sup>, Susanne K. Kjaer<sup>25,31</sup>, Jolanta Kupryjanczyk<sup>32</sup>, Diether Lambrechts<sup>33,34</sup>, Karen McLean<sup>35</sup>, Usha Menon<sup>19</sup>, Francesmary Modugno<sup>36,37,38</sup>, Kirsten Moysich<sup>39</sup>, Roberta Ness<sup>40</sup>, Susan J. Ramus<sup>41,42</sup>, Jean Richardson<sup>43</sup>, Harvey Risch<sup>44</sup>, Mary Anne Rossing<sup>23,24</sup>, Britton Trabert<sup>45</sup>, Nicolas Wentzensen<sup>45</sup>, Argyrios Ziogas<sup>4</sup>, Kathryn L. Terry<sup>12,13</sup>, Anna H. Wu<sup>46</sup>, Gillian E. Hanley<sup>47</sup>, Paul Pharoah<sup>48,49</sup>, Penelope M. Webb<sup>29,30,50</sup>, Malcolm C. Pike<sup>46,51</sup>, Celeste Leigh Pearce<sup>1,\*</sup>, for the Ovarian Cancer Association Consortium

<sup>1</sup>Department of Epidemiology, University of Michigan School of Public Health, Ann Arbor, MI, USA

<sup>2</sup>Department of Public Health, California State University Fullerton, Fullerton, CA, USA

<sup>3</sup>Department of Biostatistics, University of Michigan School of Public Health, Ann Arbor, MI, USA

<sup>4</sup>Department of Medicine, University of California Irvine, Irvine, CA, USA

<sup>5</sup>Department of Obstetrics and Gynecology, University of British Columbia, Vancouver, Canada

<sup>6</sup>Cancer Prevention and Control Program, Rutgers Cancer Institute of New Jersey, New Brunswick, NJ, USA

<sup>7</sup>Division of Gynecologic Oncology, Duke University School of Medicine, Durham, NC, USA

<sup>8</sup>Peter MacCallum Cancer Centre, Melbourne, Victoria, Australia

<sup>9</sup>Sir Peter MacCallum Department of Oncology, The University of Melbourne, Victoria, Australia

<sup>10</sup>Department of Genetics and Computational Biology, QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia

<sup>11</sup>Department of Pathology, University of Michigan Medical School, Ann Arbor, Michigan, USA

<sup>12</sup>Department of Epidemiology, Harvard TH Chan School of Public Health, Boston, MA, USA

<sup>13</sup>Obstetrics and Gynecology Epidemiology Center, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA

<sup>14</sup>Centre for Cancer Research, The Westmead Institute for Medical Research, The University of Sydney, Australia

<sup>15</sup>Department of Gynaecological Oncology, Westmead Hospital, Westmead, New South Wales, Australia

<sup>16</sup>Huntsman Cancer Institute, Department of Population Health Sciences, University of Utah, Salt Lake City, UT, USA

<sup>17</sup>Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany

<sup>18</sup>Cedars Sinai Medical Center, Los Angeles, CA, USA

<sup>19</sup>Women's Cancer, Institute for Women's Health, University College London, London, UK

- <sup>20</sup>Department of Health Science Research, Division of Epidemiology, Mayo Clinic, Rochester, MN, USA
- <sup>21</sup>Samuel Oschin Comprehensive Cancer Institute, Cancer Prevention and Genetics Program, Cedars-Sinai Medical Center, Los Angeles, CA, USA
- <sup>22</sup>Community and Population Health Research Institute, Department of Biomedical Sciences, Cedars-Sinai Medical Center, Los Angeles, CA, USA
- <sup>23</sup>Program in Epidemiology, Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA, USA
- <sup>24</sup>Department of Epidemiology, University of Washington, Seattle, WA, USA
- <sup>25</sup>Department of Virus, Lifestyle and Genes, Danish Cancer Society Research Center, Copenhagen, Denmark
- <sup>26</sup>Molecular Unit, Department of Pathology, Herlev Hospital, University of Copenhagen, Copenhagen, Denmark
- <sup>27</sup>Department of Molecular Oncology, BC Cancer Research Centre, Vancouver, Canada
- <sup>28</sup>Van Andel Research Institute (VARI), Grand Rapids, MI, USA
- <sup>29</sup>University of Queensland, School of Public Health, Brisbane, Australia
- <sup>30</sup>Department of Population Health, QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia
- <sup>31</sup>Department of Gynaecology, Rigshospitalet, University of Copenhagen, Copenhagen, Denmark
- <sup>32</sup>Department of Pathology and Laboratory Diagnostics, Maria Sklodowska-Curie National Research Institute of Oncology, Warsaw, Poland
- <sup>33</sup>Vesalius Research Center, VIB, Leuven, Belgium
- <sup>34</sup>Laboratory for Translational Genetics, Department of Oncology, University of Leuven, Leuven, Belgium
- <sup>35</sup>Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, University of Michigan Medical School, Ann Arbor, Michigan, USA
- <sup>36</sup>Womens Cancer Research Center. Magee-Women's Research Institute and Hillman Cancer Center, Pittsburgh, PA, USA
- <sup>37</sup>Division of Gynecologic Oncology, Department of Obstetrics, Gynecology and Reproductive Sciences, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA
- <sup>38</sup>Department of Epidemiology, University of Pittsburgh Graduate School of Public Health
- <sup>39</sup>Division of Cancer Prevention and Control, Roswell Park Comprehensive Cancer Center. Buffalo, NY, USA
- <sup>40</sup>School of Public Health, University of Texas Health Science Center at Houston (UTHealth), TX, USA
- <sup>41</sup>School of Women's and Children's Health, Faculty of Medicine, University of NSW Sydney, Sydney, New South Wales, Australia
- <sup>42</sup>The Kinghorn Cancer Centre, Garvan Institute of Medical Research, Sydney, New South Wales, Australia
- <sup>43</sup>Department of Preventive Medicine, University of Southern California, Los Angeles, CA, USA
- <sup>44</sup>Chronic Disease Epidemiology, Yale School of Public Health, New Haven, CT, USA
- <sup>45</sup>Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD, USA
- <sup>46</sup>Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

<sup>47</sup>University of British Columbia Faculty of Medicine, Department of Obstetrics & Gynecology, Vancouver, Canada

<sup>48</sup>Centre for Cancer Genetic Epidemiology, Department of Oncology, University of Cambridge, Cambridge, UK

<sup>49</sup>Centre for Cancer Genetic Epidemiology, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK

<sup>50</sup>Gynaecological Cancers Group, QIMR Berghofer Medical Research Institute, 300 Herston Road, Brisbane, QLD 4006 Australia

<sup>51</sup>Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, NY, USA

\*Corresponding author at: 4642 SPH Tower, 1415 Washington Heights, Ann Arbor, Michigan 48109-2029.

**ABSTRACT**

**Purpose:** Prior studies of menopausal hormone therapy (MHT) and ovarian cancer survival have been limited by lack of hormone regimen detail and insufficient sample sizes. To address these limitations, a comprehensive analysis of 6,419 post-menopausal women with pathologically confirmed ovarian carcinoma was conducted to examine the association between MHT use prior to diagnosis and survival.

**Methods:** Data from 15 studies in the Ovarian Cancer Association Consortium were included. MHT use was examined by type (estrogen-only (ET) or estrogen+progestin (EPT)), duration, and recency of use relative to diagnosis. Cox proportional hazards models were used to estimate the association between hormone therapy use and survival. Logistic regression and mediation analysis was used to explore the relationship between MHT use and residual disease following debulking surgery.

**Results:** Use of ET or EPT for at least five years prior to diagnosis was associated with better ovarian cancer survival (hazard ratio, 0.80; 95% CI, 0.74 to 0.87). Among women with advanced stage, high-grade serous carcinoma, those who used MHT were less likely to have any macroscopic residual disease at the time of primary debulking surgery (p for trend <0.01 for duration of MHT use). Residual disease mediated some (17%) of the relationship between MHT and survival.

**Conclusions:** Pre-diagnosis MHT use for 5+ years was a favorable prognostic factor for women with ovarian cancer. This large study is consistent with prior smaller studies, and further work is needed to understand the underlying mechanism.

## INTRODUCTION

Invasive epithelial ovarian cancers including ovarian, fallopian tube and primary peritoneal cancer (hereafter referred to as ovarian cancer) collectively account for more deaths than any other cancer of the female reproductive system in the United States, with a five-year survival rate of less than 50%<sup>1</sup>. There is clear evidence that menopausal estrogen-alone hormone therapy (ET) is associated with an increased risk of developing ovarian cancer<sup>2-4</sup>, but the relationship between menopausal estrogen plus progestin therapy (EPT) and risk of ovarian cancer is less clear<sup>3</sup>. Further, the relationship between menopausal hormone therapy (MHT) use and survival may not be the same as the relationship with risk.

Pre-diagnosis MHT use and ovarian cancer survival has been examined in nine population-based studies<sup>5-13</sup>. Most observed a modestly inverse association, with hazard ratios ranging from 0.23<sup>11</sup> to 1.1<sup>12</sup> (**Table S1**), but protection was statistically significant in only one study (MHT use >5 years: HR, 0.79; 95% CI, 0.55 to 0.90)<sup>5</sup>. These studies were subject to one or more of the following important limitations: they (1) lacked information about duration of use; (2) did not distinguish between types of MHT use before diagnosis (i.e., ET and/or EPT); (3) had follow-up times of only a few years; (4) had an insufficient sample size to stratify by ovarian cancer histotype; and (5) lacked information about residual disease after debulking surgery. Many women use MHT for only a short period of time, thus missing duration information is an important weakness that may have masked effects in prior studies<sup>14</sup>. Rigorously evaluating the association between pre-diagnosis MHT use and ovarian cancer survival by hormone type, duration, survival time, residual disease and cancer histotype is essential to advance our understanding of disease prognosis.



In the present analysis from the Ovarian Cancer Association Consortium (OCAC), we followed 6,419 women with ovarian cancer for up to 26 years and investigated the association between pre-diagnosis MHT use and survival. We investigated duration, type and timing of MHT use in each of the main histological subtypes. A particularly important prognostic factor in ovarian cancer survival is residual disease after initial debulking surgery. Therefore, we also considered the potential relationship of MHT use with residual disease after surgery.

## METHODS

Institutional Review Board or comparable ethics approval was received by each study and informed consent was provided by all women.

### *Study populations and exclusion criteria*

OCAC is an international, multidisciplinary collaboration of ovarian cancer research teams (<http://ocac.ccge.medschl.cam.ac.uk/>). Post-menopausal women (as defined in each study) with pathologically confirmed ovarian carcinoma and survival time available (n=10,120) were considered for our analyses. We were interested *a priori* in the potential of a duration effect of MHT use and thus three studies (n=2121) were excluded. Therefore, this analysis used pooled ovarian cancer survival data from population-based (n=14) and clinic-based (n=1) OCAC studies (**Table S2**) conducted in the United States (n=9), Europe (n=4), and Australia (n=2). Women from these studies missing MHT duration were excluded. Only those with invasive tumors, high-grade serous, low-grade serous, mucinous, endometrioid, or clear cell carcinomas, were eligible (i.e. mixed cell, undifferentiated, and non-epithelial cancers were excluded; n=1,260). Women missing data for stage at diagnosis (n=282), race/ethnicity (n=25), or time from diagnosis to interview/study enrolment (n=13) were also excluded. There was no upper or lower age limit exclusion beyond the impact of excluding women who were pre-menopausal at

diagnosis. Our final analytic sample was 6,419 ovarian cancer patients. (**Figure 1**). Survival times and proportion of deaths were comparable between women excluded and those included.

#### *Exposure and covariate assessment*

Participants provided information on their history of MHT use prior to diagnosis via phone or in-person interviews (n=10 study sites) or self-completed questionnaires (n=5 study sites) (**Table S2**) at the time of study enrollment. MHT use was categorized as exclusive use of ET, exclusive use of EPT, use of both therapies, or use of unknown type. First, exclusive ET use was examined based on (1) total duration of ET use (never (reference category), >0 to <5, 5 to <10, 10+ years) and (2) recency of ET use (within the year prior to diagnosis, 1 to <5, 5+ years prior to diagnosis). There was no additional duration effect observed after 5 years and so the categories 5 to <10 and 10+ years were combined into one. Exclusive EPT use was examined in the same manner. The reference group for both the ET and EPT analyses was never use of any type of MHT. Next, total duration of any type of MHT use prior to diagnosis was examined (ET, EPT, both, or unknown type) with the same approach. BMI (kg/m<sup>2</sup>) categories were assigned according to World Health Organization<sup>15</sup> definitions (underweight, BMI<18.5; normal weight, 18.5≤BMI<25.0; overweight, 25≤BMI<30; obese, BMI≥30), using the values reported for adult BMI one to five years prior to diagnosis. Duration of combined oral contraceptive use was coded as never, <1, 1 to <5, 5 to <10, or 10+ years. Parity was coded as 0, 1, or 2+ pregnancies. Education level was coded as less than high school, high school graduate, some college, college graduate, or graduate school. Stage was recorded as local (with no lymph node involvement), regional (direct extension and/or local lymph node involvement), and advanced (distant sites and/or distant lymph nodes involved)<sup>16</sup>. For all patients, the standard of care is assumed to have been a platinum-based regimen.

### *Outcome assessment*

Overall survival was recorded as either length of time (in days) from diagnosis to death or to date of last follow-up (for censored patients). Follow-up is largely done via linkage with national death databases. All survival models incorporated left truncation time, accounting for the difference between date of diagnosis and date of patient interview, though there was little variability in delay to patient interview and so accounting for left truncation did not affect results. Women were typically enrolled within the first sixth months following diagnosis (median 152 days). For a subset of women there was information on duration of progression-free survival (n=2,239) and presence/absence and size of residual disease after debulking surgery (n=2,056) (**Table S2**).

### *Statistical analysis*

#### Overall survival models

Cox proportional hazards models with left truncation and right censoring were used to estimate the association (hazard ratio; HR, and associated Wald-type confidence intervals) of each hormone therapy exposure on ovarian cancer survival. The exposures were modeled as categories of duration of use and recency of use, as detailed above. Exclusive use of ET or exclusive use of EPT were first examined separately to determine their association with survival. Because the hazards for the two types of hormone therapies were not statistically different and showed a similar magnitude, we combined types as an “any HT use” variable including unknown types of MHT, as these would have been either ET or EPT.

Important *a priori* variables included in all models were age at diagnosis (continuous), race/ethnicity, surgical stage at diagnosis, and OCAC study site. Sensitivity analyses adjusting

for age in five- and ten-year categories did not materially alter the HR estimates for MHT. Education level adjustment in sensitivity analyses also did not influence HR estimates. The possible confounding effects of additional exposures prior to diagnosis were examined, but none affected the association between MHT duration and survival (**Table S5**).

Separate models for each histotype were also fitted to estimate HRs for MHT duration. The adjusted survival curves presented (overall, and high-grade serous) allow for visualization of survival curves based on the Cox proportional hazard results.

#### Discrete windows of clinical interest and progression-free survival

We tested discrete windows of clinical interest following diagnosis. Although the proportional hazards assumptions were not violated for MHT use prior to diagnosis in the Cox proportional hazards models, an additional model was fit allowing the data to be split into time intervals after diagnosis. This allowed us to assess subtle variation in HR estimates at all time points after diagnosis. To assess the specificity of the protective effect of MHT, Cox proportional hazards model was fit for time to progression, treating progression of disease as the event of interest. Although ovarian cancer-specific mortality was not assessed, nearly all deaths within the first five years following diagnosis are related to ovarian cancer thus our time interval analysis provides insight into this question.

#### Residual disease in women with advanced stage, high-grade serous cancer

Among women with advanced stage (stage III or IV), high-grade serous carcinoma (HGSC, n=903), we examined possible mechanisms underlying the MHT-survival association, namely the association of MHT use with residual disease. We used logistic regression, investigating MHT use in those with and without macroscopic residual disease following primary debulking surgery. Mediation analysis was used to examine whether the relationship

between MHT use and survival was mediated by residual disease. In this analysis, the first step (mediator) model was residual disease regressed on MHT use and the covariates age, stage, histotype, education level, and race. The second step (outcome) model was modeled as survival regression on residual disease, MHT use, and the same set of covariates. Finally, mediation was assessed using 2,000 simulations to estimate the average causal mediated effect, the average causal direct effect, the total effect, and the proportion mediated, using the generalizable approach to causal mediation outlined by Imai et al.<sup>17</sup>. All statistical analysis was performed in *R*<sup>18</sup>.

## RESULTS

The analytic sample included 6,419 post-menopausal women from 15 sites in the OCAC (**Figure S1; Table S2**). A majority of the women had HGSC (68.4%) and most had advanced stage disease at diagnosis (67.7%; **Table 1**). Exclusive EPT use (18.5%) was more common than exclusive ET use (14.2%). Most women (58.9%) did not use either type and 212 (3.3%) used both ET and EPT (**Table 1**).

The median survival time was 5.4 years after diagnosis. ET and EPT use for at least five years were both associated with longer survival (**Table 2**). For exclusive ET users, lower mortality was observed for use of 5+ years (HR, 0.85; 95% CI, 0.75 to 0.96). For exclusive EPT users, the HR for use for 5+ years was similar (HR, 0.79; 95% CI, 0.70 to 0.89). Because the magnitudes of effect for ET and EPT were similar, all MHT types were combined for subsequent analyses. Significantly better survival was observed for those who had used any type of MHT for at least 5 years (HR, 0.80; 95% CI, 0.74 to 0.87) (**Table 2**). This corresponds to a median

survival of 5.75 years among women who had used MHT for 5+ years and 4.6 years for those who has not used any.

An adjusted survival curve illustrates the apparent protective benefit of MHT use was restricted to women with 5+ years use compared to those who did not use MHT and that no benefit was observed for <5 years of use (**Figure 1**). Recency of MHT use did not affect the hazard ratio estimates. The association observed for all histotypes combined was also similar for individual histotypes, with the exception of endometrioid carcinomas, but was only statistically significant for HGSC (HR, 0.78; 95% CI, 0.71 to 0.86) (**Table S3a**). Progression-free survival (time from diagnosis to first recurrence, documented by clinical, biochemical (e.g. serum CA125 levels) or radiological disease progression) was also better in those who had used MHT (**Table S4**).

Time-varying HRs were also estimated. Although the proportional hazards assumptions were not violated for the survival model of MHT use, the additional analyses allowed for finer estimation of the protective association during particular windows of interest after diagnosis. The estimated effect was protective in all time intervals. MHT use was associated with reduced risk of death significantly in the first two years after diagnosis (HR, 0.72; 95% CI, 0.62 to 0.84) and in years 2 through 5 after diagnosis (HR, 0.86; 95% CI, 0.76 to 0.97) (**Figure 2**).

Stratification by stage at diagnosis for HGSC showed a positive association with prognosis at advanced stages (III/IV) (**Table S3b**). Among women with advanced stage HGSC, MHT use was associated with improved survival both in the women with and those without residual disease (**Figure S2** and **Table S6**). MHT use prior to diagnosis was associated with lower likelihood of residual disease at the time of debulking surgery among women with advanced stage HGSC. Of women with local (stage I, n=180) and regional (stage II, n=343)

disease, only 2 women (2%) and 18 women (5.2%) respectively had residual disease after surgery, thus we cannot estimate ORs for MHT use in these strata. Among those with advanced disease (stage III/IV), MHT use was associated with significantly lower odds of having macroscopic residual disease relative to no macroscopic disease in an MHT duration-dependent manner ( $p$  for trend = 0.009), adjusted for age at diagnosis (**Table 3**). Adjusting for OCAC site and race/ethnicity did not alter the trend. Residual disease partially mediated the relationship between long-term (5+ years) MHT use and survival. Among women with advanced HGSC, the proportion mediated was 0.17 ( $p=0.04$ ).

## DISCUSSION

In this study, pre-diagnosis MHT use for at least five years was associated with better ovarian cancer survival, regardless of MHT type (ET or EPT) and recency of use relative to diagnosis. Other studies reported ever use of MHT to be associated with improved survival (**Table S1**), but this is the first study to report on the effect of duration and recency of MHT use, type of MHT use, histotype, and residual disease after debulking surgery on survival outcomes.

Women with advanced HGSC who had used MHT prior to diagnosis were less likely to have macroscopic disease following primary debulking surgery. We estimated that about 17% of the survival improvement associated with MHT use could be due to the higher proportion of MHT users with no residual disease. The mechanism of the effect of MHT on residual disease is unclear. At least one previous study has noted that MHT use was associated with optimal debulking status<sup>19</sup>. One possibility is that MHT use prior to diagnosis alters the pattern of metastatic spread, such that the disease is easier to access or less adhesive to surrounding tissues. It has been reported that tumor tissue from sub-optimally debulked patients expressed molecular signatures consistent with increased stromal activation and lymphovascular invasion<sup>20</sup>. A

predictive gene expression signature, developed for likelihood of optimal debulking, suggested that there may be a subset of tumors for which the TGF- $\beta$  activated pathway stimulates epithelial to mesenchymal transition and activation of tumor associated fibroblasts<sup>21</sup>, both of which would contribute to spread of tumor and difficulty in debulking.

Prior studies have established a complex relationship between hormonal exposures, including hormone therapy<sup>22,23</sup>, and inflammation that depends on multiple factors including the formula, dose, route of delivery, and other immune stimuli. MHT use may result in an anti-inflammatory milieu that is beneficial for resection. Particularly at high concentrations, estrogen has anti-inflammatory properties<sup>24,25</sup> in some tissues. Furthermore, evidence supports a mutually dependent relationship between inflammation and angiogenesis<sup>26</sup>. Immune cells stimulated during inflammatory reactions secrete cytokines such as IL-6, TNF- $\alpha$  and CXCR2 that promote neovascularization and thus potentially contribute to tumor establishment and growth. On the other hand, an anti-inflammatory environment would prevent this sequence. Mechanistic studies are needed to understand the relationship between MHT use and ease of debulking. Mechanistic studies are also needed to investigate whether it is primarily women with estrogen-receptor negative cancers who are driving the protective association with MHT use; indeed, the current literature suggests avoiding MHT in women with estrogen-sensitive histologic subtypes<sup>27</sup>. This may explain why the endometrioid subtype findings deviated from the other histotypes.

Pre-diagnosis use, as previously discussed and as demonstrated by this current study, appears to offer a survival benefit to women with ovarian cancer<sup>5,7-9,11,13,19</sup> (**Table S1**). The existing literature on post-diagnosis MHT use and ovarian cancer survival includes several population-based cohort studies<sup>6,8,28-30</sup> and two small randomized controlled trials<sup>31,32</sup>. The



population-based studies were largely inconclusive, but they all suggest reduced mortality in post-diagnosis MHT users<sup>6,8,28-30,33,34</sup>.

Two randomized trials have indicated survival benefits of hormone therapy use<sup>31,32</sup> after surgical debulking of the ovarian tumor. A clinical trial in 1999<sup>32</sup> randomized women with ovarian cancer of any histotype to either conjugated estrogen or to no supplementation after debulking surgery. The women who received estrogen therapy had non-significantly longer disease-free intervals and better overall survival. In a second study, Eeles et al.<sup>31</sup> randomized women who had been diagnosed with ovarian cancer within the previous 9 months to receive hormone therapy or none. The study observed a statistically significant beneficial effect of hormone therapy on overall survival (HR, 0.63; 95% CI, 0.44 to 0.90), but this likely reflects some of the general benefits of MHT on survival as this is not an ovarian cancer-specific survival estimate. However, no specific hormonal regimen was used, as individual clinicians had control over type, dose and duration.

Limitations of our results include the self-reported exposure measures. However, prior studies have documented good correlation between self-report of hormone use and prescription records<sup>35</sup>. Although our analysis was restricted to women who were classified as post-menopausal at diagnosis, some may have used MHT before menopause occurred. To address this issue, we conducted a sensitivity analysis restricting the exposure to MHT use after the age of 50, as a proxy for post-menopausal use, and the results did not change. An additional limitation was the lack of information on MHT use post-diagnosis. We cannot exclude the possibility that pre-diagnosis use predicts post-diagnosis use, conferring part of the survival benefit. Likewise, there was not a large enough sample of women with chemotherapy information to conduct analyses on any differential effects of MHT based on chemotherapy treatment. However,

regardless of prior exposures and medical history, the standard of care for the vast majority of women is a platinum-based regimen. Additionally, we assumed that the chemotherapy ordered for women who had been on MHT was comparable to that for women who were never on MHT; however, because treatment data were not available, we cannot rule out that women who were previously on MHT were better able to tolerate the full dose of chemotherapy. Finally, use of MHT could serve as a proxy for overall adherence to medical recommendations and treatment and access to specialist surgical practices. However, controlling for education, which was expected to correlate with these characteristics, did not affect results.

We observed that the association of MHT use for five or more years prior to diagnosis was protective against death at all points after diagnosis. Since the cause of death during the first five years is most commonly ovarian cancer-specific and not from other causes, this suggests that the association of survival with MHT use is at least in part due to cancer-specific protection. We also offer evidence that the relationship is partially mediated by the relationship between MHT use and optimal surgical cytoreduction.

The findings presented here, taken in context with the other literature on the topic (**Table S1**), suggest that MHT is beneficial with respect to ovarian cancer survival, particularly among women with HGSC. These findings are helpful to understand the biology of the disease, and ultimately our goal is to help women diagnosed with ovarian cancer to live both longer and with a higher quality of life. Post-menopausal symptoms, including severe vasomotor symptoms for some women, can negatively impact quality of life. It is well known that an early onset of menopause increases risks for overall mortality, particularly cardiovascular mortality, and thus ovarian cancer patients who undergo surgical menopause prior to natural menopause would likely benefit from MHT<sup>36</sup>. Whether MHT is cardioprotective in women who are

postmenopausal depends on timing of initiation in relation to onset of menopause, with women initiating MHT within 4-6 years of onset of menopause showing decreased risk for CVD. Thus, there are many ovarian cancer patients who would receive cardioprotective effects from MHT<sup>37,38</sup>. There are also important benefits of MHT use in postmenopausal women in terms of reduced risk of hip fracture<sup>39</sup> and reduced risk for colorectal cancer<sup>40</sup>. Therefore, clinician and patient confidence in using MHT offers great potential benefit to women with ovarian cancer. A large randomized clinical trial would help determine the impact of MHT on survival and quality of life for women living with ovarian cancer. Such a future trial could incorporate detailed mechanistic studies to better understand how MHT influences survival. Despite remaining questions, the current evidence should allow providers to at least discuss MHT use with ovarian cancer patients, with shared decision making regarding the benefits and limitations of therapy.

#### **Conflict of Interest Statement**

The authors declare no potential conflicts of interest.

#### **OCAC Funding**

The Ovarian Cancer Association Consortium is supported by a grant from the Ovarian Cancer Research Fund thanks to donations by the family and friends of Kathryn Sladek Smith (PPD/RPCI.07).

Funding for individual studies: **AUS:** The Australian Ovarian Cancer Study (AOCS) was supported by the U.S. Army Medical Research and Materiel Command (DAMD17-01-1-0729), National Health & Medical Research Council of Australia (199600, 400413 and 400281), Cancer Councils of New South Wales, Victoria, Queensland, South Australia and Tasmania and Cancer Foundation of Western Australia (Multi-State Applications 191, 211 and 182). AOCS gratefully

acknowledges additional support from Ovarian Cancer Australia and the Peter MacCallum Foundation; **CON:** National Institutes of Health (R01-CA063678, R01-CA074850; R01-CA080742); **DOV:** National Institutes of Health R01-CA112523 and R01-CA87538; **GER:** German Federal Ministry of Education and Research, Programme of Clinical Biomedical Research (01 GB 9401) and the German Cancer Research Center (DKFZ); **HAW:** U.S. National Institutes of Health (R01-CA58598, N01-CN-55424 and N01-PC-67001); **HOP:** University of Pittsburgh School of Medicine Dean's Faculty Advancement Award (F. Modugno), Department of Defense (DAMD17-02-1-0669) and NCI (K07-CA080668, R01-CA95023, P50-CA159981 MO1-RR000056 R01-CA126841); **MAL:** Funding for this study was provided by research grant R01-CA61107 from the National Cancer Institute, Bethesda, MD, research grant 94 222 52 from the Danish Cancer Society, Copenhagen, Denmark; and the Mermaid I project; **MAY:** National Institutes of Health (R01-CA122443, P30-CA15083, P50-CA136393); Mayo Foundation; Minnesota Ovarian Cancer Alliance; Fred C. and Katherine B. Andersen Foundation; **NEC:** National Institutes of Health R01-CA54419 and P50-CA105009 and Department of Defense W81XWH-10-1-02802; **NJO:** National Cancer Institute (NIH-K07 CA095666, R01-CA83918, NIH-K22-CA138563, and P30-CA072720) and the Cancer Institute of New Jersey; If Sara Olson and/or Irene Orlow is a co-author, please add NCI CCSG award (P30-CA008748) to the funding sources; **OPL:** National Health and Medical Research Council (NHMRC) of Australia (APP1025142) and Brisbane Women's Club; **POL:** Intramural Research Program of the National Cancer Institute; **UCI:** NIH R01-CA058860 and the Lon V Smith Foundation grant LVS-39420; **UKO:** The UKOPS study was funded by The Eve Appeal (The Oak Foundation) with investigators supported by the National Institute for Health Research University College London Hospitals Biomedical Research Centre and MRC core funding (MR\_UU\_12023); **USC:**

P01CA17054, P30CA14089, R01CA61132, N01PC67010, R03CA113148, R03CA115195, N01CN025403, and California Cancer Research Program (00-01389V-20170, 2II0200);

### **Acknowledgements**

We are grateful to the family and friends of Kathryn Sladek Smith for their generous support of the Ovarian Cancer Association Consortium through their donations to the Ovarian Cancer Research Fund. We thank the study participants, doctors, nurses, clinical and scientific collaborators, health care providers and health information sources who have contributed to the many studies contributing to this manuscript.

Acknowledgements for individual studies: **AUS:** The AOCS also acknowledges the cooperation of the participating institutions in Australia, and the contribution of the study nurses, research assistants and all clinical and scientific collaborators. The complete AOCS Study Group can be found at [www.aocstudy.org](http://www.aocstudy.org). We would like to thank all of the women who participated in this research program; **CON:** The cooperation of the 32 Connecticut hospitals, including Stamford Hospital, in allowing patient access, is gratefully acknowledged. This study was approved by the State of Connecticut Department of Public Health Human Investigation Committee. Certain data used in this study were obtained from the Connecticut Tumor Registry in the Connecticut Department of Public Health. The authors assume full responsibility for analyses and interpretation of these data; **GER:** The German Ovarian Cancer Study (GER) thank Ursula Eilber for competent technical assistance; **OPL:** Members of the OPAL Study Group (<http://opalstudy.qimrberghofer.edu.au/>); **UKO:** We particularly thank I. Jacobs, M. Widschwendter, E. Wozniak, A. Ryan, J. Ford and N. Balogun for their contribution to the study.

**Table 1:** Demographic and clinical characteristics of women with ovarian cancer from the Ovarian Cancer Association Consortium (OCAC) included in the survival analysis.

N	Pre-diagnosis MHT use duration			
	Overall <sup>1</sup> 6419	Never 3784	<5 years 1183	5+ years 1452
<b>Hormone therapy use (%)</b>				
None	3784 (58.9)	3784 (100.0)	0 (0.0)	0 (0.0)
ET only	909 (14.2)	0 (0.0)	379 (32.0)	530 (36.5)
EPT only	1188 (18.5)	0 (0.0)	561 (47.4)	627 (43.2)
ET and EPT	212 (3.3)	0 (0.0)	62 (5.2)	150 (10.3)
Unknown +/- ET/EPT	326 (5.1)	0 (0.0)	181 (15.3)	145 (10.0)
<b>Age at dx. (mean (SD))</b>	62.67 (8.71)	62.36 (9.33)	60.78 (8.16)	65.00 (6.75)
<b>Education (%)</b>				
Less than high school	1135 (20.7)	760 (23.7)	177 (17.4)	198 (15.9)
High school graduate	1567 (28.6)	948 (29.6)	272 (26.7)	347 (27.8)
Some college	1325 (24.2)	745 (23.2)	265 (26.0)	315 (25.3)
College graduate	799 (14.6)	400 (12.5)	174 (17.1)	225 (18.1)
Graduate school	646 (11.8)	353 (11.0)	132 (12.9)	161 (12.9)
<b>Race / ethnicity (%)</b>				
Non-Hispanic white	5679 (88.5)	3308 (87.4)	1042 (88.1)	1329 (91.5)
Hispanic white	198 (3.1)	126 (3.3)	45 (3.8)	27 (1.9)
Black	101 (1.6)	72 (1.9)	15 (1.3)	14 (1.0)
Asian	249 (3.9)	146 (3.9)	51 (4.3)	52 (3.6)
Other	192 (3.0)	132 (3.5)	30 (2.5)	30 (2.1)
<b>Histotype (%)</b>				
Low-grade serous	245 (3.8)	134 (3.5)	47 (4.0)	64 (4.4)
High-grade serous	4393 (68.4)	2504 (66.2)	820 (69.3)	1069 (73.6)
Mucinous	373 (5.8)	255 (6.7)	65 (5.5)	53 (3.7)
Endometrioid	925 (14.4)	552 (14.6)	168 (14.2)	205 (14.1)
Clear cell	483 (7.5)	339 (9.0)	83 (7.0)	61 (4.2)
<b>Stage (%)</b>				
Local (FIGO I)	947 (14.8)	616 (16.3)	173 (14.6)	158 (10.9)
Regional (FIGO II)	1126 (17.5)	684 (18.1)	211 (17.8)	231 (15.9)
Advanced (FIGO III/IV)	4346 (67.7)	2484 (65.6)	799 (67.5)	1063 (73.2)
<b>BMI category (%)</b>				
Underweight	117 (2.0)	71 (2.1)	18 (1.6)	28 (2.0)
Normal weight	2684 (45.7)	1424 (42.0)	515 (45.9)	745 (54.5)
Overweight	1754 (29.9)	1026 (30.3)	339 (30.2)	389 (28.5)
Obese	1320 (22.5)	866 (25.6)	249 (22.2)	205 (15.0)
<b>Family<sup>2</sup> cancer history (%)</b>				
Breast cancer	1098 (17.6)	690 (18.9)	195 (16.8)	213 (15.0)
Ovarian cancer	329 (5.3)	203 (5.6)	61 (5.3)	65 (4.6)

<b>Combined oral contraceptive use (%)</b>				
Never	3127 (49.2)	2030 (54.2)	451 (38.6)	646 (44.9)
<1 year	590 (9.3)	313 (8.4)	142 (12.1)	135 (9.4)
1 to <5 years	1209 (19.0)	659 (17.6)	265 (22.7)	285 (19.8)
5 to <10 years	773 (12.2)	390 (10.4)	176 (15.1)	207 (14.4)
10+ years	656 (10.3)	356 (9.5)	135 (11.5)	165 (11.5)
<b>Parity (%)</b>				
0 births	1223 (19.1)	738 (19.6)	232 (19.6)	253 (17.4)
1 birth	858 (13.4)	525 (13.9)	157 (13.3)	176 (12.1)
2+ births	4324 (67.5)	2508 (66.5)	794 (67.1)	1022 (70.4)
<b>Smoking (%)</b>				
Never	2910 (52.9)	1803 (55.5)	495 (48.8)	612 (49.4)
Current	700 (12.7)	445 (13.7)	126 (12.4)	129 (10.4)
Former	1891 (34.4)	998 (30.7)	394 (38.8)	499 (40.2)

<sup>1</sup> The total N for certain variables reported does not total to 6,419 because of missing data. These included variables that were not confounders and thus not needed for covariate adjustment in final models, such as family history of cancer, education, and smoking.

<sup>2</sup> First-degree family members, i.e. sister or mother.

**Table 2:** Hazards ratios for menopausal hormone therapy (MHT) use before diagnosis of ovarian cancer among women with invasive epithelial ovarian cancer in the Ovarian Cancer Association Consortium (OCAC).

MHT use	Estrogen alone (ET)		Estrogen-progestin combined therapy (EPT)		Any menopausal hormone therapy	
	N <sup>a</sup>	HR (95% CI) <sup>b</sup>	N	HR (95% CI)	N	HR (95% CI)
None (ref)	3,784	1.0	3,784	1.0	3,784	1.0
<5 years	379	0.99 (0.86, 1.15)	561	1.01 (0.89, 1.14)	1,183	0.97 (0.88, 1.06)
5+ years	530	0.85 (0.75, 0.96)	627	0.79 (0.70, 0.89)	1,452	0.80 (0.74, 0.87)

<sup>a</sup> The three analyses have different total N's because the exclusive ET analysis excluded women who had ever used EPT, and the exclusive EPT analysis excluded women who had ever used ET. Users of unknown type were also excluded from this analysis.


<sup>b</sup> Hazard ratios (HRs) are adjusted for age at diagnosis and race/ethnicity, and stratified by histotype, stage at diagnosis, and OCAC site.

**Table 3:** Odds ratios for macroscopic residual disease based on use of menopausal hormone therapy (MHT) use before diagnosis of ovarian cancer, among women with advanced, high-grade serous carcinoma in the Ovarian Cancer Association Consortium (OCAC).


MHT use	N	Residual disease	OR <sup>a</sup> (95% CI)	p for trend
None (ref)	859	574 (66%)	1.0	
<5 years	239	146 (61%)	0.79 (0.58, 1.06)	
5+ years	290	171 (59%)	0.71 (0.54, 0.93)	0.009

<sup>a</sup>ORs are adjusted for age at diagnosis. Adjusting for OCAC site and race/ethnicity did not alter the trend for inverse association.





**Figure 1: Overall survival stratified by years of menopausal hormone therapy use.** Adjusted survival curves among all women with ovarian cancer (n=6,419) and among women with advanced stage, high-grade serous cancer (n=4,393). The adjusted survival curves are generated from the hazard ratios estimated from a cox proportional hazards model of menopausal hormone therapy use and are adjusted for age at diagnosis, race/ethnicity, histotype (left panel only), stage at diagnosis, and OCAC site.



**Figure 2: Estimated time-varying hazard ratio (HR) and 95% confidence intervals (CIs) for use of menopausal hormone therapy (5+ years) relative to no use.** In a Cox proportional hazard model allowing for interaction of the effect of menopausal hormone therapy use with time since diagnosis, the estimated effect is protective at all time points. Menopausal hormone therapy use is significantly protective in the first two years after diagnosis (HR = 0.72; 95% CI = 0.62, 0.84) and in years 2 through 5 after diagnosis (HR = 0.86; 95% CI = 0.76, 0.97).

## Supplemental Appendix

## Supplemental Tables

**Table S1:** Published findings of association between menopausal hormone therapy use and ovarian cancer survival (Online only).

Author, Year	HR (95% CI) overall survival for any HT use	Type	Study design	N, population	Notes
Pre-diagnosis					
Mascarenhas, 2006 <sup>[5]</sup>	0.83 (0.65, 1.08)	Any	Cohort, follow-up on population-based cases	649, Sweden	Examination of different HRT preparations did not give different HRs
Wernli, 2008 <sup>[6]</sup>	1.1 (0.85, 1.43)	Any	Case-control, population based	751, United States	
Nagle, 2008 <sup>[7]</sup>	0.83 (0.64, 1.08)	Any	Cohort, population-based	676, Australia	
Zhang, 2012 <sup>[8]</sup>	0.23 (0.03, 1.73)	Any	Cohort	195, China	
Hein, 2013 <sup>[9]</sup>	75% (65, 86) 5-yr survival for HRT users; 43% (36, 52) for non-users	Any	Cohort	244, Bavaria	Women who had used HRT: younger, lower stage, more optimal debulking
Shafir, 2016 <sup>[10]</sup>	0.70 (0.55, 0.9)	Any	Cohort	1649, United States	
Kim, 2017 <sup>[11]</sup>	0.79 (0.62, 1.01)	Any	Cohort, registry-based	1421, Canada	
Besevic, 2015 <sup>[12]</sup>	0.80 (0.55, 1.16)	EPT	Cohort	1025, Europe	
	0.86 (0.54, 1.35)	ET			
Felix, 2015 <sup>[13]</sup>	0.97 (0.68, 1.38)	EPT	Cohort, population-based	396, United States	Noted interaction between HRT type and histology
	1.09 (0.7, 1.68)	ET			
Post-diagnosis					
Eeles, 1991 <sup>[14]</sup>	0.73 (0.44, 1.2)	Any	Cohort, retrospective	373, United Kingdom	No difference in disease-free interval
Ursic-Vrscaj,	0.67 (0.27, 1.62) (relapse)	Any	Case-control	72, Slovenia	

2001 <sup>[15]</sup>	0.90 (0.24, 5.08) (survival)	Any			
Mascarenhas, 2006 <sup>[5]</sup>	0.57 (0.42, 0.78)	Any	Cohort, follow-up on population-based cases	649, Sweden	Examination of different HRT preparations did not give different HRs
Wen, 2013 <sup>[16]</sup>	0.82 (0.48, 1.4) (relapse)	Any	Cohort, retrospective	144, China	
	0.67 (0.18, 2.5) (survival)	Any			
Eeles, 2015 <sup>[17]</sup>	0.63 (0.44, 0.90)	Any	RCT	150, United Kingdom, Spain, and Hungary	Greater relapse-free survival, HR=0.67 (0.47, 0.97)
Power, 2016 <sup>[18]</sup>	0.50 (0.23, 1.09) (<55yrs old)	Any	Cohort, retrospective	357, Manitoba, Canada	
	0.85 (0.43, 1.68) (≥55yrs old)	Any			
Li, 2012 <sup>[19]</sup>	0.88 (0.35, 2.32)	EPT	Cohort, prospective	75, China	
Guidozzi, 1999 <sup>[20]</sup>	0.97 (0.65, 1.18) (relapse)	ET	RCT	125, South Africa	Non-significantly better overall survival as well

**Table S2:** Ovarian Cancer Association Consortium (OCAC) study sites with women included in the analysis (Online only).

Site code	N (%)	Full study name	Country	Study design	Data collection	Diagnosis years	Residual Disease information
AUS	558 (8.7)	Australian Ovarian Cancer Study	Australia	Population-based	Self-completed questionnaire	2002-2006	Yes
CON	241 (3.8)	Connecticut Ovary Study	United States	Population-based	In-person interview	2002-2009	No
DOV	569 (8.9)	Diseases of the Ovary and their Evaluation	United States	Population-based	In-person interview	2002-2009	No
GER	97 (1.5)	Germany Ovarian Cancer Study	Germany	Population-based	Self-completed questionnaire	1992-1998	No
HAW	265 (4.1)	Hawaii Ovarian Cancer Study	United States	Population-based	In-person interview	1994-2007	Yes
HOP	407 (6.3)	Hormones and Ovarian Cancer Prediction	United States	Population-based	In-person interview	2003-2008	Yes
MAL	377 (5.9)	Danish Malignant Ovarian Tumor Study	Denmark	Population-based	In-person or telephone interview	1994-1999	No
MAY	638 (9.9)	Mayo Clinic Ovarian Cancer Case Control Study	United States	Clinic-based	In-person interview	1999-2008	Yes
NEC	879 (13.7)	New England Case-Control Study	United States	Population-based	In-person interview	1992-2008	Yes
NJO	118 (1.8)	New Jersey Ovarian Cancer Study	United States	Population-based	Telephone interview	2004-2008	No
OPL	478 (7.4)	Ovarian Cancer Prognosis and Lifestyle Study	Australia	Population-based	Self-completed questionnaire	2012-2015	Yes
POL	109 (1.7)	Polish Ovarian Cancer Case-Control Study	Poland	Population-based	In-person interview	2001-2003	No
UCI	221 (3.4)	UC Irvine Ovarian Cancer Study	United States	Population-based	Self-completed questionnaire	1995-2005	No
UKO	352 (5.5)	UK Ovarian Cancer Population Study	United Kingdom	Population-based	Self-completed questionnaire	2006-2007	No
USC	1110 (17.3)	Study of Lifestyle and Women's Health	United States	Population-based	In-person interview	1993-2005	No

**Table S3a:** Hazards ratios, by histotype, for menopausal hormone therapy (MHT) use before diagnosis of ovarian cancer among women with invasive epithelial ovarian cancer in the Ovarian Cancer Association Consortium (OCAC) (Online only).

	Overall	High-grade serous	Mucinous	Endometrioid	Clear cell	Low-grade serous
N	6,419	4,393	373	925	483	245
MHT use	HR (95% CI) <sup>a</sup>					
None (ref)	1.0	1.0	1.0	1.0	1.0	1.0
<5 years	0.97 (0.88, 1.06)	0.94 (0.85, 1.04)	1.77 (1.04, 3.02)	0.99 (0.71, 1.37)	0.97 (0.62, 1.51)	0.98 (0.58, 1.66)
5+ years	0.80 (0.74, 0.87)	0.78 (0.71, 0.86)	0.66 (0.34, 1.26)	1.08 (0.81, 1.43)	0.83 (0.47, 1.48)	0.76 (0.47, 1.23)

<sup>a</sup> Hazard ratios (HRs) are adjusted for age at diagnosis and race/ethnicity, and stratified by histotype (overall analysis only), stage at diagnosis, and OCAC site.

**Table S3b:** Hazard ratios (95% CIs) for MHT use and overall survival, for high-grade serous histology by stage of disease at diagnosis (Online only).

<b>High-grade serous</b>	
HR (95% CI)	
MHT use <sup>a</sup>	
<b>Stage I (local)</b>	N=205
<5 years	0.69 (0.32, 1.46)
5+ years	0.62 (0.28, 1.35)
<b>Stage II (regional)</b>	N=469
<5 years	0.83 (0.54, 1.30)
5+ years	0.72 (0.48, 1.08)
<b>Stage III and IV (advanced/distant)</b>	N=3719
<5 years	0.96 (0.86, 1.06)
5+ years	0.79 (0.71, 0.86)*

<sup>a</sup> Reference category of no use for all analyses.

<sup>b</sup> Hazard ratios (HRs) are adjusted for age at diagnosis and race/ethnicity, and stratified by OCAC site.

\* Significant at a level of  $p < 0.001$ .

**Table S4:** HR for MHT use and progression-free survival of ovarian cancer (Online only).

Progression-free survival	
N <sup>a</sup>	2,150
MHT use	HR (95% CI) <sup>b</sup>
None (ref)	1.0
<5 years	0.94 (0.81, 1.09)
5+ years	0.94 (0.82, 1.09)

<sup>a</sup> Data on presence of progression and time to progression was only available for a subset of the women.

<sup>b</sup> Hazard ratios (HRs) were adjusted for age at diagnosis and race/ethnicity, and stratified by histotype, stage at diagnosis, and OCAC site.

**Table S5:** Hazards ratios for MHT use and overall survival in a fully adjusted model (Online only).

Among women with complete information for all variables, N=4,044				
Variable included	Unadjusted	Age and stage <sup>1</sup>	Primary model	Fully adjusted <sup>2</sup> model
MHT use	--	Age at diagnosis, stage at diagnosis	Age, stage, race/ethnicity, histotype, OCAC site	HR (95% CI) <sup>a</sup>
None (ref)	1.0	1.0	1.0	1.0
<5 years	0.96 (0.87, 1.07)	0.96 (0.86, 1.06)	0.96 (0.87, 1.07)	0.96 (0.87, 1.07)
5+ years	0.98 (0.89, 1.08)	0.80 (0.73, 0.88)	0.79 (0.71, 0.87)	0.79 (0.71, 0.88)

<sup>1</sup> Change in HR estimates was primarily driven by accounting for stage, though age confounded the relationship as well.


<sup>2</sup> Stratified by stage at diagnosis, OCAC site, histotype, and adjusted for age at diagnosis, race/ethnicity, BMI, education level, tubal ligation, endometriosis, hysterectomy, combined oral contraceptive use duration (never, <1 year, 1 to <5 years, 5 to <10 years, 10+ years), parity, family history of breast cancer, family history of ovarian cancer, smoking status (never, former, current).




**Table S6:** Hazard ratios (HRs) for MHT use and overall survival, among advanced stage, high-grade serous carcinoma, stratified by macroscopic residual disease (Online only).

<b>MHT use</b>	<b>HR (95% CI)</b>	
	Residual disease N=891	No residual disease N=497
<b>None (ref)</b>	1.0	1.0
<b>&lt;5 years</b>	1.08 (0.86, 1.35)	1.07 (0.74, 1.53)
<b>5+ years</b>	0.89 (0.68, 1.17)	0.81 (0.54, 1.22)

## Supplemental Figures



**Figure S1: Study participants.** All women were from the Ovarian Cancer Association Consortium (OCAC). The population-based and clinic-based studies included were done in the United States (n=9), Europe (n=4), and Australia (n=2). Only post-menopausal women with ovarian carcinoma for whom survival time data was available were considered for this analysis. The exclusion stage “missing histotype” included exclude of those with mixed cell and undifferentiated tumors. Of the five histotypes high-grade serous was the most common (68% of cases) among these women, followed by endometrioid (14%), clear cell (7.5%), mucinous (5.8%), and low-grade serous (3.8%). Of the 6,419 women in our analytic sample, subsets with complete information were analyzed for progression-free survival (n=2,239) and for the association between hormone therapy and residual disease (n=2,056). Time of interview refers to time of study enrollment; for some studies this was an in-person interview and for some it was a self-administered questionnaire.



**Figure S2: Adjusted overall survival curves.** Survival curves, among advanced stage, high-grade serous carcinoma. Adjusted for age, race/ethnicity, and OCAC site.

## REFERENCES

1. Cronin KA, Lake AJ, Scott S, et al: Annual Report to the Nation on the Status of Cancer, part I: National cancer statistics. *Cancer* 124:2785-2800, 2018
2. Beral V, Million Women Study C, Bull D, et al: Ovarian cancer and hormone replacement therapy in the Million Women Study. *Lancet* 369:1703-10, 2007
3. Collaborative Group On Epidemiological Studies Of Ovarian C, Beral V, Gaitskell K, et al: Menopausal hormone use and ovarian cancer risk: individual participant meta-analysis of 52 epidemiological studies. *Lancet* 385:1835-42, 2015
4. Lee AW, Ness RB, Roman LD, et al: Association Between Menopausal Estrogen-Only Therapy and Ovarian Carcinoma Risk. *Obstet Gynecol* 127:828-36, 2016
5. Shafrir AL, Babic A, Tamimi RM, et al: Reproductive and hormonal factors in relation to survival and platinum resistance among ovarian cancer cases. *Br J Cancer* 115:1391-1399, 2016
6. Power L, Lefas G, Lambert P, et al: Hormone Use After Nonserous Epithelial Ovarian Cancer: Overall and Disease-Free Survival. *Obstet Gynecol* 127:837-47, 2016
7. Kim SJ, Rosen B, Fan I, et al: Epidemiologic factors that predict long-term survival following a diagnosis of epithelial ovarian cancer. *Br J Cancer* 116:964-971, 2017
8. Mascarenhas C, Lambe M, Bellocco R, et al: Use of hormone replacement therapy before and after ovarian cancer diagnosis and ovarian cancer survival. *Int J Cancer* 119:2907-15, 2006
9. Nagle CM, Bain CJ, Green AC, et al: The influence of reproductive and hormonal factors on ovarian cancer survival. *Int J Gynecol Cancer* 18:407-13, 2008
10. Felix AS, Bunch K, Yang HP, et al: Menopausal hormone therapy and mortality among women diagnosed with ovarian cancer in the NIH-AARP Diet and Health Study. *Gynecol Oncol Rep* 13:13-7, 2015
11. Zhang M, Holman CD: Tubal ligation and survival of ovarian cancer patients. *J Obstet Gynaecol Res* 38:40-7, 2012
12. Wernli KJ, Newcomb PA, Hampton JM, et al: Hormone therapy and ovarian cancer: incidence and survival. *Cancer Causes Control* 19:605-13, 2008
13. Besevic J, Gunter MJ, Fortner RT, et al: Reproductive factors and epithelial ovarian cancer survival in the EPIC cohort study. *Br J Cancer* 113:1622-31, 2015
14. Madalinska JB, van Beurden M, Bleiker EM, et al: The impact of hormone replacement therapy on menopausal symptoms in younger high-risk women after prophylactic salpingo-oophorectomy. *J Clin Oncol* 24:3576-82, 2006
15. Organization WH: BMI Classification, Global Database on Body Mass Index, 2006
16. Bhatla N, Denny L: FIGO Cancer Report 2018. *Int J Gynaecol Obstet* 143 Suppl 2:2-3, 2018
17. Imai K, Keele L, Tingley D: A general approach to causal mediation analysis. *Psychol Methods* 15:309-34, 2010
18. R Core Team: R: A language and environment for statistical computing. Vienna, Austria, R Foundation for Statistical Computing, 2019

19. Hein A, Thiel FC, Bayer CM, et al: Hormone replacement therapy and prognosis in ovarian cancer patients. *Eur J Cancer Prev* 22:52-8, 2013
20. Liu Z, Beach JA, Agadjanian H, et al: Suboptimal cytoreduction in ovarian carcinoma is associated with molecular pathways characteristic of increased stromal activation. *Gynecol Oncol* 139:394-400, 2015
21. Riestler M, Wei W, Waldron L, et al: Risk prediction for late-stage ovarian cancer by meta-analysis of 1525 patient samples. *J Natl Cancer Inst* 106, 2014
22. Georgiadou P, Sbarouni E: Effect of hormone replacement therapy on inflammatory biomarkers. *Adv Clin Chem* 47:59-93, 2009
23. Lamon-Fava S, Posfai B, Schaefer EJ: Effect of hormonal replacement therapy on C-reactive protein and cell-adhesion molecules in postmenopausal women. *Am J Cardiol* 91:252-4, 2003
24. Martin-Millan M, Castaneda S: Estrogens, osteoarthritis and inflammation. *Joint Bone Spine* 80:368-73, 2013
25. Straub RH: The complex role of estrogens in inflammation. *Endocr Rev* 28:521-74, 2007
26. Ono M: Molecular links between tumor angiogenesis and inflammation: inflammatory stimuli of macrophages and cancer cells as targets for therapeutic strategy. *Cancer Sci* 99:1501-6, 2008
27. Harris BS, Bishop KC, Kuller JA, et al: Hormonal management of menopausal symptoms in women with a history of gynecologic malignancy. *Menopause*, 2019
28. Eeles RA, Tan S, Wiltshaw E, et al: Hormone replacement therapy and survival after surgery for ovarian cancer. *BMJ* 302:259-62, 1991
29. Wen Y, Huang H, Huang H, et al: The safety of postoperative hormone replacement therapy in epithelial ovarian cancer patients in China. *Climacteric* 16:673-81, 2013
30. Li L, Pan Z, Gao K, et al: Impact of post-operative hormone replacement therapy on life quality and prognosis in patients with ovarian malignancy. *Oncol Lett* 3:244-249, 2012
31. Eeles RA, Morden JP, Gore M, et al: Adjuvant Hormone Therapy May Improve Survival in Epithelial Ovarian Cancer: Results of the AHT Randomized Trial. *J Clin Oncol* 33:4138-44, 2015
32. Guidozi F, Daponte A: Estrogen replacement therapy for ovarian carcinoma survivors: A randomized controlled trial. *Cancer* 86:1013-8, 1999
33. Bebar S, Ursic-Vrscaj M: Hormone replacement therapy after epithelial ovarian cancer treatment. *Eur J Gynaecol Oncol* 21:192-6, 2000
34. Ursic-Vrscaj M, Bebar S, Zakelj MP: Hormone replacement therapy after invasive ovarian serous cystadenocarcinoma treatment: the effect on survival. *Menopause* 8:70-5, 2001
35. Sandini L, Pentti K, Tuppurainen M, et al: Agreement of self-reported estrogen use with prescription data: an analysis of women from the Kuopio Osteoporosis Risk Factor and Prevention Study. *Menopause* 15:282-9, 2008
36. Hu FB, Grodstein F, Hennekens CH, et al: Age at natural menopause and risk of cardiovascular disease. *Arch Intern Med* 159:1061-6, 1999
37. Hodis HN, Mack WJ, Henderson VW, et al: Vascular Effects of Early versus Late Postmenopausal Treatment with Estradiol. *N Engl J Med* 374:1221-31, 2016

38. Grodstein F, Manson JE, Stampfer MJ: Hormone therapy and coronary heart disease: the role of time since menopause and age at hormone initiation. *J Womens Health (Larchmt)* 15:35-44, 2006
39. Manson JE, Chlebowski RT, Stefanick ML, et al: Menopausal hormone therapy and health outcomes during the intervention and extended poststopping phases of the Women's Health Initiative randomized trials. *JAMA* 310:1353-68, 2013
40. Rossouw JE, Anderson GL, Prentice RL, et al: Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results From the Women's Health Initiative randomized controlled trial. *JAMA* 288:321-33, 2002

**Highlights**

- Using menopausal hormone therapy prior to diagnosis extends ovarian cancer survival.
- Estrogen alone and estrogen+progestin are associated with better survival.
- Women who used hormone therapy have less residual disease after surgery.