

Table 2
SAI-ED scores in anorexia nervosa and bulimia nervosa

	AN (<i>n</i> =44)	BN (<i>n</i> =50)	<i>z</i> [†]	<i>p</i>
	Median (IQR)			
SAI-ED total	21.00 (8.00)	26.00 (5.00)	-4.99	< 0.001
Recognition of mental illness	1.00 (1.00)	2.00 (1.00)	-3.29	0.001
Awareness of need for treatment	1.00 (1.00)	2.00 (1.00)	-3.22	0.001
Recognition of body weight concerns	3.00 (0.00)	4.00 (1.00)	-3.88	< 0.001
Relabeling of body weight concerns	3.00 (1.00)	3.00 (1.00)	-4.35	< 0.001
Recognition of eating pathology	3.00 (1.00)	4.00 (1.00)	-3.37	0.001
Relabeling of eating pathology	3.00 (1.00)	3.00 (1.00)	-4.58	< 0.001
Hypothetical contradiction	2.00 (1.00)	3.00 (2.00)	-3.09	0.002
Compliance with treatment	4.00 (1.75)	5.00 (2.00)	-3.65	< 0.001

IQR: Inter-quartile range, SAI-ED: Schedule for the Assessment of Insight in Eating Disorders

[†] *p*-values are estimated with Mann-Whitney U test (standardized test statistic values are displayed)