

## REVIEW

# The role of senescent T cells in immunopathology

Luciana P. Covre<sup>1,2</sup>  | Roel P. H. De Maeyer<sup>1</sup>  | Daniel C. O. Gomes<sup>2</sup>  | Arne N. Akbar<sup>1</sup> 

<sup>1</sup>Division of Medicine, University College London, London, UK

<sup>2</sup>Núcleo de Doenças Infecciosas, Universidade Federal do Espírito Santo, Vitória, Brazil

## Correspondence

Daniel C. O. Gomes, Núcleo de Doenças Infecciosas, Universidade Federal do Espírito Santo, Vitória, Brazil.  
Email: dgomes@ndi.ufes.br

Arne N. Akbar, Division of Medicine, University College London, London, UK.  
Email: a.akbar@ucl.ac.uk

## Funding information

Fundação de Amparo a Pesquisa do Espírito Santo- FAPES/Newton Fund and Medical Research Council, Grant/Award Number: 72939273/16; Fundação de Amparo a Pesquisa do Espírito Santo- FAPES, Grant/Award Number: 90/2017; Fundação de Amparo a Pesquisa do Espírito Santo- FAPES/Ministério da Saúde - Brazil, Grant/Award Number: 83152997/2018; Coordination for the Improvement of Higher Education Personnel - CAPES- Brazil; Medical Research Council (UK), Grant/Award Number: MR/T015853/1 and MR/P00184X/1

## Abstract

The development of senescence in tissues of different organs and in the immune system are usually investigated independently of each other although during ageing, senescence in both cellular systems develop concurrently. Senescent T cells are highly inflammatory and secrete cytotoxic mediators and express natural killer cells receptors (NKR) that bypass their antigen specificity. Instead they recognize stress ligands that are induced by inflammation or infection of different cell types in tissues. In this article we discuss data on T cell senescence, how it is regulated and evidence for novel functional attributes of senescent T cells. We discuss an interactive loop between senescent T cells and senescent non-lymphoid cells and conclude that in situations of intense inflammation, senescent cells may damage healthy tissue. While the example for immunopathology induced by senescent cells that we highlight is cutaneous leishmaniasis, this situation of organ damage may apply to other infections, including COVID-19 and also rheumatoid arthritis, where ageing, inflammation and senescent cells are all part of the same equation.

## KEYWORDS

Ageing, senescence, T cell

## 1 | INTRODUCTION

As we age, we accumulate cells in many organs that exhibit signs of DNA damage, have poor proliferative capacity and are highly secretory. These cells are senescent, defined as being in a state of cell cycle arrest associated with phenotypic and functional changes (Campisi, 2013; Campisi & Fagagna, 2007). This process is primarily thought to prevent cancerous transformation of dividing cells (Braig et al., 2005; Chen et al., 2005), but senescence plays a vital role in developmental biology as well as in wound healing/tissue repair

(Demaria et al., 2014; Krizhanovsky et al., 2008; Kurundkar et al., 2016; Pitiyage et al., 2011; van Deursen, 2014). While transient senescence is a beneficial mechanism earlier in life, the accumulation of senescent cells with increasing age leads to organ dysfunction, driving inflammation and may underlie many age-related diseases such as atherosclerosis (Childs et al., 2016), osteoarthritis (Jeon et al., 2017), neurodegenerative diseases (Chinta et al., 2018; Kritsilis et al., 2018) and cirrhosis (Wiemann et al., 2002).

Senescence is triggered by activation and subsequent maintenance of DNA damage repair (DDR) signalling pathways. This can be

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Aging Cell* published by Anatomical Society and John Wiley & Sons Ltd



induced by oncogene activation, replicative stress related to telomere erosion and oxidative stress (Allsopp, 2008; Blackburn et al., 2015) that kickstart a DDR. This involves proteins such as  $\gamma$ -H2AX, ATM, ATR, p16 and p53 (Nassour et al., 2016) that leads to cycle arrest which becomes permanent if repair fails. Importantly, while senescent cells do not proliferate, they are still metabolically active and secrete a wide array of cytokines, chemokines and matrix metalloproteinases that vary according to the cell type involved (Coppé et al., 2010). It is this senescence-associated secretory phenotype (SASP) that is thought to play a crucial role in age-related pathology such as the chronic low grade inflammation observed with advanced age, called inflammaging (Franceschi et al., 2000). Therefore, two recognized characteristics of senescent cells are the lack of proliferative activity and the presence of a senescence-associated secretory phenotype (SASP, Campisi, 2013).

While senescence was first discovered in fibroblasts and extensively worked on in other non-leukocytic cells, it has become increasingly clear that immune cells undergo senescence as well. Within the immune system, the existence of non-proliferative leukocyte populations that have high capacity for biologically active mediator secretion has been recognized for many decades, albeit under a different name. These are the effector T lymphocytes that secrete pro-inflammatory cytokines and cytotoxic granules but do not proliferate after activation (Dunne et al., 2005). Recent studies show that these cells also harbour DNA damage, short telomeres, low telomerase activity and engage signalling pathways associated with cellular senescence (Callender et al., 2018; Henson et al., 2014; Lanna et al., 2014). Therefore, the terms effector T cells and senescent T cells may be synonymous and refer to the same T cell populations. Effector T cells that are expanded in number during the acute phase of a human anti-viral response to Epstein-Barr Virus (EBV) are highly expanded in number and lose the expression of the anti-apoptotic molecule Bcl-2 that makes them short-lived (Akbar et al., 1993). These cells also have relatively long telomeres due to upregulation of the enzyme telomerase and are in cell cycle (Maini et al., 1999). The difference between such short-lived effector cells and senescent T cells in the steady state is that the latter are non-proliferative, do not express telomerase after activation, exhibit short telomeres and express senescence related signalling molecules as described above. Nevertheless senescent cells in the steady state in both CD4 and CD8 compartments still express lower levels of Bcl-2 than the other subsets. Collectively this indicates that the extent of T cell proliferation the acute phase of a viral infection drives T cells to senescence. These cells are still susceptible to apoptosis but can persist if sufficient antiapoptotic cytokines are tissue niches. It can be argued that senescent T cells derive from a subpopulation of effector T cells that do not undergo apoptosis, instead becoming senescent and lingering long term. In this review, we discuss recent observations suggesting that certain infectious agents can drive the accumulation of CD8<sup>+</sup> T cells that exhibit all the hallmarks of senescence as described above (henceforth referred to as T<sub>sen</sub>). We will focus on patients infected with *Leishmania braziliensis* where the increased T<sub>sen</sub> numbers in the skin may induce the characteristic skin lesions associated with this

disease. We propose a hypothesis, based on existing data, that this pathology occurs because senescent non-lymphoid cells in the skin are killed by infiltrating T<sub>sen</sub>, a novel interaction between senescent lymphoid and non-lymphoid cells that may also have implications for ageing.

## 2 | FUNCTIONAL PROPERTIES OF CD8<sup>+</sup> T<sub>sen</sub>

CD8<sup>+</sup> T cells can be subdivided based on expression of the co-stimulatory molecules CD27 and CD28, where naïve CD8<sup>+</sup> T cells co-express both markers and as they differentiate to an effector phenotype lose CD28 expression and subsequently CD27 (Henson et al., 2012). Sometimes these will be subdivided further into T<sub>EMRA</sub> cells, so named because they re-express CD45RA (Henson et al., 2012). The CD27<sup>-</sup>CD28<sup>-</sup> compartment harbours T<sub>sen</sub> cells which can be identified further by expression of KLRG1 and CD57 (Brenchley et al., 2003; Henson et al., 2009). Moreover, many studies have shown a plethora of markers that are upregulated on T<sub>sen</sub> cells, such as downregulated telomerase, shortened telomeres, DNA damage responses and constitutive MAPK activity, features specific to T<sub>sen</sub> over effector T cells. T<sub>sen</sub> also exhibit a SASP consisting of proteases and pro-inflammatory mediators such as TNF and IL-1 $\beta$  (Callender et al., 2018). These functional changes are summarized in Table 1 (Akbar et al., 1993; Brenchley et al., 2003; Callender et al., 2018, 2019; Geginat et al., 2003; Gumá et al., 2004; Henson et al., 2009, 2012, 2014, 2015; Libri et al., 2011; Maini et al., 1999; Ouyang et al., 2003; Pereira et al., 2020; Plunkett et al., 2001; Tarazona et al., 2001, 2002; Voehringer et al., 2002; Weng et al., 2009). Single cell RNAseq analysis of peripheral blood derived CD8<sup>+</sup> T cells show that CD8<sup>+</sup> T cells identified by the above functional criteria also express multiple features associated with cellular senescence (Table 1, Pereira et al., 2020). This indicates that the terms senescent and effector CD8<sup>+</sup> T cells are synonymous and identify identical populations.

CD8<sup>+</sup> T<sub>sen</sub> have limited proliferative capacity after activation via the T cell receptor (TCR) complex (Akbar et al., 2016) that is due in part to downregulation of key TCR signalling molecules such as LCK, LAT and PLC- $\gamma$  (Pereira et al., 2020). CD4<sup>+</sup> T<sub>sen</sub> also exhibit identical markers of senescence to their CD8<sup>+</sup> counterparts (Lanna et al., 2014) but are present at much lower frequency than CD8<sup>+</sup> T<sub>sen</sub> in the peripheral blood of healthy donors. The development of senescence characteristics in T cells was initially considered to indicate the dysfunction of these cells during ageing (Akbar et al., 2004). However, CD8<sup>+</sup> T<sub>sen</sub> populations express surface receptors that are associated with NK cells such as NKG2D, NKG2C, NKG2A and Killer immunoglobulin-like receptor (KIR) families compared to undifferentiated and non-senescent CD28<sup>+</sup>CD27<sup>+</sup> CD8<sup>+</sup> T cells (Abedin et al., 2005; Michel et al., 2016; Pereira et al., 2020; Vallejo et al., 2011). This suggests rather than being dysfunctional, these cells acquire an alternative functional profile as they differentiate towards senescence. This is supported by observations that these T<sub>sen</sub> express DAP12, an NK cytotoxicity adaptor molecule, and are capable of killing



TABLE 1 List of T cell, NK cell and senescence-associated marker expression in CD8 T cell subsets

Cellular feature	Naïve	Central memory	Effector memory	EMRA/senescent	References
<b>T cell-associated</b>					
CD27	+++	++	-	-	Henson et al. (2012)
CD28	+++	++	+/-	-	Henson et al. (2012)
CD45RA	+++	++	-	++	Henson et al. (2012)
CCR7	+++	++	-	+/-	Henson et al. (2012)
CD62L	++	++	-	-	Henson et al. (2012)
Lck-LAT-Zap70	+	+	+	+/-	Pereira et al. (2020)
PI3 K-Akt-mTOR	+	+	+	+/-	Henson et al. (2014)
TCR-mediated proliferation	+++	+++	++	+/-	Henson et al. (2009) Pereira et al. (2020)
<b>NK cell-associated</b>					
CD57	-	-	+	++	Brenchley et al. (2003)
KLRG1	-	-	+	++	Voehringer et al. (2002) Ouyang et al. (2003) Henson et al. (2009)
NKG2A/C	-	-	+/-	+	Tarazona et al. (2002) Gumá et al. (2004) Pereira et al. (2020)
NKG2D	+	+	+	++	Tarazona et al. (2002) Gumá et al. (2004)
CD244	+	-	-	+++	Tarazona et al. (2002)
KIRs	-	-	-	++	Gumá et al. (2004) Pereira et al. (2020)
DAP12	-	-	-	+	Pereira et al. (2020)
<b>Senescence</b>					
Proliferative capacity	+++	+++	++	-	Weng et al. (2009)
Telomere length	+++	+++	++	+/-	Plunkett et al. (2001) Weng et al. (2009)
Telomerase	+++	++	+	-	Maini et al. (1999) Weng et al. (2009)
BCL-2	+++	++	+	+/-	Akbar et al. (1993) Geginat et al. (2003) Libri et al. (2011)
p16	-	-	-	+	Henson et al. (2015) Pereira et al. (2020)
γH2AX	-	-	-	++	Callender et al. (2019)
DDR	-	-	-	+	Callender et al. (2018) Pereira et al. (2020)
SASP	-	-	-	++	Callender et al. (2018)
IL-1β	-	-	++	+	Callender et al. (2018)
IL-18	-	-	-	++	Callender et al. (2018)
CCL16	-	-/+	++	+++	Callender et al. (2018)
ADAM28	-	-	-	+++	Callender et al. (2018)
Sestrins	-	-	+	++	Pereira et al. (2020)
MAPK (p38/Erk/Jnk)	-	-	+/-	++	Henson et al. (2014), Henson et al. (2015) Pereira et al. (2020)

tumour cell lines in an MHC-I-independent manner (Pereira et al., 2020). Mechanistically, the switch from TCR to NKR expression in CD8<sup>+</sup> T<sub>sen</sub> is regulated by stress proteins known as the sestrins in

both humans and mice (Pereira et al., 2020). Of note, CD4<sup>+</sup> T<sub>sen</sub> also express NKRs, suggesting that they may also mediate effector functions through these receptors (Pereira et al., 2020).



### 3 | CD8<sup>+</sup> T<sub>sen</sub> ACCUMULATE DURING AGEING AND PERSISTENT VIRAL INFECTION

Previous reports have shown that CD8<sup>+</sup> T<sub>sen</sub> accumulate in older humans and that this is largely driven by ongoing sub-clinical responses to persistent viruses especially cytomegalovirus (CMV) (Jackson et al., 2017; Nikolich-Zugich et al., 2020; Savva et al., 2013). Moreover, this population has increased capacity to rapidly respond to signals mediated by inflammatory cytokines (Freeman et al., 2012). Furthermore, proinflammatory cytokine production associated with persistent CMV infection may induce bystander senescence in non-CMV specific T cells (Fletcher et al., 2005). Therefore the cytotoxic CD8<sup>+</sup> T<sub>sen</sub> populations that accumulate during ageing and during persistent viral infections are now re-focused to preferentially recognize NK ligands that may be expressed by infected, malignant or "stressed" tissues (Pereira et al., 2019; Sagiv et al., 2016). Therefore, T<sub>sen</sub> can mediate NKR dependent cytotoxicity, independently of their antigen specificity.

### 4 | THE KILLING OF SENESCENT NON-LYMPHOID TISSUE CELLS BY NK AND CD8<sup>+</sup> T CELLS


Senescent non-lymphoid cells accumulate in many tissues during ageing (Campisi, 2013) and are associated with tissue dysfunction that lead to frailty. Transgenic mouse models that enable the specific removal of senescent cells from different tissues *in vivo* show that the elimination of these cells increases lifespan, improves overall fitness and reduces age-associated characteristics of the animals (Baar et al., 2017; Baker et al., 2011, 2016). An exciting observation was that cells of the immune system, including NK, CD4<sup>+</sup> and CD8<sup>+</sup> T cells, can also recognize and eliminate senescent cells *in vitro* and *in vivo* (Gorgoulis et al., 2019; Pereira et al., 2019), a new role for immunosurveillance that may be relevant for ageing. The mechanisms involved in this include the interaction between NKG2D and NKG2A/C receptors on CD8<sup>+</sup> T and NK cells, and their ligands such as MICA/B and HLA-E expressed by senescent stromal cells (Krizhanovsky et al., 2008; Pereira et al., 2019; Sagiv et al., 2016).

Despite the evidence for senescent cell clearance by the immune system, it is not yet clear why senescent cells accumulate during ageing and persist at sites of age-related pathology (Karin et al., 2019; Ovadya et al., 2018; Sharpless & Sherr, 2015). Altered immunity during ageing may contribute to this, however senescent cells also elicit evasion strategies, including the expression of inhibitory NK receptor ligands such as HLA-E (Pereira et al., 2019) and the shedding of NKG2D ligands that act as decoy receptors that interfere with cytotoxic cell recognition and killing (Muñoz et al., 2019). Therefore a balance between activating and inhibitory signals will determine the outcome of the NK and T cell cytotoxic response to senescent tissue cells. On one hand, the development of NK-like function in CD8<sup>+</sup> T<sub>sen</sub> cells that lose the capacity to proliferate may be beneficial

as it enables the retention of cytotoxic cells with broad spectrum of activity during ageing that may be important for killing of infected and malignant cells. However the acquisition of NK-like function in T cells during ageing may have negative consequences and induce pathology, for example in skin lesion formation in patients with cutaneous leishmaniasis (Figure 1).

### 5 | IMMUNITY IN PATIENTS WITH CUTANEOUS LEISHMANIASIS; THE DOUBLE-EDGED SWORD

*Leishmania braziliensis* is the main causal agent of cutaneous leishmaniasis (CL) in Brazil, where destructive cutaneous lesions develop (World Health Organization/Department of control of neglected tropical diseases, 2015). Patients with *L. braziliensis* skin infections usually develop lesions after 2–4 weeks (Oliveira-Ribeiro et al., 2017). The early stages of infection do not present with CD8<sup>+</sup> T cell infiltration, but CD8<sup>+</sup> T cell activity increases as the infection progresses and the characteristic CL skin lesions form (Faria et al., 2009). Production of inflammatory mediators such as IFN- $\gamma$  as well as CD8<sup>+</sup> cytotoxic T cell activity is observed during the acute and healing phase of *L. major* infection and is linked to



**FIGURE 1** Leishmania infection results in infiltration of T<sub>sen</sub> that aberrantly kill stressed and senescent stromal cells contributing to excessive damage. Leishmania infection of macrophages induces inflammatory cytokine production driving expression of MICA/B and other stress ligand on resident skin cells such as fibroblasts. Concurrently, these cytokines activate the endothelium which upregulates adhesion markers like E-selectin. This binds to CLA expressing senescent T cells enabling their infiltration into the skin. Here, these senescent T cells, expressing a host of NKRs including NKG2D interact with resident skin cells. The interaction between NKG2D<sup>+</sup> T<sub>sen</sub> and MICA/B<sup>+</sup> fibroblasts results in killing of the latter and contributing to off-target tissue pathology.



parasite clearance (Rossi & Fasel, 2017). There is a strong correlation between the severity of the disease and the number of CD8<sup>+</sup> T cells present in the lesion (Faria et al., 2009; Santos et al., 2013), but this is independent of the presence of parasites in the lesions (Carvalho et al., 2007; Murray et al., 2005; Pearson et al., 1996). Despite this, if untreated, the lesions increase in size progressively causing the characteristic pathology of CL. This raises the question as to the cause of the lesions in the skin suggesting the possibility that chronic inflammation and non-specific CD8<sup>+</sup> T or NK cell cytotoxic responses may lead to non-specific tissue destruction.

*Leishmania braziliensis* infected individuals have elevated numbers of circulating CD4<sup>+</sup> and CD8<sup>+</sup> T cells that possess short telomeres, decreased expression of the catalytic component of the enzyme telomerase (hTERT), have reduced proliferative activity and increased expression of the nuclear DNA-damage response protein (Covre et al., 2019). In addition, these T cells secrete SASP-like cytokines in response to *L. braziliensis* antigen (LbAg) or anti-CD3 stimulation compared to controls *in vitro*. Although T<sub>sen</sub> accumulate in patients (Covre et al., 2019), the robust proliferative response to LbAg stimulation indicates that not all the *L. braziliensis* specific T cells that are present are senescent (Covre et al., 2019). Nevertheless, the association between the frequency of both CD4<sup>+</sup> and CD8<sup>+</sup> T<sub>sen</sub> cells in the circulation and cutaneous lesion size in infected patients suggests that these cells may be recruited into the lesions and may contribute to the observed skin ulceration.

An interesting observation is that the accumulation of senescent T cells, but not NK correlates with the age of CL patients (Covre L and Gomes D, unpublished data). Compared to young subjects, elderly patients displayed larger cutaneous lesions, longer duration of illness and were more likely to develop mucocutaneous leishmaniasis, the most inflammatory and severe clinical form of the tegumentary leishmaniasis (Carvalho et al., 2015). Senescent T cells from CL patients but not healthy controls also upregulate the skin homing receptor CLA in response to activation with *L. braziliensis* antigens (Covre et al., 2019) This facilitates their homing into the skin where they can perform effector functions. Interestingly, while NK cells in the peripheral blood of have a greater cytotoxic potential compared to CD8<sup>+</sup> T<sub>sen</sub>, NK cells have reduced CLA expression and do not home to skin lesions as efficiently as CD8<sup>+</sup> T<sub>sen</sub>. This suggests that during acute infection with *L. braziliensis*, T<sub>sen</sub> within both CD4<sup>+</sup> and CD8<sup>+</sup> subsets home to the skin where they initially control the infection. However their potent inflammatory activity may also contribute to development of the skin pathology of CL even after the parasites have been cleared. This raises the question about the mechanism involved in the tissue damage within the cutaneous lesions of these patients.

## 6 | HYPOTHESIS: THE NON-SPECIFIC NATURE OF CD8<sup>+</sup> T<sub>sen</sub> INDUCES IMMUNOPATHOLOGY DURING INFECTIONS SUCH AS CUTANEOUS LEISHMANIASIS

We hypothesize that dermal macrophages infected by *Leishmania* initiate an inflammatory response that preferentially attracts

*L. braziliensis* specific CD8<sup>+</sup> T<sub>sen</sub> from the blood which clear the infection. However, the inflammation in the skin, induced by the interaction between infected macrophages and the infiltrating T cells, induces the expression of stress ligands including those that bind to NK receptors by the surrounding stroma (Gasser et al., 2005; Groh et al., 2001; Molinero et al., 2004). The interaction between these ligands and NK receptors on T<sub>sen</sub> leads to cytotoxic killing and tissue damage. In turn, this damage leads to exacerbated inflammation, inducing further expression of stress ligands and results in more destruction by CD8<sup>+</sup> T<sub>sen</sub>. This generates a destructive positive feedback loop that leads to progressively larger skin lesions if left untreated. At this advanced stage of the infection, CD8<sup>+</sup> T<sub>sen</sub> cells that infiltrate the inflamed lesion do not have to be specific for *L. braziliensis* but may be specific for other antigens (e.g. CMV) that have been shown to drive T cell senescence and therefore the acquisition of NK characteristics (Figure 1). This is supported by previous observations that the bystander activation of memory CD8<sup>+</sup> T cells in the skin may contribute to the chronic inflammation and tissue damage in an NKG2D-dependent manner during murine cutaneous leishmaniasis (Crosby et al., 2014). Furthermore, the pathogenesis of murine cutaneous leishmaniasis is exacerbated by LCMV infection-induced CD8<sup>+</sup>NKG2D<sup>+</sup> T cells, supporting the possibility that tissue damage can occur independently of the antigen specificity of the infiltrating cells (Crosby et al., 2015). Interestingly, human lesional CD4<sup>+</sup> and CD8<sup>+</sup> T cells are able to proliferate and produce IFN-γ in response not only to *Leishmania* antigens but also to other pathogens (Da-Cruz et al., 2010). This highlights that not all the T cell in the lesions are senescent but supports the role of T cells of different specificities in the promotion of cutaneous lesions in CL.

## 7 | FUTURE PERSPECTIVES

The investigation of the cellular infiltrate in patients with cutaneous leishmaniasis by gene expression (Amorim et al., 2019; Christensen et al., 2016) and histological analyses is essential to determine the presence of alternative ligands for CD8<sup>+</sup> T<sub>sen</sub> within the tissue. It is also crucial to identify the source of inflammation at these sites, are the myeloid cells, the stromal cells, the leukocytes or all of them involved? The hypothetical positive feedback loop described here that involves T<sub>sen</sub> and excessive inflammation and non-specific tissue damage may also be involved in other non-resolving inflammatory diseases where accumulation of T<sub>sen</sub> cells have been demonstrated including Chagas disease (Molina & Kierszenbaum, 1989) and malaria (Cockburn et al., 2014). The accumulation of senescent T cells with increased pro-inflammatory potential has been implicated during age-related diseases such as rheumatoid arthritis (Goronzy et al., 2013; Weyand et al., 2017), Alzheimer's (Gate et al., 2020), and cardiovascular diseases (Youn et al., 2019; Yu et al., 2015), where the non-specific tissue damage driven by T<sub>sen</sub> cells needs to be investigated and might offer new opportunities for prevention and treatment (Andersson et al., 2011). Furthermore, it would important to know if inflammation that is induced by infections such



as SARS-CoV-2 that have a disproportionate pathological impact in older individuals (Merad & Martin, 2020) results from non-specific  $T_{sen}$  recruitment and induction of non-antigen specific damage in the lung and other tissues. Furthermore, it is currently unclear how senolytic drugs that are being used to clear senescent cells will affect senescent leukocyte populations and if this may have beneficial or detrimental consequences. Collectively we suggest that T cells that have been driven to extreme differentiation and senescence by infectious agents acquire potent MHC-I-unrestricted cytotoxic and capacity for inflammatory cytokine secretion together with the ability to cause non-specific NK-related damage. Given the increased burden of tumours and infections with age, the expansion of NK cell-like functions in  $CD8^+$  T cells could be an advantageous adaptation that would enable the recognition and elimination of infected and transformed cells. However, the accumulation of senescent T cells during ageing is a double-edged sword that may induce pathology, especially in inflammatory environments.

## ACKNOWLEDGEMENTS

Authors would like to thank the Fundação de Amparo a Pesquisa do Espírito Santo- FAPES/Newton Fund and Medical Research Council (Grant 72939273/16); Fundação de Amparo a Pesquisa do Espírito Santo- FAPES (Grant 90/2017); Fundação de Amparo a Pesquisa do Espírito Santo- FAPES/Ministério da Saúde (Grant 83152997/2018); and Coordination for the Improvement of Higher Education Personnel - CAPES- Brazil; Medical Research Council (UK) (Grants MR/T015853/1 and MR/P00184X/1) for supporting this work.

## CONFLICT OF INTEREST

None declared.

## AUTHORS' CONTRIBUTIONS

All four authors conceived and wrote the manuscript.

## ORCID

Luciana P. Covre <https://orcid.org/0000-0001-5877-0137>

Roel P. H. De Maeyer <https://orcid.org/0000-0003-4041-0867>

Daniel C. O. Gomes <https://orcid.org/0000-0002-0663-0981>

Arne N. Akbar <https://orcid.org/0000-0002-3763-9380>

## REFERENCES

- Abedin, S., Michel, J. J., Lemster, B., & Vallejo, A. N. (2005). Diversity of NKR expression in aging T cells and in T cells of the aged: The new frontier into the exploration of protective immunity in the elderly. *Experimental Gerontology*, *40*, 537–548.
- Akbar, A. N., Beverley, P. C., & Salmon, M. (2004). Will telomere erosion lead to a loss of T-cell memory? *Nature Reviews Immunology*, *4*, nri1440.
- Akbar, A. N., Borthwick, N., Salmon, M., Gombert, W., Bofill, M., Shamsadeen, N., Pilling, D., Pett, S., Grundy, J. E., & Janossy, G. (1993). The significance of low bcl-2 expression by CD45RO T cells in normal individuals and patients with acute viral infections. The role of apoptosis in T cell memory. *Journal of Experimental Medicine*, *178*, 427–438.
- Akbar, A. N., Henson, S. M., & Lanna, A. (2016). Senescence of T lymphocytes: Implications for enhancing human immunity. *Trends in Immunology*, *37*, 866–876.
- Allsopp, R. (2008). *Telomeres and Telomerase in Ageing, Disease, and Cancer, Molecular Mechanisms of Adult Stem Cell Ageing*; 23–42.
- Amorim, C., Novais, F., Nguyen, B., Misic, A., Carvalho, L., Carvalho, E., Beiting, D., & Scott, P. (2019). Variable gene expression and parasite load predict treatment outcome in cutaneous leishmaniasis. *Science Translational Medicine*, *11*, eaax4204.
- Andersson, A. K., Sumariwalla, P. F., McCann, F. E., Amjadi, P., Chang, C., McNamee, K., Tornehave, D., Haase, C., Agersø, H., Stenricke, V. W., Ahern, D., Ursø, B., Trowsdale, J., Feldmann, M., & Brennan, F. M. (2011). Blockade of NKG2D ameliorates disease in mice with collagen-induced arthritis: A potential pathogenic role in chronic inflammatory arthritis. *Arthritis Rheumatism*, *63*, 2617–2629.
- Baar, M. P., Brandt, R. M. C., Putavet, D. A., Klein, J. D. D., Derks, K. W. J., Bourgeois, B. R. M., Stryeck, S., Rijksen, Y., van Willigenburg, H., Feijtel, D. A., van der Pluijm, I., Essers, J., van Cappellen, W. A., van IJcken, W. F., Houtsmuller, A. B., Pothof, J., de Bruin, R. W. F., Madl, T., Hoeijmakers, J. H. J., ... de Keizer, P. L. J. (2017). Targeted apoptosis of senescent cells restores tissue homeostasis in response to chemotoxicity and aging. *Cell*, *169*, 132.e16–147.e16.
- Baker, D. J., Childs, B. G., Durik, M., Wijers, M. E., Sieben, C. J., Zhong, J., Saltness, R. A., Jeganathan, K. B., Verzosa, G., Pezeshki, A., Khazaie, K., Miller, J. D., & van Deursen, J. M. (2016). Naturally occurring p16Ink4a-positive cells shorten healthy lifespan. *Nature*, *530*, 184–189.
- Baker, D. J., Wijshake, T., Tchkonja, T., LeBrasseur, N. K., Childs, B. G., van de Sluis, B., Kirkland, J. L., & van Deursen, J. M. (2011). Clearance of p16Ink4a-positive senescent cells delays ageing-associated disorders. *Nature*, *479*, 232–236.
- Blackburn, E. H., Epel, E. S., & Lin, J. (2015). Human telomere biology: A contributory and interactive factor in aging, disease risks, and protection. *Science*, *350*, 1193–1198.
- Braig, M., Lee, S., Loddenkemper, C., Rudolph, C., Peters, A. H. F. M., Schlegelberger, B., Stein, H., Dörken, B., Jenuwein, T., & Schmitt, C. A. (2005). Oncogene-induced senescence as an initial barrier in lymphoma development. *Nature*, *436*, 660–665.
- Brenchley, J. M., Karandikar, N. J., Betts, M. R., Ambrozak, D. R., Hill, B. J., Crotty, L. E., Casazza, J. P., Kuruppu, J., Migueles, S. A., Connors, M., Roederer, M., Douek, D. C., & Koup, R. A. (2003). Expression of CD57 defines replicative senescence and antigen-induced apoptotic death of CD8+ T cells. *Blood*, *101*, 2711–2720.
- Callender, L. A., Carroll, E. C., Beal, R. W., Chambers, E. S., Nourshargh, S., Akbar, A. N., & Henson, S. M. (2018). Human CD8+ EMRA T cells display a senescence-associated secretory phenotype regulated by p38 MAPK. *Aging Cell*, *17*, e12675.
- Callender, L. A., Carroll, E. C., Bober, E. A., Akbar, A. N., Solito, E., & Henson, S. M. (2019). Mitochondrial mass governs the extent of human T cell senescence. *Aging Cell*, *19*, e13067.
- Campisi, J. (2013). Aging, cellular senescence, and cancer. *Annual Review of Physiology*, *75*, 685–705.
- Campisi, J., & di Fagagna, F. (2007). Cellular senescence: when bad things happen to good cells. *Nature Reviews Molecular Cell Biology*, *8*, 729–740.
- Carvalho, A. M., Amorim, C. F., Lago, A. S., Barbosa, J. L. S., & Carvalho, E. M. (2015). Age Modifies the Immunologic Response and Clinical Presentation of American Tegumentary Leishmaniasis. *The American Journal of Tropical Medicine and Hygiene*, *92*, 1173–1177.
- Carvalho, L., Passos, S., Bacellar, O., Lessa, M., Almeida, R., Magalhães, A., Dutra, W., Gollob, K., Machado, P., & Jesus, A. R. D. (2007). Differential immune regulation of activated T cells between cutaneous and mucosal leishmaniasis as a model for pathogenesis. *Parasite Immunology*, *29*, 251–258.



- Chen, Z., Trotman, L. C., Shaffer, D., Lin, H.-K., Dotan, Z. A., Niki, M., Koutcher, J. A., Scher, H. I., Ludwig, T., Gerald, W., Cordon-Cardo, C., & Pandolfi, P. P. (2005). Crucial role of p53-dependent cellular senescence in suppression of Pten-deficient tumorigenesis. *Nature*, *436*, 725–730.
- Childs, B. G., Baker, D. J., Wijshake, T., Conover, C. A., Campisi, J., & van Deursen, J. M. (2016). Senescent intimal foam cells are deleterious at all stages of atherosclerosis. *Science*, *354*, 472–477.
- Chinta, S. J., Woods, G., Demaria, M., Rane, A., Zou, Y., McQuade, A., Rajagopalan, S., Limbad, C., Madden, D. T., Campisi, J., & Andersen, J. K. (2018). Cellular senescence is induced by the environmental neurotoxin Paraquat and contributes to neuropathology linked to parkinson's disease. *Cell Reports*, *22*, 930–940.
- Christensen, S., Dillon, L., Carvalho, L., Passos, S., Novais, F., Hughitt, V., Beiting, D., Carvalho, E., Scott, P., El-Sayed, N., & Mosser, D. (2016). Meta-transcriptome profiling of the human-Leishmania braziliensis cutaneous lesion. *PLoS Neglected Tropical Diseases*, *10*, e0004992.
- Cockburn, I., Tse, S., & Zavala, F. (2014). CD8 + T cells eliminate liver-stage plasmodium berghei parasites without detectable bystander effect. *Infection and Immunity*, *82*, 1460–1464.
- Coppé, J.-P., Desprez, P.-Y., Krtolica, A., & Campisi, J. (2010). The senescence-associated secretory phenotype: the dark side of tumor suppression. *Annu Rev Pathology*, *5*, 99–118.
- Covre, L. P., Martins, R. F., Devine, O. P., Chambers, E. S., Vukmanovic-Stejic, M., Silva, J. A., Dietze, R., Rodrigues, R. R., de Matos Guedes, H. L., Falqueto, A., Akbar, A. N., & Gomes, D. C. O. (2019). Circulating senescent t cells are linked to systemic inflammation and lesion size during human cutaneous leishmaniasis. *Frontiers in Immunology*, *9*, 3001.
- Crosby, E., Clark, M., Novais, F., Wherry, E., & Scott, P. (2015). Lymphocytic choriomeningitis virus expands a population of NKG2D+CD8+ T cells that exacerbates disease in mice coinfecting with leishmania major. *The Journal of Immunology*, *195*, 3301–3310.
- Crosby, E., Goldschmidt, M., Wherry, E., & Scott, P. (2014). Engagement of NKG2D on Bystander Memory CD8 T Cells Promotes Increased Immunopathology following Leishmania major Infection. *PLoS Path*, *10*, e1003970.
- Da-Cruz, A., Oliveira-Neto, M., Bertho, Á., Mendes-Aguiar, C., & Coutinho, S. (2010). T Cells specific to leishmania and other non-related microbial antigens can migrate to human leishmaniasis skin lesions. *The Journal of Investigative Dermatology*, *130*, 1329–1336.
- Demaria, M., Ohtani, N., Youssef, S. A., Rodier, F., Toussaint, W., Mitchell, J. R., Laberge, R.-M., Vijj, J., Van Steeg, H., Dollé, M. E. T., Hoeijmakers, J. H. J., de Bruin, A., Hara, E., & Campisi, J. (2014). An Essential Role for Senescent Cells in Optimal Wound Healing through Secretion of PDGF-AA. *Developmental Cell*, *31*, 722–733.
- Dunne, P. J., Belaramani, L., Fletcher, J. M., de Mattos, S. F., Lawrenz, M., Soares, M. V. D., Rustin, M. H. A., Lam, E.-W.-F., Salmon, M., & Akbar, A. N. (2005). Quiescence and functional reprogramming of Epstein-Barr virus (EBV)-specific CD8+ T cells during persistent infection. *Blood*, *106*, 558–565.
- Faria, D., Souza, P., Durães, F., Carvalho, E., Gollob, K., Machado, P., & Dutra, W. (2009). Recruitment of CD8 + T cells expressing granzyme A is associated with lesion progression in human cutaneous leishmaniasis. *Parasite Immunology*, *31*, 432–439.
- Fletcher, J. M., Vukmanovic-Stejic, M., Dunne, P. J., Birch, K. E., Cook, J. E., Jackson, S. E., Salmon, M., Rustin, M. H., & Akbar, A. N. (2005). Cytomegalovirus-specific CD4 + T cells in healthy carriers are continuously driven to replicative exhaustion. *The Journal of Immunology*, *175*, 8218–8225.
- Franceschi, C., Bonafè, M., Valensin, S., Olivieri, F., Luca, M. D., Ottaviani, E., & Benedictis, G. D. (2000). Inflamm-aging. An evolutionary perspective on immunosenescence. *Annals of the New York Academy of Sciences*, *908*, 244–254.
- Freeman, B., Hammarlund, E., Raué, H., & Slička, M. (2012). Regulation of innate CD8+ T-cell activation mediated by cytokines. *Proceedings of the National Academy of Sciences*, *109*, 9971–9976.
- Gasser, S., Orsulic, S., Brown, E. J., & Raulet, D. H. (2005). The DNA damage pathway regulates innate immune system ligands of the NKG2D receptor. *Nature*, *436*, 1186–1190.
- Gate, D., Saligrama, N., Leventhal, O., Yang, A. C., Unger, M. S., Middeldorp, J., Chen, K., Lehallier, B., Channappa, D., Santos, M. B. D. L., McBride, A., Pluvinage, J., Elahi, F., Tam, G.-K.-Y., Kim, Y., Greicius, M., Wagner, A. D., Aigner, L., Galasko, D. R., ... Wyss-Coray, T. (2020). Clonally expanded CD8 T cells patrol the cerebrospinal fluid in Alzheimer's disease. *Nature*, *577*, 399–404.
- Geginat, J., Lanzavecchia, A., & Sallusto, F. (2003). Proliferation and differentiation potential of human CD8+ memory T-cell subsets in response to antigen or homeostatic cytokines. *Blood*, *101*, 4260–4266.
- Gorgoulis, V., Adams, P. D., Alimonti, A., Bennett, D. C., Bischof, O., Bishop, C., Campisi, J., Collado, M., Evangelou, K., Ferbeyre, G., Gil, J., Hara, E., Krizhanovsky, V., Jurk, D., Maier, A. B., Narita, M., Niedernhofer, L., Passos, J. F., Robbins, P. D., ... Demaria, M. (2019). Cellular senescence: Defining a path forward. *Cell*, *179*, 813–827.
- Goronzy, J. J., Li, G., Yang, Z., & Weyand, C. M. (2013). The Janus Head of T Cell Aging – Autoimmunity and Immunodeficiency. *Frontiers in Immunology*, *4*, 131.
- Groh, V., Rhinehart, R., Randolph-Habecker, J., Topp, M., Riddell, S., & Spies, T. (2001). Costimulation of CD8 $\alpha\beta$  T cells by NKG2D via engagement by MIC induced on virus-infected cells. *Nature Immunology*, *2*, 255–260.
- Gumá, M., Angulo, A., Vilches, C., Gómez-Lozano, N., Malats, N., & López-Botet, M. (2004). Imprint of human cytomegalovirus infection on the NK cell receptor repertoire. *Blood*, *104*, 3664–3671.
- Henson, S. M., Franzese, O., Macaulay, R., Libri, V., Azevedo, R. I., Kiani-Alikhan, S., Plunkett, F. J., Masters, J. E., Jackson, S., Griffiths, S. J., Pircher, H.-P., Soares, M. V., & Akbar, A. N. (2009). KLRG1 signaling induces defective Akt (ser473) phosphorylation and proliferative dysfunction of highly differentiated CD8+ T cells. *Blood*, *113*, 6619–6628.
- Henson, S. M., Lanna, A., Riddell, N. E., Franzese, O., Macaulay, R., Griffiths, S. J., Puleston, D. J., Watson, A., Simon, A., Tooze, S. A., & Akbar, A. N. (2014). p38 signaling inhibits mTORC1-independent autophagy in senescent human CD8+ T cells. *Journal of Clinical Investigation*, *124*, 4004–4016.
- Henson, S. M., Macaulay, R., Riddell, N. E., Nunn, C. J., & Akbar, A. N. (2015). Blockade of PD-1 or p38 MAP kinase signaling enhances senescent human CD8+ T-cell proliferation by distinct pathways. *European Journal of Immunology*, *45*, 1441–1451.
- Henson, S. M., Riddell, N. E., & Akbar, A. N. (2012). Properties of end-stage human T cells defined by CD45RA re-expression. *Current Opinion in Immunology*, *24*, 476–481.
- Jackson, S. E., Redeker, A., Arens, R., van Baarle, D., van den Berg, S. P., Benedict, C. A., Čičin-Šain, L., Hill, A. B., & Wills, M. R. (2017). CMV immune evasion and manipulation of the immune system with aging. *GeroScience*, *39*, 273–291.
- Jeon, O. H., Kim, C., Laberge, R.-M., Demaria, M., Rathod, S., Vasserot, A. P., Chung, J. W., Kim, D. H., Poon, Y., David, N., Baker, D. J., van Deursen, J. M., Campisi, J., & Elisseeff, J. H. (2017). Local clearance of senescent cells attenuates the development of post-traumatic osteoarthritis and creates a pro-regenerative environment. *Nature Medicine*, *23*, 775–781.
- Karin, O., Agrawal, A., Porat, Z., Krizhanovsky, V., & Alon, U. (2019). Senescent cell turnover slows with age providing an explanation for the Gompertz law. *Nature Communications*, *10*, 5495.
- Kritsilis, M., Rizou, S. V., Koutsoudaki, P. N., Evangelou, K., Gorgoulis, V. G., & Papadopoulos, D. (2018). Ageing, cellular senescence



- and neurodegenerative disease. *International Journal of Molecular Sciences*, 19, 2937.
- Krizhanovsky, V., Yon, M., Dickins, R. A., Hearn, S., Simon, J., Miething, C., Yee, H., Zender, L., & Lowe, S. W. (2008). Senescence of activated stellate cells limits liver fibrosis. *Cell*, 134, 657–667.
- Kurundkar, A. R., Kurundkar, D., Rangarajan, S., Locy, M. L., Zhou, Y., Liu, R., Zmijewski, J., & Thannickal, V. J. (2016). The matricellular protein CCN1 enhances TGF- $\beta$ 1/SMAD3-dependent profibrotic signaling in fibroblasts and contributes to fibrogenic responses to lung injury. *The FASEB Journal*, 30, 2135–2150.
- Lanna, A., Henson, S. M., Escors, D., & Akbar, A. N. (2014). The kinase p38 activated by the metabolic regulator AMPK and scaffold TAB1 drives the senescence of human T cells. *Nature Immunology*, 15, 965–972.
- Libri, V., Azevedo, R. I., Jackson, S. E., Mitri, D., Lachmann, R., Fuhrmann, S., Vukmanovic-Stejic, M., Yong, K., Battistini, L., Kern, F., Soares, M. V., & Akbar, A. N. (2011). Cytomegalovirus infection induces the accumulation of short-lived, multifunctional CD4<sup>+</sup> CD45RA<sup>+</sup> CD27<sup>-</sup> T cells: the potential involvement of interleukin-7 in this process. *Immunology*, 132, 326–339.
- Maini, M. K., Soares, M. V., & Zilch, C. F., Akbar, A. N., & Beverley, P. C. (1999). Virus-induced CD8<sup>+</sup> T cell clonal expansion is associated with telomerase up-regulation and telomere length preservation: a mechanism for rescue from replicative senescence. *Journal of Immunology Baltimore Md 1950*, 162, 4521–4526.
- Merad, M., & Martin, J. C. (2020). Pathological inflammation in patients with COVID-19: a key role for monocytes and macrophages. *Nature Reviews Immunology*, 20, 355–362.
- Michel, J. J., Griffin, P., & Vallejo, A. N. (2016). Functionally diverse NK-like T cells are effectors and predictors of successful aging. *Frontiers in Immunology*, 7, 530.
- Molina, H. A., & Kierszenbaum, F. (1989). Interaction of human eosinophils or neutrophils with *Trypanosoma cruzi* in vitro causes bystander cardiac cell damage. *Immunology*, 66, 289–295.
- Moliner, L., Fuentès, M., Girart, M., Fainboim, L., & Rabinovich, G., Costas, M., & Zwirner, N. (2004). NF- $\kappa$ B regulates expression of the MHC class I-related chain A gene in activated T lymphocytes. *Journal of Immunology Baltimore Md (1950)*, 173, 5583–5590.
- Muñoz, D. P., Yannone, S. M., Daemen, A., Sun, Y., Vakar-Lopez, F., Kawahara, M., Freund, A. M., Rodier, F., Wu, J. D., Desprez, P.-Y., Raulet, D. H., Nelson, P. S., van 't Veer, L. J., Campisi, J., & Coppé, J. P. (2019). Targetable mechanisms driving immunoevasion of persistent senescent cells link chemotherapy-resistant cancer to aging. *JCI Insight*, 4, e124716.
- Murray, H. W., Berman, J. D., Davies, C. R., & Saravia, N. G. (2005). Advances in leishmaniasis. *Lancet*, 366, 1561–1577.
- Nassour, J., Martien, S., Martin, N., Deruy, E., Tomellini, E., Malaquin, N., Bouali, F., Sabatier, L., Wernert, N., Pinte, S., Gilson, E., Pourtier, A., Pluquet, O., & Abbadie, C. (2016). Defective DNA single-strand break repair is responsible for senescence and neoplastic escape of epithelial cells. *Nature Communications*, 7, 10399.
- Nikolich-Zugich, J., Ćicin-Šain, L., Collins-McMillen, D., Jackson, S., Oxenius, A., Sinclair, J., Snyder, C., Wills, M., & Lemmermann, N. (2020). Advances in cytomegalovirus (CMV) biology and its relationship to health, diseases, and aging. *Geroscience*, 42, 495–504.
- Oliveira-Ribeiro, C., Pimentel, M., Oliveira, R., Fagundes, A., Madeira, M., Mello, C., Mouta-Confort, E., Valette-Rosalino, C., Vasconcellos, E., Lyra, M., Quintella, L., Antonio, L., Schubach, A., & Conceição-Silva, F. (2017). Clinical and laboratory profiles of patients with early spontaneous healing in cutaneous localized leishmaniasis: a historical cohort study. *BMC Infectious Diseases*, 17, 559.
- Ouyang, Q., Wagner, W. M., Voehringer, D., Wikby, A., Klatt, T., Walter, S., Müller, C. A., Pircher, H., & Pawelec, G. (2003). Age-associated accumulation of CMV-specific CD8<sup>+</sup> T cells expressing the inhibitory killer cell lectin-like receptor G1 (KLRG1). *Experimental Gerontology*, 38, 911–920.
- Ovadya, Y., Landsberger, T., Leins, H., Vadai, E., Gal, H., Biran, A., Yosef, R., Sagiv, A., Agrawal, A., Shapira, A., Windheim, J., Tsoory, M., Schirmbeck, R., Amit, I., Geiger, H., & Krizhanovsky, V. (2018). Impaired immune surveillance accelerates accumulation of senescent cells and aging. *Nature Communications*, 9, 5435.
- Pearson, R. D., & Sousa, A. D. Q., (1996). Clinical Spectrum of Leishmaniasis. *Clinical Infectious Diseases*, 22, 1–13.
- Pereira, B. I., Devine, O. P., Vukmanovic-Stejic, M., Chambers, E. S., Subramanian, P., Patel, N., Virasami, A., Sebire, N. J., Kinsler, V., Valdovinos, A., LeSaux, C. J., Passos, J. F., Antoniou, A., Rustin, M. H. A., Campisi, J., & Akbar, A. N. (2019). Senescent cells evade immune clearance via HLA-E-mediated NK and CD8<sup>+</sup> T cell inhibition. *Nature Communications*, 10, 2387.
- Pereira, B. I., Maeyer, R. P. H. D., Covre, L. P., Nehar-Belaid, D., Lanna, A., Ward, S., Marches, R., Chambers, E. S., Gomes, D. C. O., Riddell, N. E., Maini, M. K., Teixeira, V. H., Janes, S. M., Gilroy, D. W., Larbi, A., Mabbott, N. A., Ucar, D., Kuchel, G. A., Henson, S. M., ... Akbar, A. N. (2020). Sestrins induce natural killer function in senescent-like CD8<sup>+</sup> T cells. *Nature Immunology*, 21, 684–694.
- Pitiyage, G. N., Slijepcevic, P., Gabrani, A., Chianea, Y. G., Lim, K. P., Prime, S. S., Tilakaratne, W. M., Fortune, F., & Parkinson, E. K. (2011). Senescent mesenchymal cells accumulate in human fibrosis by a telomere-independent mechanism and ameliorate fibrosis through matrix metalloproteinases. *The Journal of Pathology*, 223, 604–617.
- Plunkett, F. J., Soares, M. V. D., Anells, N., Hislop, A., Ivory, K., Lowdell, M., Salmon, M., Rickinson, A., & Akbar, A. N. (2001). The flow cytometric analysis of telomere length in antigen-specific CD8<sup>+</sup> T cells during acute Epstein-Barr virus infection. *Blood*, 97, 700–707.
- Rossi, M., & Fasel, N. (2017). How to master the host immune system? Leishmania parasites have the solutions!. *International Immunology*, 30, 103–111.
- Sagiv, A., Burton, D. G., Moshayev, Z., Vadai, E., Wensveen, F., Ben-Dor, S., Golani, O., Polic, B., & Krizhanovsky, V. (2016). NKG2D ligands mediate immunosurveillance of senescent cells. *Aging*, 8, 328–344.
- Santos, C., Boaventura, V., Cardoso, C., Tavares, N., Lordelo, M., Noronha, A., Costa, J., Borges, V., Ci, O., Weyenberg, J., Barral, A., Barral-Netto, M., & Brodskyn, C. (2013). CD8<sup>+</sup> Granzyyme B<sup>+</sup>-mediated tissue injury vs. CD4<sup>+</sup>IFN $\gamma$ <sup>+</sup>-mediated parasite killing in human cutaneous leishmaniasis. *The Journal of Investigative Dermatology*, 133, 1533–1540.
- Savva, G. M., Pachnio, A., Kaul, B., Morgan, K., Huppert, F. A., Brayne, C., & Moss, P. A. H., & Medical Research Council Cognitive Function and Ageing Study (2013). Cytomegalovirus infection is associated with increased mortality in the older population. *Aging Cell*, 12, 381–387.
- Sharpless, N., & Sherr, C. (2015). Forging a signature of in vivo senescence. *Nature Reviews Cancer*, 15, 397–408.
- Tarazona, R., DelaRosa, O., Alonso, C., Ostos, B., Espejo, J., Peña, J., & Solana, R. (2001). Increased expression of NK cell markers on T lymphocytes in aging and chronic activation of the immune system reflects the accumulation of effector/senescent T cells. *Mechanisms of Ageing and Development*, 121, 77–88.
- Tarazona, R., DelaRosa, O., Casado, J. G., Torre-Cisneros, J., Villanueva, J. L., Galiani, M. D., Peña, J., & Solana, R. (2002). NK-associated receptors on CD8 T cells from treatment-naive HIV-infected individuals: defective expression of CD56. *Aids*, 16, 197–200.
- Vallejo, A. N., Mueller, R. G., Hamel, D. L., Way, A., Dvergsten, J. A., Griffin, P., & Newman, A. B. (2011). Expansions of NK-like  $\alpha\beta$ T cells with chronologic aging: Novel lymphocyte effectors that compensate for functional deficits of conventional NK cells and T cells. *Ageing Research Reviews*, 10, 354–361.
- Van Deursen, J. M. (2014). The role of senescent cells in ageing. *Nature*, 509, 439.



- Voehringer, D., Koschella, M., & Pircher, H. (2002). Lack of proliferative capacity of human effector and memory T cells expressing killer cell lectinlike receptor G1 (KLRG1). *Blood*, *100*, 3698–3702.
- Weng, N., Akbar, A. N., & Goronzy, J. (2009). CD28<sup>-</sup> T cells: their role in the age-associated decline of immune function. *Trends in Immunology*, *30*, 306–312.
- Weyand, C. M., Zeisbrich, M., & Goronzy, J. J. (2017). Metabolic signatures of T-cells and macrophages in rheumatoid arthritis. *Current Opinion in Immunology*, *46*, 112–120.
- Wiemann, S. U., Satyanarayana, A., Tsahuridu, M., Tillmann, H. L., Zender, L., Klempnauer, J., Flemming, P., Franco, S., Blasco, M. A., Manns, M. P., & Rudolph, K. L. (2002). Hepatocyte telomere shortening and senescence are general markers of human liver cirrhosis. *The FASEB Journal*, *16*, 935–942.
- World Health Organization/Department of control of neglected tropical diseases (2015). *Investing to overcome the global impact of neglected tropical diseases*. P. Holmes, ed., WHO. Retrieved from [https://www.who.int/neglected\\_diseases/9789241564861/en/](https://www.who.int/neglected_diseases/9789241564861/en/)
- Youn, J.-C., Jung, M. K., Yu, H. T., Kwon, J.-S., Kwak, J.-E., Park, S.-H., Kim, I.-C., Park, M.-S., Lee, S. K., Choi, S.-W., Han, S., Ryu, K.-H., Kang, S.-M., & Shin, E.-C. (2019). Increased frequency of CD4<sup>+</sup>CD57<sup>+</sup> senescent T cells in patients with newly diagnosed acute heart failure: exploring new pathogenic mechanisms with clinical relevance. *Scientific Reports-uk*, *9*, 12887.
- Yu, H. T., Youn, J.-C., Lee, J., Park, S., Chi, H.-S., Lee, J., Choi, C., Park, S., Choi, D., Ha, J.-W., & Shin, E.-C. (2015). Characterization of CD8<sup>+</sup>CD57<sup>+</sup> T cells in patients with acute myocardial infarction. *Cellular & Molecular Immunology*, *12*, 466–473.

**How to cite this article:** Covre LP, De Maeyer RP, Gomes DC, Akbar AN. The role of senescent T cells in immunopathology. *Aging Cell*. 2020;00:e13272. <https://doi.org/10.1111/accel.13272>