



## Saliva Viral Load Better Correlates with Clinical and Immunological Profiles in Children with Coronavirus Disease 2019

Gilbert T Chua MBBS, FHKAM(Paed) , Joshua SC Wong MBBS, FHKAM(Paed) , Kelvin KW To MD, FRCPath , Polly PK Ho MBBS, FHKAM(Paed) , Ivan CS Lam MBChB, FHKAM(Paed) , Jaime SR Duque MD, PhD , Wai Hung Chan MBChB, FHKAM(Paed) , Felix YS Yau MBChB, FHKAM(Paed) , Cyril CY Yip PhD , Anthony CK Ng MPhil , Wilfred HS Wong PhD , Joyce HY Kwong FHKAM(Path) , Kate FS Leung FHKAM(Path) , PT Wan BSc , Kelly Lam MSc , Ian CK Wong PhD, FRCPCH(Hon) , Janette Kwok PhD, FHKAM(Path) , Marco HK Ho MD, FRCP , Godfrey CF Chan MD, FRCPCH(UK) , Yu Lung Lau MD(Hon), FRCPCH , Patrick Ip MPH, FRCPCH(UK) & Mike YW Kwan MSc(Applied Epidemiology) CUHK, FHKAM(Paed)

To cite this article: Gilbert T Chua MBBS, FHKAM(Paed) , Joshua SC Wong MBBS, FHKAM(Paed) , Kelvin KW To MD, FRCPath , Polly PK Ho MBBS, FHKAM(Paed) , Ivan CS Lam MBChB, FHKAM(Paed) , Jaime SR Duque MD, PhD , Wai Hung Chan MBChB, FHKAM(Paed) , Felix YS Yau MBChB, FHKAM(Paed) , Cyril CY Yip PhD , Anthony CK Ng MPhil , Wilfred HS Wong PhD , Joyce HY Kwong FHKAM(Path) , Kate FS Leung FHKAM(Path) , PT Wan BSc , Kelly Lam MSc , Ian CK Wong PhD, FRCPCH(Hon) , Janette Kwok PhD, FHKAM(Path) , Marco HK Ho MD, FRCP , Godfrey CF Chan MD, FRCPCH(UK) , Yu Lung Lau MD(Hon), FRCPCH , Patrick Ip MPH, FRCPCH(UK) & Mike YW Kwan MSc(Applied Epidemiology) CUHK, FHKAM(Paed) (2021): Saliva Viral Load Better Correlates with Clinical and Immunological Profiles in Children with Coronavirus Disease 2019, *Emerging Microbes & Infections*, DOI: [10.1080/22221751.2021.1878937](https://doi.org/10.1080/22221751.2021.1878937)

To link to this article: <https://doi.org/10.1080/22221751.2021.1878937>



© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group, on behalf of Shanghai Shangyixun Cultural Communication Co., Ltd



Accepted author version posted online: 19 Jan 2021.



Submit your article to this journal [↗](#)



Article views: 303



View related articles [↗](#)

---



View Crossmark data [↗](#)

---

**Publisher:** Taylor & Francis & The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group, on behalf of Shanghai Shangyixun Cultural Communication Co., Ltd

**Journal:** *Emerging Microbes & Infections*

**DOI:** 10.1080/22221751.2021.1878937



## Saliva Viral Load Better Correlates with Clinical and Immunological Profiles in

### Children with Coronavirus Disease 2019

Gilbert T Chua<sup>\*1</sup>, MBBS, FHKAM(Paed), Joshua SC Wong<sup>\*2</sup>, MBBS, FHKAM(Paed), Kelvin KW To<sup>\*3</sup>, MD, FRCPath, Polly PK Ho<sup>4</sup>, MBBS, FHKAM(Paed), Ivan CS Lam<sup>2</sup>, MBChB, FHKAM(Paed), Jaime SR Duque<sup>1</sup>, MD, PhD, Wai Hung Chan<sup>4</sup>, MBChB, FHKAM(Paed), Felix YS Yau<sup>4</sup>, MBChB, FHKAM(Paed), Cyril CY Yip<sup>3</sup>, PhD, Anthony CK Ng<sup>3</sup>, MPhil, Wilfred HS Wong<sup>1</sup>, PhD, Joyce HY Kwong<sup>5</sup>, FHKAM(Path), Kate FS Leung<sup>5</sup>, FHKAM(Path), PT Wan<sup>5</sup>, BSc, Kelly Lam<sup>5</sup>, MSc, Ian CK Wong<sup>6,7</sup>, PhD, FRCPCH(Hon), Janette Kwok<sup>8</sup>, PhD, FHKAM(Path), Marco HK Ho<sup>1</sup>, MD, FRCP, Godfrey CF Chan<sup>1</sup>, MD, FRCPCH(UK), Yu Lung Lau<sup>#1</sup>, MD(Hon), FRCPCH, Patrick Ip<sup>#1</sup>, MPH, FRCPCH(UK), Mike YW Kwan<sup>#2</sup>, MSc(Applied Epidemiology) CUHK, FHKAM(Paed)

#### Affiliations

1. Department of Paediatrics and Adolescent Medicine, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, China
2. Paediatric Infectious Disease Unit, Department of Paediatrics and Adolescent Medicine, Princess Margaret Hospital, Hong Kong SAR, China
3. Department of Microbiology, Carol Yu Centre for Infection, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, China

4. Department of Paediatrics, Queen Elizabeth Hospital, Hong Kong SAR, Hong Kong, China
5. Haematology laboratory, Department of Pathology, Princess Margaret Hospital, Hong Kong SAR, China
6. Centre for Safe Medication Practice and Research, Department of Pharmacology and Pharmacy, The University of Hong Kong, Hong Kong SAR, China
7. Research Department of Practice and Policy, UCL School of Pharmacy, University College, London, United Kingdom
8. Division of Transplantation and Immunogenetics, Department of Pathology, Queen Mary Hospital, HKSAR, China

#Co-corresponding authors: **Dr. Mike Yat Wah Kwan**, Consultant, Paediatric Infectious Disease Unit, Department of Paediatrics and Adolescent Medicine, Princess Margaret Hospital, Hong Kong SAR, China, Tel: 852-2990-3243, Email: [kwanyw1@ha.org.hk](mailto:kwanyw1@ha.org.hk)

\*Co-first authors: **Dr. Patrick Ip**, Clinical Associate Professor, Department of Paediatrics and Adolescent Medicine, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, China, Tel: 852-2255-4090, Email: [patricip@hku.hk](mailto:patricip@hku.hk)

**Prof Yu Lung Lau**

Chair Professor of Paediatrics

Doris Zimmern Professor in Community Child Health

Department of Paediatrics and Adolescent Medicine

LKS Faculty of Medicine, The University of Hong Kong

Email: [lauylung@hku.hk](mailto:lauylung@hku.hk)

Tel: (852) 2255-4481

Fax: (852) 2855-1523

Address: Department of Paediatrics & Adolescent Medicine, Room 106, 1/F, New  
Clinical Building, 102 Pokfulam Road, Queen Mary Hospital, Hong Kong

**Conflict of Interest:**

None to declare

**Authors' Contributions**

GT Chua, JSC Wong, KKW To drafted the manuscript. JSC Wong, ISC Lam, JS Rosa Duque, PPK Ho, FYS Yau, WH Chan, YW Kwan recruited the patients and collected the specimen. WHS Wong performed the statistical analysis. Cyril CY Yip and Anthony CK Ng performed the virological laboratory analysis. Kate FS Leung, Joyce HY Kwong, PT Wan and Kelly Lam performed the haematological and immunological analysis. Janette Kwok, Marco HK Ho, Godfrey CF Chan, Yu Lung Lau, Patrick Ip, and Mike YW Kwan critically appraised the manuscript as well as supervised the study. All authors agreed on and finalized the current manuscript.

**Role of Funding Source**

This study was partly supported by the Consultancy Service for Enhancing Laboratory Surveillance of Emerging Infectious Diseases and Research Capability on Antimicrobial Resistance of the Department of Health of the Hong Kong Special

Administrative Region; the donations of Richard Yu and Carol Yu, May Tam Mak Mei Yin, the Shaw Foundation Hong Kong, Michael Seak-Kan Tong.

### **Ethics Committee Approval**

This study was approved by the University of Hong Kong/Hospital Authority Hong Kong West Cluster Institutional Review Board (reference number: UW 20-292), the Kowloon West Cluster Research Ethics Committee (reference number: KW/FR-20-086) and the Kowloon Central Cluster Research Ethics Committee (reference number: KC/KE-20-0140/ER-1).

**Manuscript Word Count:** 2645

### **Abstract**

#### **Background**

Pediatric COVID-19 studies exploring the relationships between NPS and saliva viral loads, clinical and immunological profiles are lacking.

#### **Methods**

Demographics, immunological profiles, nasopharyngeal swab (NPS), and saliva samples collected on admission, and hospital length of stay (LOS) were assessed in children below 18 years with COVID-19.

#### **Findings**

91 patients were included between March and August 2020. NPS and saliva viral loads were correlated ( $r=0.315$ ,  $p=0.01$ ). Symptomatic patients had significantly higher NPS and saliva viral loads than asymptomatic patients. Serial NPS and saliva viral load measurements showed that the  $\log_{10}$  NPS ( $r=-0.532$ ,  $p<0.001$ ) and saliva ( $r=-0.417$ ,  $p<0.001$ ) viral loads for all patients were inversely correlated with the days from symptom onset with statistical significance. Patients with cough, sputum, and headache had significantly higher saliva, but not NPS, viral loads. Higher saliva, but not NPS, viral loads were associated with total lymphopenia, CD3 and CD4 lymphopenia (all  $p<0.05$ ), and were inversely correlated with total lymphocyte ( $r=-0.43$ ), CD3 ( $r=-0.55$ ), CD4 ( $r=-0.60$ ), CD8 ( $r=-0.41$ ), B ( $r=-0.482$ ), and NK ( $r=-0.416$ ) lymphocyte counts (all  $p<0.05$ ).

### **Interpretation**

Saliva viral loads on admission in children correlated better with clinical and immunological profiles than NPS.

### **Introduction**

Since the coronavirus disease 2019 (COVID-19) outbreak in late December 2019, more than 81 million people have been infected globally and more than 1.8 million people have died.<sup>1</sup> Studies in Asia reported the majority of children with COVID-19 had only mild symptoms.<sup>2,3</sup> Children with COVID-19 had various symptoms including fever, pneumonia, and low lymphocyte count that were associated with prolonged viral shedding,<sup>4</sup> whereas their immunoglobulin G (IgG) and neutrophil to lymphocyte ratio were negatively associated with liver and myocardial enzymes.<sup>5</sup> However, clinical studies focusing on the association between clinical outcomes and

immunological profiles in children with COVID-19 are lacking. Furthermore, the association between respiratory tract viral load and immunological profiles has yet to be explored.

Immunological studies showed that adults with COVID-19 with lymphopenia and suppressed T lymphocyte subsets and NK cells were predicted to have a more severe infection.<sup>6-8</sup> Furthermore, severe patients had a higher ratio of conventional dendritic cells to plasmacytoid cells.<sup>9</sup> A recent adult study demonstrated that saliva viral load was reliable for the diagnosis and monitoring of SARS-CoV-2,<sup>10,11</sup> and that saliva was possibly superior to nasopharyngeal swabs (NPS) samples with less variation and false-negative results.<sup>12</sup> However, many studies exploring the viral load burden in children have only used NPS samples.<sup>13</sup> SARS-CoV-2 RNA was detectable during the early phase of infection in mildly symptomatic and symptomatic children,<sup>14</sup> and saliva viral load was also the highest around the time of presentation.<sup>15</sup> Therefore, comparing the clinical utility and correlation of viral loads in NPS and saliva in children with COVID-19 is an important research question that needs to be answered. This study aimed to determine the relationships between the clinical symptoms, immunological profiles, viral loads in NPS and saliva, and hospital length of stay (LOS) in children with COVID-19, as well as compare symptomatic and asymptomatic patients.

## **Methods**

### **Patient recruitment and clinical data collection**



Pediatric patients below 18 years of age who tested positive for SARS-CoV-2 by reverse transcriptase polymerase chain reaction (RT-PCR) of respiratory specimens and admitted to the Princess Margaret Hospital or Queen Elizabeth Hospital in Hong Kong between 12 March and 8 August 2020 were included in the analysis.

Asymptomatic patients, detected by screening at the border or by contract tracing, were also admitted as part of the territory-wide infection control policy in Hong Kong. Demographic details including age, gender, symptoms, travel history, and length of hospital stay were collected. Laboratory tests performed on admission included complete blood count (CBC), renal and liver function tests, immunoglobulin G/A/M (IgG/IgA/IgM) levels, and lymphocyte subsets. NPS was obtained for the detection of the presence of SARS-CoV-2 RNA. NPS and saliva were also collected for viral load measurements on admission and were repeated every five to seven days until discharge. Saliva for viral load quantification was collected for children  $\geq 3$  years old by spitting saliva into a sterile bottle in the morning before brushing teeth and having breakfast. Only NPS will be collected for viral load study if they were unable to provide saliva specimens, especially for younger children.

### **Clinical definitions**

Lymphopenia or lymphocytosis, including subsets, were defined as lower than the 10<sup>th</sup> and higher than the 90<sup>th</sup> percentile based on the Chinese pediatric age and gender-matched reference ranges.<sup>16</sup> Clinical and laboratory profiles were compared between patients with lymphopenia and lymphocytosis. For discharge criteria before 6 July 2020, patients with COVID-19 should have two consecutive negative RT-PCR results for SARS-CoV-2 taken at least 24 hours apart before they could be discharged

from hospital. Patients testing positive had another repeat RT-PCR test every 2 to 3 days until the NPS results were negative. For discharge criteria after 6 July 2020, all patients with COVID-19 were allowed to be discharged from isolation facilities with detectable SARS-CoV-2 anti-nucleoprotein (NP) IgG antibodies in their serum regardless of a positive RT-PCR result of a respiratory specimen.<sup>17</sup> Anti-NP IgG measurements were performed on day 5 or day 10 after admission for asymptomatic patients and symptomatic patients, respectively. If no antibodies were detected, serology was repeated every 2 to 3 days.

### **Laboratory tests**

Nasopharyngeal swabs obtained during admission and for monitoring of the presence of SARS-CoV-2 were tested by RT-PCR using a LightMix<sup>®</sup> Modular SARS and Wuhan CoV E-gene kit (TIB Molbiol, Berlin, Germany) on a LightCycler Multiplex RNA Virus Master (Roche, Penzberg, Germany) according to the manufacturer's instructions as previously described.<sup>18</sup> SARS-CoV-2 viral load analysis of NPS and saliva samples were determined by RT-PCR assay developed by the University of Hong Kong targeting the RNA-dependent RNA polymerase /helicase gene of SARS-CoV-2 (*COVID-19-RdRp/Hel*). Details of the test have been published elsewhere.<sup>19</sup> The lymphocyte subsets were determined by immunophenotyping of peripheral blood lymphocytes using a single platform with a lyse-no-wash procedure on a FC500 flow cytometer (Beckman Coulter, Brea, California, USA). The lymphocyte subsets included total T cells (CD3+), helper T cells (CD3+CD4+), cytotoxic T cells (CD3+CD8+), total B cells (CD19+), and total NK cells (CD3-CD16/CD56+). SARS-CoV-2 anti-NP IgG was measured using a chemiluminescent microparticle immunoassay for

the qualitative detection of IgG in patient serum against the SARS-CoV-2 nucleoprotein (Abbott, Abbott Park, Illinois, USA).<sup>20</sup>

### **Ethics approval**

This study was approved by the University of Hong Kong/Hospital Authority Hong Kong West Cluster Institutional Review Board (reference number: UW 20-292), the Kowloon West Cluster Research Ethics Committee (reference number: KW/FR-20-086) and the Kowloon Central Cluster Research Ethics Committee (reference number: KC/KE-20-0140/ER-1). Individual informed consent was not required as it is a retrospective study and all patients' data were kept anonymous.

### **Statistical analysis**

Statistical analyses were performed using SPSS software version 19 (Armonk, New York, USA), Microsoft Excel (Microsoft, Redmond, Washington, USA) and Statistical Analysis System v9.4 (SAS Institute Inc., Cary, NC). Continuous variables were expressed as mean and interquartile ranges (IQR) and analyzed by independent t-test or Mann-Whitney U test where appropriate. Categorical variables were expressed as number (%) and analyzed by Chi-square test or Fisher exact test. Unpaired T-test was performed to analyze the association between the clinical symptoms and the  $\log_{10}$  of viral load. Pearson correlation was performed to analyze the relationship between the laboratory results and the  $\log_{10}$  viral load. A two-tailed p-value less than 0.05 was considered statistically significant.

### **Results**

## Demographics and Clinical Profiles

Between 12 March and 8 August 2020, 91 children with COVID-19 were admitted to the Princess Margaret Hospital and the Queen Elizabeth Hospital in Hong Kong. The mean age was 9.01 years (IQR 4 - 13 years, range 0.1 - 18 years); 48.4% (44/91) were male; 30.8% (28/91) were asymptomatic. The majority had no significant past medical history. One patient had asthma, one had attention deficit hyperactivity disorder, one had repaired coarctation of the aorta, one had epilepsy, and one had tuberous sclerosis. Most commonly reported symptoms included fever (38.5%; 35/91), cough (23.1%), rhinorrhea (19.8%), and sputum production (12.1%). Five patients reported ageusia and four patients reported anosmia, of which three patients reported suffering from both symptoms. None required intensive care or mechanical ventilation, and no mortality was reported. Among all patients, 28 (30.8%) symptomatic patients received combination antiviral therapy lopinavir/ritonavir (Kaletra®), and nine asymptomatic patients also received empirical treatment. At the time of the analysis, 27 and 56 patients were discharged before and after 6 July 2020, respectively, and eight were still in hospital. Symptomatic patients discharged before 6 July 2020 had significantly longer LOS (27.1 vs. 16.3 days,  $p=0.024$ ). There were no significant differences between symptomatic and asymptomatic patients discharged after 6 July 2020 in terms of LOS (Table 1).

## Immunological Profiles

Based on age- and gender-matched lymphocyte and lymphocyte subset references, 41 patients had total lymphopenia, 20 had CD3 lymphopenia, 15 had CD4

lymphopenia, 24 had CD8 lymphopenia, 19 had B lymphopenia, and 21 had NK lymphopenia. Children with total lymphopenia were significantly older (mean age 10.4 vs. 7.9 years,  $p=0.027$ ). Only a very small number of children had lymphocytosis. There were no significant differences between the lymphocytosis and non-lymphocytosis groups.

### **NPS and Saliva Viral Load Profiles**

On admission, 42 patients had both NPS and saliva viral loads measured. The concordance of NPS and saliva viral load was 100%. There was a significant correlation ( $r=0.315$ ,  $p=0.01$ ) between NPS and saliva  $\log_{10}$  viral loads (Figure 1). Symptomatic patients had significantly higher NPS viral loads (mean  $\log_{10}$  viral load 6.8 vs. 5.7 copies/mL,  $p=0.014$ ) and higher saliva viral loads (mean  $\log_{10}$  viral load 5.8 vs. 4.5 copies/mL,  $p=0.002$ ) compared to those of asymptomatic patients. (Table 1) Patients with cough (mean  $\log_{10}$  viral load 6.4 vs. 4.9 copies/mL,  $p=0.002$ ), sputum production (mean  $\log_{10}$  viral load 6.4 vs. 5.2 copies/mL,  $p=0.036$ ), and headache (mean  $\log_{10}$  viral load 7.3 vs. 5.2 copies/mL,  $p=0.008$ ) had higher saliva viral loads than those without the symptoms (Supplementary file).

### **Correlation between serial viral load measurements and days from symptom onset**

Figure 2 showed the serial NPS and saliva viral loads measurements of all patients against the symptom onset date presented in scatterplot manner. The  $\log_{10}$  NPS ( $r=-0.532$ ,  $p<0.001$ ) and Saliva ( $r=-0.417$ ,  $p<0.001$ ) viral loads for all patients were inversely correlated with the days from symptom onset with statistical significance.

### **Correlation between immunological profiles and viral loads**

Significantly higher saliva viral load was observed in children with total lymphopenia (mean  $\log_{10}$  viral load 5.8 vs. 4.9 copies/mL,  $p=0.031$ ), CD3 lymphopenia (mean  $\log_{10}$  viral load 6.1 vs. 5.1 copies/mL,  $p=0.031$ ), and CD4 lymphopenia (mean  $\log_{10}$  viral load 6.4 vs. 5.1 copies/mL,  $p=0.013$ ) compared to those without lymphopenia (Table 2). Saliva viral load on admission was inversely correlated with total lymphocyte ( $r=-0.43$ ,  $p=0.001$ ), CD3 cell counts ( $r=-0.55$ ,  $p=0.005$ ), CD4 cell counts ( $r=-0.60$ ,  $p=0.002$ ), CD8 ( $r=-0.41$ ,  $p=0.049$ ), B cell counts ( $r=-0.482$ ,  $p=0.017$ ) and NK cell counts ( $r=-0.416$ ,  $p=0.043$ ). There were no significant correlations in NPS viral loads across the immunological profiles or with any blood results (Supplementary File). We performed subgroup analysis for the 42 patients who had both NPS and saliva viral load measured. The correlation remained significant between saliva viral load and total lymphocyte ( $r=-0.381$ ,  $p=0.013$ ), CD3 cell counts ( $r=-0.702$ ,  $p=0.003$ ), CD4 cell counts ( $r=-0.692$ ,  $p=0.004$ ), CD8 cell counts ( $r=-0.710$ ,  $p=0.003$ ) and B cell counts ( $r=-0.515$ ,  $p=0.049$ ), but not between NPS viral load and their lymphocyte subsets.

### **Discussion**

To date, this is one of the largest pediatric COVID-19 studies to explore the relationship between the clinical outcomes, immunological profiles, and NPS and saliva viral loads. Our study demonstrated that saliva viral load, but not NPS viral load, had better correlations with clinical symptoms, hospital LOS and immunological profiles.

Children with respiratory symptoms and headache were more likely to have higher saliva viral loads, but there were no significant differences in their NPS viral loads. Pediatric cross-sectional studies have shown that approximately 30% - 50% children did not present with fever,<sup>2,21</sup> which suggested NPS viral load might not be the best to correlate with the clinical symptoms and immunological profiles. In a recent study by Chong et al with a smaller sample size of 18 children that compared cycle threshold values of SARS-CoV-2 in NPS and saliva, they concluded that saliva was not useful for the diagnosis of children with COVID-19.<sup>22</sup> Conversely, saliva was also shown to be a reliable specimen for the detection of SARS-CoV-2 by RT-PCR,<sup>12</sup> and using saliva for the detection of SARS-CoV-2 RNA had less variability and less false negatives.<sup>23</sup> Yonker et al. reported that children with COVID-19 had significantly higher viral load in NPS compared with in oropharyngeal swabs,<sup>24</sup> but data on saliva viral load was lacking in this study. Nevertheless, obtaining NPS can cause significant discomfort, especially in young children. There is also a risk of generating aerosol, which increases the infection risk in healthcare workers.<sup>25</sup> Therefore, using saliva for the detection of SARS-CoV-2 virus in children could be considered in the future.

Symptomatic patients had significantly higher NPS and saliva viral loads on admission compared to those of asymptomatic patients. Symptomatic patients who were discharged before 6 July 2020 had longer LOS compared to asymptomatic patients. These patients were discharged based on two consecutively negative RT-PCR results from NPS samples, and the asymptomatic group remained hospitalized for a mean duration of 16.3 days. Study has shown that asymptomatic patients had lower infectivity than symptomatic patients,<sup>26</sup> and data from our study also demonstrated

that both NPS and saliva viral load decreased with time. In addition, patients may not be infectious for the entire course of virus detection, as it may not represent infectious viable virus. Future studies should explore the infectivity of asymptomatic children during the later phase of infection by viral culture in order to avoid prolonged hospitalization unnecessarily.

Recent studies reported that adults with severe clinical symptoms had significantly higher viral load in NPS and longer viral shedding period than adults with milder disease.<sup>27</sup> A study also reported that NPS viral load was a predictor for mortality in adults.<sup>28</sup> These studies, however, did not address the association between viral load and clinical severity with respect to immunological profiles, especially in the pediatric population. We observed that patients with total, CD3, and CD4 lymphopenia were more likely to have higher saliva viral load on admission, but not NPS viral load on admission, when comparing with those without lymphopenia. Saliva viral load was inversely correlated with the total, CD3, CD4 and B lymphocyte counts on admission. An earlier study showed that angiotensin-converting enzyme (ACE) 2-positive epithelial cells lining salivary gland ducts were early target cells of SARS-CoV.<sup>29</sup> The salivary gland is an important immune organ that is constantly exposed to pathogens. It contains antigen-presenting dendritic cells, and is a site of B cell activation, isotype switching and the induction of antigen-specific cytotoxic T lymphocytes.<sup>30</sup> Zhou et al. reported that during the acute phase of SARS-CoV-2 infection in adults, despite neutralizing antibodies rapidly and abundantly being generated, there was impaired dendritic cell and T cell immune responses and delayed RBD- and NP-specific T-cell development.<sup>9</sup> Wang et al. demonstrated that



the total lymphocytes, CD4+ and CD8+ T cells, B cells, and NK cells were all decreased in adult COVID-19 patients, which was more prominent in severe patients than in those with mild disease.<sup>8</sup> Zhang et al. also reported that adult patients with T cell lymphopenia had worse clinical outcomes.<sup>31</sup> These clinical findings support the interaction between COVID-19 infection, especially via the salivary glands, and the evasion of the immune system.

Findings in this study need to be interpreted with the following caveats. Not all patients in our cohort received all the assessment on admission. Only 22 patients had both saliva viral load and lymphocyte subset measured, and only 33 patients had both NPS viral load and lymphocyte subset measured. Nevertheless, saliva viral load remained significantly associated with longer LOS after adjusting for treatment. In the subgroup analysis, saliva viral load remained to be significantly correlated with lymphocyte subset and performed better than NPS viral load. None of the patients in our cohort developed severe disease, therefore comparison between mild and severe COVID-19 was not possible. Multiple pediatric COVID-19 studies have reported that the majority of children with COVID-19 were either asymptomatic or suffered only mild symptoms. Saliva viral load was not measured in children younger than three years old, as well as in some pre-school children, because they were unable to provide saliva specimen. Future studies should explore whether using commercially available saliva collection devices for infants are suitable for the detection of SARS-CoV-2.

## **Conclusion**

Saliva viral load appeared to correlate better with clinical symptoms, hospital length of stay and immunological profiles than NPS viral load. Symptomatic children had higher NPS and saliva viral loads on admission, which predicted longer hospital stay regardless of the antiviral treatment. Saliva viral load correlated well with NPS viral load. Patients with total, CD3, and CD4 lymphopenia were more likely to have higher saliva viral load, but not NPS viral load. Both B lymphocyte counts and IgG levels were predictors of hospital LOS. Larger-scale studies are needed in the future to compare the sensitivity and specificity of using NPS versus saliva viral load for the detection of SARS-CoV-2 in children.

### **References**

1. WHO Coronavirus Disease (COVID-19) Dashboard: World Health Organization; 2021. Available from: [https://covid19.who.int/?gclid=CjwKCAiAirbBRBNEiwALHlnDzUZCpkqWlmb-5xmq0d0Z77zf\\_kFIDtAaayJHZhnN0gDX4TNxH6wEBoC2x8QAvD\\_BwE](https://covid19.who.int/?gclid=CjwKCAiAirbBRBNEiwALHlnDzUZCpkqWlmb-5xmq0d0Z77zf_kFIDtAaayJHZhnN0gDX4TNxH6wEBoC2x8QAvD_BwE). Accessed: Jan 1, 2021.
2. Chua GT, Xiong X, Choi EH, Han MS, Chang SH, Jin BL, et al. COVID-19 In Children Across Three Asian Cosmopolitan Regions. *Emerg Microbes Infect.* 2020;1-29.
3. Xiong X, Chua GT, Chi S, Kwan MYW, Sang Wong WH, Zhou A, et al. A Comparison Between Chinese Children Infected with Coronavirus Disease-2019 and with Severe Acute Respiratory Syndrome 2003. *J Pediatr.* 2020;224:30-6.
4. Lu Y, Li Y, Deng W, Liu M, He Y, Huang L, et al. Symptomatic Infection is Associated with Prolonged Duration of Viral Shedding in Mild Coronavirus

Disease 2019: A Retrospective Study of 110 Children in Wuhan. *The Pediatric Infectious Disease Journal*. 2020;39(7).

5. Wu H, Zhu H, Yuan C, Yao C, Luo W, Shen X, et al. Clinical and Immune Features of Hospitalized Pediatric Patients With Coronavirus Disease 2019 (COVID-19) in Wuhan, China. *JAMA Network Open*. 2020;3(6):e2010895-e.
6. Xu B, Fan C-Y, Wang A-L, Zou Y-L, Yu Y-H, He C, et al. Suppressed T cell-mediated immunity in patients with COVID-19: A clinical retrospective study in Wuhan, China. *J Infect*. 2020;81(1):e51-e60.
7. Zhao Q, Meng M, Kumar R, Wu Y, Huang J, Deng Y, et al. Lymphopenia is associated with severe coronavirus disease 2019 (COVID-19) infections: A systemic review and meta-analysis. *International Journal of Infectious Diseases*. 2020;96:131-5.
8. Wang F, Nie J, Wang H, Zhao Q, Xiong Y, Deng L, et al. Characteristics of Peripheral Lymphocyte Subset Alteration in COVID-19 Pneumonia. *J Infect Dis*. 2020;221(11):1762-9.
9. Zhou R, To KK-W, Wong Y-C, Liu L, Zhou B, Li X, et al. Acute SARS-CoV-2 Infection Impairs Dendritic Cell and T Cell Responses. *Immunity*. 2020;53(4):864-77.e5.
10. To KK-W, Tsang OT-Y, Yip CC-Y, Chan K-H, Wu T-C, Chan JM-C, et al. Consistent Detection of 2019 Novel Coronavirus in Saliva. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2020;71(15):841-3.
11. Cheuk S, Wong Y, Tse H, Siu HK, Kwong TS, Chu MY, et al. Posterior oropharyngeal saliva for the detection of SARS-CoV-2. *Clin Infect Dis*. 2020.

12. Azzi L, Carcano G, Gianfagna F, Grossi P, Gasperina DD, Genoni A, et al. Saliva is a reliable tool to detect SARS-CoV-2. *Journal of Infection*. 2020;81(1):e45-e50.
13. Heald-Sargent T, Muller WJ, Zheng X, Rippe J, Patel AB, Kociolek LK. Age-Related Differences in Nasopharyngeal Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Levels in Patients With Mild to Moderate Coronavirus Disease 2019 (COVID-19). *JAMA Pediatrics*. 2020;174(9):902-3.
14. Han MS, Seong M-W, Kim N, Shin S, Cho SI, Park H, et al. Viral RNA Load in Mildly Symptomatic and Asymptomatic Children with COVID-19, Seoul, South Korea. *Emerg Infect Dis*. 2020;26(10):2497-9.
15. To KK, Tsang OT, Leung WS, Tam AR, Wu TC, Lung DC, et al. Temporal profiles of viral load in posterior oropharyngeal saliva samples and serum antibody responses during infection by SARS-CoV-2: an observational cohort study. *Lancet Infect Dis*. 2020;20(5):565-74.
16. Ding Y, Zhou L, Xia Y, Wang W, Wang Y, Li L, et al. Reference values for peripheral blood lymphocyte subsets of healthy children in China. *Journal of Allergy and Clinical Immunology*. 2018;142(3):970-3.e8.
17. Chen P, Nirula A, Heller B, Gottlieb RL, Boscia J, Morris J, et al. SARS-CoV-2 Neutralizing Antibody LY-CoV555 in Outpatients with Covid-19. *New England Journal of Medicine*. 2020.
18. Yip CC-Y, Sridhar S, Cheng AK-W, Leung K-H, Choi GK-Y, Chen JH-K, et al. Evaluation of the commercially available LightMix® Modular E-gene kit using clinical and proficiency testing specimens for SARS-CoV-2 detection. *Journal of Clinical Virology*. 2020;129:104476.

19. Chan JF-W, Yip CC-Y, To KK-W, Tang TH-C, Wong SC-Y, Leung K-H, et al. Improved Molecular Diagnosis of COVID-19 by the Novel, Highly Sensitive and Specific COVID-19-RdRp/Hel Real-Time Reverse Transcription-PCR Assay Validated &em&gt;In Vitro&/em&gt; and with Clinical Specimens. *Journal of Clinical Microbiology*. 2020;58(5):e00310-20.
20. Bryan A, Pepper G, Wener MH, Fink SL, Morishima C, Chaudhary A, et al. Performance Characteristics of the Abbott Architect SARS-CoV-2 IgG Assay and Seroprevalence in Boise, Idaho. *Journal of Clinical Microbiology*. 2020;58(8):e00941-20.
21. Zheng Y, Huang Z, Ying G, Zhang X, Ye W, Hu Z, et al. Study of the lymphocyte change between COVID-19 and non-COVID-19 pneumonia cases suggesting other factors besides uncontrolled inflammation contributed to multi-organ injury. *medRxiv*. 2020:2020.02.19.20024885.
22. Chong CY, Kam K-Q, Li J, Maiwald M, Loo LH, Nadua KD, et al. Saliva is not a useful diagnostic specimen in children with Coronavirus Disease 2019. *Clinical Infectious Diseases*. 2020.
23. Wyllie AL, Fournier J, Casanovas-Massana A, Campbell M, Tokuyama M, Vijayakumar P, et al. Saliva or Nasopharyngeal Swab Specimens for Detection of SARS-CoV-2. *New England Journal of Medicine*. 2020.
24. Yonker LM, Neilan AM, Bartsch Y, Patel AB, Regan J, Arya P, et al. Pediatric SARS-CoV-2: Clinical Presentation, Infectivity, and Immune Responses. *The Journal of Pediatrics*.
25. Harding H, Broom A, Broom J. Aerosol-generating procedures and infective risk to healthcare workers from SARS-CoV-2: the limits of the evidence. *J Hosp Infect*. 2020;105(4):717-25.

26. Zhang W, Cheng W, Luo L, Ma Y, Xu C, Qin P, et al. Secondary Transmission of Coronavirus Disease from Presymptomatic Persons, China. *Emerg Infect Dis.* 2020;26(8):1924-6.
27. Long Q-X, Liu B-Z, Deng H-J, Wu G-C, Deng K, Chen Y-K, et al. Antibody responses to SARS-CoV-2 in patients with COVID-19. *Nature Medicine.* 2020;26(6):845-8.
28. Pujadas E, Chaudhry F, McBride R, Richter F, Zhao S, Wajnberg A, et al. SARS-CoV-2 viral load predicts COVID-19 mortality. *The Lancet Respiratory Medicine.* 2020;8(9):e70.
29. Liu L, Wei Q, Alvarez X, Wang H, Du Y, Zhu H, et al. Epithelial cells lining salivary gland ducts are early target cells of severe acute respiratory syndrome coronavirus infection in the upper respiratory tracts of rhesus macaques. *J Virol.* 2011;85(8):4025-30.
30. Ponzio TA, Sanders JW. The salivary gland as a target for enhancing immunization response. *Tropical Diseases, Travel Medicine and Vaccines.* 2017;3(1):4.
31. Zhang W, Li L, Liu J, Chen L, Zhou F, Jin T, et al. The characteristics and predictive role of lymphocyte subsets in COVID-19 patients. *International Journal of Infectious Diseases.* 2020;99:92-9.

**Table 1.** Comparison between asymptomatic and symptomatic children with COVID-19.

	<b>Asymptomatic</b> Mean (IQR) N= 28	<b>Symptomatic</b> Mean (IQR) N = 63	<b>p-value</b>

Age (years)	8.6 (4.3-11.0)	9.2 (4.0-15.0)	0.658
Male	16 (57.1%)	28 (44.4%)	0.364
Treatment (lopinavir/ritonavir)	9 (32.1%)	19 (30.6%)	1.000
Length of Stay (days)			
Discharged on or before 6 July 2020	16.3 (10.5-21.5)	27.1 (16.5-38.5)	<b>0.024*</b>
Discharged after 6 July 2020	12.4 (11.0-14.0)	11.4 (9.0-14.3)	0.296
NPS (log <sub>10</sub> copies/mL)	5.7 (4.3-7.4)	6.8 (5.6-8.3)	<b>0.014*</b>
Saliva (log <sub>10</sub> copies/mL)	4.5 (3.6-5.1)	5.8 (4.9-6.7)	<b>0.002*</b>
Total white cell count (x 10 <sup>9</sup> /L)	6.0 (7.4-4.1)	5.8 (4.2-6.4)	0.751
Hemoglobin (g/dL)	12.8 (13.5-12.0)	13.2 (14.3-12.4)	0.250
Platelets (x 10 <sup>9</sup> /L)	258.4 (298.0-210.0)	278.1 (317.0-215.0)	0.331
Neutrophil (x 10 <sup>9</sup> /L)	2.5 (1.5-2.9)	2.5 (1.7-3.2)	0.950
Lymphocytes (x 10 <sup>9</sup> /L)	2.9 (1.6-3.8)	2.7 (1.6-2.8)	0.558
Monocyte (x 10 <sup>9</sup> /L)	0.5 (0.3-0.6)	0.5 (0.4-0.6)	0.645
CD3+ T cells (cells/μL)	2049.5 (1165.—2367.0)	1557.1 (1012.0-1690.5)	0.235
CD4+ Helper T cells (cells/μL)	1020.5 (628.0-1238.0)	775.7 (439.8-849.8)	0.221
CD8+ Cytotoxic T Cells (cells/μL)	792.8 (482.0-874.0)	543.5 (358.8-592.0)	0.132
B Cells (cells/μL)	433.1 (209.0-653.0)	359.3 (185.8-372.3)	0.584
Natural Killer Cells (cells/uL)	330.1 (117.0-444.0)	288.6 (109.5-404.8)	0.647
Urea (mmol/L)	3.4 (2.6-4.2)	3.9 (3.0-4.7)	0.081
Creatinine (μmol/L)	41.6 (32.0-49.0)	44.9 (29.0-57.0)	0.383
Creatine Kinase (U/L)	122.5 (69.0-113.0)	99.7 (62.0-117.5)	0.477
Troponin I (ng/L)	1.9 (1.9-1.9)	11.3 (1.9-10.5)	0.198
Lactate Dehydrogenase (U/L)	219.5 (181.0-249.0)	220.0 (179.0-265.0)	0.967
Alanine Aminotransferase (U/L)	18.0 (13.0-22.0)	20.1 (12.0-21.0)	0.543
Alkaline Phosphatase (U/L)	180.7 (113.8-227.8)	194.8 (92.0-237.0)	0.668
Albumin (g/L)	38.9 (37.0-	40.2 (38.0-	0.064

	41.0)	42.0)	
<b>Globulin (g/L)</b>	33.0 (30.0-35.0)	34.0 (32.0-36.3)	0.268
<b>Immunoglobulin G (g/L)</b>	14.9 (9.0-13.5)	11.0 (9.5-13.3)	0.179
<b>Immunoglobulin A (g/L)</b>	1.4 (0.8-1.8)	1.5 (0.8-1.9)	0.571
<b>Immunoglobulin M (g/L)</b>	1.2 (0.7-1.5)	1.3 (0.9-1.7)	0.591
<b>Erythrocyte Sedimentation Rate (mm/hr)</b>	8.6 (2.5-13.0)	12.0 (5.0-15.3)	0.370
<b>C Reactive Protein (mg/dL)</b>	1.4 (0.7-1.0)	1.7 (0.7-2.0)	0.495

- p<0.05

**Table 2.** Comparison of nasopharyngeal swab (NPS) and saliva viral loads for patients with lymphopenia and non-lymphopenia.


		<b>Lymphopenia</b>		<b>Non-lymphopenia</b>		<b>Sig. (2-tailed)</b>
		N	Mean (IQR)	N	Mean (IQR)	
<b>Viral load (Log<sub>10</sub>/copies/mL)</b>	NPS	36	6.7 (5.5-8.3)	42	6.2 (4.3-7.8)	0.187
	Saliva	26	5.8 (4.7-7.0)	28	4.9 (3.9-5.8)	<b>0.031*</b>
		<b>T cell lymphopenia</b>		<b>Non-T cell lymphopenia</b>		<b>Sig. (2-tailed)</b>
		N	Mean (IQR)	N	Mean (IQR)	
<b>Viral load (Log<sub>10</sub>/copies/mL)</b>	NPS	15	7.1 (6.0-8.3)	63	6.3 (4.7-7.9)	0.165
	Saliva	14	6.1 (5.0-7.3)	40	5.1 (3.9-5.8)	<b>0.031*</b>
		<b>CD4 Lymphopenia</b>		<b>Non-CD4 lymphopenia</b>		<b>Sig. (2-tailed)</b>
		N	Mean (IQR)	N	Mean (IQR)	
<b>Viral load (Log<sub>10</sub>/copies/mL)</b>	NPS	11	7.3 (6.7-8.4)	67	6.4 (4.8-7.9)	0.114
	Saliva	10	6.4 (5.0-7.8)	44	5.1 (3.9-5.8)	<b>0.013*</b>

Lymphopenia refers to <10<sup>th</sup> percentile of age and gender-matched reference.

\*p<0.05


**Figure 1.** Correlation analysis between saliva and nasopharyngeal swab (NPS) viral loads





Saliva and NPS viral load were correlated with statistical significance. ( $r=0.315$ ,  $p=0.01$ )

**Figure 2.** Correlation between serial nasopharyngeal swab (NPS) and saliva viral loads, and the days from symptom onset.



The  $\log_{10}$  viral loads for all patients were negatively correlated with the days from symptom onset. (A) NPS ( $r=-0.532$ ,  $p<0.001$ ) (B) Saliva ( $r=-0.417$ ,  $p<0.001$ )

**Supplementary File**

**Table 1.** Comparison of mean nasopharyngeal swab (NPS) viral load and saliva viral load with clinical symptoms.

		Mean NPS viral load (IQR)	p-value	Mean Saliva viral load (IQR)	p-value
Fever	N	6.2 (4.3-7.8)	0.083	5.1 (3.9-5.9)	0.110

	Y	6.9 (5.7-8.3)		5.9 (4.9-6.5)	
<b>Cough</b>	N	6.5 (4.8-8.1)	0.818	4.9 (3.9-5.8)	<b>0.002</b>
	Y	6.6 (5.7-8.1)		6.4 (5.1-7.0)	
<b>Sputum</b>	N	6.4 (4.9-8.1)	0.623	5.2 (3.9-6.1)	<b>0.036</b>
	Y	6.8 (5.7-8.2)		6.4 (5.0-8.7)	
<b>Running nose</b>	N	6.6 (5.2-8.1)	0.233	5.1 (3.9-6.1)	0.055
	Y	6.0 (4.4-7.9)		6.2 (5.0-7.1)	
<b>Sneeze</b>	N	6.5 (5.0-8.1)	0.452	5.3 (4.2-6.2)	-
	Y	5.5 (4.7-)		-	
<b>Stuff nose</b>	N	6.5 (5.0-7.9)	0.486	5.3 (4.2-6.2)	-
	Y	7.2 (4.7-)		-	
<b>Ageusia</b>	N	6.5 (5.1-8.1)	0.257	5.3 (3.9-6.1)	0.322
	Y	5.5 (4.3-6.8)		6.2 (4.7-)	
<b>Anosmia</b>	N	6.5 (4.9-8.1)	0.994	5.3 (3.9-6.1)	<b>0.481</b>
	Y	6.5 (6.0-)		6.0 (4.7-)	
<b>Inappetence</b>	N	6.5 (5.0-8.1)	0.611	5.4 (4.0-6.2)	0.797
	Y	6.2 (4.6-8.3)		5.1 (4.8-)	
<b>Headache</b>	N	6.4 (4.9-8.0)	0.493	5.2 (3.9-6.1)	<b>0.008</b>
	Y	7.0 (5.6-8.2)		7.3 (5.2-9.4)	
<b>Dizziness</b>	N	6.5 (4.9-8.1)	0.745	5.4 (4.4-6.2)	0.444
	Y	6.1 (5.7-)		4.5 (3.6-)	
<b>Vomit</b>	N	6.5 (4.9-8.1)	0.946	5.3 (4.1-6.1)	0.387
	Y	6.5 (4.8-8.0)		6.7 ( )	
<b>Diarrhea</b>	N	6.5 (5.0-8.1)	0.608	5.3 (4.0-6.1)	0.897
	Y	6.1 (4.6-7.5)		5.5 (4.8-)	
<b>Stomachache</b>	N	6.5 (4.9-8.1)	0.647	5.3 (4.2-6.2)	-
	Y	7.1 (6.7-)		-	
<b>Chest pain</b>	N	6.5 (4.9-8.1)	0.917	5.3 (3.9-6.1)	0.076
	Y	6.6 (6.6-)		6.9 (5.4-)	
<b>Tiredness</b>	N	6.5 (5.1-8.1)	0.478	5.3 (3.9-6.2)	0.718
	Y	6.1 (3.8-8.1)		5.6 (4.7-5.3)	
<b>Myalgia</b>	N	6.5 (4.9-8.0)	0.315	5.3 (4.2-6.2)	-
	Y	8.3 ( )		-	


**Table 2. Correlation analysis between NPS viral load and immunological profiles**

	<b>r</b>	<b>p value</b>
<b>Total Lymphocytes</b>	-0.036	0.759
<b>Total T cells</b>	-0.187	0.298
<b>CD4 T Cells</b>	-0.196	0.274
<b>CD8 T Cells</b>	-0.158	0.379
<b>B cells</b>	-0.231	0.196
<b>NK cells</b>	0.032	0.858
<b>Immunoglobulin G</b>	-0.224	0.088
<b>Immunoglobulin A</b>	-0.191	0.147
<b>Immunoglobulin M</b>	-0.068	0.609

**Table 3. Correlation analysis between saliva viral load and immunologicla profiles**

	<b>r</b>	<b>p value</b>
<b>Total Lymphocytes</b>	-0.430	<b>0.001</b>
<b>Total T cells</b>	-0.553	<b>0.005</b>
<b>CD4 T Cells</b>	-0.603	<b>0.002</b>
<b>CD8 T Cells</b>	-0.406	<b>0.049</b>
<b>B cells</b>	-0.482	<b>0.017</b>
<b>NK cells</b>	-0.416	<b>0.043</b>
<b>Immunoglobulin G</b>	-0.236	0.127
<b>Immunoglobulin A</b>	0.062	0.692
<b>Immunoglobulin M</b>	0.032	0.843

**Supplementary Figure 1.** Correlation analysis between  $\log_{10}$  nasopharyngeal swab,  $\log_{10}$  saliva viral load and lymphocyte subsets.



Correlation analysis for (a)  $\log_{10}$  nasopharyngeal swab (NPS) viral load with total lymphocyte count ( $r=-0.36$ ,  $p=0.759$ ); (b)  $\log_{10}$  saliva viral load with total lymphocyte count [ $r=-0.43$ ,  $p=0.001$ ]; (c)  $\log_{10}$  NPS viral load with T cell Count ( $r=-0.187$ ,  $p=0.298$ ); (d)  $\log_{10}$  saliva viral load with T lymphocyte count [ $r=-0.55$ ,  $p=0.005$ ]; (e)  $\log_{10}$  NPS viral load with CD4 count ( $r=-0.196$ ,  $p=0.274$ ) (f)  $\log_{10}$  saliva viral load with CD4 lymphocyte count ( $r=-0.6$ ,  $p=0.002$ ); (g)  $\log_{10}$  NPS viral load with CD8 lymphocyte count ( $r=-0.158$ ,  $p=0.379$ ) (h)  $\log_{10}$  saliva viral load with CD8 lymphocyte count ( $r=-0.41$ ,  $p=0.049$ ) (i)  $\log_{10}$  NPS viral load with B lymphocyte count ( $r=-0.231$ ,  $p=0.196$ ); (j)  $\log_{10}$  saliva viral load with B lymphocyte count ( $r=-0.482$ ,  $p=0.017$ ); (k)  $\log_{10}$  NPS viral load with NK lymphocyte count ( $r=0.032$ ,  $p=0.858$ ) (l)  $\log_{10}$  saliva viral load with NK lymphocyte count ( $r=-0.416$ ,  $p=0.043$ )