

**The impact of COVID-19 on anxiety and wellbeing for families of individuals with
Special Education Needs and Disabilities in the UK**

Sideropoulos, V.¹, Dukes, D.^{2,3}, Hanley, M.⁴, Palikara, O.⁵, Rhodes, S.⁶, Riby, D. M.⁴,
Samson, A. C.^{2,8} & Van Herwegen, J.¹

1. Department of Psychology and Human Development, UCL, Institute of Education, University College London, London, UK
2. Institute of Special Education, University of Fribourg, Switzerland
3. Swiss Center for Affective Sciences, University of Geneva, Switzerland
4. Department of Psychology, Centre for Developmental Disorders, Durham University, UK
5. Department for Education Studies, University of Warwick, Coventry, UK
6. Centre for Clinical Brain Sciences, University of Edinburgh, UK
7. Faculty of Psychology, Unidistance Suisse, Brig, Switzerland

Word count: 8947

Correspondence to: Jo Van Herwegen, UCL Institute of Education, London, UK,
j.vanherwegen@ucl.ac.uk

Abstract

COVID-19 has affected people across the world. However, it has been suggested that individuals with Special Education Needs and Disabilities (SEND) and their families might have been particularly impacted by the first national lockdown in the UK. In contrast to previous studies, the current study examined wellbeing and anxiety at different time points and included a control group matched for family situation. Parents of 402 individuals with SEND reported on their own anxiety and wellbeing as well as that of their son/daughter at three time points (before COVID-19, when COVID-19 pandemic started, and during the national lockdown). In addition, data from 186 typically developing (TD) siblings was obtained. Repeated measures ANOVAs and regression analyses showed that, although both individuals with SEND and their TD siblings showed increased anxiety across the three time points, levels of anxiety were not predicted by age, gender or health. Instead, levels of anxiety in the SEND group, but not the TD siblings, were predicted by awareness about COVID-19, diagnosis of an existing anxiety disorder as well as parental anxiety. In addition, whilst TD individuals were reported to increasingly worry about social related issues as well as family related issues, those with SEND were reported to worry about issues related to school closures. These findings show that COVID-19 impacts the wellbeing of those with SEND differently to that of their TD siblings and that school closures have a particular effect on this group. Further implications for policy impact and interventions are discussed.

Keywords: COVID-19, special educational needs and disabilities (SEND), anxiety, worries, wellbeing, predictors

The COVID-19 pandemic has had an overwhelming impact on the global population. Like other countries across the world, the UK has experienced a significant impact of COVID-19 on people's lives. The first case was reported in the UK on January 31st, 2020 and, as of the 11th January 2021, nearly 12 months later, there have been 3,081,368 cases and 81,567 deaths in the UK alone (Johns Hopkins School of Medicine, 2021). A 'lock-down' was announced on March 23rd lasting over 3 months. During this time, schools closed for all but a few, including those with Special Educational Needs and Disabilities (SEND), and people could only leave their premises for one hour on one occasion per day for activities such as physical exercise or shopping for essential goods, e.g., food, medicines. Different parts of the UK eased lockdown at different times, but generally Scotland, England, Wales and Northern Ireland have followed similar decision-making and time scales during the first strict lockdown. Since the end of the strict lockdown, a tiered system has been introduced with areas moving from a less restrictive way of living to again restrictive lockdowns at various times. This situation means that most children and young people's lives as well as their family's lives have been significantly disrupted by COVID-19.

Special educational needs and disabilities (SEND) arise from a wide range of factors. These can include having a developmental disorder diagnosis such as Autism Spectrum Disorder (ASD), Attention Deficit Hyperactivity Disorder (ADHD), or Dyspraxia/Developmental Coordination Disorder (DCD) that impact learning or a more specific learning difficulty such as dyslexia, dysgraphia or dyscalculia. Individuals who are diagnosed with one of those conditions often show co-occurrences and at times most individuals with SEND meet criteria for more than one disorder. SEND may also arise from early adversity, prematurity or birth trauma. They may also arise from having a chromosomal disorder such as Down or Williams syndrome. In addition, there is a wide range of rare disorders with intellectual impairments for which the aetiology is currently unknown.

Reports have suggested that individuals with SEND are disproportionately impacted by COVID-19. Individuals with SEND require additional support, not only for their learning (Iovannone et al., 2003; Bowles et al., 2015; Rix et al., 2009) but often also to manage day-to-day tasks such as dressing, feeding and personal hygiene, as well as behaviours (Van Herwegen et al., 2019; Van Herwegen et al., 2018). This support is often provided by a range of

professionals within the school and the community. Due to the national lockdown in March 2020 in the UK, a significant number of these services were restricted and disrupted (Jeste et al., 2020; Rose et al., 2020), even for those children with SEND that could attend school during the COVID-19 pandemic (Van Herwegen et al., 2020). In addition, individuals with SEND often have intellectual disabilities that may impact on their understanding of significant events, e.g., COVID-19 pandemic, as a whole (Aishworiya & Kang, 2020) or the new social norms and rules related to such events, e.g., social distancing (Mutluer, Doenyas, & Aslan Genc, 2020). Hence, a number of groups with SEND are at an increased risk of anxiety and mental wellbeing difficulties (Becker et al., 2015; Harrowell, et al., 2017; Nelson & Harwood, 2011; Wigham et al., 2017). Seeing that many of those with SEND often prefer and benefit from a strict and consistent structure (Lindsay et al., 2014) and may not understand the sudden new social rules or reasons behind them (Courtenay & Perera, 2020), COVID-19 disruptions and new social rules may have provided an additional challenge for them. Finally, many groups of individuals with SEND have physical health problems, including congenital heart problems, hypothyroidism, increased risk for pneumonia (Alexander et al., 2020) and thus parents have been reluctant to send their children with SEND to school because of the increased risk of infection (Toseeb et al., 2020).

The difficulties individuals with SEND experience do not only impact themselves, but also their parents by adding to the stress that they have already been experiencing with their children with SEND (Asbury et al., 2020; Ashworth, Palikara, & Van Herwegen, 2019). Various studies report on the fact that families of individuals with SEND feel additional pressures compared to families of children with only typically developing (TD) children (Chafouleas, Iovino, & Koriakin, 2020; Dhiman et al., 2020; Neece, McIntyre, & Fenning, 2020; O'Hagan & Kingdom, 2020). The majority of the parents reported struggling to provide the main care and educational support for the individual with SEND in combination with their work and other childcare commitments. In addition, families reported still needing to receive external support and thus, relied on TD siblings to provide additional care as well (O'Hagan & Kingdom, 2020). Evidence from two different online surveys for parents of children with SEND in the UK suggested that during lockdown the families would have benefited from, among other things, appropriate educational activities set by school, specialist professional

advice for parents, regular structured activities outside home and materials or ideas for school learning (Asbury et al., 2020; Van Herwegen et al., 2020).

Seeing the difficulties experienced by both individuals with SEND and their parents because of COVID-19, it can be assumed that their quality of life is impacted upon. The “Quality-of-life” model proposed by Schalock and Verdugo (2002), includes 8 domains: Social Inclusion, Physical Well-Being, Interpersonal Relations, Material Well-Being, Emotional Well-Being, Self-Determination, Personal Development, and Rights (Schalock, 1996). Reduced quality of life (especially reduced physical wellbeing, social inclusion and reduced interpersonal relations because of COVID-19), are likely to impact on an individual’s wellbeing. Wellbeing can be defined in terms of anxiety as well as worrying about various health and family related matters.

Anxiety is a reaction to a potentially stressful or dangerous situation leading to a physiological response and cognitive processes helping us to deal with the difficult situation. However, high levels of anxiety can impact day to day functioning and ability to cope. High levels of anxiety affect between 7% to 15% of people in the typical population (Biedel & Turner, 2005). Yet, in children anxiety can vary between 2.6% to 41.2% in the general population (Cartwright-Hatton, McNicol, & Doubleday, 2006) averaging around 27% (Costello, Egger, & Angold, 2005). There is evidence that individuals with SEND are at an increased risk of mental health difficulties (Becker et al., 2015; Harrowell, et al., 2017; Nelson & Harwood, 2011; Wigham et al., 2017). This risk is likely to reflect a combination of social factors (Cooper, Smith, & Russell, 2017) and differences in cognitive functions (Rhodes et al., 2012) that are associated with mental health risk, such as difficulties in executive functions (Matthews, Coghill, & Rhodes, 2008).

Emerging evidence suggests heightened levels of anxiety and depression among TD children during the COVID-19 pandemic (Duan et al., 2020; Jiao et al., 2020). Brooks et al. (2020) conducted a review of psychological effects of being isolated during quarantine in the general population and found increased confusion, anger, frustration, boredom, financial loss, and even post-traumatic stress disorder. Orgilés and colleagues (2020), reported that 85.7% of parents surveyed mentioned changes in their TD child’s emotional state, including difficulty

concentrating, boredom, irritability, restlessness, and nervousness and that there was a relationship between parents' anxiety and the child's reported increase in emotional symptoms. Feelings of loss and worry as well as changes in their behaviour have also been reported by parents of children with SEND during the initial phases of lockdown (Asbury et al., 2020). In addition, carers of both children and adults with intellectual disability had significantly greater levels of a wish fulfilment coping style, feelings of defeat/entrapment, anxiety, and depression compared to carers of children without intellectual disability (Chafouleas et al., 2020; Dhiman et al., 2020; Willner et al., 2020).

Although some studies have examined parental anxiety, little is known about how the mental health of individuals with SEND has been affected during these unprecedented times. Combined events of routine changes, school and day activity centre closures, confinement, reduced access to external professionals and interventions are likely to increase the mental health vulnerability of these individuals and their families (Aishworiya & Kang, 2020). There have been reports of increased symptoms amongst individuals with Obsessive Compulsive Disorder (Banerjee, 2020), autistic individuals (Mutluer et al., 2020), ADHD (Zhang et al., 2020), and anxiety disorders more generally (Kwong et al., 2020) during the COVID-19 period. According to O'Hagan and Kingdom (2020), 43% of those surveyed reported that their child with SEND showed increased anxiety at the start of COVID-19 and that their child was struggling with significant mental health problems. However, it is not clear which children are affected the most. For example, 38% of families reported that their child likes to live in their own world and that their child's anxiety had improved during lockdown. In addition, levels of anxiety may also be affected by several individual factors such as age of the individual, gender, their overall health condition which may make them more at risk for COVID-19, their awareness of COVID-19 and whether the child has been diagnosed with an anxiety disorder. For example, it has been suggested that general anxiety disorder increases with age for individuals with Williams syndrome (Dodd, Schniering, & Porter, 2009; Lefeyer et al., 2006). Also, in general females are more likely to experience higher levels of anxiety and are more likely to be diagnosed with an anxiety disorder (McLean et al., 2011).

The current study is the first to our knowledge to examine the wellbeing of entire families of individuals with SEND and to compare the anxiety of carers, their children with

SEND as well as TD siblings during March to June 2020 in the UK. This will allow deeper examination into how the pandemic has affected individuals with SEND differently to TD siblings matched for familial situations and stressors. In addition, the current study did not only gather data about anxiety and worries linked to wellbeing at the time the parents or caregivers were filling out the survey, but also before the pandemic as well as at the beginning of the pandemic in the UK. This allowed examination of how anxiety and worries evolved as a direct result of the pandemic specifically.

The purpose of this paper is to explore the following specific research questions related to anxiety during three key time points; time-point 1 before COVID-19/before March 2020 (Before), time-point 2 when COVID-19 started in March 2020 (Start), and time-point 3 when the survey was completed (Now), which was between April and June 2020 during national lockdown:

1. Whether there is an effect of time (in the three time-points) as well as group (individual with SEND versus TD sibling) on reported anxiety.
2. Which factors predict anxiety when COVID-19 started for individuals with SEND and TD population at the start of the pandemic?
3. How do worries and wellbeing change over the three time-points and how do these changes compare between the groups (individual with SEND versus TD sibling)?

Based on the previous literature discussed above, it was predicted for Research Question 1 that there would be an effect of time and group on the reported anxiety, in that anxiety levels would increase in both groups across the different time points but that individuals with SEND would have higher anxiety at all time points. For Research Question 2, it was hypothesized that predictors such as age, overall health, gender, COVID-19 awareness, parental anxiety and (previous) anxiety diagnosis would predict anxiety levels for individuals with SEND and their TD siblings during the time that COVID-19 started. Lastly, related to worries and wellbeing, we hypothesized that both groups experience increased worries

concerning a range of health, social, school and familial factors across the three time points. However, it was predicted that individuals with SEND would worry more, especially with regards to worries that relate to “changes” such as in social approach and daily structure (e.g., closure of school).

Examining the differences in reported anxiety between individuals with SEND and TD siblings as well as what aspects of wellbeing the two groups worry about, allows greater insight into how the pandemic has disproportionately impacted individuals with SEND as well as which individuals with SEND were affected most. This knowledge will be informative to further tailor interventions and access to support services.

Methods

Participants

Parents and caregivers (5.72% male) of 402 individuals (64.25% male) with SEND completed the survey. The parents and caregivers were aged 21 to 73 years old ($M = 45.47$, $SD = 9.48$) and 235 (58.45%) had a university degree.

The individuals with SEND had a formal diagnosis of a range of learning and developmental difficulties as can be seen in Table 1 and about half of them ($n = 209$, 52%) had an additional diagnosis. The individuals with SEND ranged in age from 1 to 45 years old ($M = 13.04$, $SD = 7.84$). Almost all of the individuals with SEND ($n = 390$, 97%) lived at home during the COVID-19 pandemic, with the remaining 3% living in supported accommodation. Most of these individuals with SEND remained at home as schools and work placements closed during the pandemic and thus, they had their daily routines disrupted (see Table 2).

Table 1.

Overview of Diagnosis of Children with Special Education Needs and Disabilities (SEND)

Type of Diagnosis	Frequency	Percent	Valid Percent
Autism	143	35.57	35.66
Down Syndrome	103	25.62	25.69

Williams Syndrome	43	10.70	10.72
Other Genetic Syndrome	40	9.95	9.98
Intellectual Disabilities	20	4.98	4.99
Attention Deficit Hyperactivity Disorder	18	4.48	4.49
Other	13	3.23	3.24
Learning Difficulties (Dyslexia)	8	1.99	2.00
Anxiety / Mental Health	8	1.99	2.00
Speech and Language Disorder	3	0.75	0.75
Developmental Coordination Disorder	2	0.50	0.50
Missing	1	0.25	
Total	402	100.00	

A sub-sample of the respondents ($n= 186, 46\%$) also reported about a TD sibling (45% male). The TD siblings had a similar age range (1-47 years, $M= 14.84, SD= 8.96$) but were slightly older than the individuals with SEND; $t(397)= 33.179, p < .001$. Most of the TD siblings ($n= 159, 85\%$) lived at home with their family and a large proportion remained at home due to COVID-19 closures of school and workplaces (see Table 2 and Table 3).

Table 2.

Overview of Daily Routines for Individuals with Special Educational Needs and Disabilities

	Frequency	Percent	Valid Percent
At home with family	325	80.85	81.45
Mainstream School	30	7.46	7.52
Special Education School	20	4.98	5.01
Other	11	2.74	2.76
Pre-school	6	1.49	1.50
College	6	1.49	1.50
Work in Community / Volunteer Sector	1	0.25	0.25
Missing	3	0.75	
Total	402	100.00	

Table 3.

Overview of Daily Routines for Typically Developing Individuals

	Frequency	Percent	Valid Percent
At home with family	105	26.12	26.12
Mainstream school	39	9.70	9.70
Paid full time/Part time work	21	5.22	5.22
Other	14	3.48	7.33
Work in community/Volunteer sector	3	0.75	0.75
Working in protected environment	2	0.50	0.50
Pre-school	1	0.25	0.25
Special school	1	0.25	0.25
Missing	216	53.73	53.73
Total	186	100.00	

Further background about the individuals with SEND

Most individuals with SEND had mild to severe intellectual disabilities (n= 306, 76.12%) and thus, a proportion of individuals with SEND (n= 123, 30.60%) were reported to not be aware of COVID-19. Although these individuals may not have understood that any changes to routine might be caused by COVID-19, they may still have experienced higher anxiety as a result of the current pandemic and thus we included them in our analyses about anxiety (but not about worries). About a quarter of individuals with SEND (n= 133, 25.62%) had been reported to have an anxiety disorder. A minority of individuals with SEN (n= 31, 7.71%) were thought to have been affected by COVID-19 but only 2 were tested and needed hospital treatment.

Background about the TD siblings

Most of the TD siblings were healthy with only 30 TD siblings (16%) reporting any medical issues of which only 6 had severe to serious medical issues (3%). For a minority (n= 28, 15.05%) parents reported an anxiety disorder. Again, a minority of TD siblings were thought to have been affected by COVID-19 (n= 11, 5.91%) but none had been tested or needed hospital treatment. Most of the TD siblings (n= 170, 91.40%) were aware of COVID-19.

Materials

An anonymous survey was distributed using Qualtrics, an online survey tool, which contained a range of open-ended and closed questions over four sections. This survey was used in the context of a larger international collaboration (www.specialneeds covid.org) with over 10,000 families of individuals with SEND across the globe. Three of these sections were relevant to the current study. The entire survey can be accessed freely on the OSF website (Van Herwegen, Dukes, & Samson, 2020): <https://osf.io/5nkq9/>.

Section A asked questions about the respondent and their SEND child's background, including demographic information and the SEND child's medical background.

Section B included questions about the timing of the events related to COVID-19, including when the child's school closed, or their daily routine changed because of lockdown.

Section C focused on the concerns and wellbeing of the participating parent/guardian, the concerns and wellbeing of their child with SEND and (if they also had a typically developing child) the worries and wellbeing of a TD child. For each question, the respondent was asked to provide an answer related to three different time points: before the pandemic began (Before), when COVID-19 first affected them in March 2020 (Start), and at the time the participant completed the survey (Now), which was between April and June 2020. To measure anxiety, parents were asked to identify on a scale from 1 to 5 (with 1 not anxious at all to 5 being very anxious) their own anxiety, the level of anxiety that their child with SEND experienced and the anxiety of the TD child (if they had one).

The questions around worries were informed by the wellbeing categories as defined by Schalok (1996) and included worries related Social Inclusion (e.g., not being able to meet others), Physical Wellbeing (e.g., worries about catching COVID-19 and own health), Interpersonal Relations (e.g., worry about family conflict and others becoming ill), Material Wellbeing (e.g., financial worries), Emotional Wellbeing (e.g., worries about boredom), Self-Determination (e.g., loss of routine), and Personal Development (e.g., loss of institutional support). These were grouped into the following categories: Health Related Worries, Social Related Worries, School Closure Related Worries and Family Related Worries.

Procedure

Participants were recruited through social networks, social media and by emails addressed to special education institutions and support groups throughout the UK, including Williams Syndrome Foundation, Down Syndrome Association UK, ADHD Foundation. In addition, flyers and posters were also made available via Twitter and Facebook support groups.

Participants completed the survey between 8th of April 2020 and 27th of June 2020. Participants were not reimbursed for their time and participation was entirely voluntarily as well as anonymous.

Ethics

Ethical approval for the study was obtained from Ethics Commission of UniDistance, Switzerland before the start of the study. Participants provided written consent to take part in the online study.

Results

Anxiety

Effect of Time on Anxiety for Children with SEND and TD Children

To determine the effect of time on anxiety for children with SEND and TD children, a 3 (Time) x 2 (Group) mixed-model ANOVA was conducted. There was a significant main effect for Time. Mauchly's Test of Sphericity indicated that the assumption of sphericity had been violated: $\chi^2(2) = 73.61, p < .001$. Therefore, the degrees of freedom had to be adjusted by using the Huynh-Feldt correction ($\epsilon = .89$); $F(1.79, 1023.54) = 180.50, p < .001, \eta^2_p = .24$. This effect tells us that there was a difference in the reported anxiety for both groups over time. Post-hoc comparisons were computed and a difference between all time-points was evident (see Table 4).

Table 4.

Post Hoc Comparisons - Effect of Time on Anxiety for Children with SEND and TD Children

		Mean Difference	95% CI for Mean Difference		SE	t	Cohen's d	p bonf
			Lower	Upper				
Before	Start	-0.50	-0.61	-0.40	0.04	-11.64	-0.49	< .001 ***
	Now	-0.81	-0.92	-0.71	0.04	-18.83	-0.79	< .001 ***
Start	Now	-0.31	-0.41	-0.21	0.04	-7.19	-0.30	< .001 ***

*** $p < .001$

Note. Cohen's d does not correct for multiple comparisons.

Note. P-value and confidence intervals adjusted for comparing a family of 3 estimates (confidence intervals corrected using the Bonferroni method).

Note. Results are averaged over the levels of: Group

There was also a main effect for Group, $F(1,572)= 13.12, p < .001, \eta^2_p= .02$ meaning there was a difference between the groups' reported anxiety. Post-hoc analyses were conducted, individuals with SEND scored higher anxiety ($M=2.71$) compared to the TD group ($M=2.33$) (see Table 5). Figure 1 presents the mean difference over time for both groups.

Table 5.

Post Hoc Comparisons for Group: Anxiety for Children with SEND and TD Children

		Mean Difference	95% CI for Mean Difference		SE	t	Cohen's d	p bonf
SEND	TD		Lower	Upper				
		0.38	0.17	0.59	0.10	3.62	0.15	< .001 ***


*** $p < .001$

Note. Results are averaged over the levels of: Time

Note. Cohen's d does not correct for multiple comparisons.

Figure 1.

Overall Anxiety for SEND and TD groups at the three time points: Before (before March 2020), Start (from March 2020), Now (between April- June 2020)



Our mixed model indicated that there was not a significant Time*Group interaction. Mauchly's Test of Sphericity showed that the assumption of sphericity had been violated: $\chi^2(2) = 73.61, p < .001$. The degrees of freedom had to be adjusted by using the Huynh-Feldt correction ($\epsilon = .89$); $F(1.79, 1023.54) = 2.32, p = .10, \eta^2_p = 4.04e-3$.

Predictors of Anxiety

Linear Regression to Predict Anxiety for SEND

A multiple linear regression was run to predict anxiety at the start of the pandemic (time-point 2) from age, gender, health, whether they were aware of COVID-19 and whether they had a diagnosis of anxiety prior to COVID-19 as well as parental anxiety for the same time-point for the SEND group. The data was screened for assumptions and outliers and all assumptions of linearity, normality, homoscedasticity, and multicollinearity were found to be met. The multiple regression model statistically significantly predicted Anxiety levels at the beginning of the pandemic, $F(6, 385) = 43.54, p < .001$. Not all six variables added statistically significantly to the prediction. Regression coefficients and standard errors can be found in Table 6. Age, gender and health status did not add anything to the model. However, parental anxiety, anxiety disorder and COVID-19 awareness were all positively predicted Anxiety. SEND children of anxious parents were more likely to show higher anxiety as well as those who were diagnosed with an anxiety disorder and were aware of COVID-19 in the beginning of the COVID-19 pandemic.

Table 6.

Coefficients for Linear Regression to Predict Anxiety for SEND

Model		Unstandardized	Standard Error	Standardized	t	p	Collinearity Statistics	
							Tolerance	VIF
H ₀	(Intercept)	2.71	0.07		39.66	< .001		
H ₁	(Intercept)	-0.70	0.40		-1.77	0.08		
	Age	0.01	7.14e-3	0.07	1.69	0.09	0.90	1.11
	Health Status	-0.05	0.06	-0.03	-0.82	0.41	0.94	1.06

Coefficients for Linear Regression to Predict Anxiety for SEND

Model	Unstandardized	Standard Error	Standardized	t	p	Collinearity Statistics	
						Tolerance	VIF
Parental Reported Anxiety (Time-point 2)	0.20	0.05	0.18	4.47	< .001	0.98	1.02
Gender	-0.19	0.11	-0.07	-1.66	0.10	0.99	1.01
Anxiety Disorder	1.30	0.12	0.45	10.65	< .001	0.85	1.17
COVID-19 Awareness	0.80	0.13	0.27	6.32	< .001	0.83	1.20

Linear Regression to Predict Anxiety for TD

Similarly, a multiple linear regression was run to predict anxiety at the start of the pandemic (time-point 2) from age, gender, health status, whether they were aware of COVID-19, whether they had a diagnosis of an anxiety disorder prior to COVID-19 and parental anxiety for the same time-point for the TD group. The data were screened for assumptions and outliers, no outliers and all assumptions of linearity, normality, homoscedasticity and multicollinearity were met. The multiple regression model did not statistically significantly predict Anxiety levels when the pandemic started, $F(6,24) = 1.06$, $p = .41$. Table 7 presents the regression coefficients and standard errors of our model which show zero significant correlations between the variables which leads to the conclusion that none of these factors contributed to the anxiety levels of the TD group when COVID-19 started.

Table 7.

Coefficients for Linear Regression to Predict Anxiety for TD

Model	Unstandardized	Standard Error	Standardized	t	p	Collinearity Statistics	
						Tolerance	VIF
H ₀	(Intercept)	2.48	0.21	11.72	< .001		
H ₁	(Intercept)	-0.15	3.27	-0.05	0.96		
	Age	4.22e -3	0.02	0.04	0.17	0.87	0.74 1.35
	Health Status	-0.06	0.33	-0.03	-0.17	0.86	0.94 1.07
	Gender	0.40	0.50	0.16	0.81	0.43	0.88 1.14
	Anxiety Disorder	0.78	0.59	0.28	1.31	0.20	0.72 1.39
	COVID-19 Awareness	0.06	1.05	0.01	0.05	0.96	0.67 1.50
	Parental Reported Anxiety (Time-point 2)	0.36	0.21	0.35	1.76	0.09	0.85 1.18

Worries and Concerns related to Wellbeing

For the analyses related to worries, only those individuals with SEND (n= 279) and the TD siblings who were aware of COVID-19 (n= 173) were included in the repeated measures 3(Time)*2(Group) analyses, given that awareness of COVID-19 was an important predictor for anxiety in the SEND group. Sphericity violations for each model are presented in Table 8. The change over time for both SEND and TD in worries and concerns is presented in Figure 2. For a detailed overview of the mean scores for each category in the worries and concerns, see Supplementary Table 9.

Figure 2. Worries over Time for the SEND and TD groups




Table 8.

Sphericity violations for Worries and Concerns related to Wellbeing

Type of Concern	Test of Sphericity Checks	Adjustment Method and ϵ level
	Concerns about COVID-19	$\chi^2 (2) = 27.37, p < .001$ Huynh-Feldt correction ($\epsilon=.94$)
	Concerns about Others becoming ill	$\chi^2 (2) = 35.37, p < .001$ Huynh-Feldt correction ($\epsilon=.92$)
Health-related worries	Concerns about safety	$\chi^2 (2) = 27.36, p < .001$ Huynh-Feldt correction ($\epsilon=.94$)
	Concerns about health	$\chi^2 (2) = 32.66, p < .001$ Huynh-Feldt correction ($\epsilon=.92$)
	Concerns about friends	$\chi^2 (2) = 55.44, p < .001$ Huynh-Feldt correction ($\epsilon=.88$)
Social-related worries	Concerns about approach	$\chi^2 (2) = 93.78, p < .001$ Huynh-Feldt correction ($\epsilon=.82$)
	Concerns about changes in routine	$\chi^2 (2) = 37.94, p < .001$ Huynh-Feldt correction ($\epsilon=.91$)
School-related worries	Concerns about getting bored	$\chi^2 (2) = 85.18, p < .001$ Huynh-Feldt correction ($\epsilon=.83$)
	Concerns about Loss of Institution (school closure)	$\chi^2 (2) = 70.85, p < .001$ Huynh-Feldt correction ($\epsilon=.85$)
Family-related worries	Concerns about Family	$\chi^2 (2) = 121.75, p < .001$ Huynh-Feldt correction ($\epsilon=.78$)

Concerns about Finance	$\chi^2(2) = 71.98, p < .001$	Huynh-Feldt correction ($\epsilon = .85$)
------------------------	-------------------------------	---

Health-related worries

For the “Concerns about COVID-19”, our model revealed that there was an effect for Group; $F(1,565) = 4.71, p = .03, \eta_p^2 = 8.27e-3$. Similarly, there was a significant effect for Time; $F(1.87,676.11) = 174.24, p < .001, \eta_p^2 = .33$. However, the interaction for Time*Group was not significant; $F(1.87,676.11) = 0.14, p = .85, \eta_p^2 = 4.01e-4$. Concerns about COVID-19 increased over time for both groups, but those with SEND were more concerned across all time points.

With regards to “Concerns about Others Becoming Ill”, our mixed model showed that there was no effect for Group; $F(1,360) = 0.03, p = 0.86, \eta_p^2 = 9.16e-5$, while there was an effect for Time; $F(1.84,661.43) = 153.08, p < .001, \eta_p^2 = .30$. Similarly, the interaction for Time*Group was significant; $F(1.84,661.43) = 4.09, p = 0.02, \eta_p^2 = .01$. As can be seen in Figure 2, the TD group scored higher concerns about others becoming ill when the parents completed the survey (time point 3), while the SEND group showed higher concerns when the COVID-19 pandemic started (time point 2) and their concerns decreased afterwards.

When investigating for “Concerns about Safety”, no effect for Group was evident; $F(1,362) = 6.37e-3, p = 0.94, \eta_p^2 = 1.76e-5$. On the other hand, there was an effect for Time: $F(1.87,678.10) = 156.56, p < .001, \eta_p^2 = .30$. For the interaction Time*Group, there was no significant effect; $F(1.87,678.10) = 2.37, p = .10, \eta_p^2 = 6.52e-3$. While there was a difference in Concerns about Safety between time-points, yet this difference was not significant between the Groups.

Our model showed that there was no Group effect for “Concerns about Health”; $F(1,361) = 2.97, p = .09, \eta_p^2 = 8.15e-3$. Nonetheless, there was an effect for Time; $F(1.85,667.63) = 74.49, p < .001, \eta_p^2 = .17$. When examining for a Time*Group interaction, no effect was shown; $F(1.85,667.63) = 0.23, p = .78, \eta_p^2 = 6.27e-4$. Similar, to other health concerns, concerns increased with time, but there was no difference between the TD and SEND groups.

Social-related worries

Related to “Concerns about Friends”, the mixed model showed no significant effect for Group; $F(1,361) = 2.17, p = .14, \eta^2_p = 5.97e-3$. Yet, an effect for Time; $F(1.76,634.64) = 176.20, p < .001, \eta^2_p = .33$. Similarly, there was a significant interaction for Time*Group; $F(1.76,634.64) = 3.32, p < .004, \eta^2_p = 9.11e-3$. In both groups concerns about friends increased over time but worries increased more in the TD group when the pandemic started and continued to increase exponentially.

When investigating for “Concerns about Approach”, there was no overall effect for Group; $F(1,362) = 0.25, p = .61, \eta^2_p = 7.04e-4$. However, there was a Time effect; $F(1.63,591.49) = 102.57, p < .001, \eta^2_p = .22$. No significant interaction between Time*Group; $F(1.63,591.49) = 1.58, p = .21, \eta^2_p = 4.36e-3$. As can be seen in Figure 2, in both groups worries related to approaching others increased with time, but there was not a difference between the TD and SEND groups.

Worries related to school closures

For ‘Worries about Change in Routine’, there was an effect for Group; $F(1,362) = 25.03, p < .001, \eta^2_p = .06$ as well as for Time; $F(1.83,661.47) = 79.11, p < .001, \eta^2_p = .18$. However, the interaction between Time*Group did not produce an effect; $F(1.83,661.47) = .87, p = .41, \eta^2_p = 2.41e-3$. There is an increase for Concerns about Change in Routine over time in both groups but those with the SEND reported higher concerns at each of the time points.

For ‘Concern about Getting Bored’, there was not an overall effect for Group; $F(1,361) = 0.67, p = .41, \eta^2_p = 1.86e-3$. But there was an effect for Time; $F(1.66,598.74) = 134.18, p < .001, \eta^2_p = .27$. For the Time*Group interaction, no effect was noticed; $F(1.66,598.74) = 3.03, p = .06, \eta^2_p = 8.33e-3$. As can be seen in Figure 2, there is an increased concern about getting bored over time for both groups.

In terms of “Concerns about the school closure or loss of institutional support”, there was a Group effect; $F(1,360) = 9.15, p < 2.66e-3, \eta^2_p = .02$ and a Time effect; $F(1.70,631.22) = 101.21, p < .001, \eta^2_p = .22$. However, there was not an interaction effect for Time*Group;

$F(1.70,613.22) = 0.17, p = .81, \eta^2_p = 4.68e-4$. In both groups concerns about school closure increased over time, but more so for those in the SEND group.

Family related worries

Relating to “Concerns about family”, there was no Group effect; $F(1,361) = .18, p = .67, \eta^2_p = 4.99e-4$. However, there was a Time effect; $F(1.56,563.00) = 35.16, p < .001, \eta^2_p = .08$. For the Time*Group interaction; the adjusted model indicated no interaction: $F(1.56,563) = 0.52, p = .55, \eta^2_p = 1.44e-3$. While there was an increase of concerns about family over time, both Groups performed similarly.

Finally, there was a Group effect for “Concerns about Finances”; $F(1,359) = 7.96, p < 5.05e-3, \eta^2_p = .02$ as well as an effect for Time; $F(1.70,609.94) = 32.75, p < .001, \eta^2_p = .08$. Similarly, there was a significant interaction between Time*Group; $F(1,359) = 7.96, p < 5.05e-3, \eta^2_p = .02$ for concerns around finances. The TD group worried more about finances and their concerns increased more over time compared to the SEND group.

Discussion

The current study examined the impact of the early stages of COVID-19 in the UK on anxiety and wellbeing of individuals with a wide range of SEND in relation to parental anxiety and compared to that of a group of TD siblings. In addition, predictors of anxiety were also examined for both the SEND and TD groups as well as what kind of aspects related to COVID-19 both groups worried about. Together these analyses allowed for a better understanding of the specific impact of COVID-19 on the wellbeing of individuals with SEND compared to TD populations and which individuals with SEND are most affected.

Impact of COVID19 on Anxiety in SEND and TD groups

In terms of overall reported anxiety, both groups increased over time. However, whilst the SEND group worried more at all three time points, anxiety increased more in the TD group. These findings are in line with previous studies that have reported increased anxiety in those with special needs (O’Hagan & Kingdom, 2020). However, the current study adds to previous studies, in that direct comparisons to siblings, showed that anxiety tended to be higher across all time points for those with SEND.

Anxiety predictors for SEND and TD

Multiple regression analyses were run to explore what factors predicted anxiety at the start of the pandemic in March 2020. For the SEND group, higher levels of parental anxiety, a diagnosis of an anxiety disorder, and awareness of COVID-19 seemed to be strong predictors of anxiety during that time-point. Factors such as age, gender and health status did not seem to drive anxiety at all in contrast to predictions. The same analysis was run for TD group, but the results indicated that none of the factors were strong predictors of anxiety at this time point. This might suggest that anxiety during COVID-19, in contrast to general contexts, might be driven by specific factors that are not usually as important. We discuss some of the limitations of the study below, but these findings suggest that predictors of anxiety during a pandemic might be different and thus replication and further investigation of what predicts anxiety during a pandemic, including factors not yet explored (e.g., IQ and social support), is required.

The present findings also suggest that, although anxiety increases in both groups, there are differences within the SEND and TD groups concerning which individuals are at greater risk or have heightened anxiety as a result of COVID-19. Ours is the first study to provide evidence that COVID-19 awareness can predict anxiety in individuals with SEND. Similarly, our study is in line with both Platt, Williams, & Ginsburg (2016) and the recent study by Russell et al., (2020) which show that increased parental anxiety is linked with the severity of anxiety symptoms in their child, but also with the perceptions of children's stress.

Worries about the impact of COVID-19 in SEND and TD groups

In terms of worries, both groups showed increased concerns across the three time points for all types of worries. However, there were some interesting group differences and similarities. In line with the predictions, individuals with SEND worried more about COVID-19, whilst the TD group had a larger increase related to becoming ill more specifically. These differences could be explained by the fact that many individuals with SEND shielded (Van Herwegen et al., 2020) and that parents have been reluctant to send their children with SEND to school because of the increased risk of infection (Toseeb et al., 2020). As such, those with SEND might have been less worried about becoming ill but rather worried about COVID-19 more generally. Both groups showed increased worries related to general health but concerns

around safety decreased. The latter could be explained by the fact that most individuals in both groups remained at home during the lockdown.

Social related worries also increased in both groups, but the TD group showed a greater increase related to not being able to see friends. However, worries around social approach did not differ between the two groups and both groups showed increased worries over time. Finally, the TD group showed greater concerns related to family related concerns, especially finances.

In relation to school closures and closures of day or activity centres, both groups' concerns increased over the three time points. There were no differences between the two groups with regards to boredom. This finding is similar to studies in the USA related to TD individuals that reported that schools provide important structured activities that occupy TD individuals from becoming bored (Jeste et al., 2020; Orgilés et al., 2020). The current study showed that the same is true for those with SEND. However, the SEND group had higher levels of reported concerns related to a lack of structure across the three time points and showed a greater increase of concern over the time points related to loss of support because of school and activity centres' closures. This replicates the findings of O'Hagan and Kingdom, (2020) but contrasting the worries of those with SEND to TD siblings, allowed further understanding of the unique worries of those with SEND.

Limitations and future studies

The current study examined the wellbeing and anxiety levels of individuals with SEND and TD siblings through parental report rather than through the voice of the individual with SEND or TD sibling themselves. Research has shown significant links between parents' mental health and the way they perceive their children's wellbeing as well as how parents' mental health of disastrous events, e.g., COVID-19, can affect children's understanding of such situations (Neece, Green, & Baker, 2012; Orgiles et al., 2020). Therefore, there is some bias to be expected on the reported data of anxiety of the children.

In addition, although the current study examined for the first time several factors that may explain individual differences in each group related to the impact of COVID-19 on anxiety and wellbeing, it was not possible to examine differences related to the primary diagnosis of the individuals with SEND because uneven groups were recruited and 70% of the individuals with SEND had one of the following developmental disorders: autism, Down syndrome or Williams syndrome, and some conditions were under-represented (e.g., socio-emotional and

behavioral difficulties). However, previous studies that have examined outcomes in those with SEND have shown that categorical labels may not be informative with regards to the needs of the individual with SEND (Dockrell et al., 2019). Instead, examination of individual differences such as pre-existing health and anxiety diagnoses might be a more useful approach to examine those most affected or in need. Yet, our worldwide data of more than 10 000 families that completed the survey will allow further examination of individual differences related to health and physical needs as well as country specific variables (e.g., types of measures to deal with the pandemic, timing of restrictions, support available).

Impact and conclusion

Together, these findings show that COVID-19 impacts the wellbeing of those with SEND differently to that of their TD siblings and that school closures have a particular effect on those with SEND. For individuals with SEND, schools and activity centers provide an important routine and structure that helps to reduce anxiety and improves wellbeing (Van Herwegen et al., 2020). In addition, schools and activity centers provide parents with access to specialist advice that would not only benefit individuals with SEND but also their parents' anxiety (Asbury et al., 2020; Ashworth et al., 2019). Similarly, for the TD group, education provides a great opportunity for socialization with their peers, which consecutively could lead to improved overall wellbeing. The current study also showed that individuals with SEND who have a pre-existing diagnosis of an anxiety disorder and who are aware about COVID-19 are in greater need of interventions related to anxiety.

Awareness of COVID-19 is also a key predictor for SEND when looking at anxiety levels. Such evidence might be explained by the complexity of the current events and that the associated rules of these events might be more difficult to be comprehended by individuals with SEND, as research has already shown (Aishworiya & Kang, 2020), leading to more stressful and anxious feelings which in the long-term could lead to higher anxiety.

Understanding parents / carers mental health is also important; especially since it predicts anxiety of individuals with SEND. The current study provides evidence that further investigation is needed about this association as it can help understand how parental high anxiety leads to higher anxiety in their children, but also subsequently how such high-quality of stress and anxiety impacts the parent-child as has been argued (Russell et al., 2020)

Finally, our data strongly suggest that both groups would benefit from schools and activity centres such the operation of these institutes during the current ongoing pandemic could be described as vital for now, but also for future lockdowns.

Acknowledgements

We would like to thank all the children, families and professionals who took part in the research as well as members of the chEERS lab who coordinated the study and Unidistance Suisse and the Swiss National Science Foundation (SNSF PP00P1_176722) for funding. Thanks also go to Susanna Mannik who helped with the editing of the manuscript.

References

- Aishworiya, R., & Kang, Y. Q. (2020). Including Children with Developmental Disabilities in the Equation During this COVID-19 Pandemic. *Journal of Autism and Developmental Disorders*. Advance online. <https://doi.org/10.1007/s10803-020-04670-6>
- Alexander, R., Ravi, A., Barclay, H., Sawhney, I., Chester, V., Malcolm, V., Brolly, K., Mukherji, K., Zia, A., Tharian, R., Howell, A., Lane, T., Cooper, V., & Langdon, P. E. (2020). Guidance for the Treatment and Management of COVID-19 Among People with Intellectual Disabilities. *Journal of Policy and Practice in Intellectual Disabilities*. Advance online. <https://doi.org/10.1111/jppi.12352>
- Asbury, K., Fox, L., Deniz, E., Code, A., & Toseeb, U. (2020). How is COVID-19 Affecting the Mental Health of Children with Special Educational Needs and Disabilities and Their Families? *Journal of Autism and Developmental Disorders*. Advance online. <https://doi.org/10.1007/s10803-020-04577-2>
- Ashworth, M., Palikara, O., & Herwegen, J. (2019). Comparing parental stress of children with neurodevelopmental disorders: The case of Williams syndrome, Down syndrome and autism spectrum disorders. *Journal of Applied Research in Intellectual Disabilities*, 32, 1–11. <https://doi.org/10.1111/jar.12594>
- Banerjee, D. D. (2020). The other side of COVID-19: Impact on obsessive compulsive disorder (OCD) and hoarding. *Psychiatry Research*, 288, 112966. <https://doi.org/10.1016/j.psychres.2020.112966>
- Becker, S. P., Langberg, J. M., Evans, S. W., Girio-Herrera, E., & Vaughn, A. J. (2015). Differentiating Anxiety and Depression in Relation to the Social Functioning of Young Adolescents with ADHD. *Journal of Clinical Child and Adolescent Psychology*, 44(6), 1015–1029. <https://doi.org/10.1080/15374416.2014.930689>

- Beidel, D. C., & Turner, S. (2005). *Child Anxiety Disorders: A Guide to Research and Treatment*. Taylor & Francis.
- Brooks, S. K., Webster, R. K., Smith, L. E., Woodland, L., Wessely, S., Greenberg, N., & Rubin, G. J. (2020). The psychological impact of quarantine and how to reduce it: Rapid review of the evidence. *The Lancet*, 395(10227), 912–920. [https://doi.org/10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8)
- Cartwright-Hatton, S., McNicol, K., & Doubleday, E. (2006). Anxiety in a neglected population: Prevalence of anxiety disorders in pre-adolescent children. *Clinical Psychology Review*, 26(7), 817–833. <https://doi.org/10.1016/j.cpr.2005.12.002>
- Chafouleas, S., Iovino, E., & Koriakin, T. (2020). Caregivers of Children with Developmental Disabilities: Exploring Perceptions of Health-Promoting Self-Care. *Journal of Developmental and Physical Disabilities*, 32, 1–21. <https://doi.org/10.1007/s10882-019-09724-x>
- Cooper, K., Smith, L. G. E., & Russell, A. (2017). Social identity, self-esteem, and mental health in autism. *European Journal of Social Psychology*, 47(7), 844–854. <https://doi.org/10.1002/ejsp.2297>
- Costello, E., Egger, H., & Angold, A. (2005). The Developmental Epidemiology of Anxiety Disorders: Phenomenology, Prevalence, and Comorbidity. *Child and Adolescent Psychiatric Clinics of North America*, 14, 631–648, vii. <https://doi.org/10.1016/j.chc.2005.06.003>
- Courtenay, K., & Perera, B. (2020). COVID-19 and people with intellectual disability: Impacts of a pandemic. *Irish Journal of Psychological Medicine*, 37(3), 231–236.
- Dhiman, S., Sahu, P. K., Reed, W. R., Ganesh, G. S., Goyal, R. K., & Jain, S. (2020). Impact of COVID-19 outbreak on mental health and perceived strain among caregivers tending children with special needs. *Research in Developmental Disabilities*, 107, 103790. <https://doi.org/10.1016/j.ridd.2020.103790>
- Dockrell, J. E., Ricketts, J., Palikara, O., Charman, T., & Lindsay, G. A. (2019). What drives educational support for children with developmental language disorder or autism spectrum disorder: Needs, or diagnostic category?. In *Frontiers in Education* (Vol. 4, p. 29). Frontiers. 11 January 2021, 11:49

- Dodd, H., Schniering, C., & Porter, M. (2009). Beyond Behaviour: Is Social Anxiety Low in Williams Syndrome? *Journal of Autism and Developmental Disorders*, 39, 1673–1681. <https://doi.org/10.1007/s10803-009-0806-4>
- Duan, L., Shao, X., Wang, Y., Huang, Y., Miao, J., Yang, X., & Zhu, G. (2020). An investigation of mental health status of children and adolescents in china during the outbreak of COVID-19. *Journal of Affective Disorders*, 275, 112–118. <https://doi.org/10.1016/j.jad.2020.06.029>
- Harrowell, I., Hollén, L., Lingam, R., & Emond, A. (2017). Mental health outcomes of developmental coordination disorder in late adolescence. *Developmental Medicine and Child Neurology*, 59(9), 973–979. <https://doi.org/10.1111/dmcn.13469>
- Iovannone, R., Dunlap, G., Huber, H., & Kincaid, D. (2003). Effective Educational Practices for Students With Autism Spectrum Disorders. *Focus on Autism and Other Developmental Disabilities*, 18(3), 150–165. <https://doi.org/10.1177/10883576030180030301>
- Jeste, S., Hyde, C., Distefano, C., Halladay, A., Ray, S., Porath, M., Wilson, R. B., & Thurm, A. (2020). Changes in access to educational and healthcare services for individuals with intellectual and developmental disabilities during COVID-19 restrictions. *Journal of Intellectual Disability Research*, 64(11), 825–833. <https://doi.org/10.1111/jir.12776>
- Jiao, W. Y., Wang, L. N., Liu, J., Fang, S. F., Jiao, F. Y., Pettoello-Mantovani, M., & Somekh, E. (2020). Behavioral and Emotional Disorders in Children during the COVID-19 Epidemic. *The Journal of Pediatrics*, 221, 264-266.e1. <https://doi.org/10.1016/j.jpeds.2020.03.013>
- Johns Hopkins School of Medicine. (2020). COVID-19 Dashboard by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University (JHU). <https://coronavirus.jhu.edu/map.html>. Accessed 11th January 2021.
- Kwong, A. S. F., Pearson, R. M., Adams, M. J., Northstone, K., Tilling, K., Smith, D., Fawns-Ritchie, C., Bould, H., Warne, N., Zammit, S., Gunnell, D., Moran, P., Micali, N., Reichenberg, A., Hickman, M., Rai, D., Haworth, S., Campbell, A., Altschul, D., ... Timpson, N. J. (2020). Mental health during the COVID-19 pandemic in two longitudinal UK population cohorts. *MedRxiv*, 2020.06.16.20133116. <https://doi.org/10.1101/2020.06.16.20133116>

- Leyfer, O. T., Folstein, S. E., Bacalman, S., Davis, N. O., Dinh, E., Morgan, J., Tager-Flusberg, H., & Lainhart, J. E. (2006). Comorbid psychiatric disorders in children with autism: Interview development and rates of disorders. *Journal of Autism and Developmental Disorders*, 36(7), 849–861. <https://doi.org/10.1007/s10803-006-0123-0>
- Lindsay, S., Proulx, M., Scott, H., & Thomson, N. (2014). Exploring teachers' strategies for including children with autism spectrum disorder in mainstream classrooms. *International Journal of Inclusive Education*, 18(2), 101–122. <https://doi.org/10.1080/13603116.2012.758320>
- Matthews, K., Coghill, D., & Rhodes, S. (2008). Neuropsychological functioning in depressed adolescent girls. *Journal of Affective Disorders*, 111(1), 113–118. <https://doi.org/10.1016/j.jad.2008.02.003>
- McLean, C.P., Asnaani, A., Litz, B.T., Hofmann, S.G. (2011). Gender differences in anxiety disorders: Prevalence, course of illness, comorbidity and burden of illness. *Journal of Psychiatric Research*, 45(8), 1027-1035. <https://doi.org/10.1016/j.jpsychires.2011.03.006>
- Mutluer, T., Doenyas, C., & Aslan Genc, H. (2020). Behavioral Implications of the Covid-19 Process for Autism Spectrum Disorder, and Individuals' Comprehension of and Reactions to the Pandemic Conditions. *Frontiers in Psychiatry*, 11. <https://doi.org/10.3389/fpsy.2020.561882>
- Neece, C. L., McIntyre, L. L., & Fenning, R. (2020). Examining the impact of COVID-19 in ethnically diverse families with young children with intellectual and developmental disabilities. *Journal of Intellectual Disability Research*. Advance online. <https://doi.org/10.1111/jir.12769>
- Neece, C. L., Green, S. A., & Baker, B. L. (2012). Parenting stress and child behavior problems: A transactional relationship across time. *American Journal on Intellectual and Developmental Disabilities*, 117(1), 48–66. <https://doi.org/10.1352/1944-7558-117.1.48>
- Nelson, J. M., & Harwood, H. (2011). Learning Disabilities and Anxiety: A Meta-Analysis. *Journal of Learning Disabilities*, 44(1), 3–17. <https://doi.org/10.1177/0022219409359939>

- O'Hagan, B., & Kingdom, S. (2020). *Experiences of children with special educational needs and disabilities and their families in the United Kingdom during the coronavirus pandemic*. Tizard Learning Disability Review. <https://doi.org/10.1108/TLDR-09-2020-0025>
- Orgilés, M., Morales, A., Delvecchio, E., Mazzeschi, C., & Espada, J. (2020). Immediate Psychological Effects of the COVID-19 Quarantine in Youth From Italy and Spain. *Frontiers in Psychology*, 11. <https://doi.org/10.3389/fpsyg.2020.579038>
- Platt, R., Williams, S. R., & Ginsburg, G. S. (2016). Stressful Life Events and Child Anxiety: Examining Parent and Child Mediators. *Child Psychiatry and Human Development*, 47(1), 23–34. <https://doi.org/10.1007/s10578-015-0540-4>
- Radford, J., Bosanquet, P., Webster, R., & Blatchford, P. (2015). Scaffolding learning for independence: Clarifying teacher and teaching assistant roles for children with special educational needs. *Learning and Instruction*, 36, 1–10. <https://doi.org/10.1016/j.learninstruc.2014.10.005>
- Rhodes, S. M., Park, J., Seth, S., & Coghill, D. R. (2012). A comprehensive investigation of memory impairment in attention deficit hyperactivity disorder and oppositional defiant disorder. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 53(2), 128–137. <https://doi.org/10.1111/j.1469-7610.2011.02436.x>
- Rix, J., Hall, K., Nind, M., Sheehy, K., & Wearmouth, J. (2009). What pedagogical approaches can effectively include children with special educational needs in mainstream classrooms? A systematic literature review. *Support for Learning*, 24(2), 86–94. <https://doi.org/10.1111/j.1467-9604.2009.01404.x>
- Rose, J., Willner, P., Cooper, V., Langdon, P. E., Murphy, G. H., & Kroese, B. S. (2020). The effect on and experience of families with a member who has Intellectual and Developmental Disabilities of the COVID-19 pandemic in the UK: Developing an investigation. *International Journal of Developmental Disabilities*, 0(0), 1–3. <https://doi.org/10.1080/20473869.2020.1764257>
- Russell, B. S., Hutchison, M., Tambling, R., Tomkunas, A. J., & Horton, A. L. (2020). Initial Challenges of Caregiving During COVID-19: Caregiver Burden, Mental Health, and the Parent–Child Relationship. *Child Psychiatry & Human Development*, 51(5), 671–682. <https://doi.org/10.1007/s10578-020-01037-x>

- Schalock, R. L. (Ed.). (1996b). *Quality of life: Vol. 1. Conceptualization and measurement*. Washington, DC: American Association on Mental Retardation.
- Schalock, R. L., & Verdugo, M. Á. (2002). *Handbook on quality of life for human service practitioners*. American Association on Mental Retardation. Washington, DC
- Sellke, T., Bayarri, M. and Berger, J. (2001) Calibration of p-Values for Testing Precise Null Hypotheses. *The American Statistician*, 55, 62-71.
<https://doi.org/10.1198/000313001300339950>
- Toseeb, U., Asbury, D. K., Code, A., Fox, L., & Deniz, E. (2020). Supporting Families with Children with Special Educational Needs and Disabilities During COVID-19. PsyArXiv. <https://doi.org/10.31234/osf.io/tm69k>
- Van Herwegen, J., Ashworth, A., & Palikara, O. (2018). Parental views on special educational needs provision: cross-syndrome comparisons in Williams Syndrome, Down Syndrome, and Autism Spectrum Disorders. *Research in Developmental Disability*, 80, 102-111.
- Van Herwegen, J., Ashworth, A., & Palikara, O. (2019). Views of professionals about the educational needs of children with neurodevelopmental disorders. *Research in Developmental Disability*, advance online. doi: 10.1016/j.ridd.2019.05.001.
- Van Herwegen, J., Dukes, D., & Samson, A. (2020). COVID19 Crisis Response Survey for families of Individuals with Special Needs. <https://doi.org/10.17605/OSF.IO/5NKQ9>
- Van Herwegen, J., Palikara, O., Riby, Deborah., Hanley, M., Rhodes, S., Giannadou, A., Samson, A., Dukes, D. (2020). The impact of COVID-19 on education and children's services. Evidence submitted to Select Committee for the UK Government July 2020. <https://committees.parliament.uk/writtenevidence/8883/pdf/>
- Wigham, S., Barton, S., Parr, J. R., & Rodgers, J. (2017). A Systematic Review of the Rates of Depression in Children and Adults With High-Functioning Autism Spectrum Disorder. *Journal of Mental Health Research in Intellectual Disabilities*, 10(4), 267–287. <https://doi.org/10.1080/19315864.2017.1299267>
- Willner, P., Rose, J., Kroese, B. S., Murphy, G. H., Langdon, P. E., Clifford, C., Hutchings, H., Watkins, A., Hiles, S., & Cooper, V. (2020). Effect of the COVID-19 pandemic on the mental health of carers of people with intellectual disabilities. *Journal of Applied Research in Intellectual Disabilities*, 33(6), 1523–1533.
<https://doi.org/10.1111/jar.12811>

Zhang, J., Shuai, L., Yu, H., Wang, Z., Qiu, M., Lu, L., Cao, X., Xia, W., Wang, Y., & Chen, R. (2020). Acute stress, behavioural symptoms and mood states among school-age children with attention-deficit/hyperactive disorder during the COVID-19 outbreak. *Asian Journal of Psychiatry*, 51, 102077. <https://doi.org/10.1016/j.ajp.2020.102077>

Supplementary materials

Table 9

Average score for the different worries for the SEND and TD groups for each of the time points (standard deviation)

		Time-point		Before		Start		Now	
		Group	SEND	TD	SEND	TD	SEND	TD	
Type of Concerns	Diagnosis								
	Concerns about COVID-19		1.62 (1.05)	1.44 (.87)	2.54 (1.36)	2.44 (1.22)	3.00 (1.43)	2.84 (1.20)	
	Concerns about others becoming ill		1.85 (1.17)	1.73 (1.12)	2.55 (1.38)	2.49 (1.25)	2.84 (1.45)	3.09 (1.36)	
	Health-related worries	Concerns about safety	1.72 (1.13)	1.56 (1.00)	2.50 (1.43)	2.53 (1.32)	2.87 (1.49)	3.03 (1.31)	
	Concerns about health	1.88 (1.15)	1.60 (.89)	2.26 (1.33)	2.06 (1.19)	2.52 (1.42)	2.28 (1.23)		
	Social related worries	Concerns about friends	2.00 (1.21)	1.99 (1.24)	2.72 (1.35)	3.03 (1.34)	3.21 (1.48)	3.53 (1.34)	
	Concerns about approach	1.75 (1.14)	1.70 (1.13)	2.28 (1.38)	2.34 (1.32)	2.63 (1.48)	2.83 (1.53)		

	Concerns about changes in routine	2.21 (1.36)	1.5 ⁸ (1.07)	2.97 (1.40)	2.15 (1.26)	3.15 (1.46)	2.44 (1.40)
Worries related to school closures	Getting bored	1.83 (1.15)	1.78 (1.23)	2.24 (1.34)	2.38 (1.38)	2.72 (1.50)	2.99 (1.50)
	Loss of Institution (including closure of school)	1.80 (1.25)	1.35 (.80)	2.41 (1.46)	2.00 (1.30)	2.76 (1.56)	2.27 (1.46)
	Concerns about family	1.51 (.92)	1.43 (.85)	1.67 (1.06)	1.60 (.92)	1.82 (1.23)	1.82 (1.12)
Family-related worries	Finance	1.19 (.61)	1.27 (.52)	1.23 (.70)	1.46 (.78)	1.27 (.74)	1.64 (1.04)
