Open with 🔻

Adjusted Odds Ratio (95% CI)			e-mediated natory diseases		
$\begin{array}{c} 50.66 \ (11.94, 214.93) \\ 9.09 \ (1.57, 52.53) \\ 7.14 \ (3.50, 14.57) \\ 6.96 \ (1.99, 24.36) \\ 5.70 \ (4.40, 7.39) \\ 5.44 \ (0.43, 68.09) \\ 5.37 \ (2.87, 10.05) \\ 3.83 \ (1.74, 8.44) \\ 3.60 \ (2.28, 5.67) \\ 3.47 \ (1.90, 6.33) \\ 3.39 \ (1.78, 6.42) \\ 2.61 \ (1.53, 4.46) \\ 2.40 \ (0.97, 5.97) \\ 2.31 \ (1.57, 3.40) \\ 2.20 \ (1.49, 3.27) \\ 1.62 \ (0.89, 2.95) \\ 1.36 \ (0.12, 15.50) \\ 1.34 \ (1.10, 1.63) \\ 1.29 \ (0.64, 2.58) \\ 1.20 \ (0.94, 1.54) \\ 1.11 \ (0.60, 2.06) \\ 1.04 \ (0.54, 2.00) \end{array}$		Granulor Behcet's Sjogren' - Reiter's Rheuma Dermato Vasculitis Giant cel Crohn's Systemic Ankylosi Sarcoido Lymphor Polymya Ulcerativ Coeliac o SWEET Psoriasis Multiple Gout Thyroid o Vitiligo	l arteritis (GCA) disease clupus erythematosus ng spondylitis sis na Igia rheumatica e colitis disease syndrome		
4.12 (0.50, 30.33) 3.76 (0.45, 31.45) 2.56 (0.37, 17.62) 1.61 (1.34, 1.94) 1.51 (1.00, 2.27) 1.50 (1.09, 2.08) 1.35 (0.83, 2.19) 0.63 (0.14, 2.88)		Lyme dis Toxoplas Aspergill Herpetic Measles	mosis us infection Barr virus —		
2.71 (1.15, 6.38) 17.30 (13.29, 22.60)	<b>→ →</b>	- CLINICA Optic ne Uveitis	L PHENOTYPES uritis		+
.1	1	100	ا 1.	1	10 100
Decreased Risk	Increased	Risk	Decreased Ri	sk	Increased Risk

Page 1 / 1 —

Ð

+

	<b>@</b> +	Û	→	•
Adjusted Hazard Ratio (95% CI)				

Ρ

96.36 (12.99, 715.00)
 17.45 (1.94, 156.74)
 8.53 (3.35, 21.67)
 4.22 (3.32, 5.39)
 5.38 (0.33, 88.57)
 6.68 (3.22, 13.86)
 2.36 (1.03, 5.44)
 4.16 (2.90, 7.58)
 8.49 (3.35, 21.74)
 3.44 (1.21, 9.76)
 3.09 (1.15, 8.27)
 1.11 (0.70, 1.75)
 1.77 (0.99, 3.16)
 1.07 (0.43, 2.69)
 1.22 (0.92, 1.60)
 1.12 (0.41, 3.08)
 0.96 (0.73, 1.24)

 1.16 (0.45, 2.94)

 1.33 (0.55, 3.22)

 0.69 (0.08, 6.35)

 8.41 (1.43, 49.39)

 1.72 (0.17, 17.40)

 1.53 (1.16, 2.02)

 0.72 (0.08, 6.68)

 3.37 (0.82, 13.78)

 11.11 (1.05, 117.16)

20.08 (14.02, 28.76)

00 300 sk