

1

2 **Main Manuscript for**

3 LXR directly regulates glycosphingolipid synthesis and affects human
4 CD4⁺ T cell function

5

6 Kirsty E Waddington^{a,b,1}, George A Robinson^a, Beatriz Rubio-Cuesta^{b,2}, Eden Chrifi-Alaoui^a, Sara
7 Andreone^{a,3}, Kok-Siong Poon^{b,4}, Iveta Ivanova^c, Lucia Martin-Gutierrez^a, Dylan M Owen^{c,5},
8 Elizabeth C Jury^{a,#,*} and Inés Pineda-Torra^{b,#,*}.

9

10 ^a Centre for Rheumatology or ^b Centre for Cardiometabolic and Vascular Science, University
11 College London, United Kingdom. ^c Department of Physics and Randall Division of Cell and
12 Molecular Biophysics, King's College London, United Kingdom.

13 Present addresses: ¹GammaDelta Therapeutics Limited, London, UK, ²Translational Oncology
14 Laboratory, Research Institute i+12 and Spanish National Cancer Research Centre (CNIO) and
15 Universidad Autonoma de Madrid, Madrid, Spain, ³Department of Oncology and Molecular
16 Medicine, Istituto Superiore di Sanità, 00161 Rome, Italy, ⁴Department of Pathology, Yong Loo
17 Lin School of Medicine, National University Singapore and ⁵University of Birmingham, United
18 Kingdom.

19

20 # Shared senior authorship

21 * Inés Pineda-Torra and Elizabeth C. Jury

22 **Email:** i.torra@ucl.ac.uk and e.jury@ucl.ac.uk

23

24 **Classification**

25 Biological Sciences/Immunology and Inflammation

26 **Keywords**

27 liver X receptor, CD4⁺ T cell, lipid metabolism, cholesterol, glycosphingolipids

28

29 **Author Contributions**

30 KEW performed most experiments and data analysis, and prepared the figures. KEW, GAR, K-
31 SP, LMG, EC-A and SA performed flow cytometry experiments and qPCR analysis. BR-C
32 performed western blotting experiments. DMO and II acquired microscopy images for di-4-
33 ANEPPDHQ experiments and provided guidance for image analysis. KEW, IPT and ECJ
34 designed experiments, interpreted the data, and prepared the manuscript. IPT and ECJ
35 conceived the study, secured the funding and supervised all aspects of the work.

36

37 **Competing Interests**

38 No competing interest to declare

39

40 **This PDF file includes:**

41 Main Text
42 Figures 1 to 6
43

44 **Abstract**

45 The liver X receptor (LXR) is a key transcriptional regulator of cholesterol, fatty acid, and
46 phospholipid metabolism. Dynamic remodeling of immunometabolic pathways, including lipid
47 metabolism, is a crucial step in T cell activation. Here we explored the role of LXR-regulated
48 metabolic processes in primary human CD4⁺ T cells, and their role in controlling plasma
49 membrane lipids (glycosphingolipids and cholesterol) which strongly influence T cell immune
50 signaling and function. Crucially, we identified the glycosphingolipid biosynthesis enzyme
51 glucosylceramide synthase (UGCG) as a direct transcriptional LXR target. LXR activation by
52 agonist GW3965 or endogenous oxysterol ligands significantly altered the
53 glycosphingolipid:cholesterol balance in the plasma membrane by increasing glycosphingolipid
54 levels and reducing cholesterol. Consequently, LXR activation lowered plasma membrane lipid
55 order (stability), and an LXR antagonist could block this effect. LXR stimulation also reduced lipid
56 order at the immune synapse and accelerated activation of proximal T cell signaling molecules.
57 Ultimately, LXR activation dampened pro-inflammatory T cell function. Finally, compared to
58 responder T cells, regulatory T cells had a distinct pattern of LXR-target gene expression
59 corresponding to reduced lipid order. This suggests LXR-driven lipid metabolism could contribute
60 to functional specialization of these T cell subsets. Overall, we report a novel mode of action for
61 LXR in T cells involving the regulation of glycosphingolipid and cholesterol metabolism, and
62 demonstrate its relevance in modulating T cell function.

63

64 **Significance Statement**

65 This work shows for the first time that LXR regulates glycosphingolipid biosynthesis expression in
66 primary human T cells, thereby influencing T cell plasma membrane lipid composition,
67 subsequent immune synapse formation, T cell receptor-mediated signaling and function.
68 Furthermore, we show LXR actions are differentially regulated in functional T cell subsets,
69 supporting an important role for lipid metabolism in human T cell homeostasis. This new mode of
70 action for LXR could be of therapeutic relevance to disorders characterized by defects in T cell

71 signaling and metabolism, including autoimmune and neurodegenerative diseases,
72 cardiovascular disease and cancer.

73

74 **Main Text**

75

76 **Introduction**

77

78 CD4⁺ T cells (also known as T helper cells) shape the immune response by releasing cytokines
79 with both pro-inflammatory and immunomodulatory effects. A number of factors govern the
80 precise balance of pro- and anti-inflammatory mediators produced, including antigenic
81 stimulation, cell-cell signaling and micro-environmental cues. The T cell plasma membrane
82 facilitates these processes, providing a flexible interface between the cell and its
83 microenvironment, where membrane receptors integrate internal and external signals to generate
84 functional outcomes. Lipids are a key component of the plasma membrane and contribute to its
85 biophysical properties and protein receptor compartmentalization. Cholesterol and
86 glycosphingolipids are particularly enriched, forming signaling platforms known as lipid rafts which
87 play a critical role in T cell antigen receptor (TCR) signaling and T cell function(1). Cholesterol
88 maintains lipid raft structure, inhibits spontaneous TCR activation and promotes TCR clustering(2,
89 3). In addition, cholesterol has been shown to regulate T cell proliferation(4, 5), differentiation and
90 cytokine production (6). Similarly, glycosphingolipids influence TCR-mediated signaling,
91 responsiveness to cytokine stimulation and T_H17 cell differentiation(7, 8). Plasma membrane
92 cholesterol and glycosphingolipid levels influence lipid order, a measure of how tightly packed
93 lipids are in the membrane (9); notably increased cholesterol is associated with higher lipid order
94 (9-11). Variations in lipid order can influence the interaction of membrane receptors and
95 determine the strength of cell signaling (12). In particular, changes in lipid order at the T cell
96 immune synapse can alter the strength and nature of signaling events and impact T cell function
97 (9, 10, 13). Importantly, abnormal T cell plasma membrane lipids have been linked to pathogenic
98 T cell function and are attractive targets for immunotherapy in autoimmunity, viral infection and
99 cancer(14-19).

100

101 Our previous work linked pathogenic elevation of CD4⁺ T cell glycosphingolipid expression in
102 systemic lupus erythematosus (SLE) to liver X receptor (LXR) expression (7). LXR α (*NR1H3*) and
103 LXR β (*NR1H2*) are transcription factors activated by oxidized derivatives of cholesterol
104 (oxysterols)(20) and intermediates of cholesterol biosynthesis(20) to regulate gene expression.
105 The majority of LXR target genes are involved in the metabolism of lipid metabolic processes,
106 including cholesterol efflux and uptake, fatty acid biosynthesis, and phospholipid remodeling(20).
107 However, it is not known whether LXR regulates glycosphingolipid metabolism or T cell lipid rafts.
108 This prompted us to further explore the relationship between LXRs, glycosphingolipid metabolism
109 and plasma membrane lipid composition.

110

111 Here we demonstrate a new role for LXR in human CD4⁺ T cells that involves modulation of the
112 human T cell transcriptome and lipidome. We show how LXR activation modulates
113 glycosphingolipid and cholesterol homeostasis and define a new mechanism for LXR-mediated
114 effects on T cell function via regulation of plasma membrane lipid composition. Finally, we show
115 that regulatory T cells (Tregs) have a distinct plasma membrane lipid profile that corresponds to
116 differential expression of LXR target genes. We propose that regulation of membrane lipids by LXR
117 could contribute to the specialized regulatory functions of this T cell subset.

118

119

120 **Results**

121

122 ***LXR transcriptionally regulates lipid metabolic pathways in human CD4⁺ T cells***

123 To define the transcriptional effects of LXR activation in human CD4⁺ T cells, primary cells were
124 exposed to the specific LXR agonist GW3965 (GW) (21). Sixty-five LXR-responsive genes were
125 identified (Fig. 1a-b, SI Appendix-Dataset S1), and GW-treated samples were clearly
126 distinguishable from their controls by principal component analysis (PCA) (SI Appendix-Fig. S1a).
127 The majority of differentially expressed genes (DEGs) were upregulated (53 out of 65), a subset of

128 which demonstrated a very strong ligand response. These included well-characterised LXR target
129 genes (*ABCG1*, *ABCA1*, *APOC1*, *SCD* and *SREBF1* (22)) and the recently identified
130 oligodendrocyte maturation-associated long intervening non-coding RNA (*OLMALINC*) (23) (Fig.
131 1c, SI Appendix-Fig. S1b). Other previously identified LXR target genes had a more modest
132 upregulation (<5-fold) (SI Appendix-Fig S1c). The significantly enriched pathways were
133 hierarchically clustered into functionally related groups (Fig. 1d). Strikingly, all 15 clusters enriched
134 for LXR-upregulated genes were related to metabolism, the most significant of which was
135 'cholesterol metabolic process'. Only 12 genes were significantly downregulated by GW (Fig. 1a-
136 b, SI Appendix-Fig. S1b), and these were most strongly associated with the 'regulation of
137 inflammatory responses' (Fig. 1e).

138
139 LXRs can act in a subtype specific manner, and the relative expression of LXR α and β differs
140 between monocytes/macrophages and T cells(4, 24) (SI Appendix-Fig.S1d-e). Another striking
141 difference is that in monocytes/macrophages LXR α regulates its own expression via an
142 autoregulatory loop(25) which does not occur in T cells (SI Appendix-Fig. S1f). These differences
143 likely lead to cell-type specific responses to LXR activation. To identify potential T cell specific LXR
144 targets we cross-referenced our list of DEGs with two publicly available RNA-sequencing datasets
145 from murine macrophages (mM ϕ) treated with GW(25). Of the DEGs identified in T cells, 52% were
146 similarly regulated in mM ϕ , and remarkably, 29% were uniquely regulated in the T cell dataset (Fig.
147 1f). Some of these genes are known to be differentially regulated between mice and
148 humans/primates but, to our knowledge, a subset have not previously been associated with LXR
149 activation (*BRWD3*, *CHD2*, *MKNK2*, *SLC29A2*, *TDRD6*, *TKT* and *UGCG*) (SI Appendix-Table S1).
150 Overall, genes involved in lipid metabolic pathways were upregulated in both cell types, but there
151 were no shared pathways amongst the downregulated genes, which tended to be involved in the
152 regulation of immunity and inflammation (SI Appendix-Fig. S1g-h). This supports that the
153 immunomodulatory effects of LXR activation vary between cell-types and species(26).

154
155 Thus, we have identified genes responsive to LXR activation in human CD4⁺ T cells, most markedly
156 the upregulation of genes involved in lipid metabolic processes, and highlighted a subset of genes
157 that may represent human or T cell specific targets.

158 ***LXR controls transcriptional regulation of glycosphingolipid biosynthesis enzyme UGCG***

159 Since LXR activation predominantly regulated genes involved in lipid metabolism, the impact on T
160 cell lipid content was assessed using shotgun lipidomics. Although total intracellular lipid levels
161 were not affected by LXR activation (Fig. 2a), 15% of the detected lipid subspecies were
162 significantly regulated (54 out of 366, Fig. 2b). Notably, a large proportion of triacylglycerols (TAG)
163 and hexosylceramides (HexCer) were induced by LXR activation, and overall quantities of TAG
164 and HexCer were elevated (Fig. 2c-e, SI Appendix-Table S5).

165
166
167 LXR regulated many enzymes involved in fatty acid metabolic processes including synthesis,
168 desaturation and elongation (Fig 1d, SI Appendix-Dataset S1). There were no changes in total
169 levels of saturated, monounsaturated and polyunsaturated lipids. However, amongst PUFAs there
170 was an increase in degree of unsaturation, which is associated with membrane disorder(27, 28) (SI
171 Appendix-Fig. S2a). Further examination at the lipid class level revealed significant increases in
172 saturated and monounsaturated lipid species, HexCers, and TAG species with more than 4 double
173 bonds (Fig. 2c and SI Appendix-Fig.S2b).

174
175 This is the first report linking LXR activation to HexCer. We observed GW also reduced levels of
176 several ceramides (Fig. 2b-c), suggesting an accelerated conversion of ceramide to HexCer - a
177 reaction catalysed by glycosphingolipid biosynthesis enzymes UDP-glucosylceramide synthase
178 (UGCG) or UDP-glycosyltransferase 8 (UGT8) (Fig. 2f-g). In support of this, UGCG mRNA

179 expression was upregulated by LXR activation (Fig. 2h), whereas UGT8 was absent in CD4⁺ T cells
180 (SI Appendix-Fig. S2c).
181 UGCG upregulation was further amplified by co-stimulation of LXR and its heterodimeric partner
182 the retinoid X receptor (RXR) (SI Appendix-Fig. S2d), as has been reported for other LXR target
183 genes (29, 30). GW treatment also enhanced UGCG expression in other immune cell types,
184 including peripheral blood mononuclear cells (PBMCs), CD14⁺ monocytes and CD19⁺ B cells (SI
185 Appendix-Fig. S2e). However, in monocyte-derived macrophages and THP-1 macrophages,
186 UGCG was only modestly increased (<1.5-fold change, SI Appendix-Fig. S2e-f). This may explain
187 why UGCG has not been identified as an LXR target gene in previous RNA-seq and ChIP-seq
188 experiments using macrophages (31, 32), in which most of LXR biology has been reported to date.
189 The increase in UGCG expression was not a GW-specific effect, as UGCG mRNA was also
190 upregulated in response to stimulation with the endogenous LXR activators 24S, 25-
191 epoxycholesterol (24S,25-EC) and 24S-hydroxycholesterol (24S-OHC), albeit with an altered
192 kinetic (SI Appendix-Fig. S2g).

193

194 To determine whether LXR regulates UGCG expression by directly binding to the UGCG locus, we
195 screened for potential LXR response element (LXRE) sequences *in silico*. A putative DR4
196 sequence was identified upstream of the UGCG gene that coincided with an LXR-binding peak in
197 HT29 cells treated with GW(33) (SI Appendix-Fig. S2i). ChIP-qPCR experiments demonstrated
198 enrichment in LXR occupancy at this site, which increased with ligand activation (SI Appendix-Fig.
199 2i). The observed LXR occupancy at the UGCG gene followed a similar pattern to that of a reported
200 LXRE within SMPDL3A(34)(SI Appendix-Fig. 2i & S2h-j). Moreover, acetylation of histone H3K27
201 was enriched at this region compared to the IgG and a negative control sequence, suggesting this
202 site falls in an active transcriptional enhancer (SI Appendix-Fig. S2i-j).

203

204 **LXR regulates the T cell plasma membrane lipid raft profile**

205 UGCG is the rate-limiting enzyme for the biosynthesis of glycosphingolipids, important components
206 of plasma membrane lipid rafts. Indeed, LXR activation consistently upregulated T cell
207 glycosphingolipid expression measured using cholera-toxin B (CTB)(Fig. 3a), a well-established
208 surrogate glycosphingolipid marker(7). Specific pharmacological inhibition of UGCG activity
209 blocked the induction of glycosphingolipids by GW, suggesting this was UGCG-dependent (SI
210 Appendix-Fig. S2k). The increase in glycosphingolipid levels was accompanied by significant
211 downregulation of membrane cholesterol (Fig. 3b), likely due to the strong induction of cholesterol
212 efflux transporters ABCA1 and ABCG1 (Fig. 1c, SI Appendix-S2m). As expected, UGCG inhibition
213 had no effect on the reduction of cholesterol or lipid order (SI Appendix-Fig. S2k). Overall, LXR
214 activation significantly increased the ratio of glycosphingolipids to cholesterol (Fig. 3c).

215

216 The relative abundance and arrangement of lipids in the plasma membrane dictates its 'lipid order',
217 an important determinant of signalling protein localisation during immune synapse formation(10).
218 Cholesterol levels positively correlate with T-cell plasma membrane lipid order, whereas
219 glycosphingolipid levels have a negative correlation(9). LXR lowers cholesterol and raises
220 glycosphingolipids, resulting in the significant reduction of membrane lipid order by GW/ oxysterol
221 activated LXR (Fig. 3d-e, SI Appendix-S2k-l). The specific LXR antagonist GSK233 was able to
222 block the reduction of lipid order, glycosphingolipid and cholesterol levels by GW (Fig. 3e, SI
223 Appendix-S2l). Oxysterols, also activated LXR-target gene expression and reduced lipid order
224 while GSK233 only partially reversed the effect of 24S-OHC, in line with the known LXR-
225 independent actions of oxysterols (SI Appendix-Fig. S2m-o).

226

227 Furthermore, LXR target genes were differentially expressed in T cells sorted based on their
228 (high/low) plasma membrane lipid order. T cells with low membrane lipid order (low cholesterol,
229 high glycosphingolipids) had elevated expression of *ABCA1*, *ABCG1* and *UGCG* compared to T
230 cells with high membrane lipid order (high cholesterol, low glycosphingolipids) (Fig. 3f). This
231 suggests LXR ligand induced cholesterol efflux (*ABCA1/G1*) and glycosphingolipid biosynthesis
232 (*UGCG*) contribute to the generation of low membrane lipid order. In contrast, there was no

233 difference in the expression of genes controlling fatty acid synthesis (*SREBP1c*, *FASN*) in cells with
234 different membrane order (Fig. 3f).

235

236 Overall, these data suggest that LXR transcriptionally upregulates *de novo* glycosphingolipid
237 synthesis in human T cells, thereby contributing to the remodelling of plasma membrane lipid
238 composition in response to LXR activation.

239

240 ***LXR activity modulates lipid metabolism and effector function of activated T cells***

241 Next, we explored the effect of LXR on primary human T cell activation. Over 3000 genes were
242 significantly regulated by TCR activation, although most of these were regulated irrespective of
243 LXR activation with GW (Fig. 4a). Interestingly, LXR β expression was slightly increased by TCR
244 stimulation, while LXR α expression remained low (Fig. S3a). Overall, 113 genes were regulated by
245 the presence or absence of GW in activated T cells (Fig. 4b, SI Appendix-S3b & Dataset S2).
246 TCR/LXR co-stimulation upregulated genes involved in lipid metabolic processes, and
247 downregulated genes associated with immune system processes including chemokine production
248 and chemotaxis (SI Appendix-Fig. S3c). When these genes were clustered based on their
249 expression in both activated and resting cells, four major patterns of gene expression were
250 identified (Fig. 4c, SI Appendix-Table S2). Many genes upregulated by GW in resting cells were
251 upregulated to an equal or greater extent in GW/TCR co-activated cells (clusters A and C, Fig. 4c).
252 These clusters were enriched for genes involved in lipid and cholesterol metabolic processes,
253 including canonical LXR target genes *ABCA1* and *SREBF1* and the newly identified LXR-target
254 gene *UGCG*. This corresponded with changes in global plasma membrane lipid composition,
255 namely increased glycosphingolipids but reduced cholesterol in response to LXR/TCR co-
256 stimulation compared with TCR stimulation alone (Fig. 4d-f). Therefore, LXR activation continues
257 to modulate plasma membrane composition throughout the course of T cell activation.

258

259 In contrast, GW/TCR co-stimulation reduced the induction of a subset of genes involved in
260 leukocyte activation (cluster B, Fig. 4c). Interestingly, other genes were only activated (cluster C)
261 or repressed (cluster D) by LXR activation in the context of TCR stimulation (Fig. 4c). Therefore,
262 bidirectional crosstalk between LXR and TCR stimulation modulates transcription in a gene-specific
263 manner. Likely, more subtle differences did not reach statistical significance due to the
264 heterogeneous response to stimulation between the healthy donors (SI Appendix-Fig. S3d).

265

266 In murine T cells, TCR stimulation was previously reported to repress LXR transcriptional activity,
267 by reducing the availability of endogenous LXR ligands due to their modification by the
268 sulfotransferase *SULT2B1*(4). However, in the present study we observed very low levels of
269 *SULT2B1* in human CD4⁺ T cells (<11 gene counts), and *SULT2B1* was not regulated by TCR
270 activation (Fig. 4g). We considered that oxysterol levels could be controlled by an alternative
271 mechanism, for example increased efflux or metabolism. Indeed, TCR activation downregulated
272 the expression of oxysterol-binding proteins and oxysterol biosynthesis enzyme *CYP27A1*, and
273 upregulated oxysterol metabolising enzyme *CYP1B1* (Fig. 4g). Therefore concentrations of
274 endogenous LXR ligands during human T cell activation are also tightly regulated, but likely through
275 a different mechanism.

276

277 LXR and T cell co-activation had significant functional consequences including increased
278 production of interleukin (IL)-2 and IL-4, reduced IL-17A release compared to non-LXR-treated
279 controls (Fig. 4h-i, SI Appendix-Fig. S4a-b). No changes in T cell interferon- γ , tumour necrosis
280 factor- α or IL-10 production were detected (SI Appendix-Fig. S4b). Although LXR has been
281 reported to regulate the transcription of certain cytokines (6, 35), this was not observed here (SI
282 Appendix-Dataset S2). Furthermore, the expression of transcription factors which drive Th1 (Tbet),
283 Th2 (GATA3), Treg (Foxp3) and Th17 (ROR γ) polarisation were also unaffected by LXR activation
284 (SI Appendix-Fig. S4c). Proliferation was inhibited by GW-treatment (Fig. 4j, SI Appendix-S4d) and
285 importantly, addition of the UGCG inhibitor NB-DNJ countered this effect by increasing proliferation
286 and partially blocking IL-2 and IL-4 production (SI Appendix-Fig. S4e,f). Considering the preferential

287 regulation of lipid metabolism genes (SI Appendix-Fig. S3c) and observed changes in plasma
288 membrane lipid levels (Fig. 4d-f), we instead hypothesised that the effects of LXR activation on T
289 cell function could be mediated, at least in part, by an altered lipid landscape.

290

291 ***LXR-driven modification of plasma membrane lipid profile alters TCR signalling***

292 T cell activation is initiated by TCR-proximal signalling at the immune synapse, leading to
293 proliferation and cytokine production. We previously demonstrated that, compared to cells with
294 highly ordered plasma membranes, T cells with lower membrane lipid order have reduced synapse
295 area, transient synapse formation and a Th1 cytokine skew(9). These functional outcomes are
296 influenced by the localisation of TCR-signalling proteins within lipid microdomains at the immune
297 synapse(36). To examine the effect of LXR stimulation on the kinetics of lipid reorganisation during
298 the early stages of T cell activation we used di-4-ANEPPDHQ staining and TIRF microscopy to
299 assess the interaction between CD4⁺ T cells and antibody-coated glass coverslips (mimicking the
300 'immune synapse') (Fig. 5a, SI Appendix-Movies S1-2). T cells pre-treated with GW had a
301 significantly lower membrane order (generalised polarisation (GP) ratio) at the cell/coverslip
302 interface for up to 20 minutes post-activation (Fig. 5b, SI Appendix-Movies S1-2). Synapse area
303 was unaffected (SI Appendix-Fig. S4g), however, the pattern and distribution of lipid order was
304 disrupted in GW-treated T cells compared to controls (SI Appendix-Fig. S4h). This was
305 accompanied by increased levels of global tyrosine phosphorylation (Fig. 5c), increased
306 accumulation of Lck receptor tyrosine kinase at the synapse (Fig. 5c) and a preference for Lck to
307 accumulate at the synapse periphery (Fig. 5d), an area typically associated with active
308 signalling(37). Specifically, GW treatment increased phosphorylation of important proximal T cell
309 signalling molecules CD3 and the adaptor molecule linker for activation of T cells (LAT), but not
310 extracellular signal related kinase (Erk) or phospholipase (PL) C γ 1 (SI Appendix-Fig. S4i).

311 Taken together, these results suggest that plasticity in T cell function could be driven, at least in
312 part, by altered plasma membrane lipid composition controlled by LXR activation.

313

314 ***Functional T cell subsets differ in their expression of LXR-regulated genes and lipids***

315 T cells with high and low membrane lipid order are functionally distinct (9). Compared to responder
316 T cells (Tresp), regulatory T cells (Treg) (Fig. 6a) had lower membrane order increased
317 glycosphingolipid levels and reduced membrane cholesterol (Fig. 6b-d). We hypothesised that the
318 LXR pathway could contribute to these differences. LXR α mRNA expression was significantly lower
319 in Tregs, although LXR β , which is the predominant form in T cells (SI Appendix-Fig. S1d-f), tended
320 towards higher expression (p=0.06) (Fig. 6e). Corresponding to the plasma membrane lipid
321 phenotype, Treg expression of the cholesterol transporter *ABCG1* and glycosphingolipid enzyme
322 *UGCG* were increased compared to Tresp, whereas other LXR target genes were not differentially
323 expressed (*ABCA1*, *IDOL*, *SREBF1*, *FASN*) (Fig. 6e).

324

325 Interestingly, Tregs had a more variable response to LXR stimulation than Tresp in terms of
326 reduction of membrane lipid order and induction of glycosphingolipids, although downregulation of
327 cholesterol was consistently similar (Fig. 6f-h). Mirroring the regulation of glycosphingolipids and
328 cholesterol, cholesterol metabolism genes (*ABCA1*, *ABCG1*, *IDOL*) were similarly induced in both
329 subsets whereas *UGCG* mRNA was significantly upregulated in Tresp but not Treg (Fig. 6g-h).
330 Fatty acid synthesis enzymes had a similar magnitude of regulation (4-fold vs 6-fold), although
331 *FASN* levels were much higher in GW-treated Treg than Tresp (Fig. 6i).

332

333 These results demonstrate that Treg and Tresp have distinct plasma membrane lipid profiles and
334 differences in LXR ligand responses. This suggests that variation in LXR activity could influence
335 the functional specialisation of T cell subsets.

336

337

338

339

340 **Discussion**

341

342 CD4⁺ T cells provide essential protection against infection and cancer, but dysregulated T cell
343 responses contribute to the pathogenesis of many diseases. LXRs are an attractive therapeutic
344 target in many immunometabolic diseases involving T cells (38, 39). However, the actions of LXR
345 in lymphocytes have not yet been fully investigated, particularly in human cells. This is important
346 since a number of differences in LXR biology have been reported between human and rodent
347 models, including the aforementioned species-specific regulation of certain genes(23, 25, 34, 40).
348 Furthermore, in stark contrast to the anti-inflammatory effects of LXR activation in murine
349 macrophages (41-43), LXR has been shown to potentiate pro-inflammatory responses in human
350 monocytes (40, 44, 45).

351

352 Here, we have comprehensively assessed the action of LXR in human CD4⁺ T cells combining
353 transcriptomic and lipidomic analyses with cell biology approaches to study the regulation of lipid
354 metabolism and T cell function. Our findings revealed a novel regulation of glycosphingolipid
355 biosynthesis by LXR in these cells, which may be replicated in other immune cell types. The
356 combined effect of LXR activation on glycosphingolipid and cholesterol levels contributed to an
357 overall reduction in plasma membrane lipid order, which modulated immune synapse formation
358 and proximal T cell signalling in the context of TCR activation.

359

360 Whilst this work was ongoing, LXR was shown to contribute to T cell development in animal
361 models. T cell specific deletion of LXR resulted in peripheral lymphopenia, thought to be caused
362 by accumulation of plasma membrane cholesterol, heightened apoptotic signalling, and
363 subsequent enhanced negative selection(46). This supports our findings that regulation of plasma
364 membrane lipids by LXR is important for T cell function. Additionally, recent work in murine
365 models highlighted LXR β indispensable role in murine Tregs(47). LXR activation was also shown
366 to exert anti-tumour effects by reducing the Treg content of the murine tumour
367 microenvironment(48). While this work stresses the importance of LXR in T cell biology, the
368 impact of LXR on plasma membrane metabolism was not examined.

369

370 There is extensive evidence in the literature that UGCG plays an important role in T-cell immune
371 synapse formation *in vitro* and *in vivo*. Similar to other studies(49), we did not use siRNA-based
372 methods which could adversely influence membrane integrity to assess the complex changes
373 imparted by LXR on membrane lipids and order. Rather, we used pharmacological inhibitors of
374 LXR (GSK2033) and UGCG (NB-DNJ). Inhibition of UGCG has been shown to attenuate proximal
375 TCR signalling in Jurkat T cells, to reduce the production of IL-2 and IFN γ and to inhibit
376 proliferation(8, 50). Our own work also showed that inhibition of UGCG normalised T-cell
377 signalling and function in primary human T cells from SLE patients *in vitro*(7). We have also
378 shown that in human T cells, increased glycosphingolipids are associated with increased
379 accumulation of protein tyrosine phosphatase CD45 (which regulates Lck activity) within lipid rafts
380 and increased Lck phosphorylation at the immune synapse(18). Thus, changes in lipid order and
381 lipid profile could reflect an initial acceleration in signalling (seen as increased tyrosine
382 phosphorylation), altered interaction between regulatory and inhibitory molecules and altered
383 downstream signalling events. This may not result in increased overall activation, but result in
384 changes in certain cytokine levels as we and others have described previously(36, 51-57).
385 Indeed, in T cells from SLE patients, inhibition of UGCG activity increased phosphorylation of
386 TCRzeta and Erk, yet dampened proliferation and pro-inflammatory cytokine production(7).
387 Interestingly, Guy et al. (58) also demonstrated that cytokine production and proliferation can be
388 uncoupled depending on the number of phosphorylated sites on the TCR subunits, whereby weak
389 signals are sufficient to maintain cytokine production but fail to induce proliferation – similar to the
390 phenotype observed here. Furthermore, changes in sphingolipid content at the immune synapse,
391 specifically a decline in glucosylceramide (the product of UGCG), have been linked to T-cell
392 dysfunction in aged mice (59). Finally, reduced expression of the glycosphingolipid GM1 in
393 effector T cells was associated with resistance to Treg suppression(60). Overall, these studies

394 demonstrate that perturbation of UGCG activity and glycosphingolipid levels have been linked
395 with abnormal TCR signalling at the immune synapse, resulting in altered effector functions.
396 However, LXR had not previously been linked to glycosphingolipid metabolism.

397

398 It is important to note that changes in membrane lipid order, that could at first glance appear
399 modest especially when compared to changes in gene expression, can nonetheless have
400 important consequences in T cell function(9, 12, 61, 62). Changes in plasma membrane lipid
401 order, measured using phase sensitive probes, can affect T cell responses to TCR stimulation(9,
402 12). Specifically, high order cells form a more stable immune synapse, resulting in a robust
403 proliferative response and Th2 cytokine skew. In contrast, cells with lower order proliferate less
404 and produce IFN γ (Th1)(9). Furthermore, pharmacologically reducing membrane order with an
405 oxysterol is sufficient to alter the immune synapse between T cells and antigen presenting cells
406 and subsequent T cell proliferation and cytokine production(9, 10).

407 The discovery that LXR activation upregulates UGCG expression in primary human immune cells
408 provides a novel mode of action for LXR in the immune system. The magnitude of transcriptional
409 activation by LXR, as for other nuclear receptors, depends on several factors, including chromatin
410 architecture and epigenomic landscape at the specific gene that will determine cofactor recruitment
411 and corepressor release, or whether other signal dependent transcription factors are present at the
412 binding site(63). We observed that H3K27 acetylation at the UGCG site does not change in
413 response to LXR ligand activation, similar to other LXR target genes (SMPDL3A is shown). This is
414 not unusual for LXR regulation of gene expression (64). Changes in H3K27ac could be dynamic
415 and altered with kinetics different to those of LXR binding. Additionally, other chromatin acetylation
416 marks associated with transcriptional activation linked to gene activation in human CD4+ T cells or
417 in the regulation of lipid metabolism could be relevant (65, 66). Future investigations will aim to
418 characterise currently lacking global profiles of activation marks in these cells in response to LXR
419 agonist and lipid changes. Furthermore, gene regulation may be mediated by the binding of
420 additional signal-dependent transcription factors to adjacent sites(67). Finally, we and others have
421 demonstrated that LXR regulation can be gene selective (68, 69).

422 UGCG is a ubiquitously expressed and highly conserved gene. To date no post translational
423 modifications have been identified, and transcriptional regulation appears to be the main
424 determinant of its activity(70). UGCG expression has been shown to be strongly upregulated by a
425 variety of inflammatory signals(71-74), in response to inhibition of prenylation by statin treatment
426 (71-74), and by mTORC2 during tumorigenesis(75). It will be important to establish whether LXR-
427 mediated regulation of UGCG extends to other cell-types and tissues, as this could have wide-
428 reaching implications for the therapeutic activation of LXR in various contexts. For example,
429 elevated expression of UGCG has repeatedly been linked to acquisition of multi-drug resistance
430 and resistance to apoptosis in cancer models(76, 77). More recently, UGCG overexpression was
431 shown to drive enhanced glutamine and mitochondrial metabolism in breast cancer cells (78-80).

432 LXR activation can be pro- or anti-inflammatory depending on the timing of stimulation and
433 species studied (40, 44, 45). LXR activation has previously been reported to inhibit cytokine
434 production by T cells (35, 81, 82), generally attributed to repression of cytokine mRNA
435 transcription (35, 81), which we did not observe here. We confirmed inhibition of proliferation and
436 IL-17 production as previously observed(4, 35, 81, 82). However, we detected an increase in the
437 production of both IL-2 and IL-4 and, in contrast to previous studies, did not observe inhibition of
438 IFN- γ or TNF- α . Because the anti-inflammatory actions of LXR are context dependent(26, 40, 45),
439 it is likely that differences in the conditions for T cell or LXR activation could explain this
440 discrepancy. For example, LXR activation can reduce production of IL-2, TNF α and IFN γ in
441 human CD4+ T cells(81). However, in that study T cells were only briefly stimulated with anti-
442 CD3/28 (6 hours), compared to long-term (72 hours) exposure in our study. Furthermore, a
443 different LXR ligand was used (T0901317), which has also been shown to act on other nuclear
444 receptors(83). This suggests T0901317 activation could have led to LXR-independent effects on

445 T cell function which would differ from those observed with a more specific ligand such as GW. In
446 addition, the timing, duration and strength of stimulus as well as age and sex of donors can all
447 influence LXR signalling(84, 85). Future studies could explore whether these factors are relevant
448 in the LXR-dependent regulation of T cells. In our studies, LXR activation by GW did not
449 significantly alter the induction of cytokine mRNA expression. Instead, the most significantly
450 regulated transcriptional pathways were related to lipid metabolism, and we observed changes in
451 plasma membrane lipid expression early (minutes) and late (72 hours) in the course of T cell
452 activation.

453
454 We identified that LXR-regulated genes and lipids were differentially expressed in Tregs. Like other
455 nuclear receptors, LXR function is orchestrated by a complex combination of factors as mentioned
456 above. Such mechanisms could contribute to subset-specific and gene-specific regulation as we
457 have observed in human T cell subsets and will require further investigation. In murine cells, LXR
458 has been suggested to play a critical role in Treg function(47), increase Foxp3 expression, and
459 promote inducible-Treg differentiation (86). In contrast, LXR activation was recently shown to
460 decrease the frequency of a subset of T cells, intestinal ROR γ t⁺ Tregs, but this was attributed to an
461 indirect effect on myeloid cells (87). While there is currently no evidence of the regulation of Treg
462 and Tresp subsets by LXR in humans, rodent studies point to the importance of LXR β in murine T
463 regs (47). In mouse macrophages, LXR α and LXR β exert overlapping but also specific
464 transcriptional activities(69) although it is currently not known whether this also occurs in other cell
465 types. Future studies will be needed to carefully dissect the mechanisms underlying the cell and
466 LXR isotype-specific mechanisms of UGCG regulation.

467
468 In any case, a potential interaction between LXR signalling, plasma membrane lipids and Tregs
469 has not yet been explored. Murine Tregs also have low membrane order, and genetic deletion of
470 ceramide synthesizing enzyme *smpd1* increases the frequency and suppressive capacity of
471 Tregs (88). This supports a relationship between ceramide metabolism (in which UGCG plays a
472 key role), plasma membrane lipid order, and Treg function. Although plasma membrane
473 cholesterol has been shown to play an important role in the differentiation of Tregs (89),
474 increasing plasma membrane cholesterol was reported to have no effect on their suppressive
475 function (90). In contrast, reduction of intracellular cholesterol by 25-hydroxycholesterol or statin
476 treatment inhibited Treg proliferation and expression of the immune checkpoint receptor CTLA-4
477 (91). Together, this work supports the hypothesis that LXR could contribute to Treg function via
478 modulation of plasma membrane lipid order.

479
480 In addition to the changes in cholesterol and glycosphingolipid metabolism explored here,
481 triacylglycerol (TAG) levels were also substantially upregulated by LXR activation. Compared to
482 conventional T cells, Tregs are lipid-enriched and have increased TAG synthesis and a greater
483 concentration of lipid droplets which serve as a fuel source and protect against lipotoxicity (92).
484 Furthermore, TAG also promote IL-7 mediated memory CD8⁺ T cell survival (93). Thus, the role
485 of LXR-driven TAG biosynthesis in T cells also warrants further investigation, although this was
486 beyond the scope of our current study.

487
488 In their resting state, T cells express low levels of endogenous LXR ligands (94). In our
489 experiments, CYP27A1 was the only oxysterol synthesising enzyme consistently expressed in
490 these cells. However, there is evidence that certain polarisation conditions can lead to dramatic
491 regulation of oxysterol synthesis and thus endogenous modulation of LXR signalling. For
492 example, *in vitro* differentiated type 1 regulatory cells upregulate 25-hydroxycholesterol to limit IL-
493 10 production (94). In contrast, Th17 cells upregulate an enzyme that sulfates oxysterols
494 (SULT2B1), thereby inactivating them as LXR ligands and driving preferential activation of ROR γ t
495 instead of LXR (95). LXR also plays a unique role in a subset of IL-9 producing CD8⁺ T cells
496 (Tc9), in which cholesterol/oxysterol are tightly suppressed to prevent transrepression of the *Il9*
497 locus by LXR (95). Furthermore, changes in oxysterol availability have been documented in many

498 diseases, including accumulation in atherosclerotic plaques (96), production in the tumour
499 microenvironment (6), and reduced circulating levels in multiple sclerosis (97). Therefore, the new
500 mechanism described here could be of therapeutic relevance to disorders characterised by
501 defects in T cell signalling and lipid metabolism. For example, in addition to altered oxysterol
502 levels, multiple sclerosis patients are reported to have altered LXR signalling, cholesterol levels
503 and glycosphingolipid metabolism (96). However, whether plasma membrane lipid rafts contribute
504 to immune-cell dysfunction in multiple sclerosis is currently unknown.

505
506 In conclusion, our findings show for the first time that LXR regulates glycosphingolipid levels,
507 which strongly impacts plasma membrane lipid composition and T cell function. This mechanism
508 is likely to be complementary to others modes of LXR action, including the transcriptional
509 regulation of certain cytokines (6, 35) and modulation of endoplasmic reticulum cholesterol
510 content (4). However, this new mechanism could be of therapeutic relevance to disorders
511 characterised by defects in T cell signalling and metabolism, including autoimmune and
512 neurodegenerative diseases, cardiovascular disease, and cancer.

513 **Materials and Methods**

514 **Antibodies and reagents**

515
516 A detailed list of antibodies and reagents is included in the SI Appendix Methods.
517

518 **Human samples**

519
520 50 mL of peripheral blood was collected from healthy controls (HCs). Men and women aged 18-
521 60 were recruited. Exclusion criteria included current illness/infection, statin treatment,
522 pregnancy, breast-feeding, or vaccination within the past 3 months. For RNA-sequencing and
523 lipidomic analysis of T cells from HCs (Fig. 1) blood, leukocyte cones were purchased from NHS
524 Blood and Transplant. Peripheral blood mononuclear cells (PBMCs) were separated on Ficoll-
525 Paque PLUS (GE Healthcare) using SepMate tubes (StemCell Technologies). PBMCs were
526 cryopreserved in liquid nitrogen until use. Ethical approvals for this work were obtained from the
527 London - City & East Research Ethics Committee (reference 15-LO-2065), Yorkshire & The
528 Humber - South Yorkshire Research Ethics Committee (reference 16/YH/0306), South Central -
529 Hampshire B Research Ethics Committee (reference 18/SC/0323). All participants provided
530 informed written consent.

531 **Cell subset purification**

532
533 *Fluorescence activated cell sorting (FACS)*: CD3⁺ T cells for lipidomics analysis were sorted by
534 FACS. Cells were washed in MACS buffer (PBS with 2% FBS (Labtech) and 1 mM EDTA
535 (Sigma)) before staining with antibodies against surface markers for 30 minutes. Sorting was
536 performed on a BD FACSAria II.
537

538 *Magnetic assisted cell sorting (MACS)*: CD4⁺ T cells and CD19⁺ B cells were negatively isolated
539 using magnetic bead based separation (EasySep, StemCell Technologies). CD14⁺ monocytes
540 were positively selected (EasySep, StemCell Technologies). Sample purities were similar to
541 those reported by the manufacturer (95.1 ± 1.3% for negative selection and 97.6 ± 0.21% for
542 positive selection). To obtain monocyte-derived macrophages (MDMs), monocytes were plated in
543 low-serum media (1% FBS) for 1-2 hours in 12-well Nunc-coated plates (ThermoFisher Scientific)
544 to promote adherence, then cultured for 7 days in complete media (RPMI 1640 (Sigma)
545 supplemented with 10% heat-inactivated fetal bovine serum (FBS) (Labtech) and 20 µg/mL
546 gentamycin (Sigma)).
547

548 **Cell culture**

549 Full details of cell culture conditions and reagents are in the SI Appendix Methods, briefly:

550 *Culture with LXR ligands:* PBMCs or purified T cells were treated with GW3965 (GW) +/- RXR
551 agonist LG100268 (LG) or UGCG inhibitor N-Butyldeoxyojirimycin (NB-DNJ) or with oxysterols,
552 24S-hydroxycholesterol and 24S,25-epoxycholesterol and compared either to vehicle or LXR
553 antagonist GSK1440233 as control.

554 *Functional assays:* T cells were stimulated with anti-CD3 and anti-CD28. To measure
555 intracellular cytokine production cells were additionally stimulated with PMA, ionomycin and
556 GolgiPlug

557

558 **Lipidomics**

559 CD3⁺ T cells were sorted by FACS and plated at 5 x 10⁶/mL into 12 well plates in complete media
560 (n=4). A total of 10-15 x10⁶ cells were treated with DMSO (CTRL) or GW3965 (GW, 1 μM) for 36
561 hours and washed twice in PBS. Frozen cell pellets were shipped to Lipotype GmbH (Dresden,
562 Germany) for mass spectrometry-based lipid analysis as described (98) (see SI Appendix
563 Methods). Lipidomics data has been deposited at Mendeley Data: doi: 10.17632/5rzpnr7w65.1.

564

565 **RNA sequencing and analysis**

566 CD4⁺ T cells (3 x 10⁶) were treated with GW3965 (GW, 2 μM) for 24 hours. The LXR antagonist
567 GSK1440233 (CTRL, 1 μM) was used as a control to suppress baseline endogenous LXR
568 activity. For TCR stimulation, cells were transferred to anti-CD3/28 coated plates for the last 18
569 hours. Total RNA was extracted using TRIzol reagent (Life technologies) followed by DNA-free™
570 DNA Removal Kit (Invitrogen). RNA integrity was confirmed using Agilent's 2200 TapeStation.
571 UCL Genomics (London, UK) performed library preparation and sequencing (see SI Appendix
572 Methods). RNA sequencing files are available at Array Express: E-MTAB-9141.

573

574 **Analysis of gene expression**

575 Gene expression was measured by qPCR, as in (64, 99). Primers were used at a final
576 concentration of 100 nM. Sequences are provided in SI Appendix, Table S3.

577

578 **Flow cytometry**

579 Flow cytometry staining was performed as previously described (7, 9). (See SI Appendix
580 Methods).

581

582 **Immunoblotting**

583 Cells were lysed in RIPA buffer and immunoblotting was performed as previously described (99).
584 Semi-quantitative analysis was conducted using the gel analysis module in ImageJ (National
585 Institutes of Health, USA, RRID:SCR_003070).

586

587 **Chromatin immunoprecipitation**

588 Detailed description can be found in the SI Appendix Methods.

589

590 **Microscopy**

591 *Immunostaining:* CD4⁺ T cells were incubated in antibody coated chamber slides for 15 minutes
592 at 37°C, 5% CO₂ to facilitate synapse formation. Medium and non-adherent cells were discarded,
593 and wells were washed gently with PBS before fixation (4% PFA, 2% sucrose, 140 mM NaOH,
594 pH 7.2) for 20 minutes at RT. Formaldehyde was quenched with two washes in 0.1 M ammonium
595 chloride (Sigma-Aldrich), followed by a PBS wash. 0.2% Triton-X-100 was used to permeabilise
596 cells for 8 minutes at RT. Samples were blocked with 5% BSA in PBS + 0.2% fish skin gelatin
597 (Sigma-Aldrich) overnight at 4°C. Primary antibodies were added in blocking solution for 1 hour at
598 RT, followed by addition of fluorescently conjugated secondary antibodies for 30 minutes, RT.
599 Cells were preserved in Prolong Diamond mounting media with DAPI (Invitrogen). For fixed
600 synapses were stained with phalloidin-FITC conjugate (Sigma).

601

602 *Confocal microscopy*: Single slices were acquired on a Leica SPE2 confocal microscope with an
603 x63 oil-immersion objective and 488 and 633 nM excitation solid-state lasers, using the following
604 settings: 1024x1024 pixels, 600 Hz and line average of 3.

605
606 *Total Internal Reflection Fluorescence (TIRF) Microscopy*: To record live cells stained with ANE,
607 a customized two-channel set up was used as described by Ashdown et al. (61) and in the SI
608 Appendix Methods. 30-minute movies were acquired at a rate of 1 frame/minute. The background
609 MFI was based on three measurements taken from the area surrounding each cell.

610
611 *Image analysis*: Image analysis was performed using ImageJ 1.51 (National Institutes of Health,
612 USA, RRID:SCR_003070). Fluorescence intensity was analysed using the 'Analyze Particles'
613 function. Mean fluorescence intensity (MFI) was measured as mean grey scale value (between 0
614 and 255), and corrected total cell fluorescence (CTCF) was calculated as follows: $CTCF =$
615 $integrated\ density - (cell\ area \times MFI\ of\ background)$. To analyze TIRF movies of ANE-stained
616 cells ordered and disordered channels were aligned using the Cairn Image Splitter plugin.
617 Membrane lipid order was calculated as a GP ratio, using the plugin at
618 <https://github.com/quokka79/GPcalc> (GitHub, RRID: SCR_002630). Hue, saturation and
619 brightness (HSB) images were set to visualize GP and pseudocoloured using the Rainbow RGB
620 look up table.

621

622 **Statistical analysis**

623 Statistical tests were performed in GraphPad Prism 8 (GraphPad Software, La Jolla California
624 USA, RRID: SCR_002798, www.graphpad.com) unless otherwise stated. The D'Agostino &
625 Pearson omnibus K2 test was used to check whether datasets were normally distributed. In some
626 cases extreme outliers were removed based upon a ROUT test (Q=1%). Un-paired two-tailed t-
627 tests or Mann-Whitney U were used to compare between independent groups and are
628 represented as bar charts (mean \pm SD) or violin plots (median and interquartile range). In line
629 with previous studies on LXR agonism in human cells (7, 81, 100), paired two-tailed t-tests or
630 repeated measures ANOVA were used where cells from the same donor sample were exposed to
631 different treatments (e.g. GW vs CTRL). This minimizes the impact of donor-to-donor
632 heterogeneity at baseline. Where paired tests were applied, data is presented as paired line
633 graphs. Correction for multiple comparisons was made with Tukey's post-hoc test or Dunnet's
634 test (to compare all samples to vehicle), as specified. For Figure 5b, p-values from multiple un-
635 paired t-test were corrected using the two-stage linear step-up procedure of Benjamini, Krieger
636 and Yekutieli with FDR threshold of 5%.

637

638

639 **Acknowledgments**

640 We are grateful to K.R. Steffenson for the provision of the LXR antibody used for chromatin
641 immunoprecipitation and to A. Castrillo and J. Thorne for earlier discussions on LXRE
642 identification in the UGCG gene.

643
644 KEW was funded by a British Heart Foundation PhD Studentship (FS/13/59/30649) and MS
645 Society (Grant 76). IPT was funded by a Medical Research Council New Investigator Grant
646 (G0801278), a British Heart Foundation Project Grant (PG/13/10/30000) and an Academy of
647 Medical Sciences Newton Advanced Fellowship. ECJ was funded by the Innovative Medicines
648 Initiative Joint grant agreement n° 115303, as part of the ABIRISK consortium (Anti-
649 Biopharmaceutical Immunization: Prediction and analysis of clinical relevance to minimize the
650 risk), Arthritis Research UK Fellowships (20085 and 18106), Lupus UK, The Rosetrees Trust
651 (M409), and University College London Hospital Clinical Research and Development Committee
652 project grant (GCT/2008/EJ and Fast Track grant F193).

653

654 **References**

655

656

- 657 1. K. Simons, E. Ikonen, Functional rafts in cell membranes. *Nature* **387**, 569-572 (1997).
- 658 2. M. Swamy *et al.*, A Cholesterol-Based Allosteric Model of T Cell Receptor
659 Phosphorylation. *Immunity* **44**, 1091-1101 (2016).
- 660 3. F. Wang, K. Beck-García, C. Zorzín, W. W. A. Schamel, M. M. Davis, Inhibition of T cell
661 receptor signaling by cholesterol sulfate, a naturally occurring derivative of membrane
662 cholesterol. *Nature immunology* **17**, 844-850 (2016).
- 663 4. S. J. Bensinger *et al.*, LXR signaling couples sterol metabolism to proliferation in the
664 acquired immune response. *Cell* **134**, 97-111 (2008).
- 665 5. Y. Kidani *et al.*, Sterol regulatory element-binding proteins are essential for the
666 metabolic programming of effector T cells and adaptive immunity. *Nat Immunol* **14**, 489-
667 499 (2013).
- 668 6. X. Ma *et al.*, Cholesterol negatively regulates IL-9-producing CD8(+) T cell differentiation
669 and antitumor activity. *J Exp Med* **215**, 1555-1569 (2018).
- 670 7. G. McDonald *et al.*, Normalizing glycosphingolipids restores function in CD4+ T cells from
671 lupus patients. *J Clin Invest* **124**, 712-724 (2014).
- 672 8. Y. Zhu *et al.*, Lowering glycosphingolipid levels in CD4+ T cells attenuates T cell receptor
673 signaling, cytokine production, and differentiation to the Th17 lineage. *J Biol Chem* **286**,
674 14787-14794 (2011).
- 675 9. L. Miguel *et al.*, Primary human CD4+ T cells have diverse levels of membrane lipid order
676 that correlate with their function. *J Immunol* **186**, 3505-3516 (2011).
- 677 10. D. M. Owen *et al.*, High plasma membrane lipid order imaged at the immunological
678 synapse periphery in live T cells. *Mol Membr Biol* **27**, 178-189 (2010).
- 679 11. K. Simons, E. Ikonen, How Cells Handle Cholesterol. *Science* **290**, 1721-1726 (2000).
- 680 12. K. Gaus, E. Chklovskaya, B. Fazekas de St Groth, W. Jessup, T. Harder, Condensation of
681 the plasma membrane at the site of T lymphocyte activation. *J Cell Biol* **171**, 121-131
682 (2005).
- 683 13. C. Rentero *et al.*, Functional implications of plasma membrane condensation for T cell
684 activation. *PLoS One* **3**, e2262 (2008).

- 685 14. W. Wu, X. Shi, C. Xu, Regulation of T cell signalling by membrane lipids. *Nat Rev Immunol*
686 **16**, 690-701 (2016).
- 687 15. W. Yang *et al.*, Potentiating the antitumour response of CD8(+) T cells by modulating
688 cholesterol metabolism. *Nature* **531**, 651-655 (2016).
- 689 16. D. Sviridov, N. Mukhamedova, Y. I. Miller, Lipid rafts as a therapeutic target. *J Lipid Res*
690 **61**, 687-695 (2020).
- 691 17. N. S. Heaton, G. Randall, Multifaceted roles for lipids in viral infection. *Trends Microbiol*
692 **19**, 368-375 (2011).
- 693 18. E. C. Jury, P. S. Kabouridis, F. Flores-Borja, R. A. Mageed, D. A. Isenberg, Altered lipid
694 raft-associated signaling and ganglioside expression in T lymphocytes from patients with
695 systemic lupus erythematosus. *J Clin Invest* **113**, 1176-1187 (2004).
- 696 19. E. C. Jury, D. A. Isenberg, C. Mauri, M. R. Ehrenstein, Atorvastatin restores Lck
697 expression and lipid raft-associated signaling in T cells from patients with systemic lupus
698 erythematosus. *J Immunol* **177**, 7416-7422 (2006).
- 699 20. B. Wang, P. Tontonoz, Liver X receptors in lipid signalling and membrane homeostasis.
700 *Nat Rev Endocrinol* **14**, 452-463 (2018).
- 701 21. J. L. Collins *et al.*, Identification of a nonsteroidal liver X receptor agonist through parallel
702 array synthesis of tertiary amines. *J Med Chem* **45**, 1963-1966 (2002).
- 703 22. A. C. Calkin, P. Tontonoz, Transcriptional integration of metabolism by the nuclear
704 sterol-activated receptors LXR and FXR. *Nat Rev Mol Cell Biol* **13**, 213-224 (2012).
- 705 23. J. N. Benhammou *et al.*, Novel Lipid Long Intervening Noncoding RNA, Oligodendrocyte
706 Maturation-Associated Long Intergenic Noncoding RNA, Regulates the Liver Steatosis
707 Gene Stearoyl-Coenzyme A Desaturase As an Enhancer RNA. *Hepatol Commun* **3**, 1356-
708 1372 (2019).
- 709 24. E. A. DiBlasio-Smith *et al.*, Discovery and implementation of transcriptional biomarkers
710 of synthetic LXR agonists in peripheral blood cells. *J Transl Med* **6**, 59 (2008).
- 711 25. B. A. Laffitte *et al.*, Autoregulation of the human liver X receptor alpha promoter. *Mol*
712 *Cell Biol* **21**, 7558-7568 (2001).
- 713 26. K. E. Waddington, E. C. Jury, I. Pineda-Torra, Liver X receptors in immune cell function in
714 humans. *Biochem Soc Trans* **43**, 752-757 (2015).
- 715 27. K. R. Levental *et al.*, Polyunsaturated Lipids Regulate Membrane Domain Stability by
716 Tuning Membrane Order. *Biophys J* **110**, 1800-1810 (2016).
- 717 28. E. Sezgin, I. Levental, S. Mayor, C. Eggeling, The mystery of membrane organization:
718 composition, regulation and roles of lipid rafts. *Nat Rev Mol Cell Biol* **18**, 361-374 (2017).
- 719 29. A. F. Valledor *et al.*, Activation of liver X receptors and retinoid X receptors prevents
720 bacterial-induced macrophage apoptosis. *Proc Natl Acad Sci U S A* **101**, 17813-17818
721 (2004).
- 722 30. B. A. Laffitte *et al.*, The phospholipid transfer protein gene is a liver X receptor target
723 expressed by macrophages in atherosclerotic lesions. *Mol Cell Biol* **23**, 2182-2191
724 (2003).
- 725 31. E. D. Muse *et al.*, Cell-specific discrimination of desmosterol and desmosterol mimetics
726 confers selective regulation of LXR and SREBP in macrophages. *Proc Natl Acad Sci U S A*
727 **115**, E4680-e4689 (2018).
- 728 32. P. Pehkonen *et al.*, Genome-wide landscape of liver X receptor chromatin binding and
729 gene regulation in human macrophages. *BMC Genomics* **13**, 50 (2012).

- 730 33. I. Lavrnja *et al.*, Expression profiles of cholesterol metabolism-related genes are altered
731 during development of experimental autoimmune encephalomyelitis in the rat spinal
732 cord. *Sci Rep* **7**, 2702 (2017).
- 733 34. P. B. Noto *et al.*, Regulation of sphingomyelin phosphodiesterase acid-like 3A gene
734 (SMPDL3A) by liver X receptors. *Mol Pharmacol* **82**, 719-727 (2012).
- 735 35. G. Cui *et al.*, Liver X receptor (LXR) mediates negative regulation of mouse and human
736 Th17 differentiation. *J Clin Invest* **121**, 658-670 (2011).
- 737 36. D. Lingwood, K. Simons, Lipid rafts as a membrane-organizing principle. *Science* **327**, 46-
738 50 (2010).
- 739 37. R. Varma, G. Campi, T. Yokosuka, T. Saito, M. L. Dustin, T cell receptor-proximal signals
740 are sustained in peripheral microclusters and terminated in the central supramolecular
741 activation cluster. *Immunity* **25**, 117-127 (2006).
- 742 38. F. Bovenga, C. Sabbà, A. Moschetta, Uncoupling nuclear receptor LXR and cholesterol
743 metabolism in cancer. *Cell Metab* **21**, 517-526 (2015).
- 744 39. K. Mouzat *et al.*, Regulation of Brain Cholesterol: What Role Do Liver X Receptors Play in
745 Neurodegenerative Diseases? *Int J Mol Sci* **20** (2019).
- 746 40. C. Fontaine *et al.*, Liver X receptor activation potentiates the lipopolysaccharide
747 response in human macrophages. *Circ Res* **101**, 40-49 (2007).
- 748 41. S. Ghisletti *et al.*, Parallel SUMOylation-dependent pathways mediate gene- and signal-
749 specific transrepression by LXRs and PPARgamma. *Mol Cell* **25**, 57-70 (2007).
- 750 42. A. Ito *et al.*, LXRs link metabolism to inflammation through Abca1-dependent regulation
751 of membrane composition and TLR signaling. *Elife* **4**, e08009 (2015).
- 752 43. X. Rong *et al.*, LXRs regulate ER stress and inflammation through dynamic modulation of
753 membrane phospholipid composition. *Cell Metab* **18**, 685-697 (2013).
- 754 44. Y. Sohrabi *et al.*, LXR Activation Induces a Proinflammatory Trained Innate Immunity-
755 Phenotype in Human Monocytes. *Front Immunol* **11**, 353-353 (2020).
- 756 45. M. Ishibashi *et al.*, Liver x receptor regulates arachidonic acid distribution and
757 eicosanoid release in human macrophages: a key role for lysophosphatidylcholine
758 acyltransferase 3. *Arterioscler Thromb Vasc Biol* **33**, 1171-1179 (2013).
- 759 46. C. T. Chan *et al.*, Liver X receptors are required for thymic resilience and T cell output. *J*
760 *Exp Med* **217** (2020).
- 761 47. A. J. Michaels, C. Campbell, R. Bou-Puerto, A. Y. Rudensky, Nuclear receptor LXRβ
762 controls fitness and functionality of activated T cells. *Journal of Experimental Medicine*
763 **218** (2020).
- 764 48. J. M. Carbó *et al.*, Pharmacological activation of LXR alters the expression profile of
765 tumor-associated macrophages and the abundance of regulatory T cells in the tumor
766 microenvironment. *Cancer Res* 10.1158/0008-5472.Can-19-3360 (2020).
- 767 49. A. Ito *et al.*, LXRs link metabolism to inflammation through Abca1-dependent regulation
768 of membrane composition and TLR signaling. *Elife* **4**, e08009-e08009 (2015).
- 769 50. N. Blank, M. Schiller, C. Gabler, J. R. Kalden, H. M. Lorenz, Inhibition of sphingolipid
770 synthesis impairs cellular activation, cytokine production and proliferation in human
771 lymphocytes. *Biochem Pharmacol* **71**, 126-135 (2005).
- 772 51. L. Miguel *et al.*, Primary Human CD4⁺ T Cells Have Diverse Levels of
773 Membrane Lipid Order That Correlate with Their Function. *The Journal of Immunology*
774 **186**, 3505-3516 (2011).

- 775 52. I. Gombos, E. Kiss, C. Detre, G. László, J. Matkó, Cholesterol and sphingolipids as lipid
776 organizers of the immune cells' plasma membrane: their impact on the functions of
777 MHC molecules, effector T-lymphocytes and T-cell death. *Immunol Lett* **104**, 59-69
778 (2006).
- 779 53. P. S. Kabouridis, E. C. Jury, Lipid rafts and T-lymphocyte function: Implications for
780 autoimmunity. *FEBS Letters* **582**, 3711-3718 (2008).
- 781 54. R. A. Maldonado, D. J. Irvine, R. Schreiber, L. H. Glimcher, A role for the immunological
782 synapse in lineage commitment of CD4 lymphocytes. *Nature* **431**, 527-532 (2004).
- 783 55. F. Balamuth, D. Leitenberg, J. Unternaehrer, I. Mellman, K. Bottomly, Distinct patterns of
784 membrane microdomain partitioning in Th1 and th2 cells. *Immunity* **15**, 729-738 (2001).
- 785 56. E. Izsepi *et al.*, Membrane microdomain organization, calcium signal, and NFAT
786 activation as an important axis in polarized Th cell function. *Cytometry A* **83**, 185-196
787 (2013).
- 788 57. M. V. Farrell, S. Webster, K. Gaus, J. Goyette, T Cell Membrane Heterogeneity Aids
789 Antigen Recognition and T Cell Activation. *Frontiers in Cell and Developmental Biology* **8**
790 (2020).
- 791 58. C. S. Guy *et al.*, Distinct TCR signaling pathways drive proliferation and cytokine
792 production in T cells. *Nature immunology* **14**, 262-270 (2013).
- 793 59. A. Molano *et al.*, Age-dependent changes in the sphingolipid composition of mouse
794 CD4+ T cell membranes and immune synapses implicate glucosylceramides in age-
795 related T cell dysfunction. *PLoS One* **7**, e47650 (2012).
- 796 60. G. Wu, Z. H. Lu, H. J. Gabius, R. W. Ledeen, D. Bleich, Ganglioside GM1 deficiency in
797 effector T cells from NOD mice induces resistance to regulatory T-cell suppression.
798 *Diabetes* **60**, 2341-2349 (2011).
- 799 61. G. W. Ashdown *et al.*, Membrane lipid order of sub-synaptic T cell vesicles correlates
800 with their dynamics and function. *Traffic* **19**, 29-35 (2018).
- 801 62. T. Zech *et al.*, Accumulation of raft lipids in T-cell plasma membrane domains engaged in
802 TCR signalling. *EMBO J* **28**, 466-476 (2009).
- 803 63. Z. Czimmerer, L. Halasz, L. Nagy, Unorthodox Transcriptional Mechanisms of Lipid-
804 Sensing Nuclear Receptors in Macrophages: Are We Opening a New Chapter? *Front*
805 *Endocrinol (Lausanne)* **11**, 609099-609099 (2020).
- 806 64. N. Becares *et al.*, Impaired LXR α Phosphorylation Attenuates Progression of Fatty Liver
807 Disease. *Cell Rep* **26**, 984-995.e986 (2019).
- 808 65. Q. Zhang *et al.*, MS-275 induces hepatic FGF21 expression via H3K18ac-mediated CREBH
809 signal. *J Mol Endocrinol* **62**, 187-196 (2019).
- 810 66. Z. L. Cui *et al.*, Histone modifications of Notch1 promoter affect lung CD4+ T cell
811 differentiation in asthmatic rats. *Int J Immunopathol Pharmacol* **26**, 371-381 (2013).
- 812 67. Z. Czimmerer, L. Halasz, L. Nagy, Unorthodox Transcriptional Mechanisms of Lipid-
813 Sensing Nuclear Receptors in Macrophages: Are We Opening a New Chapter? *Front*
814 *Endocrinol (Lausanne)* **11**, 609099 (2020).
- 815 68. N. Becares, M. C. Gage, I. Pineda-Torra, Posttranslational Modifications of Lipid-
816 Activated Nuclear Receptors: Focus on Metabolism. *Endocrinology* **158**, 213-225 (2017).
- 817 69. A. Ramón-Vázquez *et al.*, Common and Differential Transcriptional Actions of Nuclear
818 Receptors Liver X Receptors α and β in Macrophages. *Mol Cell Biol* **39** (2019).

- 819 70. Y. Ishibashi, A. Kohyama-Koganeya, Y. Hirabayashi, New insights on glucosylated lipids:
820 Metabolism and functions. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell*
821 *Biology of Lipids* **1831**, 1475-1485 (2013).
- 822 71. M. S. Köberlin *et al.*, A Conserved Circular Network of Coregulated Lipids Modulates
823 Innate Immune Responses. *Cell* **162**, 170-183 (2015).
- 824 72. P. J. Brennan *et al.*, Invariant natural killer T cells recognize lipid self antigen induced by
825 microbial danger signals. *Nat Immunol* **12**, 1202-1211 (2011).
- 826 73. M. K. Pandey *et al.*, Complement drives glucosylceramide accumulation and tissue
827 inflammation in Gaucher disease. *Nature* **543**, 108-112 (2017).
- 828 74. A. K. Iyer, J. Liu, R. M. Gallo, M. H. Kaplan, R. R. Brutkiewicz, STAT3 promotes CD1d-
829 mediated lipid antigen presentation by regulating a critical gene in glycosphingolipid
830 biosynthesis. *Immunology* **146**, 444-455 (2015).
- 831 75. B. Binnington *et al.*, Inhibition of Rab prenylation by statins induces cellular
832 glycosphingolipid remodeling. *Glycobiology* **26**, 166-180 (2016).
- 833 76. M. W. Khan *et al.*, mTORC2 controls cancer cell survival by modulating gluconeogenesis.
834 *Cell Death Discovery* **1**, 15016 (2015).
- 835 77. Y. Guri *et al.*, mTORC2 Promotes Tumorigenesis via Lipid Synthesis. *Cancer Cell* **32**, 807-
836 823.e812 (2017).
- 837 78. Y. Y. Liu, R. A. Hill, Y. T. Li, Ceramide glycosylation catalyzed by glucosylceramide
838 synthase and cancer drug resistance. *Adv Cancer Res* **117**, 59-89 (2013).
- 839 79. Y. Lavie, H. Cao, S. L. Bursten, A. E. Giuliano, M. C. Cabot, Accumulation of
840 glucosylceramides in multidrug-resistant cancer cells. *J Biol Chem* **271**, 19530-19536
841 (1996).
- 842 80. M. S. Wegner *et al.*, UDP-glucose ceramide glucosyltransferase activates AKT, promoted
843 proliferation, and doxorubicin resistance in breast cancer cells. *Cell Mol Life Sci* **75**, 3393-
844 3410 (2018).
- 845 81. D. Walcher *et al.*, LXR activation reduces proinflammatory cytokine expression in human
846 CD4-positive lymphocytes. *Arterioscler Thromb Vasc Biol* **26**, 1022-1028 (2006).
- 847 82. L. Wu *et al.*, Activation of the liver X receptor inhibits Th17 and Th1 responses in
848 Behcet's disease and Vogt-Koyanagi-Harada disease. *Curr Mol Med* **14**, 712-722 (2014).
- 849 83. K. A. Houck *et al.*, T0901317 is a dual LXR/FXR agonist. *Mol Genet Metab* **83**, 184-187
850 (2004).
- 851 84. S. Della Torre *et al.*, An Essential Role for Liver ER α in Coupling Hepatic Metabolism to
852 the Reproductive Cycle. *Cell Rep* **15**, 360-371 (2016).
- 853 85. K. Y. DeLeon-Pennell *et al.*, LXR/RXR signaling and neutrophil phenotype following
854 myocardial infarction classify sex differences in remodeling. *Basic Res Cardiol* **113**, 40
855 (2018).
- 856 86. M. Herold *et al.*, Liver X receptor activation promotes differentiation of regulatory T
857 cells. *PLoS One* **12**, e0184985 (2017).
- 858 87. S. M. Parigi *et al.*, Liver X receptor regulates Th17 and ROR γ t(+) Treg cells by distinct
859 mechanisms. *Mucosal Immunol* 10.1038/s41385-020-0323-5 (2020).
- 860 88. C. Hollmann *et al.*, Inhibition of Acid Sphingomyelinase Allows for Selective Targeting of
861 CD4+ Conventional versus Foxp3+ Regulatory T Cells. *J Immunol* **197**, 3130-3141 (2016).
- 862 89. H. Y. Cheng *et al.*, Loss of ABCG1 influences regulatory T cell differentiation and
863 atherosclerosis. *J Clin Invest* **126**, 3236-3246 (2016).

- 864 90. J. Surls *et al.*, Increased membrane cholesterol in lymphocytes diverts T-cells toward an
865 inflammatory response. *PLoS One* **7**, e38733 (2012).
- 866 91. H. Zeng *et al.*, mTORC1 couples immune signals and metabolic programming to establish
867 T(reg)-cell function. *Nature* **499**, 485-490 (2013).
- 868 92. D. Howie *et al.*, A Novel Role for Triglyceride Metabolism in Foxp3 Expression. *Front*
869 *Immunol* **10**, 1860 (2019).
- 870 93. G. Cui *et al.*, IL-7-Induced Glycerol Transport and TAG Synthesis Promotes Memory CD8+
871 T Cell Longevity. *Cell* **161**, 750-761 (2015).
- 872 94. S. Vigne *et al.*, IL-27-Induced Type 1 Regulatory T-Cells Produce Oxysterols that
873 Constrain IL-10 Production. *Front Immunol* **8** (2017).
- 874 95. X. Hu *et al.*, Sterol metabolism controls T(H)17 differentiation by generating endogenous
875 ROR γ agonists. *Nat Chem Biol* **11**, 141-147 (2015).
- 876 96. N. J. Spann *et al.*, Regulated accumulation of desmosterol integrates macrophage lipid
877 metabolism and inflammatory responses. *Cell* **151**, 138-152 (2012).
- 878 97. K. Fellows Maxwell *et al.*, Oxysterols and apolipoproteins in multiple sclerosis: a 5 year
879 follow-up study. *J Lipid Res* **60**, 1190-1198 (2019).
- 880 98. J. L. Sampaio *et al.*, Membrane lipidome of an epithelial cell line. *Proc Natl Acad Sci U S A*
881 **108**, 1903-1907 (2011).
- 882 99. M. C. Gage *et al.*, Disrupting LXRA phosphorylation promotes FoxM1 expression and
883 modulates atherosclerosis by inducing macrophage proliferation. *Proc Natl Acad Sci U S*
884 *A* **115**, E6556-e6565 (2018).
- 885 100. D. L. Asquith *et al.*, The liver X receptor pathway is highly upregulated in rheumatoid
886 arthritis synovial macrophages and potentiates TLR-driven cytokine release. *Ann Rheum*
887 *Dis* **72**, 2024-2031 (2013).

888

889

890

891

892 **Figure Legends**

893

894 **Figure 1. LXR regulates lipid metabolism in human CD4⁺ T cells.** Primary human CD4⁺ T
895 cells (n=3) were cultured with or without LXR agonist (GW3965, GW) for 24 hours. Gene
896 expression was assessed by RNA-Seq. **(a)** Volcano plot showing fold changes and p-values.
897 Coloured points represent significantly regulated genes (p < 0.05). **(b)** Clustered heatmap of
898 normalised gene counts of all LXR-regulated genes with FDR corrected p < 0.1. **(c)** Regulation of
899 a selection of genes was confirmed by qPCR in an independent set of donors (n=3-6). Bars
900 represent mean ± SD. Unpaired two-tailed t-test; *p < 0.05, **p < 0.01, ****p < 0.0001. **(d-e)**
901 Network diagrams illustrate pathways significantly enriched for up- **(d)** or down- **(e)** regulated
902 genes. Each node represents a significantly enriched term, with node size proportional to the
903 number of contributing genes. Similar terms with a high degree of redundancy were clustered, as
904 depicted. Bar charts plot cluster significance and show enrichment ratios (ER). **(f)** Pie chart
905 showing the proportion of genes regulated by GW in human T cells that are also regulated in
906 murine bone marrow derived macrophages (BMDMs) (21) or peritoneal macrophages (pMφ) (22).
907

908 **Figure 2. LXR activation regulates the transcription of glucosylceramide synthase (UGCG)**
909 **(a-e)** Primary human CD3⁺ T cells (n=4) were sorted by FACS and treated ± GW (2 μM) for 36
910 hours and total cellular lipid content analysed by shotgun lipidomics. **(a)** Total lipids (normalized
911 to cell numbers) were unchanged (mean ± SD). **(b)** Volcano plot represents significant changes in
912 the expression of lipid sub-species, colour coded by broader lipid class (p < 0.05). **(c)** Bars show
913 the number of subspecies detected for each lipid type. The filled area represents the proportion of
914 subspecies significantly altered by GW treatment. **(d)** Unclustered heatmaps represent levels of
915 individual sub-species. **(e)** Dot plots show overall change in triacylglycerol (TAG) and
916 hexosylceramide (HexCer) levels. **(f)** Schematic illustrating the role of UGCG in the conversion of
917 ceramide to HexCer. **(g)** Pie chart showing a GW-induced shift from ceramide (Cer) to
918 hexosylceramide (HexCer). **(h)** Upregulation of UGCG mRNA expression in CD4⁺ T cells after 24
919 hours GW treatment (n=13). **(i)** Cells were treated with LXR (GW, 1 μM) and RXR (LG100268;
920 LG, 100 nM) ligands for 2 hours. LXR occupancy at the putative DR4 motif at *UGCG* compared to
921 IgG control, positive control (*SMPDL3A*) and negative control (*RPLP0*) sequences.
922 Representative of three independent experiments. **(a-h)** Two-tailed t-tests: *p < 0.05, **p < 0.01,
923 ***p < 0.001.

924 Abbreviations: Cer – ceramide, PE – phosphatidylethanolamine, DAG – diacylglycerol, HexCer –
925 hexosylceramide, SM – sphingomyelin, PE-O – phosphatidylethanolamine-ether, PI –
926 phosphatidylinositol, PC – phosphatidylcholine, PS – phosphatidylserine, PC-O –
927 phosphatidylcholine-ether, LPI – lyso-phosphatidylinositol, LPE – lyso-phosphatidylethanolamine,
928 CE – cholesterol esters, PA – phosphatidate, CL – cardiolipin, LPG – lyso-phosphatidylglycerol ,
929 LPA – lyso-phosphatidate, LPS – lyso- phosphatidylserine.

930

931

932 **Figure 3. Plasma membrane lipid order is reduced by LXR activation.** **(a-c)** Cells were
933 cultured ± LXR ligands for 24 hours (GW) or 72 hours (24S,25-EC and 24S-OHC) and flow
934 cytometry was used to identify CD4⁺ T cells and measure plasma membrane lipid expression in
935 >4 independent experiments. **(a)** Representative flow cytometry plots show the percentage of T
936 cells highly expressing cholera toxin B (CTB) and CTB gMFI as surrogate markers for
937 glycosphingolipids, as in(9). Cumulative data shows change in percentage of cells highly
938 expressing CTB. **(b)** Representative histogram of filipin staining for cholesterol, and cumulative
939 data showing change in gMFI. **(c)** Cumulative data showing GSL/cholesterol ratio as CTB/filipin
940 (n=6). **(d-f)** Magnetically purified CD4⁺ T cell membrane lipid order was measured using di-4-
941 ANEPPDHQ. **(d)** Representative confocal microscopy image and a histogram of average
942 generalised polarisation (GP) ratio per image analysed are shown (n=1 donor). **(e)** Cumulative
943 data from three experiments showing lipid order measured by flow cytometry. Cells were treated
944 with an LXR agonist (GW) or antagonist (GSK233)(n=5) for 24 hours. **(f)** di-4-ANEPPDHQ-
945 stained CD4⁺ T cells (n=4) were sorted into high or low membrane order by FACS, and gene

946 expression was compared by qPCR. Bars show mean \pm SD. **(a-f)** Two-tailed t-tests or one-way
947 ANOVA with Tukey's posthoc test; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

948 **Figure 4. LXR activation modulates T cell immune function.** **(a-c)** RNA-seq was performed
949 on magnetically-isolated CD4⁺ T cells incubated \pm GW for 6 hours, before stimulation with anti-
950 CD3/CD28 (TCR) \pm GW (n=3). **(a)** Venn diagrams compare the number of genes up or down
951 regulated by TCR stimulation in the presence (red) or absence (blue) of GW. **(b)** Volcano plot of
952 genes differentially expressed between GW+TCR and CTRL+TCR. **(c)** Normalized RNA-Seq
953 gene counts of differentially expressed genes were compared to resting T cells. Four patterns of
954 gene expression were identified by hierarchical clustering (clusters A-D). One gene from each
955 cluster is shown as an example (mean \pm SD) and the most significantly enriched gene ontology
956 (GO) term is given. **(d-f)** Representative flow cytometry plots and cumulative data from 4
957 independent experiments (n=6–9) show the effect of GW on the plasma membrane cholesterol
958 **(d)** and glycosphingolipid **(e)** content of activated CD4⁺ T cells and the ratio of GSLs to
959 cholesterol **(f)**. **(g)** Schematic illustrating enzymes controlling oxysterol metabolism. Bar charts
960 show normalized RNA-seq gene counts of enzymes significantly regulated by TCR activation
961 (mean \pm SD, n=3). *FDR < 0.1. **(h-j)** Magnetically-isolated CD4⁺ T cells were activated with anti-
962 CD3/CD28 (+TCR) for 72 hours in the presence of GW3965 (GW) or control (CTRL). **(h-j)**
963 Intracellular cytokines were analysed by flow cytometry after additional treatment with PMA and
964 ionomycin **(h-i)** and Ki67 was used as a marker of proliferation **(j)**. Representative flow cytometry
965 plots are labelled with percentage of positive cells and gMFI of both the cytokine-
966 producing/proliferating population and total T cells. Cumulative data from four independent
967 experiments shows cytokine production expressed as an integrated MFI (iMFI = gMFI*frequency
968 of cytokine producing cells (11))(i) or percentage of proliferating cells **(j)**. Two-tailed t-tests; * $p <$
969 0.05, ** $p < 0.01$, *** $p < 0.001$.

970 **Figure 5. LXR activation regulates immune synapse formation and proximal TCR**
971 **signalling.** **(a-d)** Magnetically purified CD4⁺ T cells were cultured \pm GW before addition to
972 chamber slides coated with anti-CD3/28 for immune synapse formation. **(a-b)** T cells were
973 stained with di-4-ANEPPDHQ and immune synapse formation was recorded for 30 minutes using
974 TIRF microscopy. **(a)** Representative images at 5 minute intervals, scale bar = 5 μ M. **(b)** GP ratio
975 was quantified at each minute (n= 10-12 cells/condition, mean \pm SEM). **(c-d)** Immune synapses
976 (n=2 donors) were fixed at 15 mins post activation and immunostained for Lck (CTRL=68 cells,
977 GW=52 cells) and phosphotyrosine (pY) (CTRL=59 cells, GW=52 cells). Representative images
978 and quantification of corrected total cell fluorescence (CTCF) **(c)** or classification of Lck
979 distribution patterns **(d)**. Violin plots show median and quartile values. Multiple unpaired t-tests
980 corrected for multiple comparisons **(b)** or Mann Whitney U **(c-d)**: * $p < 0.05$, ** $p < 0.01$, *** $p <$
981 0.001. Abbreviations: Lck – lymphocyte-specific protein tyrosine kinase; pY – phosphotyrosine.

982 **Figure 6. Treg and Trespander subsets have distinct lipid metabolic phenotypes.** **(a)**
983 Responder (Tresp: CD4⁺CD25^{lo}CD127⁺) and regulatory (Treg: CD4⁺CD25⁺CD127⁻) T cell subsets
984 were defined by flow cytometry. **(b-d)** Plasma membrane lipid order (GP ratio) **(b)**,
985 glycosphingolipid levels (GSL) **(c)**, and cholesterol content **(d)** were analysed using flow
986 cytometry. Lines connect matched Tresp and Treg results from the same sample. **(e)** Expression
987 of LXR and LXR-target genes that regulate cholesterol, GSL, and fatty acid levels were analysed
988 in FACS sorted T cell subsets (n=3-8). Mean \pm SD. **(f-i)** Cells were treated with GW for 24 hours.
989 Lines connect control (CTRL) and GW treated samples from the same donor. Cumulative data
990 from 3 independent experiments shows the change in membrane lipid order (GP ratio) **(f)**,
991 cholesterol **(g)**, and GSL **(h)** expression. **(g-i)** Induction of LXR target genes involved in
992 cholesterol **(g)**, GSL **(h)** and fatty acid metabolism **(i)** were analysed in FACS-sorted T cell
993 subsets (n=5-6). Gene expression is expressed relative to the average of control (CTRL) treated
994 Tresp. The average fold change (GW vs CTRL) was calculated for each subset. Two-tailed t-
995 tests; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

