

Translation of experimental cardioprotective capability of P2Y₁₂ inhibitors into clinical outcome in patients with ST-elevation myocardial infarction

Short title: Clinical impact of P2Y₁₂ inhibitors' cardioprotective capability

Marie V. Hjortbak¹; Kevin K.W. Olesen²; Jacob M. Seefeldt¹, Thomas R. Lassen¹, Rebekka V. Jensen², Alexander Perkins³, Matthew Dodd³, Tim Clayton³, Derek Yellon⁴, Derek J. Hausenloy^{4,5,6,7}, Hans Erik Bøtker^{1,2} on behalf of the CONDI-2/ERIC-PPCI investigators

Affiliations:

1. Department of Clinical Medicine, Cardiology, Aarhus University, Aarhus, Denmark
2. Department of Cardiology, Aarhus University Hospital, Aarhus, Denmark
3. London School of Hygiene and Tropical Medicine, Clinical Trials Unit, London, UK
4. The Hatter Cardiovascular Institute, University College London, London, UK
5. Cardiovascular & Metabolic Disorders Program, Duke-National University of Singapore Medical School, Singapore
6. National Heart Research Institute Singapore, National Hearts Centre, Singapore Yong Loo Lin School of Medicine, National University Singapore, Singapore
7. Cardiovascular Research Center, College of Medical and Health Sciences, Asia University, Taiwan

Address for correspondence:

Marie V. Hjortbak

Aarhus University, Department of Clinical Medicine

Palle Juul-Jensens Boulevard 82, DK-8200, Aarhus N, Denmark

Phone: 0045 78452262

Email: hjortbak@clin.au.dk

Abstract

Objectives: We studied the translational cardioprotective potential of P2Y₁₂ inhibitors against acute myocardial ischemia/reperfusion injury (IRI) in an animal model of acute myocardial infarction and in patients with ST-elevation myocardial infarction (STEMI) undergoing primary percutaneous coronary intervention (PPCI).

Background: P2Y₁₂ inhibitors have pleiotropic effects that may induce cardioprotection against acute myocardial IRI beyond their inhibitory effects on platelet aggregation.

Methods: We compared the cardioprotective effects of clopidogrel, prasugrel and ticagrelor on infarct size in an *in vivo* rat model of acute myocardial IRI, and investigated the effects of the P2Y₁₂ inhibitors on enzymatic infarct size (48-hour area-under-the-curve (AUC) troponin T release) and clinical outcomes in a retrospective study of STEMI patients from the CONDI-2/ERIC-PPCI trial using propensity score analyses.

Results: Loading with ticagrelor in rats reduced infarct size after acute myocardial IRI compared to controls (37±11% vs 52±8%, p<0.01), whereas clopidogrel and prasugrel did not (50±11%, p>0.99 and 49±9%, p>0.99, respectively). Correspondingly, troponin release was reduced in STEMI patients treated with ticagrelor compared to clopidogrel (adjusted 48-hour AUC ratio: 0.67, 95% CI 0.47-0.94). Compared to clopidogrel the composite endpoint of cardiac death or hospitalization for heart failure within 12 months was reduced in STEMI patients loaded with ticagrelor (HR 0.63; 95% CI 0.42-0.94) but not prasugrel (HR 0.84, 95% CI 0.43-1.63), prior to PPCI. Major adverse cardiovascular events did not differ between clopidogrel, ticagrelor or prasugrel.

Conclusions: The cardioprotective effects of ticagrelor in reducing infarct size may contribute to the clinical benefit observed in STEMI patients undergoing PPCI.

Key words: P2Y₁₂ inhibitor, cardioprotection, ischemic conditioning, myocardial infarction

Introduction

Acute myocardial infarction still contributes to mortality and morbidity worldwide. During myocardial infarction, the myocardium suffers ischemic damage, which can only be targeted by timely reperfusion therapy. The paradoxical myocardial reperfusion injury that may extend final infarct size [77] requires adjunctive treatment strategies beyond reperfusion to improve clinical outcome. Although remote ischemic conditioning (RIC) reduces myocardial injury by activating inherent cardioprotective mechanisms [34], verification of a clinical benefit for the patients has been challenging, mainly because clinical event rates with modern reperfusion therapy are low [27, 29, 43].

The cardiomyocyte has been the primary target of cardioprotective strategies given that final infarct size is the main predictor of cardiovascular mortality. However, increasing evidence shows that other targets might be of importance to attenuate injury during myocardial infarction. In addition to mediating the occlusive thrombus in acute myocardial infarction, platelets may also release factors that exacerbate acute myocardial ischemia and reperfusion injury [22, 79].

Loading treatment with P2Y₁₂ inhibitors is an established adjunctive therapy to invasive treatment of acute coronary syndrome because of their inhibitory effect on platelet aggregation. However, clopidogrel, prasugrel and ticagrelor have all demonstrated pleiotropic, cardioprotective effects in experimental studies [72, 74]. Observations from minor, retrospective studies indicate that the cardioprotective effects of P2Y₁₂ inhibitors may be transferrable to a clinical setting [36, 52].

The aims of the present study were to compare head-to-head loading with clopidogrel, prasugrel and ticagrelor on infarct size in an experimental rat model of myocardial ischemia and reperfusion, and subsequently study the translational potential in a cohort of STEMI patients from the CONDI-2/ERIC-PPCI trial [29].

Methods

Rat experiments

All animal experiments were performed in accordance with Danish legal and institutional guidelines (Authorization number: 2018-15-0201-01475). Male Sprague Dawley rats (Taconic, Ry, Denmark) (250-350 g) were randomized to one of the following protocols: 1) Control, 2) IPC, 3) RIC, 4) Clopidogrel, 5) Prasugrel, 6) Ticagrelor, 7) IPC+Ticagrelor or 8) RIC+Ticagrelor as specified in Figure S1. Combination therapy with ischemic conditioning and ticagrelor was investigated to determine interactions.

Delivery of P2Y₁₂ inhibitors

P2Y₁₂ inhibitors were administered by oral gavage using crushed tablets suspended in tab water; doses were adjusted to body weight of the individual rat. Clopidogrel (15 mg/kg) (Clopidogrel STADA, STADA Arzneimittel AG, Bad Vilbel, Germany) was given 4 hours prior to induction of myocardial ischemia, ticagrelor (20 mg/kg)(Brilique, AstraZeneca, Cambridge, United Kingdom) and prasugrel (10 mg/kg) (Efient, Daiichi-Sankyo Europe GmbH, Munich, Germany) were given 2 hours prior to induction of myocardial ischemia. Placebo treatment consisted of tab water only given 2 hours before myocardial ischemia.

The dosage and timing of P2Y₁₂ inhibitors were chosen from available data in the literature. Clopidogrel is a prodrug that requires enzymatic activation. The loading dose of clopidogrel must be given before reperfusion of the myocardium, but the duration of pretreatment to induce protection varies between 4 hours and two days in animal studies [66, 74]. In the present study we loaded the animals with clopidogrel 4 hours prior to induction of ischemia because the resultant plasma concentration is associated with antiplatelet efficacy [56, 66] and because the approach may have some potential for clinical translation when given before reperfusion. The dose of clopidogrel

was based on previous studies demonstrating cardioprotective effect of clopidogrel [66, 76]. Ticagrelor and prasugrel have more rapid and potent antiplatelet responses than clopidogrel. In rats, platelet aggregation is significantly inhibited one to two hours after administration of ticagrelor or prasugrel, whereas clopidogrel may require 2-4 hours [56, 57]. This pharmacologic profile may increase the cardioprotective potential of ticagrelor and prasugrel within a clinically relevant timeframe for STEMI patients. As for clopidogrel, the doses of prasugrel [25, 58] and ticagrelor [3, 66, 72, 76] were based on previous studies demonstrating cardioprotective effect.

In vivo myocardial infarction

The rats were anesthetized with an intraperitoneal injection of pentobarbiturate (100 mg/kg body weight) (Skanderborg Pharmacy, Skanderborg, Denmark). Immediately after anesthesia was achieved the rats were intubated, connected to a ventilator (UGO BASILE, Comerio, Varese, Italy), and ventilated with atmospheric air. Body temperature was maintained at 37 °C (± 0.5 °C) (CMA/150, CMA Microdialyses AB, Krista, Sweden). The heart was accessed through a left sided thoracotomy. The left anterior descending artery (LAD) was identified and ligated with a 4-0 silk suture (Sofsilk™, Covidien, Dublin, Ireland) at the level of the left atrial appendix tip. All hearts received 30 minutes of myocardial ischemia followed by 2 hours of reperfusion.

RIC was performed prior to the thoracotomy using a tourniquet around a hind leg, to induce 3 cycles of 5 minutes limb ischemia followed by 5 minutes of reperfusion. IPC was performed after the thoracotomy, using the myocardial suture around LAD to induce 3 cycles of 5 minutes of ischemia followed by 5 minutes of reperfusion.

Infarct size

After 2 hours of reperfusion, the LAD was reoccluded, and a 2 % solution of Evans Blue (Sigma-Aldrich, St. Louis, MO, USA) was injected in the inferior vena cava to visualize the area at risk. The hearts were rapidly removed and stored at -80 °C. The hearts were then sliced and stained using a 1% solution of Triphenyl Tetrazolium Chloride (Sigma-Aldrich, St Louis, MO, USA). After 24 hours in 4% formalin buffer (VWR International, Leuven, Belgium), the slices were scanned using a flatbed scanner (Epson Perfection V600 Photo scanner, Epson, Nagano, Japan). The infarct size, area at risk and area of the left ventricle were assessed using ImageJ software (NIH, Bethesda, Maryland, USA). All measurements were correlated to the wet weight of the individual slice. Final infarct size is expressed as the percent of infarcted area over the area at risk.

Statistical analyses

Statistical analyses of the rat experiments were performed using GraphPad Prism 8.2.0 (GraphPad Software, California, USA). Data are presented as mean \pm SD. One-way ANOVA with post hoc Bonferroni correction for multiple comparisons was used for all rat experimental data [13]. Sample size calculations were based on an infarct size of 50% in controls and 35% in intervention groups, with a standard deviation of 10%. A significance level $\alpha=0.05$ and a power of 95% yielded a sample size of 12 animals in each group.

We tested for interaction between type of intervention (none, IPC, and RIC) and ticagrelor on infarct size. The interaction analysis was performed in StataIC version 16 (Stata Corp, College Station, Texas, USA).

Clinical studies

The clinical part of the study was designed as a retrospective, non-prespecified post hoc sub-study of the international, multicenter, single-blind, randomized controlled CONDI-2/ERIC-PPCI trial [29]. A detailed description of the study is provided in the original publication [29]. Patients with ST-segment elevation myocardial infarction, eligible to PPCI, were randomized to standard treatment or treatment with RIC. The study included patients from 33 centers across United Kingdom, Denmark, Spain and Serbia. We analyzed the data collected for the CONDI-2/ERIC-PPC trial to investigate interaction between treatment with P2Y₁₂ receptor inhibitors and RIC in relation to PPCI for clinical outcomes.

In accordance with contemporary guidelines, patients with STEMI were loaded with a P2Y₁₂ receptor inhibitor prior to PPCI. Patients received either clopidogrel (600mg), ticagrelor (180 mg) or prasugrel (60 mg). Choice of P2Y₁₂ receptor inhibitor for loading was based on current guidelines and regional preferences. The time from administration of the chosen P2Y₁₂ inhibitor to reperfusion by PPCI was not registered.

Patient Selection

We excluded patients, who were on treatment with clopidogrel, ticagrelor, or prasugrel prior to PPCI. Patients, who were not treated with either peri-procedural clopidogrel, ticagrelor or prasugrel, were also excluded.

Infarct size

We estimated myocardial infarct size measured as area-under-the-curve (AUC) of high-sensitivity troponin T measured between 0 and 48 hours after PPCI in a subset of patients.

Clinical outcomes

The main endpoint was a composite of cardiac death or hospitalization for heart failure at 12 months. Secondary endpoints included cardiac death, hospitalization for heart failure, major cardiovascular adverse events (MACE; a composite of all-cause death, reinfarction, coronary revascularization, and stroke), myocardial infarction, stroke, revascularization, and all-cause death. A blinded independent endpoint committee reviewed all events. A detailed description of endpoint definitions has been published elsewhere [29].

Statistical analysis

Patients were stratified according to peri-procedural treatment with clopidogrel, ticagrelor or prasugrel. We used propensity score based-methods to estimate the average treatment effect of ticagrelor or prasugrel compared to clopidogrel [55].

For the infarct size calculations, we estimated 48-hour troponin T AUC for subsets of patients using multiple imputation by chained equations in case of missing data. We log-transformed AUC since distributions were skewed, and computed the AUC ratio by linear regression. AUC ratios were calculated in the propensity score cohorts characterized below. In the main analysis, we compared clopidogrel vs. ticagrelor vs. prasugrel in a combined analysis. For the sensitivity analyses, comparisons between clopidogrel vs. ticagrelor and clopidogrel vs. prasugrel were analyzed separately because the number of patients were higher than in the combined analysis.

Covariates associated with both the outcome and exposure or only the outcome were included to estimate the propensity score: age (continuous variable), sex, body mass index (<18.5 kg/m², 18.5-24.9 kg/m², 25-29.9 kg/m², ≥30 kg/m²), active smoking, hypertension, previous myocardial infarction, hypercholesterolemia, diabetes, first medical contact to balloon time (<60 minutes, 60-119 minutes, 120-179 minutes, ≥180 minutes) [59], multivessel disease, LAD stenosis, Killip class,

Thrombolysis In Myocardial Infarction Flow Grade, periprocedural heparin, and country [15]. The original CONDI-2/ERIC-PPCI trial analyses showed no interaction between treatment with ticagrelor and RIC [29]. A total of 17.3% of patients had missing values in ≥ 1 of the covariates included in the propensity score. Missing values were handled through multiple imputations using chained equations, generating 20 imputations. We used multinomial logistic regression to estimate the propensity of type of P2Y₁₂ receptor inhibitor. A Cox regression was used to estimate crude and stabilized inverse-probability-weighted (IPW) hazard ratios (HRs) using clopidogrel as reference [16, 30]. The proportional hazards assumption was evaluated by log–log plots, and found to be satisfied. Twelve-month cumulative incidence proportion was estimated, accounting for the competing risk of all-cause death, except in the case of MACE and all-cause death. Twelve-month cumulative incidence curves of the main outcome and MACE were constructed. We also estimated the 30-day risk of the main outcome and MACE.

We performed two sensitivity analyses. First, a ‘full cohort’ analysis in which all patients received ticagrelor, prasugrel or clopidogrel in relation to PPCI, including patients who were not eligible in propensity score based-analyses. We estimated adjusted HRs by multivariable Cox regression. We adjusted for the same covariates used for the propensity score. Second, a propensity-score based analysis in which we analyzed the data in two separate analyses, one comparing clopidogrel and ticagrelor, and one comparing clopidogrel and prasugrel. In the separated analyses all Spanish patients were excluded due to structural non-positivity, since all Spanish patients were treated with clopidogrel [30]. For the same reason all patients from Serbia were excluded from the analysis of prasugrel vs clopidogrel, since no patients in Serbia received prasugrel. To improve balance in distribution of propensity scores in the treatment groups, patients with a propensity score <0.1 and >0.9 were excluded [20].

All statistical analyses of clinical data were performed using StataIC version 16 (Stata Corp, College Station, Texas, USA).

Results

Animal experiments – Infarct size

IPC significantly reduced infarct size compared to controls ($26\pm 12\%$ vs $52\pm 8\%$, $p<0.0001$) (Figure 1). RIC also reduced infarct size compared to controls, but not to the same degree as IPC (41 ± 11 vs $52\pm 8\%$, $p<0.05$).

Ticagrelor reduced infarct size compared to controls ($37\pm 11\%$ vs $52\pm 8\%$, $p<0.01$). Clopidogrel or prasugrel did not affect infarct size ($50\pm 11\%$, $p>0.99$ and $49\pm 9\%$, $p>0.99$, respectively).

Combination therapy with IPC and ticagrelor resulted in a reduction in infarct size compared to controls ($25\pm 9\%$ vs $52\pm 8\%$, $p<0.0001$). The reduction in infarct size was similar to IPC treatment alone ($p>0.99$), suggesting no additive cardioprotective effect with the combination of IPC and ticagrelor.

The reduction in infarct size from combination therapy with RIC and ticagrelor was similar treatment with RIC alone (42 ± 13 , $p>0.99$), but the reduction only reached borderline statistical significance when compared to controls ($p=0.08$). Again, there was no additive cardioprotective effect with the combination of RIC and ticagrelor.

Interaction analyses of infarct size showed interaction between ticagrelor treatment and IPC ($p<0.05$) and RIC ($p<0.05$) (Table S1).

Infarct size related to left ventricle showed the same results as infarct size related to area at risk. With an average of 40% of left ventricle, area at risk did not differ between any of the intervention groups and controls.

Clinical study – infarct size

In our combined analysis, the 48-hour AUC of troponin release was reduced in patients treated with ticagrelor compared to clopidogrel (Adjusted AUC ratio: 0.67, 95% CI 0.47-0.94) (Table S2). The number of prasugrel treated patients with troponin data (n=5) did not allow sufficient statistical power to provide valid results (Table S2). The supplementary sensitivity analysis, where AUC troponin release was compared separately as clopidogrel vs. ticagrelor and clopidogrel vs. prasugrel, showed no significant reduction in troponin release from either ticagrelor or prasugrel (Table S3).

Clinical study - outcome

Out of 5115 patients included in the original CONDI-2/ERIC-PPCI study, we included a total of 1754 patients in the retrospective main analysis (Figure 2). Of these 395 patients received clopidogrel, 1210 patients received ticagrelor and 149 received prasugrel. The number of patients differs between the groups, as patients are included based on the propensity scores. Baseline characteristics of the patients included in the analysis are shown in table 1. All three P2Y₁₂ inhibitors were only prescribed in the UK. Thus, only UK patients were ultimately included in the multinomial logistic regression analysis. Baseline characteristics were generally well balanced. Patients with previous myocardial infarction were slightly more prevalent in the groups treated with ticagrelor (37.6%) and prasugrel (41.6%) compared to clopidogrel (25.6%). Nitrates were used more often in patients treated with clopidogrel (89.4%) and prasugrel (92.6%) compared to ticagrelor (81.3%), which may be due to regional differences in medication strategy.

The main composite outcome of one-year risk of cardiac death or hospitalization for heart failure occurred in 9.6% of the clopidogrel treated patients, compared to 6.5% in the ticagrelor treated patients (HR 0.63; 95% CI 0.42-0.94) and 8.1% in the prasugrel treated patients (HR 0.84,

95% CI 0.43-1.63) (Table 2) with the time course specified in Figure 3. In analyses of the individual components of the composite primary endpoint ticagrelor reduced the risk to a similar extent, but not all with statistical significance: one-year risk of hospitalization for heart failure (HR 0.57, 95% CI 0.36-0.91), cardiac death (HR 0.78, 95% CI 0.38-1.58) and all-cause death (HR 0.59, 95% CI 0.35-1.00) (Table S4). The individual components of the primary endpoint were not affected by prasugrel: one-year risk of hospitalization for heart failure (HR 0.98, 95% CI 0.48-1.99), cardiac death (HR 0.28, 95% CI 0.04-2.19) and all-cause death (HR 0.24, 95% CI 0.06-1.04) (Table S4). The one-year risk of MACE was not affected by ticagrelor (HR 0.86, 95% CI 0.56-1.32) or prasugrel (HR 0.54, 95% CI 0.24-1.23) compared to clopidogrel (Table 2 and Figure 3). Reinfarction, stroke or revascularization did not differ between groups (Table S4).

The thirty-day risk of cardiac death or hospitalization for heart failure was reduced in patients receiving ticagrelor (HR 0.64, 95% CI 0.42-0.98), but not prasugrel (HR 0.80, 95% CI 0.39-1.65) compared to clopidogrel (Figure 3 and Table S5). No reduction in MACE was found at 30 days.

The sensitivity analyses of the composite clinical endpoint yielded results consistent with the main analysis (Table S6-S10, Figure S2 and S3).

Discussion

The results of our study demonstrate that ticagrelor but not clopidogrel or prasugrel decreased infarct size in an *in vivo* rat model of ischemia reperfusion injury. Correspondingly, ticagrelor seemed to reduce infarct size as measured by troponin release in the clinical setting in a post-hoc sub-study. The results translated into a beneficial effect of ticagrelor pre-treatment in terms of a reduced incidence of a composite endpoint including cardiac death and hospitalization for heart failure with contribution of each component.

Effect of P2Y₁₂ inhibitors on infarct size

Beyond the documented beneficial antithrombotic effects on myocardial damage [19, 41, 42, 50, 53, 67, 70, 78], experimental studies have shown that second and third generation P2Y₁₂ inhibitors may be capable of reducing infarct size in experimental settings, but their cardioprotective capacity appears variable [9, 25, 66, 72, 74, 76]. Translation into a potential clinical effect was already demonstrated for the second generation P2Y₁₂ inhibitor, clopidogrel, but appears to vary as well [24, 51].

Our results confirm that ticagrelor reduces infarct size in experimental models of ischemia-reperfusion injury [9, 22, 48, 79]. Cardioprotection can be obtained by a single dose given only two hours before myocardial infarction [3, 72, 76], which potentially increases clinical translation because cangrelor, an intravenously administered equivalent to ticagrelor, has cardioprotective effects when given just before reperfusion [75]. The mechanisms behind the ticagrelor-induced cardioprotection seem not solely related to the inhibition of platelet aggregation, but also to pleiotropic effects [22].

Among the orally administered P2Y₁₂ inhibitors, ticagrelor has most convincingly demonstrated infarct size reduction in STEMI patients undergoing rapid revascularization using

measurement of troponin release [47, 52] and more reliably by magnetic resonance imaging [37, 52]. Although our statistical analyses are not completely consistent due to our retrospective design and suboptimal statistical power, our data do not dispute that the beneficial clinical outcome with ticagrelor was associated with reduced infarct size, measured by troponin release in patients. We acknowledge that the measurement of infarct size by circulating biomarkers should be interpreted with caution. Our main troponin analysis relies on 260 patients. Only 5 patients in the analyses received prasugrel, such that a valid estimate was not obtainable. The sensitivity analyses of troponin release in a pairwise comparison between clopidogrel and ticagrelor in 503 patients did not confirm the results of the main analysis. When infarct sizes are minor, i.e. in the order of magnitude of 16% of the left ventricle, as obtained by modern reperfusion therapy [14], the sensitivity of circulating biomarkers may not be optimal.

Experimental studies of the cardioprotective effect of pretreatment with prasugrel are limited and with varying results [9, 25, 44]. Despite three days of pretreatment with prasugrel, Birnbaum *et al* did not show infarct reduction after coronary occlusion [9], whereas Dost *et al* demonstrated that a single dose prasugrel reduced infarct size [25]. We found no reduction in infarct size by a single dose of prasugrel, although our experimental setup seemed similar to the approach used by Dost *et al*. in terms of dosing, timing and ischemia/reperfusion protocol. The use of two different rat strains may explain the discrepancy as sensitivity to ischemia and reperfusion injury is known to vary between rat strains [5].

Effect of P2Y₁₂ inhibitors on clinical outcome

In accordance with The Ticagrelor Therapy in STEMI Patients Planned for Percutaneous Coronary Intervention (ATLANTIC) trial [45], we observed no reduction in MACE, potentially reflecting that the benefit of ticagrelor is not caused only by a more efficient long-term platelet

inhibition than with clopidogrel. A statistically significant improvement by prasugrel compared to clopidogrel treatment was not evident from our main endpoint. Consistent with our results a prespecified substudy of the ISAR REACT 5 trial [53] in STEMI-patients demonstrated no significant difference in the primary endpoint (incidence of death, myocardial infarction, or stroke at 1 year after randomization) between prasugrel and ticagrelor [4]. The endpoint in the ISAR REACT 5 study mainly relates to the antithrombotic effect of the P2Y₁₂ inhibitors. We observed a reduction of cardiac death or hospitalization for heart failure by ticagrelor that emerged early compared to clopidogrel but compared to prasugrel most clearly after 180 days of follow-up (Figure 3a and b). The early effect may reflect a superior antithrombotic efficacy of prasugrel and ticagrelor compared to clopidogrel, whereas infarct size reduction by ticagrelor becomes evident with a delay when inappropriate remodeling due to a significant MI size translates into clinical symptoms.

Mechanistic considerations

In observational post hoc analyses, the effect of the third generation P2Y₁₂ inhibitors, ticagrelor and prasugrel, versus the second-generation inhibitor, clopidogrel, on microvascular obstruction is equivocal [36, 65]. Ticagrelor does not seem to be superior to prasugrel in reducing microvascular obstruction [62, 64]. Although experimental data suggest that P2Y₁₂-receptor inhibition using cangrelor at the onset of reperfusion can itself reduce MI size [75], it is unclear whether the cardioprotective effect is mediated on the coronary vasculature or the cardiomyocyte [28, 36, 37]. Ticagrelor increases circulating levels of adenosine in humans mainly at doses higher than standard [61]. Still, increased serum concentration of adenosine seems to be responsible for ticagrelor-related adverse effects, including dyspnea, ventricular pauses, and bradyarrhythmias. Moreover, experimental as well as human studies suggest that ticagrelor enhances the biological effects of endogenous adenosine [63, 69], implying that adenosine may serve as a mediator of some

of the pleiotropic cardioprotective effect [1, 61, 69]. Ticagrelor has a favorable effect on endothelial function after ischemia and reperfusion compared to clopidogrel in humans [68]. The A_{2A} receptor is the main adenosine receptor responsible for coronary vasodilation, mediated by both nitric oxide-dependent and -independent pathways [46]. Adenosine may also act cardioprotective by inhibiting neutrophil trafficking, granule release, and production of reactive oxygen species and inflammatory mediators [6, 21, 49].

Studies of cardioprotection by P2Y₁₂ inhibitors imply that activated platelets are involved although interference with platelet aggregation itself [18, 48, 73] or improved early coronary reperfusion [24] may not be the main targets. Despite faster P2Y₁₂ inhibition by cangrelor, compared to ticagrelor, this does not necessarily induce an increased salvage of myocardium [60]. Near-obliteration of circulating platelets either with cell poison or an antibody abrogates the cardioprotective effect of P2Y₁₂ antagonists [18]. Furthermore, P2Y₁₂ antagonists have no effect in isolated hearts perfused with platelet-free buffer [18, 73]. Platelets may be a target of P2Y₁₂ antagonists for creation of a cardioprotective effect [7]. However, it is unknown whether activated platelets release substances with protective effects on the endothelium and how events between binding of the P2Y₁₂ blocker to its platelet receptor relates to emergence of cardioprotection.

Platelet reactivity declines relatively slowly after oral administration of P2Y₁₂ inhibitors and requires several hours before reaching full effect. The profile is most favorable for oral ticagrelor or prasugrel administration as manifestation of the antiplatelet activity within 1-3 hours [2, 8, 10, 26] is less than for clopidogrel for which the effect initiates after 6-11 hours [8]. It remains unknown whether the timing of P2Y₁₂ inhibitors in the CONDI2-PPCI trial was optimal. In the original trial we only had access to data on the type of P2Y₁₂ inhibitor treatment given in relation to the PPCI and not the timing of the administration or the treatment strategy after the index event. Nonetheless, ticagrelor improved the main endpoint indicating that the effect was sufficient.

Interaction of P2Y₁₂ inhibitor treatment and ischemic conditioning

In preclinical studies, ischemic pre- and postconditioning conditioning interact with the cardioprotective effect of the P2Y₁₂ inhibitor cangrelor, with no additional effect of combination therapy of P2Y₁₂ inhibition and ischemic conditioning [72, 74]. We found a similar interaction with no additive effect between ticagrelor treatment and both IPC and RIC in our experimental data and extended the knowledge of an interaction between ischemic conditioning and P2Y₁₂ inhibitors to include RIC. Combination treatment with ticagrelor and RIC did not significantly reduce infarct size. Whether this is solely related to the larger variation in the data or whether other factors are responsible is not known. In our clinical trial, we observed no interaction between RIC and ticagrelor treatment due to the lack of effect by RIC on the main endpoints [29]. Eventually, a potential interaction may be explained by two different even opposing mechanisms, which are almost undifferentiable: 1. There is a potential recruitment of protection by the patient medications and 2. Protection can be attenuated/abrogated by the medication and in parallel [38]. These two mechanisms may interfere in different ways with IPC and RIC, since the underlying signal transduction of IPC and RIC may also differ [32, 39]. In an experimental setting, IPC improved the recovery of coronary flow and LVDP during reperfusion whereas RIC only impacted on the recovery of LVDP [40], indicating that IPC may exert stronger protective effects on the coronary vasculature than RIC [31]. While the near maximum efficacy of IPC and RIC seems to have been reached in our experimental setting, a further protection potential by an intensified stimulus [35, 40, 54] may be present in the clinical setting [71]. Deployment of the full protection capacity seems to be necessary to uncover an interaction with pharmacological treatment and should be taken into consideration when applying multitarget strategies.

Clinical implications

The signal indicating that ticagrelor has the most potent cardioprotective capacity among currently recommended oral P2Y₁₂ inhibitors seems to contribute to the improvement in clinical outcome by modern standard care of patients with STEMI. The achieved improvement in clinical outcome may challenge the ability to document adjunctive cardioprotective treatments beyond optimized standard care in future studies. We have realized this challenge when changing from clopidogrel to ticagrelor in our previous clinical studies of RIC [14, 29]. Careful selection of high-risk patients [11, 12] and multitarget cardioprotective strategies may increase the protective potential [23, 33]. Since intravenous cangrelor may have similar cardioprotective efficacy as ticagrelor, an alternative treatment strategy might be intravenous cangrelor infusion shortly prior to stenting followed by subsequent post-PCI transition to an oral agent [17].

Study limitations

To investigate the cardioprotective capacity of P2Y₁₂ inhibitors we chose an *in vivo* rat model, as the P2Y₁₂ inhibitors necessitate *in vivo* metabolization of the drugs and presence of platelets. Since IPC was induced by an invasive procedure, the experimental design was limited by the prolonged surgical procedure in these groups. Our unpublished pilot trials showed no impact on infarct size from a prolonged surgical procedure in control animals. As IPC and RIC may have different signaling profiles, we cannot exclude an effect of the timing between IPC or RIC and index ischemia. Doses of P2Y₁₂ inhibitors relied on results from other laboratories [25, 44, 66, 72, 74, 76], so we did not conduct dose-response experiments and it cannot be excluded that other doses of P2Y₁₂ inhibitors might increase the cardioprotective effects.

The main limitation of the clinical part of the study is that the patients in the original CONDI-2/ERIC-PPCI trial were not randomized by P2Y₁₂ inhibitor prescription. To reduce confounding, our statistical analyses rely on propensity scored analyses. To balance the distribution

of propensity score between treatment groups, we excluded both low and high propensity patients, hereby reducing the sample size. Also, relatively few patients in our cohort were treated with prasugrel, so estimates of clinical outcome in prasugrel treated patients are with higher statistical uncertainty. The sensitivity analyses with pairwise comparison of ticagrelor and clopidogrel and ticagrelor and prasugrel had larger cohorts. Consequently, the statistical power was increased. The sensitivity analyses confirmed the results of the main analysis for the primary clinical outcome but not infarct size data. Moreover, residual confounding may still be present, so our results should be considered exploratory and needs confirmation in a randomized trial.

Conclusion

Pre-treatment with ticagrelor reduced infarct size in rats after ischemia and reperfusion injury, whereas clopidogrel or prasugrel did not. In patients suffering from STEMI and treated with PPCI, we found that treatment with ticagrelor, but not prasugrel, reduced cardiac death and hospitalization for heart failure compared to treatment with clopidogrel. The improved clinical outcome with ticagrelor may be caused by pleiotropic effects that attenuate ischemia and reperfusion injury.

Declarations

Funding Sources

The ERIC-PPCI trial was funded by a British Heart Foundation Clinical Study Grant (CS/14/3/31002) and a University College London Hospital/University College London Biomedical Research Clinical Research grant. The CONDI-2 trial was funded by Danish Innovation Foundation grants (11-108354 and 11-115818), Novo Nordisk Foundation (NNF13OC0007447), and Trygffonden (109624). DJH was supported by the British Heart Foundation (FS/10/039/28270), Duke-National University Singapore Medical School, Singapore Ministry of Health's National Medical Research Council under its Clinician Scientist-Senior Investigator scheme (NMRC/CSA-SI/0011/2017) and its Collaborative Centre Grant scheme (NMRC/CGAug16C006). This article is based upon the work of COST Action EU-CARDIOPROTECTION (CA16225) and supported by COST (European Cooperation in Science and Technology).

Conflicts of interest

The authors declare no conflicts of interest.

Ethics approval

The ethical approval of the animal experiments of the study was approved by the Danish Veterinary and Food Administration (Authorization number: 2018-15-0201-01475). The clinical part of the study was authorized on ClinicalTrials.gov as NCT02342522.

Availability of data and material

Data can be made available if the manuscript is accepted for publication.

Code availability

Not applicable.

Author contributions

All authors have participated in parts of the conception and design of the study or analysis and interpretation of data.

Authors responsible for the animal experiments: MV Hjortbak, JM Seefeldt, TR Lassen, RV Jensen and HE Bøtker.

Authors responsible for the clinical analyses: KKW Olesen, MV Hjortbak, A Perkins, M Dodd, T Clayton, D Yellon, DJ Hausenloy and HE Bøtker.

All authors have contributed to drafting the manuscript, and all have read and approved the manuscript.

References

1. Alexopoulos D, Moulias A, Koutsogiannis N, Xanthopoulou I, Kakkavas A, Mavronasiou E, Davlouros P, Hahalis G (2013) Differential effect of ticagrelor versus prasugrel on coronary blood flow velocity in patients with non-ST-elevation acute coronary syndrome undergoing percutaneous coronary intervention: an exploratory study. *Circ Cardiovasc Interv* 6:277-283 doi:10.1161/CIRCINTERVENTIONS.113.000293
2. Alexopoulos D, Xanthopoulou I, Gkizas V, Kassimis G, Theodoropoulos KC, Makris G, Koutsogiannis N, Damelou A, Tsigkas G, Davlouros P, Hahalis G (2012) Randomized assessment of ticagrelor versus prasugrel antiplatelet effects in patients with ST-segment-elevation myocardial infarction. *Circ Cardiovasc Interv* 5:797-804 doi:10.1161/CIRCINTERVENTIONS.112.972323
3. Audia JP, Yang XM, Crockett ES, Housley N, Haq EU, O'Donnell K, Cohen MV, Downey JM, Alvarez DF (2018) Caspase-1 inhibition by VX-765 administered at reperfusion in P2Y12 receptor antagonist-treated rats provides long-term reduction in myocardial infarct size and preservation of ventricular function. *Basic Res Cardiol* 113:32 doi:10.1007/s00395-018-0692-z
4. AYTEKIN A, Ndrepepa G, Neumann FJ, Menichelli M, Mayer K, Wohrle J, Bernlochner I, Lahu S, Richardt G, Witzenbichler B, Sibbing D, Cassese S, Angiolillo DJ, Valina C, Kufner S, Liebetrau C, Hamm CW, Xhepa E, Hapfelmeier A, Sager HB, Wustrow I, Joner M, Trenk D, Fusaro M, Laugwitz KL, Schunkert H, Schupke S, Kastrati A (2020) Ticagrelor or Prasugrel in Patients With ST-Segment-Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. *Circulation* 142:2329-2337 doi:10.1161/CIRCULATIONAHA.120.050244

5. Baker JE, Konorev EA, Gross GJ, Chilian WM, Jacob HJ (2000) Resistance to myocardial ischemia in five rat strains: is there a genetic component of cardioprotection? *Am J Physiol Heart Circ Physiol* 278:H1395-1400 doi:10.1152/ajpheart.2000.278.4.H1395
6. Barletta KE, Ley K, Mehrad B (2012) Regulation of neutrophil function by adenosine. *Arterioscler Thromb Vasc Biol* 32:856-864 doi:10.1161/ATVBAHA.111.226845
7. Barrabes JA, Inseste J, Mirabet M, Quiroga A, Hernando V, Figueras J, Garcia-Dorado D (2010) Antagonism of P2Y₁₂ or GPIIb/IIIa receptors reduces platelet-mediated myocardial injury after ischaemia and reperfusion in isolated rat hearts. *Thromb Haemost* 104:128-135 doi:10.1160/TH09-07-0440
8. Bergmeijer TO, Godschalk TC, Janssen PWA, Berge KVD, Breet NJ, Kelder JC, Hackeng CM, Ten Berg JM (2017) How Long Does It Take for Clopidogrel and Ticagrelor to Inhibit Platelets in Patients Undergoing Primary Percutaneous Coronary Intervention? A Detailed Pharmacodynamic Analysis: Time Course of Platelet Reactivity in STEMI (TOPS). *Semin Thromb Hemost* 43:439-446 doi:10.1055/s-0037-1599156
9. Birnbaum Y, Birnbaum GD, Birnbaum I, Nylander S, Ye Y (2016) Ticagrelor and Rosuvastatin Have Additive Cardioprotective Effects via Adenosine. *Cardiovasc Drugs Ther* 30:539-550 doi:10.1007/s10557-016-6701-2
10. Bonello L, Laine M, Camoin-Jau L, Noirot F, Guieu R, Dignat-George F, Paganelli F, Frere C (2015) Onset of optimal P2Y₁₂-ADP receptor blockade after ticagrelor and prasugrel intake in Non-ST elevation acute coronary syndrome. *Thromb Haemost* 114:702-707 doi:10.1160/TH15-02-0149
11. Botker HE (2020) The Future of Cardioprotection-Pointing Toward Patients at Elevated Risk as the Target Populations. *J Cardiovasc Pharmacol Ther* 25:487-493 doi:10.1177/1074248420937871

12. Botker HE (2021) Searching myocardial rescue through intermittent upper arm occlusion and lizard saliva. *Basic Res Cardiol* 116:5 doi:10.1007/s00395-021-00843-1
13. Botker HE, Hausenloy D, Andreadou I, Antonucci S, Boengler K, Davidson SM, Deshwal S, Devaux Y, Di Lisa F, Di Sante M, Efentakis P, Femmino S, Garcia-Dorado D, Giricz Z, Ibanez B, Iliodromitis E, Kaludercic N, Kleinbongard P, Neuhauser M, Ovize M, Pagliaro P, Rahbek-Schmidt M, Ruiz-Meana M, Schluter KD, Schulz R, Skyschally A, Wilder C, Yellon DM, Ferdinandy P, Heusch G (2018) Practical guidelines for rigor and reproducibility in preclinical and clinical studies on cardioprotection. *Basic Res Cardiol* 113:39 doi:10.1007/s00395-018-0696-8
14. Botker HE, Kharbanda R, Schmidt MR, Bottcher M, Kaltoft AK, Terkelsen CJ, Munk K, Andersen NH, Hansen TM, Trautner S, Lassen JF, Christiansen EH, Krusell LR, Kristensen SD, Thuesen L, Nielsen SS, Rehling M, Sorensen HT, Redington AN, Nielsen TT (2010) Remote ischaemic conditioning before hospital admission, as a complement to angioplasty, and effect on myocardial salvage in patients with acute myocardial infarction: a randomised trial. *Lancet* 375:727-734 doi:10.1016/S0140-6736(09)62001-8
15. Brookhart MA, Schneeweiss S, Rothman KJ, Glynn RJ, Avorn J, Sturmer T (2006) Variable selection for propensity score models. *Am J Epidemiol* 163:1149-1156 doi:10.1093/aje/kwj149
16. Brookhart MA, Wyss R, Layton JB, Sturmer T (2013) Propensity score methods for confounding control in nonexperimental research. *Circ Cardiovasc Qual Outcomes* 6:604-611 doi:10.1161/CIRCOUTCOMES.113.000359
17. Cohen MV, Downey JM (2020) What Are Optimal P2Y12 Inhibitor and Schedule of Administration in Patients With Acute Coronary Syndrome? *J Cardiovasc Pharmacol Ther* 25:121-130 doi:10.1177/1074248419882923

18. Cohen MV, Yang XM, White J, Yellon DM, Bell RM, Downey JM (2016) Cangrelor-Mediated Cardioprotection Requires Platelets and Sphingosine Phosphorylation. *Cardiovasc Drugs Ther* 30:229-232 doi:10.1007/s10557-015-6633-2
19. Committee CS (1996) A randomised, blinded, trial of clopidogrel versus aspirin in patients at risk of ischaemic events (CAPRIE). CAPRIE Steering Committee. *Lancet* 348:1329-1339 doi:10.1016/s0140-6736(96)09457-3
20. Crump RK, Hotz VJ, Imbens GW, Mitnik OA (2009) Dealing with limited overlap in estimation of average treatment effects. *Biometrika* 96:187-199 doi:10.1093/biomet/asn055
21. Csoka B, Nemeth ZH, Rosenberger P, Eltzschig HK, Spolarics Z, Pacher P, Selmeczy Z, Koscsó B, Himer L, Vizi ES, Blackburn MR, Deitch EA, Haskó G (2010) A2B adenosine receptors protect against sepsis-induced mortality by dampening excessive inflammation. *J Immunol* 185:542-550 doi:10.4049/jimmunol.0901295
22. Davidson SM, Andreadou I, Barile L, Birnbaum Y, Cabrera-Fuentes HA, Cohen MV, Downey JM, Girao H, Pagliaro P, Penna C, Pernow J, Preissner KT, Ferdinandy P (2019) Circulating blood cells and extracellular vesicles in acute cardioprotection. *Cardiovasc Res* 115:1156-1166 doi:10.1093/cvr/cvy314
23. Davidson SM, Ferdinandy P, Andreadou I, Botker HE, Heusch G, Ibanez B, Ovize M, Schulz R, Yellon DM, Hausenloy DJ, Garcia-Dorado D, Action CC (2019) Multitarget Strategies to Reduce Myocardial Ischemia/Reperfusion Injury: JACC Review Topic of the Week. *J Am Coll Cardiol* 73:89-99 doi:10.1016/j.jacc.2018.09.086
24. De Backer O, Ratcovich H, Biasco L, Pedersen F, Helqvist S, Saunamaki K, Tilsted HH, Clemmensen P, Olivecrona G, Kelbaek H, Jorgensen E, Engstrom T, Holmvang L (2015) Prehospital administration of P2Y12 inhibitors and early coronary reperfusion in primary

PCI: an observational comparative study. *Thromb Haemost* 114:623-631

doi:10.1160/TH15-01-0026

25. Dost T (2020) Cardioprotective properties of the platelet P2Y₁₂ receptor inhibitor prasugrel on cardiac ischemia/reperfusion injury. *Pharmacol Rep* 72:672-679 doi:10.1007/s43440-019-00046-5
26. Franchi F, Rollini F, Rivas A, Wali M, Briceno M, Agarwal M, Shaikh Z, Nawaz A, Silva G, Been L, Smairat R, Kaufman M, Pineda AM, Suryadevara S, Soffer D, Zenni MM, Bass TA, Angiolillo DJ (2019) Platelet Inhibition With Cangrelor and Crushed Ticagrelor in Patients With ST-Segment-Elevation Myocardial Infarction Undergoing Primary Percutaneous Coronary Intervention. *Circulation* 139:1661-1670 doi:10.1161/CIRCULATIONAHA.118.038317
27. Hausenloy DJ, Candilio L, Evans R, Ariti C, Jenkins DP, Kolvekar S, Knight R, Kunst G, Laing C, Nicholas J, Pepper J, Robertson S, Xenou M, Clayton T, Yellon DM, Investigators ET (2015) Remote Ischemic Preconditioning and Outcomes of Cardiac Surgery. *N Engl J Med* 373:1408-1417 doi:10.1056/NEJMoa1413534
28. Hausenloy DJ, Chilian W, Crea F, Davidson SM, Ferdinandy P, Garcia-Dorado D, van Royen N, Schulz R, Heusch G (2019) The coronary circulation in acute myocardial ischaemia/reperfusion injury: a target for cardioprotection. *Cardiovasc Res* 115:1143-1155 doi:10.1093/cvr/cvy286
29. Hausenloy DJ, Kharbanda RK, Moller UK, Ramlall M, Aaroe J, Butler R, Bulluck H, Clayton T, Dana A, Dodd M, Engstrom T, Evans R, Lassen JF, Christensen EF, Garcia-Ruiz JM, Gorog DA, Hjort J, Houghton RF, Ibanez B, Knight R, Lippert FK, Lonborg JT, Maeng M, Milasinovic D, More R, Nicholas JM, Jensen LO, Perkins A, Radovanovic N, Rakhit RD, Ravkilde J, Ryding AD, Schmidt MR, Riddervold IS, Sorensen HT, Stankovic G,

- Varma M, Webb I, Terkelsen CJ, Greenwood JP, Yellon DM, Botker HE, Investigators C-E-P (2019) Effect of remote ischaemic conditioning on clinical outcomes in patients with acute myocardial infarction (CONDI-2/ERIC-PPCI): a single-blind randomised controlled trial. *Lancet* 394:1415-1424 doi:10.1016/S0140-6736(19)32039-2
30. Hernán MA, Robins JM (2020) *Causal Inference: What If*. Boca Raton: Chapman & Hall/CRC, forthcoming
31. Heusch G (2016) The Coronary Circulation as a Target of Cardioprotection. *Circ Res* 118:1643-1658 doi:10.1161/CIRCRESAHA.116.308640
32. Heusch G (2015) Molecular basis of cardioprotection: signal transduction in ischemic pre-, post-, and remote conditioning. *Circ Res* 116:674-699 doi:10.1161/CIRCRESAHA.116.305348
33. Heusch G (2020) Myocardial ischaemia-reperfusion injury and cardioprotection in perspective. *Nat Rev Cardiol* 17:773-789 doi:10.1038/s41569-020-0403-y
34. Heusch G, Botker HE, Przyklenk K, Redington A, Yellon D (2015) Remote ischemic conditioning. *J Am Coll Cardiol* 65:177-195 doi:10.1016/j.jacc.2014.10.031
35. Johnsen J, Pryds K, Salman R, Lofgren B, Kristiansen SB, Botker HE (2016) The remote ischemic preconditioning algorithm: effect of number of cycles, cycle duration and effector organ mass on efficacy of protection. *Basic Res Cardiol* 111:10 doi:10.1007/s00395-016-0529-6
36. Khan JN, Greenwood JP, Nazir SA, Lai FY, Dalby M, Curzen N, Hetherington S, Kelly DJ, Blackman D, Peebles C, Wong J, Flather M, Swanton H, Gershlick AH, McCann GP (2016) *Infarct Size Following Treatment With Second- Versus Third-Generation P2Y12 Antagonists in Patients With Multivessel Coronary Disease at ST-Segment Elevation*

Myocardial Infarction in the CvLPRIT Study. *J Am Heart Assoc* 5(6):e003403

doi:10.1161/JAHA.116.003403

37. Kim EK, Park TK, Yang JH, Song YB, Choi JH, Choi SH, Chun WJ, Choe YH, Gwon HC, Hahn JY (2017) Ticagrelor Versus Clopidogrel on Myocardial Infarct Size in Patients Undergoing Primary Percutaneous Coronary Intervention. *J Am Coll Cardiol* 69:2098-2099
doi:10.1016/j.jacc.2017.02.034
38. Kleinbongard P, Botker HE, Ovize M, Hausenloy DJ, Heusch G (2019) Co-morbidities and co-medications as confounders of cardioprotection-Does it matter in the clinical setting? *Br J Pharmacol* doi:10.1111/bph.14839
39. Kleinbongard P, Skyschally A, Heusch G (2017) Cardioprotection by remote ischemic conditioning and its signal transduction. *Pflugers Arch* 469:159-181 doi:10.1007/s00424-016-1922-6
40. Lieder HR, Irmert A, Kamler M, Heusch G, Kleinbongard P (2019) Sex is no determinant of cardioprotection by ischemic preconditioning in rats, but ischemic/reperfused tissue mass is for remote ischemic preconditioning. *Physiol Rep* 7:e14146 doi:10.14814/phy2.14146
41. Mehta SR, Tanguay JF, Eikelboom JW, Jolly SS, Joyner CD, Granger CB, Faxon DP, Rupprecht HJ, Budaj A, Avezum A, Widimsky P, Steg PG, Bassand JP, Montalescot G, Macaya C, Di Pasquale G, Niemela K, Ajani AE, White HD, Chrolavicius S, Gao P, Fox KA, Yusuf S, investigators C-Ot (2010) Double-dose versus standard-dose clopidogrel and high-dose versus low-dose aspirin in individuals undergoing percutaneous coronary intervention for acute coronary syndromes (CURRENT-OASIS 7): a randomised factorial trial. *Lancet* 376:1233-1243 doi:10.1016/S0140-6736(10)61088-4
42. Mehta SR, Yusuf S, Peters RJ, Bertrand ME, Lewis BS, Natarajan MK, Malmberg K, Rupprecht H, Zhao F, Chrolavicius S, Copland I, Fox KA, Clopidogrel in Unstable angina

to prevent Recurrent Events trial I (2001) Effects of pretreatment with clopidogrel and aspirin followed by long-term therapy in patients undergoing percutaneous coronary intervention: the PCI-CURE study. *Lancet* 358:527-533 doi:10.1016/s0140-6736(01)05701-4

43. Meybohm P, Bein B, Brosteanu O, Cremer J, Gruenewald M, Stoppe C, Coburn M, Schaelte G, Boning A, Niemann B, Roesner J, Kletzin F, Strouhal U, Reyher C, Laufenberg-Feldmann R, Ferner M, Brandes IF, Bauer M, Stehr SN, Kortgen A, Wittmann M, Baumgarten G, Meyer-Treschan T, Kienbaum P, Heringlake M, Schon J, Sander M, Treskatsch S, Smul T, Wolwender E, Schilling T, Fuernau G, Hasenclever D, Zacharowski K, Collaborators RIS (2015) A Multicenter Trial of Remote Ischemic Preconditioning for Heart Surgery. *N Engl J Med* 373:1397-1407 doi:10.1056/NEJMoa1413579
44. Mohammad HMF, Makary S, Atef H, El-Sherbiny M, Atteia HH, Ibrahim GA, Mohamed AS, Zaitone SA (2020) Clopidogrel or prasugrel reduces mortality and lessens cardiovascular damage from acute myocardial infarction in hypercholesterolemic male rats. *Life Sci* 247:117429 doi:10.1016/j.lfs.2020.117429
45. Montalescot G, van 't Hof AW, Lapostolle F, Silvain J, Lassen JF, Bolognese L, Cantor WJ, Cequier A, Chettibi M, Goodman SG, Hammett CJ, Huber K, Janzon M, Merkely B, Storey RF, Zeymer U, Stibbe O, Ecollan P, Heutz WM, Swahn E, Collet JP, Willems FF, Baradat C, Licour M, Tsatsaris A, Vicaut E, Hamm CW, Investigators A (2014) Prehospital ticagrelor in ST-segment elevation myocardial infarction. *N Engl J Med* 371:1016-1027 doi:10.1056/NEJMoa1407024
46. Mustafa SJ, Morrison RR, Teng B, Pelleg A (2009) Adenosine receptors and the heart: role in regulation of coronary blood flow and cardiac electrophysiology. *Handb Exp Pharmacol*:161-188 doi:10.1007/978-3-540-89615-9_6

47. Ozyuncu N, Goksuluk H, Tan TS, Esenboga K, Atmaca Y, Erol C (2020) Does the level of myocardial injury differ in primary angioplasty patients loaded first with clopidogrel and the ones with ticagrelor? *Anatol J Cardiol* 24:107-112 doi:10.14744/AnatolJCardiol.2020.22903
48. Penna C, Aragno M, Cento AS, Femmino S, Russo I, Bello FD, Chiazza F, Collotta D, Alves GF, Bertinaria M, Zicola E, Mercurio V, Medana C, Collino M, Pagliaro P (2020) Ticagrelor Conditioning Effects Are Not Additive to Cardioprotection Induced by Direct NLRP3 Inflammasome Inhibition: Role of RISK, NLRP3, and Redox Cascades. *Oxid Med Cell Longev* 2020:9219825 doi:10.1155/2020/9219825
49. Ramakers BP, Riksen NP, Stal TH, Heemskerk S, van den Broek P, Peters WH, van der Hoeven JG, Smits P, Pickkers P (2011) Dipyridamole augments the antiinflammatory response during human endotoxemia. *Crit Care* 15:R289 doi:10.1186/cc10576
50. Roe MT, Armstrong PW, Fox KA, White HD, Prabhakaran D, Goodman SG, Cornel JH, Bhatt DL, Clemmensen P, Martinez F, Ardissino D, Nicolau JC, Boden WE, Gurbel PA, Ruzyllo W, Dalby AJ, McGuire DK, Leiva-Pons JL, Parkhomenko A, Gottlieb S, Topacio GO, Hamm C, Pavlides G, Goudev AR, Oto A, Tseng CD, Merkely B, Gasparovic V, Corbalan R, Cinteza M, McLendon RC, Winters KJ, Brown EB, Lokhnygina Y, Aylward PE, Huber K, Hochman JS, Ohman EM, Investigators TA (2012) Prasugrel versus clopidogrel for acute coronary syndromes without revascularization. *N Engl J Med* 367:1297-1309 doi:10.1056/NEJMoa1205512
51. Roubille F, Lairez O, Mewton N, Rioufol G, Ranc S, Sanchez I, Cung TT, Elbaz M, Piot C, Ovize M (2012) Cardioprotection by clopidogrel in acute ST-elevated myocardial infarction patients: a retrospective analysis. *Basic Res Cardiol* 107:275 doi:10.1007/s00395-012-0275-

52. Sabbah M, Nepper-Christensen L, Kober L, Hofsten DE, Ahtarovski KA, Goransson C, Kyhl K, Ghotbi AA, Schoos MM, Sadjadieh G, Kelbaek H, Lonborg J, Engstrom T (2020) Infarct size following loading with Ticagrelor/Prasugrel versus Clopidogrel in ST-segment elevation myocardial infarction. *Int J Cardiol* 314:7-12 doi:10.1016/j.ijcard.2020.05.011
53. Schupke S, Neumann FJ, Menichelli M, Mayer K, Bernlochner I, Wohrle J, Richardt G, Liebetrau C, Witzenbichler B, Antoniucci D, Akin I, Bott-Flugel L, Fischer M, Landmesser U, Katus HA, Sibbing D, Seyfarth M, Janisch M, Boncompagni D, Hilz R, Rottbauer W, Okrojek R, Mollmann H, Hochholzer W, Migliorini A, Cassese S, Mollo P, Xhepa E, Kufner S, Strehle A, Leggewie S, Allali A, Ndrepepa G, Schuhlen H, Angiolillo DJ, Hamm CW, Hapfelmeier A, Tolg R, Trenk D, Schunkert H, Laugwitz KL, Kastrati A, Investigators I-RT (2019) Ticagrelor or Prasugrel in Patients with Acute Coronary Syndromes. *N Engl J Med* 381:1524-1534 doi:10.1056/NEJMoa1908973
54. Skyschally A, van Caster P, Iliodromitis EK, Schulz R, Kremastinos DT, Heusch G (2009) Ischemic postconditioning: experimental models and protocol algorithms. *Basic Res Cardiol* 104:469-483 doi:10.1007/s00395-009-0040-4
55. Spreeuwenberg MD, Bartak A, Croon MA, Hagenars JA, Busschbach JJ, Andrea H, Twisk J, Stijnen T (2010) The multiple propensity score as control for bias in the comparison of more than two treatment arms: an introduction from a case study in mental health. *Med Care* 48:166-174 doi:10.1097/MLR.0b013e3181c1328f
56. Sugidachi A, Ogawa T, Kurihara A, Hagihara K, Jakubowski JA, Hashimoto M, Niitsu Y, Asai F (2007) The greater in vivo antiplatelet effects of prasugrel as compared to clopidogrel reflect more efficient generation of its active metabolite with similar antiplatelet activity to that of clopidogrel's active metabolite. *J Thromb Haemost* 5:1545-1551 doi:10.1111/j.1538-7836.2007.02598.x

57. Sugidachi A, Ohno K, Ogawa T, Jakubowski J, Hashimoto M, Tomizawa A (2013) A comparison of the pharmacological profiles of prasugrel and ticagrelor assessed by platelet aggregation, thrombus formation and haemostasis in rats. *Br J Pharmacol* 169:82-89 doi:10.1111/bph.12108
58. Sugidachi A, Yamaguchi S, Jakubowski JA, Ohno K, Tomizawa A, Hashimoto M, Niitsu Y (2011) Selective blockade of P2Y₁₂ receptors by prasugrel inhibits myocardial infarction induced by thrombotic coronary artery occlusion in rats. *J Cardiovasc Pharmacol* 58:329-334 doi:10.1097/FJC.0b013e3182244a6f
59. Terkelsen CJ, Sorensen JT, Maeng M, Jensen LO, Tilsted HH, Trautner S, Vach W, Johnsen SP, Thuesen L, Lassen JF (2010) System delay and mortality among patients with STEMI treated with primary percutaneous coronary intervention. *JAMA* 304:763-771 doi:10.1001/jama.2010.1139
60. Ubaid S, Ford TJ, Berry C, Murray HM, Wrigley B, Khan N, Thomas MR, Armesilla AL, Townend JN, Khogali SS, Munir S, Martins J, Hothi SS, McAlindon EJ, Cotton JM (2019) Cangrelor versus Ticagrelor in Patients Treated with Primary Percutaneous Coronary Intervention: Impact on Platelet Activity, Myocardial Microvascular Function and Infarct Size: A Randomized Controlled Trial. *Thromb Haemost* 119:1171-1181 doi:10.1055/s-0039-1688789
61. van den Berg TN, El Messaoudi S, Rongen GA, van den Broek PH, Bilos A, Donders AR, Gomes ME, Riksen NP (2015) Ticagrelor Does Not Inhibit Adenosine Transport at Relevant Concentrations: A Randomized Cross-Over Study in Healthy Subjects In Vivo. *PLoS One* 10:e0137560 doi:10.1371/journal.pone.0137560
62. van der Hoeven NW, Janssens GN, Everaars H, Nap A, Lemkes JS, de Waard GA, van de Ven PM, van Rossum AC, Escaned J, Mejia-Renteria H, Ten Cate TJF, Piek JJ, von

- Birgelen C, Valgimigli M, Diletti R, Rixsen NP, Van Mieghem NM, Nijveldt R, van Leeuwen MAH, van Royen N (2020) Platelet Inhibition, Endothelial Function, and Clinical Outcome in Patients Presenting With ST-Segment-Elevation Myocardial Infarction Randomized to Ticagrelor Versus Prasugrel Maintenance Therapy: Long-Term Follow-Up of the REDUCE-MVI Trial. *J Am Heart Assoc* 9:e014411 doi:10.1161/JAHA.119.014411
63. van Giezen JJ, Sidaway J, Glaves P, Kirk I, Bjorkman JA (2012) Ticagrelor inhibits adenosine uptake in vitro and enhances adenosine-mediated hyperemia responses in a canine model. *J Cardiovasc Pharmacol Ther* 17:164-172 doi:10.1177/1074248411410883
64. van Leeuwen MAH, van der Hoeven NW, Janssens GN, Everaars H, Nap A, Lemkes JS, de Waard GA, van de Ven PM, van Rossum AC, Ten Cate TJF, Piek JJ, von Birgelen C, Escaned J, Valgimigli M, Diletti R, Rixsen NP, van Mieghem NM, Nijveldt R, van Royen N (2019) Evaluation of Microvascular Injury in Revascularized Patients With ST-Segment-Elevation Myocardial Infarction Treated With Ticagrelor Versus Prasugrel. *Circulation* 139:636-646 doi:10.1161/CIRCULATIONAHA.118.035931
65. Vannini L, Muro A, Sanchis J, Ortiz-Perez JT, Flores Umanzor E, Lopez-Lereu MP, Badimon L, Sabate M, Brugaletta S (2016) Can new generation P2Y12 inhibitors play a role in microvascular obstruction in STEMI? *Int J Cardiol* 223:226-227 doi:10.1016/j.ijcard.2016.08.182
66. Vilahur G, Gutierrez M, Casani L, Varela L, Capdevila A, Pons-Llado G, Carreras F, Carlsson L, Hidalgo A, Badimon L (2016) Protective Effects of Ticagrelor on Myocardial Injury After Infarction. *Circulation* 134:1708-1719 doi:10.1161/CIRCULATIONAHA.116.024014
67. Wallentin L, Becker RC, Budaj A, Cannon CP, Emanuelsson H, Held C, Horrow J, Husted S, James S, Katus H, Mahaffey KW, Scirica BM, Skene A, Steg PG, Storey RF, Harrington

- RA, Investigators P, Freij A, Thorsen M (2009) Ticagrelor versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med* 361:1045-1057
doi:10.1056/NEJMoa0904327
68. Weisshaar S, Litschauer B, Eipeldauer M, Hobl EL, Wolzt M (2017) Ticagrelor mitigates ischaemia-reperfusion induced vascular endothelial dysfunction in healthy young males - a randomized, single-blinded study. *Br J Clin Pharmacol* 83:2651-2660
doi:10.1111/bcp.13378
69. Wittfeldt A, Emanuelsson H, Brandrup-Wognsen G, van Giezen JJ, Jonasson J, Nylander S, Gan LM (2013) Ticagrelor enhances adenosine-induced coronary vasodilatory responses in humans. *J Am Coll Cardiol* 61:723-727 doi:10.1016/j.jacc.2012.11.032
70. Wiviott SD, Braunwald E, McCabe CH, Montalescot G, Ruzyllo W, Gottlieb S, Neumann FJ, Ardissino D, De Servi S, Murphy SA, Riesmeyer J, Weerakkody G, Gibson CM, Antman EM, Investigators T-T (2007) Prasugrel versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med* 357:2001-2015 doi:10.1056/NEJMoa0706482
71. Wu Q, Gui P, Wu J, Ding D, Purusram G, Dong N, Yao S (2011) Effect of limb ischemic preconditioning on myocardial injury in patients undergoing mitral valve replacement surgery. -A randomized controlled trial. *Circ J* 75:1885-1889 doi:10.1253/circj.cj-10-1130
72. Yang XM, Cui L, Alhammouri A, Downey JM, Cohen MV (2013) Triple therapy greatly increases myocardial salvage during ischemia/reperfusion in the in situ rat heart. *Cardiovasc Drugs Ther* 27:403-412 doi:10.1007/s10557-013-6474-9
73. Yang XM, Gadde S, Audia JP, Alvarez DF, Downey JM, Cohen MV (2019) Ticagrelor Does Not Protect Isolated Rat Hearts, Thus Clouding Its Proposed Cardioprotective Role Through ENT 1 in Heart Tissue. *J Cardiovasc Pharmacol Ther* 24:371-376
doi:10.1177/1074248419829169

74. Yang XM, Liu Y, Cui L, Yang X, Liu Y, Tandon N, Kambayashi J, Downey JM, Cohen MV (2013) Platelet P2Y(1)(2) blockers confer direct postconditioning-like protection in reperfused rabbit hearts. *J Cardiovasc Pharmacol Ther* 18:251-262
doi:10.1177/1074248412467692
75. Yang XM, Liu Y, Cui L, Yang X, Liu Y, Tandon N, Kambayashi J, Downey JM, Cohen MV (2013) Two classes of anti-platelet drugs reduce anatomical infarct size in monkey hearts. *Cardiovasc Drugs Ther* 27:109-115 doi:10.1007/s10557-012-6436-7
76. Ye Y, Birnbaum GD, Perez-Polo JR, Nanhwan MK, Nylander S, Birnbaum Y (2015) Ticagrelor protects the heart against reperfusion injury and improves remodeling after myocardial infarction. *Arterioscler Thromb Vasc Biol* 35:1805-1814
doi:10.1161/ATVBAHA.115.305655
77. Yellon DM, Hausenloy DJ (2007) Myocardial reperfusion injury. *N Engl J Med* 357:1121-1135 doi:10.1056/NEJMra071667
78. Yusuf S, Zhao F, Mehta SR, Chrolavicius S, Tognoni G, Fox KK, Clopidogrel in Unstable Angina to Prevent Recurrent Events Trial I (2001) Effects of clopidogrel in addition to aspirin in patients with acute coronary syndromes without ST-segment elevation. *N Engl J Med* 345:494-502 doi:10.1056/NEJMoa010746
79. Ziegler M, Wang X, Peter K (2019) Platelets in cardiac ischaemia/reperfusion injury: a promising therapeutic target. *Cardiovasc Res* 115:1178-1188 doi:10.1093/cvr/cvz070

Figure titles and legends

Figure 1 Infarct size in rats. Final infarct size of area at risk. IS: infarct size, AAR: area at risk, CON: control, IPC: local ischemic preconditioning, RIC: remote ischemic preconditioning. All statistical comparisons showed have controls as reference. * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$

Figure 2 Flowchart. Flowchart of patient selection and exclusions.

Figure 3 Graphical presentations. a) the composite endpoint of cardiac death and b) hospitalization for heart failure, and major adverse cardiovascular events, c) and d) display 30-day curves of the same endpoints.

Tables

Table 1. Baseline characteristics and procedural details.

	Clopidogrel	Ticagrelor	Prasugrel
	(n=395)	(n=1,210)	(n=149)
Mean age, years (SD)	65.7 (12.4)	63.7 (12.0)	61.1 (11.0)
Sex			
Male	294 (74.4%)	946 (78.2%)	125 (83.9%)
Female	101 (25.6%)	264 (21.8%)	24 (16.1%)
Current smoker	132 (33.4%)	437 (36.1%)	59 (39.6%)
Body mass index, kg/m ² (SD)	27.5 (4.8)	27.6 (5.1)	28.2 (4.5)
eGFR, µg/L/1.73 m ² (IQR)	87 (71-97)	86 (72-96)	87 (77-99)
Comorbidity			
Hypertension	175 (44.3%)	489 (40.4%)	58 (38.9%)
Previous MI	35 (8.9%)	95 (7.9%)	10 (6.7%)
Hypercholesterolaemia	121 (30.6%)	353 (29.2%)	46 (30.9%)
Peripheral vascular disease	9 (2.3%)	18 (1.5%)	1 (0.7%)
Diabetes	48 (12.2%)	131 (10.8%)	15 (10.1%)
Family history of IHD	101 (25.6%)	455 (37.6%)	62 (41.6%)
Baseline medication			
Insulin	12 (3%)	31 (2.6%)	5 (3.4%)
Metformin	39 (9.9%)	100 (8.3%)	10 (6.7%)
Sulphonylurea	14 (3.5%)	37 (3.1%)	5 (3.4%)
Other anti-diabetic medication	16 (4.1%)	29 (2.4%)	5 (3.4%)
Statin	98 (24.8%)	272 (22.5%)	35 (23.5%)
Beta-blocker	49 (12.4%)	139 (11.5%)	18 (12.1%)
ACE-inhibitor	74 (18.7%)	194 (16%)	31 (20.8%)
ARB	40 (10.1%)	106 (8.8%)	9 (6%)

Aspirin	53 (13.4%)	180 (14.9%)	15 (10.1%)
Diuretics	32 (8.1%)	85 (7%)	18 (12.1%)
Blood pressure at inclusion (mmHg)			
Systolic (SD)	132.1 (25.9)	133.1 (22.7)	126.3 (22.6)
Diastolic (SD)	78.7 (16.4)	80.2 (15.7)	73.6 (15.0)
Killip Class on admission			
Class I	389 (98.5%)	1,172 (96.9%)	145 (97.3%)
Class II	2 (0.5%)	33 (2.7%)	1 (0.7%)
Class III	0	1 (0.1%)	0
Class IV (including cardiogenic shock)	4 (1%)	4 (0.3%)	3 (2%)
Symptom to balloon time, min (IQR)	177 (129-261)	185 (138-298)	176 (121-257)
First medical contact to balloon time, min (IQR)	105 (90-127)	112 (92-137)	104 (86-132)
Culprit vessel			
Left anterior descending	160 (40.5%)	489 (40.4%)	62 (41.6%)
Circumflex	43 (10.9%)	160 (13.2%)	21 (14.1%)
Right coronary	190 (48.1%)	551 (45.5%)	63 (42.3%)
Other	0	4 (0.3%)	0
Missing	2 (0.5%)	6 (0.5%)	3 (2%)
Culprit lesion stented	367 (92.9%)	1,170 (96.7%)	139 (93.3%)
Number of vessels with angiographically significant disease			
0	0	2 (0.2%)	0
1	222 (56.2%)	639 (52.8%)	89 (59.7%)
2	125 (31.6%)	388 (32.1%)	39 (26.2%)
3	46 (11.6%)	176 (14.5%)	18 (12.1%)
Missing	2 (0.5%)	5 (0.4%)	3 (2%)
Thrombus aspiration performed	115 (29.1%)	378 (31.2%)	39 (26.2%)
TIMI flow pre-angioplasty			
TIMI 0	284 (71.9%)	944 (78%)	116 (77.9%)

TIMI 1	31 (7.8%)	72 (6%)	12 (8.1%)
TIMI 2	32 (8.1%)	88 (7.3%)	12 (8.1%)
TIMI 3	48 (12.2%)	106 (8.8%)	9 (6%)
Missing	284 (71.9%)	944 (78%)	116 (77.9%)
TIMI flow post-procedure			
TIMI 0	5 (1.3%)	14 (1.2%)	0
TIMI 1	4 (1%)	7 (0.6%)	1 (0.7%)
TIMI 2	27 (6.8%)	56 (4.6%)	5 (3.4%)
TIMI 3	351 (88.9%)	1,092 (90.2%)	139 (93.3%)
Missing	8 (2%)	41 (3.4%)	4 (2.7%)
Staged PCI performed	36 (9.1%)	108 (8.9%)	7 (4.7%)
Staged CABG performed	5 (1.3%)	21 (1.7%)	2 (1.3%)
pPCI related medication			
Opioids	0	0	0
Heparin	372 (94.2%)	1,159 (95.8%)	144 (96.6%)
Aspirin	375 (94.9%)	1,136 (93.9%)	138 (92.6%)
Glycoprotein IIb/IIIa inhibitor	94 (23.8%)	324 (26.8%)	45 (30.2%)
Bivalirudin	21 (5.3%)	35 (2.9%)	0
Protaminsulphate	2 (0.5%)	5 (0.4%)	1 (0.7%)
Nitrates	353 (89.4%)	984 (81.3%)	138 (92.6%)
Country			
UK	395 (100%)	1,210 (100%)	149 (100%)

Table 1. Baseline characteristics of the patient cohorts included in the propensity weighted analyses. On the left the cohorts included in the analyses of clopidogrel vs ticagrelor, and on the right the cohorts included in the analyses of clopidogrel vs prasugrel. IHD: Ischemic heart disease, ACE: Angiotensin converting enzyme, ARB: Angiotensin receptor blocker, TIMI: Thrombolysis in myocardial infarction, SD: Standard deviation, IQR: Interquartile range.

Table 2. One-year cardiovascular risk in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention treated with either clopidogrel, ticagrelor, or prasugrel.

	Patients	Events	Cumulative incidence proportion (95% CI)	Unadjusted HR (95% CI)	Stabilized IPW weighted HR (95% CI)
Cardiac death or hospitalization for heart failure					
Clopidogrel	395	38	9.6% (7.0-12.8)	reference	reference
Ticagrelor	1,210	78	6.5% (5.2-7.9)	0.66 (0.45-0.97)	0.63 (0.42-0.94)
Prasugrel	149	12	8.1% (4.4-13.1)	0.83 (0.44-1.57)	0.84 (0.43-1.63)
Major adverse cardiovascular events					
Clopidogrel	395	30	7.6% (5.4-10.7)	reference	reference
Ticagrelor	1,210	83	6.9% (5.6-8.4)	0.90 (0.59-1.36)	0.86 (0.56-1.32)
Prasugrel	149	8	5.4% (2.7-10.5)	0.70 (0.32-1.53)	0.54 (0.24-1.23)

Table 2 One-year cardiovascular risk. One-year risk of cardiac death or hospitalization for heart failure, or major adverse cardiovascular events (MACE) in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention treated with either ticagrelor compared to clopidogrel, or prasugrel compared to clopidogrel. HR: hazard ratios, IPW: inverse-probability-weighted.