

ORIGINAL RESEARCH

Value of imaging to guide interventional procedures in rheumatic and musculoskeletal diseases: a systematic literature review informing EULAR points to consider

Philipp Bosch ¹, Francesco Carubbi ^{2,3}, Carlo Alberto Scirè ⁴, Xenofon Baraliakos ⁵, Louise Falzon,⁶ Christian Dejaco ^{1,7}, Pedro M Machado ^{8,9,10}

To cite: Bosch P, Carubbi F, Scirè CA, et al. Value of imaging to guide interventional procedures in rheumatic and musculoskeletal diseases: a systematic literature review informing EULAR points to consider. *RMD Open* 2021;7:e001864. doi:10.1136/rmdopen-2021-001864

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/rmdopen-2021-001864>).

PB and FC are joint first authors.

Received 9 August 2021

Accepted 5 October 2021

ABSTRACT

Objectives To summarise current data on the value of imaging to guide interventional procedures in patients with rheumatic and musculoskeletal disease (RMDs) informing an European Alliance of Associations for Rheumatology taskforce.

Methods A systematic literature review was conducted to retrieve prospective and retrospective studies published in English and comparing different (imaging) techniques, different settings and procedural protocols to guide interventions in patients with RMDs. MEDLINE, EMBASE, the Cochrane Library and Epistemonikos databases were searched through October 2021. Risk of bias (RoB) was assessed using the Cochrane RoB tool for randomised trials V.2 (ROB2), the RoB tool for Non-Randomised Studies of Interventions and the appraisal tool for cross-sectional studies.

Results Sixty-six studies were included (most with moderate/high RoB); 49 were randomised controlled trials, three prospective cohort studies and 14 retrospective studies. Fifty-one studies compared either one imaging technique with another imaging technique, or with palpation-guided interventions. Ultrasound (US) was most frequently studied (49/51), followed by fluoroscopy (10/51). Higher accuracy was found for US or fluoroscopy compared with palpation-guided interventions. Studies comparing different imaging techniques (12/51) did not endorse one specific method. Different settings/equipment for imaging-guided procedures (eg, automatic vs manual syringes) were investigated in three studies, reporting heterogeneous results. Fifteen studies compared different imaging-guided procedures (eg, intra-articular vs periarticular injections).

Conclusion Higher accuracy of needle positioning at joints and periarticular structures was seen in most studies when using imaging (especially US) guidance as compared with palpation-guided interventions with the limitation of heterogeneity of data and considerable RoB.

Key messages

What is already known about this subject?

- Imaging techniques such as ultrasound, fluoroscopy, CT and MRI can be used to perform interventional procedures in patients with rheumatic and musculoskeletal diseases. However, consensus on when to use imaging, which settings should be preferred and whether certain procedural protocols may be better than others is still unclear.

What does this study add?

- Using imaging guidance leads to better accuracy compared with palpation-guided interventions.
- Positioning a needle or another instrument in, rather than outside the target not only reduces the risk of adverse events but may also have beneficial effects on short-term and long-term pain, as well as on efficacy outcomes.

How might this impact on clinical practice or further developments?

- The results of this systematic literature review may help clinicians to decide when and how to use imaging to guide interventional procedures in patients with rheumatic and musculoskeletal diseases.

INTRODUCTION

Imaging is widely used to guide interventional procedures in patients with rheumatic and musculoskeletal disease (RMDs).¹ However, consensus on when to use imaging, which settings should be preferred and whether certain procedural protocols may be better than others is still unclear.

A number of previous systematic literature reviews (SLRs) have addressed some of these aspects,^{2–7} however, the data reported were mostly limited to certain joints or diseases, or



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Christian Dejaco;
christian.dejaco@gmx.net



did not focus on studies comparing directly two or more techniques.

The objective of this SLR was to gather available data on different imaging methodologies, settings and procedural protocols in RMDs to inform an European Alliance of Associations for Rheumatology (EULAR) task force developing points to consider on the use of imaging for interventional procedures in patients with RMDs.

METHODS

Literature review

In the first EULAR task force meeting, the steering group framed three broad key questions according to the Population, Intervention, Comparator, Outcome (PICO) format related to the following aspects⁸: (1) What is the value of imaging methods to guide interventional procedures in patients with RMDs, (2) What is the value of different imaging settings and technical standards and (3) What is the value of different procedural techniques for imaging-guided interventions. See see online supplemental table 1 for details on the PICO questions. Eligible studies were full research articles, short reports and research letters of prospective and retrospective studies. The following studies were excluded from the analysis: studies of interventions on tumours, vessels or glands, studies on vertebroplasty/arthroplasty, studies of interventions with the aim to perform anaesthesia and studies investigating methods not used in routine clinical practice. We also excluded studies not answering any of the PICO questions, for example, due to wrong population or missing comparator.

Study selection, data extraction and assessment of risk of bias

Two reviewers (PB and FC) conducted the SLR under the guidance of the methodologist (PMM) and the junior methodologist (CAS). The search strategy was developed by an experienced librarian (LF). The databases MEDLINE, Embase, Epistemonikos and the Cochrane Library were searched until September 2019. An update of the original search was conducted with the same search strategy from September 2019 to October 2020. See online supplemental table 2 for the list of key words used.

The reviewers independently screened titles and abstracts to identify eligible studies, which were consecutively assessed for eligibility by reading the full text. Both reviewers independently retrieved data from included studies using a predefined data extraction sheet. The following data were extracted: study design, year of publication, number of patients included, the site where the intervention was performed, details on the procedure including equipment, imaging method(s) and individual settings, in case of injections—the compound(s) administered, data on monitoring and potential outcomes. The extracted outcomes were then assigned to the following categories: accuracy, safety, efficacy and cost/time as

detailed in online supplemental table 1. Outcome variables on pain, including procedural, short-term and long-term pain, were altogether considered safety outcomes. Data on arthroscopy were only extracted if it was compared with a second imaging technique (ie, ultrasound (US), CT, MRI or fluoroscopy).

Risk of bias (RoB) was appraised independently by the same reviewers who conducted the SLR. For randomised controlled trials (RCTs), the Cochrane risk-of-bias tool for randomised trials V.2 (RoB 2) was used.⁹ The Risk Of Bias tool for Non-Randomised Studies of Interventions (ROBINS-I) was applied to non-randomised trials and the Appraisal tool for cross-sectional studies (AXIS) was used for cross-sectional studies.^{10 11} Disagreement on study selection, data extraction and RoB assessment was solved by discussion until a consensus was achieved, the methodologist (PMM) served as a tiebreaker. To homogenise reporting of the different tools for the assessment of RoB, we transformed the items ‘serious concern’ and ‘some concern’ used in the original version of the ROBINS-I tool into ‘high’ and ‘moderate’ RoB, according to the RoB 2 classification. Since the AXIS tool does not provide a global classification of the RoB, but rather includes a series of question addressing possible quality concerns, we grouped the studies according to the number of questions with a positive answer (ie, answer indicating less concern) as follows: <10/20=high RoB, 10–15/20=moderate RoB, >15/20=low RoB. According to this approach studies were classified into the following four RoB categories: low, moderate, high and critical RoB (the latter was used for RCTs only). In tables of the main manuscript, the RoB is depicted by coloured circles in green, yellow, red and black respectively. Quality appraisal was performed for each outcome of every study separately.

RESULTS

A total of 6049 references (after duplicates removal) were screened for eligibility (see online supplemental figure 1). Sixty-six articles fulfilled the inclusion criteria, with some of them addressing multiple PICO questions. Fifty-one articles compared interventions with imaging or palpation guidance, or interventions assisted by different imaging techniques (PICO1),^{12–62} three articles assessed the influence of different settings (PICO2),^{25 63 64} while 16 articles compared the value of different procedures for imaging-guided interventions (PICO3).^{22 61 65–78} RoB assessment of the retrieved studies is detailed in online supplemental table 3.

Imaging techniques for interventional procedures (PICO1)

Large joints and periarticular structures

Twenty-nine articles (23 RCTs, 4 retrospective cohort studies, 1 retrospective cross-sectional study and 1 RCT-letter) assessed the value of imaging for interventions at large joints, with 20 studies comparing US versus palpation-guided interventions,^{18–33 55–58} 5 investigating

US versus fluoroscopy^{12–16} and 1 study each comparing fluoroscopy with palpation guidance,⁵³ fluoroscopy with US and palpation guidance,⁵⁹ or fluoroscopy with CT-guided interventions,⁵⁴ as well as one study comparing either US or fluoroscopy with palpation guidance.¹⁷ The following large joints and periarticular structures were studied: knee (n=10), shoulder (n=9), sacroiliac joint (SIJ) (n=3) and one study each for elbow, wrist, spine, piriformis muscle, hip, trochanteric bursa and ankle. All interventions were injections with one or more of the following compounds: glucocorticoids (GC), local anaesthetics, viscosupplements and/or contrast media. The most important characteristics of the 29 studies are depicted in **table 1**, full details on interventions and outcomes can be found in online supplemental table 4.

Accuracy

Nine studies reported better accuracy for needle placement/injectate location (n=8) or synovial fluid acquisition (n=1) using US as compared with palpation-guided injections (seven with moderate RoB^{23–26 29 31 56} and two with high RoB),^{33 58} while two studies concluded that there was no difference between the two methods (one moderate, one high RoB).^{20 32} Results for correct needle placement were heterogeneous for comparisons of US with fluoroscopy: one study favoured US for biceps tendon sheath injections (high RoB),¹⁴ one reported better accuracy for SIJ injections using fluoroscopy (high RoB)¹⁵ and another study reported no difference between the techniques for SIJ injections (moderate RoB).¹³ Fluoroscopy was superior to palpation-guided injections for injectate location in patients with painful SIJ (low RoB)⁵³ and superior to CT when assessing the positive rate of discovertebral biopsy cultures at the lumbar spine in patients with osteomyelitis (high RoB).⁵⁴

Safety

No difference in the overall frequency of adverse events was observed between US and palpation-guided interventions in eight studies (five high, three moderate RoB),^{19 20 22 27 33 55 56 58} while one study reported a lower rate of contact of the needle to the cartilage of the patella by using US (moderate RoB).²⁴ Four studies (two high, two moderate RoB)^{25 28 57 58} reported lower procedural pain using US as compared with palpation-guided injections, while one study (high RoB)¹⁹ found no difference between the groups. Long-term pain (ranging from 1 week to 6 months) was found to be lower in US, compared with palpation-guided interventions according to ten studies (five high, five moderate RoB),^{19 21 25 27 28 30 33 56–58} whereas four (two high, two moderate RoB)^{20 22 32 55} studies found no difference between the groups. Furthermore, five studies comparing US and fluoroscopy (three high, two moderate RoB)^{12–16} and one comparing US or fluoroscopy versus palpation-guided injections (moderate RoB)¹⁷ reported no differences in procedural,^{13 16} and long-term pain.^{12–17} One study (low RoB) comparing fluoroscopy with palpation-guided injections reported

lower pain levels at 3 months in the fluoroscopy group when adjusting data for baseline pain levels.⁵³

Efficacy

Results for efficacy outcomes comparing US and palpation-guided interventions were heterogeneous: Function, quality of life, treatment response/treatment failure and time until the next intervention were better in the US groups according to ten studies (total of 24 distinct outcomes, 7 of them with moderate, 17 with high RoB),^{18 20 21 27 28 30 33 55 57 58} while 8 studies reported no difference between these groups for 18 outcomes (3 with moderate, 15 with high RoB).^{19–22 27 28 32 55}

In one retrospective cohort study with high RoB, an even worse outcome for US versus palpation-guided injection was found with a higher percentage of patients requiring an additional knee reinjection with hyaluronic acid in the US group (20.3% vs 15%, respectively).¹⁸ In the palpation-guided group, however, more patients received an additional GC injection (34% vs 27.4%, respectively). In another study, it was observed that patients with knee osteoarthritis and US-guided injection had a worse quality of life at 3 months than patients with palpation-guided injections, while walking and standing⁵⁵ tests were better in the former than in the latter group.

All four studies comparing US and fluoroscopy guidance of intervention revealed no difference concerning the nine efficacy outcomes assessed (six outcomes with moderate, three with high RoB).^{12 13 15 16} In contrast, one study comparing US or fluoroscopy versus palpation-guided injections reported a higher patient satisfaction and patients' judgement of treatment efficacy in the US/fluoroscopy as compared with the palpation group. The need for pain medication and tolerability of the intervention was similar in both groups.¹⁷ A study comparing fluoroscopy with palpation-guided injections (low RoB) in SIJ found no difference between groups concerning function, pain medication and patient satisfaction.⁵³

Cost and time

One RCT with high RoB reported worse cost-effectiveness for US compared with palpation-guided interventions in the USA according to reimbursements by Medicare, and taking into consideration the time that was needed until the next intervention.¹⁹ In another RCT from the USA with moderate RoB, higher annual costs for US were observed only in private practice, whereas costs in hospitals' outpatient clinics were similar for US and palpation-guided interventions. Cost-effectiveness was nevertheless better for US considering a longer therapeutic effect of the injection in case US was used.²⁸ Costs for US and fluoroscopy were similar in one study (high RoB) from the USA,¹⁴ while higher costs for fluoroscopy as compared with US and/or palpation-guided interventions were reported in another study (high RoB).⁵⁹

Procedural times for US guidance were either similar,²⁴ or shorter⁵⁸ than those for palpation guidance (one high, one moderate RoB). Fluoroscopy-guided injections

Table 1 Studies comparing different imaging techniques or imaging vs palpation guidance in large joints and periarticular structures (PICO1)

Year author	Study design	Imaging	Comparator	Disease	No Imaging/ Comparator	Intervention (COMPOUND)	SITE	FAVORS imaging*	FAVORS comparator*	No difference between imaging and comparator*	Robot†
US vs palpation guidance											
2017 Raeissadat ²⁰	RCT	US	Palpation guidance	Adhesive capsulitis	20/21	Injection (contrast medium, triamcinolone, lidocaine)	shoulder joint	1x Efficacy (1.4w)	-	1x Accuracy (BSL) 2x Safety (BSL,1.4w) 5x Efficacy (BSL,1.4w)	●
2009 Lee ³⁰	RCT	US	Palpation guidance	Adhesive capsulitis	20/20	Injection (triamcinolone, lidocaine, saline hyaluronic acid)	shoulder joint	2x Safety (1.2w) 5x Efficacy (1.2,3.4w)	-	-	●
2014 Saeed ²¹	RCT	US	Palpation guidance	Shoulder impingement	41/39	Injection (methylprednisolone, lidocaine)	glenohumeral joint, biceps tendon sheath, AC joint)	1x Safety (6.12w) 2x Efficacy (6.12w)	-	1x Efficacy (12w)	●
2004 Naredo ³³	RCT	US	Palpation guidance	Periarticular shoulder disorders	21/20	Injection (triamcinolone)	SA-SD bursa	1x Accuracy (BSL) 1x Safety (6w) 1x Efficacy (6w)	-	1x Safety (BSL)	●
2011 Hashiuchi ²⁹	RCT	US	Palpation guidance	Anterior shoulder pain	15/15	Injection (contrast agent)	biceps tendon sheath	1x Accuracy (BSL)	-	-	●
2011 Zhang ²⁷	RCT	US	Palpation guidance	Biceps brachii tendinitis	53/45	Injection (lidocaine, triamcinolone)	brachial bicipital groove	1x Safety (n.a.) 2x Efficacy (n.a.)	-	1x Safety (n.a.) 1x Efficacy (n.a.)	●
2019 Yiannakopoulos ³⁸	RCT	US	Palpation guidance	Tendinitis of the long head of the biceps	22/22	Injection (triamcinolone, bipuvacaine)	long biceps tendon	1x Accuracy (BSL) 1x Safety (BSL,4w,6m) 2x Efficacy (4w,6m) 1x Cost/Time (BSL)	-	1x Safety (up to 6m)	●
2014 Chang ²²	RCT	US	Palpation guidance	Scapular pain	18/18	Injection (triamcinolone, lidocaine)	subscapularis muscle	-	-	2x Safety (BSL,1.2,3w,3m)	●
2013 Kim ²³	RCT	US	Palpation guidance	Elbow OA	40/40	Injection (dexamethasone, lidocaine, contrast medium)	Elbow joint	1x Accuracy (BSL)	-	-	●
2008 Luz ³² (letter)	RCT	US	Palpation guidance	RA with wrist synovitis	30/30	Injection (lidocaine, triamcinolone, contrast medium, air)	wrist joint	-	-	1x Accuracy (BSL) 2x Safety (BSL,1.4,8,12w) 1x Efficacy (BSL,1.4,8,12w)	●
2018 Mitchell ¹⁹	RCT	US	Palpation guidance	Trochanteric bursitis	15/15	Injection (lidocaine, methylprednisolone)	Trochanteric bursa	1x Safety (6m) (n.a.)	1x Accuracy (BSL)	1x Safety (n.a.) 2x Efficacy (6m,12 m)	●
2011 Sibbitt ²⁸	RCT	US	Palpation guidance	Knee OA	46/46	Injection (Xylocaine, triamcinolone)	knee joint	1x Safety (BSL,2w,6m) 1x Efficacy (BSL to 6m)	1x Cost/Time (BSL to 12m)	1x Efficacy (BSL to 12m) 1x Cost/Time (BSL to 12m)	●
2011 Bum Park, ²⁶	RCT	US	Palpation guidance	Knee OA	50/49	Injection (hyaluronic acid, contrast dye)	knee joint	1x Accuracy (BSL)	-	-	●
2013 Jang ³⁴	RCT	US	Palpation guidance	Knee joint OA	41/44/41	Injection (lidocaine, triamcinolone, contrast agent)	knee joint	1x Accuracy (BSL) 1x Safety (BSL)	-	1x Cost/Time (BSL)	●
2020 Cankurtaran ⁵⁵	RCT	US	Palpation guidance	Chronic knee OA	11/12	Injection (triamcinolone, lidocaine)	knee joint	2x Efficacy (1.3m)	3x Efficacy (3m)	2x Safety (BSL,1m,3m) 6x Efficacy (1m,3m)	●
2012 Sibbitt ²⁵	RCT	US	Palpation guidance	Knee effusion	42/22	aspiration, injection (lidocaine, triamcinolone)	knee joint	2x Accuracy (BSL) 1x Safety (BSL,2w)	-	-	●
2009 Im ³¹	RCT	US	Palpation guidance	Knee joint OA	45/44	injection (Hyaluronic acid, contrast dye)	knee joint	1x Accuracy (BSL)	-	-	●
2019 Lundstrom ¹⁸	Retrospective cohort study	US	Palpation guidance	Receiving knee injection	500/647	injection (hyaluronic acid formulas)	knee joint	2x Efficacy (up to 6y)	1x Efficacy (up to 6y)	(y 9 co)	●

Continued

Table 1 Continued

Year author	Study design	Imaging	Comparator	Disease	No Imaging/ Comparator	Intervention (COMPOUND)	SITE	FAVORS imaging*	FAVORS comparator*	No difference between imaging and comparator*	RBT
2020 Sheth ⁵⁷	RCT	US	Palpation guidance	Receiving knee injection	19/18	aspiration and injection (depomedrol, lidocaine)	knee joint	1x Safety (BSL, 4–6w) 6x Efficacy (BSL, 4–6w)	–	–	●
2019 Lee ⁵⁸	RCT	US	Palpation guidance	Pes anserinus tendinobursitis	27/27	injection (triamcinolone, lidocaine)	Pes anserinus bursa	1x Accuracy (BSL) 1x Safety (BSL, 1,4w)	–	1x Safety (up to 4 w)	●
US vs fluoroscopy											
2016 Peitschavage-Thomas ¹⁴	Retrospective cohort study	US	Fluo.	Anterior shoulder pain	51/50	injection (triamcinolone, lidocaine)	biceps tendon sheath	1x Accuracy (BSL)	–	2x Safety (1–3m, n.a.) 1x Cost/Time (n.a.)	
2014 Fowler ¹⁶	RCT	US	Fluo.	Piniformis syndrome	15/12	injection, nerve stimulation (lidocaine, triamcinolone)	piniformis muscle	–	–	2x Safety (BSL, 1,2w,3m) 2x Efficacy (BSL, 1,2w,3m) 1x Cost/Time (BSL, 1,2w,3m)	●
2016 Soneji ¹³	RCT	US	Fluo.	Chronic SJL arthritis	20/20	injection (methylprednisolone, bupivacaine, epinephrine)	SJL	–	–	1x Cost/Time (BSL)	●
2014 Gee ¹⁵	RCT	US	Fluo.	Non-inflamm. SJL dysfunction	55/55	injection (contrast media, lidocaine, dexamethasone)	SJL	–	1x Accuracy (BSL)	1x Safety (BSL, 3d,1w,1,3m) 2x Safety (BSL, 1,3m) 2x Efficacy (BSL, 1m)	●
2019 Kim ¹²	RCT	US	Fluo.	Chronic knee OA	38/33	injection, genicular nerve block (lidocaine, triamcinolone)	knee joint	–	–	2x Safety (BSL, 2–12w) 5x Efficacy (BSL, 3,6m)	●
US/fluoroscopy vs palpation guidance											
2016 Bossert ¹⁷	Retro spective cross sectional study	Fluo./US	Palpation guidance	Ankle OA	29/9/12	injection (viscosupplement)	ankle	2x Efficacy (BSL, n.a.)	–	1x Safety (n.a.) 3x Efficacy (n.a.)	
Fluoroscopy vs palpation guidance											
2019 Cohen ⁵³	RCT	Fluo.	Palpation guidance	Painful SJL	64/61	injection (dexamethylprednisolone, bupivacaine)	SJL	1x Accuracy (BSL) 2x Safety (1,3m)	–	3x Safety (BSL, 1,3m) 4x Efficacy (BSL, 1,3m)	
US vs fluoroscopy vs palpation guidance											
2020 Henne ⁵⁹	Retrospective cohort study	Fluo.	US, Palpation guidance	Hip OA or FAI	302/855	injection (corticosteroid, or hyaluronic acid)	Hip	–	–	2x Cost/Time (post-procedure)	
Fluoroscopy vs CT											
2020 Diffe ²⁴	retrospective cohort study	Fluo.	CT	pyogenic vertebral osteomyelitis	51/37	Biopsy	disco-vertebral tissue	1x Accuracy (BSL)	–	1x Safety (n.a.)	●

*These columns indicate the number of outcomes studied and their categories (e.g. 6 x efficacy) indicates that six different outcomes related to the efficacy category were studied, with the respective time points) when the outcome was determined (parenthesis). The term 'BSL' refers to differences concerning the outcomes at BL. A detailed description of studies and outcomes can be found in supplement.
TA, green, yellow, red and black circle indicate low, moderate, high and critical risk of bias, respectively. In case multiple circles are depicted, certain outcomes within one study had different risk of bias.
BSL, baseline; FAI, femoroacetabular impingement; n.a., not applicable; OA, osteoarthritis; PICO, Population, Intervention, Comparator, Outcome; RA, rheumatoid arthritis; RCT, randomised controlled trial; RbG, Risk of Bias; SJL, sacroiliac joint(s); US, ultrasound;



required similar or less time than procedures using US according to two studies (one high, one moderate RoB).^{13 16}

Small joints and periarticular structures

Seven studies (four RCTs, two retrospective cohort studies and one retrospective cross-sectional study) compared the value of imaging versus palpation-guided interventions at small joints: five concerned US,^{42 48–51} one US or CT⁵² and one US or fluoroscopy.⁶² The majority of studies were related to GC injections at^{48 51} or percutaneous release of^{42 49} the first annular (A1) pulley (n=4), while the effects of GC injections at the acromioclavicular joint (AC),⁵⁰ temporomandibular joint (TMJ)⁵² and the carpometacarpal joint of the thumb⁶² were investigated in one study each. A summary of study characteristics is depicted in table 2 while full details of the studies including all outcomes can be found in online supplemental table 5.

Accuracy

A single study with high RoB reported a higher accuracy (defined as the percentage of intra-articular needle placement) of US as compared with palpation-guided injections at the AC joint (96% vs 60.5%).⁵⁰

Safety

Safety outcomes were addressed in five studies. In two of them (one with moderate, one with high RoB), lower long-term pain levels were found in the groups receiving US-guided injections as compared with those receiving palpation-guided injections.^{49 50} The rate of other adverse events was comparable between groups. The other three studies (two with moderate and one with high RoB) found no difference concerning long-term pain or other adverse events when comparing US or US/CT with palpation-guided procedures.^{48 51 52}

Efficacy

One study with moderate RoB reported slightly higher patient satisfaction (4.1 ± 0.8 vs 3.7 ± 0.8 , based on a 5-point Likert scale with 1=worst, 5=best) when performing US-guided rather than palpation-guided injections,⁴⁸ while five studies (three moderate, two high RoB) revealed similar results concerning function, treatment failure and other efficacy outcomes when comparing US, and US or CT-guided interventions with palpation-guided interventions, as detailed in online supplemental table 5.^{42 48 49 51 52} One retrospective cohort study with high RoB described longer duration of treatment effects for palpation and fluoroscopy-guided injections as compared with US guidance at the carpometacarpal joint of the thumb.⁶²

Cost and time

One study with moderate RoB reported longer procedural time for TMJ injections using US or CT guidance as compared with palpation-guided injections,⁵² while in another study with high RoB, longer operational durations were found when performing an US-guided release of trigger finger using a needle-knife, as compared with

performing the procedure without imaging.⁴² One study (high RoB) described higher costs including physician and facility charges for US as compared with fluoroscopy and palpation-guided injections, however, the difference was not tested by statistical means.⁶²

Studies on multiple (small and large) joints and periarticular structures

Seven studies (four RCTs,^{34–37} one prospective cross-sectional study,²⁸ one retrospective cross-sectional study⁶⁰ and one retrospective cohort study³⁹) were identified that compared different methods to guide interventions at multiple (large and small, or unknown) joints. See table 3 and online supplemental table 6 for details. Five of them (all with moderate RoB)^{34–38} compared US with palpation-guided injections and/or aspirations. One study (with high RoB)⁶⁰ investigated the performance of US and CT to guide synovial biopsies, while one study (with critical RoB) compared US with palpation, or arthroscopic guidance for synovial tissue biopsies.³⁹

Accuracy

A better needle placement³⁶ and more successful aspirations³⁸ were observed in the US groups of two studies as compared with those undergoing palpation-guided injections, however, two other studies (one critical, one high RoB) reported similar performance of US versus palpation or arthroscopic guidance concerning the amount and quality of synovial tissue,³⁹ as well as US versus CT⁶⁰ for the yield of synovial tissue in the biopsy samples.

Safety

Two studies reported lower procedural and long-term pain in the US group as compared with palpation-guided interventions,^{35 37} while another study revealed no difference in pain between US and palpation guidance³⁶ (all moderate RoB).

Efficacy

A longer therapeutic effect after joint aspiration (defined by the time until the next intervention or a Visual Analogue Scale ≥ 2),³⁵ better quality of life³⁶ and a better treatment response³⁷ were found after US as compared with palpation-guided joint interventions. In one study, however, joint function was similar in both groups³⁶ (all moderate RoB).

Cost and time

In the USA, costs per year were higher for US as compared with palpation-guided interventions in a physician's office, while costs were similar in hospitals' outpatient clinic.³⁵

Nerves

Eight studies (seven RCTs,^{40–42 44–47 61} one prospective cohort study⁴³) compared US and palpation guidance for injections of the carpal tunnel syndrome. For details see table 4 and online supplemental table 7. Accuracy

Table 2 Studies comparing different imaging techniques or imaging versus palpation guidance at small joints and periartricular structures (PICO1)

Year author	STUDY DESIGN	Imaging	Comparator	Disease	No Imaging/ Comparator	Intervention (COMPOUND)	SITE	FAVORS imaging*	FAVORS comparator*	No difference between imaging and comparator*		RoB†
										No difference between imaging and comparator*	No difference between imaging and comparator*	
US vs palpation guidance												
2015 Park ³⁰	Retrospective cohort study	US	Palpation guidance	AC joint OA	50/50	injection (lidocaine, triamcinolone, contrast material)	AC joint	1 x Accuracy (BSL) 3 x Safety (3,6m)	–	1 x Safety (BSL,1,3,6m)	–	Red
2019 Roh ⁴⁸	RCT	US	Palpation guidance trigger finger	41/38	injection (triamcinolone)	finger flexor tendon sheath	–	2 x Safety (BSL,12,24w) 2 x Efficacy (BSL,12,24w)	–	–	Yellow	
2018 Lee ⁴⁹	RCT	US	Palpation guidance trigger finger	20, 23 fingers /21, 25 fingers	percutaneous pulley release (lidocaine)	A1 pulley	1 x Safety (2,4w)	–	–	1 x Safety (n.a.) 2 x Efficacy (BSL, n.a.)	Yellow	
2015 Cecen ⁵¹	RCT	US	Palpation guidance trigger finger	35/35	injection (methylprednisolone)	A1 pulley	–	–	–	2 x Safety (BSL,6w,6m) 2 x Efficacy (BSL,6w,6m)	Red	
2019 Pan ⁴²	RCT	US	Palpation guidance trigger finger (A1 pulley)	20/21	cutting (trigger finger release through needle knife)	A1 pulley	–	–	1 x Cost/Time (BSL)	1 x Efficacy (BSL)	Red	
US vs fluoroscopy vs palpation guidance												
2019 Gershkovich ⁶²	Retrospective cohort study	US	Fluo., Palpation guidance	Thumb CMC arthritis	8099/2533/51701	Injection (n.a.)	CMC joint of the thumb	–	–	1 x Efficacy for palpation guidance (n.a.) 1 x Efficacy for Fluo. (n.a.) 1 x Cost/Time (n.a.)	Red	
US/CT vs palpation guidance												
2017 Raenick ⁵²	Retrospective cross-sectional study	US or CT	Palpation guidance	JIA	22/23	injection (triamcinolone)	TMJ	–	–	2 x Cost/Time (BSL)	1 x Safety (n.a.) 2 x Efficacy (n.a.)	Yellow

*These columns indicate the number of outcomes studied and their categories (e.g. 6 x efficacy) indicates that six different outcomes related to the efficacy category were studied), with the respective time point(s) when the outcome was determined (parenthesis). The term 'BSL' refers to differences concerning the outcomes at BSL. A detailed description of studies and outcomes can be found in supplement.
TA, green; yellow, red and black circle indicate low, moderate, high and critical risk of bias, respectively. In case multiple circles are depicted, certain outcomes within one study had different risk of bias.

AC, acromioclavicular joint; BSLS, baseline; CMC, carpometacarpal; JIA, juvenile idiopathic arthritis; n.a., not applicable; OA, osteoarthritis; PICO, Population, Intervention, Comparator, Outcome; RCT, randomised controlled trial; RoB, Risk of Bias; TMJ, temporomandibular joint; US, ultrasound;

Table 3 Studies comparing different imaging techniques or imaging versus palpation guidance at multiple joints and periaricular structures (PICO1)

Year author	STUDY DESIGN	Imaging	Comparator	Disease	No Imaging/ Comparator	Intervention (COMPOUND)	SITE	FAVORS imaging*	FAVORS comparator*	No difference between imaging and comparator*	RoB†
US vs palpation guidance											
2011 Sibbitt ³⁵	RCT	US	Palpation guidance	Inflammatory Arthritis	124/120	aspiration	joints	1 x Safety (BSL, 2 w, 6 m) 2 x Efficacy (BSL to 12 m)	1 x Cost/Time (BSL to 12 m)	1 x Cost/Time (BSL to 12 m)	●
2010 Cunningham ³⁶	RCT	US	Palpation guidance	Inflammatory Arthritis	91/89	injection (triamcinolone, lidocaine, contrast agent iohexol)	joints	1 x Accuracy (BSL, 2 w) 1 x Efficacy (6 w)	1 x Efficacy (2 w)	1 x Safety (BSL, 2, 6 w) 3 x Efficacy (BSL, 2, 6 w)	●
2009 Sibbitt ³⁷	RCT	US	Palpation guidance	Painful arthritic joints (100 RA, 48 OA)	74/74	aspiration, injection (lidocaine, triamcinolone)	large and small joints	1 x Safety (BSL, 2 w) 2 x Efficacy (2 w)	–	–	●
2018 Nordberg ³⁴ of RCT	Secondary analysis	US	Palpation guidance	Early RA	118 (285 joints) /112 (338 joints)	injection (triamcinolone)	joints	–	–	1 x Efficacy (BSL, 12 m)	●
2002 Bainbridge ³⁸	Prospective cross-sectional study	US	Palpation guidance	Patients for joint/soft tissue aspiration	31/30	aspiration and injection	joints/soft tissue	1 x Accuracy (BSL)	–	1 x Accuracy (BSL)	●
US vs CT											
2020 McKee ⁶⁰	Retrospective cross-sectional study	US	CT	Receiving imaging-guided synovial biopsy	18/36	synovial biopsy	joints	–	–	1 x Accuracy (BSL)	●
US vs palpation guidance vs arthroscopy											
2018 Humby ³⁹	Retrospective cohort study	US-NB/ US-P&F	Palpation guidance/ Arthroscopy	Inflammatory arthritis	US-NB: 31 US-P&F: 46 Palpation guidance: 23 Arthroscopy: 25	Synovial biopsy	Small/ large joints	–	–	6 x Accuracy (BSL)	●

*These columns indicate the number of outcomes studied and their categories (eg, 6 x efficacy indicates that six different outcomes related to the efficacy category were studied), with the respective time points) when the outcome was determined (parenthesis). The term 'BSL' refers to differences concerning the outcomes at BSL. A detailed description of studies and outcomes can be found in supplement.
 †A green, yellow, red and black circle indicate low, moderate, high and critical risk of bias, respectively. In case multiple circles are depicted, certain outcomes within one study had different risk of bias.
 NB: needle biopsy; BSL, baseline; n.a., not applicable; OA, osteoarthritis; P&F, portal and forceps; PICO, Population, Intervention, Comparator, Outcome; RA, rheumatoid arthritis; RCT, randomised controlled trial; RoB, Risk of Bias; US, ultrasound.

Table 4 Studies comparing imaging versus palpation guidance at peripheral nerves (PICO1)

Year author	STUDY DESIGN	Imaging	Comparator	Disease	Intervention (COMPOUND)	SITE	FAVORS imaging*	FAVORS comparator*	No difference between imaging and comparator*	RoB†
US vs palpation guidance										
2017 Estamian ⁴⁵	RCT	US	Palpation guidance	CTS	27/20	Injection (methylprednisolone)	carpal tunnel	1x Efficacy (3m)	–	1x Safety (BSL,3 m) 1x Efficacy (BSL,3 m)
2019 Vandalpour ⁴⁰	RCT	US	Palpation guidance	CTS	29/23	injection (methylprednisolone)	carpal tunnel	1x Efficacy (4,12w)	1x Efficacy (4,12w)	2x Safety (BSL, 4, 12 w) 4x Efficacy (BSL, 4, 12 w)
2019 Roh ⁴¹	RCT	US	Palpation guidance	CTS	39/35	injection (lidocaine, triamcinolone)	carpal tunnel	1x Safety (4 w)	–	1x Safety (24 w) 2x Efficacy (BSL, 4, 12,24 w) 1x Efficacy (24w)
2018 Chen ⁴⁴	RCT	US	Palpation guidance	CTS	17 (22 wrists)/14 (17 wrists)	injection (betamethasone)	carpal tunnel	1x Safety (1 w)	–	4 x Safety (BSL,1 w,1,3,6 m) 20x Efficacy (BSL,1,3,6 m)
2013 Makhoul ⁴⁶	RCT	US	Palpation guidance	CTS	37 wrists /40 wrists	injection, hydrodissection (lidocaine, triamcinolone)	carpal tunnel	6x Safety (BSL,2 w6 m) 3x Efficacy (2w, 6 m,12 m)	1x Cost/Time (12m)	1x Safety (12m) 2 x Cost/Time (12m)
2018 Omar ⁴³	Prospective cohort study	US	Palpation guidance	CTS	15/15	injection (lidocaine, triamcinolone)	carpal tunnel	2x Efficacy (4w)	1 x Efficacy (4 w)	2 x Safety (BSL,4w) 8 x Efficacy (BSL,4w)
2013 Üstün ⁴⁷	RCT	US	Palpation guidance	CTS	23/23	injection (methylprednisolone)	carpal tunnel	1x Efficacy (6w)	–	2x Safety (BSL,6,12w) 1x Efficacy (BSL,6,12w)
2019 Rayegani ⁶¹	RCT	US	Palpation guidance	CTS	54/27	injection (triamcinolone, lidocaine)	carpal tunnel	–	–	3 x Safety (BSL,10w) 7 x Efficacy (BSL,10w)

*These columns indicate the number of outcomes studied and their categories (e.g. 6 x efficacy indicates that six different outcomes related to the efficacy category were studied), with the respective time point(s) when the outcome was determined (parenthesis). The term 'BSL' refers to differences concerning the outcomes at BSL. A detailed description of studies and outcomes can be found in supplement.

†A green, yellow, red and black circle indicate low, moderate, high and critical risk of bias, respectively. In case multiple circles are depicted, certain outcomes within one study had different risk of bias.
BSL, baseline; CTS, carpal tunnel syndrome; n.a., not applicable; PICO, Population, Intervention, Comparator, Outcome; RCT, randomised controlled trial; RoB, Risk of Bias; US, ultrasound; ;

outcomes were not assessed in these studies. No studies were found for interventions at nerves at other sites.

Safety

Procedural pain was assessed in one study (high RoB) favouring US over palpation-guided injections.⁴⁶ Long-term pain and symptom severity, mostly assessed by the Boston Carpal Tunnel Questionnaire, were similar in US and palpation groups according to six studies (four moderate, two high RoB),^{40 43–45 47 61} while higher symptom severity⁴¹ and long-term pain⁴⁶ in the palpation as compared with the US groups were observed in two studies (both with high RoB). Furthermore, one study with moderate RoB, reported a higher rate of hand weakness in patients who underwent palpation as compared with US-guided injections (18% vs 0%),⁴⁴ and another study with high RoB revealed higher cumulative adverse events including finger numbness, skin discolouration, subcutaneous fat atrophy and pain caused by steroid injection (steroid flares) in patients undergoing palpation (25%) as compared with the US-guided interventions (8%).⁴¹ Other adverse events did not differ between groups.^{43 46 47}

Efficacy

Hand function, quality of life and treatment response were similar between US and palpation-guided injections,^{41 43 44 47 61} while two studies described greater symptom relief⁴⁷ and longer duration until another intervention was required in the US groups.⁴⁶ Studies assessing electrophysiological parameters^{40 44 45 61} and median nerve thickness⁴³ revealed heterogeneous results, without a clear signal for any method.

Cost and time

Costs for US-guided injections in private practice and hospital based outpatient clinics were higher than for palpation guidance according to one study in the USA.⁴⁶ No data were available on cost-effectiveness.

Imaging settings and technical standards (PICO2)

Only three studies comparing different technical settings/equipment for imaging-guided interventions were found.^{25 63 64} Study characteristics and detailed results are depicted in table 5 and online supplemental table 8, respectively. According to one RCT with moderate RoB, a one-needle technique to dissolve calcific depositions in rotator cuff calcific tendinopathy resulted in a higher rate of needle obstruction (15% vs 3.6%) as compared with a two-needle method (one needle for injection, one for aspiration). Other outcomes such as the duration of the procedure, accuracy and safety were similar.⁶³

A second RCT with moderate RoB investigated the performance of a 25 mL mechanical as compared with a 60 mL automatic syringe for US-guided aspiration of knee effusion. No differences were found concerning pain or gathered amount of synovial fluid.²⁵

Just *et al* performed a retrospective cohort study (high RoB) to compare an US-guided needle—an US-guided

Year author	Study design	Setting	Alternative setting	Disease	No Setting/ Alternative setting	Intervention (COMPOUND)	SITE	FAVORS alternative setting*		No difference between settings* RoB†
								FAVORS setting*	FAVORS alternative setting*	
Large joints and perarticular structures										
2017 Orlandi ⁶³	RCT	One Needle	Two Needles	Rotator cuff calcific tendinopathy	100/111	US aspiration/ injection (heated saline, mepivacaine, tramcinolone)	subacromial/ subdeltoid bursa	–	2 x Accuracy (BSL)	1 x Accuracy (BSL) 3x Safety (BSL, 1,3m,1y) 1x Cost/Time (BSL)
2012 Sibbitt ²⁵	RCT	25mL mechanical syringe	60mL automatic syringe	knee effusion in rheumatoid arthritis, osteoarthritis	20/22	US joint aspiration/ injection (lidocaine, tramcaine, tramcinolone)	knee joint	–	–	1 x Accuracy (BSL) 1 x Safety (BSL,2w)
Multiple joints										
2018 Just ⁶⁴	Retrospective cohort study	US-NB	US-P&F/ Arthroscopy	Inflammatory arthritis, degenerative disease	US-NB: 402/ US-P&F: 65	Synovial biopsy	wrist, knee, MCP, ankle, elbow, MTP	–	1 x Efficacy (n.a.) 2x Safety (BSL,7-14d) 2x Efficacy (BSL-14d)	C

*These columns indicate the number of outcomes studied and their categories (eg, 6 x efficacy indicates that six different outcomes related to the efficacy category were studied, with the respective time point(s) when the outcome was determined (parenthesis)). The term 'BSL' refers to differences concerning the outcomes at BSL. A detailed description of studies and outcomes can be found in supplement.

†A green, yellow, red and black circle indicate low, moderate, high and critical risk of bias, respectively. In case multiple circles are depicted, certain outcomes within one study had different risk of bias. BSL, baseline; MCP, metacarpophalangeal joints; MTP, metatarsophalangeal joints; n.a., not applicable; NB, needle biopsy; P&F, portal and forceps; PICO, Population, Intervention, Comparator, Outcome; RCT, randomised controlled trial; RoB, Risk of Bias; US, ultrasound.

portal and forceps—and an arthroscopic approach for synovial tissue biopsy at multiple joints. They observed similar safety (pain and adverse events) and efficacy outcomes in all three groups. Slightly more patients in the portal and forceps as well as in the arthroscopic groups reported that they would eventually accept further synovial biopsies.⁶⁴ Studies comparing different settings to perform interventional procedures were not found.

Procedural protocols for imaging-guided interventions (PICO3)

Sixteen studies were found (nine on large, three on small joints and periarticular structures and four on nerves) comparing either different procedures and puncture sites (eg, in-of-plain US and out-of-plain US, injection in different knee recesses),^{22 61 67–72 74 78} or different targets (eg, periarticular vs intra-articular) for imaging-guided interventions.^{65 66 73 75–77} Table 6 and online supplemental table 9 contain an overview of studies and detailed results, respectively.

Large joints and periarticular structures

Shoulder

Two studies (one RCT with high⁶⁷ and one RCT with low⁶⁸ RoB) compared glenohumeral joints with the subacromial/subdeltoid bursa as targets for US-guided GC injections in patients with adhesive capsulitis/frozen shoulder. Patients reported higher pain levels,^{67 68} as well as increased symptom severity⁶⁸ after injections into the subacromial/subdeltoid bursa. Shoulder extension and external rotation, however, were better in the subacromial/subdeltoid bursa group in one study,⁶⁷ whereas in the other study, internal rotation and scores according to the American Shoulder and Elbow Surgeon scale were better when the glenohumeral joint was injected.⁶⁸ Another RCT (moderate RoB) focusing on patients with scapular pain and comparing injections either in the subscapular muscle or in the scapulothoracic bursa reported no differences between the groups according to safety and efficacy outcomes.²²

Spine and SIJ

A retrospective cross-sectional study (high RoB) investigated different targets at the spine when using CT to retrieve tissue samples in patients with infectious spondylodiscitis/osteomyelitis. No differences in quality and quantity of samples were observed when either the bone, disc or paravertebral tissue was targeted.⁶⁹ Two prospective cohort studies (both high RoB) compared the value of injecting GC intra-articularly or periarticularly at the SIJ via CT⁶⁵ or US⁶⁶ guidance. The CT study reported higher pain levels at 3 and 6 months in case of periarticular injections, while the US study concluded that there was no difference in pain 4 weeks after the injections.

Knee

In-of-plain US and out-of-plain US was compared in patients undergoing knee injections in one RCT with

moderate RoB. The study revealed no differences concerning accuracy, safety and duration of the intervention between groups.⁷¹ Furthermore, no differences were found for accuracy and safety when US-guided knee injections were conducted at the medial/lateral, midlateral or superolateral recess according to two RCTs with moderate RoB each.^{70 72}

Small joints and periarticular structures

US-guided periarticular injections of the AC joint resulted in lower pain reduction when performing the crossover arm test as compared with intra-articular injection, according to one RCT with low RoB. Other pain assessments (eg, pain at night or pain after pressure) and the amount of other adverse events were similar between groups.⁷⁷ Intra and extra tendon sheath injections at the A1 pulley were compared in two RCTs with high⁷⁶ and moderate⁷⁵ RoB. No differences concerning safety and efficacy outcomes were found between groups in both studies.

Nerves

Two RCT on US-guided injections in CTS, both with moderate RoB, investigated the performance of drug release above or below the median nerve,⁷⁴ or an ulnar versus a midline approach for injection.⁶¹ No differences between groups were found for safety and efficacy outcomes including long-term pain, other adverse events, function and electrophysiological parameters. Another study with moderate RoB reported less long-term pain, better function and less swelling of the median nerve after injection of the carpal tunnel from the radial rather than from the ulnar site of the wrist.⁷⁸ Furthermore, one retrospective cohort study (high RoB) assessed whether there were differences in safety or efficacy outcomes when the needle was placed in the intraepineurium as compared with the extraepineurium, when performing US-guided hydrodissection and GC injection of the median nerve in CTS.⁷³ The results for safety and clinical efficacy outcomes like function and symptom severity were comparable between groups, while patients receiving intraepineurial injections yielded a higher reduction of perineurial oedema 2 weeks after the intervention as compared with those who underwent extraepineurial interventions.

DISCUSSION

Overall, we found that imaging (especially US)-guided interventions at joints and periarticular structures were reported to lead to more accurate needle positioning as compared with palpation-guided interventions in the majority of studies. The studies themselves, as well as results for safety and efficacy were heterogeneous, and data on cost-effectiveness were absent. Another observation is that the quality of studies was generally low: 51% had a moderate and 47% a high or critical RoB assessment.

Table 6 Procedural protocols for imaging-guided interventions (PICo3)

Year author	Study design	Procedure	Alternative procedure	Disease	Intervention (COMPOUND)	SITE	FAVORS procedure*	FAVORS alternative procedure*	No difference between procedures*	RoB†
Large joints and perarticular structures										
2018 Khalil ⁶⁷	RCT	GH approach	SASD approach	adhesive capsulitis	20/20	US injection (methylprednisolone, lidocaine)	shoulder	1 x Safety (12w)	2 x Efficacy (12w)	1 x Safety (BSL_12w) 3 x Efficacy (BSL_12w)
2016 Cho ⁶⁸	RCT	Intra-articular	Subacromial Bursa or intra-articular and bursa	Frozen shoulder	36/37	US injection (triamcinolone, lidocaine)	GH/Subacromial bursa/both	1 x Safety (BSL_3,6,12w) 2 x Efficacy (BSL_3,6,12w)	1 x Safety (BSL_3,6,12w)	3 x Efficacy (BSL_3,6,12w)
2014 Chang ²²	RCT	subscapular muscle	scapulothoracic bursa	patients with scapular pain	18/18	Subscapular muscle: US injection Scapulothoracic bursa: palpation guidance injection (lidocaine, triamcinolone)	subscapularis muscle/ scapulothoracic bursa	–	–	2 x Safety (BSL_1,2,3,12w) 1 x Efficacy (1,12w)
2015 Chang ⁶⁹	Retrospective cross sectional study	Bone (Endplate) Biopsy	Disk Biopsy/ Paravertebral soft tissue Biopsy	Diskitis osteomyelitis	102	CT Biopsy	Spine	–	–	2 x Accuracy (BSL)
2015 Althoff ⁶⁵	Prospective cohort study	intra-articular	peri-articular	Sacroiliitis	23/7	CT injection (triamcinolone)	SIJ	1 x Safety (3,6m)	–	–
2009 Hartung ⁶⁶	Prospective cohort study	Intra-articular	Periarticular	Acute sacroiliitis	8/12 injections	US Injection (triamcinolone, gadolinium)	SIJ	–	–	1 x Safety (BSL_24h,4w)
2013 Jang ⁷¹	RCT	Out of plain US	In plain US	knee OA	41/44	Injection (lidocaine, triamcinolone, contrast agent)	knee joint	–	–	1 x Accuracy (BSL) 1 x Safety (BSL)
2011 Park ⁷²	RCT	Medial Portal	Midlateral Portal/ Superolateral Portal	Knee OA	40/43/43 knees	US injection (lidocaine, triamcinolone, contrast agent)	knee joint	–	–	1 x Cost/Time (BSL)
2013 Park ⁷⁰	RCT	Midmedial Portal	Midlateral Portal/ Superolateral Portal	Knee OA	40/40/40 knees	US injection (contrast medium, lidocaine, triamcinolone)	knee joint	–	–	1 x Accuracy (BSL) 1 x Safety (n.a, no difference between any group)
Small Joints and Periarticular Structures										
2013 Sabeti-Aschraf ⁷⁷	RCT	Intra-articular injection	Periarticular injection	AC joint arthralgia	53/53 shoulders	US Injection (lidocaine, betamethasone)	AC joints	1 x Safety (BSL_1h,1,3w)	–	4 x Safety (BSL_1h,1,3w)
2016 Shinomiya ⁷⁸	RCT	Intra sheath	Extra sheath	Trigger finger	27/29 fingers)/28 (32 fingers)	Injection (triamcinolone, mepivacaine)	A1 pulley middle/ ring fingers	–	–	2 x Safety (BSL_4w) 5 x Efficacy (BSL_4w)
2018 Mardani-Kivi ⁷⁵	RCT	Intra sheath	Extra sheath	trigger finger	83/83	US Injections (methylprednisolone, lidocaine)	A1 pulley flexor tendons	–	–	1 x Safety (BSL_3,6,12,24,48w) 1 x Efficacy (BSL_3,6,12,24,48w)
Nerves										
2018 Babaei-Ghazani ⁷⁴	RCT	Above Median nerve	Below Median nerve	CTS	22/22	US injection (triamcinolone)	carpal tunnel	–	–	3 x Safety (BSL_6w,12w) 6 x Efficacy (BSL_6w,12w)
2018 Hsu ⁷³	Retrospective cohort study	Intraepineurium	Extraneurium	CTS	39/62 wrists	US hydrodissection, injection (lidocaine, triamcinolone)	carpal tunnel	1 x Safety (2w) 4 x Efficacy (BSL_2w)	–	2 x Safety (BSL_2w) 3 x Efficacy (2w)
2019 Rayegani ⁶¹	RCT	US ulnar in plane	US midline in plane	CTS	27/27	injection (triamcinolone, lidocaine)	carpal tunnel	–	–	3 x Safety (BSL_10w) 7 x Efficacy (BSL_10w)

Continued

Table 6 Continued

Year author	Study design	Procedure	Alternative procedure	Disease	Intervention (COMPOUND)	SITE	FAVORS procedure*	FAVORS alternative procedure*	No difference between procedures*	RoB†
2020 Babaei-Ghazani ⁷⁸	RCT	US radial approach	US ulnar approach	CTS	30/30 injection (triamcinolone)	carpal tunnel	1x Safety (12w) 5x Efficacy (2,6,12w)	1x Efficacy (12w)	3x Efficacy (BSL, 2,6,12w)	●

*These columns indicate the number of outcomes studied and their categories (eg, 6 x efficacy indicates that six different outcomes related to the efficacy category were studied), with the respective time point(s) when the outcome was determined (parenthesis). The term BSL refers to differences concerning the outcomes at BSL. A detailed description of studies and outcomes can be found in supplement.
†A green, yellow, red and black circle indicate low, moderate, high and critical risk of bias, respectively. In case multiple circles are depicted, certain outcomes within one study had different risk of bias.
AC, acromioclavicular; BSL, baseline; CTS, carpal tunnel syndrome; GH, glenohumeral joint; n.a., not applicable; OA, osteoarthritis; RCT, randomised controlled trial; RoB, Risk of Bias; SASD, subacromial-subdeltoid; SJS, sacroiliac joint(s); US, ultrasound.

Positioning a needle or another instrument in, rather than outside the target not only reduces the risk of adverse events,^{41–44} but may also have positive effects on short-term and long-term pain, as well as on efficacy outcomes.^{65–77}

Studies comparing different imaging techniques such as US and fluoroscopy were scarce, which is related to the fact that centres usually have expertise with one rather than multiple techniques.¹

Only a few studies were available on different technical settings/equipment for imaging-guided procedures,^{25–63,64} addressing the value of different equipment, but not the role of assistance by healthcare professionals, monitoring of vital signs, the use of contrast agents or different types of aseptic/antiseptic conditions. Similarly, studies considering different levels of experience of the healthcare provider performing the intervention have not been conducted. It would for example be important to know the accuracy of a palpation-guided intervention of an experienced clinician as compared with a imaging-guided procedure of a fellow.

A Delphi-based consensus paper series on clinical indications for imaging-guided interventional procedures in the musculoskeletal system has recently been published by the European Society of Musculoskeletal Radiology.^{5–7} While this series presented important general statements on the use and the value of imaging (eg, ‘US guidance is a safe and effective method for brachial plexus block’),⁶ the present work focused on direct comparisons between methods, settings and procedures when performing imaging-guided interventions (eg, US vs fluoroscopy, or intra-articular vs periarticular injections). Certain studies only assessing one imaging method, setting or procedure are therefore not covered in this review.^{79–87}

One of the major limitations of the available studies was their low scientific quality. Missing information on the randomisation process (for RCTs), outcome assessment (RCTs and non-randomised studies) and the population of interest (cross-sectional studies) were the main reasons for moderate/high RoB of the studies retrieved. Potential confounders were also observed in a small number of non-randomised studies, which was mainly due to their retrospective design. Prospective study design as well as adequate statistical models accounting for confounding factors such as the baseline values of certain outcomes may in future lead to more robust results.

Another limitation was the heterogeneity of study designs and outcomes which prevented us to meta-analyse the available data. We nevertheless tried to categorise outcomes according to overarching groups (accuracy, efficacy, safety and cost-effectiveness) in order to facilitate comparison between studies and to enable overall conclusions. Some outcomes could have been assigned to several of these categories (eg, pain was reported under ‘safety’, even though it could also be considered an efficacy outcome), and we tried to be as consistent as possible when categorising the outcomes into these broader groups in order to provide the best information available to the EULAR task force and the scientific community.



In conclusion, the majority of studies indicated a higher accuracy of needle positioning at joints and periarticular structures when using imaging (especially US) guidance as compared with palpation-guided interventions with the limitation of heterogeneity of data and considerable RoB. Heterogeneity was also observed among studies on safety and efficacy, whereas data on cost-effectiveness were virtually absent.

Author affiliations

- ¹Department of Rheumatology and Immunology, Medical University of Graz, Graz, Austria
- ²Internal Medicine and Nephrology Unit, Department of Life, Health and Environmental Sciences, University of L'Aquila, L'Aquila, Italy
- ³Department of Medicine, ASL 1 Avezzano-Sulmona-L'Aquila, San Salvatore Hospital, L'Aquila, Italy
- ⁴University of Ferrara, Ferrara, Italy
- ⁵Rheumazentrum Ruhrgebiet Herne, Ruhr University Bochum, Bochum, Germany
- ⁶Northwell Health, New York, New York, USA
- ⁷Rheumatology Service, Hospital of Bruneck, Bruneck, Italy
- ⁸Centre for Rheumatology, University College London, London, UK
- ⁹National Institute for Health Research (NIHR) University College London Hospitals Biomedical Research Centre, University College London Hospitals NHS Foundation Trust, London, UK
- ¹⁰Department of Rheumatology, Northwick Park Hospital, London North West University Healthcare NHS Trust, London, UK

Twitter Pedro M Machado @pedrommcmachado

Contributors PB and FC drafted the first version of the manuscript. LF designed the search terms for the literature search. All authors contributed, revised critically and finally approved the final version of the manuscript. PB acted as guarantor.

Funding This work was funded by the European Alliance of Associations for Rheumatology (IMG016). PMM is supported by the National Institute for Health Research (NIHR), University College London Hospitals (UCLH), Biomedical Research Centre (BRC).

Competing interests PMM has received consulting/speaker's fees from Abbvie, BMS, Celgene, Eli Lilly, Galapagos, Janssen, MSD, Novartis, Orphazyme, Pfizer, Roche and UCB, all unrelated to this manuscript. FC has received consulting/speaker's fees from Abbvie and Celgene, all unrelated to this manuscript. XB has received consulting/speaker's fees from Abbvie, BMS, Celgene, Galapagos, Gilead, Eli Lilly, Janssen, MSD, Novartis, Pfizer and UCB, all unrelated to this manuscript. CD has received consulting/speaker's fees from Abbvie, Eli Lilly, Janssen, Novartis, Pfizer, Roche and Sanofi, all unrelated to this manuscript.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplemental information. Not applicable.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

- Philipp Bosch <http://orcid.org/0000-0002-6783-6422>
 Francesco Carubbi <http://orcid.org/0000-0003-1958-5136>
 Carlo Alberto Scirè <http://orcid.org/0000-0001-7451-0271>
 Xenofon Baraliakos <http://orcid.org/0000-0002-9475-9362>
 Christian Dejaco <http://orcid.org/0000-0002-0173-0668>
 Pedro M Machado <http://orcid.org/0000-0002-8411-7972>

REFERENCES

- 1 Carubbi F, Bosch P, Machado PM, et al. AB1083 CURRENT PRACTICE AND OPINIONS ON IMAGING-GUIDED INTERVENTIONAL PROCEDURES IN RHEUMATIC AND MUSCULOSKELETAL DISEASES: INTERIM RESULTS OF A MULTINATIONAL MULTIDISCIPLINARY SURVEY TO INFORM EULAR POINTS TO CONSIDER. *Ann Rheum Dis* 2020;79:1830.2-1.
- 2 Maricar N, Parkes MJ, Callaghan MJ, et al. Where and how to inject the knee—a systematic review. *Semin Arthritis Rheum* 2013;43:195–203.
- 3 Daley EL, Bajaj S, Bisson LJ, et al. Improving injection accuracy of the elbow, knee, and shoulder: does injection site and imaging make a difference? A systematic review. *Am J Sports Med* 2011;39:656–62.
- 4 Zabotti A, Bandinelli F, Batticciotto A, et al. Musculoskeletal ultrasonography for psoriatic arthritis and psoriasis patients: a systematic literature review. *Rheumatology* 2017;56:1518–32.
- 5 Sconfienza LM, Adriaensen M, Albano D, et al. Clinical indications for image-guided interventional procedures in the musculoskeletal system: a Delphi-based consensus paper from the European Society of Musculoskeletal Radiology (ESSR)-part I, shoulder. *Eur Radiol* 2020;30:903–13.
- 6 Sconfienza LM, Adriaensen M, Albano D, et al. Clinical indications for image guided interventional procedures in the musculoskeletal system: a Delphi-based consensus paper from the European Society of Musculoskeletal Radiology (ESSR)-part III, nerves of the upper limb. *Eur Radiol* 2020;30:1498–506.
- 7 Sconfienza LM, Adriaensen M, Albano D, et al. Clinical indications for image-guided interventional procedures in the musculoskeletal system: a Delphi-based consensus paper from the European Society of Musculoskeletal Radiology (ESSR)-Part II, elbow and wrist. *Eur Radiol* 2020;30:2220–30.
- 8 Sackett D, Richardson W, Rosenberg W. *How to practice and teach EBM*, 2005: 2.
- 9 Sterne JAC, Savović J, Page MJ, et al. Rob 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019;366:i4898.
- 10 Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016;355:i4919.
- 11 Downes MJ, Brennan ML, Williams HC, et al. Development of a critical appraisal tool to assess the quality of cross-sectional studies (axis). *BMJ Open* 2016;6:e011458.
- 12 Kim D-H, Lee M-S, Lee S, et al. A prospective randomized comparison of the efficacy of Ultrasound- vs Fluoroscopy-Guided Genicular nerve block for chronic knee osteoarthritis. *Pain Physician* 2019;22:139–46.
- 13 Soneji N, Bhatia A, Seib R, et al. Comparison of fluoroscopy and ultrasound guidance for Sacroiliac joint injection in patients with chronic low back pain. *Pain Pract* 2016;16:537–44.
- 14 Petscavage-Thomas J, Gustas C. Comparison of ultrasound-guided to Fluoroscopy-Guided biceps tendon sheath therapeutic injection. *J Ultrasound Med* 2016;35:2217–21.
- 15 Jee H, Lee J-H, Park KD, et al. Ultrasound-Guided versus fluoroscopy-guided sacroiliac joint intra-articular injections in the noninflammatory sacroiliac joint dysfunction: a prospective, randomized, single-blinded study. *Arch Phys Med Rehabil* 2014;95:330–7.
- 16 Fowler IM, Tucker AA, Weimerskirch BP, et al. A randomized comparison of the efficacy of 2 techniques for piriformis muscle injection: ultrasound-guided versus nerve stimulator with fluoroscopic guidance. *Reg Anesth Pain Med* 2014;39:126–32.
- 17 Bossert M, Bouibil D, Parisaux J-M, et al. Imaging guidance improves the results of Viscosupplementation with HANOX-M-XL in patients with ankle osteoarthritis: results of a clinical survey in 50 patients treated in daily practice. *Clin Med Insights Arthritis Musculoskelet Disord* 2016;9:195–9.
- 18 Lundstrom ZT, Sytsma TT, Greenlund LS. Rethinking Viscosupplementation: Ultrasound- versus Landmark-Guided injection for knee osteoarthritis. *J Ultrasound Med* 2020;39:113–7.
- 19 Mitchell WG, Kettwich SC, Sibbitt WL, et al. Outcomes and cost-effectiveness of ultrasound-guided injection of the trochanteric bursa. *Rheumatol Int* 2018;38:393–401.
- 20 Raeissadat SA, Rayegani SM, Langrudi TF, et al. Comparing the accuracy and efficacy of ultrasound-guided versus blind injections of steroid in the glenohumeral joint in patients with shoulder adhesive capsulitis. *Clin Rheumatol* 2017;36:933–40.
- 21 Saeed A, Khan M, Morrissey S, et al. Impact of outpatient clinic ultrasound imaging in the diagnosis and treatment for shoulder impingement: a randomized prospective study. *Rheumatol Int* 2014;34:503–9.
- 22 Chang WH, Kim YW, Choi S, et al. Comparison of the therapeutic effects of intramuscular subscapularis and scapulothoracic bursa injections in patients with scapular pain: a randomized controlled trial. *Rheumatol Int* 2014;34:1203–9.

- 23 Kim TK, Lee JH, Park KD, et al. Ultrasound versus palpation guidance for intra-articular injections in patients with degenerative osteoarthritis of the elbow. *J Clin Ultrasound* 2013;41:479–85.
- 24 Jang SH, Lee SC, Lee JH, et al. Comparison of ultrasound (US)-guided intra-articular injections by in-plain and out-of-plain on medial portal of the knee. *Rheumatol Int* 2013;33:1951–9.
- 25 Sibbitt WL, Kettwich LG, Band PA, et al. Does ultrasound guidance improve the outcomes of Arthrocentesis and corticosteroid injection of the knee? *Scand J Rheumatol* 2012;41:66–72.
- 26 Bum Park Y, Ah Choi W, Kim Y-K, et al. Accuracy of blind versus ultrasound-guided suprapatellar bursal injection. *J Clin Ultrasound* 2012;40:20–5.
- 27 Zhang J, Ebraheim N, Lause GE. Ultrasound-Guided injection for the biceps brachii tendinitis: results and experience. *Ultrasound Med Biol* 2011;37:729–33.
- 28 Sibbitt WL, Band PA, Kettwich LG, et al. A randomized controlled trial evaluating the cost-effectiveness of sonographic guidance for intra-articular injection of the osteoarthritic knee. *J Clin Rheumatol* 2011;17:409–15.
- 29 Hashiuchi T, Sakurai G, Morimoto M, et al. Accuracy of the biceps tendon sheath injection: ultrasound-guided or unguided injection? A randomized controlled trial. *J Shoulder Elbow Surg* 2011;20:1069–73.
- 30 Lee H-J, Lim K-B, Kim D-Y, et al. Randomized controlled trial for efficacy of intra-articular injection for adhesive capsulitis: ultrasonography-guided versus blind technique. *Arch Phys Med Rehabil* 2009;90:1997–2002.
- 31 Im SH, Lee SC, Park YB, et al. Feasibility of sonography for intra-articular injections in the knee through a medial Patellar portal. *J Ultrasound Med* 2009;28:1465–70.
- 32 Luz KR, Furtado RNV, Nunes CCG, et al. Ultrasound-Guided intra-articular injections in the wrist in patients with rheumatoid arthritis: a double-blind, randomised controlled study. *Ann Rheum Dis* 2008;67:1198–200.
- 33 Naredo E, Cabero F, Beneyto P, et al. A randomized comparative study of short term response to blind injection versus sonographic-guided injection of local corticosteroids in patients with painful shoulder. *J Rheumatol* 2004;31:308–14.
- 34 Nordberg LB, Lillegren S, Aga A-B, et al. The impact of ultrasound on the use and efficacy of intraarticular glucocorticoid injections in early rheumatoid arthritis: secondary analyses from a randomized trial examining the benefit of ultrasound in a clinical tight control regimen. *Arthritis Rheumatol* 2018;70:1192–9.
- 35 Sibbitt WL, Band PA, Chavez-Chiang NR, et al. A randomized controlled trial of the cost-effectiveness of ultrasound-guided intraarticular injection of inflammatory arthritis. *J Rheumatol* 2011;38:252–63.
- 36 Cunningham J, Marshall N, Hide G, et al. A randomized, double-blind, controlled study of ultrasound-guided corticosteroid injection into the joint of patients with inflammatory arthritis. *Arthritis Rheum* 2010;62:1862–9.
- 37 Sibbitt WL, Peisajovich A, Michael AA, et al. Does sonographic needle guidance affect the clinical outcome of intraarticular injections? *J Rheumatol* 2009;36:1892–902.
- 38 Balint PV, Kane D, Hunter J. Ultrasound guided versus conventional joint and soft tissue fluid aspiration in rheumatology practice: a pilot study. *J Rheumatol* 2002;29:2209–13.
- 39 Humby F, Romão VC, Manzo A, et al. A multicenter retrospective analysis evaluating performance of synovial biopsy techniques in patients with inflammatory arthritis: arthroscopic versus ultrasound-guided versus blind needle biopsy. *Arthritis Rheumatol* 2018;70:702–10.
- 40 Vahdatpour B, Haghight S, Azimi Z. Carpal tunnel syndrome treatment using ultrasound-guided versus landmark-guided corticosteroid injection: a randomized prospective trial. *Shiraz E Medical Journal* 2019;20.
- 41 Roh YH, Hwangbo K, Gong HS, et al. Comparison of ultrasound-guided versus Landmark-Based corticosteroid injection for carpal tunnel syndrome: a prospective randomized trial. *J Hand Surg Am* 2019;44:304–10.
- 42 Pan M, Sheng S, Fan Z, et al. Ultrasound-Guided percutaneous release of A1 Pulley by using a needle knife: a prospective study of 41 cases. *Front Pharmacol* 2019;10:267–67.
- 43 Omar G, Ali F, Ragaae A, et al. Ultrasound-Guided injection of carpal tunnel syndrome: a comparative study to blind injection. *The Egyptian Rheumatologist* 2018;40:131–5.
- 44 Chen P-C, Wang L-Y, Pong Y-P, et al. Effectiveness of ultrasound-guided vs direct approach corticosteroid injections for carpal tunnel syndrome: a double-blind randomized controlled trial. *J Rehabil Med* 2018;50:200–8.
- 45 Eslamian F, Eftekharadsat B, Babaei-Ghazani A, et al. A randomized prospective comparison of ultrasound-guided and Landmark-Guided steroid injections for carpal tunnel syndrome. *J Clin Neurophysiol* 2017;34:107–13.
- 46 Makhlof T, Emil NS, Sibbitt WL, et al. Outcomes and cost-effectiveness of carpal tunnel injections using sonographic needle guidance. *Clin Rheumatol* 2014;33:849–58.
- 47 Ustün N, Tok F, Yagz AE, et al. Ultrasound-Guided vs. blind steroid injections in carpal tunnel syndrome: a single-blind randomized prospective study. *Am J Phys Med Rehabil* 2013;92:999–1004.
- 48 Roh YH, Kim S, Gong HS, et al. Retracted: a randomized comparison of ultrasound-guided versus landmark-based corticosteroid injection for trigger finger. *J Hand Surg Eur Vol* 2020;45:NP1–6.
- 49 Lee SH, Choi YC, Kang HJ. Comparative study of ultrasonography-guided percutaneous A1 pulley release versus blinded percutaneous A1 pulley release. *J Orthop Surg* 2018;26:2309499018772368.
- 50 Park KD, Kim TK, Lee J, et al. Palpation versus ultrasound-guided acromioclavicular joint intra-articular corticosteroid injections: a retrospective comparative clinical study. *Pain Physician* 2015;18:333–41.
- 51 Cecen GS, Gulabi D, Saglam F, et al. Corticosteroid injection for trigger finger: blinded or ultrasound-guided injection? *Arch Orthop Trauma Surg* 2015;135:125–31.
- 52 Resnick CM, Vakilian PM, Kaban LB, et al. Is intra-articular steroid injection to the temporomandibular joint for juvenile idiopathic arthritis more effective and efficient when performed with image guidance? *J Oral Maxillofac Surg* 2017;75:694–700.
- 53 Cohen SP, Bicket MC, Kurihara C, et al. Fluoroscopically guided vs Landmark-Guided Sacroiliac joint injections: a randomized controlled study. *Mayo Clin Proc* 2019;94:628–42.
- 54 Difre C, Jousset C, Roux A-L, et al. Predictive factors for positive disco-vertebral biopsy culture in pyogenic vertebral osteomyelitis, and impact of fluoroscopic versus scanographic guidance. *BMC Infect Dis* 2020;20:512.
- 55 Cankurtaran D, Karaahmet OZ, Yildiz SY, et al. Comparing the effectiveness of ultrasound guided versus blind genicular nerve block on pain, muscle strength with isokinetic device, physical function and quality of life in chronic knee osteoarthritis: a prospective randomized controlled study. *Korean J Pain* 2020;33:258–66.
- 56 Lee JH, Lee JU, Yoo SW. Accuracy and efficacy of ultrasound-guided PEs anserinus bursa injection. *J Clin Ultrasound* 2019;47:77–82.
- 57 Sheft T, Miranda OM, Johnson B. Assessment of patient satisfaction, functionality, and quality of life after ultrasound-guided knee intervention: a prospective study. *Clin Rheumatol* 2021;40:735–40.
- 58 Yiannakopoulos CK, Megaloikonomos PD, Foufa K, et al. Ultrasound-Guided versus palpation-guided corticosteroid injections for tendinosis of the long head of the biceps: a randomized comparative study. *Skeletal Radiol* 2020;49:585–91.
- 59 Henne M, Centurion A, Zeini IM, et al. Trends in utilization of image guidance for hip joint injections. *Clin J Sport Med* 2021;31:374–8.
- 60 McKee TC, Belair JA, Sobol K, et al. Efficacy of image-guided synovial biopsy. *Skeletal Radiol* 2020;49:921–8.
- 61 Rayegani SM, Raeissadat SA, Ahmadi-Dastgerdi M, et al. Comparing the efficacy of local triamcinolone injection in carpal tunnel syndrome using three different approaches with or without ultrasound guidance. *J Pain Res* 2019;12:2951–8.
- 62 Gershkovich GE, Boyadjian H, Conti Mica M. The effect of image-guided corticosteroid injections on thumb Carpometacarpal arthritis. *Hand* 2021;16:1558944719846572.
- 63 Orlandi D, Mauri G, Lacelli F, et al. Rotator cuff calcific tendinopathy: randomized comparison of US-guided percutaneous treatments by using one or two needles. *Radiology* 2017;285:518–27.
- 64 Just SA, Humby F, Lindegaard H, et al. Patient-Reported outcomes and safety in patients undergoing synovial biopsy: comparison of ultrasound-guided needle biopsy, ultrasound-guided portal and forceps and arthroscopic-guided synovial biopsy techniques in five centres across Europe. *RMD Open* 2018;4:e000799.
- 65 Althoff CE, Bollow M, Feist E, et al. Ct-Guided corticosteroid injection of the sacroiliac joints: quality assurance and standardized prospective evaluation of long-term effectiveness over six months. *Clin Rheumatol* 2015;34:1079–84.
- 66 Hartung W, Ross CJ, Straub R, et al. Ultrasound-Guided sacroiliac joint injection in patients with established sacroiliitis: precise la injection verified by MRI scanning does not predict clinical outcome. *Rheumatology* 2010;49:1479–82.
- 67 Khalaf SF, Hussein MI, El-Barbary AM, et al. Efficacy of ultrasonography-guided intra-articular steroid injection of the shoulder and excercising in patients with adhesive capsulitis:

- Glenohumeral versus subacromial approaches. *The Egyptian Rheumatologist* 2018;40:277–80.
- 68 Cho C-H, Kim DH, Bae K-C, et al. Proper site of corticosteroid injection for the treatment of idiopathic frozen shoulder: results from a randomized trial. *Joint Bone Spine* 2016;83:324–9.
- 69 Chang CY, Simeone FJ, Nelson SB, et al. Is Biopsying the paravertebral soft tissue as effective as Biopsying the disk or vertebral endplate? 10-year retrospective review of CT-guided biopsy of Diskitis-Osteomyelitis. *AJR Am J Roentgenol* 2015;205:123–9.
- 70 Park KD, Ahn JK, Lee S-C, et al. Comparison of ultrasound-guided intra-articular injections by long axis in plane approach on three different sites of the knee. *Am J Phys Med Rehabil* 2013;92:990–8.
- 71 Jang SH, Lee SC, Lee JH, et al. Comparison of ultrasound (US)-guided intra-articular injections by in-plain and out-of-plain on medial portal of the knee. *Rheumatol Int* 2013;33:1951–9.
- 72 Park Y, Lee SC, Nam H-S, et al. Comparison of sonographically guided intra-articular injections at 3 different sites of the knee. *J Ultrasound Med* 2011;30:1669–76.
- 73 Hsu Y-C, Yang F-C, Hsu H-H, et al. Ultrasound-Guided corticosteroid injection in patients with carpal tunnel syndrome: efficacy of Intra-Epineurial injection. *Ultraschall Med* 2018;39:334–42.
- 74 Babaei-Ghazani A, Nikbakht N, Forogh B, et al. Comparison between effectiveness of ultrasound-guided corticosteroid injection above versus below the median nerve in mild to moderate carpal tunnel syndrome: a randomized controlled trial. *Am J Phys Med Rehabil* 2018;97:407–13.
- 75 Mardani-Kivi M, Karimi-Mobarak M, Babaei Jandaghi A, et al. Intra-sheath versus extra-sheath ultrasound guided corticosteroid injection for trigger finger: a triple blinded randomized clinical trial. *Phys Sportsmed* 2018;46:93–7.
- 76 Shinomiya R, Sunagawa T, Nakashima Y, et al. Impact of corticosteroid injection site on the treatment success rate of trigger finger: a prospective study comparing ultrasound-guided true Intra-Sheath and true Extra-Sheath injections. *Ultrasound Med Biol* 2016;42:2203–8.
- 77 Sabeti-Aschraf M, Stotter C, Thaler C, et al. Intra-articular versus periarticular acromioclavicular joint injection: a multicenter, prospective, randomized, controlled trial. *Arthroscopy* 2013;29:1903–10.
- 78 Babaei-Ghazani A, Forogh B, Raissi GR, et al. Ultrasound-Guided corticosteroid injection in carpal tunnel syndrome: comparison between radial and ulnar approaches. *J Pain Res* 2020;13:1569–78.
- 79 Choi CK, Lee HS, Kwon JY, et al. Clinical implications of real-time visualized ultrasound-guided injection for the treatment of ulnar neuropathy at the elbow: a pilot study. *Ann Rehabil Med* 2015;39:176–82.
- 80 Fritz J, Henes JC, Thomas C, et al. Diagnostic and interventional MRI of the sacroiliac joints using a 1.5-T open-bore magnet: a one-stop-shopping approach. *AJR Am J Roentgenol* 2008;191:1717–24.
- 81 Fritz J, Tzaribachev N, Thomas C, et al. Evaluation of Mr imaging guided steroid injection of the sacroiliac joints for the treatment of children with refractory enthesitis-related arthritis. *Eur Radiol* 2011;21:1050–7.
- 82 Klauser A, De Zordo T, Feuchtnner G, et al. Feasibility of ultrasound-guided sacroiliac joint injection considering sonoanatomic landmarks at two different levels in cadavers and patients. *Arthritis Rheum* 2008;59:1618–24.
- 83 Klauser AS, Abd Ellah MMH, Halpern EJ, et al. Meralgia paraesthetica: ultrasound-guided injection at multiple levels with 12-month follow-up. *Eur Radiol* 2016;26:764–70.
- 84 Klauser AS, De Zordo T, Feuchtnner GM, et al. Fusion of real-time us with CT images to guide sacroiliac joint injection in vitro and in vivo. *Radiology* 2010;256:547–53.
- 85 Klauser AS, Sailer-Hoeck M, Abdellah MMH, et al. Feasibility of ultrasound-guided Sacroiliac joint injections in children presenting with sacroilitis. *Ultraschall Med* 2016;37:393–8.
- 86 vanVeen KEB, Alblas KCL, Alons IME, et al. Corticosteroid injection in patients with ulnar neuropathy at the elbow: a randomized, double-blind, placebo-controlled trial. *Muscle Nerve* 2015;52:380–5.
- 87 Zacchino M, Almolla J, Canepari E, et al. Use of ultrasound-magnetic resonance image fusion to guide sacroiliac joint injections: a preliminary assessment. *J Ultrasound* 2013;16:111–8.