

1 **The Relationship between Alcohol Intake and Falls**  
2 **Hospitalization: Results from the EPIC-Norfolk**

3 Guo Jeng Tan, MB BCh BAO<sup>1</sup>; Maw Pin Tan, BMBS, MD<sup>1,2,3</sup>; Robert N Luben, MSc<sup>4</sup>;  
4 Nicholas J Wareham, MBBS, PhD<sup>5</sup>; Kay-Tee Khaw, MBBChir<sup>4</sup>; Phyo Kyaw Myint,  
5 MD<sup>6</sup>.

6  
7 <sup>1</sup>Department of Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur,  
8 Malaysia

9 <sup>2</sup>Ageing and Age-Associated Disorders Research Group, Faculty of Medicine,  
10 University of Malaya, Kuala Lumpur, Malaysia

11 <sup>3</sup>Department of Medical Sciences, Faculty of Healthcare and Medical Sciences, Sunway  
12 University, Bandar Sunway, Malaysia

13 <sup>4</sup>Clinical Gerontology Unit, Department of Public Health and Primary Care, School of  
14 Clinical Medicine, University of Cambridge, Cambridge, UK

15 <sup>5</sup>MRC Epidemiology Unit, Cambridge, UK

16 <sup>6</sup>Ageing Clinical & Experimental Research (ACER) Team, Institute of Applied Health  
17 Sciences, University of Aberdeen, Aberdeen, UK

18  
19  
20 Guo Jeng Tan

21 *Lecturer in Medicine*

22  
23 Maw Pin Tan

24 *Professor of Geriatric Medicine*

25  
26 Robert N Luben

27 *Senior Research Associate*

28  
29 Nicholas J Wareham

30 *Professor & Director of MRC Epidemiology Unit*

31  
32 Kay-Tee Khaw

33 *Professor of Clinical Gerontology*

34  
35 Phyo Kyaw Myint

36 *Professor of Medicine of Old Age*

37  
38  
39 **Corresponding author:**

40  
41 Phyo Kyaw Myint

42 Room 4:013 Polwarth Building,

43 University of Aberdeen, Foresterhill,

44 Aberdeen

45 AB25 2ZD

46 Tel:+44 (0) 1224 437957 | Fax: +44 (0) 1224 437911

47 Mail to: [phyo.myint@abdn.ac.uk](mailto:phyo.myint@abdn.ac.uk)

48

49 **Short running title: Alcohol and Falls Hospitalizations**  
50  
51 **Word Count: 3345 words (without abstract) 3567 (with abstract)**  
52

53 **Abstract:**

54 **Aim:** To evaluate the relationship between habitual alcohol consumption and risk of  
55 falls hospitalization.

56 **Methods:** The EPIC-Norfolk is a prospective population-based cohort study in Norfolk,  
57 United Kingdom. A total of 25,637 community dwelling adults aged 40-79 were  
58 recruited. Units of alcohol consumed per week were measured using a validated Food  
59 Frequency Questionnaire. The main outcome was the first hospital admission following  
60 a fall.

61 **Results:** Over a median follow-up period of 11.5 years (299,211 total person years), the  
62 cumulative incidence function (95% confidence interval, CI) of hospitalized falls at  
63 121-180 months for non-users, light (>0 to ≤7 units/week), moderate (>7 to ≤28  
64 units/week) and heavy (>28 units/week) were 11.08 (9.94-12.35), 7.53 (7.02-8.08), 5.91  
65 (5.29-6.59), and 8.20 (6.35-10.56), respectively. Moderate alcohol consumption was  
66 independently associated with a reduced risk of falls hospitalization after adjustment for  
67 most major confounders (hazard ratio=0.88;95%CI:0.79-0.99). The relationship  
68 between light alcohol consumption and falls hospitalization was attenuated by gender  
69 differences. Alcohol intake higher than the recommended threshold of 28 units/week  
70 was associated with an increased risk of falls hospitalization (HR 1.40 (1.14-1.73))

71 **Conclusions:** Moderate alcohol consumption appears to be associated with a reduced  
72 risk of falls hospitalization, intake above the recommended limit is associated with an  
73 increased risk. This provides incentive to limit alcohol consumption within the  
74 recommended range and has important implications for public health policies for ageing  
75 populations.

76 **Keywords:** alcohol, elderly, falls, geriatric, hospitalization

77

## 78 **Introduction**

79 Globally, the number of persons aged 60 years and above has been estimated to  
80 be over 900 million in 2015. This number is projected to rise to 1.4 billion by 2030 and  
81 2.1 billion by 2050.(1) Falls are commonly reported among older people, with over a  
82 quarter of the American adult population (27.7%), without severe visual impairment,  
83 reporting having had falls.(2) It is also a common cause of mortality among adults. In  
84 the United States alone, more than 25,000 deaths, among the population aged 65 years  
85 and above, is attributable to unintentional falls in 2013.(3)

86 While several studies have shown that alcohol intake is not a significant  
87 predictor of falls, (4) other studies show that there is a significant association between  
88 alcohol intake and falls. (5, 6) There are also studies showing that alcohol intake is  
89 associated with a reduced risk of falls. (7) The positive association between moderate  
90 alcohol consumption in cardiovascular disease is now considered well established (8).  
91 The relationship between moderate alcohol intake and other health outcomes such as  
92 falls, is less well researched. A better understanding of the relationship between alcohol  
93 and falls is important to determine whether current alcohol consumption guidelines are  
94 potentially beneficial in reducing falls risk.

95 Therefore, based on the current evidence for the association between alcohol  
96 intake and health, we postulate that judicious use of alcohol may be protective of falls  
97 while drinking in excess is linked to increased risk of falls. The objective of this study  
98 is, therefore, to determine the potential differential relationship between different levels  
99 of habitual alcohol consumption with falls hospitalization over long term follow up in a  
100 general population of middle and older age. Individuals in middle age were included as

101 the length of follow-up, allowed us to explore the prospective relationship between  
102 alcohol consumption in middle age and falls hospitalization in later life.

103

## 104 **Methods**

### 105 **Study design and setting**

106 The EPIC-Norfolk is a prospective population-based cohort study. Participants  
107 were recruited between 1993-1997 from the city of Norwich and its surrounding rural  
108 areas in Norfolk, United Kingdom. Norfolk is a county in the East of England and  
109 Norwich is the largest city in Norfolk and serves as its administrative capital. With a  
110 population 859,400 people; a population density of 155 persons per km square. 40% of  
111 the population live in the urban areas while 60% are in the rural areas.

### 112 **Participants**

113 Men and women aged from 40 to 79 years of age were recruited from the  
114 general practice age-sex registers into the EPIC-Norfolk study. Protocol for the EPIC-  
115 Norfolk study has been published elsewhere in detail; the characteristics of the cohort  
116 were comparable to the UK population as a whole, although the percentage of current  
117 smokers were lower. (9) All participants gave written informed consent. Ethical  
118 approval was obtained from the Norwich Local Research Ethics Committee.

### 119 **Baseline assessments and measurements**

120 At baseline, all consenting participants were completed a detailed health and lifestyle  
121 questionnaire that collected information on participants' educational status, occupation,  
122 socioeconomic status, physical activity, smoking status, prevalent illness and  
123 medications. Prevalent illnesses collected included self-reported physician-diagnosed  
124 condition of heart disease, diabetes mellitus, cancer, asthma, and chronic obstructive

125 pulmonary disease Socioeconomic status was defined according to the Registrar  
126 General's occupation-based classification scheme.

127 Trained nurses then took measurements of the weight, height, body mass index (BMI)  
128 and blood pressure, and obtained non-fasting venous blood samples. Hip circumference  
129 were measured at the widest portion of the buttocks in accordance to WHO  
130 recommendations.[\(10\)](#) Physical activity, graded according to a four-level physical  
131 activity index, was derived from the validated EPIC short physical activity  
132 questionnaire. [\(11\)](#)

133 The use of medications was ascertained by enquiring whether the participant has taken  
134 any drugs or medications either prescribed by their doctor or from the chemist. The use  
135 of aspirin, steroids or diuretics was determined by asking about continual use for three  
136 months or more. Smoking history was determined using the questions: "*Have you ever*  
137 *smoked as much as one cigarette a day for as long as a year?*" and "*Do you smoke*  
138 *cigarettes now?*". Educational status was recorded as no qualification, O-level (five  
139 years of secondary education), A-level (sixth form or college), degree or higher  
140 qualification.

141

142 Area deprivation was assessed from residential postcodes using the Townsend  
143 Deprivation Index. The Townsend Deprivation Index assessed an area to be deprived  
144 based on the percentages of the following: a) households with fewer rooms than persons  
145 b) households lacking a car c) economically active persons seeking work d) children  
146 aged 5 to 15 who received school meals free and e) households experiencing  
147 disconnection of electricity in the previous 12 months.[\(12\)](#)

148 Three separate questions were asked about the reported weekly intake of fresh fruits,  
149 green leafy vegetables, and other vegetables. The options given were never, seldom,  
150 once a week, 2-3 times a week, 5-6 times a week, once or more daily and don't know.  
151 [\(13\)](#)

### 152 **Alcohol consumption**

153 The EPIC food frequency questionnaire, mentioned above, was used to assess  
154 the amount of alcohol habitually consumed. The participants reported the type of  
155 alcoholic beverage and volume in terms of cans, pints, glasses, and shots consumed  
156 within an average week. From that information the total weekly consumption of alcohol  
157 was determined. The number of units of alcohol consumed per week was then  
158 calculated based on the UK government recommended guidelines, as this is a British  
159 population. One unit of alcohol according to the National Health Service (NHS) is  
160 defined as 10ml or 8g of pure alcohol. This is different from the United States where a  
161 "standard drink: is defined as 14g of pure alcohol. An estimated consumption of more  
162 than 28 units (>224g) of alcohol per week was considered heavy alcohol consumption.  
163 Among those who consumed 28 units or less, 7 units (56g) or less per week was  
164 considered light consumption while over 7 units to 28 units (56g-228g) was considered  
165 moderate consumption. For comparison one pint of beer with an ABV of 5% is  
166 considered 3 units.

167

### 168 **Hospitalization due to falls**

169 Hospital admission episodes were identified from the National Health Service  
170 hospital information system and ENCORE (East Norfolk Commission Record). The  
171 ENCORE system has previously been validated for other diseases such as stroke. [\(14\)](#)

172 Falls as the reason for admission was identified using the International Classification of  
173 Diseases -10<sup>th</sup> Revision (ICD-10) codes W00-W19.

174

#### 175 **Statistical methods**

176 Statistical analysis was performed using SPSS version 21.0. The baseline  
177 characteristics of different alcohol consumption categories (0, >0 to ≤7, >7 to ≤28 and  
178 >28 units/week) were compared with the analysis of variance for continuous data and  
179 Chi-squared test for categorical data. A Kaplan-Meier survival curve was produced for  
180 the various categories of alcohol usage in relationship to time to first hospitalization  
181 with a fall. The length of follow-up was censored at the date of death while the time to  
182 event was the actual date of the fall. We did not look consider fall-related death only  
183 fall-related hospitalization in relation alcohol consumption. The proportion of those  
184 hospitalized during the duration of follow-up was plotted serially. We created a graph  
185 for each of the categories of alcohol intakes in 3.5 unit increments. We then repeated the  
186 analysis for all participants but this time we divide the groups into those aged <65 years  
187 and ≥65 years to examine for other patterns of association. Cox proportional hazards  
188 regression analysis was employed to determine the hazard ratios (HR) with 95%  
189 confidence intervals (CI) for risk of falls hospitalization according to alcohol categories.  
190 Using dummy variables, with the no alcohol consumption category as the reference  
191 group, the individual models compared low (>0, ≤7 units/week), moderate (>7, ≤28  
192 units/week) and heavy (>28 units/week) against no alcohol intake. There is no specific  
193 standardized cut-off point used across different studies and our cut-off points were  
194 defined based on the visual interpretation of the relationship between fall hospitalization  
195 and alcohol consumption in our graphical representations and the UK recommendations.



196 (15) Independent variables within the Cox proportional hazard analysis were added  
197 using a hierarchical approach and the variables selected for the included within the Cox  
198 proportional hazard models were informed by available published literature and clinical  
199 experience. In previous studies falls was associated with increasing age, female gender,  
200 social deprivation (16), anti-depressant use (17), reduced physical activity (18), lower  
201 BMI (19), stroke (20), diabetes mellitus (21), and increasing vegetable intake(22). Other  
202 variables were selected on the basis on difference between groups and this included  
203 smoking and educational level.

204

## 205 **Results**

### 206 **Participants**

207 Data on alcohol consumption and the presence or absence of hospitalization from a fall  
208 was available for 25,639 participants. The participants were recruited from the years  
209 1993 to 1997. The median follow-up period was 11.5 years (mean 16.23 years; total  
210 person years 299,211). The minimum follow-up period was for one month with a  
211 maximum of 277 months.

212

213 The baseline characteristics measured at enrolment between 1993-1997 are summarized  
214 in Table 1, according to alcohol usage categories.

215 Table 1

216

217

218 The number of participants who experienced hospitalization due to a fall during  
219 the follow-up period was 700 (19.2%) for those who do not drink; 1867 (13.7%) for  
220 those who drink  $>0, \leq 7$  units/week; 494 (10.9%) who drink  $>7, \leq 14$  units/week; 207

221 (10.7%) for those who drink  $>14, \leq 21$  units/week; 84 (9.2%) for those  $>21, \leq 28$   
222 units/week; and 118 (12.3%) for those drinking  $>28$  units/week. There were significant  
223 differences between the gender, age, body-mass index, waist-hip ratio, mean diastolic  
224 and systolic blood pressure, physical activity, smoking status, education level,  
225 occupational social class, the Townsend index, prevalent major co-morbid conditions  
226 such as diabetes mellitus, stroke, cancer, medications including anti-depressant use,  
227 aspirin use, anti-hypertensive use and vitamin D supplementation as well as fruit and  
228 fish intake between the different alcohol consumption categories ( $p < 0.05$ ). Time to falls  
229 hospitalization at various time points is summarized in Table 2. The cumulative  
230 incidence function (95% confidence interval, CI) at 121-180 months was 11.08 (9.94 to  
231 12.35) % for teetotalers, 7.53 (7.02-8.08)% for  $>0$  but  $\leq 7$  units/week, 5.91 (5.29 to  
232 6.59) % for  $>7$  and  $\leq 28$  units/week and 8.20 (6.35 to 10.56) % for  $>28$  units/week.  
233 Figure 1 provides the trend for fall hospitalization for overall population according to  
234 alcohol consumption categories.

## 235 **Table 2**

## 236 **Figure 1**

## 237 **Cox proportional hazards models**

238 Table 3 shows the Cox's proportional hazards models according to alcohol  
239 consumption categories. In the unadjusted analysis, falls hospitalization was  
240 significantly less likely in those with low, moderate and high intake compared to those  
241 with no alcohol intake (Model 1 and Figure 2). Following adjustment for age  
242 differences, the low intake and moderate intake groups remained significantly less likely  
243 to experience falls hospitalization compared to the no alcohol intake group. The high  
244 intake group was now significantly more likely to experience falls hospitalization

245 compared to the no alcohol intake group (Model 2). The relationship remained  
246 unchanged after additional adjustment for physical activity (Model 3) However, after  
247 adjustment for age, physical activity, stroke and diabetes, asthma and anti-depressant  
248 use the protective relationship of low alcohol intake with number of falls hospitalization  
249 against no alcohol intake was attenuated (Model 4). Moderate alcohol intake continued  
250 to be associated with reduced risk, while high intake remained deleterious after the latter  
251 adjustments. The above models suggest that the apparent lower likelihood of falls  
252 hospitalization among those with high alcohol intake over those with no alcohol intake  
253 in the unadjusted analysis was accounted for by age and gender differences.  
254 Additionally, the association of low intake with reduced falls compared to no alcohol  
255 intake appeared to be accounted for by differences in age, gender, presence  
256 comorbidities such as stroke, diabetes and physical activity. After adjusting for age,  
257 physical activity, stroke, diabetes, asthma, antidepressant use, the Townsend index, fish  
258 intake, fruit & vegetable consumption, low alcohol usage and moderate usage was still  
259 associated with a reduced risk of falls compared to non-consumers (Model 6). However  
260 after adjusting for age the reduction in falls risk in the low consumption group is no  
261 longer significant while the moderate consumption group remained significant (Model  
262 7). For Model 8, in addition to the variables adjusted in Model 7 we also adjusted for  
263 the usage of medications such as aspirin, anti-hypertensive, and vitamin D; and the  
264 physical measurements of systolic and diastolic blood pressure, body mass index and  
265 prevalent cancer. Finally, when we further adjusted for differences in smoking status  
266 and educational level whereby moderate consumption remained significantly associated  
267 with a reduction in risk of falls hospitalization (Model 9). The sub-group analysis, as  
268 stated in the methodology, for those below 65 years and those 65 years and above,

269 found that the relationship was unchanged for those aged less than 65 but the  
270 relationship was no longer present for those aged 65 years and older (Supplementary  
271 Table 1 and Supplementary Table 2).

272 **Table 3**

273 **Figure 2**

274 **Discussion**

275 Our study has revealed a U-shaped relationship between alcohol consumption  
276 and falls hospitalization over a median follow-up period of 11.5 years, after adjustments  
277 for as many potential confounders as possible. We found that those with a moderate  
278 alcohol intake experienced a 12% relatively lower risk of falls hospitalization compared  
279 to those who abstained from alcohol after controlling for potential confounders. The  
280 effect of light alcohol usage on reduced falls risk was accounted for by gender  
281 differences, physical activity, a history of diabetes, and cerebrovascular disease. Heavy  
282 alcohol consumption, as defined by the UK Chief Medical Officer's guidelines of >28  
283 units per week, was an independent predictor (40% increase in relative risk) of falls  
284 hospitalization over those who consumed no alcohol. Though the UK Chief Medical  
285 Officer's guideline now recommends <14 units per week for both men and women, our  
286 study shows that the risk of falls hospitalization is increased in those who drink  $\leq 7$  units  
287 per week and >28 units per week. Drinking > 14 units per week can be associated with  
288 increase health risks but the risk of falls hospitalization from does not increase until >  
289 28 units per week. This must be cautiously interpreted, however in the light of current  
290 evidence.  
291

292           Agahi and colleagues have shown that in the oldest old, abstaining from alcohol  
293 and heavy usage (>30 drinks/month) was associated with reduced survival.[\(23\)](#)  
294 However, our study found that no such effect in those aged 65 years and above even if it  
295 is present in those below the age of 65 years. The rationale for the reduced survival in  
296 those who abstained were two-fold, potential drug interaction with alcohol, and the  
297 association with alcohol consumption and social interaction, with social participation  
298 now considered a major protective factor against ill-health and mortality. It could be  
299 that in those aged 65 years at the outset within this study, either the effects of even mild  
300 and moderate alcohol consumption in this age group would have contributed to falls risk  
301 and therefore cancelling out any potential benefits of social inclusion. Over the long  
302 period of follow-up provided by this study those aged 65 years and over would have  
303 acquired an increasing number of comorbidities, which could then predominate over  
304 alcohol as factors which influence falls hospitalization risk.

305

306           Our study is the first to report hospitalization from falls rather than falls events  
307 in relation to alcohol consumption. Other studies have shown conflicting evidence. This  
308 is likely to have occurred because these studies assumed a rigid dichotomy either  
309 between those who consume alcohol and those who are abstinent or between those who  
310 drink heavily and those who don't. Hospitalization could be regarded as an indicator of  
311 the severity of the falls and captures falls that lead to complications such as fracture or  
312 other injuries, which are linked to morbidity, institutionalization, and death.  
313 Furthermore, hospitalization is also associated with major cost implications, with falls  
314 requiring hospitalizations regarded as an important health outcome. As alcohol  
315 consumption was determined using the food frequency questionnaire, the amount of

316 alcohol consumed could be considered as part of habitual intake (24) and results from  
317 the Agahi and colleague's study showed that drinking habits over time do not  
318 change(25). We were also able to control robustly for potential confounders. Falls  
319 events in previous studies were mostly determined by the retrospective recall of falls,  
320 which could be liable to reporting bias. Ganz had suggested that in order to overcome  
321 this, information should be gathered weekly or monthly. (26) In this study however, the  
322 falls hospitalization event was not dependent on self-report, but on objectively  
323 documented hospital records, vetted by the clinicians, and therefore can be considered  
324 hard outcomes. This will of course miss minor falls for which the patients were not  
325 hospitalized but the main thrust of our study was falls sufficiently serious to warrant an  
326 admission.

327

328         In another Swedish cohort study involving 20,212 participants, falls was  
329 associated with increased age and reduced physical activity but the tendency to drink  
330 was not associated with increased risk of falls. In this study, alcohol consumption was  
331 categorized into those who did not drink, those who drank a little or moderately, and  
332 those who drank heavily. (27) In a longitudinal analysis of five Australian cohorts  
333 involving a total of 16,785 patients, alcohol categories were separated into abstinent,  
334 <20g per day (low risk), 20g to 40g per day (long-term risk) and >40g per day (short-  
335 term risk). Abstainers of both sexes in this study did have increased odds of falling in  
336 the unadjusted models which was accounted for by depression, diabetes and if the  
337 person was female or had musculoskeletal conditions (28) In another prospective cohort  
338 study involving 5974 men a significant association with reduction in falls was shown  
339 with light alcohol intake, which was defined as less than 14 units a week. (29) Our

340 study findings suggest that the conflicting evidence produced by previous studies could  
341 be explained by the biphasic relationship between alcohol consumption with  
342 consumption of alcohol within the recommended limits being associated with a  
343 reduction of falls hospitalization compared to no alcohol and excess alcohol intake.

344

345         The association of heavy alcohol intake with falls leading to injuries, such as  
346 fractures and subdural hemorrhage, thus leading to hospitalization is well established. In  
347 a previous study examining the etiology of traumatic brain injuries, alcohol intoxication  
348 and older age were associated with increasing likelihood of traumatic brain injury from  
349 ground level falls. (30) Alcohol intoxication also increases body movements spans and  
350 balance perturbations but reduces the ability to readjust postural alignment and this lead  
351 to an increase chance of injury. Functional magnetic resonance imaging studies have  
352 demonstrated that chronic alcohol excess is associated with structural and functional  
353 changes in areas of the brain responsible for motivation and behavioral control.

354

355         We observed the U-shaped relationship in which no or low alcohol consumption  
356 increases falls hospitalization and the reason is unclear. It could be that individuals in  
357 the no alcohol intake category had health issues or financial difficulties thus requiring  
358 the individuals to be abstinent. Conversely the observed association with falls reduction  
359 of moderate alcohol consumption may be due to a reduced level of other risk factors for  
360 falls such as polypharmacy.

361

362         Although we controlled for medical risk factors, we were not able to control  
363 adequately for residual confounding and known or unknown confounders which were

364 not adjusted for. While there are concerns of recall bias when asking for an alcohol  
365 history, according to Streppel and colleagues the use food frequency questionnaire  
366 showed high correlation with 24 hour recall.<sup>(24)</sup> To minimize this bias, participants  
367 were clearly shown what would constitute a unit of alcohol. Given the prospective  
368 relationship between exposure and outcome, it is unlikely that the results are due to  
369 reverse causality (i.e. people who fell consumed more alcohol as the result of falls  
370 history). Whilst we acknowledge the usual limitations of observational cohort studies  
371 such as potential healthy responder bias, EPIC-Norfolk sample is comparable to other  
372 representative UK cohorts and thus generalizable to the UK population as whole. The  
373 hospitalization outcomes were determined based on alcohol history obtained at baseline.  
374 Alcohol consumption may have changed over the 13-year follow-up period. However,  
375 this random over/underestimate is unlikely to have impact on the direction of results.  
376 Another limitation of this study, was the EPIC-Norfolk dataset for originally collected  
377 to determine the nutritional risk factors for cancer and because of that well known fall  
378 risk factors such as previous the history of falls, fear of falling, frailty status, physical  
379 performance and gait speed were not determined at the outset.

380

381         Our study confirms the deleterious effects of drinking to excess but it also  
382 suggests that the beneficial associations of alcohol extends beyond cardiovascular  
383 diseases. Therefore, we conclude that moderate alcohol consumption may be associated  
384 with a reduced risk hospitalization from falls but the mechanism by which it may exert  
385 this effect is still unknown. Future studies should be conducted to identify the  
386 mechanisms underlying potential beneficial effect of moderate alcohol. Our findings



387 suggest that the revised recommended safe consumption level of <14 units/week is less  
388 applicable than the previous levels of <28 units/week.

389

### 390 **Acknowledgement**

391 The EPIC-Norfolk study (DOI 10.22025/2019.10.105.00004) has received funding from  
392 the Medical Research Council (MR/N003284/1 and MC-UU\_12015/1) and Cancer  
393 Research UK (C864/A14136). We are grateful to all the participants who have been part  
394 of the project and to the many members of the study teams at the University of  
395 Cambridge who have enabled this research. All authors have no conflict of interests to  
396 declare.

397

### 398 **Disclosure Statement**

399 **All authors have no conflict of interests to declare. This manuscript has not been**  
400 **published or submitted for publication elsewhere.**

401

### 402 **REFERENCES**

- 403 1. Department of Economic and Social Affairs UN. World Population Ageing  
404 2015. United Nations, Nations DoEaSAU; 2015.
- 405 2. Crews J, Chou C, Stevens J, Saaddine J. Falls Among Persons Aged >65 years  
406 with and without severe vision impairment. 2014. Contract No.: 65.
- 407 3. Control CFD. National Vital Statistics System Mortality Data.
- 408 4. Wong H, Heuberger R, Logomarsino J, Hewlings S. Association between  
409 alcohol use, polypharmacy and falls in older adults. Nurs Older People. 2016;28(1):30-  
410 6.
- 411 5. Grundstrom A, Guse C, Layde P. Risk factors for falls and fall-related injuries in  
412 adults 85 years of age and older  
413 Arch Gerontol Geriatr. 2012;54(3):421-8.
- 414 6. Chang V, Minh T. Risk Factors for Falls Among Seniors: Implications of  
415 Gender. Am J Epidemiol. 2014;181(7):521-31.
- 416 7. Coutinho E, Fletcher A, Bloch K, Rodrigues L. Risk factors for falls with severe  
417 fracture in elderly people living in a middle-income country: a case control study. BMC  
418 Geriatr. 2008;8(21).
- 419

- 420 8. Bell S, Daskalopoulou M, Rapsomaniki E, George J, Britton A, Bobak M, et al.  
421 Association between clinically recorded alcohol consumption and initial presentation of  
422 12 cardiovascular diseases: population based cohort study using linked health records.  
423 *BMJ*. 2017;356:j909.
- 424 9. Day N, Oakes S, Luben R, Khaw KT, Bingham S, Welch A, et al. EPIC-  
425 Norfolk: study design and characteristics of the cohort. *European Prospective*  
426 *Investigation of Cancer. Br J Cancer*. 1999;80 Suppl 1:95-103.
- 427 10. Organization WH. Waist circumference and waist-hip ratio: report of a WHO  
428 expert consultation, Geneva, 8-11 December 2008. 2011.
- 429 11. Wareham NJ, Jakes RW, Rennie KL, Schuit J, Mitchell J, Hennings S, et al.  
430 Validity and repeatability of a simple index derived from the short physical activity  
431 questionnaire used in the European Prospective Investigation into Cancer and Nutrition  
432 (EPIC) study. *Public Health Nutr*. 2003;6(4):407-13.
- 433 12. Townsend P, Simpson D, Tibbs N. Inequalities in health in the city of Bristol: a  
434 preliminary review of statistical evidence. *Int J Health Serv*. 1985;15(4):637-63.
- 435 13. Sargeant LA, Khaw KT, Bingham S, Day NE, Luben RN, Oakes S, et al. Fruit  
436 and vegetable intake and population glycosylated haemoglobin levels: the EPIC-  
437 Norfolk Study. *Eur J Clin Nutr*. 2001;55(5):342-8.
- 438 14. Sinha S, Myint PK, Luben RN, Khaw KT. Accuracy of death certification and  
439 hospital record linkage for identification of incident stroke. *BMC Med Res Methodol*.  
440 2008;8:74.
- 441 15. Health Do. UK Chief Medical Officers' alcohol guidelines review: Summary of  
442 the proposed new guidelines. Department of Health and Social Care London; 2016.
- 443 16. Sharma S, Mueller C, Stewart R, Veronese N, Vancampfort D, Koyanagi A, et  
444 al. Predictors of Falls and Fractures Leading to Hospitalization in People With  
445 Dementia: A Representative Cohort Study. *J Am Med Dir Assoc*. 2018;19(7):607-12.
- 446 17. Prizer LP, Smith ML, Housman J, Ory MG. Depressive symptomology  
447 management and falls among middle aged and older adults. *Aging Ment Health*.  
448 2016;20(1):13-21.
- 449 18. Dipietro L, Campbell WW, Buchner DM, Erickson KI, Powell KE, Bloodgood  
450 B, et al. Physical Activity, Injurious Falls, and Physical Function in Aging: An  
451 Umbrella Review. *Med Sci Sports Exerc*. 2019;51(6):1303-13.
- 452 19. Mazur K, Wilczyński K, Szewieczek J. Geriatric falls in the context of a hospital  
453 fall prevention program: delirium, low body mass index, and other risk factors. *Clin*  
454 *Interv Aging*. 2016;11:1253-61.
- 455 20. Goh HT, Nadarajah M, Hamzah NB, Varadan P, Tan MP. Falls and Fear of  
456 Falling After Stroke: A Case-Control Study. *PM & R : the journal of injury, function,*  
457 *and rehabilitation*. 2016;8(12):1173-80.
- 458 21. Yokomoto-Umakoshi M, Kanazawa I, Kondo S, Sugimoto T. Association  
459 between the risk of falls and osteoporotic fractures in patients with type 2 diabetes  
460 mellitus. *Endocr J*. 2017;64(7):727-34.
- 461 22. Sim M, Blekkenhorst LC, Lewis JR, Bondonno CP, Devine A, Zhu K, et al.  
462 Vegetable and fruit intake and injurious falls risk in older women: a prospective cohort  
463 study. *Br J Nutr*. 2018;120(8):925-34.
- 464 23. Agahi N, Kelfve S, Lennartsson C, Kåreholt I. Alcohol consumption in very old  
465 age and its association with survival: A matter of health and physical function. *Drug*  
466 *Alcohol Depend*. 2016;159:240-5.

- 467 24. Streppel MT, de Vries JH, Meijboom S, Beekman M, de Craen AJ, Slagboom  
468 PE, et al. Relative validity of the food frequency questionnaire used to assess dietary  
469 intake in the Leiden Longevity Study. *Nutr J.* 2013;12:75.
- 470 25. Agahi N, Dahlberg L, Lennartsson C. Social integration and alcohol  
471 consumption among older people: A four-year follow-up of a Swedish national sample.  
472 *Drug Alcohol Depend.* 2019;196:40-5.
- 473 26. Ganz DA, Higashi T, Rubenstein LZ. Monitoring falls in cohort studies of  
474 community-dwelling older people: effect of the recall interval. *J Am Geriatr Soc.*  
475 2005;53(12):2190-4.
- 476 27. Helgadottir B, Moller J, Laflamme L. Patterns in health-related behaviours and  
477 fall injuries among older people: a population-based study in Stockholm County,  
478 Sweden. *Age Ageing.* 2015;44(4):604-10.
- 479 28. Tait RJ, French DJ, Burns RA, Byles JE, Anstey KJ. Alcohol, hospital  
480 admissions, and falls in older adults: a longitudinal evaluation. *Int Psychogeriatr.*  
481 2013;25(6):901-12.
- 482 29. Cawthon PM, Harrison SL, Barrett-Connor E, Fink HA, Cauley JA, Lewis CE,  
483 et al. Alcohol intake and its relationship with bone mineral density, falls, and fracture  
484 risk in older men. *J Am Geriatr Soc.* 2006;54(11):1649-57.
- 485 30. Poyry T, Luoto TM, Kataja A, Brander A, Tenovuo O, Iverson GL, et al. Acute  
486 assessment of brain injuries in ground-level falls. *J Head Trauma Rehabil.*  
487 2013;28(2):89-97.
- 488

## 489 **Figure Legend**

### 490 **Figure 1. Alcohol Consumption and Fall Hospitalization**

491 Alcohol consumption and hospitalization due to falls for all participants and for those  
492 aged <65 years and ≥65 years. Hospitalization from falls appeared lower among groups  
493 who consumed more than 7 units of alcohol to less than and up to 28 units of alcohol  
494 per week for categories for all categories.

495

### 496 **Figure 2. Kaplan Meier Survival Curve for Falls Hospitalization**

497 Kaplan Meier survival curve for time to hospitalization due to falls for the four different  
498 alcohol consumption categories using unadjusted statistics. Unadjusted figures suggest  
499 that individuals who consume no alcohol were most likely to be hospitalized with falls,  
500 while those who consume >7 units but ≤28 units of alcohol per week were least likely to  
501 be hospitalized for falls.

502

503 **Table Legend**

504

505 **Table 1 Basic Characteristics According to Alcohol Usage**

506

507 **Table 2: First Falls Hospitalization at Various Time Points**

508

509 **Table 3: Hazard Ratios for Falls Hospitalizations Adjusted for Various Potential**

510 **Confounders**

511

512

513 **Supplementary Tables**

514

515 **Supplementary Table 1: Hazard Ratios for Falls Hospitalizations Adjusted for**

516 **Various Potential Confounders aged below 65 Years**

517

518 **Supplementary Table 2: Hazard Ratios for Falls Hospitalizations Adjusted for**

519 **Various Potential Confounders for Those Aged 65 Years and Above**

520

Tables for Alcohol and Falls Hospitalizations

**Table 1 Basic Characteristics According to Alcohol Usage**

		Alcohol Usage				p-value
		0 units/ week  (n=3638)	>0, ≤7 units/ week  (n=1367 5)	>7, ≤28 units/ week  (n=7367)	>28 units/ week  (n=959)	
Age (years), mean (standard deviation)		62.15 (9.02)	59.17 (9.23)	58.22 (9.33)	57.07 (9.09)	<0.001
Female, n (%)		2478 (68.1%)	8577 (62.7%)	2885 (39.2%)	92 (9.6%)	<0.001
Physical activity, n (%)	Inactive	1564 (43.0%)	4076 (29.8%)	1957 (26.6%)	266 (27.7%)	<0.001
	Moderately inactive	912 (25.1%)	4035 (29.5%)	2146 (29.1%)	258 (26.9%)	
	Moderately active	660 (18.1%)	3179 (23.2%)	1744 (23.7%)	193 (20.1%)	
	Active	502 (13.8%)	2385 (17.4%)	1519 (20.6%)	242 (25.2%)	
Cigarette smoking, n (%)	Current	511 (14.0%)	1547 (11.3%)	936 (12.7%)	210 (21.9%)	<0.001
	Former	1183 (32.5%)	5151 (37.7%)	3839 (52.1%)	588 (61.3%)	
	Never	1944 (53.4%)	6977 (51.0%)	2592 (35.2%)	161 (16.8%)	
Occupation, n(%)	Professional	149 (4.1%)	833 (6.1%)	696 (9.4%)	76 (7.9%)	<0.001
	Managerial and Technical	918 (25.2%)	4504 (32.9%)	3287 (44.6%)	449 (46.8%)	
	Skilled non-manual	592 (16.3%)	2351 (17.2%)	1087 (14.8%)	109 (11.4%)	
	Skilled manual	921 (25.3%)	3293 (24.1%)	1374 (19.7%)	184 (19.2%)	
	Semi-skilled	673 (18.5%)	1908 (14.0%)	677 (9.2%)	103 (10.7%)	
	Unskilled	234 (6.4%)	496 (3.6%)	135 (1.8%)	20 (2.1%)	
Myocardial infarction, n (%)		135 (3.7%)	417 (3.0%)	223 (3.0%)	32 (3.3%)	0.197
Cerebrovascular accident, n (%)		95 (2.6%)	159 (1.2%)	96 (1.3%)	13 (1.4%)	<0.001
Cancer, n (%)		226 (6.2%)	775 (5.7%)	374 (5.1%)	35 (3.6%)	0.004
Diabetes mellitus, n (%)		149 (4.1%)	289 (2.1%)	132 (1.8%)	18 (1.9%)	<0.001

Asthma, n (%)	339 (9.3%)	1130 (8.3%)	611 (8.3%)	83 (8.7%)	0.217	
Anti-depressant, n (%)	292 (8.0%)	610 (4.5%)	255 (3.5%)	30 (3.1%)	<0.001	
Aspirin, n (%)	322 (8.9%)	929 (6.8%)	601 (8.2%)	86 (9.0%)	<0.001	
Statin, n (%)	31 (0.9%)	152 (1.1%)	69 (0.9%)	7 (0.7%)	0.325	
Anti-hypertensive, n (%)	994 (27.3%)	2506 (18.3%)	1179 (16.0%)	144 (15.0%)	<0.001	
Vitamin D, n (%)	1020 (28.0%)	4407 (32.2%)	7367 (30.7%)	224 (23.4%)	<0.001	
Education level, n (%)	None	2010 (55.4%)	5397 (39.5%)	1820 (24.7%)	229 (23.9%)	<0.001
	O-level	308 (8.5%)	1457 (10.7%)	768 (10.4%)	97 (10.1%)	
	A-level	1074 (29.5%)	5339 (39.0%)	3386 (46.0%)	459 (47.9%)	
	Degree	240 (6.6%)	1482 (10.8%)	1393 (18.9%)	174 (18.1%)	
Mean systolic blood pressure, mmHg (s.d.)	138.2 (19.3)	134.7 (18.4)	135.0 (17.9)	140.2 (17.3)	<0.001	
Mean diastolic blood pressure, mmHg (s.d.)	83.2 (11.6)	81.9 (11.1)	82.8 (11.1)	87.1 (11.3)	<0.001	
Mean body mass index (s.d.)	26.80 (4.44)	26.32 (4.01)	26.16 (3.45)	26.97 (3.60)	<0.001	
Mean waist-hip ratio (s.d.)	0.85 (0.09)	0.84 (0.09)	0.87 (0.09)	0.93 (0.07)	<0.001	
Townsend Index (s.d.)	-1.73 (2.34)	-2.08 (2.13)	-2.17 (2.08)	-1.74 (2.39)	<0.001	
Fruit intake (s.d.)	264 (220)	256 (186)	234 (172)	168 (159)	<0.001	
Vegetable intake (s.d.)	270 (163)	271 (134)	272 (123)	265 (126)	0.445	
Fish intake (s.d.)	36 (28)	37 (25)	40 (27)	39 (25)	<0.001	

s.d.=standard deviation

**Table 2: First Falls Hospitalization at Various Time Points**

Alcohol consumption (units/week)	Time Points (months)	No. at risk	No. falls hospitalization	Cumulative incidence function (%) (95%CI)
None	0-12	3,638	2	0.04 (0.01-0.22)
	12-36	3,625	21	0.64 (0.42-0.95)
	37-60	3,588	36	1.63 (1.27-2.10)
	61-120	3,551	132	5.29 (4.61-6.07)
	121-180	3,419	124	11.08 (9.94-12.35)
>0, ≤7 units/week	0-12	13,674	3	0.02 (0.01-0.07)
	13-36	13,632	27	0.22 (0.15-0.32)
	37-60	13,551	74	0.77 (0.63-0.93)
	61-120	13,477	337	3.25 (2.96-3.56)
	120-180	13,140	343	7.53 (7.02-8.08)
>7, ≤28 units/week	0-12	7,365	2	0.03 (0.01-0.11)
	13-36	7,353	14	0.22 (0.13-0.36)
	37-60	7,317	29	0.61 (0.46-0.82)
	61-120	7,284	131	2.40 (2.07-2.78)
	120-180	7,153	148	5.91 (5.29-6.59)
>28 units/week	0-12	959	0	0
	13-36	954	3	0.32 (0.10-0.98)
	37-60	944	3	0.63 (0.28-1.40)
	61-120	941	27	3.48 (2.49-4.87)
	120-180	914	26	8.20 (6.35-10.56)
Any Alcohol Use (>0 units/week)	0-12	21,998	5	0.02 (0.01-0.05)
	13-36	21,939	44	0.22 (0.17-0.30)
	37-60	21,812	106	0.71 (0.61-0.83)
	61-120	21,702	495	2.97 (2.76-3.21)
	121-180	21,207	517	7.02 (6.63-7.43)

**Table 3: Hazard Ratios for Falls Hospitalizations Adjusted for Various Potential Confounders**

	Hazards Ratio (95% Confidence Interval)			
	0 unit/week (no alcohol)	>0-7.0units/week (low)	>7.0-28 units/week (moderate)	>28 units/week (heavy)
Model 1	reference	<b>0.70 (0.64-0.77)</b>	<b>0.54 (0.49-0.60)</b>	<b>0.66 (0.54-0.80)</b>
Model 2	reference	<b>0.87 (0.80-0.95)</b>	<b>0.71 (0.64-0.79)</b>	0.95 (0.78-1.15)
Model 3	reference	<b>0.88 (0.80-0.96)</b>	<b>0.72 (0.65-0.80)</b>	0.96 (0.78-1.17)
Model 4	reference	<b>0.89 (0.81-0.97)</b>	<b>0.73 (0.66-0.81)</b>	0.97 (0.79-1.18)
Model 5	reference	<b>0.90 (0.82-0.98)</b>	<b>0.74 (0.67-0.82)</b>	0.97 (0.80-1.19)
Model 6	reference	<b>0.90 (0.82-0.98)</b>	<b>0.75 (0.67-0.83)</b>	1.00 (0.82-1.22)
Model 7	reference	0.93 (0.85-1.02)	<b>0.88 (0.79-0.98)</b>	<b>1.39 (1.13-1.17)</b>
Model 8	reference	0.94 (0.85-1.03)	<b>0.88 (0.79-0.99)</b>	<b>1.40 (1.14-1.73)</b>
Model 9	reference	0.93 (0.85-1.02)	<b>0.89 (0.79-0.99)</b>	<b>1.40 (1.14-1.72)</b>

**Bold letters indicate statistical significance**

Model 1=unadjusted

Model 2=adjusted for age

Model 3=adjusted for age, and physical activity

Model 4=adjusted for age, physical activity, stroke, diabetes, asthma and antidepressant use.

Model 5=adjusted for age, physical activity, stroke, diabetes, asthma, antidepressant use, and Townsend index.

Model 6= adjusted for age, physical activity, stroke, diabetes, asthma, antidepressant use, Townsend index, fish intake and fruit and vegetable consumption.




Model 7= adjusted for age, physical activity, stroke, diabetes, asthma, antidepressant use, Townsend index, fish intake, fruit and vegetable consumption, and gender.

Model 8= adjusted for age, physical activity, stroke, diabetes, asthma, antidepressant use, Townsend index, fish intake and fruit & vegetable consumption, gender, aspirin usage, anti-hypertensive medication use, vitamin D supplementation, systolic and diastolic blood pressure, body mass index, waist-hip ratio, and prevalent cancer.

Model 9 = adjusted for age, physical activity, stroke, diabetes, asthma, antidepressant use, Townsend index, fish intake and fruit & vegetable consumption, gender, aspirin usage, anti-hypertensive medication use, vitamin D supplementation, systolic and diastolic blood pressure, body mass index, waist-hip ratio, and prevalent cancer, smoking status, educational status.


**Figure 1 Alcohol Consumption and Falls Hospitalization**



**Figure 1. Alcohol Consumption and Fall Hospitalization**

Alcohol consumption and hospitalization due to falls for all participants and for those aged <65 years and ≥65 years. Hospitalization from falls appeared lower among groups who consumed more than 7 units of alcohol to less than and up to 28 units of alcohol per week for categories for all categories.

**Figure 2. Kaplan Meier Survival Curve for Falls Hospitalization**



**Figure 2. Kaplan Meier Survival Curve for Falls Hospitalization**

Kaplan Meier survival curve for time to hospitalization due to falls for the four different alcohol consumption categories using unadjusted statistics. Unadjusted figures suggest that individuals who consume no alcohol were most likely to be hospitalized with falls, while those who consume >7 units but ≤28 units of alcohol per week were least likely to be hospitalized for falls.

# The Relationship between Alcohol Intake and Falls Hospitalization:

## Results from the EPIC-Norfolk

### Supplementary Tables and Figures

#### Supplementary Table 1: Measurements

Smoking history	
“Have you ever smoked as much as one cigarette a day for as long as a year?”	Yes No
“Do you smoke cigarettes now?”	Yes No
EPIC short physical activity questionnaire (in the last one year)	
Occupational activity	Sedentary Standing Manual work Heavy manual work
Participation in walking, cycling, do-it-yourself, gardening, sports, household chores	None ≤ 3.5 hours/week >3.5 to ≤7.0 hours/week >7.0 hours/week
Participation in vigorous non-occupational activities	None ≤ 3.5 hours/week >3.5 to ≤7.0 hours/week >7.0 hours/week
Number of floors of stairs climbed a day	
Educational status (the highest attainment was recorded as follows)	
No qualification	
O-level (five years of secondary education)	
A-level (sixth form or college)	
Degree or higher qualification	

Townsend Deprivation Index	
Percentage of households with fewer rooms than persons and	
Percentage of households lacking a car	
Percentage of economically active persons seeking work	
Percentage of children aged 5 to 15 who received school meals free	
Percentage of households experiencing disconnection of electricity in the previous 12 months	
Weekly Intake of fresh fruits, green leafy vegetables, and other vegetables	
Weekly intake of fresh fruits	Never Seldom Once a week 2-3 times a week 5-6 times a week Once or more daily Don't know
Weekly intake of green leafy vegetables	Never Seldom Once a week 2-3 times a week 5-6 times a week Once or more daily Don't know
Weekly intake of other vegetables	Never Seldom Once a week 2-3 times a week 5-6 times a week Once or more daily Don't know

