

## The Versatile Wayfinder: Prefrontal Contributions to Spatial Navigation

Eva Zita Patai<sup>1,2</sup> and Hugo J. Spiers<sup>2</sup>

1. Centre for Neuroimaging Sciences, IoPPN, King's College London, UK

2. Institute of Behavioural Neuroscience, Department of Experimental Psychology, Division of Psychology and Language sciences, University College London, UK

\*Correspondence : [eva.patai@ucl.ac.uk](mailto:eva.patai@ucl.ac.uk) (E.Z. Patai) [h.spiers@ucl.ac.uk](mailto:h.spiers@ucl.ac.uk) (HJ Spiers)

**Keywords:** Navigation, Prefrontal Cortex, Hippocampus, Planning, Decision-Making.

### Abstract

The prefrontal cortex supports decision-making, goal tracking and planning. Spatial navigation is a behaviour that taxes these cognitive processes, yet the role of the prefrontal cortex in models of navigation has been largely overlooked. In humans, activity in dorsolateral and ventrolateral prefrontal cortex during detours reveal a role in inhibition and re-planning. Dorsal anterior cingulate is implicated in planning and spontaneous internally-generated changes of route. Orbitofrontal cortex integrates representations of the environment with the value of actions, providing a 'map' of possible decisions. In rodents, medial frontal areas interact with hippocampus during spatial decisions and switching between navigation strategies. In reviewing these advances, we provide a framework for how different prefrontal regions may contribute to different stages of navigation.

## **Prefrontal cortex: An overlooked key brain region for navigation**

Our planet offers a dazzling variety of environments that animals must navigate. For mammals, a set of four key brain regions are thought to form a core navigation network: the hippocampus, parahippocampus, retrosplenial cortex, and posterior parietal cortex [1,2]. These regions support place learning and flexible navigation requiring self-localisation (e.g. ‘where am I on this mountain?’). They are accompanied by the striatum for navigation based on actions linked to specific stimuli [3,4] (e.g. ‘turn left at the waterfall’). Here, we review evidence that the prefrontal cortex (PFC) is an overlooked key component of the mammalian navigation system which allows the navigator to adapt to changes in the environment, track progress, and plan ahead to avoid making costly mistakes.

Planning, decision-making, prospective memory, goal-coding, and adaptive behaviour are all cognitive functions synonymous with PFC function [5,6] and are often required in real-world navigation [1,2]. Indeed, many functional neuroimaging studies have reported PFC activation during navigation [7,8,17–22,9–16] (see Figure 1). So why has the PFC been neglected in theories of navigation? We suggest three main reasons. First, patients with frontal lobe damage are commonly reported to have impaired executive function [23], with navigation impairments being a corollary of a more general dysfunction in the regulation of behaviour [24,25]. Second, while brain regions in the core navigation network display spatial coding in their activity patterns, such as **grid cells** and **place cells** [26], similar coding has been only recently reported for PFC neurons [27,28]. Third, lesions to the rodent medial PFC, which receives hippocampal afferents [29], have generally been reported to have little or no impact on navigation in a widely used assay for navigation: the Morris water-maze [30,31]. These observations may explain why many reviews of navigation make little or no mention of the PFC [32–37].

More recent research has revealed that the PFC may play important roles in navigation. Rodent lesion studies have implicated medial PFC regions, infralimbic (IL) and prelimbic (PL) in aspects of navigation, such as learning multi-region environments [48], reversal learning in the water maze [49] and strategy switching [50]. Evidence of spatial metric coding in PFC has been reported in rodent [51] and human studies using both functional imaging [52] and intracranial recordings [27,28,53]. Such studies add weight to the argument that PFC is a key brain region for spatial coding in navigation. More broadly, PFC regions may aid navigation by processing the presence of others in the environment to infer popular routes or problems ahead - the social aspects of wayfinding [54]. Consistent with this, the rostral medial PFC [55] and anterior cingulate cortex (ACC) [56] are both active when tracking others in the environment, and medial regions of PFC are active when mapping social networks [58] or processing the intentions of others during navigation [57].

Here we integrate findings from humans, non-human primates, and rodents. This cross-species integration requires acknowledgement of the challenge due to the differences in neuroanatomy [38-40], navigation tasks used, and neural recording methods employed. For human studies, overlap in the nomenclature for similar regions can be confusing (see Figure 2 and Table 1). It is important not to oversimplify the PFC as a set of separate 'modules' - rather it should be considered part of an interconnected set of brain regions, within which the PFC provides a dynamic, state dependent, contribution to network operations via functional networks interacting on both local and global hierarchical architectures [37,41]. However, because of the gaps in our current knowledge, we will often be discussing specific prefrontal areas and their involvement in a specific scenario during navigation.

To examine how different PFC regions might contribute to solving the changing demands imposed by real-world navigation, we draw on the literature to describe how PFC dynamics likely operate

during an exemplar navigational journey. Because navigation ability and strategy choices vary considerably in humans [42–45] and other mammals [46], we focus on brain dynamics in successful navigators who efficiently make use of the core navigation network during navigation [18,47].

### **Selecting the goal and making a plan**

We start our navigational journey by setting a goal or goals. Sometimes there is a clear goal, such as ‘How do I get home?’. Other times, we may need to select a navigation goal from a set of possibilities, such as ‘Where can I find coffee in a hurry?’. A trip could range from involving a single familiar goal to multiple new goals. Importantly, navigating a highly familiar route will likely place little demand on PFC or indeed much of the core network for navigation [18,47]. In contrast, selecting a goal from competitors, navigating a novel region of space, or considering alternate routes, would all likely place demands on PFC. Such goal selection and comparison of alternative paths are core example of what defines planning as opposed to the moment-to-moment decision-making at waypoints [8,59–62].

Some models of navigation argue that the PFC supports initial planning [63,64] and distance to goal coding [65] by interacting with hippocampal networks to support navigational guidance [25,62]. Consistent with this, multiple lines of empirical evidence from tasks requiring learning stimulus-reward associations suggest that choosing the correct target from a set of competing alternatives involves the anterior cingulate cortex (ACC) and ventromedial (vm) PFC [66,67]. ACC might integrate the physical demands of travel to guide actions [68,69] and vmPFC might weight alternatives in terms of their potential value [67]. However, there has been surprisingly little research to verify this for navigation. Once the goal is set it may need to be visualised to aid consideration of the route [70]. vmPFC may also play a role in driving the construction of scene

imagery of the goal in posterior regions including the hippocampus [71–73], with such scene imagery forming an important part of planning complex routes [70].

Sometimes we must navigate to a set of goals, requiring the allocation of a final goal and sub-goals [74]. The neuropsychological task the ‘Zoo Map Test’ [75] probes the ability to plan a multi-step route from viewing a map and is impaired by medial frontal damage in humans [76]. Single unit recordings in non-human primates indicate that lateral PFC regions code information about multiple step decisions during action planning in on-screen mazes [77,78] (see Figure 3E). This research, combined with evidence from non-spatial studies, suggests the lateral prefrontal cortex, and its interaction with posterior parietal cortex, is important for planning routes and dealing with sub-goals [79], and is necessary for sequencing higher-order behaviour [80]. Consistent with this, London taxi drivers making an initial plan for complex routes through London (UK) show increased activity in lateral PFC regions [21].

Lateral PFC is not the only region engaged during sequential planning. Manipulating the complexity and difficulty of the maze being navigated reveals that rostro-dorsal medial (rd-m) PFC is more active when decisions are easy at a first choice point but hard at later point, whereas dorsal-ACC/pre-supplemental motor area activity is increased when choices are difficult in general [14]. Lateral frontopolar cortex (LFPC) responded to the difficulty at the second choice point, i.e. future demands only (see Table 1, Figure 3B). Follow-up psychophysiological interaction analyses showed that coupling between the hippocampus and rd-mPFC increased during sequential planning and when correct (versus incorrect) choices were made.

A key question is how might a ‘plan’ be processed at the neural level? There has been increasing evidence that during periods of relative immobility sequential **replay** of hippocampal place cells

along a future route, whether previously experienced or not, might form the basis for such planning (Box 1) [81–85]. ‘Forward sweeps’ through alternative future paths have been observed during active movements [86,87]. While these pre/replay events have been extensively studied in the rodent hippocampal formation, spatio-temporal reinstatement (‘replay’) of abstract states has been reported in humans using non-invasive imaging [88,89] and research with rodents has found increased coordinated forward replay across hippocampus and PFC resulted in better spatial memory performance [90], for review see [60]. Evidence for prefrontal and hippocampal coherence, primarily in the theta band, has also been found in human neuroimaging experiments testing planning with the recall of the spatial location of a target [91] and imaginary future thinking [92].

### **Travelling: Tracking progress, goal maintenance, and the default-mode network**

During navigation, a major part of the journey will likely be simply travelling to the next waypoint or goal [70]. During such travel there is potential for freeing up systems from navigational deliberation for other tasks, such as recalling memories, enjoying the view, mulling future life plans and general mind wandering [70]. In a recent study we found evidence of increased activity in core regions of the **default-mode network (DMN)**, including the vmPFC, when periods of ongoing travel through the environment were compared to events where the participants decided to change their plan [13]. Given its proposed role in internalized thought, this may point to a role of the DMN during navigation when no updates need to be made and the route-plan is being followed from memory [59]. Thus, during travel periods in a familiar environment, much of the time may be spent engaging regions of our default-mode network, possibly due to mind-wandering or the recall of spatial locations along our route [70]. Simultaneous fMRI and intracranial recordings in anaesthetised primates have shown that hippocampal sharp wave ripples (SWRs) - which are associated with place cell replay and memory consolidation in sleep - increase the activation of the cortical DMN [93].

However, it is unclear how planning indexed via hippocampal replay events interact with the DMN during active navigation. Moreover, PFC dynamics based on spontaneous behaviour are missing in most models of navigation. This may be because most studies are optimized to include as much time focused on navigational processing. Future research integrating across rodents and humans with longer more diverse experiences may be useful.

Sometimes however, we cannot be on 'auto-pilot' and choices are needed to reach our destination. While reports of 'goal' related activity have mainly focussed on the rodent and human hippocampus [83,94,95], there is ample evidence of a similar signal in medial prefrontal cortices [51,96,97]. In a seminal study, Ekstrom et al. [28] recorded neurons of pre-surgical epileptic patients navigating a virtual environment and found frontal neurons which were selectively active for specific goal locations and maintained their activity during the travel. This work was recently extended by showing goal selective activity in medial PFC regions, revealing the oscillatory dynamics of such regions for coding information for navigation [53]. The representation of salient locations has also been found in the rodent prelimbic area, and these cells outnumbered 'place' cells found in the same region that code for current location [51,95]. Goal trajectory information appears to be conveyed from the rodent mPFC to the CA1 region of the hippocampus via the nucleus reuniens (NR) once learning is established, but this interaction is reversed during early learning [98]. While these studies show the representation of general goal coding or sequences [99] or paths to a goal [100] they do not explicitly show the coding of the specific, current goal location. It is also unclear whether goal-coding observed in hippocampal regions is inherited from PFC regions or vice versa. Evidence suggests that the core spatial computations for distance, direction, and future trajectory come from the hippocampal formation, but that computations supporting goal selection and other aspects of navigation covered in this review likely arise in the PFC, which exert influence on the spatial computations in the hippocampal formation.

In humans, dorsomedial PFC may track progress to the goal. Dorsomedial PFC showed increased activity during periods of travel where the navigator reported looking ahead for a key landmark or junction [57], and when keeping track of self-motion relative to a previous location [101,102]. Activity in the mPFC scales with current goal distance in both life-like route planning [8,10,22] and in simplified environments [20,74] (Figure 3H). In a rare case, a patient with extensive ventromedial PFC damage was tested on real-world navigation of a familiar environment and was found to frequently forget the current goal and arrive instead at locations that were very familiar. If the patient was given reminders to maintain the goal, they were able to navigate successfully (Figure 1E), indicating that ventral regions of the PFC may play a key role in goal maintenance during travel, rather than long-term memory for the goals [24,103]. This observation may also relate to neural activity recorded in similar regions in non-human primates where progress (proximity to reward) through a set of sequential stimuli was found to be tracked by increasing activity in ACC [104]. When monkeys were tasked with moving a cursor to capture a virtual prey on a 2D computer display, dACC neurons were found to predict the future location of a target [105]. This not only extends the finding that the PFC is involved in predicting future states, but also indicates that it can represent an external entity. It should be noted that goal distance tracking can be confounded by number of other spatial factors, such as number of upcoming choices and proximity to reward [83]. Furthermore, recent evidence in rodents suggests the PFC may not purely support maintenance of the current goal location [106].

### **Choosing the correct path at waypoints**

For many terrestrial animals, navigation often occurs along distinct paths due to the vegetation, rock formations, or the structure of urban developments. A key challenge facing the navigator is choosing the optimal path when confronted with options. Increased dorsolateral PFC activity is evident when navigators are making an active choice compared to when their choice is guided at junctions [10,47]. Sometimes it may be difficult to choose the path because the two options ahead have a similar distance to the goal. Dorsolateral, ventrolateral PFC and ACC activity in humans has been reported to increase with the similarity of the distance between the alternative paths to a goal [16]. In rodents, a number of studies have shown that during active navigational decision-making PFC-hippocampal interactions increase (e.g. phase-locking of firing or increased coherence in local field potentials: [107,108]). Recent evidence suggests CA1 can share future goal information with mPFC and that mPFC may guide which goal is coded by CA1 [109].

One striking piece of evidence for path choice behaviour in rodents comes from Muenzinger (1938) who described back-and-forth looking behaviour at choice points in a maze. This **vicarious trial and error (VTE)** behaviour resembles 'planning' in animals when there is uncertainty, and has been shown to be hippocampal dependent [110] and inversely related to replay linked hippocampal sharp-wave-ripples [111]. VTE is thought to rely on hippocampal-cortical interactions [61], and recent evidence suggests that disruption of the dorsal medial PFC reduces VTE behaviour and has downstream consequences in the hippocampus [112], pointing to a more direct role in planning during memory-guided behaviour. Consistent with a role in coordinating behaviour, disruptions to rodent medial PFC also disrupts the capacity to switch between place and response strategies at junctions in a maze [50].

In terms of evaluating choices, the OFC has been argued to operate as a **cognitive map** of the **state space** required for value-based decision making, in particular in allowing inferences about **hidden states** and connections [113,114]. Evidence for this so far has been lacking for navigation, but studies exploring choices with non-spatial stimuli in humans support this view [115], including the role of the OFC in representing distant states in the future [116]. A recent study showed that OFC, but not ACC, inactivation in rodents abolished inference-based foraging on a lever-based pressing task and lead to stimulus-bound behaviour in a foraging task [117]. Similarly, OFC lesions in rodents cause deficits in choice behaviour when action-outcome associations are hidden, but not when they are directly observable [118]. Recent rodent and human evidence suggests that the OFC is involved in updating and learning values, not directly in choice behaviour itself [119,120]. Thus, it seems likely that the OFC would play a key role in assigning value to choices at junctions during navigation, via inferences about untaken but possible paths.

### **Adapting the plan: Detours and Shortcuts**

Despite meticulous planning and keeping the goal in mind, the navigator may still struggle to reach their goal. Detours are a frequent, and frustrating part of life. However, animals show a remarkable capacity for adaptation, with the earliest detour behaviour reported for spiders in 1912 [121]. How the brain adapts behaviour when detours are required has been the focus of a number of human neuroimaging studies. These have consistently reported PFC activity (for a review see [25]). Superior frontal gyrus (SFG) and (predominantly right) lateral PFC are activated when people encounter detours in their path [10–12,15,16,19,21,122,123] which suggests these regions play a role in halting behaviour when unexpected changes in the environment are detected. Such a result is consistent with this area being involved in detecting novelty [124], or updating predictions about the environment with a ‘prediction error’ [125], and response inhibition [126]. Consistent with this,

in a study involving continuous changes in possible paths, activation in lateral PFC scaled with the difference between the learned value of a route and its value after the change [15].

Some detours can be a mild inconvenience along a similar length of path, while others can cause significant additional travel. Thus, activations linked to detours may vary in relation to the future demands. To explore this in a recent study [12], participants navigated a familiar maze in order to find various goal objects. Compared to the training period, however, changes to the previously familiar layout of the maze occurred during the test phase, causing participants to encounter either a block in the path (detour) or a new opening (shortcut). Importantly, both the type (detour or shortcut) and magnitude of change (long or short) were manipulated. Firstly, prefrontal areas increased activation upon the detection of detours only, not shortcuts, and secondly, they were maximally active when a large detour was required (Figure 3F). This points to a specific role of the SFG and rIPFC in representing prediction errors that increase with demands and when a significant re-routing is required, rather than pure novelty. When a detour is encountered, planning the new route may require searching through the possible paths available to reach the goal. Javadi et al. [11] found that at detours lateral PFC activity increased when the number of future streets to consider was higher (Figure 3D), indicating this region may be performing a breadth-first-search through the possible paths.

Few human studies have explored the neural responses during shortcuts [12,15] and even fewer in rodents [127], despite evidence that de-novo paths, including shortcuts, can be generated in neural responses [84] and Tolman's emphasis on this behaviour in his conceptualization of the cognitive map [128]. There is little evidence of evoked increases in PFC activity when novel shortcuts were detected and taken [12]. This highlights that visual changes in the path and changes in the current plan do not necessarily result in observable responses in the PFC. However, increased rIPFC

activation was evident when false shortcuts were encountered (a novel path option is encountered that appears to lead to the goal, but actually adds distance). When such false shortcuts were correctly rejected, SFG was active (Figure 3F). This indicates that the rIPFC may be involved in initiating a stop signal driving re-plan, whereas the SFG is important for processing and successfully resolving conflict between route options [25]. Exploring neural oscillations during navigation revealed decreases in beta power when false shortcuts needed to be avoided [12]. Such reductions in beta power have been associated with increased mnemonic reactivation [129], suggesting a need for increased mnemonic processing of the spatial layout when unhelpful novel paths must be avoided.

### **Backtracking**

Occasionally, it is not the environment that causes a change in route, but rather ourselves, when we realise we have taken the wrong path. This can be observed when an animal turns around and 'backtracks' along the path just taken [130]. Only a couple of studies have reported the neural correlates of this process [13,131]. Javadi et al [13] found that during navigation, participants would intermittently backtrack and that in the majority of cases this was the optimal choice (Figure 3G). At the initiation of backtracking dorsal ACC (dACC) was more active and regions associated with the core DMN suppressed. This is consistent with work showing that dACC processes new information linked with uncertainty and updating beliefs or behaviour [132], choosing future options [133], increased foraging behaviour [67], and the ability to switch and maintain new strategies [134]. The idea that the dACC preferentially represents the value of the 'nondefault' option and the vmPFC represents the value of the 'default' option [135] also dovetails with our result that DMN activity, which includes the vmPFC, was decreased when this spontaneous (and non-standard) behaviour was exhibited. Leveraging the high temporal resolution of magnetoencephalography, it was possible to reveal that 700ms before the onset of backtracking, alpha power decreased, a pattern of activity

associated with anticipatory changes in attention [136]. Thus, the brain may shift between a default-state of moving through the environment following external environmental cues, to a focus on internal (memory-based) guidance systems to guide action. Other theories argue the polar frontal cortex should also be engaged for such switches [137]. However, this has not been observed yet during navigation. New evidence shows that activity in the hippocampus is increased for upcoming attentional states that facilitate spatial-relational processing based on memory-guided cueing, but this is not the case for the mPFC, which predicts attentional states from both memory or explicit cues [138]. Translating this to active wayfinding, it would be interesting to explore how externally or internally derived attentional cues modify navigation behaviour and prefrontal activity. More research exploring how changes in attention impact neural activity in navigation is needed.

### **Goal approach and memory consolidation via prefrontal-hippocampal interactions**

The last step of a journey will be approaching the goal location. In mice, such approach behaviour has been related to activation of pathways linking the hippocampus to frontal cortical regions [139]. Evidence suggests arrival at a goal may trigger consolidation of the recent experiences in existing memory networks, particularly if a novel route was required or if the new route is added to a schematic representation of the space [140,141] perhaps allowing meta-learning over multiple experiences ([142] and updating a map of the state-space for navigation [115], (see Box 1).

Directly linking PFC-hippocampal coordinated activity to measurable outcomes in navigation behaviour, a recent rodent study demonstrated two fundamental advancements on this topic: firstly, coordinated replay between CA1 and PFC was higher for trajectories that were subsequently taken than for alternative but untaken paths (both for forward and reverse replay events), and secondly, the strength of this relationship during recall of the task rule also predicted performance

on the task [90] (see also Box 1). This points to a role in PFC-hippocampal interaction, not only in terms of planning, but also for consolidation of successful routes post-arrival at the goal [143–145].

### **Concluding Remarks**

We have argued the PFC plays a pivotal role in a range of situations during navigation (See Key Figure, Table 1 for summaries). Such situations include: route planning, adapting to changes in the environment, updating route plans, tracking the goal, choosing paths and representing their value, and consolidating and abstracting from spatial memories. Future computational models that predict the engagement of PFC sub-regions during different events in navigation would be highly beneficial. This combined with empirical testing of these models across species with directly comparable tasks may provide a way forward in helping understand the full contributions of the PFC to navigation [146]. We have outlined important questions arising from this review in the Outstanding Questions box. Some key questions are: a) how do different PFC areas interact with regions in the core navigation brain network to support flexible guidance, b) what neural mechanisms does the PFC use to plan and select navigational goals and c) how does the PFC track information for successful navigation?

**CONFLICT OF INTERESTS:** None.

**AKNOWLEDGEMENTS:** We would like to thank William de Cothi and Nils Nyberg, as well as Kate Jeffrey, for helpful edits and comments on the manuscript.

### **BOX 1: The Role of Prefrontal-Hippocampal Interactions in Navigation**

Evidence supports the idea that PFC-hippocampus engage in coordinated activity [143–145], and in fact point to the role of the prefrontal cortex in driving hippocampal activity in certain situations when the goal states (e.g. current target) need to be retrieved in a context appropriate manner [109,147,148]. The importance of PFC and medial temporal interaction has long been established for memory processes such as encoding and retrieval [6] and for incorporating new information into existing memories or schemas [140], for example via immediate early genes that are activated in the prelimbic cortex of the PFC during hippocampal-dependent memory encoding [141,149]. Simultaneous recordings from the ACC and CA1 in the rodent has also shown that during remote recall the ACC drives hippocampal activity to increase representations of context to facilitate retrieval [147]. One conclusion that follows is that PFC may be important for reducing interference during hippocampal dependent tasks such as navigation. Further supporting this view, it has been found that increased mPFC-hippocampal synchrony resulted in less interference from the previous learned rule in a spatial task requiring rule switching [50]. Crucially the mPFC was not important for initial learning and retrieval, only for the maintenance of rule reversals, and was shown to precede activity in CA1 during early stages of learning, and also when this learning was rapid. This new evidence suggests that the PFC may have a role in facilitating this rule switching process, or even initiating it [147]. Similarly, during early learning of rules concerning reward locations on a W-maze, PFC to CA1 spiking synchrony was increased, even higher than during sleep [150]. For further reviews on PFC-hippocampus interactions see [59,60,62].

### **Glossary**

**Cognitive map:** a theoretical construct describing the neural representation that underlies our ability to remember the layout of the environment and store information about events occurring in places.

**Default-mode network (DMN):** Default-mode network is a set of brain regions, including the medial prefrontal cortex, hippocampus and precuneus/posterior cingulate cortex, which are more active when thought is directed internally, as opposed to towards the external environment.

**Grid cell:** a location-modulated neuron that fires at regular intervals as an animal navigates an open area. Grid cells have been most commonly found in the entorhinal cortex and are thought to form an essential part of the brain's navigation system by allowing it to understand its position in space by storing and integrating information about location, distance, and direction.

**Hidden states:** a collection of information relevant to a given decision that is difficult to distinguish based on sensory input alone and is therefore 'hidden'.

**Place cell:** neurons that become active when an animal enters a particular location within an environment, which are most commonly reported in the hippocampus.

**Replay:** During 'replay', temporally compressed sequences of place cells reactivate spatial trajectories in explored environments in either forward or reverse order. Replay is associated with high-frequency sharp-wave ripple (SWR) events prevalent during offline periods in both sleep and non-exploratory waking states ("awake replay").

**State space:** the set of all possible states of the environment that are relevant for a given task, which for navigation might be the possible paths or enclosed spaces available to navigate.

**Vicarious trial and error (VTE):** a behaviour considered an overt marker of ‘mental planning/exploration’ or deliberation, observed in rodents at a choice point in a maze, where it frequently pauses and alternately faces towards its potential goal, before deciding.

**Table 1: Variety of functions associated with the prefrontal cortex in the context of navigation behaviour.** [note: human-centric organization; italics refer to the structure in the rodent brain; see Figure 2 for more details on nomenclature used]

Brain Region (Human / <i>Rodent</i> )	Proposed Function	Reference
ACC / <i>PL</i>	Effort	[69]
	Grid coding of space	[27]
	Recalling episodic memories of the environment	[147]
	Vicarious Trial-and-Error Behaviour	[112]
dACC / <i>MCC</i>	Hierarchical coding	[8]
	Difficulty of choice (Current and Future)	[14]
	Inhibition of current plans	[12,13]
	Predictive monitoring of the environment and the agents moving in the environment	[105]
	Back-tracking	[13]
	Foraging/Counterfactuals	[133]
Frontopolar cortex (BA10)	Future choice	[14]
	Searching possible paths in future	[11,41,137]
mPFC	Proximity to the goal	[22] [16]
	Replay	[151]
	Path integration	[152]
OFC	Proximity to the goal	[74]
	Determining the decision/state space	[115]
	Grid cells	[153]
	Regret	[154]

SFG/(r)dmPFC (BA8)	Difficulty of Choice (Current - Future, Interaction); Coupling with hippocampus increased when correct choice	[14]
	Novelty detection and environmental changes leading to re-planning (e.g. long detours)	[12,25]
	Path integration (reducing error)	[102]
Right IPFC (vIPFC)	Novelty detection and environmental changes leading to re-planning (in particular long detours and false shortcuts, but not real shortcuts)	[12,25]
Left IPFC (vIPFC)	Difficulty of future choice	[14]
vmPFC	Optimizing search path	[155]
	Scene construction	[73]
	Integration with existing schema	[49,141]
	Maintaining Current Goal	[103]
	Grid-coding	[52] [153]

## REFERENCES

- 1 Epstein, R.A. *et al.* (2017) The cognitive map in humans: Spatial navigation and beyond. *Nat. Neurosci.* 20, 1504–1513
- 2 Ekstrom, A.D. *et al.* (2018) *Human spatial navigation*, Princeton University Press.
- 3 Goodroe, S.C. *et al.* (2018) The Complex Nature of Hippocampal-Striatal Interactions in Spatial Navigation. *Front. Hum. Neurosci.* 12, 1–9
- 4 Gahnstrom, C.J. and Spiers, H.J. (2020) Striatal and hippocampal contributions to flexible navigation in rats and humans. *Brain Neurosci. Adv.* 4, 1-7
- 5 Miller, E.K. and Cohen, J.D. (2001) An integrative theory of prefrontal cortex function. *Annu. Rev. Neurosci.* 24, 167-202
- 6 Simons, J.S. and Spiers, H.J. (2003) Prefrontal and medial temporal lobe interactions in long-term memory. *Nat. Rev. Neurosci.* 4, 637–48
- 7 Anggraini, D. *et al.* (2018) Neural signatures of reinforcement learning correlate with strategy adoption during spatial navigation. *Sci. Rep.* 8, 1–14
- 8 Balaguer, J. *et al.* (2016) Neural Mechanisms of Hierarchical Planning in a Virtual Subway Network. *Neuron* 90, 893–903
- 9 Dahmani, L. and Bohbot, V.D.V.D. (2015) Dissociable contributions of the prefrontal cortex to hippocampus- and caudate nucleus-dependent virtual navigation strategies.

- Neurobiol. Learn. Mem.* 117, 42–50
- 10 Howard, L.R. *et al.* (2014) The hippocampus and entorhinal cortex encode the path and euclidean distances to goals during navigation. *Curr. Biol.* 24, 1331–1340
  - 11 Javadi, A.-H. *et al.* (2017) Hippocampal and prefrontal processing of network topology to simulate the future. *Nat. Commun.* 8, 14652
  - 12 Javadi, A.-H. *et al.* (2019) Prefrontal Dynamics Associated with Efficient Detours and Shortcuts: A Combined Functional Magnetic Resonance Imaging and Magnetoencephalography Study. *J. Cogn. Neurosci.* 31, 1227–1247
  - 13 Javadi, A.-H. *et al.* (2019) Backtracking during navigation is correlated with enhanced anterior cingulate activity and suppression of alpha oscillations and the ‘default-mode’ network. *Proc. R. Soc. B Biol. Sci.* 286, 20191016
  - 14 Kaplan, R. *et al.* (2017) The Neural Representation of Prospective Choice during Spatial Planning and Decisions. *PLOS Biol.* 15, e1002588
  - 15 Simon, D.A. and Daw, N.D. (2011) Neural correlates of forward planning in a spatial decision task in humans. *J. Neurosci.* 31, 5526–5539
  - 16 Viard, A. *et al.* (2011) Anterior Hippocampus and Goal-Directed Spatial Decision Making. *J. Neurosci.* 31, 4613–4621
  - 17 Maguire, E.A. *et al.* (1998) Knowing where and getting there: A human navigation network. *Science.* 280, 921–924
  - 18 Hartley, T. *et al.* (2003) The well-worn route and the path less traveled: Distinct neural bases of route following and wayfinding in humans. *Neuron* 37, 877–888
  - 19 Iaria, G. *et al.* (2008) Detection of unexpected events during spatial navigation in humans: Bottom-up attentional system and neural mechanisms. *Eur. J. Neurosci.* 27, 1017–1025
  - 20 Sherrill, K.R. *et al.* (2013) Hippocampus and retrosplenial cortex combine path integration signals for successful navigation. *J. Neurosci.* 33, 19304–13
  - 21 Spiers, H.J. and Maguire, E.A. (2006) Thoughts, behaviour, and brain dynamics during navigation in the real world. *Neuroimage* 31, 1826–1840
  - 22 Spiers, H.J. and Maguire, E.A. (2007) A Navigational Guidance System in the Human Brain. *Hippocampus* 17, 618-626
  - 23 Shallice, T. and Burgess, P.W. (1991) Deficits in strategy application following frontal lobe damage in man. *Brain* 114, 727–741
  - 24 Spiers, H.J. (2008) Keeping the goal in mind: Prefrontal contributions to spatial navigation. *Neuropsychologia* 46, 2106–2108
  - 25 Spiers, H.J. and Gilbert, S.J. (2015) Solving the detour problem in navigation: a model of prefrontal and hippocampal interactions. *Front. Hum. Neurosci.* 9, 1–15
  - 26 Grieves, R.M. and Jeffery, K.J. (2017) The representation of space in the brain. *Behav. Processes* 135, 113–131

- 27 Jacobs, J. *et al.* (2013) Direct recordings of grid-like neuronal activity in human spatial navigation. *Nat. Neurosci.* 16, 1188–1191
- 28 Ekstrom, A.D. *et al.* (2003) Cellular networks underlying human spatial navigation. *Nature* 425, 184–187
- 29 Watrous, A.J. *et al.* (2013) Frequency-specific network connectivity increases underlie accurate spatiotemporal memory retrieval. *Nat. Neurosci.* 16, 349–56
- 30 Lacroix, L. *et al.* (2002) Effect of excitotoxic lesions of rat medial prefrontal cortex on spatial memory. *Behav. Brain Res.* 133, 69–81
- 31 Sloan, H.L. *et al.* (2006) Double dissociation between hippocampal and prefrontal lesions on an operant delayed matching task and a water maze reference memory task. *Behav. Brain Res.* 171, 116–126
- 32 Bermudez-Contreras, E. *et al.* (2020) The Neuroscience of Spatial Navigation and the Relationship to Artificial Intelligence. *Front. Comput. Neurosci.* 14, 63
- 33 Epstein, R.A. and Vass, L.K. (2014) Neural systems for landmark-based wayfinding in humans. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 369, 20120533
- 34 Hartley, T. *et al.* (2013) Space in the brain: how the hippocampal formation supports spatial cognition. *Philos. Trans. R. Soc. B Biol. Sci.* 369, 20120510
- 35 Moser, E.I. *et al.* (2008) Place cells, grid cells, and the brain's spatial representation system. *Annu. Rev. Neurosci.* 31, 69–89
- 36 Bicanski, A. and Burgess, N. (2018) A neural-level model of spatial memory and imagery. *Elife*, 7:e33752
- 37 Ekstrom, A.D. *et al.* (2014) A critical review of the allocentric spatial representation and its neural underpinnings: toward a network-based perspective. *Front. Hum. Neurosci.* 8, 803
- 38 Uylings, H.B.M. *et al.* (2003) Do rats have a prefrontal cortex? *Behav. Brain Res.* 146, 3–17
- 39 van Heukelum, S. *et al.* (2020) Where is Cingulate Cortex? A Cross-Species View. *Trends Neurosci.* 43, 285–299
- 40 Laubach, M. *et al.* (2018) What, If Anything, Is Rodent Prefrontal Cortex?. *eNeuro.* 5(5), 1–14
- 41 Badre, D. and Nee, D.E. (2018) Frontal Cortex and the Hierarchical Control of Behavior. *Trends Cogn. Sci.* 22, 170–188
- 42 Coutrot, A. *et al.* (2018) Global Determinants of Navigation Ability. *Curr. Biol.* 28, 2861–6
- 43 Brunec, I.K. *et al.* (2019) Cognitive mapping style relates to posterior-anterior hippocampal volume ratio. *29, 748–754*
- 44 Weisberg, S.M. *et al.* (2014) Variations in cognitive maps: Understanding individual differences in navigation. *J. Exp. Psychol. Learn. Mem. Cogn.* 40, 669–682

- 45 Bohbot, V.D. *et al.* (2007) Gray matter differences correlate with spontaneous strategies in a human virtual navigation task. *J. Neurosci.* 27, 10078–10083
- 46 Asem, J.S.A. and Holland, P.C. (2013) Immediate Response Strategy and Shift to Place Strategy in Submerged T-Maze. *Behav. Neurosci.* 127, 854–859
- 47 Patai, E.Z. *et al.* (2019) Hippocampal and Retrosplenial Goal Distance Coding after Long-term Consolidation of a Real-World Environment. *Cereb. Cortex* 29, 2748–2758
- 48 Churchwell, J.C. *et al.* (2010) Prefrontal and hippocampal contributions to encoding and retrieval of spatial memory. *Neurobiol. Learn. Mem.* 93, 415–421
- 49 Farina, F.R. and Commins, S. (2016) Differential expression of immediate early genes Zif268 and c-Fos in the hippocampus and prefrontal cortex following spatial learning and glutamate receptor antagonism. *Behav. Brain Res.* 307, 194–198
- 50 Guise, K.G. and Shapiro, M. (2017) Medial Prefrontal Cortex Reduces Memory Interference by Modifying Hippocampal Encoding. *Neuron* 94, 183–192
- 51 Hok, V. *et al.* (2005) Coding for Spatial Goals in the Prelimbic-Infralimbic. *Proc. Natl. Acad. Sci.* 102, 4602–7
- 52 Doeller, C.F. *et al.* (2010) Evidence for grid cells in a human memory network. *Nature* 463, 657–61
- 53 Watrous, A.J. *et al.* (2018) Phase-tuned neuronal firing encodes human contextual representations for navigational goals. *Elife* 7, 1–16
- 54 Dalton, R.C. *et al.* (2019) Wayfinding as a social activity. *Front. Psychol.* 10, 1–14
- 55 Krueger, F. *et al.* (2009) The medial prefrontal cortex mediates social event knowledge. *Trends Cogn. Sci.* 13, 103–109
- 56 Rudebeck, P.H. *et al.* (2007) Distinct contributions of frontal areas to emotion and social behaviour in the rat. *Eur. J. Neurosci.* 26, 2315–2326
- 57 Spiers, H.J. and Maguire, E.A. (2006) Spontaneous mentalizing during an interactive real world task: An fMRI study. *Neuropsychologia* 44, 1674–1682
- 58 Park, S.A. *et al.* (2020) Map Making: Constructing, Combining, and Inferring on Abstract Cognitive Maps. *Neuron* 107, 1–13
- 59 Pezzulo, G. *et al.* (2019) Planning at decision time and in the background during spatial navigation. *Curr. Opin. Behav. Sci.* 29, 69–76
- 60 Zielinski, M.C. *et al.* (2020) The role of replay and theta sequences in mediating hippocampal-prefrontal interactions for memory and cognition. *Hippocampus* 30, 60–72
- 61 Redish, A.D. (2016) Vicarious trial and error. *Nat. Rev. Neurosci.* 17, 147–159
- 62 Ito, H.T. (2018) Prefrontal–hippocampal interactions for spatial navigation. *Neurosci. Res.* 129, 2–7
- 63 Ivey, R. *et al.* (2011) A neuromorphic model of spatial lookahead planning. *Neural Networks* 24, 257–266

- 64 Gorchetchnikov, A. and Hasselmo, M.E. (2005) A biophysical implementation of a bidirectional graph search algorithm to solve multiple goal navigation tasks. *Conn. Sci.* 17, 145–164
- 65 Martinet, L.E. *et al.* (2011) Spatial learning and action planning in a prefrontal cortical network model. *PLoS Comput. Biol.* 7(5): e1002045
- 66 Economides, M. *et al.* (2014) Anterior cingulate cortex instigates adaptive switches in choice by integrating immediate and delayed components of value in ventromedial prefrontal cortex. *J. Neurosci.* 34, 3340–9
- 67 Kolling, N. *et al.* (2012) Neural mechanisms of foraging. *Science.* 335, 95–8
- 68 Croxson, P.L. *et al.* (2009) Effort-Based Cost–Benefit Valuation and the Human Brain. *J. Neurosci.* 29, 4531–4541
- 69 Porter, B.S. *et al.* (2019) Anterior cingulate cortex encoding of effortful behavior. *J. Neurophysiol.* 121, 701–714
- 70 Spiers, H.J. and Maguire, E.A. (2008) The dynamic nature of cognition during wayfinding. *J. Environ. Psychol.* 28, 232–249
- 71 Monk, A.M. *et al.* (2020) The role of hippocampal-vmPFC neural dynamics in building mental representations. *J. Cogn. Neurosci.* 33, 89-103
- 72 McCormick, C. *et al.* (2018) Comparing and Contrasting the Cognitive Effects of Hippocampal and Ventromedial Prefrontal Cortex Damage: A Review of Human Lesion Studies. *Neuroscience* 374, 295–318
- 73 Barry, D.N. *et al.* (2019) The neural dynamics of novel scene imagery. *J. Neurosci.* 39, 4375–4386
- 74 Brown, T.I. *et al.* (2016) Prospective representation of navigational goals in the human hippocampus. *Science.* 352, 1323–6
- 75 Wilson, B. *et al.* (1998) The Development of an Ecologically Valid Test for Assessing Patients with a Dysexecutive Syndrome. *Neuropsychol. Rehabil.* 8(3), 213-28
- 76 Bird, C.M. *et al.* (2004) The impact of extensive medial frontal lobe damage on ‘Theory of Mind’ and cognition. *Brain* 127, 914–928
- 77 Mushiake, H. *et al.* (2006) Activity in the Lateral Prefrontal Cortex Reflects Multiple Steps of Future Events in Action Plans. *Neuron* 50, 631–641
- 78 Saito, N. *et al.* (2005) Representation of Immediate and Final Behavioral Goals in the Monkey Prefrontal Cortex during an Instructed Delay Period. *Cereb. Cortex* 15, 1535–46
- 79 Tanji, J. and Hoshi, E. (2008) Role of the lateral prefrontal cortex in executive behavioral control. *Physiol. Rev.* 88, 37–57
- 80 Desrochers, T.M. *et al.* (2015) The Necessity of Rostrolateral Prefrontal Cortex for Higher-Level Sequential Behavior. *Neuron* 87, 1357–68
- 81 Foster, D.J. (2017) Replay Comes of Age. *Annu. Rev. Neurosci.* 40, 581–602


- 82 Ólafsdóttir, H.F. *et al.* (2015) Hippocampal place cells construct reward related sequences through unexplored space. *Elife* 4, 1–17
- 83 Spiers, H.J. and Barry, C. (2015) Neural systems supporting navigation. *Curr. Opin. Behav. Sci.* 1, 47–55
- 84 Gupta, A.S. *et al.* (2010) Hippocampal replay is not a simple function of experience. *Neuron* 65, 695–705
- 85 Pfeiffer, B.E. (2020) The content of hippocampal “replay.” *Hippocampus* 30, 6–18
- 86 Wikenheiser, A.M. and Redish, A.D. (2015) Hippocampal theta sequences reflect current goals. *Nat. Neurosci.* 18, 1–8
- 87 Kay, K. *et al.* (2020) Constant Sub-second Cycling between Representations of Possible Futures in the Hippocampus. *Cell* 180, 552–567
- 88 Kurth-Nelson, Z. *et al.* (2016) Fast Sequences of Non-spatial State Representations in Humans. *Neuron* 91, 194–204
- 89 Liu, Y. *et al.* (2019) Human Replay Spontaneously Reorganizes Experience. *Cell* 178, 640–652
- 90 Shin, J.D. *et al.* (2019) Dynamics of Awake Hippocampal-Prefrontal Replay for Spatial Learning and Memory-Guided Decision Making. *Neuron* 104, 1110–25
- 91 Kaplan, R. *et al.* (2014) Medial prefrontal theta phase coupling during spatial memory retrieval. *Hippocampus* 24, 656–65
- 92 Kaplan, R. *et al.* (2017) Medial Prefrontal–Medial Temporal Theta Phase Coupling in Dynamic Spatial Imagery. *J. Cogn. Neurosci.* 29, 509–519
- 93 Kaplan, R. *et al.* (2016) Hippocampal Sharp-Wave Ripples Influence Selective Activation of the Default Mode Network. *Curr. Biol.* 26, 686–691
- 94 Tsitsiklis, M. *et al.* (2020) Single-Neuron Representations of Spatial Targets in Humans. *Curr. Biol.* 30, 245–253
- 95 Poucet, B. and Hok, V. (2017) Remembering goal locations. *Curr. Opin. Behav. Sci.* 17, 51–56
- 96 Jung, M.W. *et al.* (1998) Firing characteristics of deep layer neurons in prefrontal cortex in rats performing spatial working memory tasks. *Cereb. Cortex* 8, 437–450
- 97 Mashhoori, A. *et al.* (2018) Rat anterior cingulate cortex recalls features of remote reward locations after disfavoured reinforcements. *Elife* 7, 1–18
- 98 Ito, H.T. *et al.* (2015) A prefrontal-thalamo-hippocampal circuit for goal-directed spatial navigation. *Nature* 522, 50–55
- 99 Ma, L. *et al.* (2014) Tracking progress toward a goal in corticostriatal ensembles. *J. Neurosci.* 34, 2244–53
- 100 Young, J.J. and Shapiro, M.L. (2011) Dynamic coding of goal-directed paths by orbital prefrontal cortex. *J. Neurosci.* 31, 5989–6000

- 101 Chrastil, E.R. *et al.* (2015) There and Back Again: Hippocampus and Retrosplenial Cortex Track Homing Distance during Human Path Integration. *J. Neurosci.* 35, 15442–452
- 102 Wolbers, T. *et al.* (2007) Differential recruitment of the hippocampus, medial prefrontal cortex, and the human motion complex during path integration in humans. *J. Neurosci.* 27, 9408–16
- 103 Ciaramelli, E. (2008) The role of ventromedial prefrontal cortex in navigation: A case of impaired wayfinding and rehabilitation. *Neuropsychologia* 46, 2099–2105
- 104 Shidara, M. and Richmond, B.J. (2002) Anterior Cingulate: Single Neuronal Signals Related to Degree of Reward Expectancy. *Science.* 296, 1709-11
- 105 Yoo, S.B.M. *et al.* (2020) The neural basis of predictive pursuit. *Nat. Neurosci.* 23,252-9.
- 106 Böhm, C. and Lee, A.K. (2020) Canonical goal-selective representations are absent from prefrontal cortex in a spatial working memory task requiring behavioral flexibility. *Elife* 9:e63035,
- 107 Jones, M.W. and Wilson, M.A. (2005) Theta rhythms coordinate hippocampal-prefrontal interactions in a spatial memory task. *PLoS Biol.* 3, e402
- 108 Tavares, L.C.S. and Tort, A.B.L. (2020) Hippocampal-Prefrontal Interactions during Spatial Decision-Making. *bioRxiv* DOI: 10.1101/2020.06.24.168732
- 109 Hasz, B.M. and Redish, A.D. (2020) Dorsomedial prefrontal cortex and hippocampus represent strategic context even while simultaneously changing representation throughout a task session. *Neurobiol. Learn. Mem.* 171, 107215
- 110 Hu, D. and Amsel, A. (1995) A simple test of the vicarious trial-and-error hypothesis of hippocampal function. *Proc. Natl. Acad. Sci. U. S. A.* 92, 5506–09
- 111 Papale, A.E. *et al.* (2016) Interplay between Hippocampal Sharp-Wave-Ripple Events and Vicarious Trial and Error Behaviors in Decision Making. *Neuron* 92, 975–982
- 112 Schmidt, B. *et al.* (2019) Disrupting the medial prefrontal cortex alters hippocampal sequences during deliberative decision making. *J. Neurophysiol.* 121, 1981–2000
- 113 Wikenheiser, A.M. and Schoenbaum, G. (2016) Over the river, through the woods: Cognitive maps in the hippocampus and orbitofrontal cortex. *Nat. Rev. Neurosci.* 17, 513–523
- 114 Sharpe, M.J. *et al.* (2019) An Integrated Model of Action Selection: Distinct Modes of Cortical Control of Striatal Decision Making. *Annu. Rev. Psychol.* 70, 53–76
- 115 Schuck, N.W. *et al.* (2016) Human Orbitofrontal Cortex Represents a Cognitive Map of State Space. *Neuron* 91, 1402–1412
- 116 Elliott Wimmer, G. and Büchel, C. (2019) Learning of distant state predictions by the orbitofrontal cortex in humans. *Nat. Commun.* 10, 1–11
- 117 Vertechi, P. *et al.* (2020) Inference-Based Decisions in a Hidden State Foraging Task: Differential Contributions of Prefrontal Cortical Areas. *Neuron* 106, 166-176

- 118 Bradfield, L.A. *et al.* (2015) Medial Orbitofrontal Cortex Mediates Outcome Retrieval in Partially Observable Task Situations. *Neuron* 88, 1268–1280
- 119 Miller, K.J. *et al.* (2020) Value Representations in the Rodent Orbitofrontal Cortex Drive Learning, not Choice. *bioRxiv* DOI: 10.1101/245720
- 120 Kennerley, S.W. *et al.* (2011) Double dissociation of value computations in orbitofrontal and anterior cingulate neurons. *Nat. Neurosci.* 14, 1581–9
- 121 Turner, C.H. (1912) Sphex Overcoming Obstacles. *Psyche (New York)* 19, 100–2
- 122 Rauchs, G. *et al.* (2008) Partially segregated neural networks for spatial and contextual memory in virtual navigation. *Hippocampus* 18, 503–518
- 123 Rosenbaum, R.S. *et al.* (2004) “I have often walked down this street before”: fMRI studies on the hippocampus and other structures during mental navigation of an old environment. *Hippocampus* 14, 826–835
- 124 Fletcher, P.C. *et al.* (2001) Responses of human frontal cortex to surprising events are predicted by formal associative learning theory. *Nat. Neurosci.* 4, 1043–8
- 125 Gläscher, J. *et al.* (2010) States versus rewards: Dissociable neural prediction error signals underlying model-based and model-free reinforcement learning. *Neuron* 66, 585–595
- 126 Aron, A.R. (2011) From reactive to proactive and selective control: Developing a richer model for stopping inappropriate responses. *Biol. Psychiatry* 69, e55–e68
- 127 Alvernhe, A. *et al.* (2008) Different CA1 and CA3 Representations of Novel Routes in a Shortcut Situation. *J. Neurosci.* 28, 7324–33
- 128 Tolman, E.C. (1948) Cognitive maps in rats and men. *Psychol. Rev.* 55, 189–208
- 129 Hanslmayr, S. *et al.* (2016) Oscillations and episodic memory – Addressing the synchronization/desynchronization conundrum. *Trends Neurosci.* 39, 16–25
- 130 Wystrach, A. *et al.* (2013) Backtracking behaviour in lost ants: an additional strategy in their navigational toolkit. *Proc. Biol. Sci.* 280, 20131677
- 131 Yoshida, W. and Ishii, S. (2006) Resolution of Uncertainty in Prefrontal Cortex. *Neuron* 50, 781–9
- 132 White, J.K. *et al.* (2019) A neural network for information seeking. *Nat. Commun.* 10, 1–19
- 133 Fouragnan, E.F. *et al.* (2019) The macaque anterior cingulate cortex translates counterfactual choice value into actual behavioral change. *Nat. Neurosci.* 22, 797–808
- 134 Schuck, N.W. *et al.* (2015) Medial prefrontal cortex predicts internally driven strategy shifts. *Neuron* 86, 331–340
- 135 Heilbronner, S.R. and Hayden, B.Y. (2016) Dorsal Anterior Cingulate Cortex: A Bottom-Up View. *Annu. Rev. Neurosci.* 39, 149–170
- 136 Nobre, A.C. *et al.* (2012) Nervous Anticipation : Top-Down Biasing across Space and


- Time. In *Cognitive neuroscience of attention* (2nd edn) (Posner, M., ed), Guilford Press
- 137 Burgess, P.W. *et al.* (2007) The gateway hypothesis of rostral prefrontal cortex (area 10) function. *Trends Cogn. Sci.* 11, 290–8
  - 138 Gunseli, E. and Aly, M. (2020) Preparation for upcoming attentional states in the hippocampus and medial prefrontal cortex. *Elife* 9, 1–33
  - 139 Sánchez-Bellot, C. and MacAskill, A.F. (2020) Two opposing hippocampus to prefrontal cortex pathways for the control of approach and avoidance behavior. *bioRxiv* 2019.12.18
  - 140 Alonso, A. *et al.* (2020) Naïve to expert : considering the role of previous knowledge in memory. *Brain and Neuroscience Advances.* 4: 1 –17
  - 141 Tse, D. *et al.* (2011) Schema-dependent gene activation and memory encoding in neocortex. *Science.* 333, 891–5
  - 142 Wang, J.X. *et al.* (2018) Prefrontal cortex as a meta-reinforcement learning system. *Nat. Neurosci.* 21, 860–8
  - 143 Tang, W. *et al.* (2017) Hippocampal-prefrontal reactivation during learning is stronger in awake compared with sleep states. *J. Neurosci.* 37, 11789–11805
  - 144 Zielinski, M.C. *et al.* (2019) Coherent coding of spatial position mediated by theta oscillations in the hippocampus and prefrontal cortex. *J. Neurosci.* 39, 4550–65
  - 145 Jadhav, S.P.P. *et al.* (2016) Coordinated Excitation and Inhibition of Prefrontal Ensembles during Awake Hippocampal Sharp-Wave Ripple Events. *Neuron* 90, 113–127
  - 146 De Cothi, W. *et al.* (2020) Predictive Maps in Rats and Humans for Spatial Navigation. *bioRxiv* at <<https://doi.org/10.1101/2020.09.26.314815>>
  - 147 Wirt, R.A. and Hyman, J.M. (2019) ACC Theta Improves Hippocampal Contextual Processing during Remote Recall. *Cell Rep.* 27, 2313-27
  - 148 Place, R. *et al.* (2016) Bidirectional prefrontal-hippocampal interactions support context-guided memory. *Nat. Neurosci.* 19, 992–4
  - 149 Wang, S.H. *et al.* (2012) Anterior cingulate cortex in schema assimilation and expression. *Learn. Mem.* 19, 315–8
  - 150 Tang, W. and Jadhav, S.P.P. (2019) Sharp-wave ripples as a signature of hippocampal-prefrontal reactivation for memory during sleep and waking states. *Neurobiol. Learn. Mem.* 160, 11–20
  - 151 Peyrache, A. *et al.* (2009) Replay of rule-learning related neural patterns in the prefrontal cortex during sleep. *Nat. Neurosci.* 12, 919–926
  - 152 Blankenship, P.A. *et al.* (2016) The medial frontal cortex contributes to but does not organize rat exploratory behavior. *Neuroscience* 336, 1–11
  - 153 Constantinescu, A.O. *et al.* (2016) Organizing conceptual knowledge in humans with a gridlike code. *Science.* 352, 1464–8

- 154 Steiner, A.P. and Redish, A.D. (2014) Behavioral and neurophysiological correlates of regret in rat decision-making on a neuroeconomic task. *Nat. Neurosci.* 17, 995–1002
- 155 Carrieri, M. *et al.* (2018) Does ventrolateral prefrontal cortex help in searching for the lost key? Evidence from an fNIRS study. *Brain Imaging Behav.* 12, 785–797
- 156 Carlén, M. (2017) What constitutes the prefrontal cortex? *Science.* 358, 478–482




### FIGURE 1: The Prefrontal Cortex is Active During Navigation

(A) Data from [17], earliest study to show increased PFC during virtual navigation. Activity in the middle and superior frontal gyrus is increased when navigating to learned goal in a virtual town where novel barriers forced detours to the goal. (B) Increased activity in medial, ventrolateral, dorsolateral, and right anterior prefrontal cortices during wayfinding to goals compared to following a well-learned route to a goal [18]. (C) Middle and left superior frontal gyrus were more active when participants thought about the route around a blocked path in their hometown [123]. (D) Prelimbic cortex (PrL), a proposed rodent analogue of part of human PFC, shows increased immediate early gene (IEG) counts when animals must integrate new memories into an existing spatial memory schema (NPA), compared to when they navigate to learned ('old') object locations (OPA) or form new maps of the space (NM). An example rodent brain (from [39]) shown with vertical slice indicating location of PrL and exact slice location are shown [141]. (E) human neuropsychological study, where a patient with ventromedial prefrontal damage was able to find his way between two locations in his hometown if he was given reminders (either of the goal location itself, or a prompt to rehearse the location), however not when given no reminders or given a meaningless one (a warning beep), pointing to a role of the vmPFC in maintaining the goal in mind while wayfinding [103].



**Figure 2: Anatomical Summary & Different Nomenclatures**


A) Functional definition of brain areas (human). B) Brodmann areas in the human frontal cortex. C) Functional definition of brain areas (rodent). Nomenclature used in text: *(d)ACC*: (dorsal) anterior cingulate cortex (also used in rodent literature - better referred to as *MCC*, see [39]), *dmPFC*: dorsal medial prefrontal cortex, *lIPFC*: left lateral prefrontal cortex, *OFC*: orbitofrontal cortex (also used in rodent literature), *PrL(PL)*: Prelimbic cortex (rodent – often also termed ‘*mPFC*’ or ‘*ACC*’, see [39]), *rIPFC*: right lateral prefrontal cortex, *SFG/rdmPFC*: superior frontal gyrus/rostradorsal medial prefrontal cortex, *vmPFC*: ventral medial prefrontal cortex. Reproduced with



permission [156]

**FIGURE 3: Examples of Prefrontal Activity to Specific Events During Navigation**  
 Neural activity changes as a function of the behavioural correlate are represented by the schematic. Route planning has been shown to involve responses in prefrontal cortex: (A) in an unstable ever-changing environment [15], a simplified maze [14], (B) film simulation of a real city [10], (C) in a virtual subway network [8] (D) or virtual render of a city [21]. (E) Macaques performed a multi-step maze task and before the onset of movement, all future actions (three cursor motions) within the maze were selectively encoded in the population firing rate of dorsolateral PFC (BA46). These neurons did not code for the physical actions (arm movements) that were decoupled from the steps within the maze [77]. (F) View in a virtual desert island, in which shifting paths of lava create shortcuts and detours to hidden treasure [18]. Activity in superior frontal gyrus and right lateral prefrontal cortex is increased when detours adding a larger distance to the route were compared with detours adding a short distance. Importantly rIPFC was also activated by false shortcuts, i.e. changes in route options that seem like a shortcut but will in fact lead the participant away from the goal [12]. (G) Dorsal anterior cingulate (dACC) was active when participants engaged in 'backtracking', i.e. voluntarily changing their path (inset shows an example of such a backtracking event overlaid on a portion of the maze from F, white: walkable paths; grey: path boundaries)

[13]. (H) The medial prefrontal cortex codes path distance to the goal location during initial planning [16] and during travel periods [22]. (I) The anterior cingulate cortex codes path distance during travel periods [10].



### KEY FIGURE: Summary of Expected Human Prefrontal Engagement During Navigation

Making a trip from a cozy pub in London’s Soho, we first recall where our goal is located - medial PFC interacts with the hippocampus to support recall and mental scene construction of the goal (1). Accessing possible routes and comparing them requires dynamic interactions within dmPFC/ dACC and lateral PFC (1). En route, a barrier is spotted that elicits enhanced activity in the superior frontal gyrus, right lateral PFC and dACC, halting the current plan and driving new planning (2). Inferior lateral PFC supports a search through the different route options to select a new path (3). We encounter a path that appears to provide a potential shortcut to the goal. However, retrieval and simulation of the future route leads to the conclusion that this is a waste of our time, with right lateral PFC engaged to suppress choosing this path (4). The OFC updates representations of the usefulness of paths (including inferred states for the false shortcut just avoided). OFC activity helps process rule-based state information about what paths are possible when we are walking rather than driving –

modulating the affordances of different paths. While coasting along the pedestrian bridge, we have no immediate actions to plan or take, so the default-mode network becomes active (6). As we approach the goal region, medial PFC and anterior cingulate cortex code the path distance to our goal (7). Having gone too far down the riverside walk, dACC activity plays a role in changing the current plan, resulting in backtracking to a previous point in order to correct our route (8). Finally, we arrive at the Hayward Gallery, and enjoy another gin and tonic before experiencing the art.