Table 1 – Included studies stratified by type of attitudes/stigma and region

		type of attitudes/st	r · · · · · · · · · · · · · · · · · · ·	T		T
	Asia and Oceania	Africa	East Mediterranean	Europe	Latin America	North America
	N = 46 studies	N = 40	N = 16	N = 44	N = 14	N = 25
Attitudes towards r	person with epilepsy and ena	acted stigma				
Sample size	Mean = 643.65 Range 10-2610	Mean = 503.03 (excluding Grunitzky) Range = 66-3256	Mean = 1754.06 Range = 121-16500	Mean = 641.29 Range = 22-3875	Mean = 396.36 Range = 18-1850	Mean = 1291.72 Range = 27-19441
Study type	Population based = 19 Non population based = 13 Unclear or Unspecified = 14	Population based = 16 Non population based = 12 Unclear or Unspecified = 12	Population based = 5 Non population based = 7 Unclear or Unspecified = 4	Population based = 12 Non population based = 17 Unclear or Unspecified = 15	Population based = 2 Non population based = 10 Unclear or Unspecified = 2	Population based = 1 Non population based = 9 Unclear or Unspecified = 15
Target respondents	Persons with epilepsy = 5 General public = 23 Students = 4 Teachers = 6 Others = 8	Persons with epilepsy = 2 General public = 13 Students = 6 Teachers = 7 Others = 12	Persons with epilepsy = 0 General public = 8 Students = 2 Teachers = 2 Others = 4	Persons with epilepsy = 2 General public = 20 Students = 4 Teachers = 6 Others = 12	Persons with epilepsy = 1 General public = 1 Students = 6 Teachers = 3 Others = 3	Persons with epilepsy = 2 General public = 4 Students = 2 Teachers = 6 Others = 11
Age group of respondents	Children = 3 Adults = 34 Both = 7 Not specified = 2	Children = 3 Adults = 25 Both = 10 Not specified = 2	Children = 2 Adults = 12 Both = 1	Children = 3 Adults = 32 Both = 7 Not Specified = 2	Children = 2 Adults = 10 Both = 2	Children = 4 Adults = 20 Both = 1
Ascertainment method	Quantitative = 43 - Validated = 15 - Not Validated/Unclear = 28 Qualitative = 3	Quantitative = 39 - Validated = 4 - Not Validated/Unclear = 35 Qualitative = 1	Quantitative = 14 - Validated = 6 - Not Validated/Unclear = 8 Qualitative = 0	Quantitative = 42 - Validated = 5 - Not Validated/Unclear = 37 Qualitative = 2	Quantitative = 14 - Validated = 5 - Not Validated/Unclear = 9 Qualitative = 0	Quantitative = 24 - Validated = 8 - Not Validated/Unclear = 16 Qualitative = 1
Validated scales used	 Public Attitudes Towards Epilepsy Scale Implicit Association Test Stigma Scale for Epilepsy Scale of Attitudes Toward Persons with Epilepsy (ATPE) Knowledge, attitude and practice towards epilepsy Malay Awareness, Knowledge, and Attitudes Epilepsy 	The semantic differential The behavioral differential of Triandis	- Epilepsy Attitude Scale	Stigmatization of patients with epilepsy questionnaire Attitudes and Beliefs about Living with Epilepsy (ABLE) scale	KAP Questionnaire (Knowledge, Attitude and Perception) Stigma Scale of Epilepsy	 Stigma Scale of Epilepsy Epilepsy Beliefs and Attitudes Scale Attitudes Towards Persons with Epilepsy (ATPE) Fishbein Expectancy-Value Model of Attitudes
Attitudes towards self	f and felt stigma					
	Asia and Oceania N = 26	Africa N = 14	East Mediterranean N = 2	Europe N = 24	Latin America N = 10	North America N = 32
Sample size	Mean = 130 Range = 12-530	Mean = 152.79 Range = 4-423	Mean = 1946 Range = 45-3847	Mean = 771.58 Range = 10-6156	Mean = 364.1 Range = 30-1850	Mean = 220.22 Range = 15-1023
Study type	Population based = 0 Non population based = 24 Unclear or Unspecified = 2	Population based = 1 Non population base = 11 Unclear or unspecified = 2	Not population based = 1 Unclear or unspecified = 1	Population based = 3 Not population based = 17 Unclear or unspecified = 4	Population based = 1 Non population based = 8 Unclear or unspecified = 1	Population based = 2 Not population based = 18 Unclear or unspecified = 12
Target respondents	Persons with epilepsy = 18 Others = 8	Persons with epilepsy = 11 Other = 3	Persons with epilepsy = 2	Persons with epilepsy = 22 Other = 2	Persons with epilepsy = 7 General Public = 1 Students = 1 Other = 1	Persons with epilepsy = 27 Other = 5

A C	Children = 4	Children = 5	Adults = 2	Adults = 20	Children = 2	Children = 6
Age group of	Adults = 19	Adults = 9	Addits = 2	Both = 4	Adults = 5	Adults = 22
respondents	Both $= 3$	Adults = 7		Doul = 4	Both = 3	Both = 4
Ascertainment	Ouantitative = 20	Ouantitative = 13	Ouantitative = 2	Ouantitative = 23	Ouantitative = 10	Quantitative = 32
	- Validated = 17	- Validated = 10	- Validated = 1	- Validated = 17	- Validated = 7	- Validated = 23
method	- Not Validated/Unclear = 3	- Not Validated/Unclear = 3	- Not Validated/Unclear = 1	- Not Validated/Unclear = 6	- Not Validated/Unclear = 3	- Not Validated/Unclear = 9
	Oualitative = 6	Oualitative = 1	Oualitative = 0	Oualitative = 1	Oualitative = 0	Oualitative = 0
Validated scales used	Stigma Scale in Epilepsy Epilepsy Stigma Scale Child Stigma Scale Parents Stigma Scale Epilepsy Surgery Inventory- 55 questionnaire	- Family Interview Schedule - Stigma Scale - Kilifi Stigma Scale - Epilepsy Stigma Scale	- Jacoby stigma scale	Stigma Scale for people with intellectual impairment Stigma Scale of Epilepsy Epilepsy Attitudes Scale Epilepsy Stigma Scale Jacoby Stigma Scale	- Stigma Scale of Epilepsy	- Child Attitude Toward Illness Scale (CATIS) - Child Stigma Scale - Epilepsy Stigma Scale - Jacoby Stigma Scale - Ryan (1980) scale - Link (1991) scale - Parent Stigma Scale - Liverpool Stigma Scale - Adult Stigma Scale - Perceived Stigma Scale - Mittan scale - Stigma Scale for Chronic Illness
Attitudes towards bot	th, and Felt/Enacted Stigma					inics
	Asia and Oceania	Africa	East Mediterranean	Europe	Latin America	North America
	N = 11	N = 7	N = 1	N = 20	N = 5	N = 5
Sample size	Mean = 579.36	Mean = 157.86	533 participants	Mean = 228.7	Mean = 243.8	Mean = 638.8
Sample Size	Range = $59-2275$	Range = 16-338		Range = 19-696	Range = 100-498	Range = 9-2931
Study type	Population based = 2 Not population based = 4 Case-control study = 1 Unclear or unspecified = 4	Not population based = 4 Unclear or unspecified = 3	Not population based = 1	Population based = 1 Not population based = 16 Unclear or unspecified = 3	Not population based = 5	Not population based = 2 Unclear or unspecified = 3
Target respondents	Persons with epilepsy = 8 Other = 3	Persons with epilepsy = 5 Other = 2	Persons with epilepsy and parents/caregivers = 1	Persons with Epilepsy = 15 Other = 5	Persons with Epilepsy = 4 Other = 1	Persons with epilepsy = 3 Other = 2
Age group of respondents	Adults = 10 Children and Adults = 1	Adults = 5 Children and Adults = 2	Children and Adults = 1	Children = 2 Adults = 10 Both = 5 Not specified = 3	Children = 1 Adults = 3 Both = 1	Adults = 3 Children and Adults = 2
Ascertainment method	Quantitative = 9 - Validated = 4 - Not Validated/Unclear = 5 Qualitative = 2	Quantitative = 6 - Validated = 2 - Not Validated/Unclear = 4 Qualitative = 1	Quantitative = 1 - Not Validated/Unclear = 1 Qualitative = 0	Quantitative = 18 - Validated = 6 - Not Validated/Unclear = 12 Qualitative = 2	Quantitative = 5 - Validated = 1 - Not Validated/Unclear = 4 Qualitative = 0	Quantitative = 3 - Validated = 0 - Not Validated/Unclear = 3 Qualitative = 2
Validated scales used	Knowledge Attitudes and Practice Questionnaire Jacoby Stigma Scale	- Jacoby Stigma Scale	None	- Self-esteem scale - Jacoby Stigma Scale - Child Stigma Scale	Stigma Scale of Epilepsy	None