

Childhood maltreatment, educational attainment and IQ: findings from a multicentric case-control study of first-episode psychosis (EU-GEI)

MALTREATMENT, EDUCATION AND IQ IN FEP AND CONTROLS

Lucia Sideli^{1,2*}, Adriano Schimmenti³, Daniele La Barbera⁴, Caterina La Cascia⁴, Laura Ferraro⁴, Monica Aas^{2,5,6}, Luis Alameda^{2,7}, Eva Velthorst^{8,9,10}, Helen L. Fisher^{11,12}, Vincenzo Caretti¹, Giulia Trotta², Giada Tripoli^{2,4}, Diego Quattrone¹¹, Charlotte Gayer-Anderson^{11,13}, Fabio Seminerio⁴, Crocettarachele Sartorio⁴, Giovanna Marrazzo⁴, Antonio Lasalvia¹⁴, Sarah Tosato¹⁴, Iaria Tarricone¹⁵, Domenico Berardi¹⁶, Giuseppe D'Andrea¹⁶, EU-GEI WP2 Group*, Celso Arango¹⁷, Manuel Arrojo¹⁸, Miguel Bernardo¹⁹, Julio Bobes²⁰, Julio Sanjuán²¹, Jose Luis Santos²², Paulo Rossi Menezes²³, Cristina Marta Del-Ben²⁴, Hannah E. Jongsma^{25,26}, Peter B. Jones^{27,28}, James B. Kirkbride²⁵, Pierre-Michel Llorca²⁸, Andrea Tortelli²⁹, Baptiste Pignon^{30,31,32}, Lieuwe de Haan³³, Jean-Paul Selten^{34,35}, Jim Van Os^{2,35,36}, Bart P. Rutten³⁵, Marta Di Forti¹¹, Craig Morgan^{12,13} and Robin M. Murray²

¹Department of Human Science, LUMSA University, Rome, Italy; ²King's College London, Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience, London, England; ³Faculty of Human and Social Sciences, UKE - Kore University of Enna, Enna, Italy; ⁴Department of Biomedicine, Neuroscience, and Advanced Diagnostic, University of Palermo, Palermo, Italy; ⁵NORMENT, Centre for Research On Mental Disorders, Oslo University Hospital and University of Oslo, Norway; ⁶Department of Mental Health Research and Development, Division of Mental Health and Addiction, Vestre Viken Hospital Trust, Norway; ⁷Centro Investigación Biomedica en Red de Salud Mental (CIBERSAM); Instituto de Biomedicina de Sevilla (IBIS), Hospital Universitario Virgen del Rocío, Departamento de Psiquiatría, Universidad de Sevilla, Sevilla, Spain; ⁸Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, USA; ⁹Seaver Autism Center for Research and Treatment, Icahn School of Medicine at Mount Sinai, New York, USA; ¹⁰Early Psychosis Section, Department of Psychiatry, Academic Medical Centre, University of Amsterdam, Amsterdam, The Netherlands; ¹¹King's College London, Social, Genetic, & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, London, England; ¹²ESRC Centre for Society and Mental Health, King's College London, London, UK; ¹³King's College London, Department of Health Services and Population Research, Institute of Psychiatry, Psychology & Neuroscience, London, England; ¹⁴Section of Psychiatry, Department of Neuroscience, Biomedicine and Movement, University of Verona, Verona, Italy; ¹⁵Department of Medical and Surgical Sciences, Alma Mater Studiorum - Bologna University; ¹⁶Department of Biomedical and NeuroMotor Sciences, Psychiatry Unit, Alma Mater Studiorum Università di Bologna, Bologna, Italy; ¹⁷Institute of Psychiatry and Mental Health, Department of Child and Adolescent Psychiatry, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, ISGM, CIBERSAM, Madrid, Spain; ¹⁸Department of Psychiatry, Psychiatric Genetic Group, Instituto de Investigación Sanitaria de Santiago de Compostela, Complejo

Hospitalario Universitario de Santiago de Compostela, Santiago, Spain; ¹⁹Barcelona Clinic Schizophrenia Unit, Hospital Clinic, Department of Medicine, Neuroscience Institute, University of Barcelona, Institut d'Investigacions Biomèdiques, August Pi I Sunyer, Centro de Investigación Biomédica en Red de Salud Mental, Barcelona, Spain; ²⁰Department of Medicine, Psychiatry Area, Universidad de Oviedo, ISPA, INEUROPA, CIBERSAM, Oviedo, Spain; ²¹Department of Psychiatry, School of Medicine, Universidad de Valencia, Centro de Investigación Biomédica en Red de Salud Mental, Valencia, Spain; ²²Department of Psychiatry, Hospital "Virgen de la Luz", Cuenca, Spain; ²³Department of Preventive Medicine, Faculdade de Medicina, Universidade de São Paulo, São Paulo, Brazil; ²⁴Division of Psychiatry, Department of Neuroscience and Behaviour, Ribeirão Preto Medical School, Universidade de São Paulo, São Paulo, Brazil; ²⁵PsyLife Group, Division of Psychiatry, University College London, London, England; ²⁶Department of Psychiatry, University of Cambridge, Cambridge, England; ²⁷CAMEO Early Intervention Service, Cambridgeshire and Peterborough National Health Service Foundation Trust, Cambridge, England; ²⁸EA 7280 Npsydo, Université Clermont Auvergne, Clermont-Ferrand, France; ²⁹Établissement Public de Santé, Maison Blanche, Paris, France; ³⁰AP-HP, Groupe Hospitalier "Mondor," Pôle de Psychiatrie, Créteil, France; ³¹Institut National de la Santé et de la Recherche Médicale, U955, Créteil, France; ³²Fondation Fondamental, Créteil, France; ³³Early Psychosis Section, Department of Psychiatry, Amsterdam UMC, Amsterdam, The Netherlands; ³⁴Institute for Mental Health, GGZ Rivierduinen, Leiden, The Netherlands; ³⁵Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, Maastricht University Medical Centre, Maastricht, The Netherlands; ³⁶Department Psychiatry, Utrecht University Medical Centre, Utrecht, The Netherlands.

*GROUP INFORMATION: The European Network of National Schizophrenia Networks Studying Gene-Environment Interactions (EU-GEI) WP2 Group members includes:

Amoretti, Silvia, PhD, Barcelona Clínic Schizophrenia Unit, Neuroscience Institute, Hospital Clínic of Barcelona, Barcelona, Spain; Bipolar and Depressive Disorder Unit, Neuroscience Institute, Hospital Clínic de Barcelona, Barcelona, Spain; Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Spain; Group of Psychiatry, Mental Health and Addictions, Psychiatric Genetics Unit, Vall d'Hebron Research Institute (VHIR), Barcelona, Spain; **Andreu-Bernabeu, Álvaro**, Child and Adolescent Psychiatry Department, Hospital General Universitario Gregorio Marañón, Universidad Complutense, Madrid, Spain; **Baudin, Grégoire**, MSc, AP-HP, Groupe Hospitalier "Mondor," Pôle de Psychiatrie, Créteil, France, Institut National de la Santé et de la Recherche Médicale, U955, Créteil, France; **Beards, Stephanie**, PhD, Department of Health Service and Population Research, Institute of Psychiatry, King's College London, London, England; **Bonetto, Chiara**, PhD, Section of Psychiatry, Department of Neuroscience, Biomedicine and Movement, University of Verona, Verona, Italy; **Bonora, Elena**, PhD, Department of Medical and Surgical Science, Genetic Unit, Alma Mater Studiorum Università di Bologna, Bologna, Italy; **Cabrera, Bibiana**, MSc, PhD, Barcelona Clinic Schizophrenia Unit, Neuroscience Institute, Hospital Clinic of Barcelona, Barcelona, Spain, CIBERSAM, Spain; **Carracedo, Angel**, MD, PhD, Fundación Pública Galega de Medicina Xenómica, Hospital Clínico Universitario, CIBERER, Santiago de Compostela, Spain; **Charpeaud, Thomas**, MD, Fondation Fondamental, Créteil, France, CMP B CHU, Clermont Ferrand, France, and Université Clermont Auvergne, Clermont-Ferrand, France; **Costas, Javier**, PhD, Fundación Pública Galega de Medicina Xenómica, Hospital Clínico Universitario, Santiago de Compostela, Spain; **Cristofalo, Doriana**, MA, Section of Psychiatry, Department of Neuroscience, Biomedicine and Movement, University of Verona, Verona, Italy; **Cuadrado, Pedro**, MD, Villa de Vallecas Mental Health Department, Villa de Vallecas Mental Health Centre, Hospital Universitario Infanta Leonor/Hospital Virgen de la Torre, Madrid, Spain; **Durán-Cutilla, Manuel**, MD, Department of Child and Adolescent Psychiatry, Hospital General Universitario Gregorio Marañón, Institute of Psychiatry and Mental Health, IISGM, School of Medicine, Universidad Complutense, Madrid, Spain; **Ferchiou, Aziz**, MD, AP-HP, Groupe Hospitalier "Mondor", Pôle de Psychiatrie,

Créteil, France, Institut National de la Santé et de la Recherche Médicale, U955, Créteil, France; **Fraguas, David**, MD, PhD, Biomedical Research Networking Center for Mental Health Network (CIBERSAM), Barcelona, Spain; Department of Child and Adolescent Psychiatry, Hospital General Universitario Gregorio Marañón, Institute of Psychiatry and Mental Health, IiSGM, School of Medicine, Universidad Complutense, Madrid, Spain; **Franke, Nathalie**, MSc, Department of Psychiatry, Early Psychosis Section, Academic Medical Centre, University of Amsterdam, Amsterdam, the Netherlands; **Frijda, Flora**, MSc, Etablissement Public de Santé Maison Blanche, Paris, France; **Llorente, Cloe**, MD, Department of Psychiatry, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, Investigación Sanitaria del Hospital Gregorio Marañón (CIBERSAM), Madrid, Spain; **García-Portilla, Paz**, MD, PhD, Department of Medicine, Psychiatry Area, School of Medicine, Universidad de Oviedo, ISPA, INEUROPA, CIBERSAM, Oviedo, Spain; **González Peñas, Javier**, Hospital Gregorio Marañón, IiSGM, School of Medicine, Madrid, Spain; **Hubbard, Kathryn**, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London; **Jamain, Stéphane**, PhD, Institut National de la Santé et de la Recherche Médicale, U955, Créteil, France, Faculté de Médecine, Université Paris-Est, Créteil, France, and Fondation Fondamental, Créteil, France; **Jiménez-López, Estela**, MSc, Department of Psychiatry, Servicio de Psiquiatría Hospital "Virgen de la Luz," Cuenca, Spain; **Leboyer, Marion**, MD, PhD, AP-HP, Groupe Hospitalier "Mondor," Pôle de Psychiatrie, Créteil, France, Institut National de la Santé et de la Recherche Médicale, U955, Créteil, France, Faculté de Médecine, Université Paris-Est, Créteil, France, and Fondation Fondamental, Créteil, France; **López Montoya, Gonzalo**, PhD, Department of Child and Adolescent Psychiatry, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, IiSGM (CIBERSAM), Madrid, Spain; **Lorente-Rovira, Esther**, PhD, Department of Psychiatry, School of Medicine, Universidad de Valencia, CIBERSAM, Valencia, Spain; **M Díaz-Caneja, Covadonga**, Child and Adolescent Psychiatry Department, Hospital General Universitario Gregorio Marañón, Universidad Complutense, Madrid, Spain; **Marcelino Loureiro, Camila**, MD, Departamento de Neurociências e Ciências do Comportamento, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, São Paulo, Brasil, and Núcleo de Pesquisa em Saúde Mental Populacional, Universidade de São Paulo, São Paulo, Brasil; **Matteis, Mario**, MD, Department of Child and Adolescent Psychiatry, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, IiSGM (CIBERSAM), Madrid, Spain; **Messchaert, Elles**, MSc, Rivierduinen Centre for Mental Health, Leiden, the Netherlands; **Moltó, Ma Dolores**, Department of Genetics, University of Valencia, Campus of Burjassot, Biomedical Research Institute INCLIVA, Valencia, Spain, Centro de Investigación Biomedica en Red de Salud Mental (CIBERSAM), Madrid, Spain; **Mezquida, Gisela**, PhD, Barcelona Clinic Schizophrenia Unit, Neuroscience Institute, Hospital Clinic of Barcelona, Barcelona, Spain; Centre for Biomedical Research in the Mental Health Network (CIBERSAM), Spain; Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona; Department of Clinical Foundations, Pharmacology Unit, University of Barcelona, Barcelona, Spain; **Moreno, Carmen**, MD, Department of Psychiatry, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, IiSGM (CIBERSAM), Madrid, Spain; **Muratori, Roberto**, MD, Department of Mental Health and Pathological Addiction, Local Health Authority, Bologna, Italy; **Nacher Juan**, Neurobiology Unit, Department of Cell Biology, Interdisciplinary Research Structure for Biotechnology and Biomedicine (BIOTECMED), Universitat de València, Valencia, Spain, Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM): Spanish National Network for Research in Mental Health, Madrid, Spain, Fundación Investigación Hospital Clínico de Valencia, INCLIVA, Valencia, Spain; **Parellada, Mara**, MD, PhD, Department of Child and Adolescent Psychiatry, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, IiSGM (CIBERSAM), Madrid, Spain; **Pignon, Baptiste**, MD, AP-HP, Groupe Hospitalier "Mondor," Pôle de Psychiatrie, Créteil, France, Institut National de la Santé et de la Recherche Médicale, U955, Créteil, France, and Fondation Fondamental, Créteil, France; **Rapado-Castro, Marta**, PhD, Department of Child and Adolescent Psychiatry, Hospital General Universitario Gregorio Marañón,

Institute of Psychiatry and Mental Health, IiSGM, School of Medicine, Universidad Complutense, Madrid, Spain; **Ruggeri, Mirella**, MD, PhD, Section of Psychiatry, Department of Neuroscience, Biomedicine and Movement, University of Verona, Verona, Italy; **Richard, Jean-Romain**, MSc, Institut National de la Santé et de la Recherche Médicale, U955, Créteil, France, and Fondation Fondamental, Créteil, France; **Rodríguez Solano, José Juan**, MD, Puente de Vallecas Mental Health Department, Hospital Universitario Infanta Leonor/Hospital Virgen de la Torre, Centro de Salud Mental Puente de Vallecas, Madrid, Spain; **Sáiz, Pilar A**, Department of Psychiatry, School of Medicine, University of Oviedo, CIBERSAM. Instituto de Neurociencias del Principado de Asturias, INEUROPA, Oviedo, Spain; Servicio de Salud del Principado de Asturias (SESPA), Oviedo, Spain; **Sánchez-Gutierrez, Teresa**, Department of Child and Adolescent Psychiatry, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, IiSGM, CIBERSAM, C/Doctor Esquerdo 46, 28007 Madrid, Spain. Faculty of Health Science. Universidad Internacional de La Rioja (UNIR), Madrid, Spain; **Sánchez, Emilio**, MD, Department of Psychiatry, Hospital General Universitario Gregorio Marañón, School of Medicine, Universidad Complutense, IiSGM (CIBERSAM), Madrid, Spain; **Schürhoff, Franck**, MD, PhD, AP-HP, Groupe Hospitalier “Mondor,” Pôle de Psychiatrie, Créteil, France, Institut National de la Santé et de la Recherche Médicale, U955, Créteil, France, Faculté de Médecine, Université Paris-Est, Créteil, France, and Fondation Fondamental, Créteil, France; **Seri, Marco**, MD, Department of Medical and Surgical Science, Genetic Unit, Alma Mater Studiorum Università di Bologna, Bologna, Italy; **Shuhama, Rosana**, PhD, Departamento de Neurociências e Ciências do Comportamento, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, São Paulo, Brasil, and Núcleo de Pesquisa em Saúde Mental Populacional, Universidade de São Paulo, São Paulo, Brasil; **Stilo, Simona A**, PhD, Department of Health Service and Population Research, and Department of Psychosis Studies, Institute of Psychiatry, King’s College London, London, England; **Termorshuizen, Fabian**, PhD, Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, South Limburg Mental Health Research and Teaching Network, Maastricht University Medical Centre, Maastricht, the Netherlands, and Rivierduinen Centre for Mental Health, Leiden, the Netherlands; **Tronche, Anne-Marie**, MD, Fondation Fondamental, Créteil, France, CMP B CHU, Clermont Ferrand, France, and Université Clermont Auvergne, ClermontFerrand, France; **van Dam, Daniella**, PhD, Department of Psychiatry, Early Psychosis Section, Academic Medical Centre, University of Amsterdam, Amsterdam, the Netherlands; **van der Ven, Elsje**, PhD, Department of Psychiatry and Neuropsychology, School for Mental Health and Neuroscience, South Limburg Mental Health Research and Teaching Network, Maastricht University Medical Centre, Maastricht, the Netherlands, and Rivierduinen Centre for Mental Health, Leiden, the Netherlands.

***Corresponding author:**

Lucia Sideli, Department of Human Science, LUMSA University, Piazza delle Vaschette, 101 – 00193 Rome, Italy +39 06 684 221, l.sideli@lumsa.it

Childhood maltreatment, educational attainment and IQ: findings from a multicentric case-control study of first-episode psychosis (EU-GEI)

Abstract

Background and hypothesis: Evidence suggests that childhood maltreatment (i.e., childhood abuse and childhood neglect) affects educational attainment and cognition. However, the association between childhood maltreatment and Intelligence Quotient (IQ) seems stronger among controls compared to people with psychosis. We hypothesised that: the association between childhood maltreatment and poor cognition would be stronger among community controls than among people with first-episode of psychosis (FEP); compared to abuse, neglect would show stronger associations with educational attainment and cognition; the association between childhood maltreatment and IQ would be partially accounted for by other risk factors; and the association between childhood maltreatment, educational attainment and IQ would be stronger among patients with affective psychoses compared to those with non-affective psychoses.

Study Design: 829 patients with FEP and 1283 community controls from 16 EU-GEI sites were assessed for child maltreatment, education attainment, and IQ.

Study Results: In both the FEP and control group, childhood maltreatment was associated with lower educational attainment. The association between childhood maltreatment and lower IQ was robust to adjustment for confounders only among controls. Whereas childhood neglect was consistently associated with lower attainment and IQ in both groups, childhood abuse was associated with IQ only in controls. Among both patients with affective and non-affective psychoses, negative associations between childhood maltreatment and educational attainment were observed, but the crude association with IQ was only evident in affective psychoses.

Conclusions: Our findings underscore the role of childhood maltreatment in shaping academic outcomes and cognition of people with FEP as well as controls.

Keywords: IQ, psychosis, schizophrenia, childhood abuse, childhood neglect

Introduction

Accumulating evidence suggests that the burden of child maltreatment is not limited to the detrimental effect on mental health.¹ Childhood maltreatment can have long lasting effects on cognitive development and the capacity to achieve expected educational outcomes.^{2,3} Notably, childhood maltreatment can deviate the typical neurodevelopment of the individual,⁴ as it might produce multiple alterations in information processing and emotion regulation,^{5,6} and the underlying brain structures, circuits and processes.^{7,8} Furthermore, childhood maltreatment has been linked to long-term changes of the hypothalamic-pituitary-adrenal axis which may affect brain regions rich in glucocorticoids receptors, such as the hippocampus and the prefrontal cortex, contributing to cognitive impairment.⁸⁻¹⁰

The effect of childhood maltreatment on cognition may be part of the developmental pathway for psychosis, especially for schizophrenia, as the disorder has been consistently associated with lower intelligence quotient (IQ).^{11,12} Cognitive impairment is already present several years prior to the first episode of psychosis (FEP)^{13,14} and significantly affects community functioning.^{15,16} Cognitive impairment may also influence academic outcomes;¹⁷ yet, longitudinal studies on school performance have led to inconsistent findings, with a few studies reporting poor academic achievement predicting psychosis onset, but also some non-significant findings.^{18,19}

Meta-analytic findings suggest that children exposed to maltreatment show poorer cognitive performance than unexposed children, even in the absence of post-traumatic stress disorder.²⁰ More recently, a meta-analysis on adults with and without psychotic disorders found a modest negative correlation between childhood maltreatment and overall cognition. Subgroup analysis revealed that the association between early adversities and cognition was stronger amongst healthy controls than amongst people with psychosis, and it was suggested that the difference might be partially explained by concurrent risk factors affecting the cognitive development and cognitive performance of people with psychosis.³ These potential confounders include socio-economic disadvantage,^{21,22} poor premorbid adjustment,²³⁻²⁵ and psychotic experiences.^{26,27} Since all these factors have been associated with both childhood adversities and cognition, they might reduce the association between early adversities and cognition. Moreover, among people with psychosis, cannabis use has been related to a higher IQ,²⁸ suggesting that the association between childhood maltreatment and cognition might be weaker among cannabis users with FEP.

Another issue to account for when examining the association between childhood maltreatment and cognition in patients with FEP is the different impact on cognition exerted by specific types of childhood adversities. A recent literature review²⁹ found that early deprivation was strongly

associated with cognitive impairment among institutionalized children. However, study findings on non-institutionalized children were less robust, and only a few of them explored the differential effects of childhood abuse and neglect, with mixed findings.²⁹ Evidence regarding a specific effect of childhood adversities on adult cognitive impairment is also limited,³⁰ but preliminary findings indicate that academic failure may be more strongly related with neglect, institutionalization, and multiple maltreatment, compared to abuse.^{2,31} It was proposed that childhood neglect, in combination or not with childhood abuse, might be related with inadequate stimulation during critical periods of brain development, insecure attachment, emotion dysregulation, and impaired sense of agency, which in turn affect cognitive development and academic success.^{2,5}

Another important factor is the heterogeneity of psychosis syndromes.^{32,33} Accumulating evidence suggests that individuals who will develop non-affective psychoses have a premorbid IQ lower than controls, while evidence about affective psychoses is mixed.³⁴⁻³⁶ At the first onset of psychosis and in the long-term course of these disorders, cognitive impairments appear more severe amongst those with non-affective psychoses than in those with affective psychoses.^{26,37} This may suggest that childhood adversities are less relevant for understanding impaired cognitive functioning among patients with non-affective psychoses.³⁸

In light of such findings, the current study aimed to better understand the association between childhood maltreatment and educational attainment and cognitive functioning in a large multicentric sample of people with FEP and community controls. We hypothesised that: a) the association between childhood maltreatment and poor cognition would be stronger among community controls than among people with FEP; b) compared to childhood abuse, childhood neglect would show stronger associations with educational attainment and cognition; c) the association between childhood maltreatment and IQ would be partially accounted for by other risk factors potentially affecting cognitive functioning; and d) the association between childhood maltreatment, educational attainment and IQ would be stronger among patients with affective psychoses compared to those with non-affective psychoses.

Methods

Participants and procedure

Study participants were recruited from May 2010 to April 2015 within the EU-GEI study, a multi-centre case-control study involving 16 study centres across five European countries and Brazil. The Internal Review Boards of the study centres approved the study and participants provided written informed consent to be interviewed and let their data be stored and analysed anonymously.³⁹

Patients were recruited among incident cases of psychosis, aged 18-64 and resident in the study catchment areas, approaching mental health services for the first time during the study period for a diagnosis of psychotic disorder (ICD-10 diagnoses: F20–F33), neither secondary to acute intoxication (ICD-10: F1X.5) nor to medical condition (ICD-10: F09), and not previously treated with antipsychotics. Diagnoses of FEP were made according to ICD-10 criteria⁴⁰ on the basis of the Operational Criteria Checklist algorithm, OPCRIT⁴¹ administered by trained researchers (interrater reliability: $k=.7$).⁴² Clinical diagnoses were used only when OPCRIT assessment was not possible (12.1%). Diagnoses were combined to form a group of non-affective (ICD-10 codes F20-F29) and affective (ICD-10 codes F30-F33) psychoses.

Community controls were recruited among people aged 18-64, resident in the same catchment areas as patients, never referred or treated for psychotic disorders. Random and quota sampling (population stratification by age, sex, and ethnicity) were used to ensure representativeness of the same population as the patients.^{42,43} Controls were excluded if they had ever received a diagnosis or treatment for psychotic disorders.

Measures

Childhood maltreatment was assessed using the Childhood Trauma Questionnaire (CTQ),⁴⁴ a 28-item self-report tool assessing the frequency of five types of childhood adversity (physical, sexual, and emotional abuse, and physical and emotional neglect) on a 5-point Likert scale (from 1 [never] to 5 [very often]). Consistent with previous studies suggesting a differential effect of childhood abuse and neglect^{29,31} on education and cognition, an overall “childhood maltreatment” score, and separate “childhood abuse” and “childhood neglect” scores were calculated on the basis of the mean score of the respective items. A second-order confirmatory factor analysis (DWLS estimation) supported the two-factor structure of CTQ, comprising neglect and abuse factors (see Supplementary data 1). Although evidence suggests the relevance of using continuous measures of childhood maltreatment,⁴⁵ in this study childhood maltreatment was operationalized as a dichotomous variable, because assumptions of homoscedasticity for linear regression were not met and in order to highlight the presence of severe instances of childhood maltreatment. Therefore, three dichotomous variables for childhood maltreatment, abuse, and neglect were calculated using the 80th percentile of the control group as a cut-off value, according to the procedure used in a previous study.³⁸ The CTQ considered exposure to experiences of abuse and neglect prior to age 18.

Cognition was estimated from overall Intelligence Quotient (IQ) assessed using an abbreviated Wechsler Adult Intelligence Scale (WAIS-III).⁴⁶ The administration and scoring

procedure of the abbreviated version have been previously described and psychometrically validated.^{21,47}

Educational attainment was assessed using a modified version of the MRC sociodemographic questionnaire⁴⁸ and defined as the highest level of education fully completed, on a scale from 1 (no education) to 6 (postgraduate education).

To account for the confounding effect of concurrent and early conditions potentially affecting cognitive functioning, the following conditions were also assessed: (a) lifetime cannabis use was assessed using a modified version of the Cannabis Experience Questionnaire (CEQmv);⁴⁹ (b) lifetime psychotic experiences were assessed using the mean score of the Community Assessment of Psychic Experiences (CAPE);⁵⁰ (c) premorbid social adjustment in childhood and adolescence was assessed using the mean score of the Premorbid Adjustment Scale (PAS);⁵¹ (d) social disadvantage was estimated by proxy from the main family social class during upbringing, assessed on a four-level scale (from long-term unemployment to salariat), using the MRC sociodemographic questionnaire.⁴⁸

Analyses

Patients and controls were compared according to the prevalence of childhood maltreatment, educational attainment, and IQ using odds ratios (OR) and t-tests. The level of educational attainment and IQ were compared between patients and controls exposed and not exposed to maltreatment using t-tests. The associations between childhood maltreatment, abuse, and neglect (independent categorical variables, IVs) and IQ and educational attainment (dependent continuous variables, DVs) were assessed separately for patients and controls using general linear regression models (model 1). The crude association (model 1) was adjusted for: study country, sex, age, ethnicity (white vs. non-white), and education (only the child maltreatment-IQ association) or IQ (only the child maltreatment-education attainment association) (model 2). Also, analyses were additionally adjusted for lifetime cannabis use and lifetime psychotic experiences (model 3); premorbid social adjustment and family social disadvantage (model 4); and current use of antipsychotics (none vs. one vs. more than one) (model 5). All categorical confounders were included as fixed factors, except country which was included as a random factor. Given the number of predictors and the limited sample size (N<50 in 56% of the study sites), analyses were not controlled for study site which is consistent with previous studies on the same sample.^{21,28}

Assumptions of normality and homoscedasticity of IQ and educational attainment between groups (i.e., exposed vs. unexposed cases, and exposed vs. unexposed controls), and lack of notable multicollinearity among childhood maltreatment and covariates were verified (see Supplementary

data 2). Interactions between case-control status and childhood maltreatment, childhood abuse, and childhood neglect were assessed using generalized linear models. Subgroup analysis was carried out to investigate the specific associations between childhood maltreatment and education/IQ among FEP patients with affective and non-affective psychosis.

Associations between childhood maltreatment and education attainment or IQ were reported as regression coefficients (B) (see Tables 3 and 4). In order to estimate effect sizes, analyses were repeated using standardized IVs and DVs. Resulting β values .1-.3, .3-.5, and $>.5$ were considered to represent small, medium, and large effect sizes (ES), respectively. β values were compared across models in order to assess the strength of the associations between different types of maltreatment and education or IQ.⁵² Only study participants with complete measures of childhood maltreatment, cognition, and educational status were included in the analyses. Study participants with missing data in one or more of the confounders were included only in the crude analyses (see Tables 3 and 4). Analyses were run using the Statistical Package for the Social Sciences (SPSS) program version 27.0.

Results

Participants

Eight hundred and twenty-nine patients with FEP and 1283 community controls with complete measures of childhood maltreatment, cognition, and educational status (i.e., 73.4% and 85.7% of eligible FEP and controls, respectively) were included in the analyses. Those with incomplete information were more often of non-white ethnicity ($\chi^2(1)=13.05, p<.001$), less frequently graduated ($\chi^2(5)=23.62, p<.001$), and from a lower social class ($\chi^2(3)=8.71, p=.033$).

Compared to community controls, patients with FEP were more often males, younger, of non-white ethnicity, and from a lower social class (all $p's \leq .001$, see Supplementary Table 1). Compared to controls, patients were about three times as likely to have been exposed to childhood maltreatment (OR=3.39, 95%CI=2.78,4.12), abuse (OR=3.17, 95%CI=2.60,3.87), and neglect (OR=3.24, 95%CI=2.66,3.93). On average, the highest educational attainment of patients was one level below the highest attainment of controls ($t(2110)=14.12, p<.001$). Patients' average IQ was about 18 points lower than controls ($t(2110)=22.55, p<.001$) (Table 1).

Childhood maltreatment, educational attainment and IQ among community controls

Controls exposed to childhood maltreatment had lower education attainment compared to those who were unexposed (Table 2). In the unadjusted model, both childhood abuse and childhood neglect

(Table 3, Supplementary Table 3, model 1) were associated with lower educational attainment, with a small ES ($\beta=-.07$ and $\beta=-.12$, respectively), but in the fully adjusted model only neglect (model 4, $\beta=-.08$) contributed to lower academic attainment.

A 5-point mean difference was observed between the IQ of controls exposed to childhood maltreatment and the IQ of those unexposed (Table 2). The small associations between abuse and IQ, as well as between neglect and IQ (Table 4, Supplementary Table 4, model 1; $\beta=-.13$, $\beta=-.12$), were both attenuated in the fully adjusted model (model 4, $\beta=-.05$; $\beta=-.05$).

Childhood maltreatment, educational attainment and IQ among patients with FEP

Patients exposed to childhood maltreatment less frequently achieved higher academic qualifications (Table 2). The crude association between abuse and educational attainment (Table 3, Supplementary table 3, model 1; $\beta=-.09$) was no longer evident after controlling for psychotic experiences and cannabis use (model 3), whereas the small size association with neglect was still evident in the fully adjusted model (model 5, $\beta=-.11$).

A 3-point mean difference was observed between the IQ of FEP patients exposed to childhood maltreatment and the IQ of those unexposed (Table 2). In the unadjusted model, only neglect was weakly associated with lower IQ (Table 4, Supplementary Table 4, model 1, $\beta=-.07$), but the association was no longer evident after controlling for sociodemographic variables and education (model 2).

Despite the association between neglect and education was more robust to adjustment for confounders than the association between abuse and education, both in the control and the case group, the overlapping 95%CI suggested that there was no evidence of a stronger effect of one type of maltreatment over the other. The same was the case for the association between childhood abuse, childhood neglect and IQ. Furthermore, non-significant differences between β values suggested similar ES of the two types of maltreatment (all $ps>.05$).

When we formally tested whether the association between childhood maltreatment and education or IQ differed between cases and controls, we found no evidence to suggest that this was the case for childhood neglect or abuse and education, or neglect and IQ. We did observe a statistically significant interaction (Wald $\chi^2=11.06$, $p=.001$) between childhood abuse and case-control status on IQ, such that the association between abuse and IQ was evident in controls (Wald $\chi^2=4.33$, $p=.037$), but not in cases (Wald $\chi^2=0.46$, $p=.461$).

Potential confounders of the association between childhood maltreatment, educational attainment, and IQ

Only among controls, socio-demographic factors and IQ reduced the association between childhood abuse and educational attainment to non-significance (Table 3, model 2). Furthermore, controls who achieved lower qualifications, reported greater frequency of psychotic experiences, and more often belonged to the lower and the intermediate social classes (see Supplementary data 3).

In both groups, IQ scores were related to male sex, age, non-white ethnicity, education, and country (see Supplementary data 3). Furthermore, social disadvantage was associated with lower IQ and slightly attenuated the association with childhood maltreatment in the control group (Table 4, model 4). Specifically, both the lower and the intermediate social classes were associated with lower IQ compared to those of higher social class. Only among patients was lifetime cannabis use associated with higher IQ (see Supplementary data 3).

Subgroup analysis: childhood maltreatment, educational attainment and IQ among patients with affective and non-affective FEP

A similar percentage of patients with non-affective (n=575) and affective (n=240) FEP reported any form of childhood maltreatment (43.7% vs. 45.0%, OR=1.06, 95%CI=0.78,1.43), and this was also found for abuse (39.3% vs. 43.8%, OR=1.20, 95%CI=0.88,1.63) and neglect (44.0% vs. 44.2%, OR=1.01, 95%CI=0.74,1.36) when considered separately. No significant difference between the two groups was found for their mean educational attainment ($t(813)=0.64, p=.522$) and mean IQ ($t(825)=-1.81, p=.071$).

Patients with non-affective FEP exposed to childhood abuse or childhood neglect achieved lower educational levels than those unexposed (Supplementary Tables 5 and 7, model 1; $\beta=-.09$ and $\beta=-.08$, respectively). The association with abuse was robust to adjustment for sociodemographic and clinical factors, except antipsychotic treatment (model 5, $\beta=-.10$). Furthermore, in this group no association between childhood maltreatment and IQ was found (Supplementary Table 6 and 8, model 1).

Among patients with affective FEP, childhood neglect was weakly associated with lower educational attainment, after accounting for potential confounders (Supplementary Tables 5 and 7, model 5, $\beta=-.15$). In this group, neglect was associated with a 5-point difference in IQ in the crude model, with a small ES (Supplementary Table 6 and 8, model 1, $\beta=-.13$), but the association was reduced in the adjusted models (Supplementary Tables 6 and 8, model 2, $\beta=-.06$). However, the

overlapping 95% CI and the non-significant difference between β values suggested that the effect of childhood neglect was similar to the effect of abuse. Furthermore, the limited sample size did not allow us to formally test the influence of potential interactions between FEP diagnosis and childhood maltreatment, abuse, or neglect, on education and IQ.

Discussion

In summary, childhood abuse and childhood neglect were associated with poorer educational attainment in both people with FEP and community controls, both with a small ES. However, the association between childhood maltreatment and IQ was more robust to adjustment for confounders in community controls, as compared with FEP patients. Furthermore, an interaction between case status and abuse was found, such that the association between abuse and IQ was only evident among controls.

Associations between childhood maltreatment, educational attainment, and IQ varied according to the FEP clinical phenotype. In the non-affective psychosis group, childhood abuse and neglect were associated with poorer achievement, and no association between any type of childhood maltreatment and IQ was observed. In the affective psychosis group, only neglect was associated with lower educational attainment and, weakly, with lower IQ.

Associations between childhood maltreatment and education and IQ among FEP patients and community controls

Across both the clinical and community groups, childhood maltreatment, especially neglect, was associated with lower educational attainment, even when the effects of IQ and social disadvantage were taken into account. To our knowledge, only the GROUP study previously investigated the effect of childhood maltreatment on education among people with psychosis controlling for a proxy of social disadvantage different from that used in this study (i.e., parental educational level), with negative findings.⁵³ Inconsistency between the two studies may be due to differences in the study population (only non-affective psychoses in the GROUP study vs. both affective and non-affective psychoses in the current study), the definition of the outcome variable (inter-generational educational difference vs. participants' education level), or the effect of other variables (i.e., the study countries and the characteristics of the different school systems). Therefore, further replication studies are warranted.

This study builds on existing literature regarding a different effect of childhood maltreatment on IQ among patients with FEP and community controls without psychosis. Consistent with previous literature,^{3,13} the association between childhood maltreatment and IQ was much more robust in the control group than in the patient group. The findings suggest that the association between childhood maltreatment and IQ may be partially confounded by lower education, social disadvantage, and cannabis use, which are also associated with psychosis.^{21,42,54} This is consistent with a recent study utilising the Dunedin and E-Risk cohorts, which found that the association between childhood maltreatment and adult cognition was attenuated after controlling for early cognitive impairment and family disadvantage.⁵⁵ Contrary to our hypotheses, we did not observe a confounding effect of premorbid social functioning. This may depend on the effect of premorbid social functioning on current IQ being partially accounted by the effect of other factors included in the model, such as education.

Association between specific types of adversities and educational attainment and IQ

Exploring different types of maltreatment, this study found that childhood abuse and neglect were associated with lower educational attainment in the crude models, with a small ES. Furthermore, in both samples the association with neglect was more robust to adjustment for confounders.

Among community controls childhood abuse and childhood neglect had a similar negative association with IQ. Among patients with FEP, only neglect was associated with IQ. Furthermore, even controlling for confounders, the association between abuse and IQ was only evident among community controls, suggesting a possible interaction.

The specific effect of different types of maltreatment on education and IQ might have been attenuated by the difficulty in disentangling childhood abuse by childhood neglect, as well as by the possible relationship with other risk factors (e.g., parental loss, poor social support)⁵⁶. However, the more consistent pattern of association between childhood neglect, education, and IQ across samples is consistent with previous studies on both community^{2,31,57,58} and psychosis sample.^{59–63}

Relationship between childhood maltreatment, educational attainment, and IQ across diagnostic groups

Subgroup analyses showed that childhood abuse and childhood neglect were related to poor educational outcomes in patients with non-affective FEP with similar ES, whereas only neglect was associated with poor educational outcomes in the affective FEP patients. This suggests that different

clinical phenotypes within the psychosis spectrum might be more sensitive to the effect of specific types of adversities.^{64,65}

Furthermore, the association between childhood neglect and IQ was only evident among patients with affective psychoses. This is in line with preliminary findings from smaller samples^{59,66} and suggests a limited or null effect of childhood adversities on cognitive functions of people with non-affective psychotic disorders, which may be due to a pre-existing cognitive impairment affected by earlier biological risk factors not assessed here (e.g., obstetrical complications).^{33,67,68} For instance, evidence has suggested that preterm birth is associated with early attentional and executive impairment.⁶⁸ The lack of association between neglect and IQ among people with non-affective psychosis may also be influenced by a floor effect related to the lower IQ of people with non-affective psychoses in comparison to the IQ of those with affective psychoses.^{38,69} The lower sensitivity to social stressors by patients with non-affective psychoses would also be compatible with the hypothesis of an affective pathways to psychosis.^{32,70,71}

Strength and limitations

This study used a large multi-centre representative sample of patients with FEP and community controls to investigate associations between childhood maltreatment and adult academic attainment and cognitive functioning. However, the findings should be considered in light of several limitations. A key limitation of this study is the cross-sectional design which prevents any conclusions being drawn about the direction of the associations found. Additionally, EU-GEI study participants with complete information about education, IQ, and childhood maltreatment were more often of white ethnicity, highly educated, and belonging to a medium-high social class. The wide age range of study sample (i.e., 18-64) might have affected some participants' capacity to accurately recall childhood experiences particularly if they happened several decades ago. Also, retrospective measures of childhood abuse have shown poor agreement with prospective measures and may be affected by recall bias.⁷² Furthermore, since childhood and adolescent adversities might have a differential impact on IQ, future studies should account for the timing of childhood maltreatment, which was not available in this study. In this study, educational attainment was measured only with reference to quantitative aspects, not accounting for qualitative aspects. Furthermore, early and recent confounders were identified on the basis of the current literature and tested through multivariate model but other potential confounders not investigated here may include: a) genetic liability for psychotic disorders; b) developmental abnormalities (e.g., preterm birth); and c) psychiatric disorders other than psychosis, which might be potentially related to childhood maltreatment (e.g., depression).

Clinical implications

The findings of this study underscore the role of childhood abuse and childhood neglect in shaping the long-term academic outcomes and the cognitive functions of both patients with psychosis and unaffected controls. This suggests that adequate clinical attention should be given, in addition to severe forms of physical and sexual abuse, to less visible types of maltreatment, such as physical and emotional neglect,⁴ as they may similarly impair the cognitive and affective development of children. Children who are victims of maltreatment could be screened for cognitive impairment, and cognitive rehabilitation programs could be implemented as part of a comprehensive treatment package. Furthermore, considering literature suggesting a protective role of education and intact cognitive functions in the course and outcome of psychosis,^{73,74} and their relevance for later occupational, social, and economic outcomes,^{22,75,76} the results of this study emphasise the relevance of cognitive rehabilitation programs, school support, and vocational interventions for people with early psychosis.^{77,78}

Conclusions

This study found that, accounting for the effect of social class and IQ, childhood maltreatment was related to poorer academic outcomes among people with FEP and community controls. We also confirmed that among community controls childhood maltreatment was negatively related with adult IQ, and this association seemed relatively independent of confounders. The association with cognitive functioning was less evident among people with psychosis, particularly among those with non-affective psychoses.

FUNDING

The EU-GEI Study is funded by grant agreement HEALTH-F2-2010-241909 (Project EU-GEI) from the European Community's Seventh Framework Programme, and Grant 2012/0417-0 from the São Paulo Research Foundation. B.P.F. Rutten is funded by a VIDI award (no. 91.718.336) from the Netherlands Scientific Organization.

H. L. Fisher, C. Gayer-Anderson, and C. Morgan are supported by the Economic and Social Research Council (ESRC) Centre for Society and Mental Health at King's College London [ES/S012567/1].

C. Arango has received support by the Spanish Ministry of Science and Innovation. Instituto de Salud Carlos III (SAM16PE07CP1, PI16/02012, PI19/024), co-financed by ERDF Funds from the European Commission, “A way of making Europe”, CIBERSAM. Madrid Regional Government (B2017/BMD-3740 AGES-CM-2), European Union Structural Funds. European Union Seventh Framework Program under grant agreements, FP7- HEALTH-2013-2.2.1-2-603196 (Project PSYSCAN) and FP7- HEALTH-2013-2.2.1-2-602478 (Project METSY); and European Union H2020 Program under the Innovative Medicines Initiative 2 Joint Undertaking (grant agreement No 115916, Project PRISM, and grant agreement No 777394, Project AIMS-2-TRIALS), Fundación Familia Alonso and Fundación Alicia Koplowitz.

J.B. Kirkbride is supported by the NIHR University College London Hospital Biomedical Research Centre.

The views expressed are those of the authors and not necessarily those of the ESRC or King’s College London.

ACKNOWLEDGEMENTS

Dr. Arango has been a consultant to or has received honoraria or grants from Acadia, Angelini, Boehringer, Gedeon Richter, Janssen Cilag, Lundbeck, Minerva, Otsuka, Pfizer, Roche, Sage, Servier, Shire, Schering Plough, Sumitomo Dainippon Pharma, Sunovion and Takeda. Prof. Kirkbride has been a consultant to Roche and the Health Services Executive, Ireland.

The other authors have declared that there are no conflicts of interest in relation to the subject of this study.

References

1. Gilbert R, Widom CS, Browne K, Fergusson D, Webb E, Janson S. Burden and consequences of child maltreatment in high-income countries. *Lancet*. 2009;373(9657):68-81. doi:10.1016/S0140-6736(08)61706-7
2. Romano E, Babchishin L, Marquis R, Fréchette S. Childhood Maltreatment and Educational Outcomes. *Trauma, Violence, Abus*. 2015;16(4):418-437. doi:10.1177/1524838014537908
3. Vargas T, Lam PH, Azis M, Osborne KJ, Lieberman A, Mittal VA. Childhood trauma and neurocognition in adults with psychotic disorders: A systematic review and meta-analysis. *Schizophr Bull*. 2019;45(6):1195-1208. doi:10.1093/schbul/sby150
4. Bifulco A, Schimmenti A. Assessing child abuse: “We need to talk!” *Child Abus Negl*. 2019;98. doi:10.1016/j.chiabu.2019.104236
5. Ford JD. Treatment implications of altered affect regulation and information processing

following child maltreatment. *Psychiatr Ann.* 2005;35(5):410-419. doi:10.3928/00485713-20050501-07

6. Hedges DW, Woon FL. Early-life stress and cognitive outcome. *Psychopharmacology (Berl).* 2011;214(1):121-130. doi:10.1007/s00213-010-2090-6
7. Hart H, Rubia K. Neuroimaging of child abuse: A critical review. *Front Hum Neurosci.* 2012;19(6):52. doi:10.3389/fnhum.2012.00052
8. McCrory E, De Brito SA, Viding E. The impact of childhood maltreatment: a review of neurobiological and genetic factors. *Front Psychiatry.* 2011;2:48. doi:10.3389/fpsy.2011.00048
9. Tarullo AR, Gunnar MR. Child maltreatment and the developing HPA axis. *Horm Behav.* 2006;50(4):632-639.
10. Carpenter LL, Shattuck TT, Tyrka AR, Geraciotti TD, Price LH. Effect of childhood physical abuse on cortisol stress response. *Psychopharmacology (Berl).* 2011;214(1):367-375. doi:10.1007/s00213-010-2007-4 [doi]
11. Heinrichs RW, Zakzanis KK. Neurocognitive deficit in schizophrenia: A quantitative review of the evidence. *Neuropsychology.* 1998;12(3):426-445. doi:10.1037/0894-4105.12.3.426
12. Reichenberg A, Harvey PD. Neuropsychological impairments in schizophrenia: Integration of performance-based and brain imaging findings. *Psychol Bull.* 2007;133(5):833-858. doi:10.1037/0033-2909.133.5.833
13. Aas M, Dazzan P, Mondelli V, Melle I, Murray RM, Pariante CM. A systematic review of cognitive function in first-episode psychosis, including a discussion on childhood trauma, stress, and inflammation. *Front Psychiatry.* 2014;4(JAN). doi:10.3389/fpsy.2013.00182
14. Mesholam-Gately RI, Giuliano AJ, Goff KP, Faraone S V., Seidman LJ. Neurocognition in First-Episode Schizophrenia: A meta-analytic review. *Neuropsychology.* 2009;23(3):315-336. doi:10.1037/a0014708
15. Fett AKJ, Viechtbauer W, Dominguez M de G, Penn DL, van Os J, Krabbendam L. The relationship between neurocognition and social cognition with functional outcomes in schizophrenia: A meta-analysis. *Neurosci Biobehav Rev.* 2011;35(3):573-588. doi:10.1016/j.neubiorev.2010.07.001
16. Rodriguez V, Aas M, Vorontsova N, et al. Exploring the interplay between adversity, neurocognition, social cognition, and functional outcome in people with psychosis: A narrative review. *Front Psychiatry.* 2021;12. doi:10.3389/fpsy.2021.596949
17. Kendler KS, Ohlsson H, Mezuk B, Sundquist K, Sundquist J. A Swedish national prospective and co-relative study of school achievement at age 16, and risk for schizophrenia, other nonaffective psychosis, and bipolar illness. *Schizophr Bull.* 2016;42(1):77-86. doi:10.1093/schbul/sbv103
18. Laurens KR, Luo L, Matheson SL, et al. Common or distinct pathways to psychosis? A systematic review of evidence from prospective studies for developmental risk factors and antecedents of the schizophrenia spectrum disorders and affective psychoses. *BMC Psychiatry.* 2015;15(1). doi:10.1186/s12888-015-0562-2
19. Welham J, Isohanni M, Jones P, McGrath J. The antecedents of schizophrenia: a review of birth cohort studies. *Schizophr Bull.* 2009;35(3):603-623. doi:10.1093/schbul/sbn084
20. Malarbi S, Abu-Rayya HM, Muscara F, Stargatt R. Neuropsychological functioning of childhood trauma and post-traumatic stress disorder: A meta-analysis. *Neurosci Biobehav*

Rev. 2017;72:68-86. doi:10.1016/j.neubiorev.2016.11.004

21. Velthorst E, Mollon J, Murray RM, et al. Cognitive functioning throughout adulthood and illness stages in individuals with psychotic disorders and their unaffected siblings. *Mol Psychiatry*. Published online 2021. doi:10.1038/s41380-020-00969-z
22. Lewis GG, Dykxhoorn J, Karlsson H, et al. Assessment of the role of IQ in associations between population density and deprivation and nonaffective psychosis. *JAMA Psychiatry*. 2020;77(7):729-736. doi:10.1001/jamapsychiatry.2020.0103
23. Cannon M, Jones P, Gilvarry C, et al. Premorbid social functioning in schizophrenia and bipolar disorder: Similarities and differences. *Am J Psychiatry*. 1997;154(11):1544-1550. doi:10.1176/ajp.154.11.1544
24. Velthorst E, Fett AKJ, Reichenberg A, et al. The 20-year longitudinal trajectories of social functioning in individuals with psychotic disorders. *Am J Psychiatry*. 2017;174(11):1075-1085. doi:10.1176/appi.ajp.2016.15111419
25. Alameda L, Ferrari C, Baumann PS, Gholam-Rezaee M, Do KQ, Conus P. Childhood sexual and physical abuse: Age at exposure modulates impact on functional outcome in early psychosis patients. *Psychol Med*. 2015;45(13):2727-2736. doi:10.1017/S0033291715000690
26. Sheffield JM, Karcher NR, Barch DM. Cognitive deficits in psychotic disorders: A lifespan perspective. *Neuropsychol Rev*. 2018;28(4):509-533. doi:10.1007/s11065-018-9388-2
27. Varese F, Smeets F, Drukker M, et al. Childhood adversities increase the risk of psychosis: A meta-analysis of patient-control, prospective-and cross-sectional cohort studies. *Schizophr Bull*. 2012;38(4):661-671. doi:10.1093/schbul/sbs050
28. Ferraro L, La Cascia C, Quattrone D, et al. Premorbid adjustment and IQ in patients with first-episode psychosis: a multisite case-control study of their relationship with cannabis use. *Schizophr Bull*. 2020;46(3):517-529. doi:10.1093/schbul/sbz077
29. Young-Southward G, Eaton C, O'Connor R, Minnis H. Investigating the causal relationship between maltreatment and cognition in children: A systematic review. *Child Abus Negl*. 2020;107. doi:10.1016/j.chiabu.2020.104603
30. Irigaray TQ, Pachecob JB, Grassi-Oliveiraa R, Fonseca RP, de Carvalho Leite JC, Kristensena CH. Child maltreatment and later cognitive functioning: A systematic review. *Psicol Reflex e Crit*. 2013;26(2):376-387. doi:10.1590/S0102-79722013000200018
31. Perfect MM, Turley MR, Carlson JS, Yohanna J, Saint Gilles MP. School-related outcomes of traumatic event exposure and traumatic stress symptoms in students: A systematic review of research from 1990 to 2015. *School Ment Health*. 2016;8(1):7-43. doi:10.1007/s12310-016-9175-2
32. Myin-Germeys I, van Os J. Stress-reactivity in psychosis: evidence for an affective pathway to psychosis. *Clin Psychol Rev*. 2007;27(4):409-424. doi:10.1016/j.cpr.2006.09.005
33. Murray RM, Sham P, Van Os J, Zanelli J, Cannon M, McDonald C. A developmental model for similarities and dissimilarities between schizophrenia and bipolar disorder. *Schizophr Res*. 2004;71(2-3):405-416. doi:10.1016/j.schres.2004.03.002
34. Arango C, Fraguas D, Parellada M. Differential neurodevelopmental trajectories in patients with early-onset bipolar and schizophrenia disorders. *Schizophr Bull*. 2014;40(SUPPL. 2). doi:10.1093/schbul/sbt198
35. Isohanni M, Isohanni I, Koponen H, et al. Developmental precursors of psychosis. *Curr Psychiatry Rep*. 2004;6(3):168-175. doi:10.1007/s11920-004-0061-5

36. Mollon J, Reichenberg A. Cognitive development prior to onset of psychosis. *Psychol Med*. 2018;48(3):392-403. doi:10.1017/S0033291717001970
37. Trotta A, Murray RM, Maccabe JH. Do premorbid and post-onset cognitive functioning differ between schizophrenia and bipolar disorder? A systematic review and meta-Analysis. *Psychol Med*. 2015;45(2):381-394. doi:10.1017/S0033291714001512
38. van Os J, Marsman A, van Dam D, et al. Evidence that the impact of childhood trauma on IQ is substantial in controls, moderate in siblings, and absent in patients with psychotic disorder. *Schizophr Bull*. 2017;43(2):316-324. doi:10.1093/schbul/sbw177
39. Van Os J, Rutten BPBP, Myin-Germeys I, et al. Identifying gene-environment interactions in schizophrenia: Contemporary challenges for integrated, large-scale investigations. *Schizophr Bull*. 2014;40(4):729-736. doi:10.1093/schbul/sbu069
40. World Health Organization. *The ICD-10 Classification of Mental and Behavioural Disorders: Clinical Descriptions and Diagnostic Guidelines*. World Health Organization; 1992.
41. McGuffin P, Farmer A, Harvey I. A polydiagnostic application of operational criteria in studies of psychotic illness: Development and reliability of the OPCRIT system. *Arch Gen Psychiatry*. 1991;48(8):764-770. doi:10.1001/archpsyc.1991.01810320088015
42. Jongsma HE, Gayer-Anderson C, Lasalvia A, et al. Treated incidence of psychotic disorders in the multinational EU-GEI study. *JAMA Psychiatry*. 2018;75(1). doi:10.1001/jamapsychiatry.2017.3554
43. Gayer-Anderson C, Jongsma HE, Di Forti M, et al. The European network of national schizophrenia networks studying Gene–Environment Interactions (EU-GEI): Incidence and First-Episode case–control programme. *Soc Psychiatry Psychiatr Epidemiol*. Published online January 23, 2020. doi:10.1007/s00127-020-01831-x
44. Bernstein DP, Ahluvalia T, Pogge D, Handelsman L. Validity of the childhood trauma questionnaire in an adolescent psychiatric population. *J Am Acad Child Adolesc Psychiatry*. 1997;36(3):340-348. doi:10.1097/00004583-199703000-00012
45. Cohodes EM, Kitt ER, Baskin-Sommers A, Gee DG. Influences of early-life stress on frontolimbic circuitry: Harnessing a dimensional approach to elucidate the effects of heterogeneity in stress exposure. *Dev Psychobiol*. 2021;63(2):153-172. doi:10.1002/dev.21969
46. Blyler CR, Gold JM, Iannone VN, Buchanan RW. Short form of the WAIS-III for use with patients with schizophrenia. *Schizophr Res*. 2000;46(2-3):209-215. doi:10.1016/S0920-9964(00)00017-7
47. Velthorst E, Levine SZ, Henquet C, et al. To cut a short test even shorter: Reliability and validity of a brief assessment of intellectual ability in Schizophrenia - A control-case family study. *Cogn Neuropsychiatry*. 2013;18(6):574-593. doi:10.1080/13546805.2012.731390
48. Mallet R. *Sociodemographic Schedule*. Section of Social Psychiatry, Institute of Psychiatry; 1997.
49. Di Forti M, Sallis H, Allegri F, et al. Daily use, especially of high-potency cannabis, drives the earlier onset of psychosis in cannabis users. *Schizophr Bull*. 2014;40(6):1509-1517. doi:10.1093/schbul/sbt181
50. Konings M, Bak M, Hanssen M, Van Os J, Krabbendam L. Validity and reliability of the CAPE: A self-report instrument for the measurement of psychotic experiences in the general

population. *Acta Psychiatr Scand*. 2006;114(1):55-61. doi:10.1111/j.1600-0447.2005.00741.x

51. Cannon-Spoor HE, Potkin SG, Jed Wyatt R. Measurement of premorbid adjustment in chronic schizophrenia. *Schizophr Bull*. 1982;8(3):470-480. doi:10.1093/schbul/8.3.470
52. Clogg CC, Petkova E, Haritou A. Statistical methods for comparing regression coefficients between models. *Am J Sociol*. 1995;100(5):1261-1293. doi:10.1086/230638
53. Frissen A, Lieveerse R, Marcelis M, Drukker M, Delespaul P. Psychotic disorder and educational achievement: a family-based analysis. *Soc Psychiatry Psychiatr Epidemiol*. 2015;50(10):1511-1518. doi:10.1007/s00127-015-1082-6
54. Di Forti M, Quattrone D, Freeman TP, et al. The contribution of cannabis use to variation in the incidence of psychotic disorder across Europe (EU-GEI): a multicentre case-control study. *The Lancet Psychiatry*. 2019;6(5). doi:10.1016/S2215-0366(19)30048-3
55. Danese A, Moffitt TE, Arseneault L, et al. The origins of cognitive deficits in victimized children: Implications for neuroscientists and clinicians. *Am J Psychiatry*. 2017;174(4):349-361. doi:10.1176/appi.ajp.2016.16030333
56. Schimmenti A. The trauma factor: Examining the relationships among different types of trauma, dissociation, and psychopathology. *J Trauma Dissociation*. 2018;19(5):552-571. doi:10.1080/15299732.2017.1402400
57. Eckenrode J, Laird M, Doris J. School performance and disciplinary problems among abused and neglected children. *Dev Psychol*. 1993;29(1):53-62. doi:10.1037/0012-1649.29.1.53
58. Geoffroy MC, Pinto Pereira S, Li L, Power C. Child neglect and maltreatment and childhood-to-adulthood cognition and mental health in a prospective birth cohort. *J Am Acad Child Adolesc Psychiatry*. 2016;55(1):33-40.e3. doi:10.1016/j.jaac.2015.10.012
59. Aas M, Steen NE, Agartz I, et al. Is cognitive impairment following early life stress in severe mental disorders based on specific or general cognitive functioning? *Psychiatry Res*. 2012;198(3):495-500. doi:10.1016/j.psychres.2011.12.045
60. Garcia M, Montalvo I, Creus M, et al. Sex differences in the effect of childhood trauma on the clinical expression of early psychosis. *Compr Psychiatry*. 2016;68:86-96. doi:10.1016/j.comppsy.2016.04.004
61. Kilian S, Asmal L, Chiliza B, et al. Childhood adversity and cognitive function in schizophrenia spectrum disorders and healthy controls: Evidence for an association between neglect and social cognition. *Psychol Med*. 2018;48(13):2186-2193. doi:10.1017/S0033291717003671
62. Li X Bin, Bo QJ, Zhang GP, et al. Effect of childhood trauma on cognitive functions in a sample of Chinese patients with schizophrenia. *Compr Psychiatry*. 2017;76:147-152. doi:10.1016/j.comppsy.2017.04.010
63. Mørkved N, Johnsen E, Kroken RA, et al. Does childhood trauma influence cognitive functioning in schizophrenia? The association of childhood trauma and cognition in schizophrenia spectrum disorders. *Schizophr Res Cogn*. 2020;21. doi:10.1016/j.scog.2020.100179
64. Alameda L, Christy A, Rodriguez V, et al. Association between specific childhood adversities and symptom dimensions in people with psychosis: Systematic review and meta-analysis. *Schizophr Bull*. 2021;47(4):975-985. doi:10.1093/schbul/sbaa199
65. Bentall RP, de Sousa P, Varese F, et al. From adversity to psychosis: pathways and

mechanisms from specific adversities to specific symptoms. *Soc Psychiatry Psychiatr Epidemiol.* 2014;49(7):1011-1022. doi:10.1007/s00127-014-0914-0

66. Aas M, Dazzan P, Fisher HL, et al. Childhood trauma and cognitive function in first-episode affective and non-affective psychosis. *Schizophr Res.* 2011;129(1):12-19. doi:10.1016/j.schres.2011.03.017
67. Richards AL, Pardiñas AF, Frizzati A, et al. The relationship between polygenic risk scores and cognition in Schizophrenia. *Schizophr Bull.* 2020;46(2):336-344. doi:10.1093/schbul/sbz061
68. Vanes LD, Murray RM, Nosarti C. Adult outcome of preterm birth: Implications for neurodevelopmental theories of psychosis. *Schizophr Res.* Published online 2021. doi:10.1016/j.schres.2021.04.007
69. Sideli L, Fisher HL, Russo M, et al. Failure to find association between childhood abuse and cognition in first-episode psychosis patients. *Eur Psychiatry.* 2014;29(1):32-35. doi:10.1016/j.eurpsy.2013.02.006
70. Sideli L, Murray RM, Schimmenti A, et al. Childhood adversity and psychosis: A systematic review of bio-psycho-social mediators and moderators. *Psychol Med.* 2020;50(11):1761-1782. doi:10.1017/S0033291720002172
71. Alameda L, Rodriguez V, Carr E, et al. A systematic review on mediators between adversity and psychosis: Potential targets for treatment. *Psychol Med.* 2020;50(12):1966-1976. doi:10.1017/S0033291720002421
72. Baldwin JR, Reuben A, Newbury JB, Danese A. Agreement between prospective and retrospective measures of childhood maltreatment: A systematic review and meta-analysis. *JAMA Psychiatry.* 2019;76(6):584-593. doi:10.1001/jamapsychiatry.2019.0097
73. Leeson VC, Sharma P, Harrison M, Ron MA, Barnes TRE, Joyce EM. IQ trajectory, cognitive reserve, and clinical outcome following a first episode of psychosis: A 3-year longitudinal study. *Schizophr Bull.* 2011;37(4):768-777. doi:10.1093/schbul/sbp143
74. Ziermans T, De Wit S, Schothorst P, et al. Neurocognitive and clinical predictors of long-term outcome in adolescents at ultra-high risk for psychosis: A 6-year follow-up. *PLoS One.* 2014;9(4). doi:10.1371/journal.pone.0093994
75. Morgan C, Fearon P, Lappin J, et al. Ethnicity and long-term course and outcome of psychotic disorders in a UK sample: The AEsOP-10 study. *Br J Psychiatry.* 2017;211(2):88-94. doi:10.1192/bjp.bp.116.193342
76. Shinn AK, Cawkwell PB, Bolton K, et al. Return to College After a First Episode of Psychosis. *Schizophr Bull Open.* 2020;1(1). doi:10.1093/schizbullopen/sgaa041
77. Arango C, Díaz-Caneja CM, McGorry PD, et al. Preventive strategies for mental health. *The Lancet Psychiatry.* 2018;5(7):591-604. doi:10.1016/S2215-0366(18)30057-9
78. Fusar-Poli P, Salazar De Pablo G, Correll CU, et al. Prevention of psychosis: advances in detection, prognosis, and intervention. *JAMA Psychiatry.* 2020;77(7):755-765. doi:10.1001/jamapsychiatry.2019.4779

Table 1: Childhood maltreatment, IQ and educational attainment of included FEP patients and controls

Variable	Total N=2112	Patients N=829	Controls N=1283	<i>t</i> / χ^2 (df)	<i>p</i>	OR (95% CI)	<i>p</i>
CTQ Mean score, range 1-4							
- Childhood Maltreatment, <i>M</i> (<i>SD</i>)	1.50 (0.51)	1.67 (0.57)	1.37 (0.43)	-14.27 (2110)	<.001	0.84 (0.82; 0.86)	<.001
- Abuse, <i>M</i> (<i>SD</i>)	1.35 (0.51)	1.50 (0.60)	1.26 (0.42)	-11.19 (2110)	<.001	0.87 (0.82; 0.92)	<.001
- Neglect, <i>M</i> (<i>SD</i>)	1.71 (0.68)	1.96 (0.74)	1.55 (0.58)	-14.02 (2110)	<.001	0.85 (0.80; 0.90)	<.001
Maltreatment exposure *							
- Childhood Maltreatment, <i>n</i> (%)	605 (28.6)	364 (43.9)	241 (18.8)	155.52 (1)	<.001	3.39 (2.78; 4.12)	<.001
- Abuse, <i>n</i> (%)	565 (26.8)	337 (40.7)	228 (17.8)	134.55 (1)	<.001	3.17 (2.60; 3.87)	<.001
- Neglect, <i>n</i> (%)	608 (28.8)	361 (43.5)	247 (19.3)	144.99 (1)	<.001	3.24 (2.66; 3.93)	<.001
IQ (N)							
- Full score, <i>M</i> (<i>SD</i>)	95.93 (19.88)	85.04 (18.18)	102.97 (17.63)	22.55 (2110)	<.001	0.99 (0.99; 0.99)	<.001
- Digit symbol, <i>M</i> (<i>SD</i>)	8.98 (3.46)	6.72 (2.91)	10.45 (2.96)	28.46 (2110)	<.001	0.92 (0.91; 0.93)	<.001
- Arithmetic, <i>M</i> (<i>SD</i>)	9.32 (3.61)	7.89 (3.45)	10.25 (3.40)	15.51 (2110)	<.001	0.94 (0.93; 0.94)	<.001
- Block design, <i>M</i> (<i>SD</i>)	9.19 (3.76)	7.70 (3.54)	10.15 (3.58)	15.43 (2110)	<.001	0.93 (0.93; 0.94)	<.001
- Information, <i>M</i> (<i>SD</i>)	9.96 (3.82)	8.78 (3.80)	10.72 (3.64)	11.73 (2110)	<.001	0.95 (0.94; 0.95)	<.001
Education							
- No qualification, <i>n</i> (%)	189 (8.9)	131 (15.8)	58 (4.5)	206.13 (5)	<.001	1	
- Compulsory, <i>n</i> (%)	387 (18.3)	216 (26.1)	171 (13.3)			1.26 (1.03; 1.54)	.022
- Tertiary, <i>n</i> (%)	542 (25.7)	199 (23.9)	344 (26.8)			0.58 (0.48; 0.69)	<.001
- Job related, <i>n</i> (%)	359 (17.0)	148 (17.9)	211 (16.4)			0.70 (0.57; 0.87)	.001
- University, <i>n</i> (%)	405 (19.2)	97 (11.7)	308 (24.0)			0.31 (0.25; 0.40)	<.001
- Post-degree, <i>n</i> (%)	230 (10.9)	39 (4.7)	191 (14.9)			0.20 (0.14; 0.28)	<.001
Mean education, range 1-6							
- <i>M</i> (<i>SD</i>)	3.52 (1.48)	2.98 (1.40)	3.87 (1.42)	14.12 (2110)	<.001	0.84 (0.82; 0.86)	<.001

CI: confidence intervals; CTQ: Childhood Trauma Questionnaire; df: degrees of freedom; FEP: first-episode psychosis; IQ: intelligence quotient; M: Mean; OR: odds ratio; SD: Standard Deviation. *defined as mean CTQ >80th percentile of the control group.

Table 2: IQ and educational attainment across group as a function of childhood maltreatment exposure*

Maltreatment exposure	Unexposed <i>M (SD)</i>	Exposed <i>M (SD)</i>	<i>t (df)</i>	<i>p</i>
IQ				
Controls (N=1283)				
Childhood maltreatment (1042 vs. 241)	103.96 (17.44)	98.69 (17.86)	4.21 (1281)	<.001
Abuse (1055 vs. 228)	104.05 (17.65)	98.01 (16.72)	4.73 (1281)	<.001
Neglect (1036 vs. 247)	103.97 (17.29)	98.80 (18.49)	4.17 (1281)	<.001
FEP Patients (N=829)				
Childhood maltreatment (465 vs. 364)	86.34 (18.47)	83.37 (17.69)	2.35 (827)	.019
Abuse (492 vs. 337)	85.33 (18.32)	84.61 (17.99)	0.56 (827)	.578
Neglect (468 vs. 361)	86.21 (18.54)	83.52 (17.62)	2.11 (827)	.035
EDUCATIONAL ATTAINMENT				
Controls (N=1283)				
Childhood maltreatment (1042 vs. 241)	3.93 (1.42)	3.59 (1.42)	3.38 (1281)	.001
Abuse (1055 vs. 228)	3.92 (1.42)	3.64 (1.45)	2.66 (1281)	.008
Neglect (1036 vs. 247)	3.95 (1.41)	3.52 (1.42)	4.32 (1281)	<.001
FEP Patients (N=829)				
Childhood maltreatment (465 vs. 364)	3.12 (1.41)	2.80 (1.37)	3.25 (827)	.001
Abuse (492 vs. 337)	3.08 (1.38)	2.82 (1.42)	2.60 (827)	.010
Neglect (468 vs. 361)	3.13 (1.45)	2.78 (1.35)	3.61 (827)	<.001

df: degrees of freedom; FEP: first-episode psychosis; IQ: intelligence quotient; M: Mean; SD: Standard Deviation; *defined as mean Childhood Trauma Questionnaire score >80th percentile of the control group.

Table 3: Associations between childhood maltreatment and educational attainment

Childhood maltreatment exposure	Model 1			Model 2			Model 3			Model 4			Model 5		
	<i>B</i>	95% CI	<i>p</i>	<i>B</i> ^a	95% CI	<i>p</i>	<i>B</i> _{a+b}	95% CI	<i>p</i>	<i>B</i> _{a+b+c}	95% CI	<i>p</i>	<i>B</i> _{a+b+c+d}	95% CI	<i>p</i>
Controls	<i>N</i> =1283			<i>N</i> =1280			<i>N</i> =1268			<i>N</i> =1145			<i>N</i> =1145		
Maltreatment	-0.34	-0.54; -0.14	.001	-0.22	-0.40; -0.04	.019	-0.18	-0.37; 0.01	.058	-0.19	-0.39; 0.00	.055			
Abuse	-0.28	-0.48; -0.07	.008	-0.11	-0.29; 0.08	.263	-0.05	-0.25; 0.14	.603	-0.10	-0.30; 0.11	.354			
Neglect	-0.43	-0.63; -0.24	<.001	-0.29	-0.47; -0.11	.002	-0.27	-0.45; -0.08	.004	-0.26	-0.45; -0.07	.006			
FEP Patients	<i>N</i> =829			<i>N</i> =829			<i>N</i> =695			<i>N</i> =599			<i>N</i> =561		
Maltreatment	-0.32	-0.51; -0.13	.001	-0.26	-0.44; -0.09	.003	-0.28	-0.47; -0.08	.006	-0.22	-0.44; -0.01	.041	-0.23	-0.45; -0.01	.040
Abuse	-0.26	-0.45; -0.06	.010	-0.22	-0.40; -0.05	.012	-0.16	-0.35; 0.04	.127	-0.17	-0.38; 0.05	.128	-0.14	-0.36; 0.08	.208
Neglect	-0.35	-0.54; -0.16	<.001	-0.28	-0.46; -0.11	.001	-0.31	-0.12; -0.26	.001	-0.29	-0.50; -0.08	.006	-0.30	-0.51; -0.08	.006

CI: confidence intervals; FEP: first-episode psychosis. ^a adjusted for sex, age, ethnicity, intelligence quotient (IQ), and study country; ^b adjusted for psychotic experiences and lifetime cannabis use; ^c adjusted for social disadvantage and premorbid social functioning; ^d adjusted for antipsychotic treatment; significant associations (*p* < .05) are shown in bold type.

Table 4: Association between childhood maltreatment and IQ

Childhood maltreatment exposure	Model 1			Model 2			Model 3			Model 4			Model 5		
	<i>B</i>	95% CI	<i>p</i>	<i>B</i> ^a	95% CI	<i>p</i>	<i>B</i> _{a+b}	95% CI	<i>p</i>	<i>B</i> _{a+b+c}	95% CI	<i>p</i>	<i>B</i> _{a+b+c+d}	95% CI	<i>p</i>
Controls	<i>N</i> =1283			<i>N</i> =1280			<i>N</i> =1268			<i>N</i> =1145			<i>N</i> =1145		
Maltreatment	-5.28	-7.73; -2.82	<.001	-2.35	-4.50; -0.19	.033	-2.62	-4.85; -0.39	.022	-2.13	-4.48; 0.21	.075			
Abuse	-6.05	-8.55; -3.54	<.001	-2.93	-5.12; -0.75	.009	-3.23	-5.52; -0.93	.006	-2.54	-4.99; -0.09	.042			
Neglect	-5.17	-7.61; -2.74	<.001	-2.95	-5.09; -0.81	.007	-3.05	-5.22; -0.89	.006	-2.36	-4.64; -0.09	.042			
FEP Patients	<i>N</i> =829			<i>N</i> =829			<i>N</i> =695			<i>N</i> =599			<i>N</i> =561		
Maltreatment	-2.98	-5.47; -0.49	.019	-1.02	-3.22; 1.19	.368	-1.43	-3.91; 1.06	.261	-0.59	-3.25; 2.08	.666	-0.72	-3.52; 2.06	.610
Abuse	-0.72	-3.24; 1.81	.578	1.07	-1.16; 3.29	.347	0.35	-2.17; 2.86	.787	0.98	-1.68; 3.64	.470	1.10	-1.67; 3.87	.436
Neglect	-2.68	-5.18; -0.19	.035	-0.96	-3.16; 1.24	.391	-0.79	-3.21; 1.63	.522	-0.45	-3.02; 2.13	.734	-0.65	-3.34; 2.04	.635

CI: confidence intervals; FEP: first-episode psychosis; IQ: intelligence quotient. ^a adjusted for sex, age, ethnicity, education, and study country; ^b adjusted for psychotic experiences and lifetime cannabis use; ^c adjusted for social disadvantage and premorbid social functioning; ^d adjusted for antipsychotic treatment; significant associations (*p* < .05) are shown in bold type.