

Near-infrared fluorescence image-guided surgery in esophageal and gastric cancer operations

Authors:

Ekman Marcin(1), Girnyi Sergii(1), Marano Luigi(2), Roviello Franco(2), Chand Manish(3), Diana Michele(4), Polom Karol(1)

1. Department of Surgical Oncology, Medical University of Gdansk, Gdansk, Poland
2. Department of Medicine, Surgery and Neurosciences, Unit of General Surgery and Surgical Oncology, University of Siena, Siena, Italy
3. Wellcome EPSRC Centre for Interventional and Surgical Sciences (WEISS), University College London, London
4. Department of General, Digestive, and Endocrine Surgery, University Hospital of Strasbourg, Strasbourg, France

Abstract

Near-infrared fluorescence image-guided surgery helps surgeons to see beyond the classical eye vision. Over the last few years, we have witnessed a revolution which has begun in the field of image-guided surgery. Fluorescence technology using indocyanine green (ICG) has shown promising results in many organs, and in this review article, we wanted to discuss the 6 main domains where fluorescence image-guided surgery is currently used for esophageal and gastric cancer surgery. Visualization of lymphatic vessels, tumor localization, fluorescence angiography for anastomotic evaluation, thoracic duct visualization, tracheal blood flow analysis, and sentinel node biopsy are discussed. It seems that this technology has already found its place in surgery. However, new possibilities and research avenues in this area will probably make it even more important in the near future.

Keywords: Anastomotic leakage; lymphography; indocyanine green fluorescence angiography; sentinel node.

Introduction

Over the last few years, near-infrared (NIR) fluorescence technology using indocyanine green (ICG) has shown promising results in image-guided surgery for many organs. Initially proposed by Kitai *et al.* in NIR fluorescence (NIRF) sentinel node biopsy (SNB) using ICG as a fluorophore, NIRF is currently used in various applications(1) worldwide. ICG is a water-soluble anionic amphiphilic tricarboyanine probe(2). Its small 1.2nm diameter is accountable for the fast migration of this fluorophore in the lymphatic system. With the use of an NIR charge-coupled device, the excitation of this fluorophore is 778nm, and its emission is 830nm. This fluorescence tool has also found a place in upper gastrointestinal surgery. In this literature review, we describe 6 main domains, namely identification of lymphatic structures, tumor localization, angiography of blood supply of different organs, thoracic duct visualization, tracheal blood flow analysis, and sentinel lymph node biopsy navigation.

Identification of lymphatic structures

A proper lymphadenectomy is a key issue associated with optimized gastric cancer outcomes since lymphadenectomy is recommended in advanced gastric cancer surgery(3) (4). Intraoperative lymphography may guide the surgeon in an appropriate lymph node (LN) dissection. The first fluorescence-guided lymphography using ICG for gastric cancer was performed by Herrera-Almario(5). It is clear that NIRF-guided lymphography has improved the visualization of LNs along gastric vessels. In a publication by Lan *et al.*,(6) more resected LNs were found in stations 4d and 6. In a randomized control trial by Chen *et al.*, it was clearly shown that a significantly higher number of LNs were dissected, and improved quality in optimized lymphadenectomy was achieved(7). Fluorescence lymphography may also be used for tailored lymphadenectomy, as presented by Baiocchi(8) where some nodes in station 11p would have been left in one patient, or in two patients. Additional potential LNs outside standard lymphadenectomy may have been stained with this fluorophore. In a study by Kim *et al.*, after complete lymphadenectomy, some ICG-stained tissue remained in 28% of patients. However, histologically confirmed LNs were only found in 4% of cases. In an article by Rho *et al.*, a mean of 2.1 non-fluorescent stations were found. This information, together with a 100% sensitivity in detecting metastatic LNs in fluorescent stations and a 100% negative predictive value in their study, may lead to the tailored extension of lymphadenectomy. However, further research is required, especially for obese patients considering the limited fluorescent penetration of ICG through adipose tissue(9). Some major limitations of this method need to be addressed. First, the problem of nodal involvement may limit the spread of ICG to

all required lymph node stations(10). Secondly, the effect of preoperative chemotherapy should be widely explored(11). Additionally, time, side, and depth of ICG injection should be standardized(12). Interestingly, Cianchi *et al.*(13) have shown that, in the case of linitis plastica, the occlusion of lymphatic vessels limits the spread of ICG. We must also recall preoperative endoscopic submucosal dissection (ESD), which may cause fibrosis and hamper proper lymphatic flow(9).

Tumor localization

Injection points for lymphography were used by Liu *et al.* as a landmark for early gastric cancer location during partial gastrectomy evaluation of the resection margin(14). Similar results of fluorescent marking with ICG were presented by Ushimaru *et al.*; in a group without ICG injection, a positive resection margin was found in 5% of cases, whereas in a group with ICG tumor position marking, it was found in 0% of cases(15). Tanaka *et al.* proposed a dosage of 0.1mL at a concentration of 0.5mg/mL for optimal visualization of injection sites(16). Diameters of ICG injection sites were much smaller (i.e. 21mm) than historical controls with India ink (52mm). Another valuable idea is the creation of a fluorescent rubber band generated from ICG mixed with a liquid rubber solution and placed endoscopically on the porcine stomach and colon. The localization of these clips was easily achieved, and the authors differentiated separate clips during observation. Additionally, the resection margins were sufficient after specimen extraction(17). Waiting for targeted fluorophores to become widely clinically available, this easy method may be used as a position of safe transection line during gastrectomy. Initial results of intraoperative imaging of colorectal cancer using a carcinoembryonic antigen (CEA)-targeting agent, namely SGM-101, were recently published, and a potential extension of these results to gastric cancer is anticipated(18).

Angiography of blood supply of different organs

After esophageal and gastric resection surgery, failure of anastomotic blood supply represents one of the main problems associated with complications. Intraoperative evaluation of blood perfusion using ICG is a new method which helps to predict anastomotic side leakage(19). Graft perfusion is an important factor of anastomotic integrity. It is a major clinical problem, with leakages in the range of 5 to 20% after esophagectomy(20–24), 1.2 to 6.7% after gastrectomy(25–29), and 0.5 to 6% after sleeve gastrectomy in bariatric surgery(30–33). In daily practice, tissue perfusion is evaluated using tissue color and vessel pulsation. Both are surgeon-dependent, and more reproducible objective methods for evaluation of graft viability

are necessary. In the literature review by Jansen *et al.*, several methods were examined, such as laser speckle contrast imaging, gastric tonometry, Doppler flowmetry, spectroscopy, infrared thermographic imaging, optical coherence tomography, and angiography(34–37). None of these approaches have been widely accepted in routine practice, mainly because of low reproducibility. NIR angiography using ICG has recently been adopted in many centers. Data from available studies are listed in Table 1 (38,39,48–57,40,58,41–47).

In a meta-analysis by Slooter *et al.*, a change in management based on ICG guidance was found in 24.55% of cases(59). The researchers also showed that less anastomotic leakage and graft necrosis were found in the ICG group, with an odds ratio (OR) of 0.30. The method of fluorescence angiography in the case of esophagectomy with gastric conduit anastomosis also revealed some important factors, which are discussed below.

- Time of visualization:

Conduit perfusion fluorescence angiography takes a median of – 37.5s(60). Shimada *et al.* described good visualization of the gastric cancer vasculature at the greater curvature after 1 min and visibility of the mucosal arterial network after 2 min(61). Karampiris *et al.*(46) presented that ICG angiography took 173s (\pm 74s), with no significant impact on operative time. In a publication by Mori *et al.* In gastric cancer patients, the time difference between ICG appearance on one or the other side of the anastomosis was an independent predictor of postoperative anastomotic leakage(62). In addition, different times of visualization for the gastric, jejunal, and duodenal sides were analyzed after total or subtotal gastrectomies(63).

- Complications:

The only temporal side effect of ICG intravenous injection and vascular angiography is a short drop in oxygenation immediately after injection. As presented by Murawa *et al.*(64), the saturation dropped to 90% without any further consequence. No other complications related to technical failures of different camera systems or other adverse events were reported.

- Quantification and validation:

Only a few studies have analyzed objective quantification of ICG angiography using an intensity and or time of fluorescence(44,45,49,52,54). The first description of different times of ICG fluorescence between the root of the right gastroepiploic artery and the tip of the conduit in patients with anastomotic leakage was described by Kumagai *et al.*(54). Inflow and outflow delays in perfusion analyzing two points of conduit vascularization was proposed by Yukaya *et al.*(52).

Koyanagi *et al.*(49) assessed fluorescence angiography flow speed by analyzing the length of the GC, the anatomical connection between the right and left gastroepiploic arteries, and the speed of NIR angiography. They found that flow speed was an independent prognostic factor for anastomotic leakage, with 1.76cm/s or less as a cut-off value. Interestingly, Noma *et al.* proposed a 30s rule as a cut-off time of demarcation for the optimal cervical anastomotic side. However, Kumagai *et al.*(44) prolonged this cut-off time up to 90 sec, obtaining promising results. In a publication by Nishikawa *et al.*, a quantitative analysis of blood flow using thermal imaging as compared to ICG angiography was presented to evaluate conduit viability(65). Both methods showed promising outcomes which need to be studied in the future.

- Intraoperative changes in surgical procedures:

In the systematic review by Van Daele *et al.* including 1,192 esophagectomy patients, ICGA was used perioperatively in 758 patients to guide esophageal reconstruction. A change in surgical strategy was found in 93 patients (12.4%). In 4 studies, the authors resected the tip of the conduit in the case of poor perfusion if sufficient length was obtained(66). Three authors changed the side of the anastomosis from end-to-side to end-to-end(43,47,64). Another 4 authors performed supercharge or super drainage arterial or venous anastomosis, increasing arterial perfusion or improving venous drainage of the cervical anastomosis(43,45,47,61). Modification in the surgical strategy led to an anastomotic leakage rate of 6.5%, which is much less than 20.5% without fluorescence angiography(66). This result is comparable to 6.3% of anastomotic leakage in the well-perfused group of patients.

- Dosage:

In the available literature, a wide range of ICG concentrations and volumes are used. In an animal model published by Diana *et al.*, a dose of 0.5mg/kg was found to be optimal using a virtual reality (VR) rendered software(67–69). Based on their studies, a dose of 0.2 to 0.5mg/kg is currently used in most studies. In the guidelines, it is proposed to use 2.5mg of ICG for colorectal and gastroesophageal anastomosis assessments(70).

Colon and small intestine conduits:

Although a gastric conduit is used after esophagectomy in some groups of patients, in most cases, other organ anastomoses with the remnant esophagus is also possible. In the case of free jejunal graft after pharyngoesophagectomy in a group of 26 patients, fluorescence angiography was performed(51). Five patients were diagnosed with venous anastomotic deficiency. In 3 cases, after fluorescence angiography, anastomosis was revised intraoperatively, and in 2 patients, an additional free jejunal graft transfer was required. The authors proposed that

T1/2max>9.6 sec might be a cutoff value to predict venous malperfusion in this group of patients.

A case report of a left colon supercharged with venous and arterial anastomoses in the neck conduit after esophagectomy has been presented(71). In a group of 23 patients, 1 patient required ileocolic graft reconstruction after esophagectomy in a report by Shimada *et al.*(61).

Tracheal blood flow analysis

Ischemia and necrosis of the tracheobronchial tree represent a severe complication which occurs after esophagectomy. This may originate from preoperative radiotherapy, trauma during surgery, or devascularization of the trachea and bronchus during esophageal resection. ICG fluorescence imaging can be used to assess tracheal blood perfusion during esophagectomy. Only one study published by Sugimura *et al.* focused on fluorescence imaging of the tracheal tree during this procedure(72). The researchers also considered the impact of preoperative treatment on the preservation of the right tracheobronchial artery. They visualized fluorescent tracheal blood flow in all included patients. Using a quantitative assessment, they found that lower trachea blood flow is linked with postoperative trachea ischemic changes in sputum discharge. Additionally, they revealed that resection of the right bronchial artery tended to be linked with a lower tracheal blood flow. In the case of resection of the right bronchial artery, preoperative neoadjuvant chemotherapy did not influence blood flow of the tracheobronchial tree.

Other methods to investigate tracheal blood flow changes during esophagectomy have been described. According to Fujita *et al.*, the hydrogen gas clearance method using electrochemically generated hydrogen in dogs showed that the ligation of the right bronchial artery is associated with 25% of tracheal blood flow. The same method used in 8 patients showed that esophagectomy decreased the tracheal blood flow by 43%, and by stopping the blood flow through the right bronchial artery, an additional 25% drop was found. Future studies are anticipated regarding this valuable topic.

Thoracic duct visualization

Chylothorax represents a major complication after esophagectomy, as reported in up to 43% of patients(73). Vecchiato *et al.* identified thoracic ducts in 19 patients undergoing minimally invasive esophageal resections(73). Percutaneous injection of ICG in the inguinal node area was performed, and the thoracic duct was visualized after 52.7 min. In 2 patients, due to oncologic radicality reasons, the successful ligation of the thoracic duct was performed under

fluorescence guidance. No postoperative chylothorax was noted. Some case reports have been published describing fluorescent thoracic duct fistula localization. In studies by Matsutani *et al.* and Kaburaki *et al.*, fluorescent node lymphography patients underwent reoperation on postoperative day 10, or in 2 patients, on postoperative day 28(74). ICG was injected into the bilateral inguinal node area subcutaneously, and the thoracic duct was visualized after 10 to 14 min(74,75). Another possibility of an injection site was proposed by Kaburaki *et al.* where 2ml of ICG were injected into the small bowel to identify the thoracic duct and to visualize the fistula in one patient.

Sentinel lymph node biopsy (SNB) navigation

Gastric cancer:

Gastrectomy with an appropriate lymphadenectomy, a procedure which has been standardized over many years, is the treatment of choice in gastric surgery(76,77). At early stages of gastric cancer, a less extensive resection and LN evaluation may be possible. Because of complex gastric lymphatic drainage, the concept of SNB is still under investigation. A systematic review by Cardoso *et al.* showed that the detection rate, sensitivity, and accuracy for the dye method and radioisotopes are similar(78). However, the negative rate was much higher for the dye method (34.7%) as compared to radioisotopes (18.5%). The meta-analysis by Wang *et al.* showed the diagnostic value of tracers, no matter what type was used(79). The dual tracer method seems to be most accurate. Looking for a new dye in 2004, for the first time, Nimura *et al.* proposed infrared electronic endoscopy for fluorescent ICG sentinel LN (SLN) visualization in GC(80). In a group of 84 patients with GC, fluorescent navigation detected all 11 patients with metastatic LNs, but four of the patients could not be visualized. The first SLN mapping with ICG which used NIR imaging was presented by Soltesz *et al.* in an animal model. Later on, mapping was reported in GC patients by Cusano *et al.*(81).

The method of ICG injection between the subserosa and submucosa represents another major issue(12). Some studies showed that a higher submucosal sensitivity could be found(12). The biggest concern about SLB in GC is that the tracer cannot come into metastatic LNs. In addition, the stomach lymphatic drainage is multidirectional and complex because cancer cell blockage of lymphatic lymph vessels and skip metastasis are possible in up to 11% of patients with LN metastasis. Another important issue associated with ICG LN biopsy is that it passes through higher tier LNs. An intraoperative injection shows a lower sensitivity as compared to a preoperative injection(82). It is an indication that we should wait about 20 min after intraoperative dye injection for optimal SLN mapping. In this study, false negative SNs and

metastatic non-SNs were found in the same group as SNs. Researchers have proposed to perform lymphatic basin dissection of LNs rather than using the pick-up method(83). An interesting possibility of CEA-targeted SLN detection in gastric cancer has been proposed by a Dutch group using SGM-101(84). The results of this study are awaited.

Esophageal cancer:

The concept of SLNs in esophageal cancer remains controversial as compared to other gastroesophageal cancers(43,85). Even in the case of T1 cancer, the rate of LN metastasis may reach up to 45%(86,87). Another challenge is a complicated lymphatic flow, which may vary from cervical to abdominal LN stations. To make it even more complicated, in about half of the patients with only one metastatic LN, skip metastasis may occur in distant lymphatic areas(88). This is even possible in the case of the negative involvement of the first regional LN compartment. The multidirectional lymphatic spread is a limitation of the usage of a blue dye. In addition, blue-stained LNs are often similar in color to antirachitic mediastinal nodes(89). In the case of the lower esophagus in the cardiac region, this is difficult to evaluate with preoperative lymphoscintigraphy with radiocolloids(90,91).

Lymphatic node mapping with radioactive tracers was first described by Kitagawa in 2000. Additionally, in their first pilot human trial, Krista *et al.* showed that ICG could be used to map LN drainage in esophageal cancer. Their study was performed in a group of 10 patients in whom a peritumoral injection of 2.5mL of ICG:HSA (indocyanine green:human serum albumin) was used to visualize the lymphatic pathway in esophageal cancer. In a study including a group of 9 patients with distal esophageal cancer, Schlottmann *et al.* presented intraoperative identification of LNs. The ICG was injected submucosally during endoscopy. Eight patients had metastatic LN along the left gastric artery, and one of them showed supradiaphragmatic LNs. In a group of 6 patients with distal esophageal cancer, Helminen *et al.* showed that intraoperative lymphatic mapping with NIR imaging could be useful to detect lymphatic stations which most likely present with metastatic disease and to guide the tailored extension of traditional lymphadenectomy. A Japanese study including 20 patients used the endoscopic submucosal injection of a dye called iopamidol with computed tomography (CT) to identify lymphatic drainage in SN(92). The comparison made in the group with ICG identification of SLN using CT lymphography was 100% and 95% in the ICG fluorescence group. One negative case of bulky metastasis was found. The authors pointed out that, during the long period between injection and LN dissection, the second and third lymph node stations were stained. The first usage of ICG LN mapping during minimally invasive esophagectomy was presented

by Hachey *et al.*(93). These authors used either ICG alone or ICG with HSA. Fluorescent LNs were not visualized in the case of three out of five patients with ICG usage without HSA. In the case of ICG with HSA, LN was fluorescent in all 4 patients. The visualization of fluorescent LN was difficult because of the shine-through effect.

A recent meta-analysis by Jimenez-Lillo *et al.* based on 6 studies revealed a high detection rate of ICG SNB in esophageal cancer of up to 89%(94). In addition, a trend towards a lower detection rate with a longer time between ICG injection and SLN biopsy was found. The authors also found a high rate of LN metastasis (84%). However, specificity was low (15%) because almost all nodes were stained with ICG. No difference was found in the type of esophageal cancer or the type of surgery performed.

Conclusion

New technology for NIR fluorescence image-guided surgery is used during gastroesophageal operations. In our review, we described 6 main domains in which research regarding the use of ICG fluorescence can be found. Although this technique is relatively new, many surgeons are looking for ways to implement such an approach in their daily practice. Surgery for gastric and esophageal diseases is complex, and it is sometimes associated with relatively significant complications. We have to look for and use all possible tools which will not only help to improve oncological results but also to decrease morbidity and mortality of such operations. Fluorescence image-guided surgery may be a good solution to improve surgical outcomes after gastroesophageal operations.

Acknowledgments

None.

Author Contributions

Study concept and design: Ekman Marcin, Girnyi Sergii, Marano Luigi, Roviello Franco, Diana Michele, Chand Manish, Polom Karol.

Acquisition of data: Ekman Marcin, Girnyi Sergii, Marano Luigi, Polom Karol

Study supervision: Roviello Franco, Diana Michele, Chand Manish, Polom Karol.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID

Marcin Ekman <https://orcid.org/0000-0003-3687-7464>

Sergii Girnyi <https://orcid.org/0000-0002-0988-5875>

Luigi Marano <https://orcid.org/0000-0002-9777-9588>

Franco Roviello <https://orcid.org/0000-0003-4122-1259>

Manish Chand <https://orcid.org/0000-0001-9086-8724>

Michele Diana <https://orcid.org/0000-0002-1390-8486>

Karol Polom <https://orcid.org/0000-0003-2011-4155>

References

1. Kitai T, Inomoto T, Miwa M, Shikayama T. Fluorescence navigation with indocyanine green for detecting sentinel lymph nodes in breast cancer. *Breast Cancer*. 2005;
2. Kirchherr A-K, Briel A, Mäder K. Stabilization of Indocyanine Green by Encapsulation within Micellar Systems. *Mol Pharm* [Internet]. 2009 Apr 6;6(2):480–91. Available from: <https://pubs.acs.org/doi/10.1021/mp8001649>
3. Sasako M, Sano T, Yamamoto S, Kurokawa Y, Nashimoto A, Kurita A, et al. D2 lymphadenectomy alone or with para-aortic nodal dissection for gastric cancer. *N Engl J Med*. 2008;
4. Songun I, Putter H, Kranenbarg EMK, Sasako M, van de Velde CJH. Surgical treatment of gastric cancer: 15-year follow-up results of the randomised nationwide Dutch D1D2 trial. *Lancet Oncol*. 2010;
5. Herrera-Almario G, Patane M, Sarkaria I, Strong VE. Initial report of near-infrared fluorescence imaging as an intraoperative adjunct for lymph node harvesting during robot-assisted laparoscopic gastrectomy. *J Surg Oncol*. 2016;
6. Lan Y-T, Huang K-H, Chen P-H, Liu C-A, Lo S-S, Wu C-W, et al. A pilot study of lymph node mapping with indocyanine green in robotic gastrectomy for gastric cancer. *SAGE Open Med*. 2017;
7. Chen QY, Xie JW, Zhong Q, Wang J Bin, Lin JX, Lu J, et al. Safety and Efficacy of Indocyanine Green Tracer-Guided Lymph Node Dissection during Laparoscopic Radical Gastrectomy in Patients with Gastric Cancer: A Randomized Clinical Trial. *JAMA Surg*. 2020;
8. Baiocchi GL, Molfino S, Molteni B, Quarti L, Arcangeli G, Manenti S, et al. Fluorescence-guided lymphadenectomy in gastric cancer: a prospective western series. *Updates Surg*. 2020;
9. Polom W, Markuszewski M, Rho YS, Matuszewski M. Use of invisible near infrared light fluorescence with indocyanine green and methylene blue in urology. Part 2. *Central European Journal of Urology*. 2014.
10. Choi HJ, Kim JM, Ryu JM, Kim I, Nam SJ, Yu J, et al. Patterns of axillary lymph node metastasis in breast cancer: A prospective single-center study. *J Breast Cancer*. 2018;
11. Tummers QRJG, Boogerd LSF, De Steur WO, Verbeek FPR, Boonstra MC, Handgraaf HJM, et al. Near-infrared fluorescence sentinel lymph node detection in gastric cancer: A pilot study *Clinical Trials Study*. *World J Gastroenterol*. 2016;
12. Yaguchi Y, Ichikura T, Ono S, Tsujimoto H, Sugasawa H, Sakamoto N, et al. How should tracers be injected to detect for sentinel nodes in gastric cancer - Submucosally from inside or subserosally from outside of the stomach? *J Exp Clin Cancer Res*. 2008;
13. Cianchi F, Indennitate G, Paoli B, Ortolani M, Lami G, Manetti N, et al. The Clinical Value of Fluorescent Lymphography with Indocyanine Green During Robotic Surgery for Gastric Cancer: a Matched Cohort Study. *J Gastrointest Surg*. 2020;
14. Liu M, Xing J, Xu K, Yuan P, Cui M, Zhang C, et al. Application of near-infrared fluorescence imaging with indocyanine green in totally laparoscopic distal gastrectomy. *J Gastric Cancer*. 2020;
15. Ushimaru Y, Omori T, Fujiwara Y, Yanagimoto Y, Sugimura K, Yamamoto K, et al. The Feasibility and Safety of Preoperative Fluorescence Marking with Indocyanine Green (ICG) in Laparoscopic Gastrectomy for Gastric Cancer. *J Gastrointest Surg*. 2019;
16. Tanaka C, Kanda M, Funasaka K, Miyahara R, Murotani K, Tanaka Y, et al. Detection of indocyanine green fluorescence to determine tumor location during laparoscopic gastrectomy for gastric cancer: Results of a prospective study. *Asian J Endosc Surg*.

- 2020;
17. Hyun JH, Kim SK, Kim KG, Kim HR, Lee HM, Park S, et al. A novel endoscopic fluorescent band ligation method for tumor localization. *Surg Endosc.* 2016;
 18. de Valk KS, Deken MM, Schaap DP, Meijer RP, Boogerd LS, Hoogstins CE, et al. Dose-Finding Study of a CEA-Targeting Agent, SGM-101, for Intraoperative Fluorescence Imaging of Colorectal Cancer. *Ann Surg Oncol.* 2020;
 19. Degett TH, Andersen HS, Gögenur I. Indocyanine green fluorescence angiography for intraoperative assessment of gastrointestinal anastomotic perfusion: a systematic review of clinical trials. *Langenbeck's Arch Surg [Internet].* 2016;401(6):767–75. Available from: <https://doi.org/10.1007/s00423-016-1400-9>
 20. Kassis ES, Kosinski AS, Ross P, Koppes KE, Donahue JM, Daniel VC. Predictors of anastomotic leak after esophagectomy: An analysis of the society of thoracic surgeons general thoracic database. *Ann Thorac Surg.* 2013;
 21. Biere SSAY, Maas KW, Cuesta MA, Van Der Peet DL. Cervical or thoracic anastomosis after esophagectomy for cancer: A systematic review and meta-analysis. *Digestive Surgery.* 2011.
 22. Sauvanet A, Mariette C, Thomas P, Lozac'h P, Segol P, Tiret E, et al. Mortality and morbidity after resection for adenocarcinoma of the gastroesophageal junction: Predictive factors. *J Am Coll Surg.* 2005;
 23. Sunpaweravong S, Ruangsri S, Laohawiriyakamol S, Mahattanobon S, Geater A. Prediction of major postoperative complications and survival for locally advanced esophageal carcinoma patients. *Asian J Surg.* 2012;
 24. Haga Y, Wada Y, Takeuchi H, Ikejiri K, Ikenaga M. Prediction of anastomotic leak and its prognosis in digestive surgery. *World J Surg.* 2011;
 25. Kim KM, An JY, Kim HI, Cheong JH, Hyung WJ, Noh SH. Major early complications following open, laparoscopic and robotic gastrectomy. *Br J Surg.* 2012;
 26. Kim MC, Kim W, Kim HH, Ryu SW, Ryu SY, Song KY, et al. Risk factors associated with complication following laparoscopy-assisted gastrectomy for gastric cancer: A large-scale Korean multicenter study. *Ann Surg Oncol.* 2008;
 27. Kitano S, Shiraishi N, Uyama I, Sugihara K, Tanigawa N, Inomata M, et al. A multicenter study on oncologic outcome of laparoscopic gastrectomy for early cancer in Japan. *Ann Surg.* 2007;
 28. Nagasako Y, Satoh S, Isogaki J, Inaba K, Taniguchi K, Uyama I. Impact of anastomotic complications on outcome after laparoscopic gastrectomy for early gastric cancer. *Br J Surg.* 2012;
 29. Lee KG, Lee HJ, Yang JY, Oh SY, Bard S, Suh YS, et al. Risk Factors Associated with Complication Following Gastrectomy for Gastric Cancer: Retrospective Analysis of Prospectively Collected Data Based on the Clavien-Dindo System. *J Gastrointest Surg.* 2014;
 30. Galloro G. Staple-line leak after sleeve gastrectomy in obese patients: A hot topic in bariatric surgery. *World J Gastrointest Endosc.* 2015;
 31. Abou Rached A, Basile M, El Masri H. Gastric leaks post sleeve gastrectomy: Review of its prevention and management. *World Journal of Gastroenterology.* 2014.
 32. Sakran N, Goitein D, Raziq A, Keidar A, Beglaibter N, Grinbaum R, et al. Gastric leaks after sleeve gastrectomy: A multicenter experience with 2,834 patients. *Surg Endosc.* 2013;
 33. Marchese M, Romano L, Giuliani A, Cianca G, Sibio A Di, Carlei F, et al. Corrigendum to “A case of intrasplenic displacement of an endoscopic double-pigtailstent as a treatment for laparoscopic sleeve gastrectomy leak” [(International Journal of Surgery Case Reports) (2018) 53 (367–369), (S2210261218305133),

- (10.1016/j.ijscr.2. International Journal of Surgery Case Reports. 2019.
34. Milstein DMJ, Ince C, Gisbertz SS, Boateng KB, Geerts BF, Hollmann MW, et al. Laser speckle contrast imaging identifies ischemic areas on gastric tube reconstructions following esophagectomy. *Med (United States)*. 2016;
 35. Linder G, Hedberg J, Björck M, Sundbom M. Perfusion of the gastric conduit during esophagectomy. *Dis Esophagus*. 2017;
 36. Pham TH, Perry KA, Enestvedt CK, Gareau D, Dolan JP, Sheppard BC, et al. Decreased conduit perfusion measured by spectroscopy is associated with anastomotic complications. *Ann Thorac Surg*. 2011;
 37. Tsekov C, Belyaev O, Tcholakov O, Tcherveniakov A. Intraoperative Doppler Assessment of Gastric Tube Perfusion in Esophagogastroplasty. *J Surg Res*. 2006;
 38. Koyanagi K, Ozawa S, Ninomiya Y, Oguma J, Kazuno A, Yatabe K, et al. Association between indocyanine green fluorescence blood flow speed in the gastric conduit wall and superior mesenteric artery calcification: predictive significance for anastomotic leakage after esophagectomy. *Esophagus*. 2020;
 39. Thammineedi SR, Patnaik SC, Saksena AR, Ramalingam PR, Nusrath S. The Utility of Indocyanine Green Angiography in the Assessment of Perfusion of Gastric Conduit and Proximal Esophageal Stump Against Visual Assessment in Patients Undergoing Esophagectomy: a Prospective Study. *Indian J Surg Oncol*. 2020;
 40. Nerup N, Svendsen MBS, Svendsen LB, Achiam MP. Feasibility and usability of real-time intraoperative quantitative fluorescent-guided perfusion assessment during resection of gastroesophageal junction cancer. *Langenbeck's Arch Surg*. 2020;
 41. Ishige F, Nabeya Y, Hoshino I, Takayama W, Chiba S, Arimitsu H, et al. Quantitative Assessment of the Blood Perfusion of the Gastric Conduit by Indocyanine Green Imaging. *J Surg Res*. 2019;
 42. Dalton BGA, Ali AA, Crandall M, Awad ZT. Near infrared perfusion assessment of gastric conduit during minimally invasive Ivor Lewis esophagectomy. *Am J Surg*. 2018;
 43. Kitagawa H, Namikawa T, Iwabu J, Fujisawa K, Uemura S, Tsuda S, et al. Assessment of the blood supply using the indocyanine green fluorescence method and postoperative endoscopic evaluation of anastomosis of the gastric tube during esophagectomy. *Surg Endosc*. 2018;
 44. Kumagai Y, Hatano S, Sobajima J, Ishiguro T, Fukuchi M, Ishibashi KI, et al. Indocyanine green fluorescence angiography of the reconstructed gastric tube during esophagectomy: Efficacy of the 90-second rule. *Dis Esophagus*. 2018;
 45. Noma K, Shirakawa Y, Kanaya N, Okada T, Maeda N, Ninomiya T, et al. Visualized Evaluation of Blood Flow to the Gastric Conduit and Complications in Esophageal Reconstruction. *J Am Coll Surg*. 2018;
 46. Karampinis I, Ronellenfisch U, Mertens C, Gerken A, Hetjens S, Post S, et al. Indocyanine green tissue angiography affects anastomotic leakage after esophagectomy. A retrospective, case-control study. *Int J Surg*. 2017;
 47. M. O, Y. T, Y. M, S. S, T. I, T. S, et al. Prevalence of anastomotic leak and the impact of indocyanine green fluorescein imaging for evaluating blood flow in the gastric conduit following esophageal cancer surgery. *Esophagus*. 2017;
 48. Schlottmann F, Patti MG. Evaluation of Gastric Conduit Perfusion during Esophagectomy with Indocyanine Green Fluorescence Imaging. *J Laparoendosc Adv Surg Tech*. 2017;
 49. Koyanagi K, Ozawa S, Oguma J, Kazuno A, Yamazaki Y, Ninomiya Y, et al. Blood flow speed of the gastric conduit assessed by indocyanine green fluorescence: New predictive evaluation of anastomotic leakage after esophagectomy. *Med (United*

- States). 2016;
50. Campbell C, Reames MK, Robinson M, Symanowski J, Salo JC. Conduit Vascular Evaluation is Associated with Reduction in Anastomotic Leak After Esophagectomy. *J Gastrointest Surg.* 2015;
 51. Kamiya K, Unno N, Miyazaki S, Sano M, Kikuchi H, Hiramatsu Y, et al. Quantitative assessment of the free jejunal graft perfusion. *J Surg Res.* 2015;
 52. Yukaya T, Saeki H, Kasagi Y, Nakashima Y, Ando K, Imamura Y, et al. Indocyanine Green Fluorescence Angiography for Quantitative Evaluation of Gastric Tube Perfusion in Patients Undergoing Esophagectomy. *J Am Coll Surg.* 2015;
 53. Zehetner J, DeMeester SR, Alicuben ET, Oh DS, Lipham JC, Hagen JA, et al. Intraoperative assessment of perfusion of the gastric graft and correlation with anastomotic leaks after esophagectomy. In: *Annals of Surgery.* 2015.
 54. Kumagai Y, Ishiguro T, Haga N, Kuwabara K, Kawano T, Ishida H. Hemodynamics of the reconstructed gastric tube during esophagectomy: Assessment of outcomes with indocyanine green fluorescence. *World J Surg.* 2014;
 55. Sarkaria IS, Bains MS, Finley DJ, Adusumilli PS, Huang J, Rusch VW, et al. Intraoperative near-infrared fluorescence imaging as an adjunct to robotic-assisted minimally invasive esophagectomy. *Innov Technol Tech Cardiothorac Vasc Surg.* 2014;
 56. Rino Y, Yukawa N, Sato T, Yamamoto N, Tamagawa H, Hasegawa S, et al. Visualization of blood supply route to the reconstructed stomach by indocyanine green fluorescence imaging during esophagectomy. *BMC Med Imaging.* 2014;
 57. Pacheco PE, Hill SM, Henriques SM, Paulsen JK, Anderson RC. The novel use of intraoperative laser-induced fluorescence of indocyanine green tissue angiography for evaluation of the gastric conduit in esophageal reconstructive surgery. *Am J Surg.* 2013;
 58. Kubota K, Yoshida M, Kuroda J, Okada A, Ohta K, Kitajima M. Application of the HyperEye Medical System for esophageal cancer surgery: A preliminary report. *Surg Today.* 2013;
 59. Slooter MD, Eshuis WJ, Cuesta MA, Gisbertz SS, van Berge Henegouwen MI. Fluorescent imaging using indocyanine green during esophagectomy to prevent surgical morbidity: a systematic review and meta-analysis. *J Thorac Dis.* 2019 Apr;11(Suppl 5):S755–65.
 60. Purich K, Dang JT, Poonja A, Sun WYL, Bigam D, Birch D, et al. Intraoperative fluorescence imaging with indocyanine green in hepatic resection for malignancy: a systematic review and meta-analysis of diagnostic test accuracy studies. *Surgical Endoscopy.* 2020.
 61. Shimada Y, Okumura T, Nagata T, Sawada S, Matsui K, Hori R, et al. Usefulness of blood supply visualization by indocyanine green fluorescence for reconstruction during esophagectomy. *Esophagus.* 2011;
 62. Mori M, Shuto K, Hirano A, Kosugi C, Narushima K, Hosokawa I, et al. A Novel Parameter Identified Using Indocyanine Green Fluorescence Angiography may Contribute to Predicting Anastomotic Leakage in Gastric Cancer Surgery. *World J Surg.* 2020;
 63. Huh YJ, Lee HJ, Kim TH, Choi YS, Park JH, Son YG, et al. Efficacy of Assessing Intraoperative Bowel Perfusion with Near-Infrared Camera in Laparoscopic Gastric Cancer Surgery. *J Laparoendosc Adv Surg Tech.* 2019;
 64. Murawa D, Hünerbein M, Spychała A, Nowaczyk P, Połom K, Murawa P. Indocyanine green angiography for evaluation of gastric conduit perfusion during esophagectomy - First experience. *Acta Clin Belg.* 2012;

65. Nishikawa K, Fujita T, Yuda M, Tanaka Y, Matsumoto A, Tanishima Y, et al. Quantitative Assessment of Blood Flow in the Gastric Conduit With Thermal Imaging for Esophageal Reconstruction. *Ann Surg.* 2020;
66. Van Daele E, Van Nieuwenhove Y, Ceelen W, Vanhove C, Braeckman BP, Hoorens A, et al. Near-infrared fluorescence guided esophageal reconstructive surgery: A systematic review. *World Journal of Gastrointestinal Oncology.* 2019.
67. Diana M, Noll E, Diemunsch P, Dallemagne B, Benahmed MA, Agnus V, et al. Enhanced-reality video fluorescence: A real-time assessment of intestinal viability. *Ann Surg.* 2014;
68. Diana M, Halvax P, Dallemagne B, Nagao Y, Diemunsch P, Charles AL, et al. Real-time navigation by fluorescence-based enhanced reality for precise estimation of future anastomotic site in digestive surgery. *Surg Endosc.* 2014;
69. Diana M, Agnus V, Halvax P, Liu YY, Dallemagne B, Schlagowski AI, et al. Intraoperative fluorescence-based enhanced reality laparoscopic real-time imaging to assess bowel perfusion at the anastomotic site in an experimental model. *British Journal of Surgery.* 2015.
70. L. van M, H.J.M. H, M. D, J. D, T. I, A.L. V, et al. A practical guide for the use of indocyanine green and methylene blue in fluorescence-guided abdominal surgery. *J Surg Oncol.* 2018;
71. Charalabopoulos A, Kordzadeh A, Sdralis E, Lorenzi B, Ahmad F. Thoracoscopic total esophagogastrectomy with supercharged colon interposition for the treatment of esophageal adenocarcinoma in situs inversus. *Acta Chir Belg.* 2019;
72. Sugimura K, Miyata H, Shinno N, Yanagimoto Y, Yamamoto K, Yasui M, et al. Indocyanine Green Fluorescence Imaging of the Tracheal Blood Flow During Esophagectomy. *J Surg Res.* 2019;
73. Vecchiato M, Martino A, Sponza M, Uzzau A, Ziccarelli A, Marchesi F, et al. Thoracic duct identification with indocyanine green fluorescence during minimally invasive esophagectomy with patient in prone position. *Dis Esophagus.* 2020;
74. Matsutani T, Hirakata A, Nomura T, Hagiwara N, Matsuda A, Yoshida H, et al. Transabdominal Approach for Chylorrhea after Esophagectomy by Using Fluorescence Navigation with Indocyanine Green. *Case Rep Surg.* 2014;
75. Kamiya K, Unno N, Konno H. Intraoperative indocyanine green fluorescence lymphography, a novel imaging technique to detect a chyle fistula after an esophagectomy: Report of a case. *Surg Today.* 2009;
76. De Manzoni G, Marrelli D, Baiocchi GL, Morgagni P, Saragoni L, Degiuli M, et al. The Italian Research Group for Gastric Cancer (GIRCG) guidelines for gastric cancer staging and treatment: 2015. *Gastric Cancer.* 2017;
77. Japanese gastric cancer treatment guidelines 2018 (5th edition). *Gastric Cancer.* 2020;
78. Cardoso R, Bocicariu A, Dixon M, Yohanathan L, Seevaratnam R, Helyer L, et al. What is the accuracy of sentinel lymph node biopsy for gastric cancer? A systematic review. *Gastric Cancer.* 2012.
79. Wang Z, Dong ZY, Chen JQ, Liu JL. Diagnostic value of sentinel lymph node biopsy in gastric cancer: A meta-analysis. *Ann Surg Oncol.* 2012;
80. Nimura H, Narimiya N, Mitsumori N, Yamazaki Y, Yanaga K, Urashima M. Infrared ray electronic endoscopy combined with indocyanine green injection for detection of sentinel nodes of patients with gastric cancer. In: *British Journal of Surgery.* 2004.
81. Soltesz EG, Kim S, Kim SW, Laurence RG, De Grand AM, Parungo CP, et al. Sentinel lymph node mapping of the gastrointestinal tract by using invisible light. *Ann Surg Oncol.* 2006;
82. Balagué C, Gomez I. Preoperative and Intraoperative Lymphatic Mapping for Sentinel

- Node Biopsy in Cancers of the Gastrointestinal Tract. In: Atlas of Lymphoscintigraphy and Sentinel Node Mapping. 2020.
83. Lee YJ, Ha WS, Park ST, Choi SK, Hong SC, Park JW. Which biopsy method is more suitable between a basin dissection and pick-up biopsy for sentinel nodes in laparoscopic sentinel-node navigation surgery (LSNNS) for gastric cancer? *J Laparoendosc Adv Surg Tech.* 2008;
 84. Vuijk FA, Hilling DE, Mieog JSD, Vahrmeijer AL. Fluorescent-guided surgery for sentinel lymph node detection in gastric cancer and carcinoembryonic antigen targeted fluorescent-guided surgery in colorectal and pancreatic cancer. *Journal of Surgical Oncology.* 2018.
 85. Schlag PM, Bembenek A, Schulze T. Sentinel node biopsy in gastrointestinal-tract cancer. *Eur J Cancer.* 2004;
 86. Hagens ERC, van Berge Henegouwen MI, Gisbertz SS. Distribution of lymph node metastases in esophageal carcinoma patients undergoing upfront surgery: A systematic review. *Cancers.* 2020.
 87. Tachibana M, Kinugasa S, Shibakita M, Tonomoto Y, Hattori S, Hyakudomi R, et al. Surgical treatment of superficial esophageal cancer. *Langenbeck's Archives of Surgery.* 2006.
 88. Ando N, Ozawa S, Kitagawa Y, Shinozawa Y, Kitajima M. Improvement in the results of surgical treatment of advanced squamous esophageal carcinoma during 15 consecutive years. *Ann Surg.* 2000;
 89. Yasuda S, Shimada H, Chino O, Tanaka H, Kenmochi T, Takechi M, et al. Sentinel lymph node detection with Tc-99m tin colloids in patients with esophagogastric cancer. *Jpn J Clin Oncol.* 2003;
 90. Hayashi H, Tangoku A, Suga K, Shimizu K, Ueda K, Yoshino S, et al. CT lymphography-navigated sentinel lymph node biopsy in patients with superficial esophageal cancer. *Surgery.* 2006;
 91. Tangoku A, Seike J, Nakano K, Nagao T, Honda J, Yoshida T, et al. Current status of sentinel lymph node navigation surgery in breast and gastrointestinal tract. *Journal of Medical Investigation.* 2007.
 92. Yuasa Y, Seike J, Yoshida T, Takechi H, Yamai H, Yamamoto Y, et al. Sentinel lymph node biopsy using intraoperative indocyanine green fluorescence imaging navigated with preoperative CT lymphography for superficial esophageal cancer. *Ann Surg Oncol.* 2012;
 93. Hachey KJ, Gilmore DM, Armstrong KW, Harris SE, Hornick JL, Colson YL, et al. Safety and feasibility of near-infrared image-guided lymphatic mapping of regional lymph nodes in esophageal cancer. *J Thorac Cardiovasc Surg.* 2016;
 94. Jimenez-Lillo J, Villegas-Tovar E, Momblan-Garcia D, Turrado-Rodriguez V, Ibarzabal-Olano A, De Lacy B, et al. Performance of Indocyanine-Green Imaging for Sentinel Lymph Node Mapping and Lymph Node Metastasis in Esophageal Cancer: Systematic Review and Meta-Analysis. *Ann Surg Oncol [Internet].* 2021; Available from: <http://europepmc.org/abstract/MED/33515138>