

Electrochemical Deconstructive Functionalization of Cycloalkanols via Alkoxy Radicals Enabled by Proton-Coupled Electron Transfer

Mishra Deepak Hareram, Albara A. M. A. El Gehani, James Harnedy, Alex C. Seastram, Andrew C. Jones, Matthew Burns, Thomas Wirth, Duncan L. Browne, and Louis C. Morrill*



Cite This: *Org. Lett.* 2022, 24, 3890–3895



Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: Herein, we report a new electrochemical method for alkoxy radical generation from alcohols using a proton-coupled electron transfer (PCET) approach, showcased via the deconstructive functionalization of cycloalkanols. The electrochemical method is applicable across a diverse array of substituted cycloalkanols, accessing a broad range of synthetically useful distally functionalized ketones. The orthogonal derivatization of the products has been demonstrated through chemoselective transformations, and the electrochemical process has been performed on a gram scale in continuous single-pass flow.




Alkoxy radicals are an important class of oxygen-centered radicals, which consist of an alkyl group bound to an electrophilic oxygen radical center.¹ They are particularly high energy species due to the lack of stabilization provided by mesomeric effects and spin density delocalization found in other O-centered radicals, such as aryloxy radicals. Despite this, alkoxy radicals exhibit well-defined yet diverse reactivity, including β -scission processes, hydrogen atom transfers (HATs), and addition to π -systems (Scheme 1A). As such, these privileged intermediates have been successfully employed across a plethora of powerful transformations spanning

selective C–H functionalization, C–C bond activation, and heterocycle synthesis to access valuable products, including within complex molecule synthesis.² Photoredox catalysis has enabled alkoxy radical generation employing various radical precursors including peroxides,³ N-alkoxyphthalimides,⁴ N-alkoxy-pyridiniums,⁵ N-alkoxybenzimidazoles,⁶ N-alkoxytriazoliums,⁷ and oximino acids.⁸ Nevertheless, the use of O-functionalized precursors incurs additional synthetic effort for installation, particularly within more complex molecules, and decreases step/atom economy.⁹ Recent advances in transition metal catalysis and photoredox catalysis have enabled alkoxy radical intermediates to be accessed directly from unprotected aliphatic alcohols (Scheme 1B).¹ However, many of these approaches require the use of stoichiometric oxidants (e.g., $K_2S_2O_8$ or hypervalent iodine reagents) and/or precious metal (photo)catalysts. As such, the development of alternative, more sustainable, approaches for the generation of alkoxy radicals from commodity aliphatic alcohols is an important and timely pursuit.


The development of electrochemical methods¹⁰ for the generation of alkoxy radicals has received surprisingly little attention to date and remains significantly underdeveloped. In 1981, Swenton and co-workers reported the dearomative dimethoxylation of electron rich naphthalenes (Scheme 1C).¹¹ The same approach was subsequently employed for the dimethoxylation of 4-methoxyanilines¹² and dimethoxyben-

Scheme 1. Context


A. Alkoxy radical reactivity modes



B. Generation of alkoxy radicals from aliphatic alcohols



C. Dearomative dimethoxylation of electron rich naphthalenes (Swenton *et al.*)



Received: May 5, 2022

Published: May 23, 2022




zenes.¹³ The authors proposed that the mechanism for these transformations involves the formation of methoxy radicals via direct anodic oxidation of methoxide anions and evidence to support this was provided through radical trapping experiments and electron spin resonance (ESR).^{13a} It was found that the broad potential window of boron-doped diamond or platinum anodes was essential for generating a sufficient concentration of methoxy radicals due to the high oxidation potential required for direct anodic oxidation of methoxide anions,^{14,15} which inherently limits the broader application of this approach in synthesis.^{16,17}

Cognizant of the opportunities to make advances in this underdeveloped domain, our research team reported the manganese-catalyzed electrochemical deconstructive chlorination of tertiary cyclopropanols and cyclobutanols.¹⁸ However, the method could not be more broadly applied toward larger and more widely commercially available ring sizes or alternative functionalizations.¹⁹ Proton-coupled electron transfer (PCET),²⁰ which involves the concerted movement of a proton and an electron in a single elementary step, can be used to overcome the energetic limitations of classical HAT reagents for the formation of alkoxy radicals. For example, spectroscopic studies by Baciocchi and co-workers described the β -scission of 1-arylalkanol radical cations,²¹ and this strategy has been applied toward the development of various photochemical transformations.²² To develop a more general electrochemical method to access and utilize alkoxy radicals, we envisaged an electrochemically driven PCET approach,²³ and herein we report the successful realization of this strategy. During the latter stages of this investigation, Onomura and co-workers reported an electrophotochemical deconstructive bromination of cycloalkanols, which involves the electrochemical generation of hypobromite intermediates via anodic bromide oxidation followed by visible light-promoted homolysis of the weak O–Br bond to generate an alkoxy radical.²⁴

To commence our studies, the deconstructive bromination of 1-(4-methoxyphenyl)cyclohexan-1-ol **1** to form ϵ -bromo ketone **2** was selected for reaction optimization (Table 1).²⁵ The proposed electrochemical generation of alkoxy radicals by PCET was contingent upon efficient electrochemical oxidation of the 4-methoxyphenyl ring present within **1** ($E_{p/2} = 1.03$ V vs Fc/Fc⁺). Employing BrCCl₃ as the brominating agent, collidine as the Brønsted base, and LiClO₄ as the supporting electrolyte in MeCN/TFE (12:1, [1] = 0.05 M) using galvanostatic conditions ($i = 10$ mA, $j_{\text{anode}} = 7.1$ mA/cm², $Q = 4.5$ F/mol), a graphite anode and a platinum foil cathode in an undivided cell at 25 °C under N₂, gave **2** in an encouraging 32% NMR yield (entry 1), with 68% recovered **1**. To improve conversion, we investigated the impact of employing a range of triarylamine²⁶ and triarylimidazole²⁷ redox mediators,²⁵ and it was found that the addition of N(4-NO₂C₆H₄)₂Ph (5 mol %) ($E_{\text{ox}} = 0.91$ V vs Fc/Fc⁺) gave quantitative conversion to **2** with a 94% isolated yield (entry 2).²⁸ The increase in conversion using a triarylamine mediator may be attributed to more efficient electron transfer and productive intermolecular reactivity through an electrochemical–chemical (EC') mechanism.²⁹ The use of a redox mediator also lowers the required applied potential below the oxidation potential of the substrate, and these milder electrochemical conditions can minimize side reactions, while improving functional group tolerance. No product formation was observed in the absence of electricity or in the absence of collidine (entries 3 and 4). Lowering the loading of collidine, BrCCl₃, or redox mediator

Table 1. Optimization of the Electrochemical Process^a




entry	variation from "standard" conditions	yield ^b (%)
1	no N(4-NO ₂ C ₆ H ₄) ₂ Ph	32
2	none	>98 (94)
3	no electricity	<2
4	no collidine	<2
5	collidine (2 equiv)	81
6	BrCCl ₃ (1.1 equiv)	49
7	N(4-NO ₂ C ₆ H ₄) ₂ Ph (1 mol %)	50
8	Ni plate cathode	63
9	[TBAPF ₆] = 0.1 M as supporting electrolyte	80
10	MeCN/HFIP (12:1) as solvent	49
11	CH ₂ Cl ₂ /TFE (12:1) as solvent	74
12	$i = 7.5$ mA, $j_{\text{anode}} = 5.3$ mA/cm ²	74
13 ^c	$Q = 2.5$ F/mol	56

^aReactions performed with 0.3 mmol of **1** using the ElectraSyn 2.0 batch electrochemical reactor. ^bAs determined by ¹H NMR analysis of the crude reaction mixture with 1,3,5-trimethoxybenzene as the internal standard. Isolated yield given in parentheses. ^c121 min reaction time.

all reduced the NMR yield of **2** (entries 5–7), as did variation of cathode material, supporting electrolyte, and solvent (entries 8–11). Reducing the current and charge passed both had a detrimental impact upon product formation (entries 12 and 13).

With optimized reaction conditions in hand, the full scope of the electrochemical process was explored (Scheme 2). Initially, it was found that the deconstructive bromination of 3-, 4-, 5-, 6-, 7-, 8-, and 12-membered cycloalkanols proceeded efficiently to access the corresponding distally brominated ketone products **2–8** in high yields, overcoming a key limitation of our previous electrochemical approach, which was restricted to cyclopropanols and cyclobutanols.¹⁸ The 4-methoxyphenyl ring can be replaced with a variety of alternative electron-rich and/or extended aromatic systems to give brominated ketone products **9–16** in good isolated yields. No conversion to product **13** was observed employing the previously optimized reaction conditions (Table 1, entry 2), which may be attributed to the higher oxidation potential of the cycloalkanol substrate ($E_{p/2} = 1.28$ V vs Fc/Fc⁺). However, it was found that employing N(4-Cl₂-NO₂C₆H₃)₃ (10 mol %) ($E_{\text{ox}} = 1.50$ V vs Fc/Fc⁺) as the redox mediator enabled access to product **13**, which was isolated in 48% yield. Employing 1-phenylcyclohexan-1-ol as substrate ($E_{p/2} = 1.60$ V vs Fc/Fc⁺), resulted in quantitative recovery of starting material under the same reaction conditions. A substituted adamantane scaffold was converted to bicyclo[3.3.1] product **17** as a single diastereoisomer in 84% isolated yield. An unsymmetrical substituted *trans*-decalin substrate underwent regioselective deconstructive bromination to access 1,2-disubstituted cyclohexane product **18** as a 1:1 mixture of diastereoisomers in 70% isolated yield. Due to the reversibility of β -scission processes involving alkoxy radicals,¹ the regioselectivity employing unsymmetrical cycloalkanols proceeds via selective cleavage


Scheme 2. Substrate Scope^m

^mReactions performed with 0.3 mmol of cycloalkanol using the ElectraSyn 2.0 batch electrochemical reactor with isolated yields after chromatographic purification quoted unless stated otherwise. ^aAs determined by ¹H NMR analysis of the crude reaction mixture with 1,3,5-trimethoxybenzene as the internal standard. ^bQ = 6 F/mol. ^cN(4-NO₂C₆H₄)₂Ph (10 mol %), i = 20 mA, Q = 9 F/mol. ^dQ = 12.4 F/mol. ^eN(4-Cl-2-NO₂C₆H₃)₃ (10 mol %), i = 20 mA, Q = 9 F/mol. ^fi = 12.5 mA, Q = 9 F/mol. ^gN(4-Cl-2-NO₂C₆H₃)₃ (5 mol %). ^hQ = 9 F/mol. ⁱCCl₄ (2 equiv.). ^jN-Iodosuccinimide (5 equiv.). ^kWithout N₂ atmosphere and in the absence of BrCCl₃.

of the more substituted β-C–C bond that produces the more stabilized C-centered radical.³⁰ Indeed, where relevant, only a single product regioisomer was observed for all substrates. A range of substituted cyclohexanols, piperidines, and an azepane were found to be compatible with the electrochemical method, providing access to the corresponding distally brominated ketones 19–26 in high yields. The high regioselectivity observed in the formation of products 20 and 26 was attributed to subtle inductive effects resulting in differences in the relative stability of radical intermediates formed upon β-C–C homolysis. Employing an acyclic substrate resulted in the formation of ketone 27 in 85% isolated yield. Deconstructive chlorination and iodination were demonstrated by employing CCl₄ and N-iodosuccinimide as electrophilic polarized SOMophiles to access products 28 and 29, respectively. Finally, in the absence of BrCCl₃ and performing the reaction under air without degassing the reaction mixture prior to electrolysis, keto-aldehyde 30 was isolated in 70% yield.


To demonstrate product utility, the orthogonal functionalization of ε-bromo ketone 2 was investigated (Scheme 3A). It was found that the C(sp³)–Br functionality could serve as a coupling partner in a nickel-catalyzed ball-milling enabled cross-electrophile coupling to access arylated ketone 31 in 78% isolated yield.³¹ Furthermore, the 4-methoxyphenyl ketone functionality within 2 was converted to the corresponding ester via Baeyer–Villiger oxidation to give 32 in 84% isolated yield. By employing a HPLC pump in combination with commercially available Ammonite8 flow electroreactor (volume = 1 mL, i = 840 mA),³² 10 mmol of 1-(4-methoxyphenyl)cyclohexan-1-ol 1 was converted to ε-bromo ketone 2 in 83% isolated yield (2.30 g) in a continuous single-pass. In comparison to batch, the flow process exhibits higher productivity (9.4 mmol/h vs 0.08 mmol/h), requires less charge for full consumption of 1 (Q = 2.9 F/mol vs Q = 4.5 F/mol), and employs an increased current density (j_{anode} = 42 mA/cm² vs j_{anode} = 7.1 mA/cm²).

Scheme 3. Product Utility and Reaction Scale Up in Flow



Using cyclic voltammetry, it was found that $N(4\text{-NO}_2\text{C}_6\text{H}_4)_2\text{Ph}$ undergoes irreversible oxidation at 0.91 V vs Fc/Fc^+ ,²⁵ which is attributed to the generation of the corresponding triarylamine radical cation. An increase in the oxidation current was observed upon addition of 1-(4-methoxyphenyl)cyclohexan-1-ol **1**, which suggested that the triarylamine radical cation is consumed by **1** to reform $N(4\text{-NO}_2\text{C}_6\text{H}_4)_2\text{Ph}$ and provided support for the proposed role of the triarylamine acting as a redox mediator as part of an electrochemical-chemical (EC') mechanism.²⁹ When methyl ether cyclohexane **33** was subjected to the standard electrochemical reaction conditions, no ketone **2** was observed, with 88% starting material recovered (Scheme 4A), which indicated that C–C bond cleavage does not proceed in the absence of a hydroxyl functional group. Deshielding of the hydroxyl proton shift within **1** upon the addition of collidine in CD_3CN was observed, which indicated the presence of a hydrogen bond

Scheme 4. Mechanistic Studies



adduct between **1** and the base²⁵ and provided support for a subsequent concerted multisite PCET.³³ Analysis of the reaction mixture produced using optimized conditions (Table 1, entry 2) by ^1H NMR in CD_3CN revealed that the majority of the trichloromethyl radical is converted to CHCl_3 (66% NMR yield),²⁵ which may occur via hydrogen atom transfer or via cathodic reduction followed by protonation of the trichloromethyl anion. As such, a plausible reaction mechanism initiates with anodic oxidation of $N(4\text{-NO}_2\text{C}_6\text{H}_4)_2\text{Ph}$ ($E_{\text{ox}} = 0.91$ V vs Fc/Fc^+) to form a triarylamine radical cation (Scheme 4B), which subsequently oxidizes the electron-rich aromatic ring within **1** ($E_{\text{p}/2} = 1.03$ V vs Fc/Fc^+). A concerted PCET involving alcohol deprotonation by collidine and oxidation by the internal aryl radical cation forms the alkoxy radical. Subsequent β -scission generates a distal alkyl radical, which is trapped with BrCCl_3 to form a new C–Br bond. The counter cathodic reaction is hydrogen gas production via proton reduction and/or chloroform production via reduction of trichloromethyl radicals and subsequent protonation.

■ ASSOCIATED CONTENT

S Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.2c01552>.

Optimization data, experimental procedures, characterization of new compounds, and spectral data (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Louis C. Morrill – Cardiff Catalysis Institute, School of Chemistry, Cardiff University, Cardiff CF10 3AT, United Kingdom; orcid.org/0000-0002-6453-7531; Email: MorrillLC@cardiff.ac.uk

Authors

Mishra Deepak Hareram – Cardiff Catalysis Institute, School of Chemistry, Cardiff University, Cardiff CF10 3AT, United Kingdom

Albara A. M. A. El Gehani – Cardiff Catalysis Institute, School of Chemistry, Cardiff University, Cardiff CF10 3AT, United Kingdom

James Harnedy – Cardiff Catalysis Institute, School of Chemistry, Cardiff University, Cardiff CF10 3AT, United Kingdom

Alex C. Seastram – Cardiff Catalysis Institute, School of Chemistry, Cardiff University, Cardiff CF10 3AT, United Kingdom

Andrew C. Jones – Cardiff Catalysis Institute, School of Chemistry, Cardiff University, Cardiff CF10 3AT, United Kingdom

Matthew Burns – Chemical Development, Pharmaceutical Technology & Development, Operations, AstraZeneca, Macclesfield SK10 2NA, United Kingdom; orcid.org/0000-0002-3298-0344

Thomas Wirth – School of Chemistry, Cardiff University, Cardiff CF10 3AT, United Kingdom

Duncan L. Browne – Department of Pharmaceutical and Biological Chemistry, University College London, School of Pharmacy, London W1CN 1AX, United Kingdom; orcid.org/0000-0002-8604-229X

Complete contact information is available at:
<https://pubs.acs.org/10.1021/acs.orglett.2c01552>

Notes

The authors declare no competing financial interest.
Data Access: The data that support the findings of this study are openly available in the Cardiff University data catalogue at <http://doi.org/10.17035/d.2022.0197737325>.

ACKNOWLEDGMENTS

We gratefully acknowledge the School of Chemistry, Cardiff University and AstraZeneca for generous support, the EPSRC for a standard research grant (M.D.H., EP/R006504/1) and doctoral training grants for PhD studentships (A.A.M.A.E.G. and A.S., EP/R513003/1), the European Union, WEFO and Cardiff University for a Knowledge Economy and Skills Scholarship (A.C.J.).

REFERENCES

- (1) For selected reviews, see: (a) Wu, X.; Zhu, C. Recent advances in alkoxy radical-promoted C–C and C–H bond functionalization starting from free alcohols. *Chem. Commun.* **2019**, *55*, 9747–9756. (b) Tsui, E.; Wang, H.; Knowles, R. R. Catalytic generation of alkoxy radicals from unfunctionalized alcohol. *Chem. Sci.* **2020**, *11*, 11124–11141. (c) Chang, L.; An, Q.; Duan, L.; Feng, K.; Zuo, Z. Alkoxy Radicals See the Light: New Paradigms of Photochemical Synthesis. *Chem. Rev.* **2022**, *122*, 2429–2486.
- (2) For selected examples, see: (a) de Armas, P.; Francisco, C. G.; Suarez, E. Fragmentation of carbohydrate anomeric alkoxy radicals. Tandem.β-fragmentation-cyclization of alcohols. *J. Am. Chem. Soc.* **1993**, *115*, 8865–8866. (b) Yamashita, S.; Naruko, A.; Nakazawa, Y.; Zhao, L.; Hayashi, Y.; Hiram, M. Total Synthesis of Limonin. *Angew. Chem., Int. Ed.* **2015**, *54*, 8538–8541.
- (3) DiRocco, D. A.; Dykstra, K.; Krska, S.; Vachal, P.; Conway, D. V.; Tudge, M. Late-Stage Functionalization of Biologically Active Heterocycles Through Photoredox Catalysis. *Angew. Chem., Int. Ed.* **2014**, *53*, 4802–4806.
- (4) Zhang, J.; Li, Y.; Zhang, F.; Hu, C.; Chen, Y. Generation of Alkoxy Radicals by Photoredox Catalysis Enables Selective C(sp³)–H Functionalization under Mild Reaction Conditions. *Angew. Chem., Int. Ed.* **2016**, *55*, 1872–1875.
- (5) Kim, I.; Park, B.; Kang, G.; Kim, J.; Jung, H.; Lee, H.; Baik, M.-H.; Hong, S. Visible-Light-Induced Pyridylation of Remote C(sp³)–H Bonds by Radical Translocation of N-Alkoxyopyridinium Salts. *Angew. Chem., Int. Ed.* **2018**, *57*, 15517–15522.
- (6) Zheng, W.; Morales-Rivera, C. A.; Lee, J. W.; Liu, P.; Ngai, M.-Y. Catalytic C–H Trifluoromethoxylation of Arenes and Heteroarenes. *Angew. Chem., Int. Ed.* **2018**, *57*, 9645–9649.
- (7) Zheng, W.; Lee, J. W.; Morales-Rivera, C. A.; Liu, P.; Ngai, M.-Y. Redox-Active Reagents for Photocatalytic Generation of the OCF₃ Radical and (Hetero)Aryl C–H Trifluoromethoxylation. *Angew. Chem., Int. Ed.* **2018**, *57*, 13795–13799.
- (8) Herron, A. N.; Liu, D.; Xia, G.; Yu, J.-Q. δ-C–H Mono- and Dihalogenation of Alcohols. *J. Am. Chem. Soc.* **2020**, *142*, 2766–2770.
- (9) Trost, B. M. On Inventing Reactions for Atom Economy. *Acc. Chem. Res.* **2002**, *35*, 695–705.
- (10) For selected reviews, see: (a) Zhu, C.; Ang, N. W. J.; Meyer, T. H.; Qiu, Y.; Ackermann, L. Organic Electrochemistry: Molecular Syntheses with Potential. *ACS Cent. Sci.* **2021**, *7*, 415–431. (b) Novaes, L. F. T.; Liu, J.; Shen, Y.; Lu, L.; Meinhardt, J. M.; Lin, S. Electrocatalysis as an enabling technology for organic synthesis. *Chem. Soc. Rev.* **2021**, *50*, 7941–8002.
- (11) (a) Dolson, M. G.; Swenton, J. S. Product and mechanistic studies of the anodic oxidation of methoxylated naphthalenes. The EECrCp mechanism. *J. Am. Chem. Soc.* **1981**, *103*, 2361–2371. (b) Rodríguez, S.; Wipf, P. Oxidative Spiroacetalizations and Spirolactonizations of Arenes. *Synthesis* **2004**, 2767–2783.
- (12) Carreño, M. C.; Ribagorda, M. Anodic Oxidation of N-Protected 4-Methoxy Anilines: Improved Synthesis of Quinone Imine Acetals. *J. Org. Chem.* **2000**, *65*, 1231–1234.
- (13) (a) Sumi, T.; Saitoh, T.; Natsui, K.; Yamamoto, T.; Atobe, M.; Einaga, Y.; Nishiyama, S. Anodic Oxidation on a Boron-Doped Diamond Electrode Mediated by Methoxy Radicals. *Angew. Chem., Int. Ed.* **2012**, *51*, 5443–5446. (b) Yajima, S.; Saitoh, T.; Kawa, K.; Nakamura, K.; Nagase, H.; Einaga, Y.; Nishiyama, S. Asymmetric induction in cyclohexadienones carrying α-d-glucopyranosyl moiety. *Tetrahedron* **2016**, *72*, 8428–8345.
- (14) Kirste, A.; Elsler, B.; Schnakenburg, G.; Waldvogel, S. R. Efficient Anodic and Direct Phenol-Arene C,C Cross-Coupling: The Benign Role of Water or Methanol. *J. Am. Chem. Soc.* **2012**, *134*, 3571–3576.
- (15) For an electrophotocatalytic method to generate methoxy radicals using CeCl₃ and n-Bu₄NCl, see Jiang, J.; Xu, K.; Zeng, C. Electrophotocatalytic Si–H Activation Governed by Polarity-Matching Effects. *CCS Chem.* **2021**, *3*, 1911–1920.
- (16) For a spiroketal formation, which is proposed to involve anodic oxidation of alkoxides to form alkoxy radicals, see Markó, I. E. Electrochemical oxidative cyclisation of ω-hydroxy-tetrahydropyrans to spiroketals. *Tetrahedron Lett.* **2000**, *41*, 4383–4387.
- (17) For an investigation of the favoured decomposition pathway of the radical anion generated upon single electron reduction of N-ethoxyphthalimide, see Syroeshkin, M. A.; Krylov, I. B.; Hughes, A. M.; Alabugin, I. V.; Nasybullina, D. V.; Sharipov, M. Y.; Gulyai, V. P.; Terent'ev, A. O. Electrochemical behavior of N-oxophthalimides: Cascades initiating self-sustaining catalytic reductive N—O bond cleavage. *J. Phys. Org. Chem.* **2017**, *30*, e3744.
- (18) (a) Kapustina, N. I.; Sokova, L. L.; Nikishin, H. I. Electrochemical oxidative ring opening of 1-methylcyclobutanol. *Russ. Chem. Bull.* **1996**, *45*, 1246–1248. (b) Allen, B. D. W.; Hareram, M. D.; Seastram, A. C.; McBride, T.; Wirth, T.; Browne, D. L.; Morrill, L. C. Manganese-Catalyzed Electrochemical Deconstructive Chlorination of Cycloalkanol via Alkoxy Radicals. *Org. Lett.* **2019**, *21*, 9241–9246.
- (19) For a regioselective electrochemical cyclobutanol ring expansion to 1-tetralones, see Petti, A.; Natho, P.; Lam, K.; Parsons, P. J. Regioselective Electrochemical Cyclobutanol Ring Expansion to 1-Tetralones. *Eur. J. Org. Chem.* **2021**, 2021, 854–858.
- (20) For selected reviews, see: (a) Weinberg, D. R.; Gagliardi, C. J.; Hull, J. F.; Murphy, C. F.; Kent, C. A.; Westlank, B. C.; Paul, A.; Ess, D. H.; McCafferty, D. G.; Meyer, T. J. Proton-Coupled Electron Transfer. *Chem. Rev.* **2012**, *112*, 4016–4093. (b) Gentry, E. C.; Knowles, R. R. Synthetic Applications of Proton-Coupled Electron Transfer. *Acc. Chem. Res.* **2016**, *49*, 1546–1556.
- (21) Baciocchi, E.; Bietti, M.; Lanzalunga, O. Mechanistic Aspects of β-Bond-Cleavage Reactions of Aromatic Radical Cations. *Acc. Chem. Res.* **2000**, *33*, 243–251.
- (22) For selected examples, see: (a) Yayla, H. G.; Wang, H.; Tarantino, K. T.; Orbe, H. S.; Knowles, R. R. Catalytic Ring-Opening of Cyclic Alcohols Enabled by PCET Activation of Strong O–H Bonds. *J. Am. Chem. Soc.* **2016**, *138*, 10794–10797. (b) Wu, X.; Wang, M.; Huan, L.; Wang, D.; Wang, J.; Zhu, C. Tertiary-Alcohol-Directed Functionalization of Remote C(sp³)–H Bonds by Sequential Hydrogen Atom and Heteroaryl Migrations. *Angew. Chem., Int. Ed.* **2018**, *57*, 1640–1644. (c) Huang, L.; Ji, T.; Rueping, M. Remote Nickel-Catalyzed Cross-Coupling Arylation via Proton-Coupled Electron Transfer-Enabled C–C Bond Cleavage. *J. Am. Chem. Soc.* **2020**, *142*, 3532–3539.
- (23) Murray, P. R. D.; Cox, J. H.; Chiappini, N. D.; Roos, C. B.; McLoughlin, E. A.; Hejna, B. G.; Nguyen, S. T.; Ripberger, H. H.; Ganley, J. M.; Tsui, E.; Shin, N. Y.; Koronkiewicz, B.; Qiu, G.; Knowles, R. R. Photochemical and Electrochemical Applications of Proton-Coupled Electron Transfer in Organic Synthesis. *Chem. Rev.* **2022**, *122*, 2017–2291.
- (24) Yamamoto, K.; Toguchi, H.; Kuriyama, M.; Watanabe, S.; Iwasaki, F.; Onomura, O. Electrophotocatalytic Ring-Opening

Bromination of tert-Cycloalkanols. *J. Org. Chem.* **2021**, *86*, 16177–16186.

(25) See the [Supporting Information](#) for full experimental details.

(26) (a) Steckhan, E. Indirect Electroorganic Syntheses—A Modern Chapter of Organic Electrochemistry [New Synthetic Methods (59)]. *Angew. Chem., Int. Ed.* **1986**, *25*, 683–701. (b) Wu, X.; Davis, A. P.; Lambert, P. C.; Steffen, L. K.; Toy, O.; Fry, A. J. Substituent effects on the redox properties and structure of substituted triphenylamines. An experimental and computational study. *Tetrahedron* **2009**, *65*, 2408–2414.

(27) Zhang, N.-t.; Zeng, C.-c.; Lam, C. M.; Gbur, R. K.; Little, R. D. Triarylimidazole Redox Catalysts: Electrochemical Analysis and Empirical Correlations. *J. Org. Chem.* **2013**, *78*, 2104–2110.

(28) For selected examples of deconstructive halogenation of cycloalkanols *via* alternative approaches, see: (a) Zhao, H.; Fan, X.; Yu, J.; Zhu, C. Silver-Catalyzed Ring-Opening Strategy for the Synthesis of β - and γ -Fluorinated Ketones. *J. Am. Chem. Soc.* **2015**, *137*, 3490–3492. (b) Fan, X.; Zhao, H.; Yu, J.; Bao, X.; Zhu, C. Regiospecific synthesis of distally chlorinated ketones via C–C bond cleavage of cycloalkanols. *Org. Chem. Front.* **2016**, *3*, 227–232.

(29) Sandford, C.; Edwards, M. A.; Klunder, K. J.; Hickey, D. P.; Li, M.; Barman, K.; Sigman, M. S.; White, H. S.; Minter, S. D. A synthetic chemist's guide to electroanalytical tools for studying reaction mechanisms. *Chem. Sci.* **2019**, *10*, 6404–6422.

(30) Hioe, J.; Zipse, H. Radical stability and its role in synthesis and catalysis. *Org. Biomol. Chem.* **2010**, *8*, 3609–3617.

(31) Jones, A. C.; Nicholson, W. I.; Leitch, J. A.; Browne, D. L. A Ball-Milling-Enabled Cross-Electrophile Coupling. *Org. Lett.* **2021**, *23*, 6337–6341.

(32) Green, R. A.; Brown, R. C. D.; Pletcher, D.; Harji, B. A Microflow Electrolysis Cell for Laboratory Synthesis on the Multigram Scale. *Org. Process Res. Dev.* **2015**, *19*, 1424–1427.

(33) Morris, W. D.; Mayer, J. M. Separating Proton and Electron Transfer Effects in Three-Component Concerted Proton-Coupled Electron Transfer Reactions. *J. Am. Chem. Soc.* **2017**, *139*, 10312–10319.

Recommended by ACS

Scalable Aerobic Oxidation of Alcohols Using Catalytic DDQ/HNO₃

Tania Katsina, Alexandre Jean, *et al.*

MARCH 12, 2020
ORGANIC PROCESS RESEARCH & DEVELOPMENT

READ 

Mild Condition for the Deoxygenation of α -Heteroaryl-Substituted Methanol Derivatives

Na Meng, Joseph L. Duffy, *et al.*

MARCH 30, 2021
THE JOURNAL OF ORGANIC CHEMISTRY

READ 

Selective Aerobic Oxidation of Csp³-H Bonds Catalyzed by Yeast-Derived Nitrogen, Phosphorus, and Oxygen Codoped Carbon Materials

Zhao-Yang Ju, Lan-Hui Ren, *et al.*

MARCH 07, 2022
THE JOURNAL OF ORGANIC CHEMISTRY

READ 

Selective Chloride-Mediated Neat Ethanol Oxidation to 1,1-Diethoxyethane via an Electrochemically Generated Ethyl Hypochlorite Intermediate

Siqi Li and Bart M. Bartlett

SEPTEMBER 23, 2021
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Get More Suggestions >