

Strong laws of large numbers for a growth-fragmentation process with bounded cell sizes

Emma Horton* Alexander R. Watson†

15th June 2022

Growth-fragmentation processes model systems of cells that grow continuously over time and then fragment into smaller pieces. Typically, on average, the number of cells in the system exhibits asynchronous exponential growth and, upon compensating for this, the distribution of cell sizes converges to an asymptotic profile. However, the long-term *stochastic* behaviour of the system is more delicate, and its almost sure asymptotics have been so far largely unexplored. In this article, we study a growth-fragmentation process whose cell sizes are bounded above, and prove for the first time the existence of regimes with differing almost sure long-term behaviour.

Keywords. Growth-fragmentation, law of large numbers, asynchronous exponential growth, cell division, ergodic theorem, spectral radius, spectral gap, intrinsic martingale, spectrally negative Lévy process, dividend process, skeleton decomposition.

2010 Mathematics Subject Classification. 60J80, 37A30, 47D06, 35Q92.

1. Introduction

Growth-fragmentation refers to a collection of mathematical models in which objects – classically, biological cells – slowly gather mass over time, and fragment suddenly into multiple, smaller offspring. In recent years, there has been a growing interest in probabilistic models, in the form of growth-fragmentation processes. These have been developed in the framework of piecewise-deterministic Markov processes by several authors [4, 20, 13], and in a very general form by Bertoin [10].

In this article, we study a particularly simple growth-fragmentation process where the long-term behaviour can be described explicitly. In this process, the cells grow

*INRIA, Bordeaux Research Centre, Talence, France. emma.horton@inria.fr

†Department of Statistical Science, University College London, UK. alexander.watson@ucl.ac.uk

exponentially (that is, the growth rate is linear) until they reach a certain mass, at which point growth stops abruptly. Fragmentation occurs at a constant rate regardless of the cell mass, and the relative masses of offspring are similarly independent of the parent mass. We are interested in understanding the long-term behaviour of these systems, and in particular whether, with appropriate rescaling, they can converge to some ‘asymptotic profile’, describing the masses of cells in the limit.

This process can be regarded as a basic model for cells with a fixed bound on their mass, and this property is attractive in terms of applications [15, 5], either to account in an indirect way for resource competition, or to model a biologically determined bound on how large cells seek to become. Alternatively, from a mathematical perspective, we can see this process as being part of a family of perturbations of a fundamental *homogeneous* growth-fragmentation process, in which all rates are independent of mass.

Work in this field often begins with the *growth-fragmentation equation*, which in general form is given by

$$\partial_t u_t(x) + \partial_x(\tau(x)u_t(x)) + (B(x) + D(x))u_t(x) = \int_{(x,\infty)} k(y,x)u_t(y) dy,$$

with some initial condition $u_0 = g_0$. The equation can be usefully rewritten in its weak form, in which we have a collection of measures $(\mu_t)_{t \geq 0}$, started from $\mu_0 = \delta_x$, satisfying

$$\partial_t \langle f, \mu_t \rangle = \langle \mathcal{A}f, \mu_t \rangle, \quad \mathcal{A}f(x) = \tau(x)f'(x) + \int_{(0,x)} f(y)k(x,dy) - (B(x) + D(x))f(x), \quad (1)$$

where $\langle f, \mu \rangle = \int f d\mu$, τ is the growth rate, $B(x) = \int_{(0,y)} \frac{y}{x} k(x,dy)$ is the fragmentation rate, D is the killing rate and k is a kernel describing the masses of offspring cells given the mass of the parent.

The specific case we will consider is when $\tau(x) = ax \mathbb{1}_{\{x < c\}}$, meaning that the constant $c > 0$ is the maximum possible cell mass, with cells growing exponentially until reaching mass c . We take the other coefficients to be independent of cell mass: $B(x) = B$ and $D(x) = \mathbf{k}$ are constant, and we express the offspring mass distribution in terms of a measure ρ on $(0, 1)$, such that $\int f(y)k(x,dy) = \int f(xy)\rho(dy)$. The measure ρ describes the *relative* masses of offspring cells, which are independent of the mass of the parent, and it satisfies $\int_{(0,1)} y \rho(dy) = B$. Since we only consider binary branching in this article, it will always hold that ρ is invariant under the transformation $y \mapsto 1 - y$ and that $\rho((0, 1)) = 2B$. The operator \mathcal{A} is then given by

$$\mathcal{A}f(x) = axf'(x) + \int_{(0,1)} f(xy)\rho(dy) - (B + \mathbf{k})f(x), \quad (2)$$

where the domain $\mathcal{D}(\mathcal{A})$ satisfies

$$\mathcal{D}(\mathcal{A}) \supset \{f \in C_c^1(0, c] : f'(c) = 0\}.$$

We will shortly see that the measures μ_t can be identified as the average of a stochastic system of cell masses, which forms our main object of study. Thus, we refer to $(\mu_t : t \geq 0)$

as the *mean measures*. One result of interest, which has received much attention in the literature of growth-fragmentation processes and, more generally, branching processes, is the Perron–Frobenius behaviour of such systems, which states that for some $\lambda \in \mathbb{R}$, $e^{-\lambda t} \mu_t$ converges, in some sense, to a steady state $h(x)\nu$. We call ν the asymptotic profile. The homogeneous case, where rather $\tau(x) = ax$ for all $x > 0$, was studied in [12], where it was observed that no such simple asymptotic profile exists. We refer the reader to Corollary 3.4 in that work, where it is shown that the appropriate scaling is instead $t^{-1/2}e^{-\lambda t} \mu_t$.

However, Cavalli [18] has shown that a simple change in the drift, making it piecewise constant, can yield convergence of the mean measures to an asymptotic profile, with scaling exponential in time. In this work, we use this approach to take the results a step further: not only do we obtain explicit expressions and show that the mean measures converge, but we are also able to prove this behaviour for the stochastic system of cell masses, giving a strong law of large numbers.

To describe this stochastic system, let us start by introducing a stochastic process X started at x , which can be thought of as the trajectory of a single cell with initial mass x . Write \mathbb{P}_x for the probability measure associated with this initial mass. First, let ξ be the process

$$\xi_t = y + at + \sum_{i=1}^{\mathcal{N}_t} J_i, \quad t < \zeta,$$

where $y = \log x$, $a > 0$ represents the (exponential) growth rate, \mathcal{N} is a Poisson process with rate B , and $(J_i)_{i \geq 1}$ are independent, identically distributed jump sizes whose support is contained in $[-\log 2, 0)$. Let ζ be an exponential random variable with rate k , and let ξ be sent to some cemetery state at time ζ . Define $\Pi(dx) = B\mathbb{P}(J_1 \in dx)$. The process ξ is a Lévy process with drift a , no Gaussian part, Lévy measure Π and killing rate k . Let $\bar{\xi}_t = \sup\{\xi_s : s \leq t\}$, take $b = \log c$ and define

$$\xi_t^b = \xi_t - (\bar{\xi}_t - b) \vee 0,$$

which is the process ξ reflected above at b .

Our single-cell trajectory is given by $X_t = \exp(\xi_t^b)$, the exponential of a reflected Lévy process, starting at $X_0 = x$. This trajectory increases with exponential rate a until reaching $c = e^b$, has negative jumps and is killed at rate k . We understand that, if ξ_t is in the cemetery state, then the same applies to X_t . Each jump of size $\Delta X_t < 0$ represents a splitting event, and is associated with the birth of another cell (i.e. a binary fission event), with initial mass $-\Delta X_t$ and where the trajectory of its mass is given by a copy of X which is conditionally independent given its starting value. By introducing these new cells at every jump of X , and iterating this process with each successive set of children, we build up the growth-fragmentation process. In terms of Bertoin’s Markovian growth-fragmentation processes [10] this is a growth-fragmentation with ‘cell process’ X .

In order to describe the masses of each cell, we define $Z_u(t)$ to be the mass of the cell with label u at time t ; the set of labels u will be described later. The *growth-fragmentation process*, $\mathbf{Z}(t) = \sum_u \delta_{Z_u(t)}$, is a point measure-valued process describing the masses of all cells alive at each time-point.

Crucially, the relation between $\mathbf{Z} = (\mathbf{Z}(t))_{t \geq 0}$ and $(\mu_t)_{t \geq 0}$ is given by

$$\langle f, \mu_t \rangle = \mathbb{E}_x[\langle f, \mathbf{Z}(t) \rangle], \quad (3)$$

for bounded measurable f , with $\rho = \Pi \circ g^{-1} + \Pi \circ \bar{g}^{-1}$, where $g(x) = e^x$ and $\bar{g}(x) = 1 - e^x$. In other words, as indicated earlier, μ_t represents the average linear behaviour of the growth-fragmentation process.

There is a vast literature on the long-term behaviour of growth-fragmentation equations and processes (see [12, 41, 24, 8] and [16, 22, 10], respectively, to name just a few references.) From the analytical perspective, the conventional approach is to find $\lambda \in \mathbb{R}$, a positive function h and a measure ν , such that

$$\nu \mathcal{A} = \lambda \nu \quad \text{and} \quad \mathcal{A}h = \lambda h, \quad (4)$$

where $\langle \nu, h \rangle = 1$, using the Krein-Rutman theorem [41, 7, 42] or the Lumer-Phillips theorem [8]. Generalised relative entropy methods [24, 40] can be used to show the long-term asynchronous exponential growth

$$\langle f, \mu_t \rangle \sim e^{\lambda t} \langle f, \nu \rangle h(x), \quad (5)$$

when $\mu_0 = \delta_x$. On the other hand, from a probabilistic perspective, it is often more convenient to work directly with the expectations of the growth-fragmentation process \mathbf{Z} to show that the process satisfies

$$\mathbb{E}_x[\langle f, \mathbf{Z}(t) \rangle] \sim e^{\lambda t} \langle \nu, f \rangle h(x), \quad (6)$$

which, thanks to (3), agrees with (5). One approach used in [12, 18] is to consider a single tagged particle via a Feynman–Kac formula, and to analyse the Laplace transform of the hitting time of points of this particle. Other recent approaches make use of Harris-style theorems for nonconservative semigroups [7], and quasi-stationary methods [19].

Finally, another approach, [5, 17], which can be seen as a hybrid of the functional analytic and stochastic perspectives, involves identifying h through some variation of the analytic approaches above, using it to derive a stochastic semigroup, and then checking the ergodic theorem to obtain a probability measure, which is a simple transformation of the eigenmeasure ν , leading to the desired long-term behaviour. This idea is the starting point for our work.

The main aim of this article is to describe the long-term *stochastic* behaviour of the growth-fragmentation process. To this end, we will prove two theorems that can be seen as stochastic analogues of the deterministic asymptotic (6), and which, indeed, imply it. The first of these theorems is the following *strong law of large numbers*.

Theorem 1. *Assume that*

$$B > \mathbf{k}, \quad a > \int_0^1 (-\log v) \rho(dv) \quad \text{and} \quad \int_0^1 v^{-r} \rho(dv) < \infty \quad \text{for some } r > 0.$$

Define $\lambda = B - \mathbf{k}$. Then, there exists a probability measure ν and a random variable M_∞ such that for all $x \in (0, \infty)$ and continuous, bounded f ,

$$e^{-\lambda t} \langle f, \mathbf{Z}(t) \rangle \rightarrow \langle f, \nu \rangle M_\infty,$$

\mathbb{P}_x -almost surely and in $L^1(\mathbb{P}_x)$.

In fact, M_∞ is the limit of the martingale $M_t = e^{-(B-\mathbf{k})t} \langle 1, \mathbf{Z}(t) \rangle$, and it has expectation constant in the starting mass of the process: $\mathbb{E}_x[M_\infty] = 1$. The $L^1(\mathbb{P}_x)$ convergence implies in particular (6) with $h = 1$, but in fact, we derive (4) and (6), including a version of (6) with exponential rate, under more general assumptions, as the forthcoming Propositions 12 and 13. These are fundamental to our method of proving Theorem 1.

An equivalent version of this theorem has been shown to hold for a wide variety of branching processes, including branching diffusions [29], superprocesses [28] and, more recently, a general class of branching Markov processes with non-local branching [33]. However, in these settings, the dependence of (6) on the initial condition is well-understood, which is not the case here. A form of this theorem with $L^1(\mathbb{P}_x)$ convergence was shown in [14], for a class of processes with size-dependent parameters and without killing. This is the first time that the almost sure mode of convergence has been shown for a growth-fragmentation process. In this article, we adapt the approaches used in [33] and [29] in order to prove this strong law, which requires precise control of the first and second order convergence of (6).

The second theorem provides a similar result in the transient case. To aid comparison with the literature, it will be useful to introduce a function κ , which in the context of homogeneous growth-fragmentations is called the ‘cumulant’ and is defined as:

$$\kappa(q) = aq + \int_{(0,1)} x^q \rho(dx) - (B + \mathbf{k}).$$

The function κ is convex and smooth. We define q_0 to be the unique solution of the equation $\kappa'(q_0) = 0$, or equivalently, $q_0 = \arg \inf \kappa$.

Theorem 2. *Assume that*

$$B > \mathbf{k}, \quad a < \int_{(0,1)} (-\log v) \rho(dv),$$

which implies that $q_0 > 0$. Define $\lambda = \kappa(q_0) = \inf \kappa$, and assume further that $\lambda > 0$. Let $f: (0, c] \rightarrow \mathbb{R}$ be continuous with $f(x) = O(x^{q_0})$ as $x \rightarrow 0$. Then,

$$e^{-\lambda t} \langle f, \mathbf{Z}(t) \rangle \rightarrow 0,$$

\mathbb{P}_x -almost surely and in $L^1(\mathbb{P}_x)$.

This result can be interestingly compared with [22, Theorem 2.3], which considers a *homogeneous* growth-fragmentation \mathbf{Z}° and takes $f(x) = f_q(x) = x^q$. Under the assumptions of the above theorem, part of Dadoun’s result is that $e^{-\kappa(q)t} \langle f_q, \mathbf{Z}^\circ(t) \rangle \rightarrow 0$ almost surely, provided that $q \geq \bar{q}$, where $\bar{q} > q_0$ is the unique solution of $\bar{q}\kappa(\bar{q}) = \kappa'(\bar{q})$.

The fact that $\bar{q} > q_0$ says, in a quantitative way, that the cell masses in the growth-fragmentation process with reflection decay to zero faster than without reflection. To our knowledge, this type of convergence to zero has not been shown previously for non-homogeneous processes, and one interesting avenue for future research would be to study the speed of cell masses in more detail, as done by Dadoun for homogeneous processes in [22, Corollary 2.4].

We would also like to note several other possible directions for future work. Here, we have taken the killing rate to be constant, but it may be possible to relax this restriction while remaining in a tractable class of processes. In particular, it would be interesting to consider killing cells, either instantaneously or at exponential rate, when their mass falls below a certain threshold, to mimic mechanisms for cell death in biological systems. An alternative direction would be to take a more general Lévy process as the underlying trajectory of the cell masses; for instance, it should be possible to incorporate Gaussian fluctuations into this model with little impact on our results. Finally, as mentioned above, we have assumed here that cell divisions are binary, but this is far from necessary. Indeed, provided that the number of cells produced at division is not too large (an ‘ $L \log L$ ’ condition will suffice), arbitrary division and repartition of mass can be incorporated into this model; one mathematical approach to this is described in [14, p. 8]. This could prove fruitful in biological models for tumour growth, since it is known that cancer cells can split into more than two daughter cells. It would also allow one to study a phenomenon known as aneuploidy, where a cell contains the wrong number of chromosomes due to non-binary fission (see [44] and references therein).

The rest of this paper is organised as follows. In section 2 we will analyse the expectation semigroup associated with the process \mathbf{Z} , as well as the strongly continuous semigroup associated with the operator \mathcal{A} . We will show that these two semigroups are equivalent, in an appropriate sense, for a certain class of functions. In section 3 we will formally define the growth-fragmentation process and consider the basic behaviour of the number of cells in the system. In section 4, we will make use of a spine method to distinguish the regimes of long-term behaviour. Finally, sections 5 and 6 are devoted to the proofs of Theorems 1 and 2, respectively.

2. The growth-fragmentation semigroup

One way to describe properties of the growth-fragmentation process is via its expectation semigroup. One can characterise the average behaviour of particles, under the action of suitable test functions, as the unique solution to an evolution equation. On the other hand, from an analytical perspective, it is common to use the generator defined in (2) to define a strongly continuous semigroup, usually written $(e^{t\mathcal{A}})_{t \geq 0}$, which uniquely solves the growth-fragmentation equation (1).

In this section, we will explore both perspectives and, in order to consolidate them, we will show that the two are equivalent for a certain class of test functions. Hence, roughly speaking, both the analytical and probabilistic semigroups characterise the average behaviour of such growth-fragmentation systems, and can therefore both be used to analyse their long-term behaviour.

We start by considering the expectation semigroup associated with the growth-fragmentation process. First recall the underlying reflected spectrally negative Lévy process, ξ^b , obtained by taking the process with Laplace exponent

$$\psi_\xi(q) = -k + aq - \int_{-\infty}^0 (1 - e^{qx}) \Pi(dx),$$

satisfying $\mathbb{E}_0[e^{q\xi_t}] = e^{t\psi_\xi(q)}$, and reflecting it from above at the level b .

Further, recall the growth-fragmentation kernel ρ defined on $(0, 1)$ by $\rho = \Pi \circ g^{-1} + \Pi \circ \bar{g}^{-1}$, where $g(x) = e^x$ and $\bar{g}(x) = 1 - e^x$.

As before, we write $\mathbf{Z} = (\mathbf{Z}(t), t \geq 0)$ for the Markovian growth-fragmentation associated with e^{ξ^b} , and let $c = e^b > 0$, which is the reflection level for cell sizes in \mathbf{Z} .

Define $\Psi_t f(x) = \mathbb{E}_x[\langle f, \mathbf{Z}(t) \rangle]$. In our first proposition, we derive an evolution equation for which Ψ_t is the unique solution. We regard Ψ as a semigroup on $L^\infty((0, c])$, the space of bounded measurable functions on $(0, c]$.

Proposition 3. *The semigroup Ψ satisfies the evolution equation*

$$\begin{aligned} \Psi_t f(x) &= f(xe^{a \cdot t \wedge T_c(x)}) - \mathbf{k} \int_0^t ds \Psi_{t-s} f(xe^{a \cdot s \wedge T_c(x)}) \\ &\quad + \int_0^t ds \int_0^1 \rho(dv) [\Psi_{t-s} f(xe^{a \cdot s \wedge T_c(x)v}) - v \Psi_{t-s} f(xe^{a \cdot s \wedge T_c(x)})], \end{aligned} \quad (7)$$

where $T_c(x) = \frac{1}{a} \log(\frac{c}{x})$ is the first time the deterministic path $t \mapsto xe^{at}$ reaches c . Moreover, this evolution equation uniquely determines Ψ as a semigroup on $L^\infty((0, c])$.

Proof. Let τ_1 denote the first time that the initial particle either dies or fragments. By considering the two cases $t < \tau_1$ and $t \geq \tau_1$ we have

$$\begin{aligned} \Psi_t f(x) &= f(xe^{a \cdot t \wedge T_c(x)}) e^{-(B+\mathbf{k})t} \\ &\quad + (B + \mathbf{k}) \int_0^t ds \frac{B}{B + \mathbf{k}} e^{-(B+\mathbf{k})s} \int_{-\infty}^0 \frac{\Pi(du)}{B} \\ &\quad \quad \times \left\{ \Psi_{t-s} f(xe^{a \cdot s \wedge T_c(x)} e^u) + \Psi_{t-s} f(xe^{a \cdot s \wedge T_c(x)} (1 - e^u)) \right\} \\ &= f(xe^{a \cdot t \wedge T_c(x)}) e^{-(B+\mathbf{k})t} \\ &\quad + \int_0^t ds e^{-(B+\mathbf{k})s} \int_{-\infty}^0 \Pi(du) \left\{ \Psi_{t-s} f(xe^{a \cdot s \wedge T_c(x)} e^u) + \Psi_{t-s} f(xe^{a \cdot s \wedge T_c(x)} (1 - e^u)) \right\}. \end{aligned}$$

Making a change of variables $v = e^u$ yields

$$\begin{aligned} \Psi_t f(x) &= f(xe^{a \cdot t \wedge T_c(x)}) e^{-(B+\mathbf{k})t} \\ &\quad + \int_0^t ds e^{-(B+\mathbf{k})s} \int_0^1 \Pi \circ g^{-1}(dv) \left\{ \Psi_{t-s} f(xe^{a \cdot s \wedge T_c(x)} v) + \Psi_{t-s} f(xe^{a \cdot s \wedge T_c(x)} (1 - v)) \right\}. \end{aligned}$$

Making another change of variables, $v \mapsto 1 - v$, for the second integrand in the final line above, and recalling that $\rho = \Pi \circ g^{-1} + \Pi \circ \bar{g}^{-1}$, we have

$$\begin{aligned} \Psi_t f(x) &= f(xe^{a \cdot t \wedge T_c(x)}) e^{-(B+\mathbf{k})t} \\ &\quad + \int_0^t ds e^{-(B+\mathbf{k})s} \int_0^1 \rho(dv) \Psi_{t-s} f(xe^{a \cdot s \wedge T_c(x)} v). \end{aligned}$$

Applying Dynkin's identity [27, Lemma 1.2, Chapter 4, Part I] and using the fact that that $\int_0^1 v \rho(dv) = B$ shows that $\Psi_t f(x)$ does indeed solve (7).

For uniqueness, suppose Ψ_t^1 and Ψ_t^2 are both solutions to (7) and let $\tilde{\Psi}$ denote their difference. Then, since B is finite, it is straightforward to show that there exists a constant $C > 0$ such that

$$|\tilde{\Psi}_t f(x)| \leq C \int_0^t \sup_{x \in (0, c]} |\tilde{\Psi}_{t-s} f(x)| ds.$$

Gronwall's inequality then yields the result. \square

With this result we have uniquely characterised the average (linear) behaviour of \mathbf{Z} using the evolution equation (7). Part of our goal in this work is to describe our probabilistic results (Theorem 1 and (6)) about the growth-fragmentation semigroup in terms of the spectral theory of the operator \mathcal{A} , defined in (2). However, if we were to define \mathcal{A} as the generator of a semigroup with the same action as Ψ (on the space of continuous functions vanishing at 0, say), then its domain would not be large enough to contain the eigenfunction we identify. To remedy this, we now show that \mathcal{A} (or a suitable extension) can be regarded as the generator of a strongly continuous semigroup on $L^2((0, c])$, which solves the growth-fragmentation equation (1) and agrees with the semigroup Ψ above on their common domain of definition.

In general, questions of existence and uniqueness of a semigroup with generator \mathcal{A} are not straightforward; we refer to [23, 38, 37] for some examples of classical and recent approaches. Our situation is fairly simple since the fragmentation rate is bounded, but nonetheless it does not seem to be easy to derive from existing results, so we develop the proof below.

We first focus our attention on the growth (i.e., transport) part of the operator. We define

$$\mathcal{G}f(x) = axf'(x) - (B + \mathbf{k})f(x), \quad f \in D_0,$$

where D_0 is the subspace of $L^2((0, c])$ consisting of functions $f: (0, c] \rightarrow \mathbb{R}$ such that

- (i) f is smooth with compact support, and
- (ii) $\lim_{x \rightarrow c} f'(x) = 0$.

Then, we have the following proposition.

Proposition 4. *The operator $\mathcal{G}: D_0 \rightarrow L^2$ is closable and its closure generates a strongly continuous semigroup on $L^2((0, c])$.*

Proof. The majority of the proof follows the method in [45] so we will only give an outline of the proof here. We start by considering the operators

$$U_t f(x) = e^{-(B+\mathbf{k})t} f(xe^{at}), \quad f \in L^2((0, c]), t \geq 0,$$

which we define as $e^{-(B+\mathbf{k})t} f(c)$ whenever $t > 0$ and $xe^{at} \geq c$.

First note that for each $t \geq 0$, U_t is bounded. Also, for any continuous function $f: (0, c] \rightarrow \mathbb{R}$ with compact support, we have

$$\|U_t f - f\|_2 \rightarrow 0, \quad \text{as } t \rightarrow 0.$$

Since the space of compactly supported continuous functions is dense in L^2 , these observations imply $(U_t)_{t \geq 0}$ is strongly continuous. Hence, its infinitesimal generator is closed and densely defined.

Now let $f \in L^2((0, c])$ be smooth with compact support, and for $h > 0$, set

$$\varphi_h(x) := \int_0^h U_s f(x) ds.$$

Due to [26, p.620], φ_h lies in the domain of the infinitesimal generator of $(U_t)_{t \geq 0}$. Denote by \mathcal{G}_1 the restriction of this generator to functions φ_h constructed as above. Then, again, by [26, p.620]

$$\mathcal{G}_1 \varphi_h = U_h f - f,$$

and the infinitesimal generator of $(U_t)_{t \geq 0}$ is $\bar{\mathcal{G}}_1$.

Our next step is to show \mathcal{G} is closable and its closure equals $\bar{\mathcal{G}}_1$. To show this, we first show that φ_h lies in D_0 . First note that from the definition of φ_h , it is also smooth with compact support. Next note that

$$\varphi_h(xe^{at}) = \int_t^{t+h} e^{-(B+\mathbf{k})(u-t)} f(xe^{au}) du. \quad (8)$$

Hence,

$$\begin{aligned} \frac{\partial}{\partial t} \varphi_h(xe^{at}) &= e^{-(B+\mathbf{k})h} f(xe^{a(t+h)}) + (B+\mathbf{k}) \int_0^{t+h} e^{-(B+\mathbf{k})(u-t)} f(xe^{at}) du \\ &\quad - f(xe^{at}) - (B+\mathbf{k}) \int_0^t e^{-(B+\mathbf{k})(u-t)} f(xe^{at}) du. \end{aligned}$$

Letting $t \rightarrow 0$, we have

$$\lim_{t \rightarrow 0} \frac{\partial}{\partial t} \varphi_h(xe^{at}) = U_h f(x) - f(x) + (B+\mathbf{k})\varphi_h(x),$$

which, in turn, implies that

$$ax\varphi_h'(x) - (B+\mathbf{k})\varphi_h(x) = U_h f(x) - f(x). \quad (9)$$

Now, from the definitions of φ_h and U_h , it is not too difficult to show that

$$\lim_{x \rightarrow c} \varphi_h(x) = \frac{f(c)}{B+\mathbf{k}} (1 - e^{-(B+\mathbf{k})h})$$

and $\lim_{x \rightarrow c} U_h f(x) = e^{-(B+\mathbf{k})h} f(c)$. Taking the limit as $x \rightarrow c$ in (9) and using these two equalities yields $\lim_{x \rightarrow c} \varphi_h'(x) = 0$. Hence $\varphi_h \in D_0$.

The above analysis also shows that

$$\mathcal{G}_1 \varphi_h(x) = ax\varphi_h'(x) - (B+\mathbf{k})\varphi_h(x) = \mathcal{G}\varphi_h(x).$$

Since $\varphi_h \in D_0$, it follows that $\mathcal{G}_1 \subset \mathcal{G}$.

Finally, for any $\varphi \in D_0$ we have that

$$U_h\varphi - \varphi = \int_0^h U_s \mathcal{G}\varphi ds.$$

Due to [34, Theorem 10.5.2], \mathcal{G} is a restriction of $\bar{\mathcal{G}}_1$. Therefore, $\bar{\mathcal{G}} = \bar{\mathcal{G}}_1$ and so \mathcal{G} is closable and generates the semigroup $(U_t)_{t \geq 0}$. \square

Let $\mathcal{F}: D_0 \rightarrow L^2$ be defined by

$$\mathcal{F}f(x) = \int_0^1 f(xy)\rho(dy).$$

Then, we can extend the result of the previous proposition to $\mathcal{G} + \mathcal{F}$.

Corollary 5. *The conclusion of Proposition 4 holds when we replace \mathcal{G} by $\mathcal{G} + \mathcal{F}$. Moreover, if we define \mathcal{A} to be the closure of $\mathcal{G} + \mathcal{F}$, then its domain $\mathcal{D}(\mathcal{A})$ is equal to the domain of the closure of \mathcal{G} .*

Proof. First note that since \mathcal{F} it is bounded, it can be extended to an operator with domain $L^2((0, c])$. Further, recall $(U_t)_{t \geq 0}$ from the proof of Proposition 4. We will use Theorem 1 of [46] to prove this corollary, which in our context, is as follows.

Suppose the following two conditions are satisfied:

(I) $U_t D_0 \subset D_0$ and for all $f \in D_0$, the function $t \mapsto \mathcal{F}U_t f$ is continuous.

(II) There exist constants $\alpha > 0$, $\gamma \in [0, 1)$ such that for all $f \in D_0$

$$\int_0^\alpha \|\mathcal{F}U_t f\| dt \leq \gamma \|f\|.$$

Then there exists a unique strongly continuous semigroup $(W_t)_{t \geq 0}$ on $L^2((0, c])$ such that

$$W_t f = U_t f + \int_0^t W_{t-s} \mathcal{F}U_s f ds \quad f \in D_0, \quad t \geq 0.$$

Moreover, the generator of $(W_t)_{t \geq 0}$ is the closure of $(\mathcal{G} + \mathcal{F})|_{D_0}$ and its domain is equal to the domain of the closure of \mathcal{G} .

Now, (I) follows easily from the definitions of U_t , D_0 and \mathcal{F} . For (II), note that for any $\alpha > 0$,

$$\int_0^\alpha \|\mathcal{F}U_t f\| dt \leq 2B \|f\| \int_0^\alpha e^{-(B+k)t} dt = \frac{2B \|f\|}{B+k} (1 - e^{-\alpha(B+k)}).$$

Choosing $\alpha > 0$ such that $\gamma := \frac{2B}{B+k} (1 - e^{-\alpha(B+k)}) \in [0, 1)$ yields the result. Hence, the operator \mathcal{A} with domain $\mathcal{D}(\mathcal{A})$ generates a strongly continuous semigroup, $(W_t)_{t \geq 0}$ on $L^2((0, c])$. Moreover, its domain, $\mathcal{D}(\mathcal{A})$, is equal to that of the domain of the closure of \mathcal{G} . \square

Therefore, the operator \mathcal{A} is the generator of a strongly continuous semigroup $W = (W_t)_{t \geq 0}$ on $L^2((0, c])$ and has D_0 as a core. Fixing $g \in L^2((0, c])$, if we let μ_t be defined, though the Riesz representation theorem, by $\langle W_t f, g \rangle_2 = \langle f, \mu_t \rangle$ (where $\langle \cdot, \cdot \rangle_2$ represents the L^2 inner product), then $(\mu_t)_{t \geq 0}$ solves (1) with $\mu_0(dy) = g(y)dy$.

In fact, W is an extension of Ψ in the following sense.

Proposition 6. *For each $f \in L^\infty((0, c])$, $\|W_t f - \Psi_t f\|_2 = 0$.*

Proof. The proof is identical to that of [21, Theorem 8.1]. □

This provides a version of (3) in the introduction, albeit restricted to bounded functions f and initial conditions having L^2 density. This will let us consolidate the behaviour shown in (4) and (6) later on.

3. The growth-fragmentation process

We now turn our attention to the growth-fragmentation process, \mathbf{Z} . Although we have already given a construction of \mathbf{Z} in the introduction, in this section we provide another construction via the Ulam-Harris tree, as it will be convenient in later sections to refer to the particles via their labels.

To this end, we denote by $\mathcal{U} = \bigcup_{n \geq 0} \{0, 1\}^n$ the Ulam-Harris tree of finite sequences of 0s and 1s. For $u \in \{0, 1\}^n$, we interpret the length $|u| = n$ as its generation. Further, for $v \in \{0, 1\}$, we write $uv \in \{0, 1\}^{n+1}$ to denote the concatenation of u and v and interpret it as the v^{th} daughter of u . The initial individual will be written as \emptyset .

Now fix $x \in (0, c]$ and let \mathbb{P}_x denote the law of the growth-fragmentation process started from an initial particle at x . Now define the random variable ζ_\emptyset by

$$\mathbb{P}_x(\zeta_\emptyset > t) = e^{-(B+k)t}, \quad t \geq 0,$$

which we see as the minimum of the fragmentation time and death time of the initial particle. Setting $b_\emptyset = 0$ and $d_\emptyset = \zeta_\emptyset$, it follows that the trajectory of the initial particle is given by

$$Z_\emptyset(t) = x e^{a \cdot t \wedge T_c(x)}, \quad t \in [b_\emptyset, d_\emptyset),$$

where we recall that $T_c(x) = \frac{1}{a} \log(c/x)$. With probability $\frac{k}{B+k}$, ζ_\emptyset denotes a killing time of \emptyset , in which case the process stops. If not, then ζ_\emptyset denotes the first fragmentation time, and then we set $b_u = d_\emptyset$ for each daughter $u \in \{0, 1\}$ of the initial particle and randomly choose $v \in (0, 1)$ according to the probability measure $\frac{\rho(dv)}{2B}$. The masses of the daughters are then given by xv and $x(1-v)$. The lifetime of each daughter is denoted by ζ_u , which has the same distribution as ζ_\emptyset under \mathbb{P}_{xv} or $\mathbb{P}_{x(1-v)}$. We also set $d_u = b_u + \zeta_u$ and the trajectory of the u^{th} daughter is given by

$$Z_u(t) = y e^{a \cdot t \wedge T_c(y)}, \quad t \in [b_u, d_u),$$

for $y = xv, x(1-v)$. The process then continues iteratively, with each cell u and the initial mass of its daughters $u0$ and $u1$ being determined in the same way, and so on; this

gives a trajectory $(Z_u(t))_{t \in [b_u, d_u]}$ for each $u \in \mathcal{U}$. Setting \mathcal{U}_t to be the set of individuals alive at time t , i.e. $\mathcal{U}_t = \{u \in \mathcal{U} : t \in [b_u, d_u]\}$, the growth-fragmentation process at time t is given by the collection of atomic measures

$$\mathbf{Z}(t) = \sum_{u \in \mathcal{U}_t} \delta_{Z_u(t)}.$$

In the introduction and in the previous section, this process was constructed in a different way, using the ‘cell process’ ξ and adding new cells at each jump. These two approaches are consistent: if (u_1, u_2, \dots) is a sequence of cell labels with the property that, for each $n \geq 1$, $Z_{u_{n+1}}(b_{u_{n+1}})/Z_{u_n}(d_{u_n}-) \geq 1/2$, then

$$\xi_t = Z_{u_n}(t), \text{ where } b_{u_n} \leq t < d_{u_n}, \quad t \geq 0,$$

is the cell process.

Let us now briefly turn our attention to the growth of the total number of cells in \mathbf{Z} . Set $\lambda_* = B - \mathbf{k}$ and define $N_t = \langle 1, \mathbf{Z}(t) \rangle$, the number of cells in the growth-fragmentation at time t . Since the branching and killing rates are constant, N is a discrete-space, continuous-time branching process.

Recall that a branching process is subcritical, critical or supercritical according to whether $\mathbb{E}[N_t]$ is negative, zero or positive, and that in the subcritical and critical cases, the process becomes extinct (eventually $N_t = 0$) with probability 1. We have the following simple consequence of branching process theory:

Proposition 7. *N is subcritical, critical or supercritical according to whether λ_* is negative, zero or positive. In the supercritical case, the extinction probability is $\frac{\mathbf{k}}{B}$.*

Proof. This follows using classical branching process results [3, §I.5]. The probability generating function of the offspring distribution is $g(z) = \frac{\mathbf{k} + Bz^2}{B + \mathbf{k}}$, and equation $g(z) = z$ has roots 1 and $\frac{\mathbf{k}}{B}$, which implies the extinction probability in the result. \square

We end this section with a short discussion of the relationship between the various parameters of the growth-fragmentation process. For the particular model we consider in this paper, all of our results can be expressed explicitly in terms of the parameters a, \mathbf{k} and ρ , due to the fact that $\|\rho\| = 2\|\Pi\| = 2B$, where $\|\cdot\|$ indicates the total mass of a measure. We emphasise that the factor of two appears here due to the fact that two particles are produced at every fragmentation event.

In a more general model, where we allow particles to fragment into more than two pieces, the total mass of ρ is given by the fragmentation rate multiplied by the average number of offspring, and λ_* is replaced by $\|\rho\| - B - \mathbf{k}$. Moreover, the results of this paper will still hold but with a slight adjustment of the conditions; the condition $B > \mathbf{k}$ in Theorem 1 would become $\|\rho\| - B - \mathbf{k} > 0$, and an ‘ $L \log L$ ’ condition on the number of offspring would be required, which is redundant in the binary case. One key difference in this more general model is that ρ (or equivalently, the linear evolution equation) no longer uniquely determines the growth-fragmentation process. For this, one needs to work with a non-linear version of the growth-fragmentation equation, as in section 5.

4. Asymptotic regimes

In the previous section, we determined conditions for the total number of cells to either reach zero with probability 1 (the (sub)critical cases) or to survive with positive probability (the supercritical case). We now consider the long-term behaviour of cell masses, with the goal of demonstrating that these settle into an equilibrium in the long-term; this will be most important in the supercritical case, where it will be the foundation of Theorem 1, but the results of this section apply regardless of (sub- or super-)criticality.

Our main tool will be the analysis of a single *tagged cell*, whose behaviour will tell us something about the average behaviour of the process \mathbf{Z} . We can think of this as following a distinguished line of descent in the cell line, where at each cell division event we uniformly pick a daughter cell to follow.

The motion of the cell mass of this tagged cell will be determined by an exponential reflected Lévy process, much like the ‘cell process’ ξ in the introduction; however, the process itself is different, since it should represent not the locally largest cell, but a typical cell.

Introduce a Lévy process η with drift a and Lévy measure $\Pi_\eta = \rho \circ r^{-1}$, where $r(x) = \log x$. Equivalently, this Lévy measure can be expressed in terms of that of ξ by $\Pi_\eta = \Pi + \Pi \circ s^{-1}$, where $s(x) = \log(1 - e^x)$. (Hence, if Π has density π , then Π_η has density $\pi_\eta(x) = \pi(x) + \pi(\log(1 - e^x))e^x(1 - e^x)^{-1}$.) Denote by ψ_η its Laplace exponent, which can be expressed as

$$\psi_\eta(q) = aq + \int_{(-\infty, 0)} (e^{qx} - 1) \Pi_\eta(dx) = aq + \int_{(0, 1)} (v^q - 1) \rho(dv).$$

Again, as with ξ , we define η^b to be this process reflected at b , i.e., $\eta_t^b = \eta_t - (\bar{\eta}_t - b) \vee 0$.

The mass of the tagged cell is given by $Y = \exp(\eta^b)$. Our goal for the remainder of this section is to investigate the long-term behaviour of Y , identify the growth-fragmentation semigroup in terms of the semigroup of Y , and hence prove the asynchronous exponential growth (6). This will be a key building block in the proof of the strong law, Theorem 1.

In past work on growth-fragmentation equations [13, 18], a key element of the analysis was an auxiliary function defined in terms of return times of Y . In our context, we may define this as follows. Let $S(y) = \inf\{t > 0 : Y_t \neq y\}$, and let $H(y) = \inf\{t > S(y) : Y_t = y\}$ be the first return time to y . Define

$$L_{x,y}(q) = \mathbb{E}_x[e^{-(q-\lambda_*)H(y)}; H(y) < \infty].$$

Let $q_* = \inf\{q \in \mathbb{R} : L_{c,c}(q) < \infty\}$ and $\lambda = \inf\{q \in \mathbb{R} : L_{c,c}(q) < 1\}$. The quantity λ is the conjectural exponential rate in (6), and in previous works, the behaviour of L was used in order to deduce asymptotic results for the growth-fragmentation semigroup. Our situation is actually a little simpler than [13], and we can avoid this step, but we provide results on L for the sake of comparison.

Our first result pertains to the transience and recurrence of Y . Roughly speaking, the classification depends on the mean of the underlying Lévy process, η . We distinguish

four possible cases:

$$a < \int_0^1 (-\log v) \rho(dv), \quad (\text{T})$$

$$a = \int_0^1 (-\log v) \rho(dv), \quad (\text{NR})$$

$$a > \int_0^1 (-\log v) \rho(dv), \quad (\text{PR})$$

$$a > \int_0^1 (-\log v) \rho(dv) \text{ and } \int_0^1 v^{-r} \rho(dv) < \infty \text{ for some } r > 0. \quad (\text{ER})$$

We note that (T), (NR) and (PR) are mutually exclusive, whereas (ER) implies (PR).

Proposition 8. *For $t \geq 0$ and $x \in (0, c]$, the many-to-one formula*

$$\Psi_t f(x) = e^{\lambda_* t} \mathbb{E}_x[f(Y_t)]$$

holds. Further, defining $q_ = \lambda_* + \max\{\inf \psi_\eta, -\|\Pi_\eta\|\}$, we have the following cases.*

- (i) *If (T) holds, then Y is transient, $\lambda = q_* = \lambda_* + \inf \psi_\eta$ and $L_{c,c}(\lambda) < 1$.*
- (ii) *If (NR) holds, then Y is null recurrent, $\lambda = \lambda_*$, $L_{c,c}(\lambda) = 1$ and $L'_{c,c}(\lambda) = -\infty$.*
- (iii) *If (PR) holds, then Y is positive recurrent, $\lambda = \lambda_*$, $L_{c,c}(\lambda) = 1$ and $L'_{c,c}(\lambda) > -\infty$.*
- (iv) *If (ER) holds, then so does (PR), and moreover, for any $0 < w < r$, there exist $k > 0$ and $C > 0$ such that for all $x \in (0, c]$, $t \geq 0$, and f such that $f_w(x) := (x/c)^w f(x)$ is continuous and bounded,*

$$|\mathbb{E}_x[f(Y_t)] - \langle f, \nu \rangle| \leq \|f_w\| ((c/x)^w + C) e^{-kt},$$

where ν is the invariant distribution of Y . Indeed, we can take $k = -\psi_\eta(-w)$ and $C = \int (c/x)^w \nu(dx) = \frac{-\psi'_\eta(0)w}{\psi_\eta(-w)}$.

Proof. The proof of the many-to-one formula, $\Psi_t f(x) = \mathbb{E}_x[f(Y_t)e^{\lambda_* t}]$, follows in a similar way to the proof of Proposition 3, and we omit the full detail. Indeed, splitting on the first time η is either killed or jumps, applying Dynkin's identity and some simple algebraic manipulations shows that $\mathbb{E}_x[f(Y_t)e^{\lambda_* t}]$ also solves (7). Since solutions of (7) are unique, it is equal to $\Psi_t f(x)$ for each $x \in (0, c]$ and each $t \geq 0$.

Next we deal with the recurrence and transience of the process Y in the different cases. We will show that Y is transient when (T) holds and that Y is recurrent when either (NR) or (PR) hold, and defer the proof of the distinction between positive and null recurrence to Proposition 10.

Differentiating ψ_η and letting $q \rightarrow 0$, we have

$$\psi'_\eta(0) = a + \int_{(0,1)} \log v \rho(dv) \quad (10)$$

The recurrence or transience of Y is equivalent to the recurrence or transience of the reflected Lévy process η^b . Considering its paths, we observe that η^b is recurrent if and only if $\limsup_{t \rightarrow \infty} \eta_t = \infty$ a.s.; and by [36, Theorem 7.1], this occurs if and only if $\psi'_\eta(0) \geq 0$. Combining this with (10), we see that η^b , and hence Y , is recurrent if and only if (PR) or (NR) holds, and transient if (T) holds.

We now consider the values of λ and $L_{c,c}(\lambda)$ for each of the first three cases. By conditioning on the first time Y jumps away from c , it is straightforward to show that

$$L_{c,c}(q) = \begin{cases} 1 - \frac{a\Phi(q - \lambda_*)}{\|\Pi_\eta\| + q - \lambda_*}, & \text{if } q > q_* \\ \infty & \text{if } q \leq q_*, \end{cases} \quad (11)$$

where Φ is the right inverse of ψ_η . Since $q_* \geq -\|\Pi_\eta\|$, it is clear that the denominator is positive whenever $q > q_*$. Hence, $L_{c,c}(q) = 1$ for some $q \geq q_*$ if and only if there is a solution to the equation $\Phi(q - \lambda_*) = 0$ with $q \geq q_*$.

If $\psi'_\eta(0) \geq 0$, then $\Phi(0) = 0$ and so $q = \lambda_*$ gives us the solution we seek. It also follows that $L_{c,c}(\lambda) = \mathbb{P}_c(H(c) < \infty) = 1$ since $\psi'(0) \geq 0$ implies that Y is recurrent.

If $\psi'_\eta(0) < 0$, then $\Phi(q - \lambda_*)$ is strictly positive for all $q \geq q_*$, and so $\lambda = q_*$ and $L_{c,c}(\lambda) < 1$. To show that $q_* = \lambda_* + \inf \psi_\eta$, we argue as follows:

$$\psi_\eta(q) + \|\Pi_\eta\| = aq + \int_{-\infty}^0 e^{qx} \Pi_\eta(dx) \geq 0,$$

and the two summands on the right are positive and do not approach zero simultaneously (if $q \geq 0$). Hence $\inf \psi_\eta = \inf_{q \geq 0} \psi_\eta(q) > -\|\Pi_\eta\|$.

To prove the statements about the derivative of L , note that

$$L'_{c,c}(q) = -\frac{a\Phi'(q - \lambda_*)(q - \lambda_* + \|\Pi_\eta\|) - a\Phi(q - \lambda_*)}{(q - \lambda_* + \|\Pi_\eta\|)^2},$$

and observe that $\Phi'(q - \lambda_*) = \infty$ if and only if $q = \lambda_* + \inf \psi_\eta$. Since $\inf \psi_\eta = 0$ if and only if (NR) holds, we obtain that $L'_{c,c}(\lambda) = -\infty$ if and only if we are in case (NR), which was the claim.

Part (iv) follows by Theorem 2 of Goffard and Sarantsev [31].

This completes the proof. \square

Remark 9. (i) As remarked in the proof, the cases (T), (NR) and (PR) are equivalent to $\mathbb{E}[\eta_1]$ being negative, zero and positive respectively, and the moment condition in (ER) is equivalent to $\mathbb{E}[e^{r\eta_1}] < \infty$.

(ii) In previous works, the value of $L_{c,c}(\lambda)$ was used to determine the transience or recurrence of Y . We do not need this for this result, but we continue the spirit of these works by making heavy use of the supermartingale function $\ell(x) = L_{x,c}(\lambda)$ in our analysis of the case (T) in section 6.

4.1. Recurrent regimes

We now consider the cases (NR), (PR) and (ER) in more detail. Our goal is to find an invariant measure for the process Y , which will turn out to be an eigenmeasure of the growth-fragmentation operator.

The methods in the section come from the theory of spectrally negative Lévy processes, of which η is an example, and we begin by defining the *scale function* $W: [0, \infty) \rightarrow [0, \infty)$ with Laplace transform

$$\int_0^\infty e^{-\beta x} W(x) dx = \frac{1}{\psi_\eta(\beta)},$$

valid for $\beta > \Phi(0)$, where $\Phi(q) = \sup\{p : \psi_\eta(p) \geq q\}$. We will also use W to denote the (Stieltjes) measure associated with the scale function.

For details of the existence and properties of scale functions, we refer to [36, §8]. One property that will be useful below is that W has a left-derivative on $(0, \infty)$, which we will write as W' .

Having recalled some of the theory of scale functions, we are in a position to identify the invariant measure we seek.

Proposition 10. *Suppose we are in the case (PR) or (NR). Let m be the pushforward of the measure $W(dx)$ by the function $x \mapsto ce^{-x}$, that is,*

$$m(dy) = \frac{1}{a} \delta_c(dy) + W'(\log(c/y)) \frac{dy}{y}.$$

Then, $\mathbb{E}_c \left[\int_0^{H(c)} f(Y_s) ds \right] = \frac{a}{2B} \langle f, m \rangle$.

In case (PR), $\langle 1, m \rangle = \left(a + \int_0^1 \log s \rho(ds) \right)^{-1}$, and $\nu = m / \langle 1, m \rangle$ is the invariant distribution of Y .

In case (NR), $\langle 1, m \rangle = \infty$, and m is a (unique up to a multiplicative constant) σ -finite invariant measure of Y . In the latter case, define $\nu = m$.

Proof. We give quite a general proof, which works under assumptions (PR) and (NR), and even under rather general assumptions about Y (or η). In the case (PR) and under our running assumption that η has no Gaussian part and finite jump rate, a simpler (if not shorter) proof is available; see Remark 11(i).

Let $R_t = \bar{\eta}_t - \eta_t$, the process η reflected in its running maximum, and define $L_t^{-1} := \inf\{s > 0 : \bar{\eta} > t\}$ to be the inverse local time of η at the maximum. Let $\mathbb{P} = \mathbb{P}_c$. Under \mathbb{P} , we have $Y_t = e^{b-R_t}$, so L^{-1} can be interpreted either as the inverse local time of R at 0 or the inverse local time of Y at c .

For arbitrary f , define $g(x) = f(ce^{-x})$. Following the proof of [9, Theorem VI.20], we obtain

$$\begin{aligned} \mathbb{E} \left[\int_0^{L_t^{-1}} f(Y_s) ds \right] &= \mathbb{E} \left[\int_0^{L_t^{-1}} g(R_s) ds \right] = \mathbb{E} \left[\int_0^\infty g(R_s) \mathbb{1}_{\{\bar{\eta}_s < t\}} ds \right] \\ &= k \int_0^\infty \hat{U}(dz) g(z) \int_0^t U(dy) \\ &= k \delta^{-1} t \langle g, \hat{U} \rangle, \end{aligned}$$

where U and \hat{U} are the ascending and descending renewal measures of η , δ is the drift coefficient of the upward ladder height process and k is some constant.

Now we note that $\hat{U}(dz) = k'e^{-\Phi(0)z}W(dz)$, where k' is another constant [9, p. 196, equation (4)]. Since we are in the case where Y is recurrent (and hence η does not drift to $-\infty$), $\Phi(0) = 0$.

In conclusion, we have shown that $\mathbb{E} \int_0^{L_t^{-1}} f(Y_s) ds = k''t \langle f, m \rangle$, with m as in the proposition and k'' a constant. Since $\mathbb{P}_c(H(c) < \infty) = 1$, [25, §XIX.3.46] implies that m is an invariant measure for Y .

The fact that $1/\langle 1, m \rangle = \mathbb{E}_0[\eta_1]$ follows from the Laplace transform of the measure W appearing on p. 238 of [36].

Finally, we deduce the claim about $\mathbb{E}_c \int_0^{H(c)} f(Y_s) ds$. It is shown in [30, Theorem 8.1] that this measure is invariant for Y (and indeed, this result is at its heart the same as that of [25] cited above). Since the invariant measure is unique up to normalisation, it suffices to compute the two measures of interest for some simple function in order to determine the correct normalisation.

Let $f = \mathbb{1}_{\{c\}}$. On the one hand, we have $\langle f, m \rangle = \frac{1}{a}$. On the other, if we define $T_1 = \inf\{t \geq 0 : Y_t < c\}$, we have

$$\mathbb{E}_c \int_0^{H(c)} f(Y_s) ds = \mathbb{E}_c[T_1] = \frac{1}{\|\Pi_\eta\|},$$

and, recalling that $\|\Pi_\eta\| = 2B$, this proves the desired normalisation. □

Remark 11. (i) It is also possible to obtain the formula for m by considering the first time η exits an interval $[-x, b]$ for $x > 0$, using the same method as [36, Theorem 8.7], and then letting $x \rightarrow \infty$. Then, the invariance of m follows, under assumption (PR), from [1, VI.1.5].

(ii) We also note that m is still an invariant measure when ξ (or η) is replaced by an arbitrary spectrally negative Lévy process, even if it has a Gaussian part or infinite jump rate; however, then m should be interpreted in terms of the excursions of Y from c .

(iii) The measure W has Laplace transform $\beta \mapsto \frac{\beta}{\psi_\eta(\beta)}$. Under assumption (PR), since ν is the unique invariant probability measure for Y , the previous result is equivalent, by Laplace inversion, to [1, Corollary IX.3.4], which gives another proof even under the general Lévy assumption. However, we are not aware of an existing proof of this result in the null recurrent case.

We are now in a position to solve the eigenproblem (4) and state the asynchronous exponential growth behaviour (6), which are interesting results in their own right, as well as stepping stones in our proof of the strong law of large numbers. It is worth pointing out that, whereas our main strong law result, Theorem 1, is stated under the assumption that $\lambda_* > 0$ and (ER) holds, these weaker results hold under more general assumptions, and in particular make no requirement on the growth or decay of the number of particles.

Proposition 12. *Assume (NR) or (PR). Let $h \equiv 1$ and ν be as defined above. Then:*

(i) $e^{\lambda t}$ is the spectral radius of Ψ_t .

(ii) $h \in \mathcal{D}(\mathcal{A})$ and $\mathcal{A}h = \lambda h$.

(iii) For every $f \in \mathcal{D}(\mathcal{A}) \cap L^\infty(0, c]$, $\langle \mathcal{A}f, \nu \rangle = \lambda \langle f, \nu \rangle$.

Proof. (i) Letting $(P_t)_{t \geq 0}$ denote the semigroup for Y , we have $\|\Psi_t\| = e^{\lambda t} \|P_t\|$. On the one hand, P_t is a contraction, so its operator norm is bounded above by 1. On the other hand, setting $f = 1$ yields $\|P_t\| = \sup_{f \in L^\infty} \|P_t f\| / \|f\| \geq 1$. Thus, $\|P_t\| = 1$ and hence the spectral radius of Ψ_t is given by

$$\lim_{t \rightarrow \infty} \|\Psi_t\|^{1/t} = e^\lambda \lim_{t \rightarrow \infty} \|P_t\|^{1/t} = e^\lambda.$$

(ii) The first claim follows from Corollary 5 and the second from the definition of λ .

(iii) First note that from the proof of Proposition 4 and Corollary 5, the set of function D_0 is a core for $(\mathcal{A}, \mathcal{D}(\mathcal{A}))$. Hence, we first show that this claim holds for $f \in D_0 \cap L^\infty(0, c]$.

Due to Proposition 6 and the fact that ν is the left eigenmeasure for Ψ_t corresponding to the eigenvalue $e^{\lambda t}$, we have $\langle \Psi_t f, \nu \rangle = \langle W_t f, \nu \rangle$. Then, due to Corollary 5, the claim follows for $f \in D_0 \cap L^\infty(0, c]$.

Now choose $f \in \mathcal{D}(\mathcal{A}) \cap L^\infty(0, c]$. Since D_0 is a core of \mathcal{A} , this means we can choose a sequence of functions $(f_n)_{n \geq 1} \subset D_0$ such that $\|f_n - f\|_{\mathcal{A}} \rightarrow 0$ as $n \rightarrow \infty$, where $\|\cdot\|_{\mathcal{A}}$ denotes the graph norm. In particular, this means that both $f_n \rightarrow f$ and $\mathcal{A}f_n \rightarrow \mathcal{A}f$ in $L^2((0, c])$ as $n \rightarrow \infty$, and hence the result follows. \square

Proposition 13. *Assume (PR). For all $f \in C_b((0, c])$, and all $x \in (0, c]$,*

$$\lim_{t \rightarrow \infty} e^{-\lambda t} \Psi_t f(x) = \langle f, \nu \rangle.$$

If (ER) holds, then there exists $k > 0$ such that for all $x \in (0, c]$ and $f \in C_b((0, c])$,

$$\lim_{t \rightarrow \infty} e^{kt} |e^{-\lambda t} \Psi_t f(x) - \langle f, \nu \rangle| = 0.$$

Proof. This is a corollary of Proposition 8 and Proposition 10. \square

The two preceding propositions prove equations (4) and (6) in the introduction; that is, they demonstrate the asynchronous exponential growth of the expectation semigroup, and its connection with the spectral theory of the operator \mathcal{A} .

5. Law of large numbers

This section is dedicated to the proof of Theorem 1, so we assume the process is supercritical (i.e., $\lambda_* = B - \mathbf{k} > 0$) and Y is exponentially recurrent. Recall that in this case $\lambda = \lambda_*$.

Let

$$M_t = e^{-\lambda t} N_t,$$

where N_t is the number of cells alive at time t . Then, M is a martingale; indeed, it is the intrinsic martingale in a continuous-time branching process (regardless of x). By [3, §III.4, p. 108], $\mathbb{E}_x[M_t^2] = 2(1 - e^{-\lambda t})$, so M is L^2 -bounded, and hence has a limit M_∞ almost surely and in L^1 . Note that Doob's martingale inequality also yields convergence in L^2 .

To prove Theorem 1, we will follow the ideas first used in [2], which were then later adapted to more general processes in [29, 33]. In particular, we use the skeleton decomposition, as in [33], in order to obtain almost sure convergence while also allowing particle death in the model. The technique of using spine methods to obtain the convergence of the mean measures is well-established (see, for instance, [6, 39]), but the strong law requires not just the additional use of the skeleton decomposition, but also fine control of the fluctuations around the spine. Because of this, our setting is not covered by the aforementioned articles, all of which require some form of uniform control with respect to the initial state when studying the convergence of the mean measures. This is not the case for our model, as seen from Proposition 8(iv), due to the behaviour of cells with size close to 0. However, we are still able to get enough control on the convergence by explicitly considering the dependence of the spine behaviour on its initial condition (see the proof of Lemma 20), which is not possible in the general setting.

5.1. Skeleton decomposition

A key element in the proof is a *skeleton decomposition* for the growth-fragmentation process. This corresponds to splitting \mathbf{Z} into a tree of particles that survive forever, dressed with trees of particles that all die out.

In order to describe this precisely, we start by specifying the non-linear semigroup of the growth-fragmentation process; this describes the behaviour of all the particles, rather than the average characterised by Ψ_t .

Recall \mathcal{U}_t , the set of labels of particles alive at time t , and for measurable functions f with $\|f\|_\infty < 1$, define the non-linear semigroup

$$u_t[f](x) := \mathbb{E}_x \left[\prod_{u \in \mathcal{U}_t} f(Z_u(t)) \right], \quad t \geq 0, x \in (0, c].$$

Proposition 14. *The semigroup $(u_t)_{t \geq 0}$ satisfies the following non-linear growth-fragmentation equation*

$$u_t[f](x) = f(xe^{a \cdot t \wedge T_c(x)}) + \int_0^t G[u_{t-s}[f]](xe^{a \cdot s \wedge T_c(x)}) ds, \quad (12)$$

where

$$G[f](x) = (B + \mathbf{k}) \left(\frac{B}{B + \mathbf{k}} \int_0^1 dv \kappa(v) f(xv) f(x(1-v)) + \frac{\mathbf{k}}{B + \mathbf{k}} - f(x) \right) \quad (13)$$

and $\kappa(v) = \pi(v)/vB$.

Proof. Conditioning on the time of the first event (fragmentation or killing), we have

$$\begin{aligned} u_t[f](x) &= f(xe^{a \cdot t \wedge T_c(x)}) e^{-(B+\mathbf{k})t} + \int_0^t ds (B + \mathbf{k}) e^{-(B+\mathbf{k})s} \\ &\quad \times \left(\frac{B}{B + \mathbf{k}} \int_{-\infty}^0 \frac{\Pi(du)}{B} u_{t-s}[f](e^{a \cdot s \wedge T_c(x)} e^u) u_{t-s}[f](e^{a \cdot s \wedge T_c(x)} (1 - e^u)) + \frac{\mathbf{k}}{B + \mathbf{k}} \right). \end{aligned}$$

Following the same steps as the linear case, namely making the change of variables $v = e^u$ and applying Dynkin's lemma, we obtain the required result. \square

For ease of notation, we will henceforth write

$$G[f](x) = (B + \mathbf{k}) \mathcal{E} [f(xV) f(x(1-V)) \mathbf{1}_{N=2} + \mathbf{1}_{N=0} - f(x)],$$

where under \mathcal{P} with expectation \mathcal{E} , $N \in \{0, 2\}$ is the number of particles produced at the first event (fragmentation or killing) with $\mathcal{P}(N = 2) = B/(B + \mathbf{k})$, and V is chosen according to κ .

We turn now to the skeleton decomposition. Let C_u denote the 'colour' of cell u . We write $C_u = b$ if the descendants of cell u survive forever (a 'blue' particle) and $C_u = r$ if the descendants of cell u eventually become extinct (a 'red' particle). Let $\mathbf{C}(t) = (C_u : u \in \mathcal{U}_t)$.

Define $\zeta := \inf\{t \geq 0 : N_t = 0\}$ to be the lifetime of the process, $p := \mathbb{P}_x(\zeta = \infty)$ to be the survival probability and $w = 1 - p$ to be the extinction probability. Note that since the fragmentation kernel and rate do not depend on the particle size, p and w do not depend on $x \in (0, c]$. By Proposition 7, $w = \frac{\mathbf{k}}{B} < 1$.

The following result about the colouring will be extremely useful in what follows:

Lemma 15. *Given \mathcal{F}_t , the colours of the cells of $\mathbf{Z}(t)$ are given by independently choosing each cell to be blue with probability p and red with probability w .*

Proof. This is shown in [33]. The independence of the colours is due to the spatial homogeneity of the branching and killing rates. \square

We denote by $(\mathbf{Z}, \mathbf{C}) = ((\mathbf{Z}(t), \mathbf{C}(t)) : t \geq 0)$ the process of cells marked by their colours.

We first describe the 'red trees'. For $A \in \mathcal{F}_t$ define

$$\mathbb{P}_x^R(A) := \mathbb{P}_x(A \mid C_\emptyset = r). \quad (14)$$

Proposition 16 (Red trees). *The dynamics of the process $(\mathbf{Z}, \mathbb{P}_x^R)$ can be described as follows. From a single particle at position $x \in (0, c]$, the particle will grow according to $s \mapsto xe^{a \cdot s \wedge T_c(x)}$, as before. Then, for $x \in (0, c]$ and bounded, measurable f , the branching generator is given by*

$$\begin{aligned} G^R[f](x) &= \frac{1}{w} (G[fw](x) - f(x)G[w](x)). \\ &= (B + \mathbf{k})\mathcal{E} \left[\frac{\mathbf{k}}{B + \mathbf{k}} f(xV)f(x(1 - V)) + \frac{B}{B + \mathbf{k}} - f(x) \right] \end{aligned} \quad (15)$$

The proof of this proposition can be found in [33, Prop 2.1]. We now consider the case where we condition on $C_\emptyset = b$. In order to do so, we first define a process $(\mathbf{Z}, \mathbb{P}^B)$, which should be thought of as a process representing the blue cells only. Started from an initial cell of size $x \in (0, c]$, the cell increases in size according to $s \mapsto xe^{a \cdot s \wedge T_c(x)}$ and will fragment according to the following operator,

$$\begin{aligned} G^B[f](x) &= \frac{1}{p} (G[pf + w] - (1 - f)G[w]) \\ &= B(p + 2w)\mathcal{E} \left[\frac{p}{p + 2w} f(xV)f(x(1 - V)) + \frac{w}{p + 2w} (f(xV) + f(x(1 - V))) - f(x) \right] \end{aligned} \quad (16)$$

Note $B(p + 2w) = B + \mathbf{k}$.

We are now ready describe the skeleton decomposition.

Proposition 17 (Skeleton decomposition). *With probability w , $(\mathbf{Z}, \mathbf{C}, \mathbb{P}_x)$ is equal in law to $(\mathbf{Z}, \mathbb{P}_x^R)$ with all cells coloured red; and with probability p , $(\mathbf{Z}, \mathbf{C}, \mathbb{P}_x)$ is equal in law to $(\mathbf{Z}, \mathbf{C}, \mathbb{P}_x(\cdot \mid C_\emptyset = b))$, which, in turn, is equal in law to a process $(\mathbf{Z}, \mathbb{P}_x^B)$ coloured blue, dressed with copies of $(\mathbf{Z}, \mathbb{P}_x^R)$ coloured red. In the latter case, the joint branching of the blue tree and dressing with red trees can be described via the following generator:*

$$\begin{aligned} H[f, g](x) &= B(2w + p)\mathcal{E} \left[\frac{p}{2w + p} f(xV)f(x(1 - V)) \right. \\ &\quad \left. + \frac{w}{2w + p} f(xV)g(x(1 - V)) + \frac{w}{2w + p} g(xV)f(x(1 - V)) - f(x) \right] \end{aligned} \quad (17)$$

The idea behind this proposition is that each particle in the growth-fragmentation process can either be coloured red, if its genealogy goes extinct, or blue, if its genealogy survives forever. In the case where the initial particle is blue, the branching operator H describes the colours of the offspring when a fragmentation occurs. In particular, the production of blue particles is described by the function f , whereas g describes the production of red particles. We can see from (17) that the first summand in the expectation on the righthand side of (17) describes the case where a particle fragments and both fragments are blue. The second (resp. third) summand then describes the case where the fragment of size xV (resp. $x(1 - V)$) is blue and the other is red.

The proof of this proposition follows from [33, Section 2.3] by setting $\zeta(x) = B + \mathbf{k}$ and using the specific form of the branching generator given in (13). In particular, the reader may find the proof for the exact formulation of H in the proof of [33, Proposition].

In order to prove the strong law, we will first prove it for the blue process $(\mathbf{Z}, \mathbb{P}^B)$, and then show that this implies the same result for (\mathbf{Z}, \mathbb{P}) .

The combination of Lemma 15 and Proposition 17 gives the following identity, which will be very useful.

$$\begin{aligned} \mathbb{E}_x^B[\langle f, \mathbf{Z}(t) \rangle] &= \mathbb{E}_x \left[\sum_u f(Z_u(t)) \mathbb{1}_{\{C_u=b\}} \mid C_\emptyset = b \right] = \frac{1}{p} \mathbb{E}_x \left[\sum_u f(Z_u(t)) \mathbb{1}_{\{C_u=b\}} \mathbb{1}_{\{C_\emptyset=b\}} \right] \\ &= \frac{1}{p} \mathbb{E}_x \left[\sum_u f(Z_u(t)) \mathbb{1}_{\{C_u=b\}} \right] \\ &= \frac{1}{p} \mathbb{E}_x[\langle f, p\mathbf{Z}(t) \rangle] \\ &= \mathbb{E}_x[\langle f, \mathbf{Z}(t) \rangle], \end{aligned} \tag{18}$$

where the first equality follows from the proposition; the third equality uses the fact that the sum is empty if the initial particle is not blue; and the penultimate equality comes from the lemma.

An immediate consequence is that if $(\lambda_*, 1, \nu)$ is the eigen-triple for (\mathbf{Z}, \mathbb{P}) , then $(\lambda_*, 1/p, p\nu)$ is the eigen-triple for the blue process $(\mathbf{Z}, \mathbb{P}^B)$. Moreover, the process Y defined by $\mathbb{E}_x^B[f(Y_t)] = e^{-\lambda_* t} \mathbb{E}_x^B[\langle f, \mathbf{Z}(t) \rangle]$ is the same in distribution as the process Y we defined earlier for (\mathbf{Z}, \mathbb{P}) .

Under the measures \mathbb{P}^B , we retain our notation N_t for the number of blue particles alive at time $t \geq 0$. Since λ is still the leading eigenvalue for the blue process, it follows that

$$M_t := e^{-\lambda t} N_t$$

is a positive martingale under \mathbb{P}_x^B , for each $x \in (0, c]$. Let M_∞ denote its limit. We will show in what follows that M is $L^2(\mathbb{P}_x^B)$ -convergent.

Our intermediate strong law is as follows.

Theorem 18. *For all $x \in (0, \infty)$ and continuous f with $\|f\| < 1$,*

$$e^{-\lambda t} \langle f, \mathbf{Z}(t) \rangle \rightarrow \langle f, \nu \rangle M_\infty, \quad t \rightarrow \infty,$$

\mathbb{P}_x^B -almost surely and in $L^1(\mathbb{P}_x^B)$.

5.2. Proof of Theorem 18

In this part, we follow closely the ideas of [29], though of course our processes are very different in nature.

Lemma 19. *Let $U_t = e^{-\lambda_* t} \langle f, \mathbf{Z}(t) \rangle$. Then, for any increasing sequence $(m_n)_{n \geq 1}$,*

$$U_{(m_n+n)\delta} - \mathbb{E}_x^B[U_{(m_n+n)\delta} \mid \mathcal{F}_{n\delta}] \rightarrow 0,$$

as $n \rightarrow \infty$, \mathbb{P}_x^B -a.s. and in $L^1(\mathbb{P}_x^B)$.

Proof. For convenience, we will write m instead of m_n . For almost sure convergence, by the Borel-Cantelli lemma, it is sufficient to show that, for all $\epsilon > 0$,

$$\sum_{n \geq 1} \mathbb{P}_x^B \left(|U_{(m+n)\delta} - \mathbb{E}_x^B[U_{(m+n)\delta} | \mathcal{F}_{n\delta}]| > \epsilon \right) < \infty.$$

We first note that

$$\mathbb{E}_x^B \left[\left| U_{(m+n)\delta} - \mathbb{E}_x^B[U_{(m+n)\delta} | \mathcal{F}_{n\delta}] \right|^2 \middle| \mathcal{F}_{n\delta} \right] = \mathbb{E}_x^B \left[\left| \sum_u e^{-n\delta\lambda} (U_{m\delta}^{(u)} - \mathbb{E}_{Z_u(n\delta)}^B[U_{m\delta}^{(u)}]) \right|^2 \middle| \mathcal{F}_{n\delta} \right],$$

where, conditional on $\mathcal{F}_{n\delta}$, the $U_{m\delta}^{(u)}$ are independent and distributed as $(U, \mathbb{P}_{Z_u(n\delta)})$ for $u \in \mathcal{U}_{n\delta}$. It follows that

$$\begin{aligned} \mathbb{E}_x^B \left[\left| U_{(m+n)\delta} - \mathbb{E}_x^B[U_{(m+n)\delta} | \mathcal{F}_{n\delta}] \right|^2 \middle| \mathcal{F}_{n\delta} \right] &= \mathbb{E}_x^B \left[\left| \sum_u e^{-n\delta\lambda} (U_{m\delta}^{(u)} - \mathbb{E}_{Z_u(n\delta)}^B[U_{m\delta}^{(u)}]) \right|^2 \middle| \mathcal{F}_{n\delta} \right] \\ &= \mathbb{E}_x^B \left[\sum_u e^{-2n\delta\lambda} |U_{m\delta}^{(u)} - \mathbb{E}_{Z_u(n\delta)}^B[U_{m\delta}^{(u)}]|^2 \middle| \mathcal{F}_{n\delta} \right] \\ &\leq 2\mathbb{E}_x^B \left[\sum_u e^{-2n\delta\lambda} (|U_{m\delta}^{(u)}|^2 + |\mathbb{E}_{Z_u(n\delta)}^B[U_{m\delta}^{(u)}]|^2) \middle| \mathcal{F}_{n\delta} \right] \\ &\leq 4e^{-2\lambda n\delta} \sum_u \mathbb{E}_{Z_u(n\delta)}^B \left[(U_{m\delta}^{(u)})^2 \right] \\ &\leq 4e^{-n\delta\lambda} \|f\|^2 \sup_{x,t} \mathbb{E}_x^B[(M_t)^2] \cdot M_{n\delta}. \end{aligned} \quad (19)$$

where, for the second equality, we used that the sum is over conditionally independent, zero-mean summands, for the first inequality we have used that $|a + b|^2 \leq 2(|a|^2 + |b|^2)$ and for the second inequality we have used Jensen's inequality.

A similar calculation to that of (18) shows that $\mathbb{E}_x^B[(M_t)^2] = p\mathbb{E}_x[M_t^2]$, and hence, M is $L^2(\mathbb{P}_x^B)$ -bounded, uniformly in $x \in (0, c]$. From Doob's inequality, it follows that M is $L^2(\mathbb{P}_x^B)$ -convergent and hence $L^1(\mathbb{P}_x^B)$ -convergent. Therefore,

$$\mathbb{E}_x^B \left[(U_{(m+n)\delta} - \mathbb{E}_x^B[U_{(m+n)\delta} | \mathcal{F}_{n\delta}])^2 \right] \leq 2\|f\|^2 \sup_{x,t} \mathbb{E}_x^B[M_t^2] \cdot e^{-n\delta\lambda},$$

which is summable, so the Markov inequality completes the proof of the almost sure convergence. \square

Lemma 20. *Theorem 18 holds when restricted to lattice times:*

$$\lim_{n \rightarrow \infty} e^{-n\delta\lambda} \langle f, \mathbf{Z}(n\delta\lambda) \rangle = \langle f, \nu \rangle M_\infty,$$

\mathbb{P}_x^B -almost surely and in $L^1(\mathbb{P}_x^B)$.

Proof. Noting that

$$\mathbb{E}_x^B [e^{-\lambda 2n\delta} \langle f, \mathbf{Z}(2n\delta) \rangle \mid \mathcal{F}_{n\delta}] = e^{-\lambda n\delta} \sum_u e^{-\lambda n\delta} \mathbb{E}_{Z_u(n\delta)}^B [\langle f, \mathbf{Z}(n\delta) \rangle],$$

we have

$$\begin{aligned} e^{-\lambda 2n\delta} \langle f, \mathbf{Z}(2n\delta) \rangle &= e^{-\lambda 2n\delta} \langle f, \mathbf{Z}(2n\delta) \rangle - e^{-\lambda 2n\delta} \mathbb{E}_x^B [\langle f, \mathbf{Z}(2n\delta) \rangle \mid \mathcal{F}_{n\delta}] \\ &\quad + e^{-\lambda n\delta} \sum_u \left(e^{-\lambda n\delta} \mathbb{E}_{Z_u(n\delta)}^B [\langle f, \mathbf{Z}(n\delta) \rangle] - \langle f, \nu \rangle \right) \\ &\quad + \langle f, \nu \rangle M_{n\delta}. \end{aligned}$$

The first term goes to zero a.s. and in $L^1(\mathbb{P}_x^B)$, by setting $m_n = n$ in the preceding lemma. The last term approaches $\langle f, \nu \rangle M_\infty$ by the martingale convergence discussed previously.

If we define

$$R_{s,t} = \sum_u e^{-\lambda t} |e^{-\lambda s} \mathbb{E}_{Z_u(t)}^B [\langle f, \mathbf{Z}(s) \rangle] - \langle f, \nu \rangle|,$$

then, using again (18), we have

$$R_{s,t} = \sum_u e^{-\lambda t} |e^{-\lambda s} \mathbb{E}_{Z_u(t)}^B [\langle f, \mathbf{Z}(s) \rangle] - \langle f, \nu \rangle| = \sum_u e^{-\lambda t} |\mathbb{E}_{Z_u(t)}^B [f(Y_s)] - \langle f, \nu \rangle|$$

it is sufficient to show that $\lim_{n \rightarrow \infty} R_{n\delta, n\delta} = 0$ almost surely. We now wish to apply Proposition 8. This was proven under the measures \mathbb{P} , but as remarked above, the distribution of the spine Y is the same under \mathbb{P}^B . By Proposition 8, choose $w \in (0, r)$ such that $\psi_\eta(-w) < 0$; then, there exist $k > 0$ and $C > 0$ such that for all $x \in (0, c]$ and $t \geq 0$,

$$|\mathbb{E}_x^B [f(Y_s)] - \langle f, \nu \rangle| \leq \|f_w\| ((c/x)^w + C) e^{-ks},$$

where we recall $f_w(x) = (x/c)^w f(x)$.

Applying this in our situation, we have

$$\mathbb{E}_x [R_{s,t}] \leq e^{-ks} \|f_w\| \mathbb{E}_x [(c/Y_t)^w + C] = e^{-ks} \|f_w\| \mathbb{E}_x [e^{w(b-\eta_t^b)} + C].$$

We estimate:

$$\begin{aligned} \mathbb{E}_x [e^{w(b-\eta_t^b)}] &= \mathbb{E}_x [e^{w(\bar{\eta}_t \vee b - \eta_t)}] \leq e^{w(b-x)} \mathbb{E} [e^{w(\bar{\eta}_t - \eta_t)}] \\ &= e^{w(b-x)} \mathbb{E} [e^{-w\eta_t}] \\ &\leq e^{w(b-x)} \mathbb{E} [e^{-w\eta_\infty}]. \end{aligned}$$

We will show that the right-hand side is finite, using the Wiener-Hopf factorisation of η .

Let \hat{H} denote the descending ladder height process of η and $\hat{\kappa}(0, \cdot)$ its Laplace exponent. By [36, p. 178], we know that $\eta_\infty \stackrel{d}{=} -\hat{\mathbb{H}}_{\mathbf{e}_\chi}$, where $\hat{\mathbb{H}}$ is an unkilled version of \hat{H} , and \mathbf{e}_χ is an independent exponential random variable with rate $\chi = \hat{\kappa}(0, 0) > 0$.

By decomposing the expectation according to the law of \mathbf{e}_x , we see that $\mathbb{E}[e^{w\hat{H}\mathbf{e}_x}] < \infty$ if and only if $\mathbb{E}[e^{w\hat{H}1}] < \infty$ and $\hat{\kappa}(0, -w) > 0$. Furthermore, by [36, Theorem 7.8], we know that the Lévy measures of \hat{H} and η are related by $\Pi_{\hat{H}}(dy) = \Pi_{\eta}(y, \infty) dy$, so if

$$\mathbb{E}[e^{w\eta_1}] < \infty, \quad (20)$$

then $\mathbb{E}[e^{w\hat{H}1}] < \infty$ also.

Now, by assumption, our choice of w satisfies (20) and $\psi_{\eta}(-w) < 0$, so the results of [36, §6.5.2] imply that $\hat{\kappa}(0, -w) = \frac{\psi_{\eta}(-w)}{-w} > 0$. We conclude that $\mathbb{E}[e^{-w\eta_{\infty}}] < \infty$.

It follows that

$$\mathbb{E}_x[R_{s,t}] \leq e^{-ks-wx} \|f_w\| C',$$

with $C' = e^{wb}\mathbb{E}[e^{-w\eta_{\infty}}] + C$. Hence, we have convergence in L^1 . Finally, by the Markov inequality,

$$\sum_{n \geq 1} \mathbb{P}_x(R_{n\delta, n\delta} > \epsilon) < \infty.$$

The Borel-Cantelli lemma implies that $\lim_{n \rightarrow \infty} R_{n\delta, n\delta} = 0$ almost surely. \square

Proof of Theorem 18. To complete the proof of Theorem 18, we need to pass from lattice time to continuous time. For the $L^1(\mathbb{P}_x^B)$ convergence, the Croft-Kingman lemma yields the result. For the almost sure part, the proof is identical to the proof given in [29], and so we omit it. \square

5.3. Proof of Theorem 1

In this section, we work under the probability measure \mathbb{P} , and will make use of both the full growth-fragmentation process \mathbf{Z} and the part of it coloured blue, which we denote by \mathbf{Z}^B (and which, if it exists, has the law of $(\mathbf{Z}, \mathbb{P}^B)$). The intrinsic martingale of the blue tree will be denoted M^B .

First note that the L^1 convergence for the full process follows easily from the relation (18). For the almost sure convergence, the proof follows the same idea as [33, §4], using the following proposition, a discussion of whose proof can also be found in that reference.

Proposition 21. *Let $(\Omega, \mathcal{F}, (\mathcal{F}_t, t \geq 0), \mathbb{P})$ be a filtered probability space and define $\mathcal{F}_{\infty} := \sigma(\cup_{i=1}^{\infty} \mathcal{F}_i)$. Suppose $(U_t, t \geq 0)$ is an \mathcal{F} -measurable non-negative process such that $\sup_{t \geq 0} U_t$ has finite expectation and $(\mathbb{E}(U_t | \mathcal{F}_t), t \geq 0)$ is càdlàg. If*

$$\lim_{t \rightarrow \infty} \mathbb{E}(U_t | \mathcal{F}_{\infty}) = Y, \text{ almost surely,}$$

then

$$\lim_{t \rightarrow \infty} \mathbb{E}(U_t | \mathcal{F}_t) = Y, \text{ almost surely.}$$

To put the above proposition in the context of the growth-fragmentation setting, set $U_t = e^{-\lambda t} \langle f, \mathbf{Z}^B(t) \rangle$, for f satisfying the conditions of Theorem 1, and recall that $(\mathcal{F}_t, t \geq 0)$ is the filtration generated by the growth-fragmentation process $(\mathbf{Z}(t), t \geq 0)$.

Note that we can easily bound $(U_t, t \geq 0)$ by a multiple of $(M_t^B, t \geq 0)$ and hence we automatically get that $\sup_{t \geq 0} U_t$ has a second, and hence first, moments thanks to (18). Due to Theorem 18 and the fact that $\mathbf{Z}^B(t)$ is \mathcal{F}_∞ -measurable, $U_t = \mathbb{E}(U_t | \mathcal{F}_\infty)$ and hence

$$\lim_{t \rightarrow \infty} \mathbb{E}(U_t | \mathcal{F}_\infty) = \langle f, \nu \rangle M_\infty^B,$$

\mathbb{P}_x -almost surely, for $x \in (0, c]$.

Using (18), we get

$$\mathbb{E}(U_t | \mathcal{F}_t) = \mathbb{E}(e^{-\lambda t} \langle f, \mathbf{Z}^B(t) \rangle | \mathcal{F}_t) = e^{-\lambda t} p \langle f, \mathbf{Z}(t) \rangle.$$

Combining this with Proposition 21 yields

$$\lim_{t \rightarrow \infty} e^{-\lambda t} \langle f, X_t \rangle = \langle f, \nu \rangle M_\infty^B / p, \quad (21)$$

\mathbb{P}_x -almost surely.

To complete the proof of almost sure convergence, we need to show that $M_\infty^B / p = M_\infty$, almost surely. To do so, take $f = 1$ in (21) and observe that the left-hand side is M_∞ . \square

6. Long-term behaviour in the transient regime

Finally, in this last section, we will prove Theorem 2. Consider the case where \mathbf{Z} is supercritical but (T) holds, so that $\lambda = q_* = \lambda_* + \inf \psi_\eta < \lambda_*$.

To simplify the exposition, in this section we will assume that $\mathbf{k} = 0$, that is, that there is no killing of cells. In the general case, the results of this section can be proved using the skeleton decomposition, as in the previous section.

Now, since we are assuming that (T) holds, we no longer have the advantage of being able to use M_t to define a change of measure, as in the previous section. However, in this case, we are able to use the function $\ell(x) = L_{x,c}(\lambda) = (x/c)^{\arg \inf \psi_\eta}$ to yield a useful *supermartingale* change of measure.

Our techniques in this section are inspired by a combination of [14] and [29].

Proposition 22. *(i) The process $S_t = e^{-(\lambda - \lambda_*)t} \frac{\ell(Y_t)}{\ell(x)} = e^{-\inf \psi_\eta \cdot t} \frac{\ell(Y_t)}{\ell(x)}$ is a \mathbb{P}_x -supermartingale for the natural filtration \mathcal{F}^Y of Y . Under the change of measure*

$$\left. \frac{d\tilde{\mathbb{P}}_x}{d\mathbb{P}_x} \right|_{\mathcal{F}_t^Y} = e^{-\inf \psi_\eta \cdot t} \frac{\ell(Y_t)}{\ell(x)},$$

Y is the exponential of a Lévy process with Laplace exponent $\tilde{\psi}_\eta(q) = \psi_\eta(q + \arg \inf \psi_\eta) - \inf \psi_\eta$, reflected in the level c and killed according to the multiplicative functional $t \mapsto e^{-a \cdot \arg \inf \psi_\eta \cdot \int_0^t \mathbb{1}_{\{Y_s=c\}} ds}$.

(ii) The process $\mathcal{S}_t = e^{-\lambda t} \frac{1}{\ell(x)} \langle \ell, \mathbf{Z} \rangle$ is a \mathbb{P}_x -supermartingale for the natural filtration of \mathbf{Z} . A measure $\tilde{\mathbb{P}}_x$ supporting \mathbf{Z} and the additional random variable ζ can be defined as follows:

$$\tilde{\mathbb{P}}_x(F_t \mathbb{1}_{\{\zeta > t\}}) = \mathbb{P}_x(F_t \mathcal{S}_t), \quad F_t \in \mathcal{F}_t.$$

Under this measure, the process \mathbf{Z} has the following decomposition. There is a single distinguished cell whose mass has the distribution of the process Y under $\tilde{\mathbb{P}}_x$, and at every jump ΔY_t of this cell, a copy of \mathbf{Z} under measure $\mathbb{P}_{-\Delta Y_t}$ is introduced, which we can denote $\mathbf{Z}^{[t]}$. That is, \mathbf{Z} has the same distribution as the process

$$t \mapsto \delta_{Y_t} + \sum_{0 < s \leq t} \mathbf{Z}^{[s]}(t - s).$$

The random variable ζ is the lifetime of Y .

(iii) Assume that $\lambda > 0$. Then, $\sup_{x \in (0, c]} \sup_{t \geq 0} \ell(x) \mathbb{E}_x[\mathcal{S}_t^2] < \infty$.

Proof. (i) Let $\hat{\ell}(y) = e^{\arg \inf \psi_\eta \cdot (y - b)}$, so that $\ell(Y_t) = \hat{\ell}(\eta_t^b)$.

The \mathbb{P}_x -supermartingale property of S is equivalent to the statement that

$$\mathbb{E}_y[e^{-\inf \psi_\eta \cdot t} \hat{\ell}(\eta_t^b) / \hat{\ell}(x)] \leq 1,$$

since η^b is Markov (where $y = \log x$.) Recall that $\mathbb{E}_y[e^{\alpha \eta_t - \psi_\eta(\alpha) t - \alpha y}] = 1$, for any $\alpha \geq 0$. Then,

$$\begin{aligned} \mathbb{E}_y[e^{\alpha \eta_t^b - \psi(\alpha) t - \alpha y}] &= e^{-\alpha y - \psi_\eta(\alpha) t} \mathbb{E}_y[e^{\alpha(\eta_t - (\bar{\eta}_t - b) \vee 0)}] \\ &= \mathbb{E}_y[e^{-\alpha(\bar{\eta}_t - b) \vee 0}] \leq 1. \end{aligned}$$

Setting $\alpha = \arg \inf \psi_\eta > 0$ completes the proof of the supermartingale property.

We turn to the characterisation of Y . It appears to be simplest to demonstrate this using evolution equations. Therefore, let $(\Phi_t)_{t \geq 0}$ represent the semigroup of Y under \mathbb{P} ., and $(\tilde{\Phi}_t)_{t \geq 0}$ the same object under $\tilde{\mathbb{P}}$..

It is simple to show, using the same ideas as in section 2, that Φ satisfies the evolution equation

$$\begin{aligned} \Phi_t f(x) &= f(x e^{a \cdot t \wedge T_c(x)}) \\ &+ \int_0^t ds \int_0^1 (\Pi_\eta \circ \exp^{-1})(dv) [\Phi_{t-s} f(x e^{a \cdot s \wedge T_c(x)} v) - \Phi_{t-s} f(x e^{a \cdot s \wedge T_c(x)})]. \end{aligned}$$

The semigroups Φ and $\tilde{\Phi}$ are related by the formula

$$\tilde{\Phi}_t f(x) = e^{-\inf \psi_\eta \cdot t} \frac{1}{\ell(x)} \Phi_t(f \ell)(x).$$

Rewriting the above evolution equation in terms of $\tilde{\Phi}$, performing some algebraic manipulation and making use of Dynkin's integral identity (Lemma 27) we obtain the following evolution equation for $\tilde{\Phi}$:

$$\begin{aligned} \tilde{\Phi}_t f(x) &= f(xe^{a \cdot t \wedge T_c(x)}) - \int_{T_c(x)}^t \alpha \tilde{\Phi}_{t-s} f(xe^{a \cdot s \wedge T_c(x)}) ds \\ &\quad + \int_0^t ds \int_0^1 (\tilde{\Pi}_\eta \circ \exp^{-1})(dv) [\tilde{\Phi}_{t-s} f(xe^{a \cdot s \wedge T_c(x)} v) - \tilde{\Phi}_{t-s} f(xe^{a \cdot s \wedge T_c(x)})], \end{aligned}$$

where $\alpha = \arg \inf \psi_\eta$ and $\tilde{\Pi}_\eta$ is the Lévy measure of the Lévy process with Laplace exponent $\tilde{\psi}_\eta$ given in the statement. It is clear that this is the evolution equation associated with the process described in the statement of the result.

It remains to show that the above evolution equation characterises $\tilde{\Phi}$. Since the killing and jump rates are bounded, this follows using Gronwall's inequality exactly as in section 2.

- (ii) This proof is very similar to the classical methods of [32], and we give only an outline. The first step is to introduce an additional (killed) process I_t on \mathcal{U} by defining

$$\tilde{\mathbb{P}}_x(F_t \mathbb{1}_{\{I_t=u\}}) = e^{-\lambda t} \frac{1}{\ell(x)} \mathbb{P}_x(F_t \ell(Z_u(t))), \quad F_t \in \mathcal{F}_t,$$

and declaring ζ to be the killing time of I . This random variable I_t indicates the index of the distinguished 'spine' cell.

It follows that

$$\tilde{\mathbb{E}}_x[f(Z_{I_s}(s), s \leq t)] = e^{-\lambda t} \frac{1}{\ell(x)} \sum_u \mathbb{E}_x[f(Z_u(s), s \leq t) \ell(t)],$$

and in particular, taking part (i) into account, we see that under $\tilde{\mathbb{P}}_x$, $t \mapsto Z_{I_t}(t)$ has the same distribution as the process Y .

Denote by $T_k(u)$ the k -th jump of the cell labelled u (or its ancestors), and by T_k the k -th jump of Y . Since all processes involved are Markov, it suffices to check the decomposition at fixed times, and we will focus first on the case where $T_1(I_t) \leq t < T_2(I_t)$; that is, between the first and second jumps of the spine. We define an operator r on \mathcal{U} which removes the prefix, i.e., if $u = u_1 u_2 u_3 \dots$,

$ru = u_2 u_3 \dots$. Now let f and g be measurable functions.

$$\begin{aligned}
& \tilde{\mathbb{E}}_x \left[f(Z_{I_t}(t)) g(Z_v(t), v \neq I_t) \mathbb{1}_{\{\zeta > t\}} \mathbb{1}_{\{T_1(I_t) \leq t < T_2(I_t)\}} \right] \\
&= \sum_u e^{-\lambda t} \frac{1}{\ell(x)} \mathbb{E}_x \left[f(Z_u(t)) g(Z_v(t), v \neq u) \ell(Z_u(t)) \mathbb{1}_{\{T_1(u) \leq t < T_2(u)\}} \right] \\
&= \sum_u e^{-\lambda t} \frac{1}{\ell(x)} \mathbb{E}_x \left[\mathbb{E}_{Z_u(T_1(u))} [f(Z_{ru}(t-s)) \ell(Z_{ru}(t-s)) \mathbb{1}_{\{t-s < T_1(ru)\}}] \right. \\
&\quad \left. \times \mathbb{E}_{-\Delta Z_u(T_1(u))} [g(Z_{rv}(t-s), v \neq u)]|_{s=T_1(u)} \mathbb{1}_{\{T_1(u) \leq t\}} \right] \\
&= \tilde{\mathbb{E}}_x \left[\tilde{\mathbb{E}}_{Y_{T_1}} [f(Y_{t-s}) \mathbb{1}_{\{t-s < T_1\}}] \mathbb{E}_{-\Delta Y_{T_1}} [g(\mathbf{Z}(t-s))] |_{s=T_1} \mathbb{1}_{\{T_1 \leq t\}} \right] \\
&= \tilde{\mathbb{E}}_x \left[f(Y_t) \mathbb{E}_{-\Delta Y_{T_1}} [g(\mathbf{Z}(t-s))] |_{s=T_1} \mathbb{1}_{\{T_1 \leq t < T_2\}} \right].
\end{aligned}$$

This proves the claim on the event $\{T_1(I_t) \leq t < T_2(I_t)\}$, and the full proof proceeds by induction, in each case conditioning at time $T_1(I_t)$.

(iii) This problem can be approached using the spine decomposition. We have that

$$\ell(x) \mathbb{E}_x [\mathcal{S}_t^2] = \ell(x) \tilde{\mathbb{E}}_x [\mathcal{S}_t] = e^{-\lambda t} \tilde{\mathbb{E}}_x [\ell(Y_t)] + \tilde{\mathbb{E}}_x \left[\sum_{s < t} e^{-\lambda s} \cdot e^{-\lambda(t-s)} \langle \ell, \mathbf{Z}^{[s]}(t-s) \rangle \right],$$

where the sum is over the jump times s of Y up until its lifetime. Since ℓ is bounded, the first term is bounded in t, x . To bound the second term, it is helpful to recall that Y is the exponential of a Lévy process η reflected in b , with extra killing. Therefore, if we write \mathcal{F}^Y for the filtration of the distinguished cell whose evolution has path Y , we have

$$\begin{aligned}
\tilde{\mathbb{E}}_x \left[\sum_{s < t} e^{-\lambda s} e^{-\lambda(t-s)} \langle \ell, \mathbf{Z}^{[s]}(t-s) \rangle \middle| \mathcal{F}_\infty^Y \right] &= \sum_{s < t} e^{-\lambda s} \ell(-\Delta Y_s) \mathbb{E}_{-\Delta Y_s} [\mathcal{S}_{t-s}] \\
&\leq \sum_{s < t} e^{-\lambda s} \ell(-\Delta Y_s).
\end{aligned}$$

Using the tower property and the predictable compensator of the jumps of Y , we obtain

$$\tilde{\mathbb{E}}_x \left[\sum_{s < t} e^{-\lambda s} e^{-\lambda(t-s)} \langle \ell, \mathbf{Z}^{[s]}(t-s) \rangle \right] \leq \tilde{\mathbb{E}}_x \left[\int_0^t e^{-\lambda s} \int_{-\infty}^0 \ell(Y_{s-}(1 - e^z)) \tilde{\Pi}_\eta(dz) \right],$$

Again, since ℓ is bounded and $\tilde{\Pi}_\eta$ is finite (because $\arg \inf \psi_\eta > 0$), this term is also bounded in t, x . □

Remark 23. (i) Though the process Y under $\tilde{\mathbb{P}}_x$ is a little unusual, it can be shown that its lifetime ζ satisfies

$$\begin{aligned}
\int_0^\infty e^{-qt} \tilde{\mathbb{P}}_c(\zeta > t) dt &= \int_0^\infty e^{-qt} \mathbb{E} [e^{-\arg \inf \psi_\eta \cdot (\bar{\eta}_t - \eta_t) - \inf \psi_\eta \cdot t}] dt \\
&= \frac{1}{\Phi(q + \inf \psi_\eta)} \frac{q}{\Phi(q + \inf \psi_\eta) - \arg \inf \psi_\eta},
\end{aligned}$$

This can be proved using the Wiener-Hopf factorisation, in particular the Laplace transform of the downward ladder height subordinator for a spectrally negative Lévy process [36, Theorem 6.15(ii) and equation (6.35)].

- (ii) The preceding proposition makes sense even in the presence of Gaussian fluctuations or infinite jump activity. The process Y is killed according to the local time at c . In the case of diffusions this is known as ‘elastic’ boundary behaviour [35, Example IV-5.5].
- (iii) Part (i) of the previous result proves that $\hat{\ell}$ is the extremal excessive function of the reflected Lévy process η_b associated with the Martin boundary point b . This may be of interest as an example in the context of spectrally negative Markov processes.

Lemma 24. *Let f be such that $\|f/\ell\|_\infty < \infty$, and fix $m \geq 0$. Define $U_t = e^{-\lambda t} \langle f, \mathbf{Z}(t) \rangle$. Then,*

$$U_{(m+n)\delta} - \mathbb{E}_x[U_{(m+n)\delta} \mid \mathcal{F}_{n\delta}] \rightarrow 0,$$

as $n \rightarrow \infty$, \mathbb{P}_x -a.s. and in $L^1(\mathbb{P}_x)$.

Proof. To be concise, let $s = m\delta$ and $t = n\delta$. Using the conditional independence of the zero-mean summands, we get

$$\begin{aligned} \mathbb{E}_x \left[(U_{s+t} - \mathbb{E}_x[U_{s+t} \mid \mathcal{F}_t])^2 \mid \mathcal{F}_t \right] &= \mathbb{E}_x \left[e^{-2\lambda t} \sum_i \left(e^{-\lambda s} \langle f, \mathbf{Z}^{(i)}(s) \rangle - \mathbb{E}_{Z_i(t)}[e^{-\lambda s} \langle f, \mathbf{Z}(s) \rangle] \right)^2 \mid \mathcal{F}_t \right] \\ &= \sum_i e^{-2\lambda t} \left(\mathbb{E}_{Z_i(t)}[(e^{-\lambda s} \langle f, \mathbf{Z}(s) \rangle)^2] - \mathbb{E}_{Z_i(t)}[e^{-\lambda s} \langle f, \mathbf{Z}(s) \rangle]^2 \right) \\ &\leq 2\|f/\ell\|_\infty^2 \sum_i e^{-2\lambda t} \mathbb{E}_{Z_i(t)}[(e^{-\lambda s} \langle \ell, \mathbf{Z}(s) \rangle)^2] \\ &\leq 2\|f/\ell\|_\infty^2 \sum_i e^{-2\lambda t} \ell(Z_i(t))^2 \mathbb{E}_{Z_i(t)}[\mathcal{S}_s^2] \\ &\leq 2\|f/\ell\|_\infty^2 \cdot \sup_{y,u} \ell(y) \mathbb{E}_y[\mathcal{S}_u^2] \cdot e^{-\lambda t} \ell(x) \mathcal{S}_t. \end{aligned}$$

Hence, using that \mathcal{S} is a supermartingale,

$$\mathbb{E}_x \left[(U_{s+t} - \mathbb{E}_x[U_{s+t} \mid \mathcal{F}_t])^2 \right] \leq \text{const} \cdot e^{-\lambda t} \ell(x).$$

Hence, the $L^1(\mathbb{P}_x)$ -convergence to zero holds. Moreover, this is summable in n (recall that $t = n\delta$) and so, using an application of Markov’s inequality and the Borel-Cantelli lemma, the almost sure convergence to zero holds too. \square

We are now able to state and prove the following rephrasing of our second main theorem. To understand the connection with the statement in the introduction, which is in terms of the ‘cumulant’ κ , observe that $\kappa(0) = \lambda_*$ and that

$$\psi_\eta(q) = \kappa(q) - \kappa(0).$$

Hence, $\inf \kappa = \inf \psi_\eta + \lambda_*$ and $\arg \inf \kappa = \arg \inf \psi_\eta$, so that in particular $\lambda = \inf \kappa$.

Theorem 25 (Rephrasing of Theorem 2). *Assume that $\lambda > 0$ and that $f: (0, c] \rightarrow \mathbb{R}$ is continuous and bounded with $f(x) = O(x^{\arg \inf \psi_\eta})$ as $x \rightarrow 0$. Then,*

$$e^{-\lambda t} \langle f, \mathbf{Z}(t) \rangle \rightarrow 0,$$

\mathbb{P}_x -almost surely and in $L^1(\mathbb{P}_x)$.

Proof. We begin with the proof for lattice times, and for simplicity, we assume $f \geq 0$. For $\delta > 0$, we have, as in the recurrent case,

$$\begin{aligned} e^{-\lambda(s+t)} \langle f, \mathbf{Z}(s+t) \rangle &= e^{-\lambda(s+t)} \langle f, \mathbf{Z}(s+t) \rangle - \mathbb{E}_x[e^{-\lambda(s+t)} \langle f, \mathbf{Z}(s+t) \rangle \mid \mathcal{F}_t] \\ &\quad + e^{-\lambda t} \sum_i \mathbb{E}_{Z_i(t)}[e^{-\lambda s} \langle f, \mathbf{Z}(s) \rangle]. \end{aligned}$$

The first term converges to zero in the sense stated along lattice times, by the preceding lemma. The second term can be expressed

$$\sum_i e^{-\lambda t} \mathbb{E}_{Z_i(t)}[e^{-\lambda s} \langle f, \mathbf{Z}(s) \rangle] = \sum_i e^{-\lambda t} \ell(Z_i(t)) \tilde{\mathbb{E}}_{Z_i(t)} \left[\frac{f(Y_s)}{\ell(Y_s)} \right] \leq \|f/\ell\|_\infty \mathcal{S}_t.$$

Since \mathcal{S} is a positive supermartingale, it converges almost surely [43, Corollary II.2.11]. Moreover, $\mathbb{E}_x[\mathcal{S}_t] = \tilde{\mathbb{P}}_x(\zeta > t) \rightarrow 0$, where ζ is the lifetime of Y . Hence, \mathcal{S}_t converges to zero in $L^1(\mathbb{P}_x)$. It follows that the almost sure limit of \mathcal{S}_t is also zero.

The extension of this limit from lattice to continuous times follows the same idea as in the recurrent regime. \square

Remark 26. If $e^{-\lambda' t} \langle f, \mathbf{Z}(t) \rangle \rightarrow 0$ in $L^1(\mathbb{P}_x)$ for some $\lambda' \in \mathbb{R}$ and $x \in (0, c]$, then the same argument as in [11, Lemma 2.4] shows that $\lambda' \geq \lambda$. In other words, λ is the best (i.e., smallest) exponential rate that can be obtained for the above theorem. However, it may be possible to obtain more precise results by considering the functional $\hat{L}_{x,y}(p) = \mathbb{E}_x[e^{-(\lambda-\lambda_*)H(y)} H(y)^{p-1}; H(y) < \infty]$ instead of L used in Proposition 8. We leave this question for future work.

A. An inhomogeneous version of Dynkin's integral identity

We make use of the following identity in the proof of Proposition 22. It is an adaptation of [27, §4, Lemma 1.2] to the context of inhomogeneous killing (or branching) rates.

Lemma 27. *Let $k: [0, \infty)^2 \rightarrow [0, \infty)$ be a function, left-differentiable in its first component, such that $k(r, t) = -k(t, r)$ and that, for $0 \leq r \leq t$, $\partial_r k(r, t)$ is independent of t . If*

$$w(r) = e^{-k(r,t)} u(t) + \int_r^t e^{-k(r,u)} v(u) du,$$

then

$$w(r) = u(t) + \int_r^t v(s) ds - \int_r^t \partial_s k(r, s) w(s) ds.$$

Acknowledgements Remark 23(iii) was kindly pointed out to us by Ronnie Loeffen. We thank an anonymous referee for their remarks which have improved several aspects of this work.

References

- [1] S. Asmussen. *Applied probability and queues*, volume 51 of *Applications of Mathematics (New York)*. Springer-Verlag, New York, second edition, 2003. ISBN 0-387-00211-1.
- [2] S. Asmussen and H. Hering. Strong limit theorems for general supercritical branching processes with applications to branching diffusions. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 36(3):195–212, 1976.
- [3] K. B. Athreya and P. E. Ney. *Branching processes*. Springer-Verlag, New York-Heidelberg, 1972. Die Grundlehren der mathematischen Wissenschaften, Band 196.
- [4] R. Azais and F. Bouguet, editors. *Statistical Inference for Piecewise-deterministic Markov Processes*. John Wiley & Sons, Ltd, 2018. ISBN 9781119507338. doi:10.1002/9781119507338.
- [5] J. Banasiak, K. Pichór, and R. Rudnicki. Asynchronous exponential growth of a general structured population model. *Acta Appl. Math.*, 119:149–166, 2012. ISSN 0167-8019. doi:10.1007/s10440-011-9666-y.
- [6] V. Bansaye, J.-F. Delmas, L. Marsalle, and V. C. Tran. Limit theorems for Markov processes indexed by continuous time Galton-Watson trees. *Ann. Appl. Probab.*, 21(6):2263–2314, 2011. ISSN 1050-5164. doi:10.1214/10-AAP757.
- [7] V. Bansaye, B. Cloez, P. Gabriel, and A. Marguet. A Non-Conservative Harris’ Ergodic Theorem. Preprint, Mar. 2019. URL <https://hal.archives-ouvertes.fr/hal-02062882>.
- [8] E. Bernard and P. Gabriel. Asynchronous exponential growth of the growth-fragmentation equation with unbounded fragmentation rate. *Journal of Evolution Equations*, 20(2):375–401, Jul 2019. ISSN 1424-3202. doi:10.1007/s00028-019-00526-4.
- [9] J. Bertoin. *Lévy processes*, volume 121 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1996. ISBN 0-521-56243-0.
- [10] J. Bertoin. Markovian growth-fragmentation processes. *Bernoulli*, 23(2):1082–1101, 2017. ISSN 1350-7265. doi:10.3150/15-BEJ770.
- [11] J. Bertoin. On a Feynman-Kac approach to growth-fragmentation semigroups and their asymptotic behaviors. *J. Funct. Anal.*, 277(11):108270, 29, 2019. ISSN 0022-1236. doi:10.1016/j.jfa.2019.06.012.

- [12] J. Bertoin and A. R. Watson. Probabilistic aspects of critical growth-fragmentation equations. *Adv. in Appl. Probab.*, 48(A):37–61, 2016. ISSN 0001-8678. doi:10.1017/apr.2016.41.
- [13] J. Bertoin and A. R. Watson. A probabilistic approach to spectral analysis of growth-fragmentation equations. *J. Funct. Anal.*, 274(8):2163–2204, 2018. ISSN 0022-1236. doi:10.1016/j.jfa.2018.01.014.
- [14] J. Bertoin and A. R. Watson. The strong Malthusian behavior of growth-fragmentation processes. *Ann. H. Lebesgue*, 3:795–823, 2020. doi:10.5802/ahl.46.
- [15] F. Campillo, N. Champagnat, and C. Fritsch. Links between deterministic and stochastic approaches for invasion in growth-fragmentation-death models. *J. Math. Biol.*, 73(6-7):1781–1821, 2016. ISSN 0303-6812. doi:10.1007/s00285-016-1012-6.
- [16] F. Campillo, N. Champagnat, and C. Fritsch. On the variations of the principal eigenvalue with respect to a parameter in growth-fragmentation models. *Communications in Mathematical Sciences*, 15(7):1801–1819, 2017. doi:10.4310/CMS.2017.v15.n7.a1.
- [17] J. A. Cañizo, P. Gabriel, and H. Yoldaş. Spectral gap for the growth-fragmentation equation via Harris’s theorem. Preprint, arXiv:2004.08343 [math.AP], 2020.
- [18] B. Cavalli. On a family of critical growth-fragmentation semigroups and refracted lévy processes. arXiv:1812.07951v2 [math.PR], 2019.
- [19] N. Champagnat and D. Villemonais. Exponential convergence to quasi-stationary distribution and Q-process. *Probability Theory and Related Fields*, 164(1):243–283, 2016. doi:10.1007/s00440-014-0611-7.
- [20] B. Cloez, R. Dessalles, A. Genadot, F. Malrieu, A. Marguet, and R. Yvinec. Probabilistic and piecewise deterministic models in biology. In *Journées MAS 2016 de la SMAI—Phénomènes complexes et hétérogènes*, volume 60 of *ESAIM Proc. Surveys*, pages 225–245. EDP Sci., Les Ulis, 2017. doi:10.1051/proc/201760225.
- [21] A. Cox, S. C. Harris, E. L. Horton, and A. E. Kyprianou. Multi-species neutron transport equation. *Journal of Statistical Physics*, pages 1–31, 2018.
- [22] B. Dadoun. Asymptotics of self-similar growth-fragmentation processes. *Electron. J. Probab.*, 22:30 pp., 2017. doi:10.1214/17-EJP45.
- [23] M. H. A. Davis. *Markov models and optimization*, volume 49 of *Monographs on Statistics and Applied Probability*. Chapman & Hall, London, 1993. ISBN 0-412-31410-X. doi:10.1007/978-1-4899-4483-2.
- [24] T. Debiec, M. Doumic, P. Gwiazda, and E. Wiedemann. Relative Entropy Method for Measure Solutions of the Growth-Fragmentation Equation. *SIAM Journal on Applied Mathematics*, 50(6):5811–5824, Nov. 2018.

- [25] C. Dellacherie, B. Maisonneuve, and P.-A. Meyer. *Probabilités et potentiel, chapitres XVII à XXIV*. Hermann, 1992.
- [26] N. Dunford and J. T. Schwartz. *Linear Operators. I. General Theory*. With the assistance of W. G. Bade and R. G. Bartle. Pure and Applied Mathematics, Vol. 7. Interscience Publishers, Inc., New York; Interscience Publishers, Ltd., London, 1958.
- [27] E. B. Dynkin. *Diffusions, superdiffusions and partial differential equations*, volume 50 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, 2002.
- [28] M. Eckhoff, A. E. Kyprianou, and M. Winkel. Spines, skeletons and the strong law of large numbers for superdiffusions. *Ann. Probab.*, 43(5):2545–2610, 2015. ISSN 0091-1798. doi:10.1214/14-AOP944.
- [29] J. Engländer, S. C. Harris, and A. E. Kyprianou. Strong law of large numbers for branching diffusions. *Ann. Inst. Henri Poincaré Probab. Stat.*, 46(1):279–298, 2010. ISSN 0246-0203. doi:10.1214/09-AIHP203.
- [30] R. K. Gettoor. Excursions of a Markov process. *Ann. Probab.*, 7(2):244–266, 1979. ISSN 0091-1798.
- [31] P.-O. Goffard and A. Sarantsev. Exponential convergence rate of ruin probabilities for level-dependent Lévy-driven risk processes. *J. Appl. Probab.*, 56(4):1244–1268, 2019. ISSN 0021-9002. doi:10.1017/jpr.2019.71.
- [32] R. Hardy and S. C. Harris. A spine approach to branching diffusions with applications to \mathcal{L}^p -convergence of martingales. In *Séminaire de Probabilités XLII*, volume 1979 of *Lecture Notes in Math.*, pages 281–330. Springer, Berlin, 2009. doi:10.1007/978-3-642-01763-6_11.
- [33] S. C. Harris, E. Horton, and A. E. Kyprianou. Stochastic methods for the neutron transport equation II: Almost sure growth. Preprint, arXiv:1901.00220v3 [math.PR], 2019.
- [34] E. Hille. *Functional analysis and semi-groups*. American mathematical society colloquium publications ; v.31. American Mathematical Society, Providence, R.I., rev. ed. edition, 1957. ISBN 0821810316.
- [35] N. Ikeda and S. Watanabe. *Stochastic differential equations and diffusion processes*, volume 24 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam; Kodansha, Ltd., Tokyo, second edition, 1989. ISBN 0-444-87378-3.
- [36] A. E. Kyprianou. *Fluctuations of Lévy processes with applications*. Springer, second edition, 2014. ISBN 978-3-642-37631-3. doi:10.1007/978-3-642-37632-0.
- [37] P. Maillard and E. Paquette. Interval fragmentations with choice: equidistribution and the evolution of tagged fragments. Preprint, arXiv:2006.16932 [math.PR], 2020.

- [38] A. Marguet. Uniform sampling in a structured branching population. *Bernoulli*, 25 (4A):2649–2695, 2019. ISSN 1350-7265. doi:10.3150/18-BEJ1066.
- [39] A. Marguet and C. Smadi. Parasite infection in a cell population with deaths. Preprint, arXiv:2010.16070v1 [math.PR], 2020.
- [40] P. Michel, S. Mischler, and B. Perthame. General relative entropy inequality: an illustration on growth models. *Journal de Mathématiques Pures et Appliquées*, 84 (9):1235 – 1260, 2005. ISSN 0021-7824. doi:10.1016/j.matpur.2005.04.001.
- [41] S. Mischler and J. Scher. Spectral analysis of semigroups and growth-fragmentation equations. *Annales de l’Institut Henri Poincaré (C) Non Linear Analysis*, 33(3): 849–898, June 2016.
- [42] B. Perthame. *Transport equations in biology*. Frontiers in Mathematics. Birkhäuser Verlag, Basel, 2007. ISBN 978-3-7643-7841-7; 3-7643-7841-7.
- [43] D. Revuz and M. Yor. *Continuous martingales and Brownian motion*, volume 293 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, third edition, 1999. ISBN 3-540-64325-7.
- [44] H. T. K. Tse, W. M. Weaver, and D. Di Carlo. Increased asymmetric and multi-daughter cell division in mechanically confined microenvironments. *PLOS ONE*, 7 (6):1–8, 06 2012. doi:10.1371/journal.pone.0038986.
- [45] I. Vidav. Existence and uniqueness of nonnegative eigenfunctions of the boltzmann operator. *Journal of Mathematical Analysis and Applications*, 22(1):144–155, 1968. ISSN 0022-247X.
- [46] J. Voigt. On the perturbation theory for strongly continuous semigroups. *Mathematische Annalen*, 229(2):1432–1807, 1977.