

1 Isolation and characterization of 50 microsatellite loci for two
2 shearwater species, *Ardenna pacifica* and *Puffinus bailloni*

3

4 Helena Teixeira ^{1,*}, Matthieu Le Corre ¹, Audrey Jaeger ¹, Arthur Choeur ¹, Merlène Saunier ¹,
5 François-Xavier Couzi ², Vikash Tatayah ³, Nirmal Jivan Shah ⁴, Malcolm A C Nicoll ⁵, Naïs
6 Avargues ¹, Laurence Humeau ⁶

7 ¹ UMR ENTROPIE (Université de La Réunion, IRD, CNRS, IFREMER, Université de Nouvelle-
8 Calédonie), 15 Avenue René Cassin, CS 92003, 97744 Saint Denis Cedex 9, Ile de La Réunion,
9 France.

10 ² Société d'Etudes Ornithologiques de La Réunion (SEOR), 13 ruelle des Orchidées, 97440, Saint
11 André, Réunion Island, France

12 ³ Mauritian Wildlife Foundation, Grannum Road, Vacoas, Mauritius, Indian Ocean

13 ⁴ Nature Seychelles, Roche Caiman, Mahe, Seychelles

14 ⁵ Institute of Zoology, Zoological Society of London, Regent's Park, London, NW1 4RY, UK

15 ⁶ UMR PVBMT (Université de La Réunion, CIRAD), 15 Avenue René Cassin, CS 92003, 97744
16 Saint Denis Cecodex 9, Ile de La Réunion, France

17

18 *corresponding author: helena-marisa.osorio-teixeira@univ-reunion.fr (HT)

19

20 Helena Teixeira: <https://orcid.org/0000-0001-9032-4575>

21 Matthieu Le Corre: <https://orcid.org/0000-0002-1194-351X>

22 Audrey Jaeger: <https://orcid.org/0000-0002-5649-0315>

23 Arthur Choeur: <https://orcid.org/0000-0002-1651-3797>

24 Merlène Saunier: <https://orcid.org/0000-0002-9654-8585>

25 François-Xavier Couzi: <https://orcid.org/0000-0001-8569-2541>

26 Vikash Tatayah: <https://orcid.org/0000-0003-0759-254X>

27 Nirmal Jivan: Shah <https://orcid.org/0000-0003-2114-4786>

28 Malcolm Nicoll: 0000-0001-6212-4787

29 Naïs Avargues: <https://orcid.org/0000-0003-0432-8436>

30 Laurence Humeau: <https://orcid.org/0000-0002-3690-2159>

31

32 **Abstract**

33 **Background:** Shearwaters (order Procellariiformes) are an excellent study system to investigate
34 the genetic consequences of the co-called “seabird paradox”, as they are able to disperse long
35 distances but many species exhibit natal and breeding philopatry. However, few microsatellite
36 markers are currently available for these taxa, hampering genetic inferences.

37

38 **Methods and Results:** In this study, twenty-five novel microsatellite loci were isolated and
39 characterized for each of two distantly related shearwater species: the wedge-tailed shearwater
40 (*Ardenna pacifica*) and the tropical shearwater (*Puffinus bailloni*). Polymorphism tests were
41 performed for a total of 91 *A. pacifica* individuals sampled at Reunion and Round Island, and 48
42 *P. bailloni* individuals from Reunion and Europa Island, in the western Indian Ocean. The analyses
43 revealed twenty-three polymorphic loci for *A. pacifica*, with the number of alleles per locus (N_a)
44 ranging from two to eight (mean = 3.957 ± 0.364). Nineteen polymorphic loci were found for *P.*
45 *bailloni*, with N_a varying from two to five (mean = 3.053 ± 0.247). The observed heterozygosity
46 (H_o) was relatively low for the two species, with H_o ranging from 0.022 to 0.725 (mean = $0.326 \pm$
47 0.044) for *A. pacifica* and from 0.021 to 0.688 (mean = 0.271 ± 0.051) for *P. bailloni*, but
48 comparable to the estimates available for other *Puffinus* species.

49

50 **Conclusions:** The new microsatellite loci provide a valuable tool for further population genetic
51 studies, and will allow for design of effective conservation and management plans for *A. pacifica*,
52 *P. bailloni* and other closely-related species.

53

54 **Keywords:** seabirds, wedge-tailed shearwater, tropical shearwater, microsatellites, population
55 genetics, conservation.

56

57 **Introduction**

58 Seabirds, one of the most diverse group of avifauna, are distributed across all oceans
59 and marine ecosystems of the world, and are renowned for their ability to migrate long distances
60 [1]. Shearwaters (order Procellariiformes) are highly pelagic seabirds, which undertake long-
61 distance non-breeding migrations and large-scale movements when central placed foragers
62 during breeding [2]. In contrast, many shearwater species exhibit natal philopatry and return
63 repeatedly to the same colony for breeding, often close to their natal nest site. This is often
64 described as the “seabird paradox” [3]. Although this behavior may increase the probability that
65 adequate resources and mates are available for reproduction [4], it may also result in limited gene
66 flow between colonies, high levels of endemism, and in an increased risk of inbreeding depression
67 and vulnerability to stochastic demographic fluctuations [5].

68 The taxonomic history of shearwaters has been revised repeatedly [6–8]. A recent study
69 using genomic data from 25 of the 32 recognized shearwaters species provided the first well-
70 resolved phylogeny for this taxa [9]. The study supported the monophyly of the three genera of
71 shearwaters, *Ardenna*, *Calonectris* and *Puffinus*, and positioned *Ardenna* and *Calonectris* as
72 sister genera [9]. The three genera exhibit differences in species richness [10], and are an ideal
73 study system to investigate the genetic processes underlying species diversification in marine
74 ecosystems. However, molecular nuclear markers are still unavailable for most of the shearwaters
75 species (but see [11–13]), which limits our ability to explore these genetic processes.

76 In this study, we identify and characterize new polymorphic microsatellite loci for two
77 distantly related shearwater species: the wedge-tailed shearwater (*Ardenna pacifica*, Gmelin
78 1789) and tropical shearwater (*Puffinus bailloni*, Bonaparte 1857). These microsatellite markers
79 will provide a valuable tool for population genetic studies and allow researchers to identify
80 conservation or management units, made up of one or more connected breeding colonies, and
81 hence develop effective conservation actions for *A. pacifica* and *P. bailloni*, and potentially for
82 other closely-related species.

83

84 **Materials and methods**

85 *A. pacifica* ranges throughout the tropical and subtropical waters of the Indian and Pacific
86 Oceans (latitudes 35°N and 35°S) and is known to breed on a large number of oceanic islands

87 and on the east and west coast of Australia [14]. In contrast, the congener *P. bailloni* is widely
88 distributed in the western tropical Indian and Pacific Oceans. In the western Indian Ocean the
89 main colonies are in the Seychelles, the Chagos Archipelago, Reunion and Europa Island [15].
90 For the present study, biological samples were collected in two study sites for each species
91 between 2009 and 2019. *A. pacifica* individuals were sampled in Reunion (21°70 S, 55°31 E) and
92 Round Island (Mauritius; 19°51 S, 57°47 E), in the western Indian Ocean. *P. bailloni* were sampled
93 in Reunion and Europa Island (southern Mozambique channel; 21°21 S, 40°22 E). Two distinct
94 breeding colonies were selected for each species to allow for potential variation in microsatellite
95 genotypes. Shearwaters were captured at breeding colonies by hand. Additionally, samples from
96 all *P. bailloni* from Reunion and some *A. pacifica* were collected from dead birds, fatally injured
97 as a consequence of light pollution [16]. Blood samples (approx. 0.5 ml) were collected from live
98 birds from the medial metatarsal or basilic veins, while muscle samples (approx. 10 g) were
99 collected from dead animals. All samples were preserved in 70% ethanol. A total of 91 *A. pacifica*
100 (55 from Reunion and 36 from Round Island) and 48 *P. bailloni* (30 from Reunion and 18 from
101 Europa Island) samples were available for genetic analyses. Additionally, three *A. pacifica* and
102 five *P. bailloni* blood samples from Cousin Island (Seychelles; 4°20 S, 55°40 E) were available
103 from previous field expeditions and included in the microsatellite library.

104 A microsatellite library was designed and established for each shearwater species by
105 GenoScreen (Lille, France). Genomic DNA was extracted from whole blood and muscle samples
106 using the NucleoSpin Tissue kit (Macherey-Nagel), following the manufacturer's protocol. A total
107 of nine individuals for *A. pacifica* and ten for *P. bailloni* were equimolarly pooled and 1µg of DNA
108 was used for the development of the genomic library. The libraries were analyzed on an Illumina
109 MiSeq platform to generate 250bp paired-end reads using the MiSeq Reagent Nano Kit v2. A
110 total of 1,128,338 reads were obtained for *A. pacifica* and 1,211,564 for *P. bailloni*. The resulting
111 sequences were assembled using Usearch software, producing 131,356 contigs for *A. pacifica*
112 and 161,218 contigs for *P. bailloni*. The bioinformatics program QDD v3 [17] was used to analyze
113 the sequences. All bioinformatic steps from the raw sequences until obtaining the primers were
114 performed using this software, including the removal of adapters/vectors, the detection of
115 microsatellites, the detection of redundancy/possible mobile element association, the selection of
116 sequences with target microsatellites, and the primer design using BLAST, ClustalW and Primer3

117 software. A total of 2,524 primer sets were finally designed on suitable microsatellite flanking
118 regions for *A. pacifica* and 3,102 for *P. bailloni*. Of these, 421 primer pairs were selected for *A.*
119 *pacifica* and 507 for *P. bailloni*, by keeping only perfect di/tri/tetra motives, with A and B quality
120 design (from internal parameters of QDD), and with at least 20bp between primer and
121 microsatellite. Finally, 95 microsatellite loci were randomly selected for preliminary molecular
122 tests for each shearwater, using DNA of eight individuals. Thereafter, a subset of 48 microsatellite
123 loci were tested for polymorphism for each species, using DNA from an additional fifteen
124 individuals (2 from Round Island, 10 from Reunion and 3 from Cousin Island for *A. pacifica*, and
125 one from Europa Island, 9 from Reunion and 5 from Cousin Island for *P. bailloni*). Polymerase
126 Chain Reaction (PCR) was performed for each microsatellite locus in a 10 µl reaction containing
127 0.5 U of TAQ DNA polymerase, 6 pmol of dNTP, 37.5 pmol of MgCl₂, 10 pmol of each forward
128 and reverse primer, and 1 µl (~ 10 ng) of DNA template. PCR amplifications were carried out
129 using the following thermal conditions: initial denaturation step at 95 °C for 15 min, followed by 40
130 cycles of 95 °C for 30 s, 55 °C for 30 s, and 72 °C for 60 s, and a final extension step at 72 °C for
131 10 min. PCR products were separated on an ABI 3730XL DNA analyzer (Applied Biosystems)
132 and sized with GeneScan-500 LIZ size standard. All molecular analyses were performed by
133 GenoScreen (Lille, France).

134 Microsatellite alleles were scored using GeneMapper® v4.0 (Applied Biosystems). The
135 twenty-five best microsatellite loci for each study species (di- and trinucleotide repeats) were then
136 selected for multiplex development, based on the allelic diversity, fragment size and peak patterns
137 in the electropherograms. Five multiplex reactions were developed for each shearwater species
138 with the help of the Multiplex Manager v1.2 [18]. Each multiplex included primers that amplify
139 fragments of different size and were labelled with four fluorescent dyes (6-FAM, PET, VIC, NED)
140 to enable fragment analysis multiplexing. Multiplex conditions were optimized using seven
141 individuals per study species (5 individuals from Reunion and 2 from Cousin Island for *A. pacifica*,
142 and 7 individuals from Reunion for *P. bailloni*). PCR were subsequently performed for the entire
143 *A. pacifica* (n = 91) and *P. bailloni* (n = 48) dataset, considering the same conditions and thermal
144 cycle as for the simplex PCR, but with an uneven quantity of primer per multiplex (see Table 1 for
145 details).

146 Evidence of null alleles, large-allele dropout, and stutter bands were examined for each
147 locus with MicroChecker 2.2.3 [19]. Genepop 4.7.5 [20, 21] was used to test for deviations from
148 Hardy–Weinberg equilibrium (HWE) for each locus, and to test for linkage disequilibrium for each
149 locus-pair combination. The p-values were adjusted using the Benjamini and Yekutieli correction
150 method for multiple comparisons [22, 23]. The mean observed number of alleles per locus (N_a),
151 observed heterozygosity (H_o), and unbiased expected heterozygosity (uH_e) estimated according
152 to Nei [24] were computed using GenAEx 6.5 [25]. To ensure that the set of microsatellite loci
153 could reliably identify unique genotypes for the two species, the probability of identity (pID ; [26])
154 was computed using GenAEx 6.5 [25].

155

156 **Results and discussion**

157 The initial microsatellite primer screening revealed twenty-three polymorphic loci for *A.*
158 *pacifica* (Table 2) and nineteen for *P. bailloni* (Table 3), which amplified a total of 91 and 58 alleles
159 for each study species. Despite the uneven sample sizes ($n = 91$ for *A. pacifica* and $n = 48$ for *P.*
160 *bailloni*), the number of alleles per locus (N_a), and observed (H_o) and expected unbiased (uH_e)
161 heterozygosities were comparable for the two shearwater species. The N_a for *A. pacifica* ranged
162 from two to eight (mean = 3.957 ± 0.364), and the N_a for *P. bailloni* from two to five (mean = 3.053
163 ± 0.247). For *A. pacifica*, the H_o varied from 0.022 to 0.725 (mean = 0.326 ± 0.044) and the uH_e
164 from 0.022 to 0.730 (mean = 0.343 ± 0.044 ; Table 2). Likewise, the H_o for *P. bailloni* ranged from
165 0.021 to 0.688 (mean = 0.271 ± 0.051) and the uH_e from 0.021 to 0.709 (mean = 0.298 ± 0.053 ;
166 Table 3). No heterozygote deficit was detected when combining the two study sites for each
167 shearwater species (i.e.; global F_{IS} was not significantly different from zero for both *A. pacifica*
168 and *P. bailloni*), suggesting that we did not violate the assumption of population panmixia in the
169 present study.

170 No large-allele dropout and stutter peaks were detected for the two shearwaters.
171 However, two loci (Pb72 and Pb83) exhibited null alleles for *P. bailloni* and three loci (Ap19, Ap23,
172 Ap36) for *A. pacifica*. No significant linkage disequilibrium was detected among any locus-pairs
173 for the two study species, after correcting for multiple tests comparison. Significant deviations
174 from HWE were found for three loci (Ap19, Ap21, Ap36) for *A. pacifica*, after Benjamini and
175 Yekutieli correction (p -value > 0.05). All the loci developed for *P. bailloni* were in HWE (Table 2

176 and Table 3). The probability of identity (pID) using the present set of loci was 9.27×10^{-9} for *A.*
177 *pacifica* and 3.01×10^{-6} for *P. bailloni*, suggesting that the probability of two unrelated individuals
178 drawn randomly from our dataset sharing the same multilocus genotype is approximately zero.

179 Overall, the genetic diversity indices for *A. pacifica* and *P. bailloni* were comparable to
180 previous estimates for other *Puffinus* species (e.g., $H_o = 0.377 \pm 0.241$ for *Puffinus mauretanicus*
181 [11]; $H_o = 0.436 \pm 0.257$ for *Puffinus yelkouan* [11]; $H_o = 0.549 \pm 0.175$ for *Puffinus carneipes*
182 [12]), but lower than those reported for other local seabirds using microsatellites datasets (e.g.,
183 *Onychoprion fuscatus* and *Pseudobulweria aterrima*; [27, 28]).

184 The deficit of heterozygotes observed in *P. bailloni* may reflect the smaller distribution of
185 the species and the potential isolation of the colonies, as *Puffinus* are known to be extremely
186 philopatric [11]. The results are, however, remarkable for the widely-distributed and abundant *A.*
187 *pacifica*. It is important to emphasize that the polymorphism of each 25 microsatellite loci was
188 evaluated with individuals from only two study sites per species. Microsatellite loci may exhibit
189 new alleles when genotyping individuals from other geographic areas and, consequently, display
190 a higher genetic variance than revealed in this study.

191 Altogether, the newly developed microsatellite loci provide a valuable tool to investigate
192 genetic diversity, population genetic structure, population connectivity and reconstruct the
193 demographic history of *A. pacifica* and *P. bailloni* at a large geographic scale. Although the new
194 primer pairs have not been tested for cross-amplification in both of our study species or in other
195 seabird species, there is potential application of these markers in other closely related species,
196 for which genetic studies are a priority and microsatellite markers are not yet available (e.g., [27,
197 29]). Future molecular studies are needed to investigate the cross-species transferability and
198 genetic variability of the newly described microsatellites loci in other shearwater species.

199

200 **References**

- 201 1. Egevang C, Stenhouse IJ, Phillips RA, et al (2010) Tracking of Arctic terns *Sterna paradisaea*
202 reveals longest animal migration. *Proc Natl Acad Sci* 107:2078–2081.
203 <https://doi.org/https://doi.org/10.1073/pnas.0909493107>
- 204 2. Weimerskirch H, de Grissac S, Ravache A, et al (2020) At-sea movements of wedge-tailed
205 shearwaters during and outside the breeding season from four colonies in New Caledonia. *Mar Ecol*
206 *Prog Ser* 633:225–238. <https://doi.org/https://doi.org/10.3354/meps13171>
- 207 3. Antaky CC, Young L, Ringma J, Price MR (2021) Dispersal under the seabird paradox: Probability,
208 life history, or spatial attributes? *Mar Ornithol* 49:1–8.
- 209 4. Morris-Pocock JA, Anderson DJ, Friesen VL (2016) Biogeographical barriers to dispersal and rare
210 gene flow shape population genetic structure in red-footed boobies (*Sula sula*). *J Biogeogr* 43:2125–

211 2135. <https://doi.org/10.1111/jbi.12780>

212 5. Frankham R, Briscoe DA, Ballou JD (2002) Introduction to conservation genetics. Cambridge
213 university press

214 6. Austin JJ, Bretagnolle V, Pasquet E (2004) A global molecular phylogeny of the small *Puffinus*
215 shearwaters and implications for systematics of the Little-Audubon's Shearwater complex. *Auk*
216 121:847–864. <https://doi.org/https://doi.org/10.1093/auk/121.3.847>

217 7. Penhallurick J, Wink M (2004) Analysis of the taxonomy and nomenclature of the Procellariiformes
218 based on complete nucleotide sequences of the mitochondrial cytochrome b gene. *Emu* 104:125–
219 147. <https://doi.org/https://doi.org/10.1071/MU01060>

220 8. Torres L, Pante E, González-Solís J, et al (2021) Sea surface temperature, rather than land mass
221 or geographic distance, may drive genetic differentiation in a species complex of highly dispersive
222 seabirds. *Ecol Evol* 11:14960–14976. <https://doi.org/10.1002/ece3.8180>

223 9. Ferrer Obiol J, James HF, Chesser RT, et al (2021) Integrating Sequence Capture and Restriction
224 Site-Associated DNA Sequencing to Resolve Recent Radiations of Pelagic Seabirds. *Syst Biol*
225 70:976–996. <https://doi.org/10.1093/sysbio/syaa101>

226 10. Ferrer Obiol J, James HF, Chesser RT, et al (2022) Palaeoceanographic changes in the late
227 Pliocene promoted rapid diversification in pelagic seabirds. *J Biogeogr* 49:171–188.
228 <https://doi.org/10.1111/jbi.14291>

229 11. Gonzalez EG, Genovart M, Oro D, et al (2009) Polymorphic microsatellite markers for the critically
230 endangered Balearic shearwater, *Puffinus mauretanicus*. *Mol Ecol Resour* 9:1044–1046.
231 <https://doi.org/https://doi.org/10.1111/j.1755-0998.2009.02563.x>

232 12. Hardesty BD, Metcalfe SS, Wilcox C (2013) Genetic variability and population diversity as revealed
233 by microsatellites for Flesh-footed shearwaters (*Puffinus carneipes*) in the southern hemisphere.
234 *Conserv Genet Resour* 5:27–29. <https://doi.org/10.1007/s12686-012-9721-7>

235 13. Herman RW, Winger BM, Dittmann DL, Harvey MG (2022) Fine-scale population genetic structure
236 and barriers to gene flow in a widespread seabird (*Ardenna pacifica*). 1–12.
237 <https://doi.org/10.1093/biolinnean/blac091>

238 14. Del Hoyo J, Del Hoyo J, Elliott A, Sargatal J (1992) Handbook of the birds of the world. Lynx edicions
239 Barcelona

240 15. Kirwan, G. M., J. del Hoyo and NC (2020) Tropical Shearwater (*Puffinus bailloni*), version 1.0. In
241 Birds of the World. Cornell Lab of Ornithology, Ithaca, NY, USA.
242 <https://doi.org/10.2173/bow.troshe5.01>

243 16. Chevillon L, Tourmetz J, Dubos J, Soulaïmana-Mattoir Y, Hollinger C, Pinet P, Couzi FX ,
244 Riethmuller M LCM (2022). 25 years of light-induced petrel groundings at Reunion Island:
245 retrospective analysis and predicted trend. *Global Ecology and Conservation*.
246 <https://doi.org/10.1016/j.gecco.2022.e02232>

247 17. Megléc E, Costedoat C, Dubut V, et al (2010) QDD: a user-friendly program to select microsatellite
248 markers and design primers from large sequencing projects. *Bioinformatics* 26:403–404.
249 <https://doi.org/10.1093/bioinformatics/btp670>

250 18. Holleley CE, Geerts PG (2009) Multiplex Manager 1.0: a cross-platform computer program that plans
251 and optimizes multiplex PCR. *Biotechniques* 46:511–517.
252 <https://doi.org/https://doi.org/10.2144/000113156>

253 19. Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P (2004) MICRO-CHECKER: Software for
254 identifying and correcting genotyping errors in microsatellite data. *Mol Ecol Notes* 4:535–538.
255 <https://doi.org/10.1111/j.1471-8286.2004.00684.x>

256 20. Rousset F (2008) genepop'007: a complete re-implementation of the genepop software for Windows
257 and Linux. *Mol Ecol Resour* 8:103–106. <https://doi.org/https://doi.org/10.1111/j.1471-8286.2007.01931.x>

259 21. Raymond M (1995) Population genetics software for exact test and ecumenicism. *J Hered* 86:248–
260 249

261 22. Benjamini Y, Yekutieli D (2001) The control of the false discovery rate in multiple testing under
262 dependency. *Ann Stat* 1165–1188

263 23. Narum SR (2006) Beyond Bonferroni: less conservative analyses for conservation genetics.
264 *Conserv Genet* 7:783–787. <https://doi.org/https://doi.org/10.1007/s10592-005-9056-y>

265 24. Nei M (1978) Estimation of average heterozygosity and genetic distance from a small number of
266 individuals. *Genetics* 89:583–590. <https://doi.org/https://doi.org/10.1093/genetics/89.3.583>

267 25. Peakall ROD, Smouse PE (2006) GENALEX 6: genetic analysis in Excel. Population genetic

268 software for teaching and research. Mol Ecol Notes 6:288–295.
 269 <https://doi.org/https://doi.org/10.1111/j.1471-8286.2005.01155.x>

270 26. Taberlet P, Luikart G (1999) Non-invasive genetic sampling and individual identification. Biol J Linn
 271 Soc 68:41–55. <https://doi.org/https://doi.org/10.1111/j.1095-8312.1999.tb01157.x>

272 27. Danckwerts DK, Lebarbenchon C, Le Corre M, Humeau L (2019) Isolation and characterisation of
 273 16 polymorphic microsatellite loci for the sooty tern (*Onychoprion fuscatus*; Sternidae), a super-
 274 abundant pan-tropical seabird, including a test of cross-species amplification using two noddies
 275 (*Anous spp.*). Mar Biodivers 49:509–513. <https://doi.org/10.1007/s12526-017-0773-5>

276 28. Lopez J, Nikolic N, Riethmuller M, et al (2021) High genetic diversity despite drastic bottleneck in a
 277 critically endangered, long-lived seabird, the Mascarene Petrel *Pseudobulweria aterrima*. Ibis (Lond
 278 1859) 163:268–273. <https://doi.org/10.1111/ibi.12864>

279 29. Techow NMSM, O’Ryan C (2004) Characterization of microsatellite loci in White-chinned Petrel
 280 (*Procellaria aequinoctialis*) and cross-amplification in six other procellariiform species. Mol Ecol
 281 Notes 4:33–35. <https://doi.org/10.1046/j.1471-8286.2003.00560.x>

282

283 **Statements & Declarations:**

284

285 **Funding:** The research was supported by the FEDER Smac (2020-2022, N° RE0022954) and
286 the Bertarelli Foundation as part of the Bertarelli Programme in Marine Science (GRANT
287 AGREEMENT REUNION-2021-822916). Field work at Europa and Round Island was funded by
288 the Consortium Iles Eparses (ECOMIE programme) and the Pew Fellowship Award in Marine
289 Conservation of MLC, respectively. Field work in the Seychelles was conducted under the Project
290 “Wild Fauna Indian Ocean” (NRDC/0293). Arthur Choer and Merlène Saunier were supported
291 by Region Reunion, and European Social Fund (ESF) PhD studentships.

292

293 **Competing Interests:** The authors have no relevant financial or non-financial interests to
294 disclose.

295

296 **Author Contributions:** The study was conceptualized and designed by Laurence Humeau and
297 Matthieu Le Corre. Field work was organized by Matthieu Le Corre, Vikash Tatayah, Nirmal Jivan
298 Shah and Malcolm Nicoll. Data collection were performed by Matthieu Le Corre, Audrey Jaeger,
299 Arthur Choer, Merlène Saunier, François-Xavier Couzi, Vikash Tatayah, and Naïs Avargues.
300 Data analysis was performed by Helena Teixeira and Laurence Humeau. The first draft of the
301 manuscript was written by Helena Teixeira and all authors revised the manuscript. All authors
302 read and approved the final manuscript.

303

304 **Ethics approval:** Bird capture, handling and sample collection were approved and carried out in
305 concordance with the principles of the Centre de Recherche sur la Biologie des Populations
306 d'Oiseaux (PP 616 and banding authorization 44 of MLC; CRBPO, National Museum of Natural
307 History, Paris). In addition, sample collection in Europa was conducted under the approval of the
308 Terres Australes et Antarctiques Françaises (TAAF). Sample collection and exportation permits
309 at Round Island were granted by the National Parks and Conservation Service (NPCS)
310 (Government of Mauritius). Sample collection and exportation permit at Cousin Island were
311 granted by the Seychelles Bureau of Standards.

312

313 **Data availability:** Sequences of the twenty-five microsatellites markers developed for *Ardenna*
314 *pacifica* and *Puffinus bailloni* are publicly available in GenBank under the accession numbers
315 ON854684 – ON854733.

316

317

318 **Acknowledgments:**

319 The authors are very grateful to Patrick Pinet for assistance and logistical support during sample
320 collection at Round Island. We also thank Lucie Gauchet, Céline Toty, Mickael Baumann, Nicolas
321 Ligerica and seven anonymous military for their help in the field at Europa Island. We are also
322 grateful to the Conservatoire du Littoral Antenne Océan Indien for authorization to conduct field
323 work at Grande Anse, Reunion, the TAAF for field work assistance at Europa, and NPCS for field
324 work at Round Island. We thank Helena Fontaine and Gwennaïs Fustemberg for double-checking
325 the genotypes. Finally, we are very grateful to the two anonymous reviewers whose comments
326 improved the quality of our manuscript.

327 **Table 1.** Characteristics of the five multiplexes developed for *Ardenna pacifica* and *Puffinus*
 328 *bailloni*.

<i>Ardenna pacifica</i>				<i>Puffinus bailloni</i>			
Locus	Multiplex	Dye	Quantity (pmol)	Locus	Multiplex	Dye	Quantity (pmol)
Ap19	M1	6-FAM	1.6	Pb53	M1	6-FAM	5.2
Ap83	M1	VIC	0.8	Pb26	M1	6-FAM	2.6
Ap06	M1	VIC	2.5	Pb80	M1	VIC	1.2
Ap26	M1	NED	1.0	Pb22	M1	NED	1.8
Ap58	M1	PET	2.5	Pb56	M1	PET	1.5
Ap07	M2	6-FAM	2.6	Pb06	M2	6-FAM	1.6
Ap03	M2	VIC	0.8	Pb43	M2	VIC	5.2
Ap08	M2	VIC	3.0	Pb05	M2	VIC	3.4
Ap73	M2	NED	1.4	Pb28	M2	NED	1.0
Ap92	M2	PET	4.0	Pb40	M2	PET	2.2
Ap21	M3	6-FAM	2.0	Pb46	M3	6-FAM	2.0
Ap23	M3	VIC	1.0	Pb72	M3	VIC	5.8
Ap32	M3	NED	0.8	Pb90	M3	NED	0.8
Ap36	M3	PET	5.6	Pb48	M3	NED	2.0
Ap09	M3	PET	5.2	Pb66	M3	PET	2.6
Ap48	M4	6-FAM	1.6	Pb84	M4	6-FAM	2.0
Ap15	M4	6-FAM	5.0	Pb16	M4	VIC	1.6
Ap29	M4	VIC	1.6	Pb64	M4	NED	1.0
Ap79	M4	PET	1.4	Pb83	M4	NED	1.0
Ap57	M4	PET	3.0	Pb93	M4	PET	1.6
Ap25	M5	6-FAM	3.0	Pb55	M5	6-FAM	1.6
Ap44	M5	6-FAM	8.0	Pb63	M5	6-FAM	5.4
Ap27	M5	NED	2.0	Pb21	M5	VIC	1.4
Ap37	M5	NED	6.4	Pb92	M5	NED	1.2
Ap10	M5	PET	1.4	Pb82	M5	PET	1.6

329 Multiplex, multiplex mix in which the locus was amplified; Dye, fluorescent label of the microsatellite primers;

330 Quantity, Quantity of primers used per PCR reaction.

331

332 **Table 2.** Results of the initial microsatellite primer screening in *Ardenna pacifica*. Analyses were performed using 91 individuals from Reunion and Round
 333 (Mauritius) Island.

Locus	Accession no.	Primer sequences (5' – 3')	Repeat motif	Fragment size (bp)	N	Na	Ho	uHe	HWE	Null alleles
Ap19	ON854691	F: ACCAGCCTGCAGATTTAGGA R: TGAAACAGGCAAATGAGCGC	(AC)10	140 - 142	91	2	0.088	0.161	0.010**	Yes
Ap83	ON854707	F: GGGTGCAAAGGACCTTCAGT R: AGGATCTTTGTTCCCTCAAGCA	(AC)7	200 - 204	90	3	0.067	0.065	1.000	No
Ap06	ON854685	F: GTGAGGTGTTCTCCAGGAGG R: GAGAAGTGAGGAGATGGGCT	(AG)13	301 - 321	91	7	0.495	0.562	0.945	No
Ap26	ON854695	F: ACAGAGGACAAAGCAAATATTAGCC R: AACTGAATTGTTTGTACAGGCC	(AG)9	197 - 203	91	3	0.495	0.475	1.000	No
Ap58	ON854704	F: TCCATGGATTTGTTACAGGAGT R: ACAGAGAATGCCTGACTTTGGT	(AGC)8	204 - 207	91	2	0.352	0.357	1.000	No
Ap07	ON854686	F: ACCTGACCAACTTCGAAGCA R: GCATCTGTGCCAGTGAGATT	(AG)13	267 - 275	87	5	0.678	0.638	1.000	No
Ap03	ON854684	F: AGTGCCTACGCCATCATAGC R: TGGAGATTTGCACCCTTCCC	(AAC)16	168 - 189	91	8	0.725	0.730	1.000	No
Ap08	ON854687	F: ACTGTGAAGTGAGCACTGAGA R: GGAACACCCTACAGCAGATCA	(AC)12	238 - 248	91	5	0.341	0.295	1.000	No
Ap73	ON854705	F: AGTAGAGGAATATGCACCTGGT R: CGCTTCCGTGTACTTCTTCC	(AG)8	218	91	1	—	—	—	—
Ap92	ON854708	F: CCTCCTCTTAAGATTACCTACCAAT R: GCCCACCATGATAATGCGGT	(AT)7	163 - 189	82	7	0.305	0.332	0.302	No
Ap21	ON854692	F: TCCTCCCAGATCTTCTCCGG R: AGAATTGCGAGCCTGGATCC	(AC)10	231 - 239	91	5	0.505	0.453	0.000***	No
Ap23	ON854693	F: GAACAGATTATTCCTTTCCACTCTGC R: GTTGACCAGCTGGAGAGTCC	(AC)9	144 - 148	90	3	0.300	0.407	0.056	Yes

Ap32	ON854698	F: AAGCAGCTTCTAATGCAGGT R: AACCTTGCGTGTGTAGGCT	(AC)9	151 - 155	89	3	0.225	0.252	0.991	No
Ap36	ON854699	F: TTGCCTTTCTCTCTGCTTCTG R: AACTGCATAAGTATAGAACATTGTGC	(AG)8	194 - 200	91	4	0.275	0.401	0.010**	Yes
Ap09	ON854688	F: CCAAGTTTCAGTTACCTATCTGGG R: TCCTTGAGATCTTGTGAACAGT	(AC)11	295 - 299	89	3	0.056	0.076	0.465	No
Ap48	ON854702	F: TGGGTAACAGGTCTCCAGGT R: CTGGCCGAGAAAGCATACCA	(ACC)8	169	91	1	—	—	—	—
Ap15	ON854690	F: TCCTGCAACAGAACGTCTCA R: TGCTGTGACTTTACAAGAACTGG	(AG)10	295 - 299	90	3	0.267	0.292	0.485	No
Ap29	ON854697	F: CAAGCTCAGTTGAAACGGGC R: TAGACCAAGCCCTGACCCTT	(AC)9	224 - 228	90	3	0.544	0.565	0.868	No
Ap79	ON854706	F: CCTGGAGTGAGCAGATAGGTG R: GCCAAATTAAGGCCAACTGCA	(AC)7	140 - 142	90	2	0.433	0.451	1.000	No
Ap57	ON854703	F: GAGGGCAAAGGCAATTTCCGG R: GGAACACTCTTCTGGCTCCC	(AC)8	235 - 237	90	2	0.044	0.044	1.000	No
Ap25	ON854694	F: TGTCATGGCATTGAACTGCTG R: AGGAAGACTCTGTCTATATGCCT	(AC)9	184 - 194	91	4	0.154	0.166	0.945	No
Ap44	ON854701	F: TCCCATGGATAACTGCACGC R: GGCGTAAACACTGGCAACTT	(AAT)8	291 - 318	90	5	0.156	0.148	1.000	No
Ap27	ON854696	F: CTCCTTAATAGGCTATCGCAGTCC R: AGGAATTGTCTTGATTTCCGGTGT	(AC)9	243 - 253	91	6	0.670	0.713	1.000	No
Ap37	ON854700	F: AAGAGCTAGTGCAGAAGTGAC R: ACATCCTGGAGTCAGAGATGA	(AC)8	303 - 307	91	3	0.022	0.022	1.000	No
Ap10	ON854689	F: GCTCACCACCATTTATCCACC R: GCAACTGGGAGGAACACCAT	(AC)11	221 - 225	91	3	0.297	0.275	1.000	No

334 F, sequence of forward primer, R, sequence of reverse primer; Fragment size, observed allele size range; N, sample sizes per locus; Na, number of alleles per locus across
335 samples; Ho, observed heterozygosity; uHe, unbiased expected heterozygosity; HWE, p-value for departure from Hardy–Weinberg equilibrium test; Null alleles, presence (Yes)
336 or absence (No) of null alleles.

337 * p-value < 0.5, ** p-value < 0.1, *** p-value < 0.01

338

339 **Table 3.** Results of the initial microsatellite primer screening in *Puffinus bailloni*. Analyses were performed using 48 individuals from Reunion and Europa
 340 Island.

Locus	Accession no.	Primer sequences (5' - 3')	Repeat motif	Fragment size (bp)	N	Na	Ho	uHe	HWE	Null alleles
Pb53	ON854720	F: TGAATTTGGGTTGTCTTGGTCA R: CAGCTTTACGCATGCACTGT	(AT)8	244 - 248	48	3	0.458	0.424	1.000	No
Pb26	ON854714	F: GGGAAGCTGGGAAGATGAAAGA R: ACAGAGCTGATATAAGGTGCTAAA	(AC)9	286 - 296	48	4	0.271	0.259	0.440	No
Pb80	ON854727	F: TCAAGCAACACAGGGTACGG R: AGATTCTGTGCTTCTGCCCA	(AG)7	189	48	1	—	—	—	—
Pb22	ON854713	F: AGCTGGTTGGCATCTCACAA R: GTGCACAAGTCCAGCAATGG	(AC)9	248	48	1	—	—	—	—
Pb56	ON854722	F: CCCAGAGAATACCACTGACCG R: CCACTGGAGAAGGTTGCAGA	(AC)8	250 - 258	48	3	0.458	0.503	0.829	No
Pb06	ON854710	F: GGCTTCCTAGGAACACCTGA R: AGCATCCCTATCAGAATGGCC	(AC)12	153 - 161	48	5	0.125	0.121	1.000	No
Pb43	ON854717	F: TCTTTGGGTTACGGAGAATTCA R: AGACCCACAGGCTCTGATGT	(AG)8	246	48	1	—	—	—	—
Pb05	ON854709	F: GGGTTACCATGTCTGAGACCA R: TGCTCTACCTGGAATGGGA	(AC)13	289 - 301	48	5	0.688	0.666	1.000	No
Pb28	ON854715	F: AGCTGTGCTCTGTTAGGTCTC R: GCGTGCACAAAGGCTGTAAG	(AC)9	126 - 128	48	2	0.083	0.118	0.440	No
Pb40	ON854716	F: GGGTCACGTAAAGTATCTCCTAACA R: CCTAGTCTTCATGGTGCCCT	(AG)8	271 - 273	48	2	0.042	0.041	1.000	No
Pb46	ON854718	F: GCCCAGAATGCTGAAACTGT R: TGGAGATCAGTAAGTGTCTGTCA	(AG)8	144	48	1	—	—	—	—
Pb72	ON854726	F: TTCCTTCAGCTGCCTTGGTG R: TGGCTTTCAGTTTAGGTGACCA	(AC)7	281 - 303	48	3	0.021	0.062	0.188	Yes

Pb90	ON854731	F: GGAGTCAGCTGCATCTCGAG R: GTGTCTAGTTCTGGACCGC	(AC)7	221 - 223	48	2	0.021	0.021	NA	No
Pb48	ON854719	F: GGTATGCATTGCTAATGTGCCT R: TCTCCTCATTCTCTTCTCTGCA	(AC)8	296 - 300	48	3	0.500	0.509	1.000	No
Pb66	ON854725	F: ACGGACGTGAATTTCTTCCT R: GCATATGCATTGAGCTGAAGC	(AC)7	187	48	1	—	—	—	—
Pb84	ON854730	F: CAGAGGACACAGATGTTGCA R: CTGCTGTCAGTTTCTCTGGA	(AG)9	250 - 524	48	2	0.021	0.021	NA	No
Pb16	ON854711	F: TGGTTGAATGTCAGAGAAATACAGA R: TGTACTIONGACCTGCCATCGG	(AC)10	201 - 207	48	4	0.646	0.709	0.188	No
Pb64	ON854724	F: TGCCAAATTTATGTGTCTGATGT R: AGGTTGGGTCTTCTAGCACA	(AT)8	133 - 135	48	2	0.208	0.281	0.411	No
Pb83	ON854729	F: AGAAGCCAGGTTCCCAACAC R: TCCGTTTATGTTATCAGCAGATCCT	(AC)10	277 - 287	48	4	0.354	0.493	0.214	Yes
Pb93	ON854733	F: CAAGCCAGACCTTGCTGAGA R: CGTGAGGCAATTTGATAGGACC	(AG)7	239 - 243	48	2	0.021	0.021	NA	No
Pb55	ON854721	F: CCAAGTGACTGTGTCGGGTT R: TCAGCCTGACACTGAAAGTCG	(AC)8	189 - 195	48	4	0.417	0.455	0.290	No
Pb63	ON854723	F: AGTGACCAGGCTTGTGTCAC R: CTTCCACACCGTAGCAGGAG	(AC)8	272 - 274	48	2	0.354	0.468	0.414	No
Pb21	ON854712	F: GCTAAGAAGCTCCTCCAGTCT R: GGCTTGGGATACATAGGCACA	(AC)9	145 - 151	48	4	0.375	0.410	1.000	No
Pb92	ON854732	F: GCTCAGAACTGGCTAGAGGC R: AGGGATCGCGATAGATGGGT	(AG)7	300	48	1	—	—	—	—
Pb82	ON854728	F: CAGCTGGCAAGACCTTGAGA R: GATGGCAGGACACGTACCTC	(AC)11	216 - 2018	48	2	0.083	0.081	1.000	No

341 F, sequence of forward primer, R, sequence of reverse primer; Fragment size, observed allele size range; N, sample sizes per locus; Na, number of alleles per locus across
342 samples; Ho, observed heterozygosity; uHe, unbiased expected heterozygosity; HWE, p-value for departure from Hardy–Weinberg equilibrium test; Null alleles, presence (Yes)
343 or absence (No) of null alleles.

344 * p-value < 0.5, ** p-value < 0.1, *** p-value < 0.01

345