

## 1 **Abstract**

2 Microbiome-Inspired Green Infrastructure (MIGI) was recently proposed as an integrative  
3 system to promote healthy urban ecosystems, through multidisciplinary design. Specifically,  
4 MIGI is defined as nature-centric infrastructure restored and/or designed and managed to  
5 enhance health-promoting interactions between humans and environmental microbiomes,  
6 whilst sustaining microbially-mediated ecosystem functionality and resilience. MIGI also  
7 aims to stimulate a research agenda that focuses on considerations for the importance of  
8 urban environmental microbiomes. In this paper we provide details of what MIGI entails  
9 from a bioscience and biodesign perspective, highlighting the potential dual benefits for  
10 human and ecosystem health. We present ‘what is known’ about the relationship between  
11 urban microbiomes, green infrastructure and environmental factors that may affect urban  
12 ecosystem health (ecosystem functionality and resilience as well as human health). We  
13 discuss how to start operationalising the MIGI concept based on current available knowledge,  
14 and present a horizon scan of emerging and future considerations in research and practice.  
15 We conclude by highlighting challenges to the implementation of MIGI and propose a series  
16 of workshops to discuss multi-stakeholder needs and opportunities. This article will enable  
17 urban landscape managers to incorporate initial considerations for the microbiome in their  
18 development projects to promote human and ecosystem health. However, overcoming the  
19 challenges to operationalising MIGI will be essential to furthering its practical development.  
20 Although the research is in its infancy, there is considerable potential for MIGI to help  
21 deliver sustainable urban development driven by considerations for reciprocal relations  
22 between humans and the foundations of our ecosystems — the microorganisms.

23

24

## 25        **1. Introduction**

26    Microbial communities play vital roles in ecosystem processes and provisions including  
27    carbon and nutrient cycling, climate regulation, animal and plant health, and global food  
28    security <sup>1,2</sup> (Cavicchioli et al. 2019; Guerra et al. 2020). The ongoing degradation of, and  
29    climate-associated changes in microbial communities (structure, complexity and  
30    composition) pose a considerable threat to global macro-level biodiversity across the planet  
31    <sup>3,4,5</sup> (Bach et al. 2020; Greenspan et al. 2020; Tibbett et al. 2020). In parallel with these  
32    environmental concerns, noncommunicable diseases (chronic non-infectious diseases) are on  
33    the rise <sup>6,7</sup> (Smith et al. 2014; Jairath et al. 2020). For example, in recent decades the  
34    prevalence of asthma <sup>8,9</sup> (El-Gamal et al. 2017; Borna et al. 2019), diabetes <sup>10</sup> (Holman et al.  
35    2010), allergic rhinoconjunctivitis <sup>11</sup> (Kainu et al. 2013), and autoimmune disorders <sup>12,13</sup>  
36    (Dinse et al. 2020; Paramasivan et al. 2020) has increased worldwide. Growing evidence  
37    suggests that the global trends of ecosystem degradation, urbanisation, and noncommunicable  
38    diseases are deeply interconnected <sup>14</sup> (Haahtela, 2019).

39  
40    Exposure to diverse environmental microbiomes—the complex network of microorganisms  
41    in a given environment—is thought to play an important role in human health <sup>15,16</sup> (Rook,  
42    2013; Roslund et al. 2020). Environmental microorganisms support the development and  
43    regulation of the human immune system <sup>16,17</sup> (Renz and Skevani, 2020; Roslund et al. 2020).  
44    Evidence has shown that degraded habitats may harbour a greater relative abundance and  
45    diversity of opportunistic human pathogens, and ecological restoration may restore health-  
46    regulating assemblages <sup>18,19</sup> (Liddicoat et al. 2019; Robinson et al. 2020a). Moreover,  
47    microbial exposure in urban green/blue spaces could improve our health but may depend  
48    heavily upon environmental and design factors including vertical stratification (layering of  
49    microbes in the near-surface atmosphere), vegetation presence, complexity and management

50 <sup>16,20</sup> (Robinson et al. 2020b; Roslund et al. 2020), airflow, and soil management. However,  
51 with appropriate restoration, design and management strategies, these factors could be  
52 optimised to create healthy urban ecosystems, to benefit both human and environmental  
53 health.

54

55 Ensuring long-term urban ecosystem resilience to environmental challenges will depend on  
56 our ability to restore and manage the landscape with considerations for the unseen  
57 foundations of our ecosystems — the microorganisms. Complex microbial interactions are  
58 involved in maintaining the health of urban plant and animal populations <sup>21</sup> (Berg et al.  
59 2017). Considering these microbial interactions as part of any long-term urban development  
60 vision will be essential to ensure urban ecosystems can flourish and maintain resilience.  
61 However, there are currently few considerations for the role of microbial communities in  
62 urban development and landscape design, and multispecies frameworks are rarely used to  
63 inform the management of urban ecosystems. Indeed, recognising its importance in  
64 sustainability, the recently proposed ‘multispecies urbanism’ concept puts forward a  
65 framework for urban development, driven by considerations for reciprocal relationships  
66 between humans and non-humans (including microbes) <sup>22,23</sup> (Rupprecht et al. 2020; Sharma  
67 et al. 2021).

68

69 Sharing similar principles to multispecies urbanism, Microbiome-Inspired Green  
70 Infrastructure, also known as ‘MIGI’, was recently proposed as an integrative system to  
71 promote healthy urban ecosystems <sup>24,25</sup> (Robinson et al. 2018; Watkins et al. 2020). MIGI can  
72 be defined as nature-centric infrastructure that is restored and/or designed and managed to  
73 promote interactions between humans and environmental microbiomes, with explicit  
74 considerations for sustaining microbially-mediated ecosystem functionality and resilience. A

75 considerable challenge to operationalising MIGI is a lack of awareness of the imperative for  
76 urban microbiome research, and the translation of existing research into intelligible and  
77 practicable outputs. Another challenge is addressing the complex needs and constraints of  
78 multiple stakeholders involved in urban landscape management <sup>26</sup> (Marzano et al. 2021).

79

80 In this paper, our primary objectives are to: **(a)** present what is known about the relationship  
81 between urban microbiomes, green infrastructure and environmental processes that affect  
82 urban ecosystem health (ecosystem functionality and resilience, and human health); **(b)**  
83 discuss how we can operationalise the MIGI concept i.e. actionable insights; **(c)** present a  
84 horizon scan of developmental interdisciplinary considerations for MIGI; and, **(d)** highlight  
85 challenges to the implementation of MIGI, whilst proposing a series of multi-stakeholder  
86 engagement workshops.

87

88 This article will help to encourage urban landscape managers to incorporate initial  
89 considerations for MIGI in their development projects to promote healthy urban ecosystems  
90 for humans and the wider biotic community. Although the research is in its infancy, there is  
91 considerable potential for MIGI to help deliver complex ecological and modern urban  
92 societal needs.

93

## 94 **2. MIGI: the relationship between environmental microbiomes,** 95 **ecosystem functionality, and human health**

96

### 97 ***2.1. Ecosystem functionality and resilience context***

98 Microbial communities can be considered the foundations of our ecosystems <sup>1</sup> (Cavicchioli et  
99 al. 2019) (Fig. 1). Soil-microbe-plant interaction studies have demonstrated that plants rely

100 on microbial communities for favourable health <sup>27</sup> (Nazli et al. 2020). Microbial communities  
101 are integral to nutrient and water absorption <sup>28</sup> (Trivedi et al. 2020), and phytohormone  
102 production and regulation activities <sup>29</sup> (ur Rehman et al. 2020). These microbial communities  
103 include arbuscular mycorrhizal fungi and algae, along with symbiotic, associative symbiotic  
104 and free-living plant growth promoting bacteria <sup>27</sup> (Nazli et al. 2020). Endophytes (microbes  
105 living in plant tissues) also benefit plants by enhancing competitive abilities and increasing  
106 resistance to pathogens and other abiotic stressors <sup>30</sup> (Pavithra et al. 2020). Soil microbiomes  
107 are essential to long-term ecosystem resilience in the face of global challenges such as  
108 climate change and degradation <sup>31</sup> (Dubey et al. 2019). In addition to plant health,  
109 microorganisms play roles in carbon sequestration and biogeochemical cycling <sup>31</sup> (Dubey et  
110 al. 2019). It can be further argued that the health of all organisms is interrelated through the  
111 cycling of environmental microorganisms from soils, to plants, animals, and back into the  
112 environment <sup>32</sup> (van Bruggen et al. 2019) (Fig. 1).

113

114 It has been demonstrated that microbiome diversity and network complexity drive multiple  
115 ecosystem functions related to nutrient cycling <sup>33</sup> (Wagg et al. 2019). For instance, grasslands  
116 with poorly-developed microbial networks and reduced microbial richness have low  
117 multifunctionality due to fewer taxa that support functional redundancy and uniqueness <sup>33</sup>  
118 (Wagg et al. 2019). With minimal considerations for microbiomes in urban landscape design  
119 and management projects, it is likely that poor ecosystem multifunctionality and stability will  
120 continue. To promote long-term urban ecosystem health, it is imperative that this trajectory  
121 changes. MIGI provides a framework to operationalise this change.

122


123 ***2.2. Human health context***

124 Growing evidence suggests that exposure to naturally-diverse environmental microbiomes  
125 can improve human health. For example, studies highlight the importance of green space  
126 microbiomes to immunoregulation (Fig. 1). Lehtimäki et al. (2021) showed that risks of  
127 asthma and aeroallergen sensitisation are reduced in rural infants due to exposure to more  
128 biodiverse microbiomes (compared to urban)<sup>34</sup>, and Riskumäki et al. (2021) identified  
129 several environmental taxa that are important in augmenting and/or suppressing systemic  
130 inflammatory immune responses<sup>35</sup>. A loss of biodiversity in urban areas reduces exposure to  
131 diverse environmental microbiomes, whilst increasing exposure to pathogenic microbes<sup>36</sup>  
132 (Parajuli et al. 2018). This is corroborated by a review investigating rural vs. urban  
133 environmental aerobiomes (microbiome of a given airspace) that showed rural-mediated  
134 beneficial immune responses<sup>37</sup> (Flies et al. 2020). Recently, a 28-day biodiversity  
135 intervention demonstrated that inoculating a schoolyard environment with biodiverse features  
136 (e.g., soil and plants from local forest habitats) significantly altered the microbiome of the  
137 children and enhanced important immunoregulatory pathways<sup>16</sup> (Roslund et al. 2020).  
138

139 Other studies indicate the importance of butyrate-producing bacteria which may be promoted  
140 in biodiverse plant-soil systems<sup>18,38</sup> (Liddicoat et al. 2020; Brame et al. 2021), for example  
141 where organic-rich soils experience low redox conditions consistent with fermentative  
142 decomposition of organic matter. Butyrate is a short chain fatty acid associated with gut  
143 health, immunoregulation, and mental health, and exposure to trace level dust containing the  
144 putative spore-forming butyrate-producing bacteria *Kineothrix alysoides* is linked to reduced  
145 anxiety-like behaviour in mice<sup>18</sup> (Liddicoat et al. 2020). Another recent study showed that  
146 spending a short period of time in green spaces can significantly change the human nasal and  
147 respiratory microbiome<sup>39</sup> (Selway et al. 2020). Indeed, microbiota-mediated environmental  
148 health can be thought of as two layers of protective biodiversity<sup>40</sup> (Ruokolainen et al 2017).

149 The first layer, our personal microbiome, is key to health. The second layer, the  
 150 environmental microbiome, represents an important source for replenishing the first;  
 151 therefore safeguarding it represents a critical health insurance policy. It is also important to  
 152 note that a plethora of other potential health benefits are associated with engaging with urban  
 153 nature, including reduced blood pressure <sup>41</sup> (Ideno et al. 2017), lower levels of stress, anxiety  
 154 <sup>42,43</sup> (Birch et al. 2020a; Robinson et al. 2021) and increasing positive affect <sup>44</sup> (Cameron et  
 155 al. 2020).

156



158 **Fig. 1.** Urban multispecies health. Environmental microbiomes are the foundations of our  
 159 ecosystems, and are essential to plant and animal health (including humans).  
 160

### 161 **3. MIGI: actionable insights for landscape managers**

#### 162 **3.1. Vegetation, microbiomes, and the built environment**

163 To operationalise the MIGI framework, we can draw upon several relevant studies. For  
164 example, one important factor is to ensure humans (and other species) are exposed to high  
165 microbial alpha diversity associated with naturally-biodiverse environments from a young  
166 age, which is important for immunoregulation <sup>45,46</sup> (Mulder et al. 2011; Zhang et al. 2020). It  
167 has been demonstrated that air samples downwind from biodiverse sources (e.g., species-rich  
168 plant communities) contain more diverse microbial communities compared to upwind, with  
169 ~50% of airborne bacteria in downwind samples deriving from local plant sources <sup>46</sup>  
170 (Lymperopoulou et al. 2016). Therefore, a relatively simple intervention for urban designers  
171 could be to develop public spaces and buildings downwind from (macro)biodiverse sources,  
172 and to integrate local biodiverse sources within building structures and spaces (Fig. 2, a).  
173 Recently, it was shown that urban green space aerobiomes are vertically stratified, with an  
174 altitudinal decay in bacterial alpha diversity, and possibly a higher relative abundance of  
175 pathogenic taxa at higher altitudes <sup>19</sup> (Robinson et al. 2020a). This reflects a transition from  
176 local plant and soil-related microbiomes at low heights into a broader urban (typically non-  
177 green space) airshed <sup>20</sup> (Robinson et al. 2020b). A potential mitigation measure for this could  
178 be to augment vertical planting in urban areas, allowing exposure to higher natural microbial  
179 alpha diversity in the vertical dimension (Fig. 2, b). It would also be prudent to design urban  
180 areas with greater consideration for inclusive and direct ‘hands-on’ human engagement with  
181 natural features to promote interactions between humans and diverse environmental  
182 microbiomes, and to foster long-term pro-ecological behaviours (Fig 2. c).  
183  
184 Mills et al. (2020) provided evidence that revegetation, particularly with native species, can  
185 improve urban soil microbiome functional diversity <sup>47</sup>. Other studies show that diverse  
186 vegetation communities promote below-ground functional richness, diversity and resilience  
187 <sup>48,49</sup> (Eisenhauer et al. 2018; Canals et al. 2019). Promoting diversity of local vegetation



188 communities is considered a robust strategy to maintain multifunctional processes under  
189 current and future environmental conditions<sup>48</sup> (Eisenhauer et al. 2018). As such, MIGI  
190 strategies could include the planting of diverse, and where possible, native, vegetation  
191 communities to sustain urban ecosystem functionality and resilience (Fig. 2, e). However, it  
192 is not yet clear to what extent locally native plant populations will be able to tolerate future  
193 climate conditions. Studies on woody plants offer conflicting views, with some research  
194 suggesting that intra-population genetic variation may provide sufficient resilience<sup>50</sup> (Borrell  
195 et al., 2018), whilst others argue that given the range in possible climate futures, including  
196 species beyond those that are locally native will be essential in urban environments<sup>51,52</sup>  
197 (Sjöman et al., 2016; Cameron & Blanus, 2016). As such, further research is required to  
198 understand the relationships between locally-native microbial populations and non-local/non-  
199 native plant species, including outcomes for stress tolerance, nutrient acquisition, and  
200 reproduction.

201

202 MIGI strategies should also include the promotion of urban-rural habitat connectivity via  
203 contiguous vertical and horizontal natural corridors (Fig. 3, f). Many urban environments are  
204 ‘patchy’ in terms of quality and connected nature-centric features. To ensure long-term  
205 healthy urban ecosystems, we should aim to connect all natural habitats within towns/cities  
206 and provide essential biophysical corridors to the wider landscape. This action has high  
207 biodiversity conservation value by providing multispecies resources and improving species  
208 interactions and long-term resilience across the landscape, irrespective of species dispersal  
209 abilities or population sizes<sup>53</sup> (Christie and Knowles, 2015). Enhancing networks of  
210 biologically and functionally diverse urban habitats with high vegetation complexity, also has  
211 the potential to improve the distribution of quality aerobiomes, and augment diverse

212 macroscopic species (animals and plants) that contribute to the collective urban  
213 environmental microbiome, and broader ecosystem complexity and resilience.

214

### 215 **3.2. Soil microbiomes**

216 Soil properties will have a key influence on environmental microbiomes associated with  
217 MIGI developments. Soil organic matter and clay-content (proportion of clay-sized particles)  
218 are associated with structure, aggregation, nutrient and water-holding properties, and  
219 therefore the habitat and diversity of microbes <sup>54,55,56</sup> (Jastrow and Miller, 1998; Young and  
220 Crawford, 2004; Torsvik and Øvreås, 2002). A key decision during the establishment phase  
221 of MIGI will be whether to use in-situ or imported soils. Where feasible, using soils with a  
222 moderate amount of clay-content, e.g. sandy loams (10-20% clay content) to loams (around  
223 25% clay content) would be expected to promote microbial diversity. By comparison, sandy  
224 soils provide suboptimal microbial habitat, while heavier texture clay soils (often capable of  
225 forming suitable aggregation and structure) may be more prone to poorer drainage in wet  
226 climates, or greater plant stress in dry climates or during dry periods due to higher wilting-  
227 point moisture content (where water is unavailable to plants). Whether using existing in-situ  
228 soils, or importing new bulk soil, it may be necessary to examine constraints to plant growth.  
229 Sometimes these constraints will be naturally occurring (e.g. shallow depth, impermeable  
230 layers, presence of toxic or nutrient-limiting subsoil conditions), while other times they can  
231 result from management history (e.g. compaction, acidification).

232

233 Where appropriate, addressing soil constraints will help optimise the biological activity and  
234 microbial diversity of the plant-soil system. If organic matter is being applied, ideally this  
235 should be in a nutrient-balanced, or pre-composted form, so that microbial activity and  
236 available nutrients can be harnessed to support the growing vegetation. Ongoing management

237 and human interaction should also be considered. For example, high levels of foot/vehicle  
238 traffic can lead to compaction and degradation, and may create zones of poor soil  
239 microbiome conditions with sub-optimal health influence (e.g. along paths) within a natural  
240 space that is offering health-promoting microbial exposures. Exposure pathways to permit  
241 beneficial human-soil microbiome contact also remains an area for research. Based on  
242 available knowledge, low-level exposure to soil (e.g. dust) with biodiverse content to help  
243 contain potential pathogenic activity, would represent a reasonable starting point to  
244 supporting immune fitness. It is also expected that soils will gain maximum health-promoting  
245 potential by spending the majority of time covered with biodiverse vegetation, which is  
246 another consideration for designing MIGI exposure pathways.


247

248 There is also growing evidence to show that plant communities require complex mycorrhizal  
249 networks, acting as conduits of inter-plant communication, and facilitating pathogen defence,  
250 adaptation, growth, and memory <sup>57,58</sup> (Filotas et al. 2014; Birch et al. 2020). Indeed, healthy  
251 vegetation community phenotypes at higher levels emerge from a multitude of localised and  
252 often subterranean entities interacting at lower levels <sup>59</sup> (Ibarra et al. 2020). For example,  
253 Birch et al. (2020) recently demonstrated that plant growth was significantly associated with  
254 the number of ectomycorrhizal connections to other plants, and the number of genetically  
255 distinct fungi that were present <sup>58</sup>. Therefore, we must at least consider the condition and  
256 ecology of the substrate and its role in sustaining ecosystem functionality and resilience. This  
257 will involve viewing habitat conservation and restoration through the lens of complex  
258 systems science. However, many experts in this field could provide appropriate consultation  
259 at each stage of a development project <sup>25</sup> (Watkins et al. 2020).

260

261 Interestingly, a recent study showed that landscaping materials (e.g. compost and mulch)  
 262 have a 'microbial shelf life', and long-term storage can significantly reduce the availability of  
 263 bacterial taxa linked to human health and degradation of pollution<sup>60</sup> (Soininen et al. 2021).  
 264 This suggests that as part of a holistic MIGI strategy, short-term storage times should be  
 265 considered when planning the utilisation of landscaping materials (Fig. 2, d).

266



267

268 **Fig. 2.** Actionable insights for MIGI, including vegetation complexity, downwind  
 269 development and local integration of biodiverse source (a); a solution to the concept of  
 270 vertical stratification (b); hands-on engagement with natural features to promote  
 271 immunoregulation (c); recommended soil types to promote diverse microbial habitat and  
 272 short-term storage of landscaping materials (d); revegetation with diverse native plants to  
 273 promote functional diversity (e); the concept of habitat connectivity via contiguous natural  
 274 corridors to promote long-term multispecies health (f).

275

276

277 **4. MIGI: Horizon scan of emerging research and practice**

#### 278 **4.1. Bioreceptive materials and bio-integrated design**

279 Bio-Integrated Design refers to interdisciplinary methodologies that merge applied  
280 biotechnology, architecture and design, in order to create sustainable systems for the built  
281 environment. It encompasses a range of biologically-mediated processes, such as  
282 biosilification, biomineralization and bioremediation, as well as the development of material  
283 substrates for living systems. Guillitte defined the term ‘bioreceptivity’ to describe the ability  
284 of a building material to be colonised by living organisms <sup>61</sup> (Guillitte, 1995). In the context  
285 of bio-integrated design to enable MIGI, bioreceptivity has been explored through the design  
286 of architectural scaffolds with the goal of creating self-regulating systems which are host to  
287 cryptogrammic species as well as microbial biodiversity <sup>62</sup> (Cruz & Beckett, 2016).

288

289 Whilst horizontal surfaces including roofs, terraces or pavements offer scope for plant  
290 growth, vertical surfaces (e.g. building facades and infrastructural walls) offer far harsher  
291 environments due to excessive water run-off, strong exposure to winds and lack of nutrient  
292 rich substrates. But cities have vast areas of xeric surfaces that offer opportunities to be  
293 photosynthetically active. For an accelerated creation of primary bioreceptivity on vertical  
294 surfaces, a number of design steps can accelerate this process. Porosity and surface roughness  
295 are two vital functions. Firstly, the calibration of pore size can enable water absorption and  
296 retention, reaching colonising organisms through capillary or surface-binding effects, and  
297 secondly as a means to exploit extrinsic factors for bioreceptivity through the collection of  
298 organic material and fixation of cryptogamic surface cover. Different compositions of  
299 bioreceptive cementitious materials have been explored based on Magnesium Phosphate  
300 Concrete (Manso et al., 2015) and other Ordinary Portland Concrete (OPC) mixes with the  
301 aim to create long-term carbon offset <sup>63</sup>. Studies of OPC have shown that apart from altering  
302 the physico-chemical properties of materials, morphological variations explored via

303 computational design strategies are a powerful means to reduce water run-off and increase  
304 moisture retention, extending the residence time to create zones for accelerated growth <sup>64</sup>  
305 (Cruz, 2021). In future, quaternary bioreceptive strategies may be used <sup>65</sup> (Sanmartín et al.,  
306 2021) whereby surface additives to a material scaffold such as hydrogels or humic material  
307 are applied to enhance colonisation. Further studies are needed to test the role of pH on  
308 material substrates in biofilm formation and the establishment of microbial communities  
309 which are vital to establish cryptogamic growth.

310

311 Lichens and, primarily bryophytes play key roles in design of bioreceptive structures within  
312 architecture, provoking a biophilic response <sup>66</sup> (Wilson, 1984) through their aesthetic  
313 appearance and tactility. However, they are also important components due to their capacity  
314 to regulate their photosynthetic activity depending on moisture availability while surviving  
315 for long periods without water - *poikilohydry*. In relation to MIGI, bryophytes harbour  
316 ecologies with prokaryotic and eukaryotic algae, bacteria and fungi. Cyanobacteria are  
317 keystone species in other nutrient limited environments such as desert crusts <sup>67</sup> (Yeager et al.,  
318 2007), where their ability to fix nitrogen and carbon enables succession by other organisms  
319 through exchange of metabolites. However, our understanding of the exact ecological roles of  
320 microorganisms associated with bryophyte hosts is nascent. For example, while the presence  
321 of certain bacteria varies according to the bryophyte species, in one study *Proteobacteria*,  
322 *Actinobacteria*, *Acidobacteria*, *Bacteroidetes*, *Armatimonadetes* and *Planctomycetes* were  
323 detected in all moss microbiomes <sup>68</sup> (Tang et al., 2016). Analogous to the urban environment,  
324 during early stages of habitat restoration it has been shown that bryophyte communities  
325 enrich populations of microbial life on calcareous rocks <sup>69</sup> (Cao et al., 2020). It may be  
326 speculated that colonisation by bryophytes and their associated microbiota could have  
327 advantageous effects for growth promotion in other plants due to the presence of bacteria

328 containing genes for production of indole acetic acid, siderophores or solubilisation of  
329 phosphate <sup>70</sup> (Insuk et al., 2020). Through this application of bio-integrated design to produce  
330 poikilohydric living walls, it is possible to employ more of the surface area that is  
331 underutilised within our urban environments to deliver MIGI.

332

#### 333 **4.2. Microbial inoculants**

334 Microbial inoculants have recently been used to shift microbiota in landscaping materials  
335 towards an immuno-protective assemblage <sup>71</sup> (Hui et al. 2019). The authors developed a  
336 microbial inoculant from biodiverse sources (e.g., forest materials), resembling the  
337 microbiome of organic soils. After the study subjects made contact with these inoculated  
338 materials, the relative abundance of opportunist pathogens on the skin significantly  
339 decreased. Furthermore, Roslund et al. (2020) demonstrated that a biodiversity intervention  
340 using microbial inoculants from forest floor materials changed the skin microbiome of  
341 children and enhanced immunoregulatory pathways <sup>16</sup>. Several other studies show that  
342 microbial inoculants can be beneficial for plant health, for example, via Plant Growth  
343 Promoting Rhizobacteria (PGPR) <sup>72</sup> (Sacristán-Pérez-Minayo et al. 2020). PGPRs have the  
344 potential to protect plants from drought and metal stresses and play important roles in plant  
345 growth, which itself could minimise the use of harmful synthetically produced chemical  
346 fertilisers <sup>73</sup> (Kumar et al. 2019). Therefore, MIGI strategies could incorporate microbial  
347 inoculants to enhance ecosystem health (Fig. 3, b).

348

#### 349 **4.3. Supportive tools**

350 Useful tools are being developed that could help with MIGI interventions in the near future.  
351 For example, Saleem et al. (2019) produced a framework to model the environmental  
352 microbiome's influence on plant traits and ecosystem functionality, highlighting the

353 possibility of creating an index to monitor and enhance plant growth and soil/ecosystem  
354 health <sup>74</sup>. Along similar lines, it could be valuable to develop a form of 'Health Promotion  
355 Potential Index' for human health. This could be based on known combinations of  
356 environmental microbial factors that promote (or demote) immunoregulation and  
357 homeostasis, such as alpha and functional diversity and beneficial taxa that produce health-  
358 regulating compounds (Fig. 3, c).

359

#### 360 **4.4. Bioremediation**

361 Emerging bioremediation research could also be considered in MIGI strategies. In-situ  
362 bioremediation enables ongoing decontamination or degradation of pollutants without  
363 complex excavation or transportation. This could play a role in sequestering metals such as  
364 lead and zinc—present in many cities as components of urban dusts <sup>75</sup> (Alharbi et al., 2019).  
365 Bacteria, fungi and microalgae have evolved several mechanisms to adsorb or absorb heavy  
366 metals. In a study investigating river sediments contaminated with cadmium, copper, lead and  
367 zinc, it was hypothesised that species richness may be a function of “public goods” within the  
368 microbial community, such as metallophores, EPC, biogenic sulphides or calcite. Bacteria  
369 such as *Pseudomonas* and *Bacillus* may precipitate metal and thus benefit other organisms  
370 with spatial proximity by creating detoxified regions <sup>76</sup> (Jacquiod et al. 2018).

371

372 Kang et al. (2016) showed that the synergistic combination of bacterial strains including  
373 *Viridibacillus arenosi* B-21, *Sporosarcina soli* B-22, *Enterobacter cloacae* KJ-46 and *E.*  
374 *cloacae* KJ-47 was effective at sequestering Pb (98.3% effective) and Cd (85.4%) in soils <sup>77</sup>.  
375 Biofilters are being developed that embed bacterial biofilms to absorb heavy metal ions  
376 (Priyadarshane and Das, 2020) <sup>78</sup>. In terms of MIGI, this kind of strategy could be  
377 developed with Sustainable Drainage Systems in mind, e.g., a biofilter-embedded rain garden



378 (Fig. 3, d). There are numerous organic compounds that pose a threat to human health, found  
379 in elevated concentrations. Microbial mechanisms that may be employed in MIGI are  
380 hydrolysis and oxidation, with the goal of producing benign compounds through metabolic  
381 activity. For instance, endocrine disruptors such as phthalates and alkylphenols are ubiquitous  
382 in water systems as a result of human activity<sup>79</sup> (Bergé et al., 2014) but there are microbial  
383 mechanisms to break these down under certain conditions<sup>80</sup> (Boll et al., 2020). Indeed, many  
384 bacterial taxa have been identified that have significant pollutant degradation properties<sup>81</sup>  
385 (Ojuederie and Babalola, 2017). As an alternative to contained bioreactor systems, creating  
386 stable synthetic ecologies and applying eco-evolutionary principles to enhance  
387 bioremediation is compatible with MIGI principles<sup>82</sup> (Borchert et al., 2021).

388

#### 389 ***4.5. Plant nutrition, soil issues, and anti-microbial resistance***

390 It is suspected that high-dosage artificial agricultural fertilisers are detrimental to mycorrhizal  
391 networks, for example, by promoting taxa with pathogenic traits<sup>83</sup> (Paungfoo-Lonhienne et  
392 al. 2015). Studies have suggested that organic or ‘natural’ fertilisers and plant conditioners  
393 outperform chemically synthetic N, P and K types in promoting plant health/quality<sup>84,85</sup>  
394 (Hammad et al. 2020; Dahunsi et al. 2021). Additional research in this area could bring value  
395 to the MIGI concept, particularly research focusing on the application of fertilisers  
396 sympathetic to soil-plant microbial interactions. There are also physical soil issues to  
397 consider in urban landscape management. The loss of organic matter, compaction, excessive  
398 disturbance will likely damage microbial communities<sup>86</sup> (Gregory et al. 2015). Research to  
399 fully understand the implications of these factors could enhance urban ecosystem  
400 management. The loss of soil microbial diversity has also been linked to the exacerbation of  
401 the spread of antimicrobial resistance<sup>87</sup> (Chen et al. 2019). Antimicrobial resistance has  
402 important implications for human health by making infections harder to treat and increasing

403 risks of disease spread <sup>88</sup> (WHO, 2020). MIGI researchers aiming to reduce the abundance  
404 and diversity of antimicrobial resistant genes in urban environments could explore the  
405 strategy of increasing soil microbial diversity.

406

#### 407 ***4.6. Emerging biosecurity considerations***

408 Alongside the positive opportunities presented by a multispecies approach to green  
409 infrastructure, the threats need to be equally researched and mitigated against within the  
410 MIGI framework. Many of the most pressing biosecurity threats to ecosystems are microbial  
411 in nature (in the UK, for example, *Hymenoscyphus fraxineus*, *Phytophthora ramorum* and  
412 *Candidatus liberibacter solanacearum*), threatening urban green infrastructure as well as  
413 agricultural crops that urban populations depend upon. It is increasingly recognised, however,  
414 that most biosecurity research and regulations focus on impacts to agricultural and forestry  
415 sectors. Work is urgently required to understand the extent to which urban ecosystems are  
416 threatened and what tools are most effective at safeguarding them. Key questions exist at a  
417 societal level, concerning the mismatch between public perception of biosecurity risk and  
418 expert assessment, the capacity for codes of conduct to influence behaviour and the ability to  
419 communicate information between policy makers, researchers and the public. Industry-  
420 specific technical questions also exist concerning the risks associated with importing soil for  
421 planting, the materials used in packaging for construction projects <sup>89</sup> (Kemp et al. 2021), or to  
422 what extent novel microbial pests and diseases might influence assisted colonisation  
423 programmes or substitutions for keystone species die-off.

424

425 Regulation and guidance exists to combat and mitigate these threats have been developed at  
426 international (e.g. European and Mediterranean Plant Protection Organisation <sup>90</sup> (EPPO,  
427 2020), national (Defra Plant Health Risk Register <sup>91</sup> (Defra, 2014)) and regional levels <sup>92</sup>


428 (Public Health Agency of Canada, 2018), complemented by industry-specific guidance in  
429 many countries <sup>93</sup> (Watkins & Arkell, 2019). Nevertheless, further work is required to  
430 address the open questions and operationalise policy, and MIGI offers an opportunity to  
431 address these in a holistic manner. One of the core challenges presented by biosecurity threats  
432 is that the technical understanding of biosecurity in the construction and development  
433 industries is currently at an early stage, with many practitioners confusing biosecurity with  
434 concepts such as biodiversity, and seeing threats posed by biosecurity risks as someone else's  
435 responsibility. Integrating existing schemes such as Plant Healthy  
436 (<https://planthealthy.org.uk>) within MIGI will be essential, not only to ensure that best  
437 practice is developed but also so that practitioners can clearly understand how their actions  
438 reinforce the biosecurity continuum <sup>94</sup> (Sequeira and Griffin 2014). Biosecurity is one of  
439 many layers that prove challenging to translate between research and practice, and to this  
440 end, delivering the plant selection database proposed in the MIGI toolkit <sup>19</sup> (Watkins et al.,  
441 2020) will provide not only a common understanding of the tools required to deliver MIGI  
442 but also a shared vocabulary for different sectors to draw upon when discussing projects.

443

#### 444 ***4.7. Social innovation: promoting stronger human-nature relationships***

445 Other important factors within the MIGI framework include social innovation, education and  
446 stimulating awe for nature, with emphasis on the foundations of our ecosystems—the diverse  
447 microbial communities. Strategies such as ‘learning about the land, on the land’ <sup>95</sup> (Learning  
448 the Land, 2021) could help to inspire pro-ecological behaviours that reinforce a sense of  
449 stewardship for our diverse and complex ecosystems. To paraphrase Simard (2018) “*viewing*  
450 *[ecosystems] through the lens of cognition, microbiome collaborations, and intelligence may*  
451 *contribute to a more holistic approach to studying ecosystems and a greater human empathy*  
452 *and caring for the health of our [landscapes]”* <sup>96</sup>. Various campaigns also promote the

453 concept of ‘nature connectedness’ (emotional and cognitive connection with the natural  
 454 world). Studies in this area show increases in wellbeing and pro-ecological behaviour as a  
 455 result of enhanced nature connectedness<sup>97,98</sup> (Capaldi et al. 2014; Capaldi et al. 2015).  
 456  
 457 For the technologically-minded, virtual reality systems could be developed to facilitate urban  
 458 habitat tours. These could include interactive macroscopic displays of microbial  
 459 communities, whilst providing information on the composition and functional roles that  
 460 microbes play in the local ecosystems (Fig. 3, f).  
 461



462  
 463 **Fig. 3.** Horizon scan of developmental considerations for MIGI, including interventions (b  
 464 and d), design and supportive features (a and e), and applications for engagement and to  
 465 acquire useful urban ecosystem health information (c and f).  
 466

467

## 468 5. *MIGI: Challenges to operationalisation*

469 Watkins et al. (2020) identify barriers to operationalising MIGI, relating to the complexity of  
470 urban development projects, and communicating the benefits of MIGI interventions to  
471 stakeholders involved in urban planning <sup>19</sup>. To ensure the implementation of green  
472 infrastructure strategies, stakeholder buy-in is required throughout green infrastructure  
473 planning, design, operation and management <sup>99</sup> (Smith, 2020). The range of stakeholders  
474 include: local authorities, developers and private clients, planning professionals, landowners,  
475 landscape specialists, architects, ecologists, statutory agencies, contractors, local businesses  
476 and community groups. In the UK, there are a number of government-funded research  
477 projects into green infrastructure, biosecurity, climate change-readiness, and supply chains  
478 (e.g. BRIGIT, Future Oak, Plant Health Centre). Learning from these projects is prudent as  
479 there is already evidence of stakeholder fatigue and pre-existing challenges of engaging with  
480 industry sectors that see these aspects as someone else's problem or not aligned with  
481 commercial goals. This suggests new approaches are needed and highlights the importance of  
482 internationalising research projects so that robust data can be gathered from diverse  
483 stakeholders.

484

485 Although fundamental, adding the lens of microbial ecology to an already expansive multi-  
486 stakeholder initiative, MIGI has to reckon with the “perception that multi-stakeholder  
487 initiatives slow down urban planning and policy development processes due to a lack of  
488 consensus and different sectoral interests” <sup>100</sup> (Ferreira et al. 2020). Further development  
489 should align with the priorities of the stakeholder groups and generate clear, actionable points  
490 overlaid onto existing frameworks, rather than increasing complexity. For instance, in the  
491 UK, some MIGI considerations overlap with the policies laid out in the London Plan 2021.  
492 The focus on connected landscapes and biodiversity corridors supports the value of  
493 connected nature-centric features. Policy G5 Urban Greening asks that new developments

494 incorporate high quality landscaping, green roofs, green walls and nature-based sustainable  
495 drainage and introduces an Urban Greening Factor to evaluate the quality and quantity of  
496 green space design and delivery <sup>101</sup> (Greater London Authority, 2021). This guidance is  
497 currently under consultation and could benefit from MIGI-related input to aid with the  
498 ambition of delivering biodiversity net gain. Overlaps could be identified through direct  
499 communication with existing built environment biodiversity-centric networks.

500

## 501 **6. MIGI: Workshop series proposal**

502 To determine clear action points for future research and address the challenges related to  
503 divergence of interests among stakeholders, it is vital to ascertain diverse priorities and  
504 concerns through early consultation <sup>102</sup> (Khoshkar et al. 2017). This could also aid the  
505 development of a “common language” <sup>103</sup> (Ugolini et al. 2018) that translates researchers’  
506 findings into verbal and graphic outputs relevant to non-expert audiences. We are currently  
507 developing a series of workshops to discuss what is known about the microbiome in a health  
508 and ecosystem functionality context, and reveal tangible opportunities to include MIGI in  
509 urban planning. These workshops provide an opportunity to engage with reflective  
510 stakeholders in identifying not only challenges but also specific factors (e.g. MIGI toolkits,  
511 portfolio of illustrative examples) and alignments between current requirements/protocols,  
512 and how MIGI could be integrated. During the workshops we will discuss ‘what researchers  
513 should be working on’, and opportunities and constraints by drawing together the  
514 perspectives and needs of different stakeholders. These workshops will form part of a process  
515 of long-term engagement and partnership to enhance urban ecosystem health via MIGI  
516 strategies.

517

518 Developing the MIGI concept has the potential to enhance urban ecosystem functionality and  
519 resilience as well as human health. In this paper, we have provided several examples of MIGI  
520 actionable insights in addition to a horizon scan of emerging MIGI-related research and  
521 practice. A greater emphasis on the roles of microbial communities (from below-ground and  
522 up) in our urban ecosystems is needed. Understanding microbial dynamics will likely have an  
523 important role to play in the efficacy of our adaptability and long-term resilience to ongoing  
524 global environmental change. MIGI research agendas aim to promote this realm of thinking  
525 with considerations for multispecies health. However, overcoming the challenges to  
526 operationalising MIGI will be essential to furthering its practical development.

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541 **References**

- 542 1. Cavicchioli, Ricardo, William J. Ripple, Kenneth N. Timmis, Farooq Azam, Lars R.  
543 Bakken, and others. 2019. “Scientists’ Warning to Humanity: Microorganisms and  
544 Climate Change,” *Nature Reviews. Microbiology*, 17.9: 569–86
- 545 2. Guerra, Carlos A., Anna Heintz-Buschart, Johannes Sikorski, Antonis Chatzinotas,  
546 Nathaly Guerrero-Ramírez, and others. 2020. “Blind Spots in Global Soil Biodiversity  
547 and Ecosystem Function Research,” *Nature Communications*, 11.1: 3870
- 548 3. Bach, Elizabeth M., Kelly S. Ramirez, Tandra D. Fraser, and Diana H. Wall. 2020.  
549 “Soil Biodiversity Integrates Solutions for a Sustainable Future,” *Sustainability*, 12.7:  
550 2662
- 551 4. Greenspan, Sasha E., Gustavo H. Migliorini, Mariana L. Lyra, Mariana R. Pontes,  
552 Tamílie Carvalho, and others. 2020. “Warming Drives Ecological Community  
553 Changes Linked to Host-Associated Microbiome Dysbiosis,” *Nature Climate Change*,  
554 10.11: 1057–61
- 555 5. Tibbett, Mark, Tandra D. Fraser, and Sarah Duddigan. 2020. “Identifying Potential  
556 Threats to Soil Biodiversity,” *PeerJ*, 8.e9271: e9271
- 557 6. Smith, Katherine F., Michael Goldberg, Samantha Rosenthal, Lynn Carlson, Jane  
558 Chen, and others. 2014. “Global Rise in Human Infectious Disease  
559 Outbreaks,” *Journal of the Royal Society, Interface*, 11.101: 20140950
- 560 7. Jairath, Vipul, and Brian G. Feagan. 2020. “Global Burden of Inflammatory Bowel  
561 Disease,” *The Lancet. Gastroenterology & Hepatology*, 5.1: 2–3
- 562 8. El-Gamal, Yehia M., Elham M. Hossny, Zeinab A. El-Sayed, and Shereen M. Reda.  
563 2017. “Allergy and Immunology in Africa: Challenges and Unmet Needs,” *The  
564 Journal of Allergy and Clinical Immunology*, 140.5: 1240–43



- 565 9. Borna, Eivind, Bright I. Nwaru, Anders Bjerg, Roxana Mincheva, Madeleine  
566 Rådinger, and others. 2019. “Changes in the Prevalence of Asthma and Respiratory  
567 Symptoms in Western Sweden between 2008 and 2016,” *Allergy*, 74.9: 1703–15
- 568 10. Holman, N., N. G. Forouhi, E. Goyder, and S. H. Wild. 2011. “The Association of  
569 Public Health Observatories (APHO) Diabetes Prevalence Model: Estimates of Total  
570 Diabetes Prevalence for England, 2010-2030: Estimates of Total Diabetes Prevalence  
571 for England, 2010-2030,” *Diabetic Medicine: A Journal of the British Diabetic  
572 Association*, 28.5: 575–82
- 573 11. Kainu, Annette, Paula Pallasaho, Päivi Piirilä, Ari Lindqvist, Anssi Sovijärvi, and  
574 others. 2013. “Increase in Prevalence of Physician-Diagnosed Asthma in Helsinki  
575 during the Finnish Asthma Programme: Improved Recognition of Asthma in Primary  
576 Care? A Cross-Sectional Cohort Study,” *Primary Care Respiratory Journal: Journal  
577 of the General Practice Airways Group*, 22.1: 64–71
- 578 12. Dinse, Gregg E., Christine G. Parks, Clarice R. Weinberg, Carroll A. Co, Jesse  
579 Wilkerson, and others. 2020. “Increasing Prevalence of Antinuclear Antibodies in the  
580 United States,” *Arthritis & Rheumatology*, 72.6: 1026–35
- 581 13. Paramasivam, Arumugam, Jayaseelan Vijayashree Priyadharsini, and Subramanian  
582 Raghunandhakumar. 2020. “Implications of M6A Modification in Autoimmune  
583 Disorders,” *Cellular & Molecular Immunology*, 17.5: 550–51
- 584 14. Haahtela, Tari. 2019. “A Biodiversity Hypothesis,” *Allergy*
- 585 15. Rook, Graham A. 2013. “Regulation of the Immune System by Biodiversity from the  
586 Natural Environment: An Ecosystem Service Essential to Health,” *Proceedings of the  
587 National Academy of Sciences of the United States of America*, 110.46: 18360–67
- 588 16. Roslund, Marja I., Riikka Puhakka, Mira Grönroos, Noora Nurminen, Sami  
589 Oikarinen, and others. 2020. “Biodiversity Intervention Enhances Immune Regulation

- 590 and Health-Associated Commensal Microbiota among Daycare Children,” *Science*  
591 *Advances*, 6.42
- 592 17. Renz, Harald, and Chrysanthi Skevaki. 2021. “Early Life Microbial Exposures and  
593 Allergy Risks: Opportunities for Prevention,” *Nature Reviews. Immunology*, 21.3:  
594 177–91
- 595 18. Liddicoat, Craig, Philip Weinstein, Andrew Bissett, Nicholas J. C. Gellie, Jacob G.  
596 Mills, and others. 2019. “Can Bacterial Indicators of a Grassy Woodland Restoration  
597 Inform Ecosystem Assessment and Microbiota-Mediated Human  
598 Health?,” *Environment International*, 129: 105–17
- 599 19. Robinson, Jake M., Christian Cando-Dumancela, Rachael E. Antwis, Ross Cameron,  
600 Craig Liddicoat, and others. 2020a. “Exposure to Airborne Bacteria Depends upon  
601 Vertical Stratification and Vegetation Complexity,” *Scientific Reports*.
- 602 20. Berg, Gabriele, Martina Köberl, Daria Rybakova, Henry Müller, Rita Grosch, and  
603 others. 2017. “Plant Microbial Diversity Is Suggested as the Key to Future Biocontrol  
604 and Health Trends,” *FEMS Microbiology Ecology*, 93.5
- 605 21. Rupprecht, Christoph D. D., Joost Vervoort, Chris Berthelsen, Astrid Mangnus,  
606 Natalie Osborne, and others. 2020. “Multispecies Sustainability,” *Global*  
607 *Sustainability*, 3.e34
- 608 22. Sharma, A. 2021. The City under Stress: Waking to a Multispecies Urban. *Urban*  
609 *Environments as Spaces of Living in Transformation*, p.7. Online. Available at:  
610 <https://urbanenv.org/the-city-under-stress/> (Accessed on 8<sup>th</sup> March 2021).
- 611 23. Robinson, Jake, Jacob Mills, and Martin Breed. 2018. “Walking Ecosystems in  
612 Microbiome-Inspired Green Infrastructure: An Ecological Perspective on Enhancing  
613 Personal and Planetary Health,” *Challenges*, 9.2: 40

- 614 24. Watkins, Harry, Jake M. Robinson, Martin F. Breed, Brenda Parker, and Philip  
615 Weinstein. 2020. "Microbiome-Inspired Green Infrastructure: A Toolkit for  
616 Multidisciplinary Landscape Design," *Trends in Biotechnology*, 38.12: 1305–8
- 617 25. Karlsdóttir, B., Pollard, C., Paterson, A., Watkins, H., Marzano, M. 2021. Assessment  
618 of large-scale plant biosecurity risks to Scotland from large scale plantings for  
619 landscaping and infrastructure projects: Project Final Report. PHC2019/05.
- 620 26. Nazli, F., Khan, M.Y., Jamil, M., Nadeem, S.M. and Ahmad, M. 2020. Soil microbes  
621 and plant health. In *Plant Disease Management Strategies for Sustainable Agriculture*  
622 *through Traditional and Modern Approaches* (pp. 111-135). Springer, Cham.
- 623 27. Trivedi, Pankaj, Jan E. Leach, Susannah G. Tringe, Tongmin Sa, and Brajesh K.  
624 Singh. 2020. "Plant-Microbiome Interactions: From Community Assembly to Plant  
625 Health," *Nature Reviews. Microbiology*, 18.11: 607–21
- 626 28. ur Rehman, F., Kalsoom, M., Nasir, T.A., Adnan, M., Anwar, S. and Zahra, A. 2020.  
627 Chemistry of Plant–Microbe Interactions in Rhizosphere and Rhizoplane. *Ind J Pure*  
628 *App Biosci*, 8, pp.11-19. 10.3389/fpls.2018.00112
- 629 29. Pavithra, Sumant Bindal, Meenakshi Rana, and Seweta Srivastava. 2019. "Role of  
630 Endophytic Microbes against Plant Pathogens: A Review," *Asian Journal of Plant*  
631 *Sciences*, 19.1: 54–62
- 632 30. Dubey, Anamika, Muneer Ahmad Malla, Farhat Khan, Kanika Chowdhary, Shweta  
633 Yadav, and others. 2019. "Soil Microbiome: A Key Player for Conservation of Soil  
634 Health under Changing Climate," *Biodiversity and Conservation*, 28.8–9: 2405–29
- 635 31. van Bruggen, Ariena H. C., Erica M. Goss, Arie Havelaar, Anne D. van Diepeningen,  
636 Maria R. Finckh, and others. 2019. "One Health - Cycling of Diverse Microbial  
637 Communities as a Connecting Force for Soil, Plant, Animal, Human and Ecosystem  
638 Health," *The Science of the Total Environment*, 664: 927–37

- 639 32. Wagg, Cameron, Klaus Schlaeppli, Samiran Banerjee, Eiko E. Kuramae, and Marcel  
640 G. A. van der Heijden. 2019. “Fungal-Bacterial Diversity and Microbiome  
641 Complexity Predict Ecosystem Functioning,” *Nature Communications*, 10.1: 4841
- 642 33. Lehtimäki, Jenni, Jonathan Thorsen, Morten Arendt Rasmussen, Mathis Hjelmsø,  
643 Shiraz Shah, and others. 2020. “Urbanized Microbiota in Infants, Immune  
644 Constitution, and Later Risk of Atopic Diseases,” *The Journal of Allergy and Clinical  
645 Immunology* <<https://doi.org/10.1016/j.jaci.2020.12.621>>
- 646 34. Riskumäki, Matilda, Ioannis Tassis, Noora Ottman, Alina Suomalainen, Paulina  
647 Werner, and others. 2021. “Interplay between Skin Microbiota and Immunity in  
648 Atopic Individuals,” *Allergy*, 76.4: 1280–84
- 649 35. Parajuli, Anirudra, Mira Grönroos, Nathan Siter, Riikka Puhakka, Heli K. Vari, and  
650 others. 2018. “Urbanization Reduces Transfer of Diverse Environmental Microbiota  
651 Indoors,” *Frontiers in Microbiology*, 9
- 652 36. Flies, Emily J., Laurence J. Clarke, Barry W. Brook, and Penelope Jones. 2020.  
653 “Urbanisation Reduces the Abundance and Diversity of Airborne Microbes - but  
654 What Does That Mean for Our Health? A Systematic Review,” *The Science of the  
655 Total Environment*, 738.140337: 140337
- 656 37. Brame, Joel E., Craig Liddicoat, Catherine A. Abbott, and Martin F. Breed. 2021.  
657 “The Potential of Outdoor Environments to Supply Beneficial Butyrate-Producing  
658 Bacteria to Humans,” *The Science of the Total Environment*, 777.146063: 146063
- 659 38. Selway, Caitlin A., Jacob G. Mills, Philip Weinstein, Chris Skelly, Sudesh Yadav,  
660 and others. 2020. “Transfer of Environmental Microbes to the Skin and Respiratory  
661 Tract of Humans after Urban Green Space Exposure,” *Environment International*,  
662 145.106084: 106084

- 663 39. Ruokolainen, Lasse, Jenni Lehtimäki, Antti Karkman, Tari Haahtela, Leena von  
664 Herten, and others. 2017. “Holistic View on Health: Two Protective Layers of  
665 Biodiversity,” *Annales Zoologici Fennici*, 54.1–4: 39–49
- 666 40. Birch, Jo, Clare Rishbeth, and Sarah R. Payne. 2020. “Nature Doesn’t Judge You -  
667 How Urban Nature Supports Young People’s Mental Health and Wellbeing in a  
668 Diverse UK City,” *Health & Place*, 62.102296: 102296
- 669 41. Robinson, Jake M., Paul Brindley, Ross Cameron, Danielle MacCarthy, and Anna  
670 Jorgensen. 2021. “Nature’s Role in Supporting Health during the COVID-19  
671 Pandemic: A Geospatial and Socioecological Study,” *International Journal of  
672 Environmental Research and Public Health*, 18.5: 2227
- 673 42. Cameron, Ross W. F., Paul Brindley, Meghann Mears, Kirsten McEwan, Fiona  
674 Ferguson, and others. 2020. “Where the Wild Things Are! Do Urban Green Spaces  
675 with Greater Avian Biodiversity Promote More Positive Emotions in  
676 Humans?,” *Urban Ecosystems*, 23.2: 301–17
- 677 43. Mulder, Imke E., Bettina Schmidt, Marie Lewis, Margaret Delday, Christopher R.  
678 Stokes, and others. 2011. “Restricting Microbial Exposure in Early Life Negates the  
679 Immune Benefits Associated with Gut Colonization in Environments of High  
680 Microbial Diversity,” *PloS One*, 6.12: e28279
- 681 44. Zhang, Jinya, Fernando Rodríguez, Maria Jesus Navas, Mar Costa-Hurtado, Vanessa  
682 Almagro, and others. 2020. “Fecal Microbiota Transplantation from Warthog to Pig  
683 Confirms the Influence of the Gut Microbiota on African Swine Fever  
684 Susceptibility,” *Scientific Reports*, 10.1: 17605
- 685 45. Lymperopoulou, Despoina S., Rachel I. Adams, and Steven E. Lindow. 2016.  
686 “Contribution of Vegetation to the Microbial Composition of Nearby Outdoor  
687 Air,” *Applied and Environmental Microbiology*, 82.13: 3822–33

- 688 46. Mills, Jacob G., Andrew Bissett, Nicholas J. C. Gellie, Andrew J. Lowe, Caitlin A.  
689 Selway, and others. 2020. "Revegetation of Urban Green Space Rewilds Soil  
690 Microbiotas with Implications for Human Health and Urban Design: Revegetation  
691 Rewilds Urban Soil Microbiotas," *Restoration Ecology*
- 692 47. Eisenhauer, N., Hines, J., Isbell, F., van der Plas, F., Hobbie, S.E., Kazanski, C.E.,  
693 Lehmann, A., Liu, M., Lochner, A., Rillig, M.C. and Vogel, A. 2018. Plant diversity  
694 maintains multiple soil functions in future environments. *Elife*, 7, p.e41228.  
695 10.7554/eLife.41228.001
- 696 48. Canals, R. M., L. Múgica, M. Durán, and L. San Emeterio. 2019. "Soil Bacterial  
697 Functional Diversity Mirrors the Loss of Plant Diversity by the Expansion of a Native  
698 Tall-Grass in High Mountain Grasslands," *Plant and Soil*, 445.1–2: 243–57
- 699 49. Borrell, James S., Nian Wang, Richard A. Nichols, and Richard J. A. Buggs. 2018.  
700 "Genetic Diversity Maintained among Fragmented Populations of a Tree Undergoing  
701 Range Contraction," *Heredity*, 121.4: 304–18
- 702 50. Sjöman, Henrik, Justin Morgenroth, Johanna Deak Sjöman, Arne Sæbø, and Ingo  
703 Kowarik. 2016. "Diversification of the Urban Forest—Can We Afford to Exclude  
704 Exotic Tree Species?," *Urban Forestry & Urban Greening*, 18: 237–41
- 705 51. Cameron, Ross W. F., and Tijana Blanuša. 2016. "Green Infrastructure and  
706 Ecosystem Services - Is the Devil in the Detail?," *Annals of Botany*, 118.3: 377–91
- 707 52. Christie, Mark R., and L. Lacey Knowles. 2015. "Habitat Corridors Facilitate Genetic  
708 Resilience Irrespective of Species Dispersal Abilities or Population  
709 Sizes," *Evolutionary Applications*, 8.5: 454–63
- 710 53. Jastrow, J. D., and R. M. Miller. 2018. "Soil Aggregate Stabilization and Carbon  
711 Sequestration: Feedbacks through Organomineral Associations," in *Soil Processes  
712 and the Carbon Cycle* (CRC Press), pp. 207–23

- 713 54. Young, I. M., and J. W. Crawford. 2004. "Interactions and Self-Organization in the  
714 Soil-Microbe Complex," *Science (New York, N.Y.)*, 304.5677: 1634–37
- 715 55. Torsvik, Vigdis, and Lise Øvreås. 2002. "Microbial Diversity and Function in Soil:  
716 From Genes to Ecosystems," *Current Opinion in Microbiology*, 5.3: 240–45
- 717 56. Filotas, Elise, Lael Parrott, Philip J. Burton, Robin L. Chazdon, K. David Coates, and  
718 others. 2014. "Viewing Forests through the Lens of Complex Systems  
719 Science," *Ecosphere (Washington, D.C.)*, 5.1: art1
- 720 57. Birch, Joseph D., Suzanne W. Simard, Kevin J. Beiler, and Justine Karst. 2021.  
721 "Beyond Seedlings: Ectomycorrhizal Fungal Networks and Growth of Mature  
722 *Pseudotsuga Menziesii*," *The Journal of Ecology*, 109.2: 806–18
- 723 58. Ibarra, José Tomás, Kristina L. Cockle, Tomás A. Altamirano, Yntze van der Hoek,  
724 Suzanne W. Simard, and others. 2020. "Nurturing Resilient Forest Biodiversity: Nest  
725 Webs as Complex Adaptive Systems," *Ecology and Society: A Journal of Integrative  
726 Science for Resilience and Sustainability*, 25.2 <[https://doi.org/10.5751/es-11590-  
727 250227](https://doi.org/10.5751/es-11590-250227)>
- 728 59. Soininen, Laura, Mira Grönroos, Marja I. Roslund, and Aki Sinkkonen. 2021. "Long-  
729 Term Storage Affects Resource Availability and Occurrence of Bacterial Taxa Linked  
730 to Pollutant Degradation and Human Health in Landscaping Materials," *Urban  
731 Forestry & Urban Greening*, 60.127065: 127065
- 732 60. Guillitte, O. 1995. "Bioreceptivity: A New Concept for Building Ecology  
733 Studies," *The Science of the Total Environment*, 167.1–3: 215–20
- 734 61. Cruz, Marcos, and Richard Beckett. 2016. "Bioreceptive Design: A Novel Approach  
735 to Biodigital Materiality," *Architectural Research Quarterly*, 20.1: 51–64
- 736 62. Manso, Sandra, María Ángeles Calvo-Torras, Nele De Belie, Ignacio Segura, and  
737 Antonio Aguado. 2015. "Evaluation of Natural Colonisation of Cementitious

- 738 Materials: Effect of Bioreceptivity and Environmental Conditions,” *The Science of the*  
739 *Total Environment*, 512–513: 444–53
- 740 63. Cruz, M. 2021. *Poikilohydric Living Walls - a Bartlett Research Folio*.
- 741 64. Sanmartín, P., A. Z. Miller, B. Prieto, and H. A. Viles. 2021. “Revisiting and  
742 Reanalysing the Concept of Bioreceptivity 25 Years On,” *The Science of the Total*  
743 *Environment*, 770.145314: 145314
- 744 65. Wilson, E. O. 1984. *Biophilia / Edward O. Wilson*. [Book]. Harvard University Press.
- 745 66. Yeager, Chris M., Jennifer L. Kornosky, Rachael E. Morgan, Elizabeth C. Cain,  
746 Ferran Garcia-Pichel, and others. 2007. “Three Distinct Clades of Cultured  
747 Heterocystous Cyanobacteria Constitute the Dominant N<sub>2</sub>-Fixing Members of  
748 Biological Soil Crusts of the Colorado Plateau, USA: N<sub>2</sub>-Fixing Cyanobacteria in  
749 Soil Crusts,” *FEMS Microbiology Ecology*, 60.1: 85–97
- 750 67. Tang, Jing Yan, Jing Ma, Xue Dong Li, and Yan Hong Li. 2016. “Illumina  
751 Sequencing-Based Community Analysis of Bacteria Associated with Different  
752 Bryophytes Collected from Tibet, China,” *BMC Microbiology*, 16.1: 276
- 753 68. Cao, Wei, Yuanxin Xiong, Degang Zhao, Hongying Tan, and Jiaojiao Qu. 2020.  
754 “Bryophytes and the Symbiotic Microorganisms, the Pioneers of Vegetation  
755 Restoration in Karst Rocky Desertification Areas in Southwestern China,” *Applied*  
756 *Microbiology and Biotechnology*, 104.2: 873–91
- 757 69. Insuk, Chadabhorn, Nattakorn Kuncharoen, Naowarat Cheeptham, Somboon  
758 Tanasupawat, and Wasu Pathom-Aree. 2020. “Bryophytes Harbor Cultivable  
759 Actinobacteria with Plant Growth Promoting Potential,” *Frontiers in Microbiology*,  
760 11: 563047



- 761 70. Hui, Nan, Mira Grönroos, Marja I. Roslund, Anirudra Parajuli, Heli K. Vari, and  
762 others. 2019. “Diverse Environmental Microbiota as a Tool to Augment Biodiversity  
763 in Urban Landscaping Materials,” *Frontiers in Microbiology*, 10: 536
- 764 71. Sacristán-Pérez-Minayo, Gonzalo, Domingo Javier López-Robles, Carlos Rad, and  
765 Luis Miranda-Barroso. 2020. “Microbial Inoculation for Productivity Improvements  
766 and Potential Biological Control in Sugar Beet Crops,” *Frontiers in Plant Science*, 11:  
767 604898
- 768 72. Kumar, A., J. S. Patel, V. S. Meena, and P. W. Ramteke. 2019. “Plant Growth-  
769 Promoting Rhizobacteria: Strategies to Improve Abiotic Stresses under Sustainable  
770 Agriculture,” *Journal of Plant Nutrition*, 42.11–12: 1402–15
- 771 73. Saleem, Muhammad, Jie Hu, and Alexandre Jousset. 2019. “More than the Sum of Its  
772 Parts: Microbiome Biodiversity as a Driver of Plant Growth and Soil Health,” *Annual  
773 Review of Ecology, Evolution, and Systematics*, 50.1: 145–68
- 774 74. Alharbi, Badr H., Mohammad J. Pasha, and Mohammed Ahmad S. Al-Shamsi. 2019.  
775 “Influence of Different Urban Structures on Metal Contamination in Two  
776 Metropolitan Cities,” *Scientific Reports*, 9.1: 4920
- 777 75. Jacquiod, Samuel, Valentine Cyriaque, Leise Riber, Waleed Abu Al-soud, David C.  
778 Gillan, and others. 2018. “Long-Term Industrial Metal Contamination Unexpectedly  
779 Shaped Diversity and Activity Response of Sediment Microbiome,” *Journal of  
780 Hazardous Materials*, 344: 299–307
- 781 76. Kang, Chang-Ho, Yoon-Jung Kwon, and Jae-Seong So. 2016. “Bioremediation of  
782 Heavy Metals by Using Bacterial Mixtures,” *Ecological Engineering*, 89: 64–69
- 783 77. Priyadarshane, Monika, and Surajit Das. 2021. “Biosorption and Removal of Toxic  
784 Heavy Metals by Metal Tolerating Bacteria for Bioremediation of Metal

- 785 Contamination: A Comprehensive Review,” *Journal of Environmental Chemical*  
786 *Engineering*, 9.1: 104686
- 787 78. Bergé, A., J. Gasperi, V. Rocher, L. Gras, A. Coursimault, and others. 2014.  
788 “Phthalates and Alkylphenols in Industrial and Domestic Effluents: Case of Paris  
789 Conurbation (France),” *The Science of the Total Environment*, 488–489: 26–35
- 790 79. Boll, Matthias, Robin Geiger, Madan Junghare, and Bernhard Schink. 2020.  
791 “Microbial Degradation of Phthalates: Biochemistry and Environmental  
792 Implications,” *Environmental Microbiology Reports*, 12.1: 3–15
- 793 80. Ojuederie, Omena, and Olubukola Babalola. 2017. “Microbial and Plant-Assisted  
794 Bioremediation of Heavy Metal Polluted Environments: A Review,” *International*  
795 *Journal of Environmental Research and Public Health*, 14.12: 1504
- 796 81. Borchert, Erik, Katrin Hammerschmidt, Ute Hentschel, and Peter Deines. 2021.  
797 “Enhancing Microbial Pollutant Degradation by Integrating Eco-Evolutionary  
798 Principles with Environmental Biotechnology,” *Trends in Microbiology*
- 799 82. Paungfoo-Lonhienne, Chanyarat, Yun Kit Yeoh, Naga Rup Pinaki Kasinadhuni,  
800 Thierry G. A. Lonhienne, Nicole Robinson, and others. 2015. “Nitrogen Fertilizer  
801 Dose Alters Fungal Communities in Sugarcane Soil and Rhizosphere,” *Scientific*  
802 *Reports*, 5.1: 8678
- 803 83. Hammad, Hafiz Mohkum, Abdul Khaliq, Farhat Abbas, Wajid Farhad, Shah Fahad,  
804 and others. 2020. “Comparative Effects of Organic and Inorganic Fertilizers on Soil  
805 Organic Carbon and Wheat Productivity under Arid Region,” *Communications in Soil*  
806 *Science and Plant Analysis*, 51.10: 1406–22
- 807 84. Dahunsi, S. O., S. Oranusi, V. E. Efeovbokhan, A. T. Adesulu-Dahunsi, and J. O.  
808 Ogunwole. 2021. “Crop Performance and Soil Fertility Improvement Using Organic

- 809 Fertilizer Produced from Valorization of Carica Papaya Fruit Peel,” *Scientific*  
810 *Reports*, 11.1: 4696
- 811 85. Gregory, A. S., K. Ritz, S. P. McGrath, J. N. Quinton, K. W. T. Goulding, and others.  
812 2015. “A Review of the Impacts of Degradation Threats on Soil Properties in the  
813 UK,” *Soil Use and Management*, 31.Suppl Suppl 1: 1–15
- 814 86. Chen, Qing-Lin, Xin-Li An, Bang-Xiao Zheng, Michael Gillings, Josep Peñuelas, and  
815 others. 2019. “Loss of Soil Microbial Diversity Exacerbates Spread of Antibiotic  
816 Resistance,” *Soil Ecology Letters*, 1.1–2: 3–13
- 817 87. WHO. 2020. *Anti-microbial resistance*. Online. Available at:  
818 [https://www.who.int/news-room/fact-sheets/detail/antimicrobial-](https://www.who.int/news-room/fact-sheets/detail/antimicrobial-resistance#:~:text=What%20is%20antimicrobial%20resistance%3F,sread%2C%20s)  
819 [resistance#:~:text=What%20is%20antimicrobial%20resistance%3F,sread%2C%20s](https://www.who.int/news-room/fact-sheets/detail/antimicrobial-resistance#:~:text=What%20is%20antimicrobial%20resistance%3F,sread%2C%20s)  
820 [evere%20illness%20and%20death..](https://www.who.int/news-room/fact-sheets/detail/antimicrobial-resistance#:~:text=What%20is%20antimicrobial%20resistance%3F,sread%2C%20s) (Accessed on 20 March 2021).
- 821 88. Kemp, Luke, David C. Aldridge, Olaf Booy, Hilary Bower, Des Browne, and others.  
822 2021. “80 Questions for UK Biological Security,” *PloS One*, 16.1: e0241190
- 823 89. EPPO. 2020. *European and Mediterranean Plant Protection Organisation*. Online.  
824 Available at: <https://www.eppo.int/index>. (Accessed on 28 March 2021).
- 825 90. Department for Environment, Food and Rural Affairs. 2014. UK Plant Health Risk  
826 Register. Online. Available at:  
827 <https://secure.fera.defra.gov.uk/phiw/riskRegister/downloadEntireRiskRegister.cfm>.  
828 (Accessed on 28 March 2021).
- 829 91. Public Health Agency of Canada 2018. *Biosecurity Program Annual Report 2017-*  
830 *2018*. Online. Available at: [https://www.canada.ca/en/public-](https://www.canada.ca/en/public-health/services/publications/health-risks-safety/centre-biosecurity-annual-report-2017-2018.html)  
831 [health/services/publications/health-risks-safety/centre-biosecurity-annual-report-](https://www.canada.ca/en/public-health/services/publications/health-risks-safety/centre-biosecurity-annual-report-2017-2018.html)  
832 [2017-2018.html](https://www.canada.ca/en/public-health/services/publications/health-risks-safety/centre-biosecurity-annual-report-2017-2018.html). (Accessed on 8 March 2021).

- 833 92. Watkins, H., and Arkell, T. 2019. *The Landscape Institute (LI) Biosecurity Toolkit for*  
834 *Landscape Consultants*. Online. Available at:  
835 <https://www.landscapeinstitute.org/technical-resource/biosecurity-toolkit/>. (Accessed  
836 on 8 March 2021).
- 837 93. Sequeira, Ron, and Robert Griffin. 2014. “The Biosecurity Continuum and Trade:  
838 Pre-Border Operations,” in *The Handbook of Plant Biosecurity* (Dordrecht: Springer  
839 Netherlands), pp. 119–48
- 840 94. Learning the Land. 2021. *Learning the Land*. Online. Available at:  
841 <https://www.learningtheland.ca>. (Accessed on 28 March 2021).
- 842 95. Simard, Suzanne W. 2018. “Mycorrhizal Networks Facilitate Tree Communication,  
843 Learning, and Memory,” in *Memory and Learning in Plants* (Cham: Springer  
844 International Publishing), pp. 191–213
- 845 96. Capaldi, Colin A., Raelyne L. Dopko, and John M. Zelenski. 2014. “The Relationship  
846 between Nature Connectedness and Happiness: A Meta-Analysis,” *Frontiers in*  
847 *Psychology*, 5: 976
- 848 97. Capaldi, Colin, Holli-Anne Passmore, Elizabeth Nisbet, John Zelenski, and Raelyne  
849 Dopko. 2015. “Flourishing in Nature: A Review of the Benefits of Connecting with  
850 Nature and Its Application as a Wellbeing Intervention,” *International Journal of*  
851 *Wellbeing*, 5.4: 1–16
- 852 98. Smith, B. 2020. *Planning with Nature: How can Authorities develop and implement a*  
853 *green infrastructure strategy?*. Practice Note. Online. London: Public Practice.  
854 Available at: <<http://www.publicpractice.org.uk/resources>> (Accessed 28 March  
855 2021).

- 856 99. Ferreira, Vera, Ana Barreira, Luís Loures, Dulce Antunes, and Thomas Panagopoulos.  
857 2020. “Stakeholders’ Engagement on Nature-Based Solutions: A Systematic Literature  
858 Review,” *Sustainability*, 12.2: 640
- 859 100. Greater London Authority. 2021. The London Plan: The Spatial Development  
860 Strategy For Greater London. Online. London: Greater London Authority, pp.312-333.  
861 Available at:  
862 <[https://www.london.gov.uk/sites/default/files/the\\_london\\_plan\\_2021.pdf](https://www.london.gov.uk/sites/default/files/the_london_plan_2021.pdf)> (Accessed  
863 28 March 2021).
- 864 101. Khoshkar, Sara, Berit Balfors, and Antoinette Wärnbäck. 2018. “Planning for Green  
865 Qualities in the Densification of Suburban Stockholm – Opportunities and  
866 Challenges,” *Journal of Environmental Planning and Management*, 61.14: 2613–35
- 867 102. Ugolini, Francesca, Giovanni Sanesi, Anna Steidle, and David Pearlmutter. 2018.  
868 “Speaking ‘Green’: A Worldwide Survey on Collaboration among Stakeholders in  
869 Urban Park Design and Management,” *Forests*, 9.8: 458

870

## 871 **Figure captions**

872

873 **Fig. 1.** Urban multispecies health. Environmental microbiomes are the foundations of our  
874 ecosystems, and are essential to plant and animal health (including humans).

875

876 **Fig. 2.** Actionable insights for MIGI, including vegetation complexity, downwind  
877 development and local integration of biodiverse source (a); a solution to the concept of  
878 vertical stratification (b); hands-on engagement with natural features to promote  
879 immunoregulation (c); recommended soil types to promote diverse microbial habitat and  
880 short-term storage of landscaping materials (d); revegetation with diverse native plants to

881 promote functional diversity (e); the concept of habitat connectivity via contiguous natural  
882 corridors to promote long-term multispecies health (f).

883

884 **Fig. 3.** Horizon scan of developmental considerations for MIGI, including interventions (b  
885 and d), design and supportive features (a and e), and applications for engagement and to  
886 acquire useful urban ecosystem health information (c and f).