

Omicron infection enhances Delta antibody immunity in vaccinated persons

<https://doi.org/10.1038/s41586-022-04830-x>

Received: 27 January 2022

Accepted: 4 May 2022

Published online: 6 May 2022

Open access

 Check for updates

Khadija Khan^{1,2,30}, Farina Karim^{1,2,30}, Sandile Cele^{1,2}, Kajal Reedy¹, James Emmanuel San³, Gila Lustig⁴, Houriiyah Tegally^{3,5}, Yuval Rosenberg⁶, Mallory Bernstein¹, Zesuliwe Jule¹, Yashica Ganga¹, Nokuthula Ngcobo¹, Matilda Mazibuko¹, Ntombifuthi Mthabela¹, Zoey Mhlane¹, Nikiwe Mbatha¹, Yoliswa Miya¹, Jennifer Giandhari³, Yajna Ramphal³, Taryn Naidoo¹, Aida Sivro^{4,7}, Natasha Samsunder⁴, Ayesha B. M. Kharsany⁴, Daniel Amoako⁸, Jinal N. Bhiman⁸, Nithendra Manickchund⁹, Quarraisha Abdool Karim^{4,10}, Nombulelo Magula¹¹, Salim S. Abdool Karim^{4,10}, Glenda Gray¹², Willem Hanekom^{1,13}, Anne von Gottberg^{8,14}, COMMIT-KZN Team*, Ron Milo⁶, Bernadett I. Gosnell⁹, Richard J. Lessells^{3,4}, Penny L. Moore^{4,8,15,16}, Tulio de Oliveira^{3,4,5,17}, Mahomed-Yunus S. Moosa⁹ & Alex Sigal^{1,2,4,18}✉

The extent to which Omicron infection^{1–9}, with or without previous vaccination, elicits protection against the previously dominant Delta (B.1.617.2) variant is unclear. Here we measured the neutralization capacity against variants of severe acute respiratory syndrome coronavirus 2 in 39 individuals in South Africa infected with the Omicron sublineage BA.1 starting at a median of 6 (interquartile range 3–9) days post symptom onset and continuing until last follow-up sample available, a median of 23 (interquartile range 19–27) days post symptoms to allow BA.1-elicited neutralizing immunity time to develop. Fifteen participants were vaccinated with Pfizer's BNT162b2 or Johnson & Johnson's Ad26.CoV2.S and had BA.1 breakthrough infections, and 24 were unvaccinated. BA.1 neutralization increased from a geometric mean 50% focus reduction neutralization test titre of 42 at enrolment to 575 at the last follow-up time point (13.6-fold) in vaccinated participants and from 46 to 272 (6.0-fold) in unvaccinated participants. Delta virus neutralization also increased, from 192 to 1,091 (5.7-fold) in vaccinated participants and from 28 to 91 (3.0-fold) in unvaccinated participants. At the last time point, unvaccinated individuals infected with BA.1 had low absolute levels of neutralization for the non-BA.1 viruses and 2.2-fold lower BA.1 neutralization, 12.0-fold lower Delta neutralization, 9.6-fold lower Beta variant neutralization, 17.9-fold lower ancestral virus neutralization and 4.8-fold lower Omicron sublineage BA.2 neutralization relative to vaccinated individuals infected with BA.1. These results indicate that hybrid immunity formed by vaccination and Omicron BA.1 infection should be protective against Delta and other variants. By contrast, infection with Omicron BA.1 alone offers limited cross-protection despite moderate enhancement.

The Omicron variant of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), first identified in November 2021 in South Africa and Botswana¹⁰, has been shown by us¹ and others^{2–9} to have extensive but incomplete escape from neutralizing immunity elicited by vaccines and previous infection, with boosted individuals showing better neutralization. In South Africa, Omicron infections led to a lower incidence

of severe disease relative to other variants^{11,12}, although this can be at least partly explained by pre-existing immunity¹³. The first Omicron sublineage to appear was BA.1, which was supplanted by the BA.2 sublineage in many countries¹⁴.

How Omicron BA.1 infection will interact with vaccination to protect against the previously dominant Delta variant, emerging variants such

¹Africa Health Research Institute, Durban, South Africa. ²School of Laboratory Medicine and Medical Sciences, University of KwaZulu-Natal, Durban, South Africa. ³KwaZulu-Natal Research Innovation and Sequencing Platform, Durban, South Africa. ⁴Centre for the AIDS Programme of Research in South Africa, Durban, South Africa. ⁵Centre for Epidemic Response and Innovation, School of Data Science and Computational Thinking, Stellenbosch University, Stellenbosch, South Africa. ⁶Department of Plant and Environmental Sciences, Weizmann Institute of Science, Rehovot, Israel. ⁷Department of Medical Microbiology, University of KwaZulu-Natal, Durban, South Africa. ⁸National Institute for Communicable Diseases of the National Health Laboratory Service, Johannesburg, South Africa. ⁹Department of Infectious Diseases, Nelson R. Mandela School of Clinical Medicine, University of KwaZulu-Natal, Durban, South Africa. ¹⁰Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, NY, USA. ¹¹Department of Internal Medicine, Nelson R. Mandela School of Medicine, University of KwaZulu-Natal, Durban, South Africa. ¹²South African Medical Research Council, Cape Town, South Africa. ¹³Division of Infection and Immunity, University College London, London, UK. ¹⁴School of Pathology, Faculty of Health Sciences, University of Witwatersrand, Johannesburg, South Africa. ¹⁵SAMRC Antibody Immunity Research Unit, School of Pathology, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa. ¹⁶Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Cape Town, South Africa. ¹⁷Department of Global Health, University of Washington, Seattle, WA, USA. ¹⁸Max Planck Institute for Infection Biology, Berlin, Germany. ³⁰These authors contributed equally: Khadija Khan, Farina Karim.

*A list of authors and their affiliations appears at the end of the paper. ✉e-mail: alex.sigal@ahri.org

as BA.2 and other variants is still unclear. We isolated live Omicron BA.1, Omicron BA.2, ancestral, Beta and Delta viruses and neutralized viruses with plasma from participants enrolled and longitudinally sampled during the Omicron BA.1 infection wave in South Africa, with all participants having a confirmed diagnosis of SARS-CoV-2 by quantitative PCR. To quantify neutralization, we used a live virus neutralization assay and calculated the focus reduction neutralization test (FRNT₅₀) titre, the inverse of the plasma dilution required for 50% neutralization, as measured by the reduction in the number of infection foci. We enrolled 41 participants who reported symptoms from late November 2021 to January 2022. We successfully sequenced the infecting virus in 26 participants, and all sequences corresponded to Omicron BA.1 (Extended Data Table 1). Two participants had advanced human immunodeficiency virus (HIV) disease on the basis of a low CD4 count (<200 cells μl⁻¹ throughout the study) and unsuppressed HIV infection, and we excluded these participants because of our previous data showing an atypical response to SARS-CoV-2 in advanced HIV disease¹⁵. Extended Data Table 2 summarizes the characteristics of the remaining 39 participants.

Of the 39 participants, 27 were admitted to hospital because of coronavirus disease 2019 symptoms. Seven required supplemental oxygen and one died. Fifteen participants were vaccinated and had a breakthrough Omicron BA.1 infection. The median time post vaccination was 139 days (interquartile range (IQR) 120–178), a time interval that would predict considerable waning of the vaccine-elicited immune response¹⁶, which may have contributed to the breakthrough infections. Eight participants were vaccinated with two doses of Pfizer's BNT162b2 and seven were vaccinated with Johnson & Johnson's Ad26. CoV2.S (six with a single dose and one with two doses; Extended Data Table 1). The length of hospital stay was shorter in the vaccinated (3.5 days) relative to unvaccinated (8 days; Extended Data Table 2) participants. Three participants self-reported having a previous SARS-CoV-2 infection (Extended Data Table 1).

Participants were sampled at enrolment at a median of 6 days (IQR 3–9 days) after symptom onset, and again at weekly follow-up visits that were attended as practicable because of the Christmas holidays in South Africa. The last follow-up visit was a median of 23 days (IQR 19–27 days) post-symptom onset (Extended Data Table 1). Examining neutralization at all available time points per study participant showed that neutralization of the Omicron BA.1 variant increased substantially in most participants from enrolment to the time of the last follow-up (Extended Data Fig. 1), consistent with developing a neutralizing antibody response to Omicron BA.1 infection. We therefore analysed neutralization at enrolment (baseline for the study) and the last follow-up visit to quantify the increase in neutralization capacity after Omicron infection.

We observed that Omicron BA.1 neutralization increased in vaccinated individuals from a low geometric mean titre (GMT) FRNT₅₀ of 42 at the enrolment visit to 575 at the last follow-up visit about 2 to 3 weeks later, a 13.6-fold change (95% CI confidence interval (CI) 3.7–50.2; Fig. 1a). The samples from unvaccinated participants at the study baseline neutralized Omicron BA.1 at a similar starting level of 46 and reached a final level of 272 at the last follow-up, a 6.0-fold increase (95% CI 2.2–16.1; Fig. 1b). Neutralization of the Delta virus also increased during this period. At enrolment, neutralization capacity against the Delta virus was 192 and reached a final level of 1,091 at the last follow-up visit in vaccinated participants, a 5.7-fold increase (95% CI 1.7–18.4; Fig. 1c). Unvaccinated participants had lower Delta neutralization at baseline with Delta virus FRNT₅₀ = 28, and reached FRNT₅₀ = 91, a 3.2-fold increase (95% CI 1.3–8.1; Fig. 1d).

We next compared Omicron BA.1 to Omicron BA.2, Delta, Beta (ref.¹⁷) and ancestral virus neutralization at the last available follow-up visit in three sets of paired experiments, each comparing Omicron BA.2, Delta or ancestral and Beta virus neutralization to Omicron BA.1 neutralization. The range of Omicron BA.1 neutralization shown in Fig. 2a




Fig. 1 | Enhancement of Delta neutralization by Omicron infection.

a, b, Neutralization of the Omicron BA.1 virus over time for $n = 15$ vaccinated (**a**) and $n = 24$ unvaccinated (**b**) participants infected with Omicron BA.1. **c, d,** Neutralization of the Delta virus over time for the same vaccinated ($n = 15$; **c**) and unvaccinated ($n = 24$; **d**) participants as in **a, b**. For each participant, the sample collected at the initial enrolment visit (median 6 days post symptom onset) was compared with that collected at the last follow-up visit (median 23 days post symptom onset). The neutralization capacity per participant was determined in two independent experiments, and the numbers and horizontal bars are GMTs over all participants per group of the reciprocal plasma dilution (FRNT₅₀) resulting in 50% neutralization. The fold change was calculated by dividing the GMT from the follow-up by the GMT from the enrolment visit. The dashed line is the most concentrated plasma tested. The P values were determined by a left-sided Wilcoxon rank sum test measuring the significance of the increase; ** $P = 0.01$ – 0.001 ; NS, not significant. The exact P values are 0.0012 (**a**), 0.0081 (**b**), 0.0021 (**c**) and 0.11 (**d**).

for different experiments (FRNT₅₀ = 516 to 646 for vaccinated samples and 266 to 271 for unvaccinated samples) is the result of experimental variation. BA.2 neutralization was moderately and not significantly lower relative to BA.1 neutralization in both vaccinated and unvaccinated participants. Testing only participants with sequence-confirmed Omicron BA.1 infection gave a similar result (Extended Data Fig. 2). The trend for the other variants and the ancestral virus was that neutralization was higher relative to Omicron BA.1 in vaccinated participants but lower relative to Omicron BA.1 in unvaccinated participants, although the differences were mostly not significant (Fig. 2a). As a result of the relatively moderate fold change, higher participant numbers would probably be required to make the trends statistically significant.

The comparison of the other variants to Omicron BA.1 within the vaccinated or unvaccinated group does not indicate the differences in neutralization capacity elicited by Omicron BA.1 between the vaccinated and unvaccinated participants. We therefore compared neutralization of each variant between the vaccinated and unvaccinated groups at the last time point directly (Fig. 2b). The smallest difference between vaccinated and unvaccinated participants was in neutralization of Omicron BA.1, the infecting variant, with the vaccinated participants showing 2.2-fold higher neutralization. For the other variants, neutralization was higher in vaccinated participants by a factor of 4.8-fold for Omicron BA.2, 9.6-fold for Beta, 12.0-fold for Delta and 17.9-fold for ancestral (Fig. 2b). All differences were significant, and the 95% CIs for the GMT FRNT₅₀ of vaccinated and unvaccinated participants did not overlap for BA.2, Beta, Delta or ancestral virus neutralization (Fig. 2b). For the unvaccinated participants, the absolute neutralization capacity against the BA.2, Beta, Delta and ancestral viruses was low¹⁸, with GMT FRNT₅₀ being about or below FRNT₅₀ = 100 (Fig. 2b).




Fig. 2 | Gap in neutralizing immunity between vaccinated and unvaccinated participants infected with Omicron BA.1. **a**, Neutralization of Omicron BA.2, Beta, Delta and ancestral (with the D614G substitution) viruses compared to the Omicron BA.1 virus at the last available follow-up time point in $n = 15$ vaccinated or $n = 24$ unvaccinated participants infected with Omicron BA.1. The neutralization capacity per participant was determined in two independent experiments, and the numbers and horizontal bars are GMT $FRNT_{50}$. The fold change was calculated by dividing the larger by the smaller GMT. The dashed line is the most concentrated plasma tested. The P values were determined by a two-sided Wilcoxon rank sum test; * $P = 0.05-0.01$; NS, not significant. The exact P values (vaccinated/unvaccinated) are: 0.22/0.087

for BA.2, 0.36/0.071 for Beta, 0.15/0.25 for Delta and 0.014/0.20 for ancestral. **b**, Comparison of the neutralization capacity against the Omicron BA.1, Omicron BA.2, Beta, Delta and ancestral (D614G) viruses in vaccinated ($n = 15$) versus unvaccinated ($n = 24$) participants infected with Omicron BA.1. The neutralization capacity per participant was determined in two independent experiments for all strains except for Omicron BA.1, for which six experiments were available and were used in the calculation. The points represent GMT $FRNT_{50}$ per group and the error bars are GMT 95% CIs. The P values were determined by a two-sided Wilcoxon rank sum test; * $P = 0.05-0.01$; ** $P = 0.01-0.001$; *** $P = 0.001-0.0001$. The exact P values are 0.025 (BA.1), 0.0026 (BA.2), 4.1×10^{-4} (Beta), 0.0012 (Delta) and 3.3×10^{-4} (ancestral).

We also tested neutralization of Omicron BA.1 by Delta-variant-elicited immunity. We collected 18 plasma samples from 14 participants (including pre-vaccination and post-vaccination samples from 4 participants) previously infected in the Delta variant wave in South Africa, 8 of whom were vaccinated either before or after infection (Extended Data Table 3). Confirming previously reported results¹⁹, we observed similar escape of Omicron BA.1 from Delta-elicited immunity across all samples tested, manifested as a 22.5-fold decrease (95% CI 14.4–35.0) in Omicron BA.1 neutralization compared to Delta virus neutralization (Fig. 3).

The large fold drop in Delta-infection-elicited neutralization capacity against Omicron BA.1 contrasts with the moderate and nonsignificant fold drops, or even fold increases, in neutralization of other variants relative to Omicron BA.1 in individuals infected with Omicron BA.1. However, in unvaccinated individuals, even though fold drops in neutralization were moderate and nonsignificant, the absolute levels of neutralization of the other variants, and of Omicron BA.1 itself, were low and on a similar scale to the cross-neutralization capacity against Omicron in Delta-infection-elicited immunity. This is consistent with other recently reported results²⁰, and possibly indicates that Omicron is poorly immunogenic. In agreement with recent reports^{21,22}, our observations show moderately and nonsignificantly lower neutralization of BA.2 by BA.1-elicited immunity. The results explain epidemiological observations showing that Omicron BA.2 reinfection is relatively rare soon after Omicron BA.1 infection^{23,24}.

Our results may be supportive of a scenario in which hybrid immunity formed by Omicron infection combined with vaccination protects as well or better against reinfection with variants such as Delta relative to reinfection with Omicron itself. By contrast, unvaccinated participants infected with Omicron BA.1 only, have low neutralization capacity against the Omicron BA.2, Beta, Delta and ancestral viruses.

Limitations of this study include heterogeneity in participant immune history, including two vaccination types and one boost. On the basis of the high seroprevalence observed in South Africa^{25,26}, some participants may also have had unreported previous infection. However, including two vaccine types did not mask the differences between vaccinated and unvaccinated participants, and the low levels of neutralization in unvaccinated participants against the ancestral, Beta and Delta viruses (the dominant strains in the preceding South African infection waves) support the notion that these participants were either not previously infected, or that immunity has waned completely. Participants were also mostly hospitalized, which may not be typical of Omicron infection^{13,27}. Increased disease severity has been shown to lead to higher anti-SARS-CoV-2 antibody titres²⁸. This should help in the detection of the neutralization response, but whether it would affect the trend we observed is unclear. Omicron infection is unlikely benign to the extent that hospitalization is an outlier outcome: in the USA, the number of individuals with coronavirus disease 2019 who died in the Omicron wave was similar to the number who died in the Delta




Fig. 3 | Escape of Omicron virus from Delta-infection-elicited immunity. Neutralization of Delta compared to the Omicron BA.1 virus by Delta-infection-elicited plasma immunity in vaccinated and unvaccinated participants. A total of 18 samples were tested from $n = 14$ participants infected during the Delta infection wave in South Africa. The neutralization capacity per participant was determined in two independent experiments, and the numbers and horizontal bars are GMT FRNT₅₀. The fold change was calculated by dividing the larger by the smaller GMT. The dashed line is the most concentrated plasma tested. **** $P = 3.2 \times 10^{-7}$ as determined by a two-sided Wilcoxon rank sum test.

wave²⁷. Neutralizing immunity may have increased further in some participants had we sampled later: the neutralizing capacity did not plateau at the last time point in 8 of the 24 (33%) unvaccinated participants (participants 10, 14, 21, 27, 30, 31, 34 and 38; Extended Data Fig. 1) and 6 of the 15 (40%) vaccinated participants (participants 4, 6, 15, 16, 25 and 26). Therefore, the temporal dynamics give no clear indication that the immunity in the unvaccinated participants was delayed and would have reached similar levels to that of vaccinated participants if sampled later. However, the consequences of waning immunity several months post Omicron infection should be investigated.

The gap in immunity between unvaccinated individuals infected with Omicron BA.1 and vaccinated individuals with BA.1 breakthrough infection is concerning. Especially as immunity wanes, unvaccinated individuals post Omicron infection are likely to have poor cross-protection against existing and possibly emerging SARS-CoV-2 variants, despite acquiring some neutralizing immunity to the infecting Omicron BA.1 sub-lineage variant. The implication may be that Omicron BA.1 infection alone is not sufficient for protection and vaccination should be administered even in areas with a high prevalence of Omicron infection to protect against other variants.

Online content

Any methods, additional references, Nature Research reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41586-022-04830-x>.

- Cele, S. et al. Omicron extensively but incompletely escapes Pfizer BNT162b2 neutralization. *Nature* <https://doi.org/10.1038/s41586-021-04387-1> (2021).
- Andrews, N. et al. Covid-19 vaccine effectiveness against the omicron (B.1.1.529) variant. *N. Engl. J. Med.* **386**, 1532–1546 (2022).
- Cao, Y. et al. Omicron escapes the majority of existing SARS-CoV-2 neutralizing antibodies. *Nature* **602**, 657–663 (2022).
- Dejnirattisai, W. et al. SARS-CoV-2 Omicron-B.1.1.529 leads to widespread escape from neutralizing antibody responses. *Cell* **185**, 467–484 (2022).
- Dejnirattisai, W. et al. Reduced neutralisation of SARS-CoV-2 omicron B.1.1.529 variant by post-immunisation serum. *Lancet* **399**, 234–236 (2022).
- Garcia-Beltran, W. F. et al. mRNA-based COVID-19 vaccine boosters induce neutralizing immunity against SARS-CoV-2 Omicron variant. *Cell* **185**, 457–466 (2022).

- Lu, L. et al. Neutralization of SARS-CoV-2 Omicron variant by sera from BNT162b2 or Coronavac vaccine recipients. *Clin. Infect. Dis.* <https://doi.org/10.1093/cid/ciab1041> (2021).
- Planas, D. et al. Considerable escape of SARS-CoV-2 Omicron to antibody neutralization. *Nature* **602**, 671–675 (2022).
- VanBlargan, L. A. et al. An infectious SARS-CoV-2 B.1.1.529 Omicron virus escapes neutralization by therapeutic monoclonal antibodies. *Nat. Med.* **28**, 490–495 (2022).
- Viana, R. et al. Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in southern Africa. *Nature* <https://doi.org/10.1038/d41586-021-03832-5> (2021).
- Davies, M.-A. et al. Outcomes of laboratory-confirmed SARS-CoV-2 infection in the Omicron-driven fourth wave compared with previous waves in the Western Cape Province, South Africa. *Trop. Med. Int. Health* <https://doi.org/10.1111/tmi.13752> (2022).
- Wolter, N. et al. Early assessment of the clinical severity of the SARS-CoV-2 omicron variant in South Africa: a data linkage study. *Lancet* **399**, 437–446 (2022).
- Sigal, A. Milder disease with Omicron: is it the virus or the pre-existing immunity? *Nat. Rev. Immunol.* **22**, 69–71 (2022).
- Lyngse, F. P. et al. Transmission of SARS-CoV-2 Omicron VOC subvariants BA.1 and BA.2: evidence from Danish households. Preprint at *medRxiv* <https://doi.org/10.1101/2022.01.28.222770044> (2022).
- Cele, S. et al. SARS-CoV-2 prolonged infection during advanced HIV disease evolves extensive immune escape. *Cell Host Microbe* **30**, 154–162 (2022).
- Goldberg, Y. et al. Waning immunity after the BNT162b2 vaccine in Israel. *N. Engl. J. Med.* **385**, e85 (2021).
- Cele, S. et al. Escape of SARS-CoV-2 501Y.V2 from neutralization by convalescent plasma. *Nature* **593**, 142–146 (2021).
- Khouri, D. S. et al. Neutralizing antibody levels are highly predictive of immune protection from symptomatic SARS-CoV-2 infection. *Nat. Med.* **27**, 1205–1211 (2021).
- Rössler, A., Riepler, L., Bante, D., von Laer, D. & Kimpel, J. SARS-CoV-2 omicron variant neutralization in serum from vaccinated and convalescent persons. *N. Engl. J. Med.* **386**, 698–700 (2022).
- Servellita, V. et al. Neutralizing immunity in vaccine breakthrough infections from the SARS-CoV-2 Omicron and Delta variants. *Cell* **185**, 1539–1548 (2022).
- Yu, J. et al. Neutralization of the SARS-CoV-2 Omicron BA.1 and BA.2 Variants. *N. Engl. J. Med.* **386**, 1579–1580 (2022).
- Richardson, S. I. et al. SARS-CoV-2 Omicron triggers cross-reactive neutralization and Fc effector functions in previously vaccinated, but not unvaccinated, individuals. *Cell Host Microbe* <https://doi.org/10.1016/j.chom.2022.03.029> (2022).
- Chermaitelly, H. et al. Protection of Omicron sub-lineage infection against reinfection with another Omicron sub-lineage. Preprint at *medRxiv* <https://doi.org/10.1101/2022.02.24.22271440> (2022).
- Stegger, M. et al. Occurrence and significance of Omicron BA.1 infection followed by BA.2 reinfection. Preprint at *medRxiv* <https://doi.org/10.1101/2022.02.19.22271112> (2022).
- Khan, K. et al. Immunogenicity of SARS-CoV-2 infection and Ad26.CoV2.S vaccination in people living with HIV. *Clin. Infect. Dis.* <https://doi.org/10.1093/cid/ciab1008> (2021).
- Madhi, S. A. et al. Population immunity and Covid-19 severity with Omicron Variant in South Africa. *N. Engl. J. Med.* <https://doi.org/10.1056/NEJMOa2119658> (2022).
- Sigal, A., Milo, R. & Jassat, W. Estimating disease severity of Omicron and Delta SARS-CoV-2 infections. *Nat. Rev. Immunol.* **22**, 267–269 (2022).
- Smith, M. et al. Age, disease severity and ethnicity influence humoral responses in a multi-ethnic COVID-19 cohort. *Viruses* **13**, 786 (2021).

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022

COMMIT-KZN Team

Rohen Harrichandparsad¹⁹, Kobus Herbst¹²⁰, Prakash Jeena²¹, Thandeka Khoza¹, Henrik Kleverpris¹²², Alasdair Leslie¹¹³, Rajhmun Madansein²³, Mohlopheni Marakalala¹¹³, Mosa Moshabela²⁴, Kogie Naidoo⁴, Zaza Ndhlovu¹²⁵, Thumbi Ndung'u^{118,25,26}, Kennedy Nyamande²⁷, Vinod Patel²⁸, Theresa Smit¹, Adrie Steyn¹²⁹ & Emily Wong¹²⁹

- ¹⁹Department of Neurosurgery, University of KwaZulu-Natal, Durban, South Africa. ²⁰South African Population Research Infrastructure Network, Durban, South Africa. ²¹Department of Paediatrics and Child Health, University of KwaZulu-Natal, Durban, South Africa. ²²Department of Immunology and Microbiology, University of Copenhagen, Copenhagen, Denmark. ²³Department of Cardiothoracic Surgery, University of KwaZulu-Natal, Durban, South Africa. ²⁴College of Health Sciences, University of KwaZulu-Natal, Durban, South Africa. ²⁵Ragon Institute of MGH, MIT and Harvard, Boston, MA, USA. ²⁶HIV Pathogenesis Programme, The Doris Duke Medical Research Institute, University of KwaZulu-Natal, Durban, South Africa. ²⁷Department of Pulmonology and Critical Care, University of KwaZulu-Natal, Durban, South Africa. ²⁸Department of Neurology, University of KwaZulu-Natal, Durban, South Africa. ²⁹Division of Infectious Diseases, University of Alabama at Birmingham, Birmingham, AL, USA.

Article

Methods

Informed consent and ethics

Blood samples and the Delta isolate were obtained after written informed consent from adults with PCR-confirmed SARS-CoV-2 infection who were enrolled in a prospective cohort study at the Africa Health Research Institute approved by the Biomedical Research Ethics Committee at the University of KwaZulu-Natal (reference BREC/00001275/2020). Omicron BA.1 was isolated from a residual swab sample for SARS-CoV-2 where isolation from the sample was approved by the University of the Witwatersrand Human Research Ethics Committee (HREC; reference M210752). The sample to isolate Omicron BA.2 was collected after written informed consent as part of the study “COVID-19 transmission and natural history in KwaZulu-Natal, South Africa: Epidemiological Investigation to Guide Prevention and Clinical Care” of the Centre for the AIDS Programme of Research in South Africa (CAPRISA) and isolation from the sample approved by the Biomedical Research Ethics Committee at the University of KwaZulu-Natal (reference BREC/00001195/2020, BREC/00003106/2021).

Reagent availability

Virus isolates and the cell line are available from the corresponding author. A Biosafety Level 3 facility is required for laboratories receiving live SARS-CoV-2.

Whole-genome sequencing, genome assembly and phylogenetic analysis

RNA was extracted on an automated Chemagic 360 instrument, using the CMG-1049 kit (Perkin Elmer). The RNA was stored at -80°C before use. Libraries for whole-genome sequencing were prepared using either the Oxford Nanopore Midnight protocol with Rapid Barcoding or the Illumina COVIDseq Assay. For the Illumina COVIDseq assay, the libraries were prepared according to the manufacturer's protocol. Briefly, amplicons were tagged, followed by indexing using the Nextera UD Indexes Set A. Sequencing libraries were pooled, normalized to 4 nM and denatured with 0.2 N sodium acetate. An 8 pM sample library was spiked with 1% PhiX (PhiX Control v3 adaptor-ligated library used as a control). We sequenced libraries on a 500-cycle v2 MiSeq Reagent Kit on the Illumina MiSeq instrument (Illumina). On the Illumina NextSeq 550 instrument, sequencing was performed using the Illumina COVIDseq protocol (Illumina), an amplicon-based next-generation sequencing approach. The first-strand synthesis was carried using random hexamer primers from Illumina, and the synthesized cDNA underwent two separate multiplex PCR reactions. The pooled PCR-amplified products were processed for tagmentation and adaptor ligation using IDT for Illumina Nextera UD Indexes. Further enrichment and cleanup was performed as per protocols provided by the manufacturer (Illumina). Pooled samples were quantified using a Qubit 3.0 or 4.0 fluorometer (Invitrogen) through the Qubit dsDNA High Sensitivity Assay according to the manufacturer's instructions. The fragment sizes were analysed using TapeStation 4200 (Invitrogen). The pooled libraries were further normalized to 4 nM concentration and 25 μl of each normalized pool containing unique index adaptor sets was combined in a new tube. The final library pool was denatured and neutralized with 0.2 N sodium hydroxide and 200 mM Tris-HCl (pH 7), respectively. A 1.5 pM sample library was spiked with 2% PhiX. Libraries were loaded onto a 300-cycle NextSeq 500/550 HighOutput Kit v2 and run on the Illumina NextSeq 550 instrument (Illumina). For Oxford Nanopore sequencing, the Midnight primer kit was used: cDNA synthesis was performed on the extracted RNA using LunaScript RT mastermix (New England BioLabs) followed by gene-specific multiplex PCR using the Midnight Primer pools that produce 1,200-base-pair amplicons that overlap to cover the 30-kb SARS-CoV-2 genome. Amplicons from each pool were pooled and used neat for barcoding with the Oxford Nanopore Rapid Barcoding kit as per the manufacturer's protocol. Barcoded

samples were pooled and bead purified. After the bead cleanup, the library was loaded on a prepared R9.4.1 flow cell. A GridION XS or MinION sequencing run was initiated using MinKNOW software with the base-call setting switched off. We assembled paired-end and nanopore.fastq reads using Genome Detective 1.132 (<https://www.genomedetective.com>), which was updated for the accurate assembly and variant calling of tiled primer amplicon Illumina or Oxford Nanopore reads, and the Coronavirus Typing Tool. For Illumina assembly, the GATK HaploTypeCaller `-min-pruning 0` argument was added to increase mutation calling sensitivity near sequencing gaps. For Nanopore, low-coverage regions with poor alignment quality (<85% variant homogeneity) near sequencing/amplicon ends were masked to be robust against primer drop-out experienced in the spike gene, and the sensitivity for detecting short inserts using a region-local global alignment of reads was increased. In addition, we also used the `wf_artic` (ARTIC SARS-CoV-2) pipeline as built using the nextflow workflow framework. In some instances, mutations were confirmed visually with .bam files using Geneious software V2020.1.2 (Biomatters). The reference genome used throughout the assembly process was NC_045512.2 (numbering equivalent to MN908947.3). For lineage classification, we used the widespread dynamic lineage classification method from the Phylogenetic Assignment of Named Global Outbreak Lineages software suite (<https://github.com/hCoV-2019/pangolin>).

Cells

Vero E6 cells (originally ATCC CRL-1586, obtained from Cellnex in South Africa) were propagated in complete growth medium consisting of Dulbecco's modified Eagle medium with 10% fetal bovine serum (Hyclone) containing 10 mM HEPES, 1 mM sodium pyruvate, 2 mM L-glutamine and 0.1 mM nonessential amino acids (Sigma-Aldrich). Vero E6 cells were passaged every 3–4 days. The H1299-E3 cell line (H1299 originally from ATCC as CRL-5803) was propagated in growth medium consisting of complete Roswell Park Memorial Institute 1640 medium with 10% fetal bovine serum containing 10 mM HEPES, 1 mM sodium pyruvate, 2 mM L-glutamine and 0.1 mM nonessential amino acids. Cells were passaged every second day. The H1299-E3 (H1299-ACE2, clone E3) cell line was derived from H1299 as described in our previous work^{1,17}. Cell lines were not authenticated. Cell lines tested negative for mycoplasma contamination.

Virus expansion

All work with live virus was performed in Biosafety Level 3 containment using protocols for SARS-CoV-2 approved by the Africa Health Research Institute Biosafety Committee. ACE2-expressing H1299-E3 cells were seeded at 4.5×10^5 cells in a 6-well plate well and incubated for 18–20 h. After one Dulbecco's phosphate-buffered saline (PBS) wash, the subconfluent cell monolayer was inoculated with 500 μl universal transport medium diluted 1:1 with growth medium filtered through a 0.45- μm filter. Cells were incubated for 1 h. Wells were then filled with 3 ml complete growth medium. After 4 days of infection (completion of passage 1 (P1)), cells were trypsinized, centrifuged at 300 RCF for 3 min and resuspended in 4 ml growth medium. All cells from the P1 infection were added to Vero E6 cells that had been seeded at 2×10^5 cells ml^{-1} , 20 ml total, 18–20 h earlier in a T75 flask for cell-to-cell infection. The coculture of ACE2-expressing H1299-E3 and Vero E6 cells was incubated for 1 h, and the flask was filled with 20 ml of complete growth medium and incubated for 4 days. The viral supernatant from this culture (passage 2 (P2) stock) was used for experiments.

Live virus neutralization assay

H1299-E3 cells were plated in a 96-well plate (Corning) at 30,000 cells per well 1 day pre-infection. Plasma was separated from EDTA-anticoagulated blood by centrifugation at 500 RCF for 10 min and stored at -80°C . Aliquots of plasma samples were heat-inactivated at 56°C for 30 min and clarified by centrifugation at 10,000 RCF for

5 min. Virus stocks were used at approximately 50–100 focus-forming units per microwell and added to diluted plasma. Antibody–virus mixtures were incubated for 1 h at 37 °C, 5% CO₂. Cells were infected with 100 µl of the virus–antibody mixtures for 1 h, and then 100 µl of a 1× Roswell Park Memorial Institute 1640 (Sigma-Aldrich, R6504), 1.5% carboxymethylcellulose (Sigma-Aldrich, C4888) overlay was added without removing the inoculum. Cells were fixed 18 h after infection using 4% paraformaldehyde (Sigma-Aldrich) for 20 min. Foci were stained with a rabbit anti-spike monoclonal antibody (BS-R2B12, GenScript A02058) at 0.5 µg ml⁻¹ in a permeabilization buffer containing 0.1% saponin (Sigma-Aldrich), 0.1% BSA (Sigma-Aldrich) and 0.05% Tween-20 (Sigma-Aldrich) in PBS. Plates were incubated with primary antibody overnight at 4 °C, and then washed with wash buffer containing 0.05% Tween-20 in PBS. Secondary goat anti-rabbit antibody conjugated to horseradish peroxidase (Abcam ab205718) was added at 1 µg ml⁻¹ and incubated for 2 h at room temperature with shaking. TrueBlue peroxidase substrate (SeraCare 5510-0030) was then added at 50 µl per well and incubated for 20 min at room temperature. Plates were imaged in an ImmunoSpot Ultra-V S6-02-6140 Analyzer ELISPOT instrument with BioSpot Professional built-in image analysis (Cellular Technology Limited).

Statistics and fitting

Statistical methods were not used to predetermine sample size, and blinding and randomization were not used. All statistics and fitting were performed using custom code in MATLAB v.2019b. Neutralization data were fitted to:

$$Tx = 1/1 + (D/ID_{50}).$$

Here Tx is the number of foci normalized to the number of foci in the absence of plasma on the same plate at dilution *D*, and ID₅₀ is the plasma dilution giving 50% neutralization. FRNT₅₀ = 1/ID₅₀. Values of FRNT₅₀ < 1 are set to 1 (undiluted), the lowest measurable value. We note that the most concentrated plasma dilution was 1:25 and therefore FRNT₅₀ < 25 data were extrapolated. To calculate CIs, FRNT₅₀ or fold change in FRNT₅₀ per participant was log transformed and the arithmetic mean plus 2 s.d. and the arithmetic mean minus 2 s.d. were

calculated for the log-transformed values. These were exponentiated to obtain the upper and lower 95% CIs on the geometric mean FRNT₅₀ or the fold change in FRNT₅₀ geometric means.

Reporting summary

Further information on research design is available in the Nature Research Reporting Summary linked to this paper.

Data availability

Sequences of outgrown Omicron sublineages have been deposited in GISAID (<https://www.gisaid.org/>) with accessions EPI_ISL_7886688 (Omicron BA.1), EPI_ISL_9082893 (Omicron BA.2) and EPI_ISL_602626.1 (ancestral/D614G). Delta (EPI_ISL_3118687) and Beta (EPI_ISL_678615) isolates have been described previously¹⁵. Raw images of the data are available upon reasonable request. Source data are provided with this paper.

Acknowledgements This study was supported by the Bill and Melinda Gates award INV-018944 (A.Sigal), the National Institutes of Health award R01 AI138546 (A.Sigal), South African Medical Research Council awards including 6084COAP2020 (A.Sigal, T.d.O. and P.L.M.), the UK Foreign, Commonwealth and Development Office and the Wellcome Trust (grant number 221003/Z/20/Z; P.L.M.). P.L.M. is also supported by the South African Research Chairs Initiative of the Department of Science and Innovation and the NRF (grant number 98341). The funders had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

Author contributions A.Sigal., K.K. and F.K. conceived the study and designed the study and experiments. A.v.G., Q.A.K., S.S.A.K., G.L., A.Sivro and N.S. identified and provided virus samples. M.-Y.S.M., F.K., B.I.G., M.B., K.K., T.N., M.M., N.Mthabela, Z.M., N.N., Y.M., N.Mbatha, N.Manickchund, N.Magula, Z.J., K.R. and Y.G. set up and managed the cohort and cohort data. K.K., Z.J., K.R., S.C., H.T., J.E.S., Y.G., J.G., Y. Ramphal, A.B.M.K., D.A. and J.N.B. performed experiments and sequence analysis with input from A.Sigal, T.d.O., R.J.L. and J.N.B., A.Sigal, K.K., F.K., R.M. and Y. Rosenberg interpreted data with input from M.-Y.S.M., G.G., S.S.A.K., W.H., T.d.O., N. Magula, R.J.L. and P.L.M., A.Sigal, K.K., G.L., F.K. and M.B. prepared the manuscript with input from all authors.

Competing interests The authors declare no competing interests.


Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41586-022-04830-x>.

Correspondence and requests for materials should be addressed to Alex Sigal.


Peer review information Nature thanks the anonymous reviewers for their contribution to the peer review of this work.

Reprints and permissions information is available at <http://www.nature.com/reprints>.



Extended Data Fig. 1 | Longitudinal Omicron/BA.1 and Delta neutralization capacity in Omicron/BA.1 infected participants. Neutralization of Omicron (blue) and Delta (red) at all study visits. Participant number is as in Extended Data Table 1. Top three rows are participants vaccinated with Pfizer BNT162b2 (n = 8) or Johnson and Johnson Ad26.CoV2.S (n = 7) and bottom five rows are

unvaccinated participants (n = 24). X-axis is the time post-symptom onset when sample was collected, and y-axis is neutralization as FRNT₅₀. Dashed line is the most concentrated plasma tested (LOQ, limit of quantification below which FRNT₅₀ values are extrapolated). All participants recovered except participant 29, who died.



Extended Data Fig. 2 | Fold-drop in BA.2 versus BA.1 neutralization in all and sequence confirmed samples. Neutralization of Omicron BA.2 compared to BA.1 in participants described in Extended Data Table 1, excluding participant 14 for technical reasons and participants 40 and 41 because of advanced HIV disease. Left panel shows neutralization capacity in all n = 38

participants and right panel shows neutralization capacity for n = 25 participants where infection was successfully sequenced and determined to be BA.1. Dashed line is the most concentrated plasma tested. p-values were 0.077 for all and 0.15 for BA.1 sequence confirmed participants as determined by a two-sided Wilcoxon rank sum test. ns, not significant.

Article

Extended Data Table 1 | Characteristics of Omicron/BA.1 infected participants

| # | Age | Sex | Vacc. type | Date of vacc. | Vacc. to enrol. (days) | Date symp. onset | Ct at enrol. | Symp. to enrol. (days) | Symp. to last follow-up (days) | Max CD4 | Sub-lineage | Seq. GISAID ID | Supp. O ₂ | Hosp. |
|----|-------|-----|-------------------------|---------------|------------------------|-----------------------|--------------|------------------------|--------------------------------|---------|-------------|-----------------|----------------------|-------|
| 1 | 30-39 | M | AD26.COV2 AD26.COV2* | MAR-2021 | 274 | DEC-2021* | 24.9 | 1 | 23 | 1071 | BA.1 | EPI_ISL_9967759 | No | No |
| 2 | 30-39 | M | * | NOV-2021 | 14 | NOV-2021 | 14.5 | 1 | 22 | 789 | BA.1 | EPI_ISL_9967761 | No | No |
| 3 | 50-59 | F | BNT162b2 | JUL-2021 | 138 | DEC-2021 | 16.8 | 4 | 27 | 777 | BA.1 | EPI_ISL_8604915 | No | No |
| 4 | 30-39 | F | AD26.COV2 | MAY-2021 | 210 | DEC-2021 | 30.7 | 3 | 13 | 1169 | BA.1 | EPI_ISL_8604910 | No | No |
| 5 | 20-29 | F | AD26.COV2 | SEP-2021 | 89 | DEC-2021 | 23.9 | 5 | 27 | 1220 | BA.1 | EPI_ISL_9967760 | No | Yes |
| 6 | 10-19 | F | BNT162b2 | JUL-2021 | 157 | DEC-2021 | 23.1 | 6 | 12 | 732 | BA.1 | EPI_ISL_8604906 | No | Yes |
| 7 | 20-29 | F | | | | NOV-2021 | UND | 3 | 24 | 712 | N/A | N/A | No | Yes |
| 8 | 30-39 | M | | | | DEC-2021 | 18.2 | 1 | 23 | 847 | BA.1 | EPI_ISL_8604919 | No | Yes |
| 9 | 40-49 | F | | | | DEC-2021 | 32.3 | 6 | 28 | 1032 | BA.1 | EPI_ISL_8604901 | No | Yes |
| 10 | 20-29 | M | | | | DEC-2021 | 30.4 | 2 | 13 | 1197 | BA.1 | EPI_ISL_8604908 | No | Yes |
| 11 | 20-29 | F | | | | DEC-2021 | 28.3 | 8 | 22 | 863 | BA.1 | EPI_ISL_8604913 | No | No |
| 12 | 20-29 | F | | | | DEC-2021* | UND | 7 | 22 | 1259 | BA.1 | EPI_ISL_8604912 | No | Yes |
| 13 | 30-39 | M | BNT162b2 | JUL-2021 | 129 | NOV-2021 | 31.6 | 6 | 28 | 1069 | BA.1 | EPI_ISL_8604916 | No | Yes |
| 14 | 20-29 | M | | | | NOV-2021 | 30.8 | 8 | 15 | 1225 | N/A | N/A | No | Yes |
| 15 | 60-69 | F | BNT162b2 | JUL-2021 | 139 | DEC-2021 | 24.6 | 4 | 25 | 345 | BA.1 | EPI_ISL_8604920 | No | Yes |
| 16 | 60-69 | M | BNT162b2 | DEC-2021 | 15 | DEC-2021 | 24.6 | 2 | 10 | 904 | BA.1 | EPI_ISL_8604911 | No | No |
| 17 | 30-39 | M | | | | DEC-2021 | 37.0 | 5 | 19 | 1008 | BA.1 | EPI_ISL_8604923 | No | No |
| 18 | 60-69 | F | | | | DEC-2021 | 26.8 | 8 | 23 | 1111 | BA.1 | EPI_ISL_8578312 | Yes | Yes |
| 19 | 30-39 | M | | | | DEC-2021* | 30.7 | 13 | 27 | 1077 | BA.1 | EPI_ISL_8604924 | No | Yes |
| 20 | 20-29 | F | | | | DEC-2021 | 35.9 | 8 | 24 | 533 | BA.1 | EPI_ISL_8604911 | No | Yes |
| 21 | 20-29 | M | | | | DEC-2021 | 29.1 | 7 | 21 | 225 | BA.1 | EPI_ISL_8604922 | No | No |
| 22 | 30-39 | F | AD26.COV2 | AUG-2021 | 120 | DEC-2021 | 33.4 | 9 | 23 | 777 | BA.1 | EPI_ISL_8693907 | No | Yes |
| 23 | 20-29 | F | | | | DEC-2021 | 35.8 | 3 | 10 | 1167 | BA.1 | EPI_ISL_8604902 | No | No |
| 24 | 50-59 | M | BNT162b2 | AUG-2021 | 128 | DEC-2021 | 36.6 | 4 | 18 | 605 | N/A | N/A | No | Yes |
| 25 | 30-39 | F | AD26.COV2 | APR-2021 | 237 | DEC-2021 | 23.5 | 3 | 24 | 640 | BA.1 | EPI_ISL_8578311 | No | No |
| 26 | 50-59 | F | AD26.COV2 | JUL-2021 | 150 | DEC-2021 | UND | 5 | 23 | 716 | N/A | N/A | No | No |
| 27 | 50-59 | F | | | | DEC-2021 | 32.4 | 12 | 28 | 625 | N/A | N/A | Yes | Yes |
| 28 | 80-89 | F | BNT162b2 | JUL-2021 | 177 | JAN-2022 | 30.8 | 7 | 22 | 407 | N/A | N/A | Yes | Yes |
| 29 | 60-69 | M | BNT162b2 | JUL-2021 | 178 | DEC-2021 [§] | UND | 25 | 32 | 351 | N/A | N/A | Yes | Yes |
| 30 | 40-49 | M | | | | DEC-2021 | 20.2 | 0 | 13 | 844 | BA.1 | EPI_ISL_8604909 | No | No |
| 31 | 30-39 | F | | | | DEC-2021 | 34.8 | 1 | 22 | 647 | N/A | N/A | Yes | Yes |
| 32 | 50-59 | F | | | | DEC-2021 | 28.2 | 15 | 36 | 620 | BA.1 | EPI_ISL_8578347 | No | Yes |
| 33 | 20-29 | F | | | | DEC-2021 | UND | 3 | 18 | 902 | N/A | N/A | No | Yes |
| 34 | 30-39 | F | | | | DEC-2021 | 34.8 | 9 | 30 | 1363 | N/A | N/A | No | Yes |
| 35 | 50-59 | F | | | | DEC-2021 | 26.6 | 4 | 27 | 766 | BA.1 | EPI_ISL_8578342 | Yes | Yes |
| 36 | 20-29 | F | | | | DEC-2021 | UND | 9 | 23 | 1212 | N/A | N/A | No | Yes |
| 37 | 50-59 | F | | | | DEC-2021 | UND | 12 | 30 | 995 | N/A | N/A | No | Yes |
| 38 | 30-39 | M | | | | DEC-2021 | UND | 9 | 31 | 746 | N/A | N/A | No | Yes |
| 39 | 50-59 | F | | | | DEC-2021 | UND | 13 | 30 | 840 | N/A | N/A | Yes | Yes |
| 40 | 30-39 | F | | | | DEC-2021 | 22.5 | 5 | 19 | 61*** | BA.1 | EPI_ISL_8578314 | Yes | Yes |
| 41 | 40-49 | F | | | | NOV-2021 | 29.8 | 17 | 24 | 53*** | N/A | N/A | No | Yes |

Ct enrol.: qPCR cycle threshold for SARS-CoV-2 at enrollment. Symptoms to enrol.: time between symptoms onset and study enrolment. Symp. to last follow-up: time between symptoms onset and last follow-up visit. Max CD4: maximum CD4 count per microliter blood across all study visits. Supp O₂: participant required supplemental oxygen during the study. Hosp.: participant hospitalized during the study. UND: Undetectable Ct. N/A: Not available; sequencing failed, usually due to insufficient virus substrate. *Reported previous infection. **Boosted with Ad26.Cov2.S in Nov-2021. ***Participants with persistent low CD4 count and uncontrolled HIV viremia indicative of advanced HIV disease and immune suppression. Excluded from analysis. §Deceased.

Extended Data Table 2 | Summary characteristics of Omicron/BA.1 infected participants

| | All 39 | Vaccinated 15 (38%) | Unvaccinated 24 (62%) |
|--|------------------|-------------------------------|---------------------------------|
| Age | 35 (27-55) | 37 (32-60) | 31.5 (26-49) |
| Female | 25 (64%) | 9 (60%) | 16 (67%) |
| Vaccination to enrollment (days) | - | 139 (120-178) | - |
| Symptom onset to enrolment (days) | 6 (3-9) | 4 (3-6) | 7.5 (3-9) |
| Symptom onset to last follow-up (days) | 23 (19-27) | 23 (18-27) | 23 (20-28) |
| Maximum CD4 count (cell/ μ L) | 844 (647-1077) | 777 (605-1069) | 882.5 (729-1139) |
| Required supp. O ₂ | 7 (18%) | 2 (13%) | 5 (21%) |
| Hospitalized | 27 (69%) | 8 (53%) | 19 (79%) |
| Duration of hospitalization (days) | 7 (3-11) | 3.5 (2.5-14.5) | 8 (3-11) |

Values are median (IQR). Hospital stay calculated to last inpatient study visit.

Article

Extended Data Table 3 | Characteristics of Delta infected participants

| # | Age | Sex | Vacc. type | Date of vacc. | Vacc. to enrol. (days) | Date symp. onset | Ct at enrol. | Symp. to collection (days) | Seq. GISAID ID |
|---------|-------|-----|------------|---------------|---------------------------|------------------|--------------|-------------------------------|-----------------|
| 1 | 40-49 | F | | | | JUL-2021 | 26 | 26 | EPI_ISL_3722338 |
| 2 | 40-49 | M | | | | JUL-2021 | 31 | 23 | EPI_ISL_3722335 |
| 3 | 50-59 | M | | | | JUL-2021 | 30 | 31 | N/A |
| 4 | 50-59 | M | | | | JUN-2021 | 27 | 37 | N/A |
| 5 | 40-49 | M | | | | JUL-2021 | 35 | 44 | N/A |
| 6 | 30-39 | M | | | | JUL-2021 | 37 | 32 | N/A |
| 7 | 70-79 | M | BNT162b2 | JUN-2021 | 37 | JUL-2021 | 37 | 15 | N/A |
| 8 | 60-69 | F | BNT162b2 | NOV-2021 | 14 | AUG-2021 | UND | 116 | N/A |
| 9 | 40-49 | F | AD26.COV | MAY-2021 | 117 | JUL-2021 | UND | 31 | N/A |
| 10 | 50-59 | F | AD26.COV | APR-2021 | 147 | JUL-2021 | UND | 57 | N/A |
| 11 Pre | 40-49 | M | | | | AUG-2021* | 35 | 13* | N/A |
| 11 Post | 40-49 | M | BNT162b2 | OCT-2021 | 18 | AUG-2021 | UND | 83 | N/A |
| 12 Pre | 40-49 | M | | | | JUL-2021 | 23 | 24 | EPI_ISL_3939068 |
| 12 Post | 40-49 | M | AD26.COV | SEP-2021 | 32 | JUL-2021 | UND | 92 | N/A |
| 13 Pre | 30-39 | M | | | | JUL-2021 | 27 | 24 | EPI_ISL_3939088 |
| 13 Post | 30-39 | M | AD26.COV | SEP-2021 | 32 | JUL-2021 | UND | 94 | N/A |
| 14 Pre | 50-59 | F | | | | JUL-2021* | 27 | 23* | EPI_ISL_3447779 |
| 14 Post | 50-59 | F | BNT162b2 | OCT-2021 | 22 | JUL-2021 | UND | 93 | N/A |

*Asymptomatic, date of diagnostic swab used instead of symptoms onset. Ct enrol: qPCR cycle threshold for SARS-CoV-2 at enrollment. UND: undetectable. Pre: sample taken pre-vaccination. Post: sample taken post-vaccination for participants with a pre-vaccination sample. N/A: not available.

Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

- | | |
|-----|-----------|
| n/a | Confirmed |
|-----|-----------|
- The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
 - A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
 - The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
 - A description of all covariates tested
 - A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
 - A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
 - For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
 - For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
 - For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
 - Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection

Data analysis

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our [policy](#)

Sequences of outgrown Omicron sub-lineages have been deposited in GISAID with accession EPI_ISL_7886688 (Omicron/BA.1) and EPI_ISL9082893 (Omicron/BA.2). Delta, Beta, and ancestral SARS-CoV-2 isolates have been previously described (15). Raw images of the data are available upon reasonable request.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

Life sciences Behavioural & social sciences Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/documents/nr-reporting-summary-flat.pdf](https://www.nature.com/documents/nr-reporting-summary-flat.pdf)

Life sciences study design

All studies must disclose on these points even when the disclosure is negative.

| | |
|-----------------|--|
| Sample size | Samples size was not predetermined. We offered study enrollment to all participants who met the inclusion/exclusion criteria |
| Data exclusions | We excluded two samples from participants who were immune suppressed due to advanced HIV disease based on a low CD4 count and uncontrolled HIV viremia. |
| Replication | Repeated in independent experiments on different days in sets of paired experiments to which always included Omicron/BA.1. Two experiments performed for all variants and 6 experiments performed for BA.1. Geometric mean of measurements per participant was used. |
| Randomization | No randomization. |
| Blinding | No blinding |

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

| n/a | Involved in the study |
|-------------------------------------|---|
| <input type="checkbox"/> | <input checked="" type="checkbox"/> Antibodies |
| <input type="checkbox"/> | <input checked="" type="checkbox"/> Eukaryotic cell lines |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> Palaeontology and archaeology |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> Animals and other organisms |
| <input type="checkbox"/> | <input checked="" type="checkbox"/> Human research participants |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> Clinical data |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> Dual use research of concern |

Methods

| n/a | Involved in the study |
|-------------------------------------|---|
| <input checked="" type="checkbox"/> | <input type="checkbox"/> ChIP-seq |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> Flow cytometry |
| <input checked="" type="checkbox"/> | <input type="checkbox"/> MRI-based neuroimaging |

Antibodies

| | |
|-----------------|---|
| Antibodies used | Foci were stained with a rabbit anti-spike monoclonal antibody (BS-R2B12, GenScript A02058) at 0.5 µg/mL. Secondary goat anti-rabbit horseradish peroxidase (Abcam ab205718) antibody was added at 1 µg/mL |
| Validation | Information sheet for A02058 at https://www.genscript.com/antibody/A02058-MonoRab_SARS_CoV_2_Spike_S1_Antibody_BS_R2B12_mAb_Rabbit.html . Information sheet for ab205718: https://www.abcam.com/goat-rabbit-igg-hl-hrp-ab205718.html |

Eukaryotic cell lines

Policy information about [cell lines](#)

| | |
|---|---|
| Cell line source(s) | Vero E6 cells (ATCC CRL-1586) obtained from Cellonex in South Africa. The H1299-E3 (H1299-ACE2, clone E3) cell line was derived from H1299 as described in our previous work. H1299 was a gift from M. Oren, originally obtained from ATCC (CRL-5803) |
| Authentication | Cell lines have not been authenticated. |
| Mycoplasma contamination | The cell lines have been tested for mycoplasma contamination and are mycoplasma negative. |
| Commonly misidentified lines (See ICLAC register) | None |

Human research participants

Policy information about [studies involving human research participants](#)

| | |
|----------------------------|---|
| Population characteristics | Participant characteristics are listed per participant in Table S1 and S3 and summarized in Table S2. |
| Recruitment | Blood samples were obtained from hospitalized adults with PCR-confirmed SARS-CoV-2 infection and/or vaccinated individuals who were enrolled in a prospective cohort study approved by the Biomedical Research Ethics Committee at the University of KwaZulu–Natal. Investigators were blinded to participant information. |
| Ethics oversight | Study approved by the Biomedical Research Ethics Committee at the University of KwaZulu–Natal (reference BREC/00001275/2020). Use of residual Omicron/BA.1 swab sample was approved by the University of the Witwatersrand Human Research Ethics Committee (HREC) (ref. M210752). Use of swab sample to isolate Omicron/BA.2 was collected as part of the “COVID-19 transmission and natural history in KwaZulu-Natal, South Africa: Epidemiological Investigation to Guide Prevention and Clinical Care” Centre for the AIDS Programme of Research in South Africa (CAPRISA) study and approved by the Biomedical Research Ethics Committee at the University of KwaZulu–Natal (reference 201 BREC/00001195/2020, BREC/00003106/2021). |

Note that full information on the approval of the study protocol must also be provided in the manuscript.