



Contents lists available at ScienceDirect

Vaccine

journal homepage: www.elsevier.com/locate/vaccine

A case-crossover study of the effect of vaccination on SARS-CoV-2 transmission relevant behaviours during a period of national lockdown in England and Wales

Aimee Serisier^a, Sarah Beale^{a,b,*}, Yamina Boukari^b, Susan Hoskins^a, Vincent Nguyen^{a,b}, Thomas Byrne^b, Wing Lam Erica Fong^b, Ellen Fragaszy^{b,c}, Cyril Geismar^{a,b}, Jana Kovar^a, Alexei Yavlinsky^b, Andrew Hayward^a, Robert W. Aldridge^a

^a Institute of Epidemiology and Health Care, University College London, London WC1E 7HB, UK

^b Centre for Public Health Data Science, Institute of Health Informatics, University College London, NW1 2DA, UK

^c Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK

ARTICLE INFO

Article history:

Received 29 August 2022

Received in revised form 28 November 2022

Accepted 29 November 2022

Available online xxxx

Keywords:

COVID-19
Vaccine
Behaviour
Mitigations

ABSTRACT

Background: Studies of COVID-19 vaccine effectiveness show increases in COVID-19 cases within 14 days of a first dose, potentially reflecting post-vaccination behaviour changes associated with SARS-CoV-2 transmission before vaccine protection. However, direct evidence for a relationship between vaccination and behaviour is lacking. We aimed to examine the association between vaccination status and self-reported non-household contacts and non-essential activities during a national lockdown in England and Wales.

Methods: Participants (n = 1154) who had received the first dose of a COVID-19 vaccine reported non-household contacts and non-essential activities from February to March 2021 in monthly surveys during a national lockdown in England and Wales. We used a case-crossover study design and conditional logistic regression to examine the association between vaccination status (pre-vaccination vs 14 days post-vaccination) and self-reported contacts and activities within individuals. Stratified subgroup analyses examined potential effect heterogeneity by sociodemographic characteristics such as sex, household income or age group.

Results: 457/1154 (39.60 %) participants reported non-household contacts post-vaccination compared with 371/1154 (32.15 %) participants pre-vaccination. 100/1154 (8.67 %) participants reported use of non-essential shops or services post-vaccination compared with 74/1154 (6.41 %) participants pre-vaccination. Post-vaccination status was associated with increased odds of reporting non-household contacts (OR 1.65, 95 % CI 1.31–2.06, p < 0.001) and use of non-essential shops or services (OR 1.50, 95 % CI 1.03–2.17, p = 0.032). This effect varied between men and women and different age groups.

Conclusion: Participants had higher odds of reporting non-household contacts and use of non-essential shops or services within 14 days of their first COVID-19 vaccine compared to pre-vaccination. Public health emphasis on maintaining protective behaviours during this post-vaccination time period when individuals have yet to develop full protection from vaccination could reduce risk of SARS-CoV-2 infection.

© 2022 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The coronavirus disease 2019 (COVID-19) pandemic has had a devastating impact on global public health since the causative agent, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was first identified in late 2019 [1,2]. SARS-CoV-2 is transmitted through direct or indirect contact with infected respiratory droplets or aerosols [3–5] and consequently public settings and

Abbreviations: COVID-19, Coronavirus disease 2019; SARS-CoV-2, Severe acute respiratory syndrome coronavirus 2; UK, United Kingdom; IMD, Index of Multiple Deprivation; VoC, Variant of Concern.

* Corresponding author at: Institute of Epidemiology and Health Care, University College London, London WC1E 7HB, UK.

E-mail address: sarah.beale.19@ucl.ac.uk (S. Beale).

<https://doi.org/10.1016/j.vaccine.2022.11.073>

0264-410X/© 2022 The Authors. Published by Elsevier Ltd.

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

activities that involve direct and indirect contact may promote SARS-CoV-2 transmission [4,6]. These were the target of non-pharmaceutical interventions (NPIs) introduced by many governments worldwide to control the spread of the virus, including social distancing measures to minimise contacts with other people, face mask mandates and closure of non-essential public venues. In the UK, the most stringent levels of national restrictions included periods of 'lockdown', which broadly involved strict restrictions on non-household mixing, closure or restrictions on usage of non-essential public spaces, advice to stay at home and work from home where possible, and stringent isolation and quarantine protocols [7]. Lockdowns were introduced three times in England and Wales, between March and May 2020 (first national lockdown), October and November 2020 (depending on nation, second lockdown), and January to March 2021 (third national lockdown).

COVID-19 vaccination programmes are now a cornerstone of pandemic response in the UK and worldwide, allowing for relaxation of lockdown restrictions with varying degrees of remaining NPIs. In the UK, COVID-19 vaccinations began in December 2020 with the emergency licencing of first the Pfizer BioNTech and then AstraZeneca vaccines. Delivered as two doses with additional booster doses, first and subsequent doses are proven to be effective in reducing symptomatic and asymptomatic infections, hospitalisations, and deaths from COVID-19 [8–11]. However, the protection provided by vaccines is not immediate and infection with SARS-CoV-2 after vaccination is possible [12]. Studies using data from Israel and the UK suggest an increased risk of symptomatic infection within 14 days of the first dose of a COVID-19 vaccine [13,14]; the reduction in risk of symptomatic infection is seen after 14 days. It is possible that increased risk in the 14 days after vaccination may reflect changes in behaviour associated with SARS-CoV-2 transmission during the period of time in which immunologic protection is building [15–17], particularly given the extent to which human behaviour is known to influence infectious disease dynamics [18–21]. Furthermore, concerns have been raised that a reduction in behaviours protecting against SARS-CoV-2 infection could be seen if the perceived risk of infection is reduced after vaccination against COVID-19 [22].

Empirical evidence regarding the effect of vaccination on infection prevention behaviour is limited. One study examining protective behaviours before and after vaccination against Lyme disease found that people who were vaccinated reduced protective behaviours and believed that they were at less risk of infection than unvaccinated people [23]. There is evidence in the context of the COVID-19 pandemic that suggests changes in protective behaviours may occur. In February 2021 the Office for National Statistics (ONS) reported that 41 % of over 80 s met a person who was not a part of their household or support bubble indoors within 21 days post-vaccination [24]. A December 2020 YouGov survey poll found that of the 1706 people surveyed, 29 % said that they would follow public health restrictions less strictly after receiving a vaccine [25]; this poll was conducted prior to widespread availability of COVID-19 vaccination in the UK. Reductions in compliance with mask use and handwashing post-vaccination were found amongst healthcare workers in Ethiopia after the first dose of a COVID-19 vaccine [26]. Contradictory to these findings, a longitudinal analysis investigating compliance with protective behaviours found increases in self-reported compliance with public health guidance and social distancing in vaccinated and unvaccinated individuals from October 2020 to February 2021 [27]. However, variation in context over this time could have influenced these results - for example, the introduction of the third UK-wide national lockdown in early January 2021 [7] or changes in the perceived risk of infection as case numbers rose.

This study aimed to quantify the effect of a change in COVID-19 vaccination status on transmission-relevant behaviours during a

period of national lockdown using data from Virus Watch, a large prospective cohort study based in England and Wales. We set out to investigate whether participants' self-reported levels of non-household contacts and retail and social activities classed as non-essential under lockdown restrictions (such as use of a hairdressers or other services for personal care, attending a party, or dining at a restaurant or café) changed within 14 days of their first dose of a COVID-19 vaccination compared to pre-vaccination.

2. Methods

2.1. Study design

A case-crossover design was used to examine the association between vaccination status (pre-vaccination versus ≤ 14 days post first dose) and self-reported contacts and activities within individuals. Case-crossover designs are appropriate to examine the association between transient exposures and acute outcomes [28] and eliminate measured and unmeasured time-invariant confounding when comparing within-person exposed and unexposed periods [28–31].

To maximise the comparability of referent periods and reduce the potential for confounding by temporal or spatial trends in contacts and activities, the study timeframe was limited to include survey responses from within the third national lockdown in England and Wales (6 January–29 March 2021). Each survey was open for 7 days and concerned non-household contacts and activities in the week prior to its start date - surveys from 9 to 16 February 2021 and 9–16 March 2021 could therefore be included, and the pre-vaccination period comprised responses from 2 to 9 February 2021 and the post-vaccination period from 2 to 9 March 2021 (Fig. 1).

2.2. Study setting and population

Data analysed in this project were collected as part of Virus Watch, a household community cohort study of SARS-CoV-2 transmission and COVID-19 in England and Wales. Details of Virus Watch relevant to the present study are described briefly here, with further information on its full scope and methodology described in the study protocol [32].

Whole households who met the following inclusion criteria were recruited for voluntary participation through social media and General Practice supported postal and SMS recruitment campaigns: resident in England or Wales, household size 1–6 people (due to survey infrastructure limitations), internet and email access, and ability of at least one household member to complete online surveys in English [32]. All participants completed a baseline survey upon study registration that collected information on sociodemographic factors for each household member. Participants reported on changes in vaccination status in weekly web surveys, which were weekly completed by approximately 50 % of the recruited cohort [33]. Monthly web surveys collected health-related and behavioural/psychosocial factors, including information about participants' activities and contacts, with responses to the monthly surveys varying, with responses to relevant monthly surveys reported in the results.

Inclusion criteria for the present study were having received the first dose of a COVID-19 vaccine and responded to a monthly survey regarding activities and contacts both pre-vaccination and post-vaccination ($n = 1154$). All surveys were administered online and survey completion was voluntary and not incentivised. Participants under 18 years old were excluded ($n = 4$) as this age group was not the focus of vaccinations during the study timeframe. Participants with missing data on sociodemographic covariates (see




Fig. 1. Participants reported contacts and activities in the previous week in monthly web surveys, which were open for completion for 7 days. Vaccination status was reported in weekly web surveys. The post-vaccination monthly web survey was therefore selected to cover a period of time within 7–14 days post-vaccination.

below) to be used in stratified analyses were excluded from the study sample following initial descriptive statistics ($n = 23$). The final study population for analysis included 1141 participants (Fig. 2).

2.3. Exposure

Self-reported vaccination status was the exposure of interest, defined as 'pre-vaccination' and 'within 14-days post-vaccination' of a first dose of a COVID-19 vaccine and derived from data on the date and dose of vaccinations reported in weekly surveys.

2.4. Outcome

The following contact and activity outcomes were derived based on responses to monthly surveys, and were binary coded (yes or no during survey week): [1] any close contact with non-household members ('face-to-face contact with another person <1 m away, spending more than 15 min within 2 m of another per-

son, or travelling in a car or other small vehicle with another person' [34], [2] any social or leisure activity (attending a theatre, cinema, concert, or sports event; attending a party; going to a restaurant, café, or canteen; going to a bar, pub, or club; and/or use of a gym or indoor sports facility), and [3] using non-essential shops or services (retail venues or services not required to meet basic needs such as food and medicine, e.g., hairdressers, barbers, or beauty salons). Such behaviours were targeted by social distancing restrictions and/or public venue closures under lockdown restrictions in place in February–March 2021. Due to low numbers of participants reporting social activities in both pre- and post-vaccination surveys during this lockdown period, social and leisure activities could not be included in this analysis.

2.5. Demographic characteristics

Data on sociodemographic characteristics were collected from study baseline responses. Age group was defined according to reported age of participants at baseline: <60 years and 60 years




Fig. 2. Exclusion criteria for the analysis of the effect of vaccination status on contacts and activities.

or more. Participants were categorised as White British and Minority Ethnic according to self-reported ethnicity. Sex was categorised as male and female by self-reported sex. Household size was defined as households of 1 person, 2–3 people and 4–6 people. The Index of Multiple Deprivation (IMD) for each participant was categorised using household postcodes recorded on registration, comprising 1 (least deprived) through 5 (most deprived). Region was measured using participant's postcodes and categorised according to ONS national regions. Observations with missing data on sex (1.12 %) and age (0.43 %) which were included as potential effect modifiers were excluded following descriptive statistics in order to perform a complete cases analysis.

2.6. Analysis

The frequency with which non-household contacts and use of non-essential shops or services were reported in surveys pre- and post-vaccination were calculated and stratified by age and sex only as group sizes amongst other participant subgroups were too small. Conditional logistic regression was used to assess the odds of reporting contacts and activity outcomes within 14 days after the first dose of a COVID-19 vaccine compared to the earlier pre-vaccination time point within individuals. This approach accounts for non-independence in responses within individuals over time by conditioning the participants responses using an individual-specific fixed effect [35]. Each participant contributes two observations occupying a single stratum in the model. As the model estimates within-individual changes, strata that do not vary (i.e., people who reported contacts and activities in neither time period, or in both) do not contribute to the model [30]. Stratification of conditional logistic regression models by age and sex was then performed. Sociodemographic covariates were collected at baseline and considered time-invariant confounders, and consequently were not entered into the models. Cluster robust standard errors were used to account for household-level clustering. The alpha level was set at $p = 0.05$.

2.7. Ethics

Ethical approval for Virus Watch was obtained from the Hampstead National Health Service (NHS) Health Research Authority Ethics Committee (ethics approval number 20/HRA/2320). All participants provided online informed consent. All analyses were conducted using the University College London Data Safe Haven.

3. Results

Table 1 presents the sociodemographic characteristics of Virus Watch participants in this analysis ($n = 1154$ from 1031 households). The median age in years was 60 (interquartile range: 56,62). The majority of participants were over 60 years of age (58.12 %). Participants were mostly white (95.93 %), from the East of England (24.61 %) or South East (20.10 %), lived in households with 2–3 people (62.48 %), in IMD 5 (least deprived), (31.63 %) or with an annual household income of £50,000 or more (38.88 %). There was a greater proportion of men in the sample (54.59 %) compared to women (44.37 %).

Table 2 shows the frequency of reporting non-household contacts and use of non-essential shops or services. Pre-vaccination surveys asked participants to record activities from 2 to 9 February 2021, with post-vaccination surveys recording activities from 2 to 9 March 2021. 371/1,154 (32.15 %) participants reported contact with a person outside of their household or support bubble pre-vaccination compared with 457/1,154 (39.60 %) participants in the 14 days after the first dose of a COVID-19 vaccine. 74/1154

Table 1
Characteristics of study participants.

Characteristic	<i>n</i>	%
Age		
<60	480	41.45
≥60	673	58.12
Missing	<5	0.43
Sex		
Male	630	54.59
Female	512	44.37
Missing	12	1.04
Ethnicity		
White	1,069	92.63
Minority ethnic	72	6.24
Missing	13	1.13
Region		
East Midlands	104	9.01
East of England	284	24.61
London	101	8.75
North East	50	4.33
North West	132	11.44
South East	232	20.10
South West	90	7.80
Wales	19	1.65
West Midlands	65	5.63
Yorkshire and the Humber	64	5.55
Missing	13	1.13
Household size		
1 person	223	19.32
2–3 people	721	62.48
4–6 people	210	18.20
Missing	0	0
IMD		
1 (most deprived)	78	6.76
2	173	14.99
3	254	22.01
4	271	23.48
5 (least deprived)	365	31.63
Missing	13	1.13
Household income		
£0–£24,999	249	25.75
£25,000–£49,999	342	35.37
≥£50,000	376	38.88
Missing	187	16.20
Total	1,154	

(6.41 %) participants reported using non-essential shops or services pre-vaccination, while 100/1,154 (8.67 %) participants did so in the 14 days after their first COVID-19 vaccine.

Table 3 shows the frequency with which participants reported activities in pre-vaccination and post-vaccination surveys stratified by age group and sex. A greater proportion of participants reported non-household contacts in both pre-vaccination and post-vaccination surveys compared to use of non-essential shops and services, but both activities were reported more frequently post-vaccination. Post-vaccination, reporting of non-household contacts was greatest amongst male participants and those aged <60 (257/630, 40.79 % and 199/480, 41.46 % respectively), but the greatest increase in reported non-household contacts was seen when comparing post-vaccination with pre-vaccination surveys from male participants and those aged 60 years or more. Greater proportions of female participants (47/512, 9.18 %) and participants aged 60 years or more (70/673, 10.40 %) reported use of non-essential shops or services within 14 days post-vaccination. However, reported use of non-essential shops or services by male participants increased by more than female participants post-vaccination.

Table 4 presents the results of conditional logistic models for the within-individual effects of vaccination on non-household close contacts and use of non-essential shops or services. Odds of reporting non-household close contacts (OR 1.65, 95 % CI 1.31–

Table 2

Frequency of reporting non-household contacts and use of non-essential shops or services over 7 days in pre- and post-vaccination surveys (n = 1154).

Activity	Pre-vaccination		≤14 days post-vaccination	
	No n (%)	Yes n (%)	No n (%)	Yes n (%)
Contacts	783 (67.85)	371 (32.15)	697 (60.40)	457 (39.60)
Use of non-essential shops or services	1,080 (93.59)	74 (6.41)	1,054 (91.33)	100 (8.67)

The pre-vaccination survey and post-vaccination surveys asked participants to report activities from 2 to 9 February 2021 and 2–9 March 2021 respectively.

Table 3

The frequency of reporting non-household contacts and use of non-essential shops or services over 7 days in pre-vaccination and post-vaccination surveys by participant characteristics (n = 1154).

Characteristic	n	Contacts				Use of non-essential shops or services			
		Pre-vaccination		≤14 days post-vaccination		Pre-vaccination		≤14 days post-vaccination	
		No n (%)	Yes n (%)	No n (%)	Yes n (%)	No n (%)	Yes n (%)	No n (%)	Yes n (%)
Age									
<60	480	310 (64.58)	170 (35.42)	281 (58.54)	199 (41.46)	457 (95.21)	23 (4.79)	450 (93.75)	30 (6.25)
≥60	673	472 (70.13)	201 (29.87)	415 (61.66)	258 (38.34)	622 (92.42)	51 (7.58)	603 (89.60)	70 (10.40)
Missing	<5	<5	0	<5	0	<5	0	<5	0
Sex									
Male	630	425 (67.46)	205 (32.54)	373 (59.21)	257 (40.79)	597 (94.76)	33 (5.24)	579 (91.90)	51 (8.10)
Female	512	351 (68.55)	161 (31.45)	317 (61.91)	195 (38.09)	472 (92.19)	40 (7.81)	465 (90.82)	47 (9.18)
Missing	12	7 (58.33)	5 (41.67)	7 (58.33)	5 (41.67)	11 (91.67)	<5	10 (83.33)	<5

The pre-vaccination survey and post-vaccination surveys asked participants to report activities from 2 to 9 February 2021 and 2–9 March 2021 respectively.

Table 4

Conditional logistic regression models for the effect of vaccinations status on contacts and use of non-essential shops or services.

Vaccination status	n	Paired OR	Std. Err.	95 % CI	P-value
Contacts					
≤14 days post-vaccination	352	1.65	0.19	1.31–2.06	<0.001
Pre-vaccination			-	-	-
Use of non-essential shops or services					
≤14 days post-vaccination	125	1.50	0.28	1.03–2.17	0.032
Pre-vaccination			-	-	-

Models adjusted for household structure with cluster-robust standard errors.

2.06, $p < 0.001$) and use of non-essential shops or services (OR 1.50, 95 % CI 1.03–2.17, $p = 0.032$) were higher within 14 days post-vaccination compared to pre-vaccination.

Table 5 shows the results of stratified conditional logistic models examining heterogeneity of the within-individual effect of a change in vaccination status for contacts and use of non-essential shops or services by sociodemographic characteristics. All analyses compare time periods within 14 days post-vaccination to time periods pre-vaccination.

After stratification by sex, greater odds of reporting non-household close contacts within the 14 days post-vaccination compared to pre-vaccination were consistent for both men (OR 1.71, 95 % CI 1.28–2.29, $p < 0.001$) and women (OR 1.57, 95 % CI 1.13–2.17, $p = 0.007$). However, overlapping confidence intervals indicate a lack of effect heterogeneity by sex in the study population. There was evidence of effect of vaccination status on use of non-essential shops or services amongst male participants, who had greater odds of reporting these activities post-vaccination (OR 1.86 95 % CI 1.14–3.32, $p = 0.015$). Female participants were not at increased odds of reporting use of non-essential shops or services (OR 1.24, 95 % CI 0.76–2.03, $p < 0.39$).

The odds of reporting non-household contacts and use of non-essential shops or services were respectively 1.48 (95 % CI 1.05–2.08, $p = 0.027$) and 1.37 (95 % CI 0.75–2.47, $p = 0.30$) amongst par-

ticipants aged <60 years. Elevated odds were seen in those aged 60 years or more, with greater odds for both non-household close contacts (OR 1.80, 95 % CI 1.33–2.41, $p < 0.001$) and use of non-essential shops or services (OR 1.58, 95 % CI 1.01–2.48, $p = 0.046$).

4. Discussion

Our findings indicate that within 14 days of their first dose of a COVID-19 vaccine, participants were more likely to report non-household close contacts (OR = 1.65, 1.31–2.06) and using non-essential shops and services (OR = 1.50, 1.03–2.17) compared to pre-vaccination. There was no substantial heterogeneity in the effect of vaccination status on non-household close contacts between men and women, but men had higher odds of reporting use of non-essential shops or services. Participants aged 60 years or more had greater odds of reporting both non-household close contacts and use of non-essential shops or services post-vaccination. Increased close contact with people outside of the household and in non-essential retail activities may contribute to the increased risk of infection seen within 14 days of a first dose of a COVID-19 vaccine [13,14], although directly measuring the risk of infection associated with these behaviours was outside the scope of this analysis.

Table 5

Conditional logistic analyses stratified by sociodemographic characteristics for the effect of vaccination status on self-reported contacts and use of non-essential shops or services within 14 days of the first dose of a COVID-19 vaccine compared to pre-vaccination.

Characteristic	Contacts					Use of non-essential shops or services				
	n	Paired OR	Std. Err.	95 % CI	P-value	n	Paired OR	Std. Err.	95 % CI	P-value
Sex										
Male	198	1.71	0.25	1.28–2.29	<0.001	60	1.86	0.51	1.08–3.19	0.025
Female	154	1.57	0.26	1.13–2.17	0.007	65	1.24	0.31	0.76–2.03	0.39
Age group										
<60	151	1.48	0.26	1.05–2.08	0.027	45	1.37	0.41	0.75–2.47	0.30
≥60	201	1.80	0.27	1.33–2.41	<0.001	80	1.58	0.36	1.01–2.48	0.046

Models adjusted for household structure with cluster-robust standard errors.

The greatest effect was seen for contacts with people outside of the household. Few participants reported use of non-essential shops or services or social activities, leading to the exclusion of social activities from this analysis due to small group sizes. This is largely due to closure of public venues (e.g., non-essential retail or indoor dining venues) under national regulations in England and Wales at the time of the survey. Results from stratified analyses must also be interpreted with caution due to small subgroup sizes (Table 5), which likely substantially reduced power for these analyses.

These results highlight the need to maintain protective behaviours within 14 days of the first dose of a COVID-19 vaccine while immunological protection builds. This could reduce the risk of infection in this time and is a relevant target for public health messaging in future outbreaks of infectious diseases for which vaccines are available. Although this analysis focused specifically on non-essential contacts and activities during the third national lockdown, it is possible that the effect of vaccination on activity levels at times when social distancing and other public health measures are relaxed may be even greater. Additionally, there is some evidence to suggest that older people are more likely to adopt or adhere to protective behaviours than younger people during pandemics [36]. As most participants in this project were older than 50 years of age, our findings may be different or magnified in younger age groups. Our findings may also be of relevance for current and future booster doses of COVID-19 vaccines considering reported reductions in vaccine effectiveness over time [37–40]. This may be of particular importance to booster programmes given recent evidence for immune evasion of both natural and vaccine-acquired immunity by the Omicron Variant of Concern (VoC) [41], currently the most dominant strain in the UK [42]. Longer-term, quantitative data on behaviour related to vaccination could be used to parametrize and validate models of infectious disease dynamics used in public health decision making and policy [20,21], assisting public health planning in response to future outbreaks of infectious diseases.

Our findings are supported by an analysis of social contacts between school children in the United States during the COVID-19 pandemic, which reported increased non-household contacts amongst children and adults in a household where at least one adult was vaccinated between February and April 2021 [43]. While we found no differences in the effect of vaccination status between men and women, previous research has suggested that women may be more likely to adopt protective behaviours than men during pandemics, which is potentially related to increased perceived risk [36]. A study examining adherence to public health guidance in the UK during the COVID-19 pandemic found that women reported making fewer trips outside of the home than men and less use of non-essential shops [44]. Although our findings are supported by evidence that a reduction in protective behaviours may be seen after the introduction of vaccines during the COVID-19 pandemic [26,45,46], they are not consistent with those from a

Virus Watch study using GPS location data to examine changes in travel distance pre- and post- the first dose of a COVID-19 vaccine during the third national lockdown [47]. In this analysis, Nguyen et al. found no evidence for an increase in the rate of change in the distance travelled by participants post-vaccination in the 30 days before and after vaccination, suggesting that participants may not have altered behaviours leading to an increased distance travelled during this time. However, participants in the geolocation tracking arm of the study were aware that they were being monitored and so may have been more likely to modify their behaviour. Our results differ from those of Wright et al. [27], whose research showed that compliance with public health guidelines increased in both unvaccinated and vaccinated individuals from October 2020 to February 2021. Context-specific psychological predictors of behaviour could account for this difference. A scoping review of 149 studies of behaviour change during pandemics (including the COVID-19 pandemic) found higher perceived risk predicted greater levels of adherence to a number of protective behaviours, including social distancing and avoidance of non-essential shopping [48]. A number of contextual factors could have caused changes in perceived risk between late 2020 and early 2021. Perceived risk may have been greater during the national lockdown beginning 6 January 2021 at the peak of the second wave of COVID-19 cases [49], which was larger and associated with more deaths from COVID-19 than the first wave [50,51]. It also occurred at a time when the SARS-CoV-2 Alpha VoC was rapidly expanding [52]. Comparison of longer-term changes in compliance with guidelines or protective behaviours post-vaccination with changes over a shorter period of time may therefore not be appropriate.

Alternatively, perceived risk may contribute to the changes in behaviour seen in this analysis independently of vaccination status. Following the peak in January 2021, COVID-19 cases decreased from February – March 2021 [48], and so over time participants may have felt at less risk of COVID-19 and gradually increased their contacts and activities. As our analysis did not directly examine the perceived risk of infection further investigation of the relationship between psychological predictors of behaviour, vaccination and infectious disease dynamics is warranted.

A strength of the case-crossover design is that self-matching controls for time-invariant confounders, which may be unknown or difficult to quantify. Furthermore, by selecting pre- and post-vaccination time periods to be during a national lockdown, temporal and spatial variation in behaviours was likely minimised. The potential for information bias is minimised as survey questions remained the same over time, and self-matching means that the same individuals reported information for both pre- and post-vaccination time periods.

An important limitation of this analysis is the lack of representativeness of the study population with the general population in England and Wales. Participants who responded to surveys were predominantly male, of White ethnicity, living in the East or South

East of England and living in less deprived environments with high household income. This is significant given the unequal burden of disease and mortality from COVID-19 in the UK population. People living in overcrowded households are at greater risk of SARS-CoV-2 infection [53], and rates of diagnosis and death are higher in people who live in more deprived areas and are from minority ethnic backgrounds [54]. Most participants were over 60 years of age, likely due to the phases of COVID-19 vaccine delivery, with older age groups being prioritised in the early stages of the COVID-19 vaccination programme [55]. The age structure of the cohort may therefore have been subject to eligibility bias. It is possible that a change in vaccination status may have had a greater effect on participants aged 60 years or more who were less worried about COVID-19 than those under 60 years of age who were eligible for vaccination during the same time period. Confounders such as household income, IMD and region were only available at baseline, however these were unlikely to vary over the study period. Self-reported contacts and activities may have been affected by recall bias, although this may have been minimised by survey timing (i.e., the following week). Social desirability bias may have been reduced by recording survey responses online. Importantly, the infection risk of contacts and activities in this project could not be directly measured using survey responses – this limits inferences about the impact of changes in behaviour on infection risk and is recommended as a focus for future research.

This analysis provides quantitative evidence for an association between vaccination status and transmission-relevant behaviour using data gathered during the third national lockdown in England and Wales. Our findings suggest that changes in protective behaviours occur while immunological protection is building and may contribute to risk of infection in the 14 days after vaccination, and therefore interventions emphasising the need to maintain protective behaviours in the recently vaccinated could reduce the risk of infection during time periods while immunity is building. It is possible that such an effect could exist during outbreaks of other vaccine preventable infectious diseases, and therefore quantitative data on behaviour relevant to vaccination could be of use in policymaking for not only current and future booster doses of COVID-19 vaccines but also vaccines for emerging and future pathogens.

Funding

The Virus Watch study is supported by the MRC Grant Ref: MC_PC 19,070 awarded to UCL on 30 March 2020 and MRC Grant Ref: MR/V028375/1 awarded on 17 August 2020. The study also received \$15,000 of Facebook advertising credit to support a pilot social media recruitment campaign on 18th August 2020. This study was also supported by the Wellcome Trust through a Wellcome Clinical Research Career Development Fellowship to RA [206602]. SB and TB are supported by an MRC doctoral studentship (MR/N013867/1). The funders had no role in study design, data collection, analysis and interpretation, in the writing of this report, or in the decision to submit the paper for publication.

Conflict of Interest Statement

AH serves on the UK New and Emerging Respiratory Virus Threats Advisory Group and is a member of the COVID-19 transmission sub-group of the Scientific Advisory Group for Emergencies (SAGE). [17:09] Boukari, Yamina. YB's spouse is employed by Elsevier as a Software Engineer. The other authors report no conflicts of interest.

Data availability

We aim to share aggregate data from this project on our website and via a "Findings so far" section on our website - [https://ucl-](https://ucl-virus-watch.net/)

[virus-watch.net/](https://ucl-virus-watch.net/). We also share some individual record level data on the Office of National Statistics Secure Research Service. In sharing the data we will work within the principles set out in the UKRI Guidance on best practice in the management of research data. Access to use of the data whilst research is being conducted will be managed by the Chief Investigators (ACH and RWA) in accordance with the principles set out in the UKRI guidance on best practice in the management of research data. We will put analysis code on publicly available repositories to enable their reuse.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] Wu F, Zhao S, Yu B, et al. A new coronavirus associated with human respiratory disease in China. *Nature* 2020;579:265–9. <https://doi.org/10.1038/s41586-020-2008-3>.
- [2] Zhou P, Yang XL, Wang XG, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020;579:270–3. <https://doi.org/10.1038/s41586-020-2012-7>.
- [3] Falahi S, Kenarkoobi A. Transmission routes for SARS-CoV-2 infection: review of evidence. *New Microbes New Infect* 2020;38:. <https://doi.org/10.1016/j.nmni.2020.100778>.
- [4] Government of the United Kingdom (EMG and NERVTAG). SARS-Cov-2 Transmission Routes and Environments SAGE - 22 October 2020; 2020. <<https://www.gov.uk/government/publications/sars-cov-2-transmission-routes-and-environments-22-october-2020>>.
- [5] World Health Organisation. Transmission of SARS-CoV-2: implications for infection prevention precautions; 2020. <<https://www.who.int/news-room/commentaries/detail/transmission-of-sars-cov-2-implications-for-infection-prevention-precautions>>.
- [6] Hayward AC, Beale S, Johnson AM, Fragaszy EB, Cowling BJ, Uzicanin A. Public activities preceding the onset of acute respiratory infection syndromes in adults in England-implications for the use of social distancing to control pandemic respiratory infections. [version 1; peer review: 2 approved]. *Wellcome Open Research* 2020;5:54. <<https://doi.org/10.12688/wellcomeopenres.15795.1>>.
- [7] Brown J., Kirk-Wade E, Barber S. Coronavirus: A history of 'Lockdown law' in England. House of Commons Library Research Briefing, 22 December 2021; 2021. <<https://researchbriefings.files.parliament.uk/documents/CBP-9068/CBP-9068.pdf>>.
- [8] Public Health England. Public Health England vaccine effectiveness report: March 2021; 2021. <https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/989360/PHE_COVID-19_vaccine_effectiveness_report_March_2021_v2.pdf>.
- [9] Public Health England. Impact of COVID-19 vaccines on mortality in England: December 2020 to March 2021; 2021. <https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/977249/PHE_COVID-19_vaccine_impact_on_mortality_March.pdf>.
- [10] Public Health England. PHE monitoring of the early impact and effectiveness of COVID-19 vaccination in England; 2021. <https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/968977/COVID-19_vaccine_effectiveness_surveillance_report_February_2021.pdf>.
- [11] Public Health England. Direct and indirect impact of the vaccination programme on COVID-19 infections and mortality; 2021. <https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/997495/Impact_of_COVID-19_vaccine_on_infection_and_mortality.pdf>.
- [12] Antonelli M, Penfold RS, Merino J, Sudre CH, Molteni E, Berry S, et al. Risk factors and disease profile of post-vaccination SARS-CoV-2 infection in UK users of the COVID Symptom Study app: a prospective, community-based, nested, case-control study. *Lancet Infect Dis* 2021;22(1):43–55. [https://doi.org/10.1016/S1473-3099\(21\)00460-6](https://doi.org/10.1016/S1473-3099(21)00460-6).
- [13] Bernal JL, Andrews N, Gower C, Stowe J, Robertson C, Cottrell S, et al. Early effectiveness of COVID-19 vaccination with BNT162b2 mRNA vaccine and ChAdOx1 adenovirus vector vaccine on symptomatic disease, hospitalisations and mortality in older adults in England. *MedRxiv* 2021. <https://doi.org/10.1101/2021.03.01.21252652>.
- [14] Hunter PR, Brainard J. Estimating the effectiveness of the Pfizer COVID-19 BNT162b2 vaccine after a single dose. A reanalysis of a study of 'real-world' vaccination outcomes from Israel. *MedRxiv* 2021. <https://doi.org/10.1101/2021.02.01.21250957>.
- [15] Shrotri MA, Fragaszy E, Geismar C, Nguyen V, Beale S, Braithwaite I, et al. Spike-antibody responses following first and second doses of ChAdOx1 and BNT162b2 vaccines by age, gender, and clinical factors - a prospective

- community cohort study (Virus Watch). MedRxiv 2021. <https://doi.org/10.1101/2021.05.12.21257102>.
- [16] Folegatti PM, Ewer KJ, Aley PK, Angus B, Becker S, Belij-Rammerstorfer S, et al. Safety and immunogenicity of the ChAdOx1 nCoV-19 vaccine against SARS-CoV-2: a preliminary report of a phase 1/2, single-blind, randomised controlled trial. *Lancet* 2020;396(10249):467–78. [https://doi.org/10.1016/S0140-6736\(20\)31604-4](https://doi.org/10.1016/S0140-6736(20)31604-4).
- [17] Sahin U, Muik A, Derhovanessian E, Vogler I, Kranz LM, Vormehr M, et al. COVID-19 vaccine BNT162b1 elicits human antibody and TH1 T cell responses. *Nature* 2020;22(586):594–9. <https://doi.org/10.1038/s41586-020-2814-7>.
- [18] Funk S, Salathé M, Jansen VAA. Modelling the influence of human behaviour on the spread of infectious diseases: a review. *J R Soc Interface* 2010;7(50):1247–56. <https://doi.org/10.1098/rsif.2010.0142>.
- [19] Funk S, Bansal S, Bauch CT, Eames KTD, Edmunds WJ, Galvani AP, et al. Nine challenges in incorporating the dynamics of behaviour in infectious diseases models. *Epidemics* 2015;10:21–5. <https://doi.org/10.1016/j.epidem.2014.09.005>.
- [20] Weston D, Hauck K, Amlôt R. Infection prevention behaviour and infectious disease modelling: a review of the literature and recommendations for the future. *BMC Public Health* 2018;18:336. <https://doi.org/10.1186/s12889-018-5223-1>.
- [21] Yan QL, Tang SY, Xiao YN. Impact of individual behaviour change on the spread of emerging infectious diseases. *Stat Med* 2018;37(6):948–69. <https://doi.org/10.1002/sim.7548>.
- [22] Scientific Pandemic Insights Group on Behaviours. SPI-B: Possible impact of the COVID-19 vaccination programme on adherence to rules and guidance about personal protective behaviours aimed at preventing spread of the virus – 7 December 2020; 2021. <https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/950723/s0978-spi-b-possible-impact-covid-19-vaccination-programme-adherence-to-rules-guidance.pdf>.
- [23] Brewer NT, Cuite CL, Herrington JE, Weinstein ND. Risk compensation and vaccination: can getting vaccinated cause people to engage in risky behaviors? *Ann Behav Med* 2007;34(1):95–99. <https://doi.org/10.1007/bf02879925>.
- [24] Office for National Statistics. Coronavirus and vaccine attitudes and behaviours in England: over 80s population, 15 February to 20 February 2021; 2021. <>.
- [25] YouGov. YouGov / Sky Survey Results; 2020. <https://docs.cdn.yougov.com/k6h1g211k5/Sky_CoronaBrexit_201005.pdf>.
- [26] Zewude B, Addis E. Changing patterns of compliance with protective behavioral recommendations in the post first-round COVID-19 vaccine period among healthcare workers in Southern Ethiopia 2021;14:3575–3587. <<https://doi.org/10.2147/RMHP.S325699>>.
- [27] Wright L, Steptoe A, Mak HW, Fancourt D. Do people reduce compliance with COVID-19 guidelines following vaccination? A longitudinal analysis of matched UK adults. *J Epidemiol Commun Health* 2021;76:109–15. <https://doi.org/10.1136/jech-2021-217179>.
- [28] Maclure M. The case-crossover design: a method for studying transient effects on the risk of acute events. *Am J Epidemiol* 1991;133(2):144–53. <https://doi.org/10.1093/oxfordjournals.aje.a115853>.
- [29] Maclure M, Mittleman MA. Should we use a case-crossover design? *Annu Rev Public Health* 2000;21:193–221. <https://doi.org/10.1146/annurev.publhealth.21.1.193>.
- [30] Mittleman MA, Mostofsky E. Exchangeability in the case-crossover design. *Int J Epidemiol* 2014;43(5):1645–55. <https://doi.org/10.1093/ije/dyu081>.
- [31] Mostofsky E, Coull BA, Mittleman MA. Analysis of observational self-matched data to examine acute triggers of outcome events with abrupt onset. *Epidemiology* 2018;29(6):804–16. <https://doi.org/10.1097/ede.0000000000000904>.
- [32] Hayward A, Fragaszy E, Kovar J, Nguyen V, Beale S, Byrne T, et al. Risk factors, symptom reporting, healthcare-seeking behaviour and adherence to public health guidance: protocol for Virus Watch, a prospective community cohort study. *BMJ Open* 2021;11(6):1–8. <https://doi.org/10.1136/bmjopen-2020-048042>.
- [33] UCL Virus Watch. [Internet]. [accessed 24 Nov 2022]. <<https://ucl-virus-watch.net>>.
- [34] UK Health Security Agency. Guidance for contacts of people with confirmed coronavirus (COVID-19) infection who do not live with the person; 2021. <<https://www.gov.uk/government/publications/guidance-for-contacts-of-people-with-possible-or-confirmed-coronavirus-covid-19-infection-who-do-not-live-with-the-person>>.
- [35] Warner L, Macaluso M, Austin HD, Kleinbaum DK, Artz L, Fleenor ME, et al. Application of the case-crossover design to reduce unmeasured confounding in studies of condom effectiveness. *Am J Epidemiol* 2005;15:161(8):765–773. <<https://doi.org/10.1093/aje/kwi094>>.
- [36] Bish A, Michie S. Demographic and attitudinal determinants of protective behaviours during a pandemic: a review. *Br J Health Psychol* 2010;5(4):797–824. <https://doi.org/10.1348/135910710X485826>.
- [37] Andrews N, Tessier E, Stowe J, Gower C, Kirsebom F, Simmons R, et al. Vaccine effectiveness and duration of protection of Comirnaty, Vaxzevria and Spikevax against mild and severe COVID-19 in the UK. MedRxiv 2021. <https://doi.org/10.1101/2021.09.15.21263583>.
- [38] Mizrahi B, Lotan R, Kalkstein N, Peretz A, Perez Galit, Ma MN, et al. Correlation of SARS-CoV-2 breakthrough infections to time-from-vaccine; preliminary study. MedRxiv 2021. <https://doi.org/10.1101/2021.07.29.21261317>.
- [39] Goldberg Y, Mandel M, Bar-On YM, Bodenheimer O, Freedman L, Haas EJ, et al. Waning immunity of the BNT162b2 vaccine: a nationwide study from Israel. MedRxiv 2021. <https://doi.org/10.1101/2021.08.24.21262423>.
- [40] GOV.UK. JCVI statement regarding a COVID-19 booster vaccine programme for winter 2021 to 2022; 2021. <<https://www.gov.uk/government/publications/jcvi-statement-september-2021-covid-19-booster-vaccine-programme-for-winter-2021-to-2022/jcvi-statement-regarding-a-covid-19-booster-vaccine-programme-for-winter-2021-to-2022>>.
- [41] Ferguson N, Ghani A, Cori A, Hogan A, Hinsley W, Volz E. population distribution and immune escape of Omicron in England. Imperial College London; 2021. <<https://doi.org/10.25561/93038>>.
- [42] Hodcroft EB. CoVariants: SARS-CoV-2 Mutations and Variants of Interest; 2021. <>.
- [43] Andrejko KL, Head JR, Lewnard JA, Remais JV. Longitudinal social contacts among school-aged children during the COVID-19 pandemic: the Bay Area Contacts among Kids (BACK) study. *BMC Infectious Diseases* 2022;22:242. <https://doi.org/10.1186/s12879-022-07218-4>.
- [44] Smith LE, Amlôt R, Lambert H, Oliver L, Robin C, Yardley L, Rubin GJ. Factors associated with adherence to self-isolation and lockdown measures in the UK: a cross-sectional survey. *Public Health* 2020;187:41–52. <https://doi.org/10.1016/j.puhe.2020.07.024>.
- [45] Smith LE, Mottershaw AL, Egan M, Waller J, Marteau TM, Rubin GJ. The impact of believing you have had COVID-19 on self-reported behaviour: Cross-sectional survey. *PLoS One* 2020;15:1–13. <https://doi.org/10.1371/journal.pone.0240399>.
- [46] Yan Y, Bayham J, Richter A, Fenichel EP. Risk compensation and face mask mandates during the COVID-19 pandemic. *Sci Rep* 2021;11(1):1–11. <https://doi.org/10.1038/s41598-021-82574-w>.
- [47] Nguyen V, Liu Y, Mumford R, Flanagan B, Patel P, Braithwaite I, et al. Changes in mobility pre and post first SARS-CoV-2 vaccination: findings from a prospective community cohort study including GPS movement tracking in England and Wales (Virus Watch). MedRxiv 2021. <https://doi.org/10.1101/2021.06.21.21259237>.
- [48] Majid U, Wasim A, Bakshi S, Truong J. Knowledge, (mis-)conceptions, risk perception, and behavior change during pandemics: a scoping review of 149 studies. *Public Underst Sci* 2020;29(8):777–99. <https://doi.org/10.1177/0963662520963365>.
- [49] Coronavirus (COVID-19) Infection Survey technical article waves and lags of COVID-19 in England, June 2021; 2021. <<https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/articles/coronaviruscovid19infectionsurveytechnicalarticle/wavesandlagsocovid19inenglandjune2021>>.
- [50] Cases in England. Coronavirus in the UK [Internet]. [cited 2022 Nov 13]. <<https://coronavirus.data.gov.uk/details/cases>>.
- [51] Deaths from Covid-19 (Coronavirus): The King's Fund [Internet]. [cited 2022 Nov 13]. <<https://www.kingsfund.org.uk/publications/deaths-covid-19>>.
- [52] Mishra S, Mindermann S, Sharma M, Whittaker C, Mellan TA, Wilton T, et al. Changing composition of SARS-CoV-2 lineages and rise of Delta variant in England. *eClinicalMedicine* 2021;39(101064). <<https://doi.org/10.1016/j.eclinm.2021.101064>>.
- [53] Aldridge RW, Pineo H, Fragaszy E, Eyre M, Kovar J, Nguyen V, et al. Household overcrowding and risk of SARS-CoV-2: analysis of the Virus Watch prospective community cohort study in England and Wales. MedRxiv 2021. <https://doi.org/10.1101/2021.05.10.21256912>.
- [54] Public Health England. Disparities in the risk and outcomes of COVID-19, 2020. <https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/908434/Disparities_in_the_risk_and_outcomes_of_COVID_August_2020_update.pdf>.
- [55] Joint Committee on Vaccination and Immunisation: advice on priority groups for COVID-19 vaccination, 2021. <<https://www.gov.uk/government/publications/priority-groups-for-coronavirus-covid-19-vaccination-advice-from-the-jcvi-30-december-2020/joint-committee-on-vaccination-and-immunisation-advice-on-priority-groups-for-covid-19-vaccination-30-december-2020>>.