

Optimal Treatment of the Symptomatic Smoker: Make a Diagnosis Not Empiric Treatment

Authors: Marc Miravittles (1), John R Hurst (2)

Center: 1. Pneumology Department, Hospital Universitari Vall d'Hebron/Vall d'Hebron Institut de Recerca (VHIR), Vall d'Hebron Barcelona Hospital Campus; CIBER de Enfermedades Respiratorias (CIBERES), Barcelona; Spain. **2 Include affiliation**

Correspondence: Marc Miravittles

Pneumology Department, Vall d'Hebron Barcelona Hospital Campus.

P. Vall d'Hebron 119-129. 08035 Barcelona, Spain.

Email: marcm@separ.es

Word count: 1038 (limit 1200)

A general paradigm of treatment in chronic diseases is that early diagnosis and treatment will improve long-term outcomes. In chronic obstructive pulmonary disease (COPD) this improvement in outcomes has been very difficult to demonstrate with current therapies such that pharmacologic treatments are currently prescribed with the principal aims of improving symptoms and preventing exacerbations (1). In part, this reflects the challenge of diagnosing 'early COPD', and the variable definitions that have been applied. Nevertheless, following this argument, if the objectives of COPD treatment are to mitigate symptoms and exacerbations, why not treat smokers or former smokers with these manifestations irrespective of pulmonary function impairment or diagnostic label? Current evidence supporting the efficacy and safety of inhaled medications in COPD derives from clinical trials in patients with a robust diagnosis of COPD; i.e. adult smokers or former smokers with poorly reversible airflow limitation. To date, there has been no evidence for treating symptomatic patients without airflow limitation (2,3).

The prevalence of respiratory symptoms is very frequent in the adult general population, the EPI SCAN 2 study on 9092 individuals older than 40 years in Spain reported at least one chronic respiratory symptom in 47.5% of participants and in 55% of those who reported being smokers (4). Similarly, in the SPIROMICS cohort (5), 50% of smokers with preserved pulmonary function had respiratory symptoms, and they experienced a mean of 0.27 respiratory infections ('exacerbations') per year. Woodruff et al (5) also reported that among symptomatic current and former smokers, 42% and 23% used off label bronchodilators and inhaled glucocorticoids respectively, without any evidence of efficacy and safety in this population.

In order to investigate the efficacy of dual bronchodilation (long-acting anticholinergic agents -LAMA- and long-acting beta-2 agonists -LABA) in symptomatic smokers or ex-smokers with preserved lung function, Han et al (6) conducted an important double-blind,

randomised, placebo controlled trial of indacaterol/glycopyrrolate. The study was conducted in persons who had a tobacco-smoking history of at least 10 pack-years, respiratory symptoms as defined by a COPD Assessment Test (CAT) score of at least 10 and preserved lung function on spirometry (ratio of forced expiratory volume in 1 second [FEV1] to forced vital capacity [FVC] ≥ 0.70 and FVC $\geq 70\%$ of the predicted value after bronchodilator use). The results showed that after 12 weeks of treatment there was not a significant effect of dual bronchodilation on the primary outcome, the St George's Respiratory Questionnaire (SGRQ); in fact, the proportion of participants with a clinically significant improvement of 4 units or greater in the SGRQ was 56.4% in the treatment group and 59% in the placebo group (adjusted odds ratio, 0.91; 95% confidence interval 0.60 to 1.37; $p=0.65$). The dual bronchodilator did have a physiological effect as demonstrated by an improvement in FEV1(% predicted) of 2.48% in the treatment arm versus -0.09% in the placebo arm, and the mean change in inspiratory capacity was 0.12 litres in the treatment arm versus 0.02 litres on placebo. However, these changes were not perceived clinically in terms of improvements in symptoms by the participants (6).

There are several aspects of this trial that merit discussion. Firstly, the population included is unusual, they had a high level of respiratory symptoms despite preserved spirometry. Their mean SGRQ score was 38.2 units, while the mean SGRQ in an adult population of smokers of more than 20 pack-years is 10.7 and the 95th percentile at ages 50 to 60 years is only 29.9 (7). Regarding dyspnea, they had a mean BDI score of 7.6 units, whereas patients in the TONADO trial with a mean FEV1(%) of only 50% had a BDI of 6.54 (8). Similarly, the mean CAT score was 17.6 units, similar to the 17 units observed in the POPE study in 3452 COPD patients with a mean FEV1(%) of 50% (9) and much worse than the 11 units in the study by Casanova et al (10) on 768 COPD patients with a mean FEV1(%) of 60%. Since dyspnea, SGRQ and CAT scores can be

influenced by several factors beyond COPD (11), co-morbidity such as cardiovascular disease, for example, these data suggest that the notable impairment in symptoms in this population with preserved lung function may be due, at least in part, to factors beyond the lungs, which may not be influenced using bronchodilators.

Another important aspect of the study is that patients already on inhaled medication and who were unable to undergo a 30-day medication washout were excluded. It has been argued that this strategy excluded possible responders to bronchodilators, which may be true. Nevertheless, the remaining subjects (those who were included) constitute the real target population: subjects with significant symptoms who are treatment naïve, and results in these patients were clearly negative.

Finally, the selection of SGRQ as the main outcome is controversial; SGRQ scores may be influenced by several factors, including psychological aspects. This can be illustrated by the mean improvement of almost 9 units in the placebo arm (more than twice the minimal clinically important difference of 4 units), and higher than the 7.7 unit improvement observed in the active treatment arm (6). However, the TDI breathlessness score, which may be more “respiratory specific” did not show any difference (0.92 units improvement with placebo and 0.93 with treatment) (6).

We can't rule out that the improvements in lung function parameters observed in the trial may translate to improved longer-term outcomes such as a reduction in the rate of lung function decline or prevention of exacerbations. Secondary analysis of the UPLIFT trial suggested that tiotropium significantly reduced the rate of decline in FEV1 in milder patients (GOLD 2) (12). Confirmation of this hypothesis in patients with preserved lung function would require larger and prolonged clinical trials. However, it may be challenging for patients to adhere to inhaled therapy for years without the perception of any symptomatic improvement.

The recent epidemiological study EPI SCAN 2 has shown a prevalence of COPD among the general population older than 40 years in Spain of 11.8%, with underdiagnosis still concerningly high at 74.7% (13). A recent survey showed that only 59% of adults with chronic respiratory symptoms consulted a physician and, of these, only two thirds underwent diagnostic spirometry (14). With this extremely high rate of underdiagnosis and the established, demonstrated efficacy of treatments in COPD patients, we believe that efforts should be directed to early diagnosis and treatment of airflow limitation; i.e. early COPD, whereas individuals with symptoms and preserved lung function should be encouraged to quit smoking and possible causes of respiratory symptoms beyond the lungs should be explored. We now know that bronchodilator therapy is, in general, not indicated in these patients.

Competing interests

Marc Miravittles has received speaker or consulting fees from AstraZeneca, Boehringer Ingelheim, Chiesi, Cipla, CSL Behring, Kamada, GlaxoSmithKline, Grifols, Menarini, Mereo Biopharma, Novartis, Palobiofarma SL, TEVA, Spin Therapeutics, and Zambon, and research grants from Grifols.

John Hurst has received support to attend meetings, and personal payment and payment to his employer (UCL), for educational and advisory work from pharmaceutical companies that make medicines to treat COPD.

References

1. Barrecheguren M, Gonzalez C, Miravittles M. What have we learned from observational studies and clinical trials of mild to moderate COPD? *Respir Res* 2018; 19: 177.
2. Nici L, Mammen MJ, Charbek E, Alexander PE, Au DH, Boyd CM, et al. Pharmacologic Management of Chronic Obstructive Pulmonary Disease. An Official

American Thoracic Society Clinical Practice Guideline. *Am J Respir Crit Care Med*. 2020; 201(9): e56-e69.

3. Miravittles M, Calle M, Molina J, Almagro P, Gómez JT, Trigueros JA, et al. Spanish COPD Guidelines (GesEPOC) 2021: Updated Pharmacological treatment of stable COPD. *Arch Bronconeumol*. 2022; 58: T69-T81.

4. Miravittles M, Soler-Cataluña JJ, Soriano JB, García-Río F, de Lucas P, Alfajeme I, et al. Respiratory symptoms and their determinants in the general Spanish population: Changes over 20 years. *ERJ Open Res* 2022 (accepted for publication).

5. Woodruff PG, Barr RG, Bleecker E, Christenson SA, Couper D, Curtis JL, et al. Clinical Significance of Symptoms in Smokers with Preserved Pulmonary Function. *N Engl J Med*. 2016 May 12;374(19):1811-21

6. Han MK, Ye W, Wang D, White E, Arjomandi M, Barjaktarevic IZ, et al. Bronchodilators in Tobacco-Exposed Persons with Symptoms and Preserved Lung Function. *N Engl J Med*. 2022 Sep 29;387(13):1173-1184.

7. Ferrer M, Villasante C, Alonso J, Sobradillo V, Gabriel R, Vilagut G, et al. General population norms of the St. George's respiratory questionnaire: a strategy to facilitate interpretation of quality of life scores. *Eur Respir J* 2002; 19: 405-413.

8. Buhl R, Maltais F, Abrahams R, Bjermer L, Derom E, Ferguson G, et al Tiotropium and olodaterol fixed-dose combination versus mono-components in COPD (GOLD 2-4). *Eur Respir J*. 2015 Apr;45(4):969-79.

9. Miravittles M, Koblizek V, Esquinas C, Milenkovic B, Barczyk A, Tkacova R, et al. Determinants of CAT (COPD Assessment Test) scores in a population of patients with COPD in central and Eastern Europe: The POPE study. *Respir Med* 2019; 150: 141-148.

10. Casanova C, Marin JM, Martinez-Gonzalez C, de Lucas-Ramos P, Mir-Viladrich I, Cosio B, et al. Differential Effect of Modified Medical Research Council Dyspnea, COPD Assessment Test, and Clinical COPD Questionnaire for Symptoms Evaluation Within the New GOLD Staging and Mortality in COPD. *Chest*. 2015 Jul; 148(1): 159-168.

11. Miravittles M, Molina J, Quintano JA, Campuzano A, Pérez J, Roncero C. Depressive status explains a significant amount of the variance in COPD assessment test (CAT) scores. *Int J Chron Obstruct Pulmon Dis* 2018; 13: 823-831.

12. Decramer M, Celli B, Kesten S, Lystig T, Mehra S, Tashkin DP; UPLIFT investigators. Effect of tiotropium on outcomes in patients with moderate chronic obstructive pulmonary disease (UPLIFT): a prespecified subgroup analysis of a randomised controlled trial. *Lancet*. 2009 Oct 3;374(9696):1171-8

13. Soriano JB, Alfajeme I, Miravittles M, de Lucas P, Soler-Cataluña JJ, García-Río F, et al. Prevalence and Determinants of COPD in Spain: EPISCAN II. *Arch Bronconeumol* 2021; 57: 61-69.

14. Calle Rubio M, Rodríguez Hermosa JL, Miravittles M, López-Campos JL. Knowledge of Chronic Obstructive Pulmonary Disease, Presence of Chronic Respiratory Symptoms and Use of Spirometry Among the Spanish Population: CONOCEPOC 2019 study. Arch Bronconeumol. 2021; 57(12): 741-749.