

TABLE 2. Histologic features of Wilms tumors in the cohort

| Histologic features | Wilms tumor type | |
|--------------------------|--------------------|--------------------|
| | n (%) | |
| | ET-WT (N = 118) | ST-WT (N = 149) |
| Epithelial component | | |
| 67-80% | 18 (15.3) | - |
| 81-100% | 100 (84.7) | - |
| Stromal component | | |
| 67-80% | - | 55 (36.9) |
| 81-100% | - | 94 (63.1) |
| Regressive / CIC changes | | |
| Absent | 24 (20.3) | 13 (8.7) |
| 1-10% | 58 (49.2) | 48 (32.2) |
| 11-50% | 33 (28) | 80 (53.7) |
| >50% | 3 (2.5) | 8 (5.4) |
| Blastemal component | | |
| Absent | 70 (59.3) | 43 (28.8) |
| 1-5% | 38 (32.2) | 86 (57.8) |
| 6-10% | 10 (8.5) | 20 (13.4) |
| Nephrogenic rests* | | |
| None | 69 (55.6) | 78 (50) |
| PLNR | 44 (35.5) | 16 (10.3) |
| ILNR | 3 (2.4) | 58 (37.2) |
| PLNR + ILNR | 7 (5.6) | 4 (2.6) |

ET-WT - epithelial type Wilms tumor; ST-WT - stromal type Wilms tumor; CIC - chemotherapy-induced changes

* Nephrogenic rests in 124 ET-WT and 156 ST-WT