

British Journal of Ophthalmology

Effects of socioeconomic status on baseline values and outcomes at 24 months in the Treatment of Advanced Glaucoma Study (TAGS) Randomized Controlled Trial

Journal:	<i>British Journal of Ophthalmology</i>
Manuscript ID	bjophthalmol-2022-321922.R1
Article Type:	Clinical science
Date Submitted by the Author:	n/a
Complete List of Authors:	King, Anthony; Nottingham University Hospitals NHS Trust, Ophthalmology Hudson, Jemma; University of Aberdeen Azulara-Blanco, Augusto; Queen's University Belfast, Centre for Public Health; Centre for Public Health, Queen's University Belfast Kirwan, James; Portsmouth Hospitals, ; Queen Alexandra Hospital, Goyal, Saurabh; Guy's and Saint Thomas' NHS Foundation Trust, Ophthalmology Lim, K Sheng; St Thomas Hospital, Ophthalmology Maclennan, Graeme; University of Aberdeen Health Services Research Unit, TAGS research group, TAGS research group; NUH
Keywords:	Glaucoma, Clinical Trial, Epidemiology

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7 **Effects of socioeconomic status on baseline values and outcomes at 24 months in**
8
9 **the Treatment of Advanced Glaucoma Study (TAGS) Randomized Controlled Trial**
10
11
12
13
14
15
16

17 **Authors**

18 Anthony J King¹ MD
19 Jemma Hudson² MSc
20 Augusto Azuara-Blanco³ PhD
21 James F Kirwan⁴ FRCOphth
22 Saurabh Goyal⁵ FRCOphth,
23 Kin Sheng Lim⁵ MD
24 Graeme MacClennan⁶ MSc
25
26 And the TAGS research group
27
28
29
30
31
32
33

34 **Corresponding Author**

35 Anthony J King
36 Department of Ophthalmology
37 Nottingham University Hospital
38 Nottingham, NG7 2UH
39 anthony.king@nottingham.ac.uk
40
41
42
43
44
45
46
47

- 48 1 Department of Ophthalmology, Nottingham University Hospital, Nottingham,
49 UK
50
51 2 Health Services Research Unit, University of Aberdeen, Health Services
52 Research Unit, University of Aberdeen, Aberdeen, UK.
53
54 3 Centre for Public Health, Queen's University Belfast, Royal Victoria Hospital,
55 Belfast, UK
56
57
58 4 Department of Ophthalmology, Queen Alexandra Hospital, Portsmouth, UK
59
60

- 1
2
3
4 5 Ophthalmology Department, St Thomas' Hospital, London, UK
5 6 Centre for Healthcare Randomised Trials (CHaRT), Health Services Research
6 7 Unit, University of Aberdeen, UK.
8
9
10
11

12 Word Count: 3125
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

TAGS Study Group:

31 Anthony King, Pavi Agrawal, Richard Stead, David C Broadway, Nick Strouthidis, Shenton
32 Chew, Chelvin Sng, Marta Toth, Gus Gazzard, David Garway-Heath, Keith Barton, Ahmed
33 Elkarmouty, Eleni Nikita, Giacinto Triolo, Soledad Aguilar-Munoa, Saurabh Goyal, Sheng
34 Lim, Velota Sung, Imran Masood, Nicholas Wride, Amanjeet Sandhu, Elizabeth Hill, John
35 Sparrow, Fiona Grey, Rupert Bourne, Gnanapragasam Nithyanandarajah, Catherine
36 Willshire, Philip Bloom, Faisal Ahmed, Francesca Cordeiro, Laura Crawley, Eduardo
37 Normando, Sally Ameen, Joanna Tryfinopoulou, Alistair Porteous, Gurjeet Jutley, Dimitrios
38 Bessinis, James Kirwan, Shahiba Begum, Anastasios Sepetis, Edward Rule, Richard
39 Thornton, Andrew McNaught, Nitin Anand, Anil Negi, Obaid Kousha, Marta Hovan, Roshini
40 Sanders, Pankaj Kumar Agarwal, Andrew Tatham, Leon Au, Cecelia Fenerty, Tanya
41 Karaconji, Brett Drury, Duya Penmol, Ejaz Ansari, Albina Dardzhikova, Reza Moosavi, Richard
42 Imonikhe, Prodromos Kontovourikis, Luke Membrey, Goncalo Almeida, James
43 Tildsley, Augusto Azuara-Blanco, Angela Knox, Simon Rankin, Sara Wilson, Avinash
44 Prabhu, Subhanjan Mukherji, Amit Datta, Alisdair Fern, Joanna Liput, Tim Manners, Josh
45 Pilling, Clare Stemp, Karen Martin, Tracey Nixon, Caroline Cobb, Alan Rotchford, Sikander
46 Sidiki, Atul Bansal, Obaid Kousha, Graham Auger, Mary Freeman, Gordon Fernie, Alison
47 MacDonald, Ashleigh Kernohan, Jennifer Burr, Tara Homer, Hosein Shabaninejad, Luke Vale,
48 John Norrie.
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Confidential: For Review Only

Precis

For patients presenting with advanced glaucoma SES influences disease severity at presentation but has no effect on success of either medical or surgical treatment interventions.

Key Messages

What is already known on this topic

Poorer SES is known to increase the risk of presenting with advanced glaucoma.

What this study adds

This study demonstrates that those presenting with already advanced glaucoma are from poorer SES background and have worse visual field loss in both eyes and poorer vision related QoL, however SES does not affect the success of treatment.

How this study may affect research, practice or policy

This study further emphasises the need to increase awareness of glaucoma and improve health seeking behaviour amongst those from poorer SES.

Abstract

Background/Aims: Socio-economic status (SES) is associated with late disease presentation and poorer outcomes. We evaluate the effect of SES on treatment outcomes and report the correlation between SES and baseline characteristics of participants in the Treatment of Advanced Glaucoma Study.

Methods: Pragmatic multicentre randomised control trial. Four hundred and fifty-three patients presenting with advanced open angle glaucoma in at least one eye [Hodapp-Parrish-Anderson classification]. Participants were randomised to either glaucoma drops (medical arm) or trabeculectomy (surgery arm). Clinical characteristics, Quality of life measurement (QoL) and SES defined by the index of multiple deprivation (IMD) are reported. Subgroup analysis explored treatment effect modifications of SES at 24 months. Correlation between SES and baseline characteristics was tested with the Chi-squared test of association for dichotomous variables and Pairwise Pearson's correlation for continuous variables.

Results: The mean visual field MD was -17.2(6.7)dB for the most deprived quintile of participants and -13.0(5.5) for the least deprived quintile in the index eye. At diagnosis there was a strong correlation between SES and ethnicity, age, extent of visual field loss and number of visits to opticians prior to diagnosis. At 24 months there was no evidence that the treatment effect was moderated by SES.

Conclusions: In patients presenting with advanced glaucoma, SES at baseline is correlated with poorer visual function, poorer VFQ-25 quality of life, ethnicity, age and number visits to an optician in the years preceding diagnosis. SES at baseline does not have an effect of the success of treatment at 24 months

Trial registration: Health Technology Assessment (NIHR-HTA) Programme (Project number: 12/35/38). ISRCTN registry: ISRCTN56878850

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Confidential: For Review Only

Introduction

Socioeconomic inequality is recognised to contribute to poorer health outcomes for those from poorer socioeconomic backgrounds in the UK (1) and North America (2-4).

Glaucoma is a chronic progressive eye disease, with substantial and detrimental effects on numerous aspects of daily living(5) a major cause of disability in the elderly (6,7) and worsening of Health-Related Quality of Life HRQoL (5). It is the second most common cause of irreversible blindness in the UK, North America and Europe(8,9). The number of patients with glaucoma is predicted to increase substantially as the result of an ageing population(10, 11).

While the incidence and prevalence of glaucoma are commonly reported(11) more granular data regarding the severity of glaucoma at the time of diagnosis is rarely available. In North America, Hattenhauer(12) reported 10% (29/295) of his cohort was blind from glaucoma in at least one eye at diagnosis, while Buys(13) reported 21% (60/290) of a cohort of newly diagnosed glaucoma had severe glaucoma in at least one eye at diagnosis. In the UK the approximately 1 in 4 patients present with advanced disease(14 – 19). , Previous reports from the UK and North America have shown that presentation with advanced glaucoma at diagnosis is more common amongst those people from a poorer socioeconomic back ground(13, 18, 20-23) and it is also recognised that health care usage may be affected by racial or SES background(24 – 25). However, there has been no previous opportunity to specifically look at the characteristics of a large cohort of patients who present with advanced disease or to establish if SES has an effect on treatment outcomes.

Understanding factors that influence the severity of disease presentation in those presenting with already advanced glaucoma is important as presentation with advanced glaucoma is a major risk factor for lifetime blindness and those presenting with advanced glaucoma are those most at risk of developing blindness in their

lifetime(26 – 32). Understanding if there is an effect of SES on treatment outcomes is important as it may influence the choice of treatment recommended.

Effective treatment for glaucoma stops or delays disease progression(33). The Treatment of Advanced Glaucoma Study (TAGS) is a pragmatic randomised controlled trial (RCT) designed to determine whether primary medicine or primary surgery is more effective for patients presenting with advanced glaucoma(34, 35).

The TAGS study has provided an opportunity to evaluate at a more granular level the SES and racial background of those who present with advanced glaucoma in a large cohort of advanced glaucoma patients. It also provides an opportunity to explore these factors on the outcomes of treatment which has not been previously undertaken.

The aim of this report therefore is to explore the effect of SES on quality of life and clinical measures for those presenting with advanced glaucoma and to explore whether SES affects the effectiveness of either medical or surgical management in these patients at 24 months.

Methods

TAGS is a pragmatic multicentre RCT; the design of the study has been described in detail elsewhere(34). Briefly, eligible patients with advanced POAG in either eye were randomised to have augmented trabeculectomy or IOP lowering drops as their primary intervention and followed up for 24 months. Randomisation was based on the participant (not the eye), for those where both eyes were eligible, clinical outcomes are based on the index eye defined as the eye with better mean deviation (MD) value. Research was undertaken in compliance with the tenets of the Declaration of Helsinki and Ethics for this research granted by East Midlands – Derby Research Ethics Committee (reference number 13/EM/0395).

Disease Classification: Eligible patients had primary open angle glaucoma (including pigment dispersion and pseudoexfoliation). Advanced glaucoma was defined according to the Hodapp-Parrish-Anderson (HPA) classification of glaucoma(36). The HPA classification is commonly used in glaucoma research and uses both the position and extent of visual field loss to categorise severity(36). Advanced disease was classified according to the “severe” category of visual field loss using the HPA classification [has **any** of the following]:

1. Mean Deviation (MD) < -12.00dB
2. More than 50% of points depressed below the 5% level on the pattern deviation probability plot
3. More than 20 points depressed below the 1% level on the pattern deviation probability plot
4. A point in the central five degrees has a sensitivity of 0-dB
5. Points within five degrees of fixation under 15 dB sensitivity in both upper and lower hemi-fields.

1
2
3
4
5
6
7
8 *Outcome Measures:* At baseline clinical measurements (visual field loss Mean
9 Deviation VFMD), Logarithm of the mean angle of resolution (logMAR) visual acuity
10 (VA), intraocular pressure (IOP), incidence of blindness(37), family history of
11 glaucoma and self-reported frequency of contact with primary care optometry in years
12 prior to diagnosis were recorded. In addition, Health Related Quality of Life (HRQoL)
13 generic health status [EuroQual-5 dimension – 5 level (EQ-5D-5L)](38) and Health
14 Utility Index (HUI-3)(39), visual health status [National Eye Institute Visual Function
15 Questionnaire 25 (VFQ-25)](40), glaucoma health status [glaucoma utility index
16 (GUI)](41) and patient experience were also recorded at baseline. Outcomes
17 examined and reported at 24 months were VFMD, logMAR VA, IOP and HRQoL.
18
19 SES was defined using the Index of Multiple Deprivation (IMD). The IMD is the official
20 measure of relative deprivation for small areas in England. The IMD ranks every small
21 area in England based on postcode from 1 (most deprived area) to 32,844 (least
22 deprived area). The IMD combines information from the seven domains (Income
23 Deprivation, Employment Deprivation, Education, Skills and Training Deprivation,
24 Health Deprivation and Disability, Crime, Barriers to Housing and Services and Living
25 Environment Deprivation) to produce an overall relative measure of deprivation
26
27 ([https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attach
28 ment_data/file/853811/loD2019_FAQ_v4.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/853811/loD2019_FAQ_v4.pdf)).

29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49 *Statistical analysis:* Baseline characteristics are described using numbers and
50 percentages for dichotomous variables, numbers, median and interquartile range
51 (IQR) for the number of times the participant visited the optometrist in the last 10 years
52 and mean and standard deviation (SD) for all remaining continuous variables. For
53 participants in whom both eyes were eligible, data are summarised for both the index
54 and non-index eye. The Chi-squared test of association was used to assess
55
56
57
58
59
60

1
2
3
4 correlation between IMD scores derived from participants' postcodes and
5 dichotomous variables (gender, ethnicity and family history), and Pairwise Pearson's
6 correlation for continuous variables. As TAGS recruited from England, Scotland and
7 Northern Ireland, we used the Abel et al method to convert the IMD to be on the same
8 scale centred on English scores as 89% of participants were recruited in England(42),
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

IMD is reported in quintiles from 1 (most deprived) through to 5 (least deprived). EQ-5D-5L was calculated following the method by Van Hout et al(43) and GUI was calculated following the method by Burr et al(41). Outcomes at 24 months were analysed using mixed effects linear model correcting for baseline score, bilateral disease and including a random effect for surgery and treatment using restricted maximum likelihood following the partially nested heteroscedastic method by Candlish et al(44) Subgroup analysis explored treatment effect modifications of IMD on IOP, logMAR, VFMD, NEI-VFQ-25, HUI-3 and GUI at 24 months. We used a stricter level of significance (two-sided 1% significance level) and 99% CIs. Subgroup by treatment interaction was assessed by included interaction terms in the model outlined above. Due to imbalances between IMD group as baseline for gender, age and ethnicity a sensitivity analysis adjusted for these covariates. As conclusions did not change, results are not provided. Outcomes by IMD were also analysed using linear regression adjusting for baseline score for overall cohort and by treatment arm at 24 months. All analyses were performed in Stata 16 software. (45)

Results

All patient recruited to TAGS had advanced glaucoma in at least one eye according to the Hodapp classification at diagnosis. The IMD quintiles for the randomised participants were roughly equal between the two treatment arms (Supplementary Table 1) and across each quintile.

Table 1 shows the patient characteristic values for baseline data for each quintile of deprivation for the whole cohort. It can be seen that on average there is nearly an eight-year difference in age of diagnosis between those in the lowest and highest deprivation groups with those in the most deprived group being younger. Similarly, there is a much higher proportion of non-Caucasian patients in the lower socioeconomic groups.

For patient reported outcomes, there is a 5-point difference between the upper and lower quintiles for composite vision specific VFQ score. This is represented most markedly in the near activities, dependency, driving, general health, role difficulties, mental health, general vision and ocular pain domains. There was no material difference for either of the generic HRQoL measures or the glaucoma specific GUI between quintiles.

Table 2 reports the baseline clinical characteristics and IMD for the index eye. It can be seen that on average those in the lowest SE quintile have 4dB more VF loss at presentation than those with highest SES. Similarly, those in the lowest quintile have on average a higher IOP by about 4mmHg at diagnosis. Similar observations are made for the non-index eye (supplementary Table 2). The IOP was highest at diagnosis in the lowest SES quintile, the lower IOP noted in quintile 5 is a reflection that a higher proportion of patients presented with IOP < 22mmHg (IOP<22mmHG, 29, 29,27,29,35% for quintiles 1 – 5 respectively) suggesting a higher proportion of normal tension glaucoma in this group. However, at baseline (following initial medications) the

1
2
3 IOP is reduced to a similar level across the quintile spectrum for both index and non-
4 index eyes.
5
6

7
8 Table 3 shows that the SES differences at baseline are highly statistically significant
9 for age, ethnicity, number of visits to optometrists, vision specific health (VFQ-25) and
10 visual field MD for both the index and non-index eye.
11
12

13
14 Subgroup analysis for IOP, VF MD and logMAR visual acuity and QoL measures at
15 24 months are shown in Figure 1 and Supplementary Figure 1. The trabeculectomy
16 arm had lower IOP for all IMD quintile groups with the biggest difference for the second
17 (MD -3.67 95% CI (-7.05, -0.30); p-value 0.005) and third quintile (MD -3.67, 95% CI
18 (-6.68, -0.83); p-value 0.001. However, there was no evidence that the treatment effect
19 was moderated by the quintiles. There was no material difference for VA or VF loss
20 between the trabeculectomy or medical management arms based on IMD score and
21 no evidence that the treatment effect was moderated by the quintiles (Figure 1).
22 Subgroup analysis for IMD and quality of life at 24 months showed there was no
23 evidence that the treatment effect was moderated by the quintiles (Supplementary
24 Figure 1)
25
26
27
28
29
30
31
32
33
34
35
36
37

38 There were no material differences between quintiles for outcomes based on
39 intervention either for the cohort as a whole (Supplementary Table 3) or for either the
40 trabeculectomy or medical management arms (Tables 4).
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

TAGS was designed to be a pragmatic trial comparing established options, medications or surgery, as initial treatment for people diagnosed with severe glaucoma. Only the primary intervention was dictated by the trial protocol(34). The effect of SES on health outcomes is well recognised in terms of mortality(1) and for specific diseases such as cancer(1, 46 – 51) and cardiovascular disease(52, 53), with those from more deprived backgrounds having poorer outcomes.

TAGS is the largest RCT comparing treatments for advanced glaucoma and offered a unique opportunity to explore how outcomes for patients with advanced glaucoma are affected by SES and whether there is a difference of outcomes based on SES for medical or surgical interventions. In addition, the large number of patients, all with advanced glaucoma provided a unique opportunity for a more in-depth exploration of the effects of SES on patients presenting with advanced glaucoma using prospectively collected data in a representative sample of a population with advanced glaucoma(54). The analysis used participants' postcode as a proxy for SES to look at this important dimension *within* those presenting with advanced disease.

Several previous studies have similarly indicated that for glaucoma advanced presentation compared to non-advanced disease is linked to poorer socioeconomic status(18, 20-22), however unlike TAGS this has been data collected from retrospectively evaluated cohorts.

At baseline in our cohort, poorer socioeconomic background correlates with more advanced disease at presentation in terms of visual field loss in the index and non-index eye, indicating that the relationship between more advanced visual field loss and poorer SES exists even in a cohort with advanced glaucoma and is present in both eyes. This may explain the correlation between poorer socioeconomic status and poorer vision related QoL, the subscales most affected are those of near activities, role

1
2
3
4 limitation and dependency. Near activities refer to limited ability to read small print and
5
6 undertake some personal tasks such as shaving and putting on makeup which require
7
8 near vision clarity. Role limitations refers to requirement for more help from others in
9
10 undertaking everyday tasks and limiting activities because of vision and dependency
11
12 specifically refers to limiting activity outside the home because of vision (Table 1). It is
13
14 reasonable to assume that these activities will be affected by more advanced visual
15
16 field loss particularly when affecting both eyes(55). It is likely that patients with
17
18 advanced glaucoma have some awareness of vision deterioration especially if
19
20 affecting both eye(55). One possibility that this reduced vision may not have prompted
21
22 patients to seek attention earlier, is a resignation among older people that poorer
23
24 vision is a natural consequence of ageing(56) and they may not therefore pay much
25
26 attention to the subtle and slowly developing deterioration associated with visual field
27
28 loss, especially if only affecting one eye. Previous studies have identified older
29
30 age(19, 20) and ethnicity as risk factors for presentation with advanced glaucoma.
31
32 However further evaluation of this in the TAGS cohort of patients presenting with
33
34 advanced disease reveals that patients with lower SES are younger, have more
35
36 advanced disease at presentation in both eyes and are more commonly from a non-
37
38 Caucasian ethnic background. This is important in terms of lifetime vision loss
39
40 prevention as those patients with the most severe vision loss are most likely to lose
41
42 vision during their lifetime even with treatment and as these patients are the youngest
43
44 they are more likely to spend a longer period of their life with severe visual disability.
45
46 Indeed over 6% of the cohort were eligible for sight impairment registration at the time
47
48 of diagnosis(37) and 10.6% did not achieve the visual standards for driving, once again
49
50 these were seen more commonly in those with lower SES.

51
52
53 One mechanism for minimising risk of presentation with advanced glaucoma is a
54
55 regular visit to an eye care professional. In England, current policy facilitates visits to
56
57 a community optometrist annually for those over 40 years with a family history of
58
59 glaucoma. In TAGS, there is a correlation between fewer visits to the optician in the
60

1
2
3
4 previous 10 years and SES. In the UK and many countries glaucoma diagnosis is a
5
6 consequence of opportunistic case finding, fewer visits means less opportunity for
7
8 those of lower SES to have their glaucoma picked up through routine optometrist eyes
9
10 visits, thus resulting in more advanced presentation. Indeed, Shickle (57) observed
11
12 that “A person aged 60 or over living in the least deprived quintile is 71% more likely
13
14 to attend for an eye examination than someone in that age group in the most deprived
15
16 quintile in Leeds, even though both have the same entitlement”. This may be
17
18 representative of the UK as a whole explaining this observation. The reason these
19
20 opportunities to diagnose glaucoma earlier are missed is unknown. Several
21
22 possibilities exist, It is possible that even though entitlement is equal for all SES groups
23
24 the messaging around need to attend for routine testing or awareness of these
25
26 services may be less for those from lower SES groups(58 – 60) and therefore improved
27
28 awareness and education of those at risk may help to overcome this. It is also possible
29
30 that participants have some recall bias and over-estimated the frequency of visits to
31
32 their optometrists prior to diagnosis or that they were rapid progressors as previously
33
34 suggested by Fraser(20). It has also been suggested that delays in diagnosis may
35
36 occur at several distinct points in a patient pathway, from failure of
37
38 recognition/diagnosis of glaucoma by optometrists, to failure to refer appropriately or
39
40 delays in this process occurring(61).

41
42
43 Nearly one third of our cohort reported a family history of glaucoma which is similar to
44
45 three previous primary intervention studies of patients with early glaucoma(62-64),
46
47 suggesting that having a family history of glaucoma does not reduce your risk of
48
49 presenting with advanced disease. This is a disappointing observation as in the UK
50
51 patients 40 years of age with a family history of glaucoma are entitled to a free
52
53 glaucoma screening eye exam annually and it would be hoped that this would reduce
54
55 the risk of presenting with advanced disease.

56
57
58 These findings suggest that, despite a robust public health provision in the UK to
59
60 prevent diagnosis with advanced disease, a large number of patients are still not being

1
2
3
4 diagnosed at an early stage and this disproportionately affects those from poorer SES
5
6 backgrounds who have more severe visual field loss and more reduction in visual QoL
7
8 at diagnosis. Addressing this should be a research priority.
9
10

11
12 TAGS employed two current standard interventions for lowering IOP. There was no
13
14 difference between SES quintiles in outcomes at 24 months for the cohort as a whole
15
16 (Supplementary Table 3). There was a significant difference between treatment arms
17
18 for the IOP measurement for all quintiles of SES, however this difference did not
19
20 appear to vary between different quintiles of SES and represents the general
21
22 difference in IOP lowering achieved in the study between the medical and surgical
23
24 arms of the study (Figure 1). There was no difference in any outcomes between
25
26 quintiles within either the medically treated arm or the trabeculectomy arm of the study.
27
28 It is not surprising following surgery that IOP remains relatively equivalent between
29
30 quintiles as there is generally no reliance on further treatment delivered by the patient,
31
32 however in the medicine group a reliance on use of drops remains and is ongoing.
33
34 Adherence is a recognised problem for glaucoma patients using drops(65) and poorer
35
36 adherence is correlated with greater glaucoma progression(66). Both age and
37
38 educational achievement have been implicated as barriers to adherence(65) and
39
40 these differ between the quintiles of the medical arm of the TAGS cohort, potentially
41
42 leading to a difference in adherence to drops(67). However, there is no difference
43
44 detected in IOP between quintiles in the TAGS medical arm to suggest that SES has
45
46 an influence on the level of IOP achieved and maintained.
47
48
49
50
51
52

53
54 In conclusion in patients presenting with advanced glaucoma, SES influences the
55
56 severity of presentation with those from lower SES having more advanced visual field
57
58 loss in both eyes at presentation and worse vision related quality of life, they also tend
59
60 to be younger and more commonly from non-Caucasian ethnicity. Following treatment

1
2
3
4 SES does not appear to influence the success of either medical or surgical outcomes
5
6 at 24 months.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Confidential: For Review Only

1
2
3 **Competing Interest Statement**
4
5

6 None of the authors have any competing interests relating to this project
7
8
9

10
11
12
13
14
15 **Contributorship Statement**
16

17
18 AJK - study design, data collection, manuscript writing and review,
19

20
21 JH - study design, data analysis, manuscript writing and review
22

23
24 AAB - study design, data collection, manuscript review,
25

26
27 GMc - study design, data analysis, manuscript review
28

29
30 JK - study design, data collection, manuscript review
31

32
33 SG - study design, data collection, manuscript review
34

35
36 SL - study design, data collection, manuscript review
37

38
39 All authors confirm approval of final submitted manuscript
40
41
42
43
44
45

46 **Ethics Statement**
47

48
49 Ethics for this research granted by East Midlands – Derby Research Ethics Committee
50

51
52 (reference number 13/EM/0395).
53
54
55
56
57
58
59
60

References

1. Marmot M, Allen J, Boyce T, Goldblatt P, J M. Strategic Review of Health Inequalities in England post-2010. (2010). Fair society, healthier lives: The Marmot review.; 2020.
2. Bashinskaya B, Nahed BV, Walcott BP, Coumans JV, Onuma OK. Socioeconomic status correlates with the prevalence of advanced coronary artery disease in the United States. *PLoS One*. 2012;7(9):e46314.
3. Golombos DM, O'Malley P, Lewicki P, Nguyen DP, Stone BV, Al Hussein Al Awamlh B, et al. The impact of socioeconomic status on perioperative complications and oncologic outcomes in patients undergoing radical cystectomy. *World J Urol*. 2017;35(7):1063-71.
4. Nicklett EJ. Socioeconomic status and race/ethnicity independently predict health decline among older diabetics. *BMC Public Health*. 2011;11:684.
5. Sotimehin AE, Ramulu PY. Measuring Disability in Glaucoma. *J Glaucoma*. 2018;27(11):939-49.
6. Haymes SA, Leblanc RP, Nicolela MT, Chiasson LA, Chauhan BC. Risk of falls and motor vehicle collisions in glaucoma. *Invest Ophthalmol Vis Sci*. 2007;48(3):1149-55.
7. Haymes SA, LeBlanc RP, Nicolela MT, Chiasson LA, Chauhan BC. Glaucoma and on-road driving performance. *Invest Ophthalmol Vis Sci*. 2008;49(7):3035-41.
8. Flaxman SR, Bourne RRA, Resnikoff S, Ackland P, Braithwaite T, Cicinelli MV, et al. Global causes of blindness and distance vision impairment 1990-2020: a systematic review and metaanalysis. *Lancet Glob Health*. 2017;5(12):e1221-e34.
9. Quartilho A, Simkiss P, Zekite A, Xing W, Wormald R, Bunce C. Leading causes of certifiable visual loss in England and Wales during the year ending 31 March 2013. *Eye (Lond)*. 2016;30(4):6027.
10. The Way Forward - Glaucoma. London: Royal College of Ophthalmologists; 2015.

11. Tham YC, Li X, Wong TY, Quigley HA, Aung T, Cheng CY. Global prevalence of glaucoma and projections of glaucoma burden through 2040: a systematic review and meta-analysis. *Ophthalmology*. 2014;121(11):2081-90.
12. Hattenhauer MG, Johnson DH, Ing HH, Herman DC, Hodge DO, Yawn BP, et al. The probability of blindness from open-angle glaucoma. *Ophthalmology*. 1998;105(11):2099-104.
13. Buys YM, Jin YP, Canadian Glaucoma Risk Factor Study G. Socioeconomic status as a risk factor for late presentation of glaucoma in Canada. *Can J Ophthalmol*. 2013;48(2):83-7.
14. Coffey M, Reidy A, Wormald R, Xian WX, Wright L, Courtney P. Prevalence of glaucoma in the west of Ireland. *Br J Ophthalmol*. 1993;77(1):17-21.
15. Sheldrick JH, Ng C, Austin DJ, Rosenthal AR. An analysis of referral routes and diagnostic accuracy in cases of suspected glaucoma. *Ophthalmic Epidemiol*. 1994;1(1):31-9.
16. Elkington AR, MacKean J, P. S. A collaborative hospital glaucoma survey. *Res Clin Forums*. 1982;4:31-40.
17. Boodhna T, Crabb DP. Disease severity in newly diagnosed glaucoma patients with visual field loss: trends from more than a decade of data. *Ophthalmic Physiol Opt*. 2015;35(2):225-30.
18. Sukumar S, Spencer F, Fenerty C, Harper R, Henson D. The influence of socioeconomic and clinical factors upon the presenting visual field status of patients with glaucoma. *Eye*. 2009;23(5):1038-44.
19. Ng WS, Agarwal PK, Sidiki S, McKay L, Townend J, Azuara-Blanco A. The effect of socioeconomic deprivation on severity of glaucoma at presentation. *British Journal of Ophthalmology*. 2010;94(1):85-7.
20. Fraser S, Bunce C, Wormald R. Risk factors for late presentation in chronic glaucoma. *Invest Ophthalmol Vis Sci*. 1999;40(10):2251-7.
21. Fraser S, Bunce C, Wormald R. Retrospective analysis of risk factors for late presentation of chronic glaucoma. *Br J Ophthalmol*. 1999;83(1):24-8.
22. Fraser S, Bunce C, Wormald R, Brunner E. Deprivation and late presentation of glaucoma: case-control study. *BMJ*. 2001;322(7287):639-43.
23. Shweikh Y, Ko F, Chan MP, Patel PJ, Muthy Z, Khaw PT, et al. Measures of socioeconomic status and self-reported glaucoma in the U.K. Biobank cohort. *Eye (Lond)*. 2015;29(10):1360-7.
24. Halawa OA, Kolli A, Oh G, Mitchell WG, Glynn RJ, Kim DH, et al. Racial and Socioeconomic Differences in Eye Care Utilization among Medicare Beneficiaries with Glaucoma. *Ophthalmology*. 2021.
25. Delavar A, Radha Saseendrakumar B, Weinreb RN, Baxter SL. Racial and Ethnic Disparities in

- 1
2
3 Cost-Related Barriers to Medication Adherence Among Patients With Glaucoma Enrolled in the
4 National Institutes of Health All of Us Research Program. *JAMA Ophthalmol.* 2022.
- 5
6 26. Peters D, Bengtsson B, Heijl A. Factors associated with lifetime risk of open-angle glaucoma
7 blindness. *Acta Ophthalmol.* 2014;92(5):421-5.
8
9 27. Grant WM, Burke JF, Jr. Why do some people go blind from glaucoma? *Ophthalmology.*
10 1982;89(9):991-8.
11
12 28. Mokhles P, Schouten JS, Beckers HJ, Azuara-Blanco A, Tuulonen A, Webers CA. A Systematic
13 Review of End-of-Life Visual Impairment in Open-Angle Glaucoma: An Epidemiological Autopsy. *J*
14 *Glaucoma.* 2016;25(7):623-8.
15
16 29. Odberg T. Visual field prognosis in advanced glaucoma. *Acta Ophthalmol* 1987;65 (suppl):27- 9.
17
18 30. Oliver JE, Hattenhauer MG, Herman D, Hodge DO, Kennedy R, Fang-Yen M, et al. Blindness
19 and glaucoma: a comparison of patients progressing to blindness from glaucoma with patients
20 maintaining vision. *Am J Ophthalmol.* 2002;133(6):764-72.
21
22 31. Mikelberg FS, Schulzer M, Drance SM, Lau W. The rate of progression of scotomas in
23 glaucoma. *Am J Ophthalmol.* 1986;101(1):1-6.
24
25 32. Wilson R, Walker AM, Dueker DK, Crick RP. Risk factors for rate of progression of
26 glaucomatous visual field loss: a computer-based analysis. *Arch Ophthalmol.* 1982;100(5):737-41.
27
28 33. Maier PC, Funk J, Schwarzer G, Antes G, Falck-Ytter YT. Treatment of ocular hypertension
29 and open angle glaucoma: meta-analysis of randomised controlled trials. *BMJ.* 2005;331(7509):134.
30
31 34. King AJ, Fernie G, Azuara-Blanco A, Burr JM, Garway-Heath T, Sparrow JM, et al. Treatment
32 of Advanced Glaucoma Study: a multicentre randomised controlled trial comparing primary medical
33 treatment with primary trabeculectomy for people with newly diagnosed advanced glaucoma-study
34 protocol. *Br J Ophthalmol.* 2018;102(7):922-8.
35
36 35. King AJ, Hudson J, Fernie G, Kernohan A, Azuara-Blanco A, Burr J, et al. Primary
37 trabeculectomy for advanced glaucoma: pragmatic multicentre randomised controlled trial (TAGS).
38 *BMJ.* 2021;373:n1014.
39
40 36. Hodapp E, Parrish RK, Anderson DR. Clinical decision in glaucoma. St Louis MO: Mosby;
41 1993.
42
43 37. Department of Health. Certificate of Vision Impairment: Explanatory Notes for Consultant
44 Ophthalmologists and Hospital Eye Clinic Staff in England. Leeds: Department of Health; 2017
45 [Available from:
46 [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/637590/CVI_guidance.pdf)
47 [/637590/CVI_guidance.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/637590/CVI_guidance.pdf).
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 38. Herdman M, Gudex C, Lloyd A, Janssen M, Kind P, Parkin D, et al. Development and
4 preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res.*
5 2011;20(10):1727-36.
6
7
8 39. Tosh J, Brazier J, Evans P, Longworth L. A review of generic preference-based measures of
9 health-related quality of life in visual disorders. *Value Health.* 2012;15(1):118-27.
10
11 40. Mangione CM, Lee PP, Gutierrez PR, Spritzer K, Berry S, Hays RD, et al. Development of the
12 25-item National Eye Institute Visual Function Questionnaire. *Arch Ophthalmol.* 2001;119(7):1050-8.
13
14 41. Burr JM, Kilonzo M, Vale L, Ryan M. Developing a preference-based Glaucoma Utility Index
15 using a discrete choice experiment. *Optom Vis Sci.* 2007;84(8):797-808.
16
17 42. Abel GA, Barclay ME, Payne RA. Adjusted indices of multiple deprivation to enable
18 comparisons within and between constituent countries of the UK including an illustration using
19 mortality rates. *BMJ Open.* 2016;6(11):e012750.
20
21 43. van Hout B, Janssen MF, Feng YS, Kohlmann T, Busschbach J, Golicki D, et al. Interim scoring
22 for the EQ-5D-5L: mapping the EQ-5D-5L to EQ-5D-3L value sets. *Value Health.* 2012;15(5):708-
23 15.
24
25 44. Candlish J, Teare MD, Dimairo M, Flight L, Mandefield L, Walters SJ. Appropriate statistical
26 methods for analysing partially nested randomised controlled trials with continuous outcomes: a
27 simulation study. *BMC Med Res Methodol.* 2018;18(1):105.
28
29 45. StataCorp. *Stata Statistical Software*. College Station, TX: StataCorp LLC; 2019.
30
31 46. Nur U, Quaresma M, De Stavola B, Peake M, Rachet B. Inequalities in non-small cell lung
32 cancer treatment and mortality. *J Epidemiol Community Health.* 2015;69(10):985-92.
33
34 47. Morris M, Woods LM, Rogers N, O'Sullivan E, Kearins O, Rachet B. Ethnicity, deprivation and
35 screening: survival from breast cancer among screening-eligible women in the West Midlands
36 diagnosed from 1989 to 2011. *Br J Cancer.* 2015;113(3):548-55.
37
38 48. Luque-Fernandez MA, Redondo-Sanchez D, Rodriguez-Barranco M, Chang-Chan YL,
39 Salamanca-Fernandez E, Nunez O, et al. Socioeconomic Inequalities in Colorectal Cancer Survival in
40 Southern Spain: A Multilevel Population-Based Cohort Study. *Clin Epidemiol.* 2020;12:797-806.
41
42 49. Nunez O, Rodriguez Barranco M, Fernandez-Navarro P, Redondo Sanchez D, Luque
43 Fernandez MA, Pollan Santamaria M, et al. Deprivation gap in colorectal cancer survival attributable
44 to stage at diagnosis: A population-based study in Spain. *Cancer Epidemiol.* 2020;68:101794.
45
46 50. Wan YI, McGuckin D, Fowler AJ, Prowle JR, Pearse RM, Moonesinghe SR. Socioeconomic
47 deprivation and long-term outcomes after elective surgery: analysis of prospective data from two
48 observational studies. *Br J Anaesth.* 2021;126(3):642-51.
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 51. Zhou K, Shi H, Chen R, Cochuyt JJ, Hodge DO, Manochakian R, et al. Association of Race,
4 Socioeconomic Factors, and Treatment Characteristics With Overall Survival in Patients With
5 Limited-Stage Small Cell Lung Cancer. *JAMA Netw Open*. 2021;4(1):e2032276.
6
7
8 52. Zhang YB, Chen C, Pan XF, Guo J, Li Y, Franco OH, et al. Associations of healthy lifestyle and
9 socioeconomic status with mortality and incident cardiovascular disease: two prospective cohort
10 studies. *BMJ*. 2021;373:n604.
11
12
13 53. Lindenauer PK, Lagu T, Rothberg MB, Avrunin J, Pekow PS, Wang Y, et al. Income inequality
14 and 30 day outcomes after acute myocardial infarction, heart failure, and pneumonia: retrospective
15 cohort study. *BMJ*. 2013;346:f521.
16
17
18 54. King AJ, Hudson J, Fernie G, Burr J, Azuara-Blanco A, Sparrow JM, et al. Baseline
19 Characteristics of Participants in the Treatment of Advanced Glaucoma Study: A Multicenter
20 Randomized Controlled Trial. *Am J Ophthalmol*. 2020;213:186-94.
21
22
23 55. McKean-Cowdin R, Varma R, Wu J, Hays RD, Azen SP, Los Angeles Latino Eye Study G.
24 Severity of visual field loss and health-related quality of life. *Am J Ophthalmol*. 2007;143(6):1013-
25 23.
26
27
28 56. Patel D BH, Murdoch I. Barriers to uptake of eye careservices by the Indian population living
29 in Ealing, WestLondon. . *Health Educ J* 2006;65(3):267-76.
30
31
32 57. Shickle D, Farragher TM. Geographical inequalities in uptake of NHS-funded eye
33 examinations: small area analysis of Leeds, UK. *J Public Health (Oxf)*. 2015;37(2):337-45.
34
35
36 58. Baker H, Cousens SN, Murdoch IE. Poor public health knowledge about glaucoma: fact or
37 fiction? *Eye (Lond)*. 2010;24(4):653-7.
38
39
40 59. Baker H, Murdoch IE. Can a public health package on glaucoma reach its target population?
41 *Eye (Lond)*. 2004;18(5):478-82.
42
43
44 60. Baker H, Murdoch IE. Can a public health intervention improve awareness and healthseeking
45 behaviour for glaucoma? *Br J Ophthalmol*. 2008;92(12):1671-5.
46
47
48 61. Prior M, Francis JJ, Azuara-Blanco A, Anand N, Burr JM, Glaucoma screening Platform Study
49 g. Why do people present late with advanced glaucoma? A qualitative interview study. *Br J*
50 *Ophthalmol*. 2013;97(12):1574-8.
51
52
53 62. Lascaratos G, Garway-Heath DF, Burton R, Bunce C, Xing W, Crabb DP, et al. The United
54 Kingdom Glaucoma Treatment Study: a multicenter, randomized, double-masked, placebocontrolled
55 trial: baseline characteristics. *Ophthalmology*. 2013;120(12):2540-5.
56
57
58 63. Musch DC, Lichter PR, Guire KE, Standardi CL. The Collaborative Initial Glaucoma Treatment
59 Study: study design, methods, and baseline characteristics of enrolled patients. *Ophthalmology*.
60

1
2
3 1999;106(4):653-62.
4

5 64. Konstantakopoulou E, Gazzard G, Vickerstaff V, Jiang Y, Nathwani N, Hunter R, et al. The
6 Laser in Glaucoma and Ocular Hypertension (LiGHT) trial. A multicentre randomised controlled trial:
7 baseline patient characteristics. *Br J Ophthalmol*. 2018;102(5):599-603.
8

9 65. Davis SA, Sleath B, Carpenter DM, Blalock SJ, Muir KW, Budenz DL. Drop instillation and
10 glaucoma. *Curr Opin Ophthalmol*. 2018;29(2):171-7.
11

12 66. Newman-Casey PA, Niziol LM, Gillespie BW, Janz NK, Lichter PR, Musch DC. The Association
13 between Medication Adherence and Visual Field Progression in the Collaborative Initial Glaucoma
14 Treatment Study. *Ophthalmology*. 2020;127(4):477-83.
15

16 67. Leung VC, Jin YP, Hatch W, Mammo Z, Trope GE, Buys YM, et al. The relationship between
17 sociodemographic factors and persistence with topical glaucoma medications. *J Glaucoma*.
18 2015;24(1):69-76.
19
20
21
22
23
24
25
26
27
28
29

30 **Funding**

31
32
33 The project was funded by the NIHR Health Technology Assessment Programme
34 (Project Number 12/35/38). The Health Services Research Unit are funded by the
35 Chief Scientist Office of the Scottish Government Health and Social Care Directorates.
36
37 The funders had no role in considering the study design or in the collection, analysis,
38 interpretation of data, writing the report, or decision to submit the article for publication.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Confidential: For Review Only

Table 1. Baseline characteristics by IMD for overall cohort

	IMD 1 N=106	IMD 2 N=67	IMD 3 N=88	IMD 4 N=93	IMD 5 N=96
Age - mean (SD); n	62.3 (14.4); 106	66.0 (12.3); 67	68.0 (11.5); 88	69.5 (10.8); 93	70.1 (10.6); 96
Gender - n (%)					
Male	77 (72.6)	49 (73.1)	56 (63.6)	58 (62.4)	60 (62.5)
Female	29 (27.4)	18 (26.9)	32 (36.4)	35 (37.6)	36 (37.5)
Ethnicity - n (%)					
Caucasian	61 (57.5)	50 (74.6)	75 (85.2)	90 (96.8)	94 (97.9)
Afro-Caribbean	36 (34.0)	12 (17.9)	8 (9.1)	3 (3.2)	0 (0)
Asian - India/Pakistan/Bangladesh	5 (4.7)	3 (4.5)	3 (3.4)	0 (0)	1 (1.0)
Asian - Oriental	0 (0)	1 (1.5)	1 (1.1)	0 (0)	0 (0)
Mixed heritage	0 (0)	0 (0)	1 (1.1)	0 (0)	0 (0)
Other	3 (2.8)	1 (1.5)	0 (0)	0 (0)	1 (1.0)
Missing	1 (0.9)	0 (0)	0 (0)	0 (0)	0 (0)
Advanced glaucoma in both eyes - n (%)					
No	88 (83.0)	45 (67.2)	64 (72.7)	78 (83.9)	87 (90.6)
Yes	18 (17.0)	22 (32.8)	24 (27.3)	15 (16.1)	9 (9.4)
Glaucoma in both eyes – n (%)					
No	20 (18.9)	13 (19.4)	19 (21.6)	26 (28.0)	26 (27.1)
Yes	86 (81.1)	54 (80.6)	69 (78.4)	67 (72.0)	70 (72.9)
Eligible to be registered as sight impaired - n (%)					
No	97 (91.5)	57 (85.1)	84 (95.5)	92 (98.9)	93 (96.9)

Sight impaired	8 (7.5)	9 (13.4)	2 (2.3)	1 (1.1)	2 (2.1)
Severe sight impaired	1 (0.9)	1 (1.5)	2 (2.3)	0 (0)	1 (1.0)
Glaucoma diagnosis - n (%)					
Primary open angle glaucoma (including NTG)	105 (99.1)	61 (91.0)	85 (96.6)	92 (98.9)	93 (96.9)
Pigment dispersion syndrome	0 (0)	2 (3.0)	3 (3.4)	1 (1.1)	3 (3.1)
Psuedoexfoliation syndrome	1 (0.9)	4 (6.0)	0 (0)	0 (0)	0 (0)
Family history of glaucoma - n (%)					
Yes	35 (33.0)	12 (17.9)	31 (35.2)	29 (31.2)	34 (35.4)
No	67 (63.2)	48 (71.6)	54 (61.4)	58 (62.4)	55 (57.3)
Missing	4 (3.8)	7 (10.4)	3 (3.4)	6 (6.5)	7 (7.3)
Number of times visited the optician in the last 10 year – median [IQR]; n	4 [2, 5]; 98	5 [2.5, 10]; 60	4 [2, 6]; 86	5 [3, 8]; 85	5 [4, 7]; 91
NEI-VFQ-25 - mean (SD); n	83.9 (15.8); 104	81.5 (16.2); 66	89.6 (11.5); 88	89.5 (11.2); 93	89.5 (10.7); 96
NEI-VFQ-25 subscales - mean (SD); n					
Near activities	80.6 (21.8); 103	79.5 (17.6); 66	86.7 (16.8); 88	87.9 (13.8); 93	85.6 (16.0); 96
Distance activities	87.6 (18.2); 104	83.1 (17.6); 66	91.7 (12.9); 88	90.3 (13.9); 93	91.0 (12.3); 96
Dependency	90.7 (21.2); 104	91.3 (21.7); 65	97.0 (10.7); 87	96.0 (12.6); 93	96.8 (13.6); 96
Driving	82.6 (32.6); 65	71.2 (37.4); 37	89.8 (18.5); 70	87.7 (24.1); 73	88.1 (20.8); 81
General health	57.3 (25.6); 103	60.4 (22.5); 65	63.4 (21.1); 88	62.1 (22.9); 93	67.7 (21.4); 96
Role difficulties	83.1 (22.2); 104	80.6 (24.7); 65	89.7 (16.1); 87	90.2 (19.3); 93	91.0 (17.5); 96

Mental health	75.0 (24.4); 104	73.2 (25.1); 66	84.9 (14.9); 88	86.0 (16.6); 93	86.1 (17.3); 96
General vision	69.9 (15.7); 101	69.7 (14.6); 66	76.6 (14.0); 88	77.4 (12.9); 93	74.7 (13.1); 95
Social function	94.1 (13.5); 104	90.5 (15.4); 66	95.9 (12.1); 88	96.7 (9.1); 92	96.7 (9.1); 96
Colour vision	94.5 (14.8); 104	94.9 (11.9); 64	98.3 (10.0); 86	97.8 (8.0); 92	97.9 (8.6); 96
Peripheral vision	85.1 (22.2); 104	80.8 (24.5); 65	89.2 (19.3); 88	88.6 (19.0); 92	88.8 (17.3); 96
Ocular pain	79.7 (21.4); 103	80.3 (19.6); 66	88.1 (16.0); 88	86.0 (16.7); 93	86.8 (15.0); 96
EQ-5D-5L - mean (SD); n	0.827 (0.219); 101	0.820 (0.157); 65	0.862 (0.171); 88	0.838 (0.168); 92	0.852 (0.173); 95
HUI-3 - mean (SD); n	0.803 (0.248); 98	0.789 (0.187); 61	0.830 (0.200); 84	0.808 (0.202); 90	0.820 (0.175); 92
GUI - mean (SD); n	0.883 (0.148); 101	0.857 (0.124); 63	0.902 (0.114); 86	0.899 (0.119); 93	0.905 (0.103); 96
Participant experience (glaucoma getting worse) - n (%)					
Yes	42 (39.6)	31 (46.3)	27 (30.7)	35 (37.6)	36 (37.5)
No	56 (52.8)	30 (44.8)	53 (60.2)	51 (54.8)	54 (56.3)
Missing	8 (7.5)	6 (9.0)	8 (9.1)	7 (7.5)	6 (6.3)
Visual standards for driving - n (%)					
Pass	89 (84.0)	50 (74.6)	73 (83.0)	81 (87.1)	87 (90.6)
Fail	12 (11.3)	14 (20.9)	12 (13.6)	5 (5.4)	5 (5.2)
Missing	5 (4.7)	3 (4.5)	3 (3.4)	7 (7.5)	4 (4.2)
Diamox ^A - n (%)	5 (4.7)	1 (1.5)	2 (2.3)	0 (0)	0 (0)

^Ataken orally. SD standard deviation

Table 2. Baseline clinical characteristics for index eye for overall cohort

	IMD 1 N=106	IMD 2 N=67	IMD 3 N=88	IMD 4 N=93	IMD 5 N=96
Lens status - n (%)					
Phakic	100 (94.3)	64 (95.5)	82 (93.2)	84 (90.3)	88 (91.7)
Pseudophakic	6 (5.7)	3 (4.5)	6 (6.8)	9 (9.7)	8 (8.3)
Central corneal thickness - - mean (SD); n	541.7 (38.5); 105	543.1 (33.5); 67	540.9 (36.3); 86	537.9 (33.9); 93	539.3 (35.9); 95
Glaucoma drops - n (%)					
Pg analogue	89 (84.0)	54 (80.6)	73 (83.0)	72 (77.4)	77 (80.2)
β-blocker	31 (29.2)	26 (38.8)	21 (23.9)	9 (9.7)	17 (17.7)
CA inhibitor	23 (21.7)	20 (29.9)	16 (18.2)	9 (9.7)	10 (10.4)
Agonist	6 (5.7)	2 (3.0)	0 (0)	2 (2.2)	1 (1.0)
Ocular co-morbidity - n (%)					
Yes	17 (16.0)	13 (19.4)	23 (26.1)	23 (24.7)	24 (25.0)
No	89 (84.0)	54 (80.6)	65 (73.9)	70 (75.3)	72 (75.0)
Ocular co-morbidity details ^A					
AMD	1 (5.9)	2 (15.4)	4 (17.4)	1 (4.3)	2 (8.3)
Cataract	12 (70.6)	11 (84.6)	19 (82.6)	21 (91.3)	21 (87.5)
Vascular occlusion	1 (5.9)	0 (0)	1 (4.3)	1 (4.3)	0 (0)
Diabetic retinopathy	2 (11.8)	0 (0)	0 (0)	0 (0)	0 (0)
Other	1 (5.9)	2 (15.4)	2 (8.7)	6 (26.1)	4 (16.7)
LogMAR visual acuity - mean (SD); n	0.2 (0.4); 105	0.1 (0.2); 66	0.1 (0.2); 88	0.1 (0.2); 92	0.2 (0.2); 96

Visual fields mean deviation (dB) - mean (SD); n	-17.2 (6.7); 106	-16.9 (6.6); 67	-15.3 (5.4); 88	-13.5 (6.4); 93	-13.0 (5.5); 96
Intraocular pressure (mmHg) - mean (SD); n					
Diagnosis	28.2 (10.6); 105	27.1 (9.0); 66	26.6 (9.0); 88	25.8 (7.2); 92	24.5 (7.1); 95
Baseline	19.0 (5.8); 105	19.8 (6.5); 67	19.2 (5.3); 86	20.1 (6.6); 88	18.4 (5.4); 94

^Aparticipants can have more than one. SD standard deviation

Table 3 - Index of Multiple Deprivation correlations for overall cohort

	Index of Multiple Deprivation	
	Pearson's correlation	p-value
Gender		0.30
Ethnicity		<0.001
Family history of glaucoma		0.19
Age	0.23	<0.001
Number of times visited the optician in the last 10 year	0.10	0.028
NEI-VFQ-25	0.19	<0.001
EQ-5D-5L	0.05	0.29
HUI-3	0.03	0.48
GUI	0.09	0.06
LogMAR visual acuity for index eye	-0.12	0.014
Log MAR visual acuity for non-index eye	-0.02	0.75
Log MAR f visual acuity or both eyes combined	0.02	0.69
Visual fields mean deviation (dB) for index eye	0.27	<0.001
Visual fields mean deviation (dB) for non-index eye	0.23	<0.001
Intraocular pressure (mmHg) for index eye	-0.02	0.64

Chi-squared test for association was used for dichotomous variables and Pearson's correlation with p-value for continuous

Table 4 - difference in the outcome by IMD for participants randomised to trabeculectomy and medical treatment

Trabeculectomy		Baseline	24 months	Interaction	95% CI	p-value
Glaucoma drops						
IMD 1		1.41 (0.94); 54	0.55 (0.94); 51			
IMD 2		1.50 (1.25); 30	0.73 (1.12); 26	0.15	(-0.27, 0.58)	0.480
IMD 3		1.24 (0.93); 45	0.41 (0.87); 41	-0.11	(-0.49, 0.26)	0.546
IMD 4		1.04 (0.81); 50	0.33 (0.94); 46	-0.18	(-0.55, 0.18)	0.319
IMD 5		1.28 (0.85); 47	0.43 (0.78); 46	-0.09	(-0.45, 0.27)	0.618
IOP (mmHg)						
IMD 1		18.94 (6.27); 51	13.45 (5.57); 50			
IMD 2		20.69 (7.94); 26	12.24 (4.09); 25	-1.39	(-3.61, 0.83)	0.220
IMD 3		19.65 (5.75); 41	11.02 (3.76); 39	-2.50	(-4.43, -0.57)	0.011
IMD 4		20.33 (5.85); 46	12.38 (4.67); 46	-1.19	(-3.04, 0.66)	0.207
IMD 5		18.16 (5.72); 46	12.61 (4.69); 45	-0.78	(-2.64, 1.07)	0.409
VF MD (dB)						
IMD 1		-16.11 (6.16); 51	-16.11 (6.66); 50			
IMD 2		-19.00 (6.44); 26	-18.68 (6.02); 24	-0.54	(-1.97, 0.88)	0.454
IMD 3		-15.67 (5.43); 41	-15.72 (5.54); 39	0.06	(-1.16, 1.28)	0.918
IMD 4		-12.80 (6.68); 46	-13.31 (7.05); 44	0.01	(-1.18, 1.21)	0.982
IMD 5		-12.53 (5.55); 46	-13.53 (6.60); 44	-0.79	(-2.00, 0.41)	0.196
LogMAR visual acuity						
IMD 1		0.20 (0.35); 51	0.22 (0.31); 49			
IMD 2		0.14 (0.15); 26	0.21 (0.32); 24	0.02	(-0.09, 0.14)	0.687
IMD 3		0.15 (0.26); 41	0.22 (0.29); 38	0.02	(-0.09, 0.12)	0.730
IMD 4		0.07 (0.16); 46	0.23 (0.29); 45	0.07	(-0.03, 0.17)	0.153
IMD 5		0.13 (0.19); 46	0.19 (0.19); 42	0.02	(-0.08, 0.12)	0.743
NEI-VFQ-25						
IMD 1		82.39 (16.00); 50	81.91 (15.82); 50			
IMD 2		82.82 (16.72); 26	80.41 (15.32); 26	-1.78	(-6.54, 2.98)	0.463

1						
2						
3						
4	IMD 3	88.56 (13.36);	83.71 (16.52);	-2.30	(-6.52, 1.93)	0.287
5		40	40			
6	IMD 4	91.10 (9.33); 47	88.93 (10.64);	1.21	(-2.92, 5.33)	0.566
7			46			
8						
9	IMD 5	90.83 (10.78);	90.20 (7.85); 44	2.67	(-1.49, 6.84)	0.208
10		44				
11						
12	EQ-5D-5L					
13	IMD 1	0.81 (0.24); 50	0.78 (0.21); 50			
14	IMD 2	0.83 (0.18); 26	0.80 (0.17); 25	0.02	(-0.05, 0.08)	0.582
15	IMD 3	0.85 (0.18); 40	0.81 (0.22); 40	0.01	(-0.05, 0.07)	0.761
16	IMD 4	0.87 (0.13); 47	0.84 (0.14); 46	0.03	(-0.02, 0.09)	0.252
17	IMD 5	0.87 (0.18); 44	0.82 (0.15); 44	0.01	(-0.05, 0.06)	0.784
18						
19	HUI-3					
20	IMD 1	0.79 (0.24); 50	0.76 (0.26); 46			
21	IMD 2	0.79 (0.21); 26	0.76 (0.27); 24	0.00	(-0.08, 0.09)	0.928
22	IMD 3	0.80 (0.24); 40	0.75 (0.28); 38	-0.01	(-0.08, 0.07)	0.844
23	IMD 4	0.84 (0.15); 47	0.83 (0.17); 45	0.03	(-0.04, 0.11)	0.375
24	IMD 5	0.85 (0.13); 44	0.81 (0.17); 44	0.01	(-0.07, 0.08)	0.868
25						
26	GUI					
27	IMD 1	0.87 (0.15); 50	0.85 (0.17); 50			
28	IMD 2	0.88 (0.11); 26	0.81 (0.16); 26	-0.04	(-0.11, 0.02)	0.171
29	IMD 3	0.89 (0.14); 40	0.82 (0.19); 39	-0.04	(-0.10, 0.01)	0.142
30	IMD 4	0.92 (0.11); 47	0.91 (0.10); 46	0.03	(-0.02, 0.08)	0.277
31	IMD 5	0.93 (0.10); 44	0.89 (0.10); 43	0.02	(-0.04, 0.07)	0.543
32						
33						
34						
35						
36						
37						
38	Medical					
39	Management					
40						
41		Baseline	24 months	Interaction	95% CI	p-value
42						
43	Glaucoma drops					
44	IMD 1	1.40 (1.16); 52	1.68 (1.22); 44			
45	IMD 2	1.54 (0.77); 37	2.00 (1.10); 34	0.29	(-0.22, 0.80)	0.261
46	IMD 3	1.26 (0.76); 43	1.40 (1.15); 40	-0.27	(-0.76, 0.21)	0.274
47	IMD 4	0.93 (0.70); 43	1.71 (1.13); 42	0.07	(-0.41, 0.56)	0.773
48	IMD 5	0.92 (0.61); 49	1.46 (1.13); 46	-0.19	(-0.66, 0.29)	0.440
49						
50	IOP (mmHg)					
51	IMD 1	18.43 (4.17); 44	15.32 (3.32); 43			
52	IMD 2	19.29 (5.62); 34	15.92 (7.91); 34	0.46	(-1.65, 2.57)	0.672
53	IMD 3	19.04 (4.95); 40	14.78 (3.84); 38	-0.65	(-2.69, 1.40)	0.537
54	IMD 4	20.18 (7.53); 42	14.85 (4.56); 40	-0.71	(-2.74, 1.31)	0.490
55	IMD 5	18.30 (5.18); 46	14.72 (3.96); 45	-0.60	(-2.55, 1.36)	0.551
56						
57						
58						
59						
60						

1						
2						
3						
4	VF MD (dB)					
5	IMD 1	-18.56 (6.87);	-17.26 (6.85);			
6		44	41			
7						
8	IMD 2	-15.56 (6.05);	-16.72 (6.06);	-1.45	(-2.99, 0.10)	0.066
9		34	34			
10						
11	IMD 3	-14.49 (5.61);	-14.96 (5.97);	-1.01	(-2.53, 0.50)	0.190
12		40	38			
13						
14	IMD 4	-13.99 (6.13);	-14.47 (7.20);	-0.64	(-2.14, 0.87)	0.406
15		42	40			
16						
17	IMD 5	-12.94 (5.48);	-13.99 (5.55);	-1.05	(-2.53, 0.43)	0.164
18		46	45			
19						
20	LogMAR visual					
21	acuity					
22	IMD 1	0.24 (0.39); 44	0.22 (0.38); 42			
23						
24	IMD 2	0.13 (0.22); 34	0.14 (0.19); 33	-0.02	(-0.10, 0.06)	0.639
25						
26	IMD 3	0.14 (0.21); 40	0.14 (0.17); 38	-0.01	(-0.09, 0.07)	0.770
27						
28	IMD 4	0.16 (0.18); 42	0.18 (0.27); 41	-0.01	(-0.08, 0.07)	0.867
29						
30	IMD 5	0.17 (0.20); 46	0.13 (0.20); 45	-0.06	(-0.13, 0.01)	0.104
31						
32	NEI-VFQ-25					
33	IMD 1	86.21 (15.23);	86.25 (15.36);			
34		42	42			
35						
36	IMD 2	81.00 (16.09);	77.40 (19.71);	-4.15	(-9.26, 0.95)	0.111
37		34	34			
38						
39	IMD 3	90.80 (10.08);	86.47 (13.26);	-3.92	(-8.83, 1.00)	0.118
40		39	39			
41						
42	IMD 4	88.82 (11.10);	82.55 (19.03);	-6.05	(-10.85, -	0.014
43		42	42		1.24)	
44						
45	IMD 5	89.52 (10.18);	87.66 (12.66);	-1.57	(-6.27, 3.13)	0.513
46		46	46			
47						
48	EQ-5D-5L					
49	IMD 1	0.83 (0.20); 42	0.80 (0.23); 41			
50						
51	IMD 2	0.83 (0.12); 34	0.77 (0.18); 33	-0.01	(-0.08, 0.06)	0.819
52						
53	IMD 3	0.89 (0.14); 39	0.84 (0.12); 39	0.02	(-0.05, 0.09)	0.508
54						
55	IMD 4	0.83 (0.16); 42	0.74 (0.22); 42	-0.04	(-0.11, 0.03)	0.218
56						
57	IMD 5	0.85 (0.17); 46	0.83 (0.17); 46	0.03	(-0.04, 0.10)	0.387
58						
59	HUI-3					
60	IMD 1	0.81 (0.25); 42	0.76 (0.29); 38			
	IMD 2	0.80 (0.17); 34	0.74 (0.24); 33	0.01	(-0.07, 0.09)	0.806
	IMD 3	0.88 (0.10); 39	0.79 (0.19); 39	-0.00	(-0.08, 0.08)	0.960
	IMD 4	0.78 (0.23); 42	0.67 (0.29); 39	-0.03	(-0.11, 0.05)	0.480

	IMD 5	0.81 (0.19); 46	0.79 (0.21); 42	0.05	(-0.03, 0.12)	0.254
GUI						
	IMD 1	0.90 (0.12); 42	0.89 (0.15); 40			
	IMD 2	0.85 (0.11); 34	0.78 (0.19); 34	-0.06	(-0.12, 0.01)	0.086
	IMD 3	0.91 (0.09); 39	0.86 (0.14); 39	-0.03	(-0.09, 0.03)	0.318
	IMD 4	0.88 (0.12); 42	0.82 (0.23); 42	-0.05	(-0.11, 0.01)	0.114
	IMD 5	0.89 (0.10); 46	0.87 (0.14); 45	-0.01	(-0.07, 0.05)	0.692

Values are mean (standard deviation); n. CI Confidence Interval

Figure Legends

Figure 1: Subgroup for Trabeculectomy versus Medical management: (a) IOP; (b) VF MD; (c) logMAR visual acuity

Supplementary Figure 1 – Subgroup for Trabeculectomy versus Medical management: (d) NEI-VFQ-25; (e) EQ-5D-5L; (f) HUI-3; (g) GUI. 1st quintile – most deprived, 5th quintile -least deprived




Figure 1: Subgroup for Trabeculectomy versus Medical management: (a) IOP; (b) VF MD; (c) logMAR visual acuity

90x194mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Supplementary Table 1. Index Multiple Deprivation (IMD)

	Trabeculectomy N=227	Medical management N=226
1 st quintile (most deprived) (IMD 1)	54 (23.8)	52 (23.0)
2 nd quintile (IMD 2)	30 (13.2)	37 (16.4)
3 rd quintile (IMD 3)	45 (19.8)	43 (19.0)
4 th quintile (IMD 4)	50 (22.0)	43 (19.0)
5 th quintile (least deprived) (IMD 5)	47 (20.7)	49 (21.7)
Missing	1 (0.4)	2 (0.9)

Values are numbers (percent).

Confidential: For Review Only

Supplementary Table 2 - Baseline clinical characteristics for non-index eye for overall cohort by IMD

	IMD 1 N=106	IMD 2 N=67	IMD 3 N=88	IMD 4 N=93	IMD 5 N=96
Lens status - n (%)					
Phakic	99 (93.4)	61 (91.0)	84 (95.5)	83 (89.2)	88 (91.7)
Pseudophakic	6 (5.7)	6 (9.0)	4 (4.5)	10 (10.8)	8 (8.3)
Missing	1 (0.9)	0 (0)	0 (0)	0 (0)	0 (0)
Central corneal thickness - n; mean (SD)	541.2 (40.9); 104	543.9 (32.5); 67	541.6 (37.9); 86	539.5 (34.7); 93	539.9 (36.9); 95
Glaucoma drops - n (%)					
Pg Analogue	80 (75.5)	46 (68.7)	71 (80.7)	57 (61.3)	64 (66.7)
β -blocker	25 (23.6)	20 (29.9)	18 (20.5)	6 (6.5)	13 (13.5)
CA Inhibitor	13 (12.3)	13 (19.4)	13 (14.8)	7 (7.5)	8 (8.3)
α - Agonist	4 (3.8)	1 (1.5)	0 (0)	1 (1.1)	1 (1.0)
Ocular co-morbidity - n (%)					
Yes	16 (15.1)	14 (20.9)	24 (27.3)	23 (24.7)	21 (21.9)
No	90 (84.9)	53 (79.1)	64 (72.7)	70 (75.3)	75 (78.1)
Ocular co-morbidity details ¹					
AMD	1 (5.9)	2 (15.4)	4 (17.4)	1 (4.3)	2 (8.3)
Cataract	11 (64.7)	11 (84.6)	17 (73.9)	17 (73.9)	19 (79.2)
Diabetic retinopathy	2 (11.8)	0 (0)	0 (0)	0 (0)	0 (0)
Other	1 (5.9)	1 (7.7)	5 (21.7)	7 (30.4)	2 (8.3)
LogMAR visual acuity - mean (SD); n	0.1 (0.2); 105	0.1 (0.3); 65	0.1 (0.3); 87	0.1 (0.2); 92	0.1 (0.2); 96
Visual fields mean deviation (dB) - mean (SD); n	-7.4 (8.2); 104	-9.2 (9.3); 67	-6.7 (7.7); 88	-4.4 (5.6); 93	-3.9 (4.9); 96
Intraocular pressure (mmHg) - mean (SD); n					
Diagnosis	24.1 (7.9); 104	24.2 (8.9); 66	23.3 (6.8); 88	22.3 (6.0); 92	21.0 (5.0); 95
Baseline	18.0 (4.7); 104	18.6 (6.0); 67	17.8 (4.4); 86	18.1 (4.1); 88	17.4 (4.2); 94


¹participants can have more than one. SD standard deviation

Supplementary Table 3 - difference in the outcome by IMD for the overall cohort

	Baseline	24 months	Interaction	95% CI	p-value
Glaucoma drops					
IMD 1	1.41 (1.05); 106	1.07 (1.21); 95			
IMD 2	1.52 (1.01); 67	1.45 (1.27); 60	0.36	(-0.02, 0.75)	0.063
IMD 3	1.25 (0.85); 88	0.90 (1.12); 81	-0.16	(-0.51, 0.19)	0.357
IMD 4	0.99 (0.76); 93	0.99 (1.25); 88	-0.07	(-0.41, 0.28)	0.712
IMD 5	1.09 (0.76); 96	0.95 (1.09); 92	-0.11	(-0.45, 0.23)	0.519
IOP (mmHg)					
IMD 1	18.70 (5.38); 95	14.31 (4.73); 93			
IMD 2	19.90 (6.70); 60	14.36 (6.78); 59	-0.10	(-1.69, 1.49)	0.902
IMD 3	19.35 (5.35); 81	12.87 (4.22); 77	-1.52	(-2.99, -0.05)	0.043
IMD 4	20.26 (6.67); 88	13.53 (4.76); 86	-0.94	(-2.37, 0.49)	0.197
IMD 5	18.23 (5.42); 92	13.66 (4.44); 90	-0.61	(-2.02, 0.80)	0.398
VF MD (dB)					
IMD 1	-17.24 (6.58); 95	-16.63 (6.73); 91			
IMD 2	-17.05 (6.40); 60	-17.53 (6.07); 58	-1.06	(-2.10, -0.02)	0.045
IMD 3	-15.09 (5.52); 81	-15.34 (5.73); 77	-0.48	(-1.44, 0.49)	0.331
IMD 4	-13.37 (6.41); 88	-13.87 (7.10); 84	-0.28	(-1.23, 0.67)	0.564
IMD 5	-12.74 (5.49); 92	-13.77 (6.06); 89	-0.90	(-1.85, 0.05)	0.062
LogMAR visual acuity					
IMD 1	0.21 (0.37); 95	0.22 (0.34); 91			
IMD 2	0.13 (0.19); 60	0.17 (0.25); 57	-0.00	(-0.08, 0.07)	0.911
IMD 3	0.14 (0.23); 81	0.18 (0.24); 76	-0.00	(-0.07, 0.07)	0.972
IMD 4	0.11 (0.17); 88	0.20 (0.28); 86	0.04	(-0.03, 0.10)	0.241
IMD 5	0.15 (0.19); 92	0.16 (0.20); 87	-0.02	(-0.09, 0.04)	0.497
NEI-VFQ-25					
IMD 1	84.13 (15.69); 92	83.89 (15.68); 92			
IMD 2	81.79 (16.25); 60	78.70 (17.86); 60	-3.35	(-6.87, 0.17)	0.062
IMD 3	89.66 (11.83); 79	85.08 (14.97); 79	-3.15	(-6.43, 0.13)	0.060
IMD 4	90.02 (10.21); 89	85.89 (15.48); 88	-2.62	(-5.82, 0.57)	0.108
IMD 5	90.16 (10.44); 90	88.90 (10.60); 90	0.29	(-2.89, 3.47)	0.860
EQ-5D-5L					
IMD 1	0.82 (0.22); 92	0.78 (0.22); 91			
IMD 2	0.83 (0.14); 60	0.79 (0.17); 58	0.00	(-0.05, 0.05)	0.926
IMD 3	0.87 (0.16); 79	0.82 (0.18); 79	0.02	(-0.03, 0.06)	0.489
IMD 4	0.85 (0.15); 89	0.79 (0.19); 88	-0.00	(-0.05, 0.04)	0.828
IMD 5	0.86 (0.17); 90	0.82 (0.16); 90	0.02	(-0.03, 0.06)	0.420
HUI-3					
IMD 1	0.80 (0.24); 92	0.76 (0.27); 84			
IMD 2	0.79 (0.18); 60	0.75 (0.25); 57	0.00	(-0.06, 0.06)	0.992
IMD 3	0.84 (0.19); 79	0.77 (0.24); 77	-0.01	(-0.06, 0.05)	0.857
IMD 4	0.81 (0.19); 89	0.76 (0.24); 84	-0.00	(-0.06, 0.05)	0.944
IMD 5	0.83 (0.16); 90	0.80 (0.19); 86	0.02	(-0.03, 0.07)	0.471
GUI					
IMD 1	0.88 (0.14); 92	0.87 (0.17); 90			
IMD 2	0.86 (0.11); 60	0.79 (0.18); 60	-0.06	(-0.11, -0.01)	0.011
IMD 3	0.90 (0.12); 79	0.84 (0.17); 78	-0.04	(-0.08, 0.00)	0.079
IMD 4	0.90 (0.11); 89	0.86 (0.18); 88	-0.02	(-0.06, 0.02)	0.422
IMD 5	0.91 (0.10); 90	0.88 (0.12); 88	-0.00	(-0.05, 0.04)	0.823

Values are mean (standard deviation); n. CI Confidence Interval

Supplementary Figure 1 – Subgroup for Trabeculectomy versus Medical management: (d) NEI-VFQ-25; (e) EQ-5D-5L; (f) HUI-3; (g) GUI. 1st quintile – most deprived, 5th quintile -least deprived



90x67mm (300 x 300 DPI)