

Height, social position, and coronary heart disease incidence: the contribution of genetic and environmental factors

Karri Silventoinen*^{1,2}, Hannu Lahtinen¹, George Davey Smith^{3,4}, Tim Morris^{3,4}, Pekka Martikainen^{1,5,6}

1. University of Helsinki, Faculty of Social Sciences, Population Research Unit, Finland
2. University of Helsinki, Faculty of Medicine, Department of Public Health, Finland
3. Bristol Medical School, University of Bristol, Population Health Sciences, UK
4. MRC Integrative Epidemiology Unit at the University of Bristol, Bristol, UK
5. Stockholm University, Centre for Health Equity Studies, Sweden
6. Max-Planck-Institute for Demographic Research, Germany

Running head: Height, CHD and social position

Word count: 250 (abstract) + 2998 (main text)

*Correspondence address:

Karri Silventoinen, PhD

University of Helsinki, Population Research Unit, Faculty of Social Sciences

P.O. Box 18, FIN-00014 University of Helsinki, Finland

GSM: +358400-620726

E-mail: karri.silventoinen@helsinki.fi

Abstract

Background: The associations between height, socioeconomic position (SEP) and coronary heart disease (CHD) incidence are well established, but the contribution of genetic factors to these associations is still poorly understood. We used a polygenic score (PGS) for height to shed light on these associations.

Methods: Finnish population-based health surveys in 1992–2011 (response rates 65–93%) were linked to population registers providing information on SEP and CHD incidence up to 2019. The participants (N=29,996; 54% women) were aged 25–75 at baseline, and there were 1,767 CHD incident cases (32% in women) during 472,973 person years of follow-up. PGS-height was calculated based on 33,938 single nucleotide polymorphisms, and residual height was defined as the residual of height after adjusting for PGS-height in a linear regression model. Hazard ratios (HR) of CHD incidence were calculated using Cox regression.

Results: PGS-height and residual height showed clear gradients for education, social class and income, with a larger association for residual height. Residual height also showed larger associations with CHD incidence (HRs per 1 SD 0.94 in men and 0.87 in women) than PGS-height (HRs per 1 SD 0.97 and 0.99, respectively). Only a small proportion of the associations between SEP and CHD incidence was statistically explained by the height indicators (6% or less).

Conclusions: Residual height associations with SEP and CHD incidence were larger than for PGS-height. This supports the role of material and social living conditions in childhood as contributing factors to the association of height with both SEP and CHD risk.

What is already known on this topic

- Height is associated with social position and CHD incidence.

What this study adds

- Genetic propensity for height – as measured by a polygenic score for height – showed a robust social gradient but only a weak association with CHD incidence.

- Height residuals adjusted for the genetic propensity showed stronger associations with social position and CHD incidence than the genetic propensity for height.

How this study might affect research

- Our results of strong associations of residual height with social position and CHD incidence are consistent with the idea that height reflects living conditions during neonatal development, childhood and adolescence.

- Even though both PGS-height and residual height showed clear SEP gradients and residual height was also associated with CHD incidence, they only made a negligible contribution to social inequalities in CHD risk.

Introduction

The higher risk of coronary heart disease (CHD) in those of lower socioeconomic position (SEP) is well established in high income societies [1]. In addition to behavioral factors [2], material deprivation in childhood may affect CHD risk and thus contribute to social inequalities in CHD [3]. Since reliable direct measures of early life material deprivation are seldom available in epidemiological studies, an important part of this evidence is based on surrogate measures, such as height. At the population level, adult height strongly correlates with the standard of living and population nutrition [4]. In addition, those of lower SEP have, on average, shorter stature than those of higher SEP [5]. Short stature is also associated with increased CHD risk [6,7]. This suggests that height can capture environmental variation that is potentially important for the explanation of social inequalities in CHD. However, a limitation in many previous studies looking at height, SEP and CHD risk is that they ignore the role of genetic factors. Between 65 to 85% of population variation in adult height at a particular place and time can statistically be explained by genetic factors when estimated using twin designs [8], and thousands of genetic variants have been identified that associate with adult stature in genome-wide-association studies (GWAS) [9].

Height has thus attracted interest both in social research focusing on secular trends and SEP differences [5] and in genetic research focusing on heritability estimated by twin [8] and GWAS designs [9]. Even when these approaches may be seen as counter intuitive to each other, both of them can shed light to the association between height and CHD. Indeed, previous studies have shown that polygenic scores (PGS) of height associate with SEP [10] and CHD risk [11]. Figure 1 summarizes the underlying causal pathways behind the associations between height, SEP and CHD risk expected based on previous studies. Childhood environment can affect height [12], SEP [13] and CHD risk [3]. Genetic factors can affect height directly through, e.g., growth plate

chondrocytes of long bones [14] but also indirectly through influencing childhood nutrition, as indicated by genetic correlations between childhood height and skinfold thickness measures [15]. Genetic factors can also affect SEP [10] and CHD risk [11]. Finally, height can affect SEP [16] which can in turn affect CHD risk [17].

In this study, we summarized the genetic factors affecting height as the PGS of height (hereafter referred to as PGS-height). The childhood environment cannot, however, be estimated directly since it is not possible to measure all potential environmental exposures affecting height (18). Thus, we calculated the residuals of height adjusted for PGS-height (hereafter referred to as residual height), which is designed to index environmental variation of height. Residual height will also capture measurement error and some genetic influences which is not captured by PGS-height. Based on these assumptions, we made the following study hypotheses: i) Both PGS-height and residual height are associated with SEP and CHD risk. ii) These associations are stronger for residual than PGS-height indicating that environmental factors affecting growth make a major contribution to both SEP and increased CHD risk. iii) Adjusting for these height indicators will attenuate SEP inequalities in CHD risk, with greater attenuation resulting from adjustment for residual compared to PGS-height.

Data and methods

Several Finnish population-based health surveys – FINRISK surveys conducted in 1992, 1997, 2002, 2007 and 2012 and Health 2000/2011 surveys – were pooled together to create the baseline study cohort [19]. The response rates varied between 65% and 93%. Height was measured at the baseline health examinations, and at the same time the participants gave a blood sample used for genotyping and metabolic measures. PGS-height was based on the LASSO-weighted GWAS scores

from the GWAS of height in a study without sample overlap with our current study [20], which included 33,938 single nucleotide polymorphisms (SNP) selected using the p-value threshold of 0.05. Birth year and the square of birth year explained 8% of height variation in men and 9% in women. After these adjustments, the PGS-height explained an additional 29% of height variation both in men and women (r^2 totaling 0.37 and 0.38). Residual height was calculated from a regression model for height with PGS-height as the independent variable for men and women separately.

The baseline measurements were linked to Finnish population registers using unique personal identification codes that were pseudonymized before releasing data to the research team. We used education, occupation-based social class and income as the SEP indicators. Education and social class were derived from the Finnish Population Register. Education was based on the highest completed degree up to the end of 2019 and classified into four categories (basic, secondary, lower tertiary and higher tertiary education). Social class was measured at the age of 40 or if missing at the most recent previous measurement when the individual was employed and classified into five categories (manual workers, lower non-manual workers, upper non-manual workers, entrepreneurs and farmers) [21]. Income was based on personal taxable income from the Tax Register. We first calculated the yearly income percentiles among the 35–40 year old population for each year an individual belonged within this age group. Then, we took the mean of these percentile ranks and split them further into quintiles to also allow non-linear associations.

We restricted the participants to those born between 1935 and 1980 due to the availability of the measures of SEP indicators given the age restrictions. Thus, the participants were between 25 and 76 years of age at baseline of each data collection. Information for education and income was available at every five years between 1970 and 1985 and yearly between 1987 and 2019.

Information on social class was available at every five years between 1970 and 2005 and yearly between 2006 and 2018. Together, we had 36,418 participants in the selected birth cohorts, among whom 32,074 had genotype data available. However, 643 participants had missing information on height or on SEP indicators and were thus removed. Further, we removed randomly one individual from pairs with identity-by-descent (IBD) value ≥ 0.178 (N=1,435), corresponding to the expected lower bound of second-degree relatives. Thus, in the analyses on the association between height and the SEP indicators, we had 29,996 participants (54% women). In the main analyses, the height indicators were standardized (mean=0, standard deviation (SD)=1) in men and women separately.

The longitudinal information on CHD incidence cases was based on the Hospital Discharge Register for non-fatal (ICD-9 codes 410 or 4110 and ICD-10 codes I20.0 and I21–I22) and the National Mortality Register for fatal cases without previous hospitalization (ICD-9 codes 410–414 and 798, excluding 7980A and ICD-10 codes I20–I25, I46, R96 and R98) covering the whole Finnish population. When analyzing CHD incidence, we also used information on a number of risk factors of CHD. Body mass index (BMI), systolic and diastolic blood pressure, total cholesterol and high-density lipoprotein cholesterol were measured in the baseline clinical examination. Smoking classified as current, former and never smokers was asked in a self-administrated questionnaire at baseline. In these analyses, we removed additional 327 participants with a pre-baseline CHD event and 170 participants with missing information on any of the CHD risk factors. Thus, we had 29,499 participants for the CHD incidence analyses among whom there were 1,767 incident CHD cases (27% fatal cases and 32% in women) during the 472,973 person years until the end of follow-up on December 31st 2019. The distributions of participants and CHD incidence by the SEP indicators are presented in Supplementary table 1 and the means of the three height indicators in Supplementary table 2.

We first estimated the associations of SEP indicators (education, social class and income) with height indicators (height, PGS-height and residual height) using linear regression models. The results can be interpreted as the difference in height indicators between the SEP categories, but they do not imply causal associations between the variables. Second, we analyzed how these SEP and height indicators were associated with CHD incidence using Cox regression. Those who died from causes of death other than CHD during the follow-up were censored at the time of death. Cox proportional hazards assumptions appeared not to be violated when examined graphically (Kaplan-Meier curves available from the corresponding author). Additionally, we calculated population attributable fractions (PAF), which indicate the proportion of CHD cases that would be avoided had the higher risk SEP categories had the same CHD risk as the lowest risk SEP category. Comparing PAFs between two models would indicate how much the covariates added into the model would explain the SEP gradient.

All statistical models were adjusted for the first 10 principal components of genetic population structure, five geographic areas of residence, and the combination of baseline year and genotyping batch dummies. These adjustments help to minimize the impact of geographical differences in genetic structure, CHD incidence, height and SEP in the Finnish population. In the linear regression models, we adjusted for birth year and the square of birth year to account for the secular increase in height, change in social class structure and educational expansion in the Finnish population. In Cox regression models, the analyses were adjusted for age and squared age at baseline as well as baseline year dummies. The genetic principal components, IBD values and PGS-height were calculated using the PLINK 1.9 software. The statistical models were conducted using Stata, version 16.1.

Results

Tables 1 and 2 present the mean differences of height indicators in SD units between SEP categories calculated by linear regression models in men and women, respectively. Height, PGS-height and residual height were greatest in the uppermost SEP categories, i.e., in those with higher tertiary education, upper non-manual workers and the highest income quintile when adjusted for birth year and region indicators (Model 1). Generally, all height indicators showed SEP gradients systematically decreasing to the lowest categories. However, residual height showed systematically larger differences between the SEP categories as compared to PGS-height. For example, when residual height was 0.34 SD (95% CI 0.28–0.40) higher in those men and 0.33 SD (95% CI 0.27–0.38) higher in those women having higher tertiary education as compared to the basic education category, the differences in PGS-height were only 0.18 SD (95% CI 0.12–0.24) and 0.20 SD (95% CI 0.15–0.26), respectively. Similar differences between residual height and PGS-height were found for all SEP indicators in men and women. Model 2 includes SEP indicators simultaneously adjusted for each other. The parameter estimates decreased as compared to Model 1, though residual height again showed larger SEP differences than PGS-height. All SEP indicators still showed gradients for height, PGS-height and residual height.

After the analyses of SEP differences of height indicators, we studied how these SEP and height indicators were associated with CHD incidence. We found that in men (Table 3) and women (Table 4) there were clear gradients in CHD incidence across all SEP indicators. When modelled separately, height and residual height were associated with CHD incidence with similar effect sizes among both men and women; for PGS-height, this association was weak (Model 1). The adjustment for genetic and residual height (Model 2) explained only a small proportion of social inequalities in CHD incidence (0 – 6% indexed by PAF) whereas other risk factors for CHD (Model 3) explained a more substantial proportion (19 – 39%). When all SEP indicators were included in the model

simultaneously (Model 4), education in men and social class in women showed the largest inequalities in CHD risk. Metabolic risk factors of CHD showed negative correlations with all height indicators (Supplementary table 3). Adjusting the results for metabolic and social risk factors of CHD diminished the associations of height and residual height with CHD incidence whereas for PGS-height the association was essentially null (Model 4).

Discussion

We found that both PGS-height and residual height adjusted for PGS-height showed clear gradients according to education, occupation-based social class and income. The gradients were also found when mutually adjusting for all SEP indicators suggesting that they capture different social dimensions relevant for height. Residual height was also inversely associated with CHD incidence. For PGS-height, this association was weaker and totally explained by metabolic and social risk factors of CHD. Our results concerning PGS-height are consistent with the studies based on the UK Biobank also finding associations between PGS-height and several SES indicators [10]. However, we could not replicate the results in the UK Biobank on the association between PGS-height and CHD incidence [11] found also within sibling pairs [22]. Uniquely, we showed that these height associations were systematically larger when measured height was adjusted for the known genetic variants for height, supporting the role of environmental factors behind the associations of height with SEP and CHD risk. However, PGS-height explained only around a third of height variation, which is half or less of the twin heritability in these birth cohorts [8]. Residual height thus also partly reflects unknown genetic variation as well as measurement error, in addition to environmental variation. Nevertheless, the stronger residual height than PGS-height associations with SEP and CHD incidence suggest that environmental factors are more strongly driving associations of short height with lower SEP and increased CHD risk than genetic factors.

The reasons why PGS-height is associated with SEP is not clear. A large study of Swedish conscripts found that height in early adulthood predicted further achievement of higher education after adjustment for parental social position and own cognitive ability, a result which is consistent with the causal hypothesis [16]. The association between height and social position can originate from positive stereotypes related to tall stature which could help proceed in educational and occupational careers; however, empirical evidence is mixed [23]. Previous twin studies have found genetic correlations between height and cognitive ability [24–26] as well as education [27]. It is thus possible that there are, for example, hormonal mechanisms jointly affecting growth and cognitive development. It is also possible that PGS-height captures social or demographic variation which underlies the associations with social position and CHD incidence [28]. However, this explanation was not supported in a recent study reporting that the association between PGS-height and measured height was roughly similar within sibling pairs than it was at population level [29].

Although we could not empirically assess factors driving the associations of residual height with SEP and CHD risk, their effect is likely to start early in the life course or even prenatally. In a study including undernourished pregnant mothers, energy supplementation increased the length of neonates [30]. Further, environmental factors during the first two years of life are important for growth strongly affecting later height differences [31]. For nutrition, protein is considered the most important macro-nutrient needed for growth, but it also requires balanced micronutrients, and childhood infections can also delay growth [12]. Thus, detailed longitudinal measures of childhood nutrition and health would be crucial to better understand the environmental factors behind height variation.

Even when we found that both PGS-height and residual height showed clear SEP gradients and residual height was also associated with CHD incidence, they had only a negligible contribution to social inequalities in CHD risk. Basic metabolic risk factors and smoking measured at baseline explained nearly ten times more of SEP inequalities in CHD risk than these two height indicators together. Thus, even when there is a large body of literature concerning social inequalities in CHD risk [1], social height differences [5], and the associations between height and CHD risk [6,7], caution is needed when evaluating whether height can provide meaningful additional information on early life to explain social inequalities in CHD risk.

Our data have both strengths and limitations. The strength of our data was the large sample size that allowed us to estimate the associations reliably. Information on all SEP indicators was register based and therefore not prone to recall bias. Our data also allowed us to index SEP at the same age, thus decreasing heterogeneity in our analyses. The main limitation of our data was that we had only information on adult stature. The association between height in childhood and CHD in adulthood becomes weaker when children become older because of higher CHD incidence among children experiencing catch-up growth [32]. Thus, height measured in early childhood may capture the effect of childhood environmental factors better than adult height. Further, there is evidence that leg length is more sensitive to the effects of environmental factors [33] and predict CHD risk better than overall stature [34]; thus, this measure may have produced stronger associations. Since height was measured at baseline, shrinkage may have affected the results. However, only 2% of participants were older than 70 years at baseline, and only after this age is any substantial shrinking observed [35]. Selective mortality according to SEP and height and typically higher participation rates of those with high SEP may have led to sample selection bias in our data [36]. However, the high response rates for the baseline surveys and lack of loss to follow-up because of relying on register-based CHD incidence data helped to minimize other influences of selection bias in our

study. Finally, our results should not be directly generalize to other populations having a different distribution of stressors in childhood. Also the studied birth cohorts were nearly totally European ancestry and both population level incidence and SEP inequalities in CHD are high in the Finnish population [37]. Thus, replication of these results in populations with various standards of living, nutrition and other environmental factors in childhood as well as non-European ancestry and different population level risk of CHD would be important.

In conclusion, we found strong evidence that genetic polymorphisms associated with height are also associated with social position. We also found suggestive evidence that environmental factors affecting height are more common in disadvantages SEP groups and increase CHD risk. However, since these associations were not very strong, height did not explain much of the social inequalities in CHD risk. Overall, our finding of an association of residual height with socioeconomic indicators and CHD is consistent with the idea that height reflects material and social living conditions in childhood.

Ethics approval

The Finnish Social and Health Data Permit Authority (Findata) has accepted the use of clinical data (THL/1423/14.06.00/2022) and the data linkage to the Finnish population registers (TK/2654/07.03.00/2021). All participants gave informed consent when participating in the study. The samples/data used for the research were obtained from THL Biobank (study number: THLBB2020_8). We thank all study participants for their generous participation at THL Biobank.

Funding

This work was supported by the Academy of Finland grant (#345219) for Hannu Lahtinen. Pekka Martikainen was supported by the Academy of Finland (#308247, # 345219), the European Research Council under the European Union's Horizon 2020 research and innovation programme (grant agreement No 101019329), and the Max Planck – University of Helsinki Center for Social Inequalities in Population Health. George Davey Smith and Tim Morris work within the MRC Integrative Epidemiology Unit at the University of Bristol, which is supported by the Medical Research Council (MC_UU_00011/1).

Sample Data Availability Statements

The data underlying this article were provided by third party by permission. Data will be shared on request to the corresponding author with permission of third party.

Conflicts of interests

None declared.

Author contributions

All authors contributed to the study conception and design. HL performed the analyses. KS prepared the first draft of the manuscript. HL, GDS, TM and PM revised the manuscript critically for important intellectual content. All authors approved the final version of the manuscript and agree to be accountable for all aspects of the work thereby ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

References

1. Manrique-Garcia E, Sidorchuk A, Hallqvist J, Moradi T. Socioeconomic position and incidence of acute myocardial infarction: a meta-analysis. *J Epidemiol Community Health*. 2011;65(4):301–9.
2. Martikainen P, Mäkelä P, Peltonen R, Myrskylä M. Income differences in life expectancy: the changing contribution of harmful consumption of alcohol and smoking. *Epidemiology*. 2014;25(2):182–90.
3. Davey Smith G, Hart C, Blane D, Hole D. Adverse socioeconomic conditions in childhood and cause specific adult mortality: prospective observational study. *BMJ*. 1998;316(7145):1631–5.
4. Grasgruber P, Hrazdíra E. Nutritional and socio-economic predictors of adult height in 152 world populations. *Econ Hum Biol*. 2020;37:100848.
5. Steckel RH. Heights and human welfare: recent developments and new directions. *Explorations in Economic History*. 2009;46:1–23.
6. Paajanen TA, Oksala NKJ, Kuukasjärvi P, Karhunen PJ. Short stature is associated with coronary heart disease: a systematic review of the literature and a meta-analysis. *Eur Heart J*. 2010;31(14):1802–9.
7. Emerging Risk Factors Collaboration. Adult height and the risk of cause-specific death and vascular morbidity in 1 million people: individual participant meta-analysis. *Int J Epidemiol*. 2012;41(5):1419–33.
8. Jelenkovic A, Hur YM, Sund R, Yokoyama Y, Siribaddana SH, Hotopf M, et al. Genetic and environmental influences on adult human height across birth cohorts from 1886 to 1994. *Elife*. 2016;5:e20320.
9. Yengo L, Vedantam S, Marouli E, Sidorenko J, Bartell E, Sakaue S, et al. A saturated map of common genetic variants associated with human height. *Nature*. 2022 Oct;610(7933):704–12.
10. Tyrrell J, Jones SE, Beaumont R, Astley CM, Lovell R, Yaghootkar H, et al. Height, body mass index, and socioeconomic status: Mendelian randomisation study in UK Biobank. *BMJ*. 2016;352:i582.
11. Nüesch E, Dale C, Palmer TM, White J, Keating BJ, van Iperen EP, et al. Adult height, coronary heart disease and stroke: a multi-locus Mendelian randomization meta-analysis. *Int J Epidemiol*. 2016;45(6):1927–37.
12. Millward DJ. Nutrition, infection and stunting: the roles of deficiencies of individual nutrients and foods, and of inflammation, as determinants of reduced linear growth of children. *Nutr Res Rev*. 2017;30(1):50–72.
13. Erola J, Jalonen S, Lehti H. Parental education, class and income over early life course and children's achievement. *Res Soc Stratif Mobil*. 2016;44:33–43.

14. Renthal NE, Nakka P, Baronas JM, Kronenberg HM, Hirschhorn JN. Genes with specificity for expression in the round cell layer of the growth plate are enriched in genomewide association study (GWAS) of human height. *J Bone Miner Res.* 2021;36(12):2300–8.
15. Silventoinen K, Maia J, Li W, Sund R, Gouveia ÉR, Antunes A, et al. Genetic regulation of body size and morphology in children: a twin study of 22 anthropometric traits. *Int J Obes.* 2023 (online ahead of print).
16. Magnusson PKE, Rasmussen F, Gyllensten UB. Height at age 18 years is a strong predictor of attained education later in life: cohort study of over 950,000 Swedish men. *Int J Epidemiol.* 2006;35(3):658–63.
17. Mackenbach JP, Rubio Valverde J, Bopp M, Brønnum-Hansen H, Costa G, Deboosere P, et al. Progress against inequalities in mortality: register-based study of 15 European countries between 1990 and 2015. *Eur J Epidemiol.* 2019 Dec;34(12):1131–42.
18. Perkins JM, Subramanian SV, Davey Smith G, Özaltin E. Adult height, nutrition, and population health. *Nutr Rev.* 2016;74(3):149–65.
19. Paalanen L, Härkänen T, Tolonen H. Protocol of a research project “Projections of the burden of disease and disability in Finland - health policy prospects” using cross-sectional health surveys and register-based follow-up. *BMJ Open.* 2019 Jun 20;9(6):e029338.
20. Lu T, Forgetta V, Wu H, Perry JRB, Ong KK, Greenwood CMT, et al. A polygenic risk score to predict future adult short stature among children. *J Clin Endocrinol Metab.* 2021;106(7):1918–28.
21. Statistics Finland. Classification of Socio-economic Groups 1989 [Internet]. 2022. Available from: https://www2.tilastokeskus.fi/en/luokitukset/sosioekon_asema/sosioekon_asema_1_19890101/
22. Howe LJ, Brumpton B, Rasheed H, Åsvold BO, Davey Smith G, Davies NM. Taller height and risk of coronary heart disease and cancer: A within-sibship Mendelian randomization study. *Elife.* 2022;11:e72984.
23. Bittmann F. The relationship between height and leadership: Evidence from across Europe. *Econ Hum Biol.* 2020 Jan;36:100829.
24. Silventoinen K, Posthuma D, van Beijsterveldt T, Bartels M, Boomsma DI. Genetic contributions to the association between height and intelligence: Evidence from Dutch twin data from childhood to middle age. *Genes Brain Behav.* 2006;5(8):585–95.
25. Keller MC, Garver-Apgar CE, Wright MJ, Martin NG, Corley RP, Stallings MC, et al. The genetic correlation between height and IQ: shared genes or assortative mating? *PLoS Genet.* 2013;9(4):e1003451.
26. Silventoinen K, Iacono WG, Krueger R, McGue M. Genetic and environmental contributions to the association between anthropometric measures and IQ: a study of Minnesota twins at age 11 and 17. *Behav Genet.* 2012;42(3):393–401.

27. Silventoinen K, Krueger RF, Bouchard TJ, Kaprio J, McGue M. Heritability of body height and educational attainment in an international context: comparison of adult twins in Minnesota and Finland. *Am J Hum Biol.* 2004;16(5):544–55.
28. Morris TT, Davies NM, Hemani G, Smith GD. Population phenomena inflate genetic associations of complex social traits. *Sci Adv.* 2020;6(16):eaay0328.
29. Howe LJ, Nivard MG, Morris TT, Hansen AF, Rasheed H, Cho Y, et al. Within-sibship genome-wide association analyses decrease bias in estimates of direct genetic effects. *Nat Genet.* 2022;
30. Kusin JA, Kardjati S, Houtkooper JM, Renqvist UH. Energy supplementation during pregnancy and postnatal growth. *Lancet.* 1992;340(8820):623–6.
31. Leroy JL, Ruel M, Habicht JP, Frongillo EA. Linear growth deficit continues to accumulate beyond the first 1000 days in low- and middle-income countries: global evidence from 51 national surveys. *J Nutr.* 2014;144(9):1460–6.
32. Silventoinen K, Baker JL, Sørensen TIA. Growth in height in childhood and risk of coronary heart disease in adult men and women. *PLoS One.* 2012;7(1):e30476.
33. Gunnell DJ, Davey Smith G, Frankel SJ, Kemp M, Peters TJ. Socio-economic and dietary influences on leg length and trunk length in childhood: a reanalysis of the Carnegie (Boyd Orr) survey of diet and health in prewar Britain (1937-39). *Paediatr Perinat Epidemiol.* 1998;12 Suppl 1:96–113.
34. Gunnell DJ, Davey Smith G, Frankel S, Nanchahal K, Braddon FE, Pemberton J, et al. Childhood leg length and adult mortality: follow up of the Carnegie (Boyd Orr) Survey of Diet and Health in Pre-war Britain. *J Epidemiol Community Health.* 1998;52(3):142–52.
35. Sorkin JD, Muller DC, Andres R. Longitudinal change in height of men and women: implications for interpretation of the body mass index: the Baltimore Longitudinal Study of Aging. *Am J Epidemiol.* 1999;150(9):969–77.
36. Reinikainen J, Tolonen H, Borodulin K, Härkänen T, Jousilahti P, Karvanen J, et al. Participation rates by educational levels have diverged during 25 years in Finnish health examination surveys. *Eur J Public Health.* 2018;28(2):237–43.
37. Kulhánová I, Bacigalupe A, Eikemo TA, Borrell C, Regidor E, Esnaola S, et al. Why does Spain have smaller inequalities in mortality? An exploration of potential explanations. *Eur J Public Health.* 2014;24(3):370–7.

Figure legends

Figure 1. Possible pathways between height, socio-economic position (SEP) and coronary heart disease (CHD).

Table 1. Differences by 1 standard deviation of height, PGS-height and residual height between classes of socioeconomic position indicators in men.

	Model 1									Model 2								
	Height			PGS-height			Residual height			Height			PGS-height			Residual height		
	β	95% CI		β	95% CI		β	95% CI		β	95% CI		β	95% CI		β	95% CI	
		LL	UL		LL	UL		LL	UL		LL	UL		LL	UL		LL	UL
Education																		
Basic	ref.			ref.			ref.			ref.			ref.			ref.		
Secondary	0.10	0.06	0.14	0.07	0.03	0.11	0.07	0.03	0.11	0.08	0.04	0.13	0.07	0.03	0.11	0.06	0.02	0.10
Lower tertiary	0.28	0.23	0.32	0.16	0.11	0.21	0.23	0.18	0.27	0.19	0.14	0.25	0.15	0.09	0.20	0.13	0.08	0.19
Higher tertiary	0.38	0.32	0.44	0.18	0.12	0.24	0.34	0.28	0.40	0.24	0.17	0.31	0.13	0.06	0.20	0.20	0.13	0.27
Social class																		
Manual	ref.			ref.			ref.			ref.			ref.			ref.		
Lower non-manual	0.12	0.07	0.16	0.01	-0.03	0.05	0.13	0.09	0.17	0.02	-0.03	0.06	-0.05	-0.10	-0.01	0.05	0.01	0.10
Upper non-manual	0.30	0.26	0.35	0.14	0.10	0.19	0.27	0.23	0.31	0.11	0.05	0.16	0.03	-0.03	0.09	0.11	0.05	0.16
Entrepreneurs	0.08	0.02	0.14	0.08	0.02	0.14	0.04	-0.01	0.10	0.05	-0.01	0.11	0.06	0.00	0.12	0.02	-0.04	0.08
Farmers	0.13	0.07	0.19	0.12	0.06	0.19	0.08	0.01	0.14	0.16	0.09	0.22	0.14	0.07	0.21	0.10	0.03	0.16
Income																		
Lowest quintile	ref.			ref.			ref.			ref.			ref.			ref.		
4. quintile	0.00	-0.07	0.06	-0.01	-0.07	0.05	0.00	-0.06	0.06	0.01	-0.05	0.07	0.00	-0.06	0.07	0.01	-0.05	0.07
3. quintile	0.00	-0.06	0.06	-0.01	-0.07	0.05	0.01	-0.05	0.06	0.03	-0.03	0.09	0.02	-0.04	0.08	0.02	-0.04	0.08
2. quintile	0.11	0.06	0.16	0.05	-0.01	0.10	0.10	0.05	0.15	0.12	0.06	0.17	0.07	0.01	0.13	0.09	0.04	0.15
Highest quintile	0.26	0.21	0.31	0.12	0.06	0.17	0.23	0.18	0.28	0.17	0.12	0.23	0.09	0.04	0.15	0.15	0.09	0.20

Abbreviations: β =regression coefficient, CI=confidence interval, LL=lower limit, UL=upper limit

Model 1: SEP indicators separately adjusted by year of birth (1935=0), year of birth squared, region of residence, ten first principal components of the genetic structure and genotyping batch-data collection round combination. Model 2: Model 1+ education, social class and income.

Table 2. Differences by 1 standard deviation of height, PGS-height and residual height between classes of socioeconomic position indicators in women.

	Model 1									Model 2								
	Height			PGS-height			Residual height			Height			PGS-height			Residual height		
	β	95% CI		β	95% CI		β	95% CI		β	95% CI		β	95% CI		β	95% CI	
		LL	UL		LL	UL		LL	UL		LL	UL		LL	UL		LL	UL
Education																		
Basic	ref.			ref.			ref.			ref.			ref.			ref.		
Secondary	0.12	0.08	0.16	0.08	0.04	0.13	0.09	0.05	0.13	0.11	0.07	0.15	0.08	0.04	0.12	0.08	0.03	0.12
Lower tertiary	0.28	0.23	0.32	0.14	0.09	0.19	0.24	0.20	0.29	0.22	0.17	0.26	0.13	0.08	0.18	0.17	0.13	0.22
Higher tertiary	0.39	0.33	0.44	0.20	0.15	0.26	0.33	0.27	0.38	0.27	0.20	0.34	0.17	0.10	0.24	0.21	0.14	0.28
Social class																		
Manual	ref.			ref.			ref.			ref.			ref.			ref.		
Lower non-manual	0.13	0.10	0.17	0.06	0.03	0.10	0.12	0.08	0.15	0.06	0.02	0.09	0.03	-0.01	0.07	0.04	0.01	0.08
Upper non-manual	0.29	0.24	0.33	0.13	0.09	0.18	0.26	0.21	0.30	0.10	0.04	0.15	0.05	-0.01	0.10	0.09	0.03	0.14
Entrepreneurs	0.15	0.08	0.21	0.10	0.04	0.17	0.11	0.04	0.17	0.11	0.04	0.18	0.09	0.02	0.16	0.07	0.00	0.14
Farmers	0.04	-0.03	0.11	0.08	0.01	0.15	-0.01	-0.08	0.07	0.05	-0.02	0.12	0.09	0.01	0.16	0.00	-0.07	0.07
Income																		
Lowest quintile	ref.			ref.			ref.			ref.			ref.			ref.		
4. quintile	0.04	0.00	0.08	0.02	-0.02	0.07	0.03	-0.01	0.07	0.04	0.00	0.08	0.03	-0.01	0.08	0.03	-0.01	0.07
3. quintile	0.12	0.07	0.16	0.04	-0.01	0.08	0.11	0.07	0.16	0.08	0.04	0.13	0.03	-0.02	0.07	0.08	0.04	0.12
2. quintile	0.20	0.15	0.24	0.04	-0.01	0.09	0.21	0.16	0.25	0.11	0.06	0.16	0.00	-0.05	0.05	0.13	0.08	0.18
Highest quintile	0.30	0.25	0.36	0.14	0.08	0.19	0.27	0.22	0.33	0.16	0.10	0.22	0.06	0.00	0.13	0.15	0.08	0.21

Abbreviations: β =regression coefficient, CI=confidence interval, LL=lower limit, UL=upper limit

Model 1: SEP indicators separately adjusted by year of birth (1935=0), year of birth squared, region of residence, ten first principal components of the genetic structure and genotyping batch-data collection round combination. Model 2: Model 1+ education, social class and income.

Table 3. Hazard ratios of socioeconomic position indicators, PGS-height and residual height for coronary heart disease incidence in men.

	Model 1			Model 2			Model 3			Model 4		
	HR	95% CI		HR	95% CI		HR	95% CI		HR	95% CI	
		LL	UL		LL	UL		LL	UL		LL	UL
Education												
Basic	1.88	1.44	2.47	1.86	1.41	2.44	1.49	1.13	1.96	1.46	1.05	2.03
Secondary	1.71	1.31	2.24	1.69	1.29	2.21	1.41	1.07	1.85	1.39	1.01	1.92
Lower tertiary	1.21	0.91	1.62	1.20	0.90	1.61	1.07	0.80	1.43	1.11	0.81	1.50
Higher tertiary	ref.			ref.			ref.			ref.		
PAF ^a	0.38	0.21	0.51	0.37	0.20	0.51	0.26	0.05	0.42	0.25	0.01	0.44
Social class												
Manual	1.46	1.23	1.73	1.44	1.21	1.71	1.26	1.06	1.51	0.97	0.77	1.22
Lower non-manual	1.12	0.91	1.37	1.11	0.90	1.35	1.02	0.84	1.25	0.89	0.71	1.11
Upper non-manual	ref.			ref.			ref.			ref.		
Entrepreneurs	1.47	1.14	1.88	1.45	1.13	1.86	1.26	0.98	1.62	0.95	0.72	1.26
Farmers	1.25	0.96	1.61	1.23	0.95	1.59	1.19	0.92	1.54	0.80	0.59	1.08
PAF ^a	0.22	0.11	0.32	0.21	0.10	0.31	0.14	0.01	0.25	-0.06	-0.28	0.12
Income												
Lowest quintile	1.79	1.49	2.15	1.77	1.47	2.13	1.57	1.31	1.89	1.47	1.20	1.80
4. quintile	1.42	1.17	1.72	1.39	1.15	1.70	1.28	1.06	1.56	1.18	0.95	1.45
3. quintile	1.16	0.97	1.39	1.15	0.96	1.37	1.06	0.89	1.27	0.96	0.79	1.16
2. quintile	1.25	1.07	1.46	1.24	1.06	1.45	1.18	1.01	1.38	1.09	0.92	1.28
Highest quintile	ref.			ref.			ref.			ref.		
PAF ^a	0.18	0.11	0.25	0.18	0.10	0.24	0.14	0.06	0.21	0.08	-0.01	0.17
Height ^b	0.94	0.88	1.00							0.98	0.92	1.04
PGS-height ^b	0.99	0.93	1.05							1.02	0.96	1.08
Residual height ^b	0.94	0.88	1.00							0.97	0.91	1.03

Abbreviations: HR=hazard ratio, CI=confidence interval, LL=lower limit, UL=upper limit, PAF= population attributable fraction

Model 1: SEP and height indicators added into the model separately+ age at baseline + age at baseline squared, region of residence + ten first principal components of the genetic structure + genotyping batch-data collection round (=baseline year) combination. Model 2: Model 1 + PGS-height + residual height. Model 3: Model 2 + smoking status+ body mass index + systolic blood pressure + diastolic blood pressure + high-density lipoprotein (HDL) cholesterol + total cholesterol. Model 4: Model 3+all SEP indicators (height indicators)/all SEP indicators+PGS-height+residual height (SEP indicators)

^a PAFs for CHD incidence for the socioeconomic position indicators indicate the proportion of CHD cases that would be avoided had the higher risk SEP categories the same CHD risk as the lowest risk SEP category.

^bHRs calculated per 1 standard deviation.

Table 4. Hazard ratios of socioeconomic position indicators, PGS-height and residual height for coronary heart disease incidence in women.

	Model 1			Model 2			Model 3			Model 4		
	HR	95% CI		HR	95% CI		HR	95% CI		HR	95% CI	
		LL	UL		LL	UL		LL	UL		LL	UL
Education												
Basic	1.85	1.21	2.83	1.77	1.16	2.71	1.44	0.94	2.22	0.98	0.58	1.68
Secondary	1.44	0.94	2.20	1.40	0.91	2.13	1.20	0.79	1.84	0.84	0.50	1.41
Lower tertiary	1.07	0.68	1.67	1.06	0.67	1.66	0.98	0.62	1.54	0.78	0.47	1.30
Higher tertiary	ref.			ref.			ref.			ref.		
PAF ^a	0.33	0.01	0.54	0.31	-0.02	0.53	0.20	-0.18	0.46	-0.12	-0.81	0.31
Social class												
Manual	1.96	1.40	2.72	1.88	1.35	2.62	1.65	1.18	2.30	1.43	0.94	2.18
Lower non-manual	1.62	1.18	2.22	1.58	1.15	2.18	1.45	1.05	2.00	1.34	0.91	1.98
Upper non-manual	ref.			ref.			ref.			ref.		
Entrepreneurs	1.41	0.86	2.32	1.37	0.83	2.26	1.16	0.70	1.92	1.06	0.62	1.84
Farmers	1.90	1.26	2.88	1.83	1.21	2.77	1.63	1.07	2.48	1.47	0.90	2.41
PAF ^a	0.39	0.20	0.53	0.38	0.18	0.52	0.31	0.10	0.48	0.25	-0.06	0.47
Income												
Lowest quintile	1.64	1.08	2.50	1.58	1.04	2.41	1.43	0.94	2.17	1.19	0.74	1.92
4. quintile	1.83	1.21	2.78	1.76	1.16	2.67	1.61	1.06	2.44	1.33	0.82	2.14
3. quintile	1.58	1.03	2.42	1.54	1.01	2.36	1.45	0.95	2.23	1.25	0.78	2.01
2. quintile	1.19	0.74	1.89	1.17	0.74	1.87	1.12	0.70	1.79	1.03	0.63	1.69
Highest quintile	ref.			ref.			ref.			ref.		
PAF ^a	0.36	0.08	0.56	0.35	0.05	0.55	0.29	-0.03	0.51	0.18	-0.26	0.46
Height ^b	0.87	0.79	0.96							0.91	0.82	1.00
PGS-height ^b	0.97	0.89	1.06							0.99	0.91	1.09
Residual height ^b	0.87	0.79	0.95							0.90	0.82	0.99

Abbreviations: HR=hazard ratio, CI=confidence interval, LL=lower limit, UL=upper limit, PAF= population attributable fraction

Model 1: SEP and height indicators added into the model separately+ age at baseline + age at baseline squared, region of residence + ten first principal components of the genetic structure + genotyping batch-data collection round (=baseline year) combination. Model 2: Model 1 + PGS-

height + residual height. Model 3: Model 2 + smoking status+ body mass index + systolic blood pressure + diastolic blood pressure + high-density lipoprotein (HDL) cholesterol + total cholesterol. Model 4: Model 3+all SEP indicators (height indicators)/all SEP indicators+PGS-height+residual height (SEP indicators)

^a PAFs for CHD incidence for the socioeconomic position indicators indicate the proportion of CHD cases that would be avoided had the higher risk SEP categories the same CHD risk as the lowest risk SEP category.

^bHRs calculated per 1 standard deviation.