

Metastatic Sites' Location and Impact on Patient Management After the Introduction of Prostate-specific Membrane Antigen Positron Emission Tomography in Newly Diagnosed and Biochemically Recurrent Prostate Cancer: A Critical Review

Francesco Mattana ^{a †}, Lorenzo Muraglia ^{a †}, Pawel Rajwa ^{b c}, Fabio Zattoni ^d, Giancarlo Marr ^{a e}, Peter K.F. Chiu ^f, Isabel Heidegger ^g, Veeru Kasivisvanathan ^h, Claudia V. Kesch ^{i j}, Jonathan Olivier ^k, Felix Preisser ^l, Constance Thibault ^m, Massimo Valerio ⁿ, Roderick C.N. van den Bergh ^o, Giorgio Gandaglia ^{p q ‡}, Francesco Ceci ^{a r ‡}, European Association of Urology Young Academic Urologists Prostate Cancer Working Party

a

Division of Nuclear Medicine, IEO European Institute of Oncology IRCCS, Milan, Italy

b

Department of Urology, Medical University of Vienna, Vienna, Austria

c

Department of Urology, Medical University of Silesia, Zabrze, Poland

d

Department of Surgery, Oncology and Gastroenterology, Urology Clinic, University of Padova, Padova, Italy

e

Division of Urology, Department of Surgical Sciences, AOU Città della Salute e della Scienza di Torino, Università di Torino, Turin, Italy

f

SH Ho Urology Centre, Department of Surgery, The Chinese University of Hong Kong, Hong Kong

g

Department of Urology, Medical University Innsbruck, Innsbruck, Austria

h

Division of Surgery and Interventional Science, University College London, London, UK

i

Department of Urology, University Hospital Essen, Essen, Germany

j

German Cancer Consortium (DKTK)-University Hospital Essen, Essen, Germany

k

Department of Urology, Lille University Hospital, Lille, France

l

Department of Urology, University Hospital Frankfurt, Frankfurt am Main, Germany

m

Department of Medical Oncology, European Georges Pompidou Hospital, Assistance Publique des Hôpitaux de Paris, Paris Descartes University, Paris, France

n

Department of Urology, Lausanne University Hospital, Lausanne, Switzerland

o

Department of Urology, Antonius Hospital, Utrecht, The Netherlands

p

Unit of Urology/Division of Oncology, IRCCS Ospedale San Raffaele, Milan, Italy

q

Vita-Salute San Raffaele University, Milan, Italy
r

Department of Oncology and Hemato-Oncology, University of Milan, Milan, Italy

Abstract

Context

The introduction of prostate-specific membrane antigen positron emission tomography (PSMA-PET) had a substantial impact on the management of prostate cancer (PCa) patients with a stage migration phenomenon and consequent treatment changes.

Objective

To summarise the role of PSMA-PET to define the burden of disease through an accurate location of metastatic site(s) in PCa patients, describing the most common locations at PSMA-PET in the primary staging and recurrence setting, and to assess the clinical impact in the decision-making process.

Evidence acquisition

A comprehensive nonsystematic literature review was performed in April 2022. Literature search was updated until March 2022. The most relevant studies have been summarised, giving priority to registered clinical trials and multicentre collaborations.

Evidence synthesis

PSMA-PET showed higher diagnostic accuracy than conventional imaging both in newly diagnosed PCa and in recurrent disease. This greater accuracy led to a migration of a higher proportion of patients identified with metastatic disease. Bone metastases were reported as the most frequent site of metastatic spread in staging (up to 17%) and restaging (up to 18%). In staging, considering the suboptimal sensitivity in lymph node metastasis detection prior to radical surgery, PSMA-PET should be performed in patients with high risk or unfavourable intermediate risk only, and it is not recommended to routinely avoid pelvic lymph node dissection in case of a negative scan. In case of prostate-specific antigen relapse, PSMA-PET had higher diagnostic accuracy than other diagnostic procedures in the early detection of the sites of recurrence, thus influencing the therapy decision-making process.

Conclusions

PSMA-PET detects a higher number of lesions than conventional imaging or other PET radiotracers, especially metastatic lesions unseen with other modalities. The high diagnostic accuracy of PSMA-PET leads to a significant patient upstage and thus an impact in clinical management, even if the overall impact on cancer mortality is still to be assessed.

Patient summary

Prostate-specific membrane antigen positron emission tomography (PSMA-PET) identifies metastatic lesions with higher accuracy than conventional imaging, both in primary prostate cancer and during disease recurrence. Skeletal metastasis and extrapelvic lymph nodes are the most common sites of metastatic spread. The high accuracy of PSMA-PET in the detection of metastatic disease led to a significant impact on patient management, even if the overall impact on cancer mortality is still to be assessed.

1. Introduction

Prostate-specific membrane antigen (PSMA) is considered a key target for molecular imaging in prostate cancer (PCa) patients. Several PSMA inhibitors have been proposed for positron emission tomography (PET) imaging [1], and phase III clinical studies proved the superiority of PSMA-PET to computed tomography (CT) or bone scan (BS) in patients affected by PCa in different clinical settings [2]. PSMA-PET is currently considered the standard of care in patients with biochemical relapse after radical treatment due to the potential implications for patient management and the administration of PSMA-guided therapy including metastasis-directed therapies (MDTs) [2]. Similarly, prospective trials demonstrated the superiority of PSMA-PET to conventional imaging both as a staging procedure prior to surgery and in the diagnostic/biopsy setting, as well as for the identification of candidates for radioligand therapy in advanced PCa. In this context, the results derived by clinical trials (eg, proPSMA and PRIMARY) [3], [4] highlighted the advantage of having a new-generation imaging procedure to assess the disease burden more accurately prior to surgery, but also emphasise the potential role of characterising the intraprostatic lesion(s). Nonetheless, the routine use of PSMA-PET in all newly diagnosed PCa patients is not recommended by clinical guidelines due to the lack of evidence on long-term oncological control and the stage migration phenomenon, where men who were staged as M0 at conventional imaging would now be considered as M1 at advanced imaging with consequent changes in the treatment paradigm [2]. Similarly, there is a lack of evidence regarding the role of PSMA-PET as a baseline procedure prior to the administration of systemic therapies in case of metastatic disease. Thus, a clinical decision to switch to subsequent lines in advanced PCa cannot be taken relying on PSMA-PET only. Although data regarding the impact of PSMA-PET on PCa cancer-specific and overall mortality are still awaited [5], its implementation in the management of PCa patients is already of high clinical interest.

The aim of this critical review was summarising the role of PSMA-PET to define the burden of disease through an accurate location of metastatic site(s) in PCa patients, to describe the most common sites of positive spots at PSMA-PET in newly diagnosed PCa and during biochemical recurrence (BCR), and to assess its clinical impact in the decision-making process.

2. Evidence acquisition

A comprehensive literature review was performed in April 2022 with a nonsystematic approach. The search is updated until March 2022, and was performed using the Ovid platform and a comparison of the Embase and Medline databases, using the following string: ("prostate specific membrane antigen" OR "PSMA") AND ("Positron Emission Tomography" OR PET) AND ("prostate cancer" OR PCa). The most relevant studies have been summarised, giving priority to registered clinical trials and multicentre collaborations. Four authors (F.M., L.M., P.R., and F.Z.) performed the literature research. Disagreements have been resolved by consensus. All the original articles published in English over the past 10 yr were considered. Retrospective and prospective series, as well as randomised and nonrandomised clinical trials reporting data about metastatic sites location and impact on patient management have been considered. Abstract, narrative review, case reports or case series, editorials, and letter to editors have been excluded. In the study selection process, priority was given to randomised clinical trials, prospective academic studies, or retrospective multicentre collaborations involving high-volume centres with proven expertise in PSMA-PET. For clinical studies, all PSMA radiopharmaceuticals were considered;

the most frequent PET tracer used was ^{68}Ga -PSMA-11, followed by ^{18}F -DCFPyL and ^{18}F -PSMA-1007. The literature search was updated until March 2022. After the first literature screening, a total of 38 studies have been selected. Authors tabulated and organised relevant studies and performed a comprehensive qualitative narrative synthesis of both tabulated studies and nontabulated articles.

3. Evidence synthesis

3.1. Staging prostate cancer with PSMA-PET: distribution of metastatic sites

Accurate staging of PCa is crucial to correct planning of curative-intent therapeutic strategies after disease diagnosis. PSMA-PET could be considered to stage high-risk localised or locally advanced PCa. A cross-sectional abdominopelvic imaging study for lymph node (LN) evaluation, such as CT, and a BS are still considered the staging pathway of reference and generally performed in clinical practice. According to the European Association of Urology (EAU) guidelines [2], PSMA-PET is more accurate for staging than CT and BS in high-risk disease (level of evidence 1b). However, even if PSMA-PET is performed as an imaging study with high diagnostic accuracy, data regarding long-term oncological control generated by the stage migration phenomenon are still missing.

Four studies ($n = 4$) reporting the distribution sites of positive spots at PSMA-PET were identified (Table 1). In a cohort of 691 consecutive high-risk PCa patients who performed PSMA-PET as a staging procedure, Klingenberg et al. [6] observed a disease with nodal involvement (N1/M1a) in 31% of patients, while skeletal involvement (M1b) was observed in 17%. The reported risk of advanced disease for potential clinically confined cancer (cT2a, cT2b, and cT2c) was almost equal (24%, 28%, and 22%, respectively). Sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), and accuracy for LN metastasis (LNM) detection were 31%, 97%, 69%, 85%, and 83%, respectively. However, there is heterogeneity regarding validation criteria commonly used to confirm PSMA-PET findings, thus reflecting heterogeneous results concerning its diagnostic performance among different studies (Table 1). A composite standard of truth has recently been accepted by the Food and Drug Administration (FDA) [7], [8]. These reference standards were defined as (1) evaluable histopathology results from prostatectomy, salvage pelvic LN dissection, or targeted biopsy; (2) correlative follow-up imaging findings using ^{18}F -fluciclovine or ^{11}C -choline PET, or focused magnetic resonance imaging (MRI) or CT; or (3) if neither of the above was available or informative, confirmed prostate-specific antigen (PSA) response up to 9 mo after radiation initiation (without concomitant androgen deprivation therapy [ADT]) of all PET-positive lesions. PSA response was defined as a PSA decline of $\geq 50\%$ from baseline.

In this clinical setting, a randomised controlled phase III trial has recently been published, comparing PSMA-PET with the standard of care (CT and BS). The proPSMA [3] trial reported higher specificity and PPV for LNM and bone metastasis localisation. Sensitivity remains suboptimal but significantly higher if compared with conventional imaging. First-line PSMA-PET ($n = 148$ patients) detected pelvic nodal disease (N) in 20% of cases, extrapelvic nodes (M1a) in 9%, bone metastases (M1b) in 10%, and visceral metastases (M1c) in 1%. Considering that 30% of the patients presented local or distant metastasis, PSMA-PET had a 27% (95% confidence interval [CI] 23–31, $p < 0.001$) absolute greater area under the curve (AUC) for accuracy when compared with conventional imaging (92% [88–95%] vs 65% [60–69%]), higher sensitivity (85% [74–96%] vs 38% [24–52%]), higher impact on clinical management (28% [21–36%] vs 15% [10–22%]; $p = 0.008$), and lower percentage of

inconclusive findings (7% [4–13%] vs 23% [17–21%]). Following first-line PSMA-PET, 14% patients shifted from curative to palliative-intent treatment, 11 (7%) had a change in radiotherapy technique, and 11 (7%) in surgical technique. In patients with fewer than three distant metastases on first-line imaging who crossed over to second-line imaging, conventional imaging had a high or medium effect in 5% (95% CI 2–10%) compared with 27% (20–35%) with PSMA-PET.

Further phase III studies assessed PSMA-PET accuracy in correctly detecting metastatic LNs using a histopathological analysis as reference standards. One of the largest prospective FDA-registered studies enrolled 764 patients, with 277 patients being eligible for a primary endpoint analysis and thus referred to radical prostatectomy and pelvic LN dissection, with histopathological validation [9]. In the per-patient analysis, PSMA-PET holds sensitivity of 44%, specificity of 97%, and a PPV of 88%. Interestingly, in the subcohort of patients not referred to surgery (nonsurgery cohort), skeletal metastases (M1b) have been observed in 18% of patients, while non-nodal visceral metastases (M1c) have been reported in 14%. The OSPREY trial [7] enrolled 252 patients studied with ¹⁸F-DCFPyL-PET and described sensitivity that ranged from 31% to 42% among three independent readers with high specificity (median 98%). Notably, these diagnostic performances differ from proPSMA results. Among 126 patients treated with radical prostatectomy, only 83 received histopathological validation. The diagnostic accuracy for this subgroup of patients was not reported, but lower sensitivity than the overall reported 85% might be expected, considering the data derived from literature. Recently, a meta-analysis reported overall sensitivity of 59% for PSMA-PET to detect pelvic LN involvement (range 23–100%) [10]. Patients' selection (high-risk vs high-risk and unfavourable intermediate-risk PCa), study design (prospective vs retrospective, consecutive vs nonconsecutive patients), sample size, and incidence of nodal metastasis in the selected cohort [11] are parameters related to this broad variability. Reader experience might also affect the overall diagnostic accuracy: equivocal findings on PSMA-PET, even if with a lower incidence than conventional imaging, remain challenging [12]. At present, a standardised reporting system (E-PSMA) has been published by the European Association of Nuclear Medicine developed to improve scan interpretation reproducibility [1]. However, external validation of these criteria is still awaited. Hence, the potential improvement on the diagnostic accuracy derived by the application of this reporting system still needs to be confirmed.

The higher NPV for nodal disease detection might contribute to select patients in which bilateral pelvic lymphadenectomy might be avoided. Roscigno et al. [13] retrospectively evaluated 630 consecutive intermediate- and high-risk PCa patients, reporting that nodal metastases were present in 133 patients. Out of these patients, 64 (48%), 58 (44%), 53 (40%), 16 (12%), and 20 (15%) had nodal metastases in the internal iliac, external iliac, obturator, common iliac, and presacral regions, respectively. Metastases in common iliac nodes were always associated with concomitant involvement of lower pelvic chains, confirming the theory of nodal metastasis ascending pathway. An extended pelvic lymphadenectomy (ePLND) would have removed all pathological nodes in 73% of patients only. Yaxley et al. [14] investigated the predictive value of preoperative PSMA-PET on LNM, concluding that by omitting ePLND based on negative preoperative PSMA-PET, nodal metastasis would have been missed in 20% of men. On the contrary, data from the proPSMA study and by Kulkarni et al. [15] showed a high NPV for PSMA-PET, missing LNM in 5% and 13%, respectively, in a high-risk setting.

Finally, according to the most updated literature, it is not recommended to routinely avoid pelvic LN dissection in case of negative PSMA-PET.

The results presented in this section about metastatic site location have been summarised in Table 1 and Figure 1.

3.2. Staging prostate cancer with PSMA-PET: impact on clinical management

The higher accuracy of PSMA-PET than that of conventional imaging generates a migration of patients towards different stages, generally leading to disease upstage. However, this more accurate staging is not sufficient to fully support its implementation as a standard of care procedure prior to primary therapy [3], [6], as its clinical net benefit should be assessed through the impact on treatment decisions and survival outcomes. While there is still a lack of strong evidence on the survival benefits due to the introduction of PSMA-PET in the management of newly diagnosed PCa, as PSMA-PET is still a novel technique, several studies analysed the impact of PSMA-PET on the decision-making process [3], [16], [17], [18]. In the proPSMA trial, first- and second-line PSMA-PET changed intent management in 28% and 27% of patients, respectively. Conventional imaging impacted the management in 15% (first line) and 8% (second line) of patients only [3]. Furthermore, the lower incidence of equivocal findings in PSMA-PET than in CT/BS (23% vs 7%) strengthens its reliability in daily clinical practice.

Real-world evidence from another prospective phase III trial [9] showed that urologists changed their management from radical prostatectomy to nonsurgical treatment options due to disease upstaging on molecular imaging. In detail, only 14% of D'Amico intermediate- and high-risk patients who underwent surgery had cN1 disease in PSMA-PET, while 52% with cN1 on PSMA-PET underwent different treatments. While the immediate impact on clinical management is evident, no data are supporting the hypothesis that sparing the patient from surgery will improve the overall survival. However, MDT can be offered to oligometastatic patients and a PSMA-guided therapy is a feasible approach [5].

PSMA-PET is a useful tool for treatment planning and guiding surgery or radiation therapy [18], [19], [20], [21]. Grubmüller et al. [17] found that PSMA-PET/MRI had sensitivity of 85% and specificity of 85% for detecting organ-confined disease, possibly allowing for nerve-sparing surgery. In total 29% of surgeons changed their surgical approach according to PSMA-PET/MRI results. In patients referred to radiotherapy as the primary intervention [18], the intended treatment planned prior to the PET scan was adapted in approximately 60% of cases [4], [22]. Moreover, PSMA-PET allows an accurate detection of metastatic lesions at presentation and has the potential to guide MDT, as it is an emerging treatment modality in low-volume metastatic patients, potentially improving the oncological outcomes [22], [23], [24]. Results about the impact of PSMA-PET in the initial management of PCa are summarised in Table 1.

3.3. Restaging prostate cancer with PSMA-PET during BCR: distribution of metastatic sites

BCR occurs in up to 50% of PCa patients within 10 yr from radical therapy and is defined as a rise in PSA levels (>0.2 ng/ml after radical prostatectomy and >2 ng/ml above the nadir after radiotherapy). The EAU guidelines suggest the use of PSMA-PET in all proven cases of BCR if the results are likely to influence the treatment strategy, thus letting the clinicians take the final decision on whether to refer the patient to PSMA-PET or not in case of recurrence [25]. In this setting, PSMA-PET can address subsequent treatment (eg, MDTs) to delay the initiation of ADT. Several prospective registry studies [26] confirmed the superiority of PSMA-PET imaging to conventional imaging and other PET tracers (eg, choline or

fluciclovine). Furthermore, PSMA-PET generally holds higher inter-reader agreement than other diagnostic procedures [3], and it is currently available a structured reporting system (E-PSMA) to improve its repeatability and reproducibility among different centres [1]. In this critical review, we considered studies exploring cohorts of biochemically recurrent patients. In the selected studies, the median PSA value at the time of the scan ranged from 0.32 to 2.1 ng/ml, and the median PSA doubling time (PSAdt) ranged from 4 to 11.18 mo. Metastasis detection at any site (M1) ranged from 8.8% to 40%, and the most frequent metastatic site was the bone: M1b stage was reported in 7.8% to 18% of patients analysed. The results presented in these studies have been summarised in Table 2 and Figure 2. In this scenario, by detecting more lesions in earlier stages, PSMA-PET has the potential to improve patients' outcome. However, information derived by randomised control trials, specifically designed to assess the impact of PSMA-PET on survival surrogate endpoints (eg, BCR-free survival), is still pending [5].

A prospective, single-arm, registry clinical trial assessing the accuracy of PSMA-PET in localising recurrent PCa has been published by Fendler et al. [27]. Considering a subpopulation of 223 BCR patients who had a lesion validation, authors observed different PSMA-PET performance at different PSA level thresholds, resulting in a positivity rate of 38% in patients with PSA <0.5 ng/ml and 97% in patients with PSA >5 ng/ml ($p < 0.001$). The PPV value was 84% (95% CI 0.75–0.90), with most false positive findings detected in the prostate bed. The sensitivity was 92% (95% CI 0.84–0.94) on a per-patient basis and 90% (95% CI 0.82–0.95) on a per-lesion basis, and extrapelvic disease was detected in 40% of patients. Another prospective multicentre collaboration coordinated by the International Agency for Atomic Energy (IAEA) [28] investigated the PSMA-PET positivity rate in a large cohort of patients ($n = 1004$). The main findings were the following: positivity rates stratified for a Gleason score were 61% for International Society of Urological Pathology (ISUP) grade score ≤ 3 and ranging from 66% to 87% for ISUP grade score ≥ 4 ; positivity rates stratified according to anatomical regions were 21% in pelvic LN (N1) and 27% in any metastatic sites (M1a, b, c); overall positivity rate was positively influenced by higher PSA levels, a shorter PSAdt, a higher Gleason score, and radiotherapy as the primary treatment; and disease management changed in 56.8% of cases, suggesting a very promising potential of a PSMA-guided salvage therapy. These results were consistent with the findings of other studies on this topic. Ceci et al. [29] demonstrated that PSMA-PET was able to identify the site of recurrence in 64% of patients with PSA persistence after radical prostatectomy and in 45% of patients with first-time BCR after radical prostatectomy, of whom 90% were oligorecurrent (five or fewer lesions) and ideal candidates for salvage therapy. Moreover, these results suggested that many patients were not identified as metastatic at initial staging, especially in case of persistent PSA after surgery, thus confirming the leading role of PSMA-PET in high-risk PCa prior to primary therapy. In a large retrospective international multicentre analysis, Bianchi et al. [30] validated a clinical-based nomogram developed to predict the PSMA-PET positivity rate in the recurrent setting [31]. The authors confirmed that the original nomogram retained excellent performance characteristics (AUC = 0.82) in the external validation, and PSA, PSAdt, and the clinical setting (clinical indication to request the PET scan) were the most important predictors.

These results have also been confirmed by other prospective studies [32], and the importance of PSMA-PET scans performed in early stages to detect oligorecurrent and/or oligometastatic disease has been highlighted as well. Finally, these studies highlighted that

the presence of metastatic lesions (M1a, M1b, or M1c) can be detected in up to 20–30% of patients presenting with early recurrence, namely, in patients with first-time BCR or PSA persistence after primary treatment.

3.4. Restaging prostate cancer with PSMA-PET during BCR: impact on clinical management

The decision to offer additional therapies in recurrent PCa is challenging. The rate of stage migration derived from the use of PSMA-PET in this setting generates important consequences on patients' management, and the proper patient selection is essential to ensure favourable outcomes. Men with recurrent/persistent disease reflect different clinical settings and represent a highly heterogeneous population, carrying different prognoses and different profiles of disease aggressiveness. Therefore, selection of the most suitable candidates for additional therapies is crucial. Different sites of recurrence can be identified by PSMA-PET, even if its diagnostic accuracy is still suboptimal in case of less aggressive recurrence at low PSA levels [29]. Clinical management changes are frequent, and at least half of patients will change the intended planned treatment after a PSMA-PET scan [33]. EAU recommends performing PSMA-PET in case of PSA recurrence when the scan influences the treatment decisions. However, patients identified with oligometastatic disease at PSMA-PET during the first BCR can be treated with MTD or ADT, without definitive evidence in favour of the PSMA-guided approach.

In the CONDOR trial involving 208 patients with BCR and negative standard imaging, PSMA-PET was able to change the intended management in the 64% of evaluated patients [34]. Similarly, Farolfi et al. [35] explored a cohort of patients at a very early stage of recurrence (PSA <0.5 ng/ml) and observed a change in the intended treatment in one-third of patients. These results further support the hypothesis that PSMA-PET is a valid procedure in the management of patients with recurrent PCa and low PSA levels after primary treatment, and support the implementation of this imaging procedure in the routine clinical practice [8]. In the very early stage of recurrence, PSMA-PET may allow personalised salvage radiotherapy (SRT) plans, adapting the irradiated volumes and thus leading to a potential individualised therapy [35]. In these patients, a change in the planned treatment volumes can be observed in up to one-third of patients, with a potential positive impact on progression-free survival [36]. Detection of the site of recurrence is crucial for successful treatment planning. In the event of an exclusive locoregional recurrence, long-term ADT could be avoided or at least delayed by SRT or salvage LN dissection [37]. However, a negative PSMA-PET scan should not delay the early administration of SRT, as the sensitivity of PSMA-PET for the detection of micrometastases is low and, therefore, early SRT should be offered anyway [38]. Conversely, in case of distant metastases detected, MDT as well as androgen receptor targeted therapy (abiraterone, enzalutamide, apalutamide, or darolutamide) should be considered instead [39].

During the first BCR, SRT is the treatment of choice in most patients. In a retrospective international multicentre study, Calais et al. [40] demonstrated that PSMA-PET had a major impact on a significant number of patients investigated prior to the radiotherapy planning, as 20% of patients had at least one PSMA-PET-positive lesion not covered by the clinical target volume. A randomised, controlled clinical trial investigating the outcomes of patients undergoing SRT with or without planning based on PSMA-PET findings is currently ongoing [41].

Finally, the overall impact of PSMA-PET on the oncological outcomes of patients who received this new-generation imaging in case of disease recurrence is still to be determined. Interestingly, in a retrospective study by Wenzel et al. [42], PSMA-PET locoregional positivity

has been associated, at 5 yr of follow-up, with shorter metastasis-free survival (MFS) in a group of 155 patients who underwent the scan prior to SRT, while PSMA-PET negativity has shown comparable MFS to patients who had not performed the scan. Moreover, in the Cox-regression modelling, positive PSMA-PET has been confirmed as an independent predictor of unfavourable MFS. Similar results have recently been published by Ceci et al. [43]. The authors reported prospective data about the incidence of clinically relevant events during follow-up in patients who performed PSMA-PET for PSA relapse after radical treatment, using the event-free survival (EFS) as the primary endpoint median (median follow-up of 35.4 [interquartile range: 26.5–40.3] mo). Low PSA and long PSAdt were significant predictors of EFS. Furthermore, a lower incidence of events was also observed in patients having negative PSMA-PET, since longer EFS was significantly more probable in case of a negative scan (hazard ratio 1.53; 95% CI 0.91–2.55; $p = 0.108$).

These findings suggest that PSMA-PET can be used as prognostic biomarkers as well, helping to identify patients at a higher versus lower risk of disease relapse and leading to cost-effective management of patients in early stages of disease recurrence. The results presented in this section about metastatic site location have been summarised graphically in Table 2 and Figure 2.

4. Conclusions

PSMA-PET detects a higher number of lesions than conventional imaging or other PET radiopharmaceuticals. Bone is the most frequent site of metastatic spread (M1b) followed by extrapelvic LNs (M1a), both in primary PCa and during disease recurrence. Although visceral involvement is reported with a lower incidence, it is still higher than that of conventional imaging, with a significant impact on patient's management as non-nodal visceral involvement is associated with a poorer outcome.

The high diagnostic accuracy of PSMA-PET generally leads to patients' upstaging and generates a migration of patients towards different clinical settings (nonmetastatic hormone-sensitive prostate cancer [HSPC] to metastatic HSPC or nonmetastatic castration-resistant prostate cancer [CRPC] to metastatic CRPC). Overall, approximately half of patients who performed PSMA-PET will change their therapeutic strategy according to new-generation imaging results. In case of positive PSMA-PET during recurrence, a personalised therapy approach can be adopted, with MDT generally preferred over ADT only in case of oligometastatic disease.

Finally, the clinical significance of this high diagnostic accuracy is still debated as data regarding cancer-specific mortality are still awaited. Considering the presence of new drugs able to improve patients' survival in the nonmetastatic setting (according to CT or BS), data from on-going phase III randomised controlled trials are crucial to understand whether the PSMA-guided approach holds significance in delaying the castration-resistant condition and in improving patients' overall survival.

References

[1]

F. Ceci, D.E. Oprea-Lager, L. Emmett, *et al.*
E-PSMA: the EANM standardized reporting guidelines v1.0 for PSMA-PET
Eur J Nucl Med Mol Imaging, 48 (2021), pp. 1626-1638

[2]

EAU. EAU guidelines. Presented at the EAU Annual Congress Amsterdam; 2022.

[3]

M.S. Hofman, N. Lawrentschuk, R.J. Francis, *et al.*
Prostate-specific membrane antigen PET-CT in patients with high-risk prostate cancer before curative-intent surgery or radiotherapy (proPSMA): a prospective, randomised, multicentre study
Lancet, 395 (2020), pp. 1208-1216

[4]

L. Emmett, J. Buteau, N. Papa, *et al.*
The additive diagnostic value of prostate-specific membrane antigen positron emission tomography computed tomography to multiparametric magnetic resonance imaging triage in the diagnosis of prostate cancer (PRIMARY): a prospective multicentre study
Eur Urol, 80 (2021), pp. 682-689

[5]

J. Calais, W.R. Armstrong, A.U. Kishan, *et al.*
Update from PSMA-SRT trial NCT03582774: a randomized phase 3 imaging trial of prostate-specific membrane antigen positron emission tomography for salvage radiation therapy for prostate cancer recurrence powered for clinical outcome
Eur Urol Focus, 7 (2021), pp. 238-240

[6]

S. Klingenberg, M.R. Jochumsen, B.P. Uhløi, *et al.*
⁶⁸Ga-PSMA PET/CT for primary lymph node and distant metastasis NM staging of high-risk prostate cancer
J Nucl Med, 62 (2021), pp. 214-220

[7]

K.J. Pienta, M.A. Gorin, S.P. Rowe, *et al.*
A phase 2/3 prospective multicenter study of the diagnostic accuracy of prostate specific membrane antigen PET/CT with ¹⁸F-DCFPyL in prostate cancer patients (OSPREY)
J Urol, 206 (2021), pp. 52-61

[8]

M.J. Morris, S.P. Rowe, M.A. Gorin, *et al.*
Diagnostic performance of ¹⁸F-DCFPyL-PET/CT in men with biochemically recurrent prostate cancer: results from the CONDOR phase III, multicenter study
Clin Cancer Res, 27 (2021), pp. 3674-3682

[9]

T.A. Hope, M. Eiber, W.R. Armstrong, *et al.*
Diagnostic accuracy of ⁶⁸Ga-PSMA-11 PET for pelvic nodal metastasis detection prior to radical prostatectomy and pelvic lymph node dissection: a multicenter prospective phase 3 imaging trial
JAMA Oncol, 7 (2021), pp. 1635-1642

[10]

L.J. Petersen, H.D. Zacho

PSMA PET for primary lymph node staging of intermediate and high-risk prostate cancer: an expedited systematic review

Cancer Imaging, 20 (2020), p. 10

[11]

A.W. Rutjes, J.B. Reitsma, M. Di Nisio, N. Smidt, J.C. van Rijn, P.M. Bossuyt

Evidence of bias and variation in diagnostic accuracy studies

CMAJ, 174 (2006), pp. 469-476

[12]

L.J. Petersen, M.N. Johansen, J. Strandberg, L. Stenholt, H.D. Zacho

Reporting and handling of equivocal imaging findings in diagnostic studies of bone metastasis in prostate cancer

Acta Radiol, 61 (2020), pp. 1096-1104

[13]

M. Roscigno, M. Nicolai, G. La Croce, *et al.*

Difference in frequency and distribution of nodal metastases between intermediate and high risk prostate cancer patients: results of a superextended pelvic lymph node dissection

Front Surg, 5 (2018), p. 52

[14]

J.W. Yaxley, S. Raveenthiran, F.X. Nouhaud, *et al.*

Outcomes of primary lymph node staging of intermediate and high risk prostate cancer with ⁶⁸Ga-PSMA positron emission tomography/computerized tomography compared to histological correlation of pelvic lymph node pathology

J Urol, 201 (2019), pp. 815-820

[15]

M. Kulkarni, S. Hughes, A. Mallia, *et al.*

The management impact of ⁶⁸gallium-tris(hydroxypyridinone) prostate-specific membrane antigen (⁶⁸Ga-THP-PSMA) PET-CT imaging for high-risk and biochemically recurrent prostate cancer

Eur J Nucl Med Mol Imaging, 47 (2020), pp. 674-686

[16]

F. Ceci, G. Musi, O. De Cobelli

Prostate-specific membrane antigen positron emission tomography, not conventional imaging, should be performed for primary staging of high-risk prostate cancer

Eur Urol Open Sci, 34 (2021), pp. 17-18

[17]

B. Grubmüller, P. Baltzer, S. Hartenbach, *et al.*

PSMA Ligand PET/MRI for primary prostate cancer: staging performance and clinical impact

Clin Cancer Res, 24 (2018), pp. 6300-6307

[18]

V. Karagiannis, V. Wichmann, J. Saarinen, N. Eigeliene, H. Andersen, A. Jekunen

Radiotherapy treatment modification for prostate cancer patients based on PSMA-PET/CT

Radiat Oncol, 17 (2022), p. 19

[19]

D.A. Ferraro, F. Lehner, A.S. Becker, *et al.*

Improved oncological outcome after radical prostatectomy in patients staged with (68)Ga-PSMA-11 PET: a single-center retrospective cohort comparison

Eur J Nucl Med Mol Imaging, 48 (2021), pp. 1219-1228

[20]

M. Chen, Q. Zhang, C. Zhang, *et al.*

Comparison of (68)Ga-prostate-specific membrane antigen (PSMA) positron emission tomography/computed tomography (PET/CT) and multi-parametric magnetic resonance imaging (MRI) in the evaluation of tumor extension of primary prostate cancer

Transl Androl Urol, 9 (2020), pp. 382-390

[21]

U.J. Muehlematter, I.A. Burger, A.S. Becker, *et al.*

Diagnostic accuracy of multiparametric MRI versus (68)Ga-PSMA-11 PET/MRI for extracapsular extension and seminal vesicle invasion in patients with prostate cancer

Radiology, 293 (2019), pp. 350-358

[22]

J. Calais, A.U. Kishan, M. Cao, *et al.*

Potential impact of 68Ga-PSMA-11 PET/CT on the planning of definitive radiation therapy for prostate cancer

J Nucl Med, 59 (2018), pp. 1714-1721

[23]

P. Rogowski, C. Trapp, R. von Bestenbostel, *et al.*

Outcomes of metastasis-directed therapy of bone oligometastatic prostate cancer

Radiat Oncol, 16 (2021), p. 125

[24]

M.J. Connor, A. Smith, S. Miah, *et al.*

Targeting oligometastasis with stereotactic ablative radiation therapy or surgery in metastatic hormone-sensitive prostate cancer: a systematic review of prospective clinical trials

Eur Urol Oncol, 3 (2020), pp. 582-593

[25]

N. Mottet, R.C.N. van den Bergh, E. Briers, *et al.*

EAU-EANM-ESTRO-ESUR-SIOG guidelines on prostate cancer—2020 update. Part 1: screening, diagnosis, and local treatment with curative intent

Eur Urol, 79 (2021), pp. 243-262

[26]

J. Calais, F. Ceci, M. Eiber, *et al.*

¹⁸F-fluciclovine PET-CT and ⁶⁸Ga-PSMA-11 PET-CT in patients with early biochemical recurrence after prostatectomy: a prospective, single-centre, single-arm, comparative imaging trial [published correction appears in Lancet Oncol. 2019 Nov; 20(11):e613; published correction appears in Lancet Oncol. 2020 Jun; 21(6):e304]

Lancet Oncol, 20 (2019), pp. 1286-1294

[27]

W.P. Fendler, J. Calais, M. Eiber, *et al.*

Assessment of 68Ga-PSMA-11 PET accuracy in localizing recurrent prostate cancer: a prospective single-arm clinical trial

JAMA Oncol, 5 (2019), pp. 856-863

[28]

J.J. Cerci, S. Fanti, E.E. Lobato, *et al.*

Diagnostic performance and clinical impact of ⁶⁸Ga-PSMA-11 PET/CT imaging in early relapsed prostate cancer after radical therapy: a prospective multicenter study (IAEA-PSMA Study)

J Nucl Med, 63 (2022), pp. 240-247

[29]

F. Ceci, P. Castellucci, T. Graziani, *et al.*

⁶⁸Ga-PSMA-11 PET/CT in recurrent prostate cancer: efficacy in different clinical stages of PSA failure after radical therapy

Eur J Nucl Med Mol Imaging, 46 (2019), pp. 31-39

[30]

Bianchi L, Castellucci P, Farolfi A, *et al.* Multicenter external validation of a nomogram for predicting positive prostate-specific membrane antigen/positron emission tomography scan in patients with prostate cancer recurrence. Eur Urol Oncol. In press. <https://doi.org/10.1016/j.euo.2021.12.002>.

[31]

F. Ceci, L. Bianchi, M. Borghesi, *et al.*

Prediction nomogram for ⁶⁸Ga-PSMA-11 PET/CT in different clinical settings of PSA failure after radical treatment for prostate cancer

Eur J Nucl Med Mol Imaging, 47 (2020), pp. 136-146

[32]

D. Deandreis, A. Guarneri, F. Ceci, *et al.*

⁶⁸Ga-PSMA-11 PET/CT in recurrent hormone-sensitive prostate cancer (HSPC): a prospective single-centre study in patients eligible for salvage therapy

Eur J Nucl Med Mol Imaging, 47 (2020), pp. 2804-2815

[33]

A. Eissa, A. Elsherbiny, R.F. Coelho, *et al.*

The role of ⁶⁸Ga-PSMA PET/CT scan in biochemical recurrence after primary treatment for prostate cancer: a systematic review of the literature

Minerva Urol Nefrol, 70 (2018), pp. 462-478

[34]

W.P. Fendler, J. Ferdinandus, J. Czernin, *et al.*

Impact of ⁶⁸Ga-PSMA-11 PET on the management of recurrent prostate cancer in a prospective single-arm clinical trial

J Nucl Med, 61 (2020), pp. 1793-1799

[35]

A. Farolfi, F. Ceci, P. Castellucci, *et al.*

⁶⁸Ga-PSMA-11 PET/CT in prostate cancer patients with biochemical recurrence after radical prostatectomy and PSA <0.5 ng/ml. Efficacy and impact on treatment strategy

Eur J Nucl Med Mol Imaging, 46 (2019), pp. 11-19

[36]

N.S. Schmidt-Hegemann, W.P. Fendler, H. Ilhan, *et al.*

Outcome after PSMA PET/CT based radiotherapy in patients with biochemical persistence or recurrence after radical prostatectomy

Radiat Oncol, 13 (2018), p. 37

[37]

D. Bottke, J. Miksch, R. Thamm, *et al.*

Changes of radiation treatment concept based on ⁶⁸Ga-PSMA-11-PET/CT in Early PSA-recurrences after radical prostatectomy

Front Oncol, 11 (2021), Article 665304

[38]

G. Ploussard, G. Gandaglia, H. Borgmann, *et al.*

Salvage lymph node dissection for nodal recurrent prostate cancer: a systematic review

Eur Urol, 76 (2019), pp. 493-504

[39]

F. Zattoni, I. Heidegger, V. Kasivisvanathan, *et al.*

Radiation therapy after radical prostatectomy: what has changed over time?

Front Surg, 8 (2021), Article 691473

[40]

J. Calais, J. Czernin, M. Cao, *et al.*

⁶⁸Ga-PSMA-11 PET/CT mapping of prostate cancer biochemical recurrence after radical prostatectomy in 270 patients with a PSA level of less than 1.0 ng/ml: impact on salvage radiotherapy planning

J Nucl Med, 59 (2018), pp. 230-237

[41]

J. Calais, J. Czernin, W.P. Fendler, D. Elashoff, N.G. Nickols

Randomized prospective phase III trial of ⁶⁸Ga-PSMA-11 PET/CT molecular imaging for prostate cancer salvage radiotherapy planning [PSMA-SRT] [published correction appears in BMC Cancer. 2019 Jan 21;19(1):97]

BMC Cancer, 19 (2019), p. 18

[42]

M. Wenzel, R. Hussein, T. Maurer, *et al.*

PSMA PET predicts metastasis-free survival in the setting of salvage radiotherapy after radical prostatectomy

Urol Oncol, 40 (2022), pp. 7.e1-7.e8

[43]

F. Ceci, G. Rovera, G.C. Iorio, *et al.*

Event-free survival after ⁶⁸Ga-PSMA-11 PET/CT in recurrent hormone-sensitive prostate cancer (HSPC) patients eligible for salvage therapy [published online ahead of print, 2022 Feb 26]

Eur J Nucl Med Mol Imaging, 49 (2022), pp. 3257-3268