

# **Examining readmission factors in psychiatric emergency care for individuals with personality disorders: a 6-year retrospective study**

## **Abstract**

People with personality disorders (PD) are often admitted to psychiatric emergency services due to the frequent repetition of acute crises. This study drew on the ICD diagnostic records of 2634 individuals with personality disorders who were admitted in a specialized inpatient psychiatric crisis unit over a 6-year period. Multiple logistic regressions and survival regressions were performed to examine whether personality disorder categories, gender, and other individual, interpersonal and precipitating factors were associated with readmission and time-to-readmission. The results showed a 16.1% readmission rate. Of these, 99.5% of readmissions occurred within 4 years following first admission. Gender was the main factor associated with both readmission and time-to-readmission : while men were readmitted faster, more women in total were readmitted for a second psychiatric emergency hospitalization. Findings also indicated that readmission rate and time-to-readmission differed following the category of personality disorder : readmission rate in a ratio of 1 to 2 (from 8 and 10% for dissocial and paranoid PD up to 19 to 21% for impulsive and borderline PD), and time-to-readmission in a ratio of 1 to 5 (from one month for anankastic and dependent, to 5 months for impulsive, histrionic and anxious-avoidant PD). Limitations of this naturalistic study include a lack of self-reported measures and generalizability to less specialized emergency settings. Future research should include a prospective longitudinal design using standardized scalable measurement tools to improve the completeness and accuracy of the data concerning the psychological processes involved in risk and time-to-readmission after brief hospitalizations in emergency psychiatry.

**Keywords:** personality disorder, mental health crisis, psychiatric emergency hospitalization, readmission, resilience.

## **1. Introduction**

The term 'personality disorder' (PD) refers to a diversity of symptoms presented by people who experience difficulties in forming and maintaining satisfying interpersonal relationships. Personality disorders are thought to reflect personal distress and psychosocial dysfunction that can have a serious impact on sufferers' well-being and health, in particular due to recurrent crises and a high frequency of self-harming behavior that increase the risk of suicide (Chan et al., 2016; Flynn et al., 2020; Grenyer et al., 2018; Tyrer et al., 2015). With an onset in adolescence and young adulthood, personality disorders are assumed to be chronic disorders, and their overall prevalence is estimated at 7.8% internationally (Winsper et al., 2020). As the presentation of symptoms in people with PD fluctuates over time, it is difficult for health practitioners to diagnose and care for these individuals, and individuals with PD are frequent users of the health care system, with prevalence estimated at 25% in primary care and 50% in outpatient psychiatry (Tyrer et al., 2015). Regarding inpatient care, a diagnosis of PD would increase by 2.75 the risk of admission in the general hospital (Fok et al., 2019), while prevalence of PD among psychiatric inpatients is estimated at 40 to 50% (Grenyer et al., 2018). Research also indicates that PD is associated with higher risk for readmission, with a risk ratio compared to other mental disorders ranging from 1.55 at one month to 8.7 at 36 months (Evans et al., 2017; Lewis et al., 2019). Individuals with PD are also more likely to be readmitted in shorter time than those with other mental disorders with a median time-to-readmission reduced by 1.53, and a readmission rate at one month that is 2.3 times higher than for other psychiatric disorders (Lewis et al., 2019; Lin et al., 2007). Consistent with both higher risk and shorter time-to-readmission, PD appears to be the only psychiatric diagnosis associated with rapid readmission in psychiatric emergency settings (Barr et al., 2020; Chang et al., 2014; Evans et al., 2017; Richard-Lepouriel et al., 2015; Tulloch et al., 2016). However, nosographies distinguish about ten different categories of personality disorder historically grouped into 3

clusters (Tyrer et al., 2015), and it is not clear whether the risk for and time-to-readmission differ between these categories. In fact, most of the studies focus on the borderline PD, or tend to assimilate borderline PD to the personality disorders in general. Furthermore, literature indicates that gender has a significant impact on both the symptomatology and the clinical presentation of individuals with PD, and that men and women suffering of a PD experience different behavioral, social and medical outcomes (Grant, 2009; Sansone & Sansone, 2011; Schulte Holthausen & Habel, 2018). For example, 77% of men in prison were found to have a PD (Slade & Forrester, 2013) while 59% of psychiatric inpatients with a PD would be women (Björkenstam et al., 2015). Beside personality disorders, published findings indicate that psychiatric emergency readmission is associated with numerous factors : psychiatric and somatic comorbidities (Agnafors et al., 2019; Gili et al., 2010; Golay et al., 2019; Iacovides & Siamouli, 2008; Lewis et al., 2019), self-harming behavior (Carroll et al., 2014; Gunnell et al., 2008), substance use (Degenhardt & Hall, 2001; Puddephatt et al., 2021), interpersonal and social factors (Bentley et al., 2019; Beutel et al., 2017; Donisi et al., 2016; Dube & Rishi, 2017; Lakey & Orehek, 2011; Mushtaq, 2014; Pernice-Duca, 2010; Rajani et al., 2016; Schmutte et al., 2010; Scott et al., 2014; Wang et al., 2018; Wilhelm et al., 2004), and precipitating factors (Gårdvik et al., 2021; James, 2016; Kaplan et al., 2012; Reiland & Clark, 2017; Wamser-Nanney et al., 2018). The literature usually only examines one or two factors at a time, making findings heterogeneous and poorly consistent (Donisi et al., 2016).

In this context, the aim of this study is to examine whether and how readmission and time-to-readmission to psychiatric emergency vary by category of personality disorder and gender, while considering other individual, interpersonal/social or precipitating factors known to be associated with this risk.

Based on published findings, it was hypothesized that the risk for readmission would be higher for women and young adults compared to men and middle-aged adults (Clements et al., 2006;

Gunnell et al., 2008; Polling et al., 2021; Tulloch et al., 2016). Considering PD categories as per ICD, it was hypothesized that anxious-avoidant, dependent, and anankastic categories would be associated with more readmission than the impulsive, histrionic and dissocial ones, while schizoid and paranoid with less readmissions (Fok et al., 2019). Borderline PD was expected to be both the most prevalent and with the highest risk for readmission (Lewis et al., 2019). Concerning time-to-readmission, it was expected that male gender would be associated with a decrease in time-to-readmission although in proportion that might not be statistically significant (Evans et al., 2017; Olfson et al., 2013; Tulloch et al., 2016). Psychiatric and somatic comorbidities were expected to increase the risk of readmission (Fok et al., 2019; Gunnell et al., 2008; Rosca et al., 2006), together with self-harming behavior (Flynn et al., 2020; Gunnell et al., 2008; Olfson et al., 2013; Perlman et al., 2015) and substance use, especially alcohol (Chang et al., 2014; Kaplan et al., 2012; Morel et al., 2020). Interpersonal and social factors were expected to be related with higher risk of readmission (Donisi et al., 2016; Evans et al., 2017; Rosca et al., 2006).

This study uses data collected during patients hospitalizations. Following the French regulation “Code de la santé publique – article L1121-1”, patients have been informed that their data will be used for medical research purposes, in a strictly controlled manner, and in compliance with the French General Data Protection Regulation (RGPD) and Data Protection Act (LIL). Patients were informed about their rights regarding their data. Patients who objected to the use of their data were not included in the study.

## **2. Materials and Methods**

### ***2.1 Study population***

This study draws from the medical records of individuals with a diagnosis of personality disorder who were admitted or readmitted from 1<sup>st</sup> January 2015 to 31<sup>st</sup> December 2020 at the


“Unité Psychiatrique de Crise” (UPC), a specialized psychiatric crisis unit located within the main Lyon university hospital, and covering the greater Lyon catchment area of 1,600,000 people. Patients admitted at UPC require urgent psychiatric inpatient care. They are referred by the psychiatric emergency departments of the other hospitals of the catchment area, by the primary health care system, by mobile emergency teams, or present themselves spontaneously. There are four criteria for admission at UPC : 1. acute crisis state in the context of personality disorders, anxiety disorders, depression or substance use disorders ; 2. suicidal crisis ; 3. post-traumatic crisis ; 4. emerging psychotic spectrum disorders only in individuals under 25 years of age. Therefore, this includes patients in crisis with known personality disorders as well as first-time admitted patients without a previous psychiatric diagnosis. In both undiagnosed and previously diagnosed cases, the diagnosis is established, respectively confirmed at the end of the hospitalization by the specialized medical team by Mini 5.0 or SCID-2, similarly to other practices in different countries (Bach et al., 2015; Buer Christensen et al., 2018). As this study covers all categories of personality disorder, all hospitalizations records of UPC’s patients with a personality disorder diagnosis were included. Individuals with known diagnosis of bipolar, psychotic and autism spectrum disorders are not admitted to UPC but referred to specialized institutions. Other exclusion criteria for admission at UPC relate to the presence of severe delusion symptoms, to high impulsivity with repetition of auto- or hetero-aggressivity, to manic state, and to the inability to give a clear consent for hospitalization. The intended length of stay at UPC is 4 days, and the purpose of the intervention is threefold : 1. Secure and stabilize the patient, 2. Evaluate clinical state and potential diagnosis(es), 3. Refer him/her to the further appropriate inpatient or outpatient treatment.

## ***2.2 Variables***

In addition to gender, age and personality disorders, other individual, interpersonal/social and precipitating factors assumed to be associated with readmission were collected from the ICD

codes of the patients' hospitalization records as independent variables. The outcome variables of brief psychiatric hospitalization, namely readmission and time-to-readmission, were calculated independently of the ICD codes (figure 1).

**Figure 1 :** ICD-based factors (left) assumed to be associated with readmission and time-to-readmission.



Hospitalization records provided following data : patient permanent identification code, gender, birth date, entry date, discharge date, destination of discharge, ICD codes of the diagnosed disorder(s) and up to twenty additional ICD codes. The additional ICD codes are used by trained psychiatrists who assess patients to describe their condition in terms of somatic and mental health but also in terms of private and social situation, and recent negative life events.

Factors were managed as categorical variables, i.e. for a given patient, each factor may possibly be valued among a defined list of categories (see table 1 for a summary of factors and categories). A simple example is the factor “personality disorder” which may be valued with one or several of the categories of personality disorder. ICD codes from hospitalization records served to populate categories, e.g. categories of personality disorder were directly populated from ICD codes F60.x. Categories of psychiatric comorbidity were matched with sub-sections of ICD chapter V (e.g. all F40.xx and F41.xx codes were grouped under the category "anxiety/phobia") with an exception of F1x.x codes related to substance use because “substance use” was managed as an independent factor. Categories of the factor “somatic comorbidity” were matched with ICD chapters related to somatic diseases (Fok et al., 2019), etc. (see

supplemental material 1 for correspondence rules, and details from all ICD-codes collected to categories and factors). When a patient record reported several ICD codes for the same category, those codes were grouped into this category. For example, a patient record indicated codes F10.01, F10.21, F10.240, F10.30, F14.05, F17.22, and F17.25, which respectively correspond to several categories of substance use : F10.xx to alcohol, F14.05 to cocaine, and F17.2x to tobacco. Thus, for this patient, the factor “substance use” was valued with the 3 categories “alcohol”, “tobacco” and “other substance”. Then, numerical variables were calculated to indicate how many categories were present for each factor. For the patient in above example, the numerical variable for the factor “Substance use” was equal to 3. These numerical variables are meaningful to weight the severity of the factors and are useful to perform multiple logistic regressions.

Readmission was measured by searching for patients with multiple hospitalization records based on their permanent identification code. These patients were identified with a binary variable “readmission” set at 1, and the time-to-readmission was calculated as the time elapsed between the discharge date of one hospitalization and the entry date of the successive one.

### ***2.3 Statistical analysis***

Three steps of analyses were planned : 1. Description of the sample ; 2. Analyses of the risk for readmission ; 3. Analyses of the time to readmission. First, it was aimed to describe the sample by the following characteristics : distribution of patients per number of admissions ; distribution of admissions and readmission rate as a function of age and gender ; prevalence of factors and categories. The second step of analyses aimed to investigate if and how readmission varied with factors and categories. Chi-square tests were planned to identify which factors, respectively categories, would be associated with readmission. Odds ratios at 95% confidence intervals were planned to quantify how factors and categories would be associated with readmission. Multiple logistic regressions were planned to model how factors, respectively



categories, would combine to influence the risk of readmission. The third step of analyses aimed to focus on the time-to-readmission. It was planned to display the distribution of the number of readmissions according to the time-to-readmission, and to check if this distribution varied with gender. Non-parametric Median and Kruskal-Wallis tests were planned to examine whether the time-to-readmission varied by gender, personality disorder categories and factors. Survival regressions were planned to model associations between factors, respectively categories, and time-to-readmission : a Cox regression to identify factors possibly associated with time-to-readmission, then a Kaplan-Meier regression to assess the effect of gender on the time-to-readmission.

Considering the 7 factors mentioned above plus gender and age, and assuming an effect size of 0.01 for each (Rosca et al., 2006), a statistical power of 80% at probability level of 0.05 requires a sample size of 1566 patients. Following previous research (Lewis et al., 2019), time-to-readmission may last 4 to 5 years. We thus retrospectively sampled patients with personality disorders admitted at UPC over a 6-year period (actual sample 2634).

Descriptive and statistical analysis employed Excel and SPSS 25.

### **3. Results**

#### ***3.1 Description of the study sample***

The selection of all patients with diagnosis of PD admitted at UPC from 1<sup>st</sup> January 2015 to 31<sup>st</sup> December 2020 provided 3289 hospitalization records. Due to patients with multiple admissions, those 3289 records corresponded to 2634 individuals, among whom 2209 were admitted once and 425 several times, which gives a readmission rate of 16.1%. The distribution of the number of patients according to their number of admissions and per gender is displayed on figure 2. The sex-ratio observed in the study sample was 2 women for 1 man (1770 for 864).

**Figure 2 :** Distribution of patients by number of admissions during the 6-year study period, per gender.




Figure 3 gives the number of admissions and readmission rate by age range, for women and men. It shows that readmission rate is higher for women and that, over the course of lifetime, it decreases for women and peaks around 40 for men.

**Figure 3 :** number of admissions (left scale) and readmission rate (right scale) by age range, for women and men.




Table 1 summarizes the factors and categories observed in the study sample, and indicates the size and prevalence for each sub-group.

**Table 1 :** Number of individuals in sub-groups by factors and categories, and prevalence observed in the study sample (note that n's by categories do not add to n's by factors due to comorbidity within factors).

FACTORS / categories	n	preval.	FACTORS / categories	n	preval.	FACTORS / categories	n	preval.
<b>PERSONALITY DISORDER</b>	<b>2634</b>	<b>100,0%</b>	<b>PSYCHIATRIC COMORBIDITY</b>	<b>2172</b>	<b>82,5%</b>	<b>SOMATIC COMORBIDITY</b>	<b>1037</b>	<b>39,4%</b>
Anankastic	282	10,7%	Anxiety/phobia	416	15,8%	Articul., muscul., skeleton	65	2,5%
Anxious-avoidant	102	3,9%	Autism	2	0,1%	Blood & immunity	37	1,4%
Borderline	817	31,0%	Bipolar	102	3,9%	Circulatory	112	4,3%
Dependent	234	8,9%	Conduct disorder	197	7,5%	Digestive	58	2,2%
Dissocial	92	3,5%	Depression	1457	55,3%	Ear	15	0,5%
Histrionic	354	13,4%	Developmental	11	0,4%	Endo., nutri., metabolism	553	20,9%
Impulsive	481	18,3%	Dissociative	28	1,1%	Eye	13	0,5%
Other/Unspecified	161	6,1%	Eating	181	6,9%	Factors motivating care	336	12,8%
Paranoid	102	3,9%	Neurodegenerative	16	0,6%	Genito-urinary	51	1,9%
Schizoid	77	2,9%	OCD	52	2,0%	Infectious parasitary	42	1,6%
<b>INTERPERSONAL /SOCIAL FACTORS</b>	<b>603</b>	<b>22,9%</b>	Other	35	1,3%	Injury	117	4,4%
Family difficulties	264	10,0%	Psychotic	82	3,1%	Neoplasms	11	0,4%
Loneliness	148	5,6%	PTSD	273	10,4%	Nervous system	105	4,0%
Precarity	175	6,6%	Somatoform	49	1,9%	Not classified	212	8,0%
Professional diffic.	80	3,0%	<b>SUBSTANCE USE</b>	<b>897</b>	<b>34,1%</b>	Respiratory	103	3,9%
Social difficulties	154	5,8%	Alcohol	594	22,6%	Skin	21	0,8%
<b>PRECIPITATING FACTORS</b>	<b>321</b>	<b>12,2%</b>	Other substance	467	17,7%	<b>SELF-HARMING BEHAVIOR</b>	<b>1012</b>	<b>38,4%</b>
Aggression	149	5,7%	Tobacco	405	15,4%	Self-injury	267	10,1%
Other negative life event	195	7,4%	Withdrawal	287	10,9%	Self-poisoning	786	29,8%

### 3.2 Factors and categories associated with readmission


Readmission rates are compared between sub-groups of patients presenting or not each factor, and odds ratio (OR) calculated. Results (table 2) indicate that only gender was significantly associated with readmission : while women showed a readmission rate of 18.3%, men's readmission rate was 11.7% ( $\chi^2(1,N=2634)=18.78, p<0.001, OR 1.69, 95\% CI 1.33-2.15$ ).

**Table 2 :** Readmission rate compared between sub-groups with and without factors for gender (women vs men), psychiatric comorbidity, somatic comorbidity, substance use, self-harming behavior, interpersonal/social factors, precipitating factors.

FACTOR (n)	Rate with factor	Rate w/o factor	$\chi^2$	$p \chi^2$	OR	95%CI
<b>Gender Women (1770) vs Men</b>	<b>18.3%</b>	<b>11.7%</b>	<b>18.78</b>	<b>0.000</b>	<b>1.69</b>	<b>1.33-2.15</b>
Psychiatric comorbidity (2172)	16.2%	15.8%	0.046	0.830	1.03	0.78-1.36
Somatic comorbidity (1037)	15.2%	16.7%	1.021	0.312	0.89	0.72-1.11
Substance use (897)	17.4%	15.5%	1.586	0.208	1.15	0.93-1.43
Self-harming behavior (1012)	16.1%	16.2%	0.001	0.975	1.00	0.81-1.23
Interpersonal/social factors (603)	15.8%	16.2%	0.084	0.772	0.96	0.75-1.24
Precipitating factors (321)	17.1%	16.0%	0.269	0.604	1.09	0.80-1.48

Considering personality disorders, Chi-square test support that readmission rate depends on the personality disorder category ( $\chi^2(9,n=2634)=35.12, p<0.001$ ), as shown on figure 4.

**Figure 4 :** number of patients with single vs multiple admissions per PD category, and readmission rate.



Controlling for gender (table 3), it appears that, for women, borderline PD is associated with more readmission ( $\chi^2(1,n=1770)=10.442, p=0.001, OR 1.49, 95\%CI 1.17-1.90$ ) while other / unspecified PDs are associated with less readmission ( $\chi^2(1, n=1770)=5.674, p=0.017, OR 0.40, 95\%CI 0.18-0.87$ ). For men, readmission is independent from the category of PD.

**Table 3 :** Readmission rate compared between sub-groups with and without PD categories, for women and men.

Women (n=1770)							
PD Category (n)	Rate with PD category	Rate w/o PD category	$\chi^2$	<i>p</i> $\chi^2$	OR	95%CI	
Anankastic (113)	13.3%	18.6%	2.043	0.153	0.67	0.38-1.17	
Anxious-av. (61)	11.3%	18.65%	2.114	0.146	0.56	0.25-1.24	
<b>Borderline (657)</b>	<b>22.1%</b>	<b>16.0%</b>	<b>10.442</b>	<b>0.001</b>	<b>1.49</b>	<b>1.17-1.90</b>	
Dependent (160)	17.2%	18.4%	0.164	0.686	0.91	0.60-1.39	
Dissocial (10)	0.0%	18.4%	2.253	0.520	NA	NA	
Histrionic (296)	15.6%	18.9%	1.770	0.183	0.80	0.57-1.11	
Impulsive (351)	21.7%	17.5%	3.280	0.070	1.30	0.98-1.74	
Paranoid (30)	13.3%	18.4%	0.504	0.478	0.68	0.24-1.97	
<b>Other/ Unspecified (65)</b>	<b>8.4%</b>	<b>18.8%</b>	<b>5.674</b>	<b>0.017</b>	<b>0.40</b>	<b>0.18-0.87</b>	
Schizoid (27)	25.0%	18.2%	0.853	0.356	1.50	0.63-3.56	
Men (n=864)							
PD Category (n)	Rate with PD category	Rate w/o PD category	$\chi^2$	<i>p</i> $\chi^2$	OR	95%CI	
Anankastic (169)	12.4%	11.5%	0.110	0.740	1.09	0.65-1.82	

Anxious-av. (37)	10.0%	11.8%	0.116	0.733	0.83	0.29-2.39
Borderline (152)	13.7%	11.3%	0.746	0.388	1.25	0.75-2.11
Dependent (63)	15.4%	11.4%	0.929	0.335	1.41	0.70-2.87
Dissocial (80)	8.5%	12.0%	0.873	0.350	0.68	0.31-1.53
Histrionic (46)	10.6%	11.8%	0.053	0.820	0.89	0.35-2.32
Impulsive (129)	13.1%	11.4%	0.285	0.590	1.16	0.67-2.03
Paranoid (72)	8.3%	12.0%	0.857	0.355	0.67	0.28-1.58
Other/ Unspecified (68)	9.0%	12.0%	0.612	0.434	0.73	0.32-1.62
Schizoid (48)	6.1%	12.0%	1.560	0.212	0.48	0.15-1.56

Concerning categories of psychiatric comorbidity (see supplemental material 2 table R1), for women somatoform disorders are associated with more readmission ( $\chi^2(1, n=1770)=9.208$ ,  $p=0.002$ , OR 2.84, 95% CI 1.41-5.73), while for men no association was found. Concerning categories of somatic comorbidity (supplemental material 2 table R2), for women, circulatory disorders are associated with less readmission ( $\chi^2(1, n=1770)=4.991$ ,  $p=0.025$ , OR 0.39, 95% CI 0.17-0.92). Concerning substance use (supplemental material 2 table R3), for women, alcohol is associated with more readmission ( $\chi^2(1, n=1770)=6.2$ ,  $p=0.013$ , OR 1.44, 95% CI 1.08-1.91). In contrast, among men, none of the categories of substance use yield significant correlation with readmission. Concerning self-harming behaviors, they appear independent of readmission both for women and men (supplemental material 2 table R4). Lastly, none of the categories of interpersonal/social and precipitating factors as assessed in this study were found to be associated with readmission, either for women or for men (supplemental material 2 tables R5 and R6).

### ***3.3 Logistic regressions for readmission***

A multiple logistic regression was performed on the factors age, gender, personality disorder, psychiatric comorbidity, somatic comorbidity, substance use, self-harming behavior, interpersonal/social and precipitating factors to investigate their contributions to the risk for readmission. Assumptions for multiple logistic regression were checked and met, namely the independence of errors, the absence of multicollinearity, the absence of significant outliers

(Stoltzfus, 2011; Tabachnick et al., 2019). Factors were entered stepwise forward, and removed by the conditional method. After 3 iterations, the model obtained is significant ( $\chi^2=31.913$  (df=3),  $p<0.001$ ) (Hosmer et al., 2013), and its values are consistent with the observed values ( $\chi^2=11.174$ , (df=8),  $p= 0.192$ ). The parameters defining this model (table 4) indicate that age (OR 0.993, 95%CI 0.986-1.000) and somatic comorbidity (OR 0.881, 95%CI 0.793-0.980) are associated with less readmission, while female gender with an increase in readmission (OR 1.686, 95%CI 1.325-2.145). Other variables are not significant.

**Table 4 :** parameters of the factors in the logistic regression for readmission.

Factor	$\beta$	Std err.	Wald	Df	Sig.	Exp( $\beta$ ) OR	95%CI
<b>Age</b>	<b>-0.007</b>	<b>0.004</b>	<b>4.355</b>	<b>1</b>	<b>0.037</b>	<b>0.993</b>	<b>0.986-1.000</b>
<b>Gender</b>	<b>0.522</b>	<b>0.123</b>	<b>18.082</b>	<b>1</b>	<b>0.000</b>	<b>1.686</b>	<b>1.325-2.145</b>
Personality disorder	2.519	-	-	1	0.113	-	-
Psychiatric comorbidity	0.026	-	-	1	0.872	-	-
<b>Somatic comorbidity</b>	<b>-0.126</b>	<b>0.054</b>	<b>5.462</b>	<b>1</b>	<b>0.019</b>	<b>0.881</b>	<b>0.793-0.980</b>
Substance use	2.078	-	-	1	0.149	-	-
Self-harming behavior	0.000	-	-	1	0.997	-	-
Interpersonal/social factors	0.247	-	-	1	0.619	-	-
Precipitating factors	0.025	-	-	1	0.873	-	-
Constant	-2.179	0.261	69.360	1	0.000	0.113	-

The initial and final Log-Likelihood of the logistic regression are respectively equal to 2327.95 and 2296.04, which gives a value of the pseudo- $R^2$  of 0.014, i.e. an estimate of the variance of the risk of readmission explained by this model of 1.4%. The Area Under the Receiver Operating Characteristic (AUROC) is equal to 0.574, indicating that this model is significant with low discriminatory power. This is consistent with previous studies underlining the large number of factors to be taken into account to model risk of psychiatric readmission, their low individual significance, and the modest strength to expect from multiple logistic regressions (Perlman et al., 2015; Tulloch et al., 2016).

A multiple logistic regression was performed on the sub-group of women on categories. After 5 runs, the model obtained is significant ( $\chi^2=40.709$  (df=5),  $p<0.001$ ) (Hosmer et al., 2013),

and its values are consistent with the observed values ( $\chi^2=2.002$  (df=4),  $p= 0.735$ ). The parameters defining this model (table 5) indicate that borderline PD (OR 1.776, 95%CI 1.34-2.35), impulsive PD (OR 1.738, 95%CI 1.25-2.41), alcohol use (OR 1.345, 95%CI 1.00-1.80), and somatoform disorders (OR 3.781, 95%CI 1.84-7.78) are associated with increased risk of readmission.

**Table 5 :** parameters of significant categories in the logistic regression in the sub-group of women (non-significant categories are given in supplemental materiel table R7).

Category	$\beta$	Std err.	Wald	Df	Sig.	Exp( $\beta$ ) OR	95%CI
Borderline Personality disorder	0.574	0.142	16.271	1	0.000	1.776	1.34-2.35
Impulsive Personality disorder	0.553	0.168	10.879	1	0.001	1.738	1.25-2.41
Alcohol use	0.296	0.149	3.947	1	0.047	1.345	1.00-1.80
Somatoform disorders	1.330	0.368	13.033	1	0.000	3.781	1.84-7.78
Circulatory	-0.958	0.441	4.725	1	0.030	0.384	0.16-0.91
Constant	-1.916	0.111	298.346	1	0.000	0.147	-

The initial and final Log-Likelihood of the logistic regression calculation are respectively equal to 1685.00 and 1644.29, which gives a value of the pseudo- $R^2$  of 0.025 i.e. an estimate of the variance of the risk of readmission explained by this model of 2.5%. The AUROC is equal to 0.599 indicating that this model is significant with low discriminatory power.

For men, logistic regression calculations on categories stopped after 5 iterations with no significant model ( $\chi^2=2.624$  (df=1),  $p=0.105$ ). This is consistent with the here above chi-square and odds ratio results suggesting that for men, categories as measured in this study are not associated with risk for readmission.


### ***3.4 Time-to-readmission and association with factors and PD categories***

Among the 3289 admissions of patients with PD at the UPC in the study period, 2289 were discharged home and 1000 (30.4%) were transferred to another hospital unit. Patients who were transferred to another hospital unit were excluded for the following analyses. The reason for this is that their final hospital discharge dates were not available, which made it impossible to accurately calculate their time-to-readmission. The sample of patients discharged to home

included 1819 patients, out of which 305 with multiple admissions. Transfer to hospital was independent of the readmission rate (14.7% vs 16.8%,  $\chi^2=1.737$  (df=1),  $p=0.188$ ). In terms of gender, both groups were comparable (women ratio 67.9% vs 65.5%,  $\chi^2=1.505$  (df=1),  $p=0.22$ ), as well as regarding the prevalence of factors and categories, with a diagnosis of depression being the main characteristic associated with a transfer to another hospital unit (supplemental material table R8).

The distribution of readmission according to the time-to-readmission, in number and cumulative percentage, is displayed on figure 5. Since this distribution is not normal, we considered the median time-to-readmission and not the mean (Howell, 2010). As shown on figure 5, the median time-to-readmission is 110 days, and 99.5% of readmissions occur within a 4-year horizon.


**Figure 5 :** distribution of readmissions according to time-to-readmission, in number and cumulative percentage.



Controlling for gender, the speed of readmission (supplemental material 2 figure S1) is different between men and women (Kruskal-Wallis test ( $H(1)=4.574$ ,  $p=0.032$ ). Indeed, as displayed on figure 6, the median time-to-readmission for men is 46 days while for women it is 125 days (median test  $p = 0.049$ ).


**Figure 6 :** cumulative percentage of readmission according to time-to-readmission, for men and women.





As suggested by figure 7, PD categories are associated with differences in time-to-readmission, which is supported by a Kruskal-Wallis test ( $H(9)=19.90, p=0.019$ ).

**Figure 7 :** Median time-to-readmission per category of personality disorder.



Median time-to-readmission are compared between sub-groups of patients presenting or not each factor, and median tests are performed (table 6). Results suggest that the presence of psychiatric comorbidity is associated with a decrease of the median time-to-readmission from 170.5 days to 93 days ( $p=0.042$ ). In contrast, the presence of self-harming behavior is associated with an increase of the median time-to-readmission from 78.5 to 146 days ( $p=0.03$ ).

Time-to-readmission appears independent of somatic comorbidity, interpersonal/social and precipitating factors.

**Table 6** : comparison of median time-to-readmission between sub-groups of patients without and with factors : women (vs men), psychiatric comorbidity, somatic comorbidity, substance use, self-harming behavior, interpersonal/social factors, and precipitating factors.

FACTORS (n)	Median Time w/o factor (days)	Median Time with factor (days)	p median test
<b>Gender : Women (231) vs Men</b>	<b>46.5</b>	<b>125</b>	<b>0.049</b>
<b>Psychiatric comorbidity (255)</b>	<b>170.5</b>	<b>93</b>	<b>0.042</b>
Somatic comorbidity (115)	128	80	0.108
Substance use (103)	79	165	0.083
<b>Self-harming behavior (115)</b>	<b>78.5</b>	<b>146</b>	<b>0.03</b>
Interpersonal/social factors (67)	112.5	103	0.806
Precipitating factors (40)	105	128	0.595

### 3.5 Survival analyses on time-to-readmission

A Cox survival regression, which does not assume normal distribution (Bewick et al., 2004), is performed on age, gender, psychiatric comorbidity, somatic comorbidity, substance use, self-harming behavior, interpersonal/social and precipitating factors. The model obtained ( $\chi^2=3.995$ , (df=1),  $p=0.046$ ) supports that gender is the only factor associated with time-to-readmission ( $\beta=-0.268$ , Wald =3.971,  $p=0.046$ ). This corresponds to a proportional risk ratio for women compared to men of 0.765 (95%CI 0.588-0.996), suggesting that for each day after hospital discharge, the proportional risk of a readmission is 30.7% higher for men than for women. A Kaplan-Meier logistic regression is performed to model survival probability according to gender. The model obtained is significant (Mantel-Cox Log-Rank  $\chi^2=4.027$  (df=1),  $p=0.045$ ), and suggests that, during the 3 years following discharge, the risk of readmission displays a quadratic trend for men while it is more linear for women (figure 8), which is consistent with earlier readmission for men shown on figure 6.

**Figure 8** : Readmission risk function for women (dotted red) and men (dashed blue).




Figure 9 shows that the survival probability curves for women and men have horizontal asymptotes, which is consistent with the risk of readmission tending towards 0 at 1460 days (4 years) after discharge.

**Figure 9 :** Survival probability curves for women (dotted red) and men (dashed blue).




Table 7 gives the time distribution of patients readmitted according to the Kaplan-Meier model. It indicates that after 365 days, in the subsample of readmitted patients only, 84% of the men

were readmitted while only 75% of women. This is consistent with men’s faster readmission observed, and higher proportional risk found from Cox regression.

**Table 7 :** in the sub-sample of readmitted patients : number and percentage of patients readmitted each year after hospital discharge, for women and men.

	Days after discharge	0	365	730	1095	1460	1825
Cumulative number of women readmitted over time (total 231)		0	174	206	220	230	231
Cumulative number of men readmitted (total 74)		0	62	70	72	74	74
Cumulative percentage of women readmitted		0%	75%	89%	95%	100%	100%
Cumulative percentage of men readmitted		0%	84%	95%	97%	100%	100%

#### 4. Discussion

The aim of this study was to examine whether and how personality disorder categories, as well as individual, interpersonal/social and precipitating factors may in part explain readmission rate in psychiatric emergency care, and the quickness of time-to-readmission. We observed a 16.1% readmission rate, with women more likely to be readmitted in comparison to men. Indeed, female gender was the strongest factor associated with readmission with an odds ratio of 1.69. When controlling for gender, readmission was independent of PDs categories for men, while among women, borderline category was associated with increased readmissions. Psychiatric and somatic comorbidities show only a few significant associations with readmission, which were also gender dependent. Substance use, especially alcohol, was associated with higher risk of readmission for women, but not for men. Concerning time-to-readmission, gender was found to be the first factor to consider with a median time-to-readmission of 46 days for men against 125 days for women. Categories of PD yielded different time-to-readmission, with Anankastic, Dependent and Schizoid categories being associated with a median time-to-readmission as short as 29 to 48 days, while this time is between 159 and 169 for impulsive, anxious-avoidant and histrionic categories. Psychiatric comorbidity was associated with a shorter median time-to-readmission (from 170.5 to 93 days), while self-harming behavior was associated with a longer median time-to-readmission (from 78.5 to 146

days). Lastly, both for women and men, no evidence was found to support hypotheses on association between the interpersonal/social and precipitating factors, and the risk for and time-to-readmission. In terms of statistical modelling, the factors examined here yield significant logistic regression models of the risk of readmission, though with low power with an AUROC ranging from 0.574 to 0.599. These modest results are in line with comparable studies on the risk of psychiatric readmission, which found AUROC from 0.55 to 0.74 (Morel et al., 2020; Perlman et al., 2015; Tulloch et al., 2016). Cox survival regression supports that gender is the only factor associated with the time-to-readmission, showing a time-based proportional risk of being readmitted 30.7% higher for men in comparison with women. Kaplan-Meier survival regression suggests that, both for women and men, the risk of readmission spans over 4 years, which is consistent with previous results (Lewis et al., 2019).

In our sample, the 2:1 woman to man ratio in admitted patients with PD diverges from previous studies suggesting that PD prevalence is balanced between men and women (Lewis et al., 2019; Winsper et al., 2020). However studies indicate a higher incidence of PD in women in comparison to men (Newton-Howes et al., 2021; Silberschmidt et al., 2015; Volkert et al., 2018). Cultural differences may partly explain these discrepancies (Gawda, 2018; Ronningstam et al., 2018; Winsper et al., 2020); in the socio-cultural context of our study, men may be less frequently diagnosed with a personality disorder than in other countries. The measured readmission rate is lower than findings from comparable recent publication (Lewis et al., 2019). In our view, several elements contribute to diminished readmission rate. With regards with the health-system organization, a specific intensive outpatient care is offered to UPC patients in the 3 to 6 months following their discharge, and outpatient consultations are available for known patients consulting psychiatric emergency departments in the area in order to limit the need for hospitalization. In addition, measurement of readmission made here may have missed patients who would have been hospitalized in other territories, a challenge which

we presume would affect similar studies internationally. The finding that female gender is the main risk factor for psychiatric emergency readmission differs from the results of transdiagnostic studies which indicate that this risk is independent of gender (Chang et al., 2014; Evans et al., 2017; Perlman et al., 2015; Schmutte et al., 2010) but is consistent with the observation that, among people with personality disorders, women are more likely to seek help than men (Tyrrer et al., 2015). Among women, the higher risk of readmission associated with substance use, particularly alcohol, is consistent with previous findings (Gunnell et al., 2008; Perlman et al., 2015; Schmutte et al., 2010). This underlines the importance of a thorough assessment of substance use, followed by careful management to limit the risk for readmission to psychiatric emergencies (Chang et al., 2014). Indeed, literature suggest that substance use is associated with both a detrimental increase of self-harming behavior and an increased risk of readmission, especially if it was not assessed as a primary diagnosis and consequently would not have been treated with sufficient consideration (Gunnell et al., 2008; Perlman et al., 2015). The increase in time-to-readmission associated with self-harming behavior should be considered in the context of the outpatient post-hospitalization suicide prevention programmes developed since 2015 by UPC. Although no detailed quantitative data were available for this study, it is likely that follow-up of patients which is done through phone calls or community network may contribute to reduce the risk of readmission, and in the event of a relapse, to delay its occurrence in accordance with recent results on the benefits of post-hospitalization brief contact interventions (Milner et al., 2016; Riblet et al., 2017). Considering the differences in time-to-readmission per categories of PD, ancillary analysis indicates that those differences may be related to the association between psychiatric comorbidity and shorter time-to-readmission (table 6). At last, our time-to-readmission analysis could be influenced by human factors which were not assessed in our research and may be of future interest when examining the clinical presentation of psychopathology and the decisions related to hospitalization.

There are a number of limitations to this study. First, it draws on an opportunistic, ecological sample of clinical data of individuals with a personality disorder diagnosis, collected in a specialized psychiatric crisis unit. It therefore lacks the power to generalize to other contexts, and to other diagnostic profiles including psychotic, bipolar or autistic spectrum conditions. Furthermore, even though the specialized psychiatric crisis unit at center of this study covers a large catchment area with a stable population, this population is skewed by its main sociological parameters (western european country, large urban area), and post-hospitalization trajectories are likely impacted by the follow-up care system implemented after hospitalization at UPC. These prompt caution in generalizing the results before they are confirmed with studies in centres operating in different systems, and in different socio-cultural and geographical contexts (Gawda, 2018; Ronningstam et al., 2018; Winsper et al., 2020). As with other ecological retrospective studies, another important limitation is the loss to readmission to other territories or to other parts of the health system, which we estimate as relatively limited given that UPC is the referring psychiatric emergency unit for people with personality disorder in the area. In terms of methodology, the frame for diagnosis is clinical, and although it is based on standardized tools (Mini 5.0 or SCID-2), a systematical thorough assessment with semi-structured interview is difficult to achieve with 100% of the patients in the context of a psychiatric emergency intervention. Concerning the null results obtained on associations between readmission and interpersonal/social and precipitating factors, this may be attributable to an insufficiently systematic and exhaustive assessment of these factors. Also, our goal was to examine the explanatory power of a large scope of variables on the risk for readmission, and results do not support this expectation. The poor results obtained may in part be attributed to the use of categorical variables, but they also indicate the limitations of logistic regression techniques for assessing the risk of readmission. In order to gain further knowledge and explanatory power on readmission and time-to-readmission, this orientates future research to

employ dimensional instead of categorical measures, and to use different statistical techniques such as cluster analysis or structural equations modelling, (Kupek, 2006; Oladottir et al., 2022). We also suggest that it would be fruitful to investigate psychological processes such as emotion regulation or cognitive flexibility (Bryan & Rozek, 2018; Dutcher et al., 2017; Martin et al., 2017; Wolff et al., 2018), and personality traits such as impulsivity (Auerbach et al., 2017; Moeller et al., 2001) known to be involved in mental health crisis. The identification of individual, interpersonal and precipitating factors associated with crisis occurrence and repetition may contribute to the foster individualized treatments and more efficient care during and after the crisis period to sustain resilience (Feldman, 2020; Kalisch et al., 2015).



## References

- Agnafors, S., Norman Kjellström, A., Torgerson, J., & Rusner, M. (2019). Somatic comorbidity in children and adolescents with psychiatric disorders. *European Child & Adolescent Psychiatry*, 28(11), 1517–1525. <https://doi.org/10.1007/s00787-019-01313-9>
- Auerbach, R. P., Stewart, J. G., & Johnson, S. L. (2017). Impulsivity and Suicidality in Adolescent Inpatients. *Journal of Abnormal Child Psychology*, 45(1), 91–103. <https://doi.org/10.1007/s10802-016-0146-8>
- Bach, B., Markon, K., Simonsen, E., & Krueger, R. F. (2015). Clinical Utility of the DSM-5 Alternative Model of Personality Disorders: Six Cases from Practice. *Journal of Psychiatric Practice*, 21(1), 3–25. <https://doi.org/10.1097/01.pra.0000460618.02805.ef>
- Barr, K. R., Jewell, M., Townsend, M. L., & Grenyer, B. F. S. (2020). Living with personality disorder and seeking mental health treatment: Patients and family members reflect on their experiences. *Borderline Personality Disorder and Emotion Dysregulation*, 7(1), 21. <https://doi.org/10.1186/s40479-020-00136-4>
- Bentley, R., Baker, E., & Aitken, Z. (2019). The ‘double precarity’ of employment insecurity and unaffordable housing and its impact on mental health. *Social Science & Medicine*, 225, 9–16. <https://doi.org/10.1016/j.socscimed.2019.02.008>
- Beutel, M. E., Klein, E. M., Brähler, E., Reiner, I., Jünger, C., Michal, M., Wiltink, J., Wild, P. S., Münzel, T., Lackner, K. J., & Tibubos, A. N. (2017). Loneliness in the general population: Prevalence, determinants and relations to mental health. *BMC Psychiatry*, 17(1), 97. <https://doi.org/10.1186/s12888-017-1262-x>
- Bewick, V., Cheek, L., & Ball, J. (2004). Statistics review 12: Survival analysis. *Critical Care*, 8(5), 389. <https://doi.org/10.1186/cc2955>
- Björkenstam, E., Björkenstam, C., Holm, H., Gerdin, B., & Ekselius, L. (2015). Excess cause-specific mortality in in-patient-treated individuals with personality disorder: 25-year nationwide population-based study. *British Journal of Psychiatry*, 207(4), 339–345. <https://doi.org/10.1192/bjp.bp.114.149583>

- Bryan, C. J., & Rozek, D. C. (2018). Suicide prevention in the military: A mechanistic perspective. *Current Opinion in Psychology*, 22, 27–32. <https://doi.org/10.1016/j.copsyc.2017.07.022>
- Buer Christensen, T., Paap, M. C. S., Arnesen, M., Koritzinsky, K., Nysaeter, T.-E., Eikenaes, I., Germans Selvik, S., Walther, K., Torgersen, S., Bender, D. S., Skodol, A. E., Kvarstein, E., Pedersen, G., & Hummelen, B. (2018). Interrater Reliability of the Structured Clinical Interview for the *DSM-5* Alternative Model of Personality Disorders Module i: Level of Personality Functioning Scale. *Journal of Personality Assessment*, 100(6), 630–641. <https://doi.org/10.1080/00223891.2018.1483377>
- Carroll, R., Metcalfe, C., & Gunnell, D. (2014). Hospital Presenting Self-Harm and Risk of Fatal and Non-Fatal Repetition: Systematic Review and Meta-Analysis. *PLoS ONE*, 9(2), e89944. <https://doi.org/10.1371/journal.pone.0089944>
- Chan, M. K. Y., Bhatti, H., Meader, N., Stockton, S., Evans, J., O'Connor, R. C., Kapur, N., & Kendall, T. (2016). Predicting suicide following self-harm: Systematic review of risk factors and risk scales. *British Journal of Psychiatry*, 209(4), 277–283. <https://doi.org/10.1192/bjp.bp.115.170050>
- Chang, G., Weiss, A. P., Orav, E. J., & Rauch, S. L. (2014). Predictors of frequent emergency department use among patients with psychiatric illness. *General Hospital Psychiatry*, 36(6), 716–720. <https://doi.org/10.1016/j.genhosppsych.2014.09.010>
- Clements, K. M., Murphy, J. M., Eisen, S. V., & Normand, S.-L. T. (2006). Comparison of Self-report and Clinician-rated Measures of Psychiatric Symptoms and Functioning in Predicting 1-year Hospital Readmission. *Administration and Policy in Mental Health and Mental Health Services Research*, 33(5), 568–577. <https://doi.org/10.1007/s10488-006-0066-y>
- Degenhardt, L., & Hall, W. (2001). The relationship between tobacco use, substance-use disorders and mental health: Results from the National Survey of Mental Health and Well-being. *Nicotine & Tobacco Research*, 3, 225–234.

- Donisi, V., Tedeschi, F., Wahlbeck, K., Haaramo, P., & Amaddeo, F. (2016). Pre-discharge factors predicting readmissions of psychiatric patients: A systematic review of the literature. *BMC Psychiatry*, *16*(1), 449. <https://doi.org/10.1186/s12888-016-1114-0>
- Dube, S. R., & Rishi, S. (2017). Utilizing the salutogenic paradigm to investigate well-being among adult survivors of childhood sexual abuse and other adversities. *Child Abuse & Neglect*, *66*, 130–141. <https://doi.org/10.1016/j.chiabu.2017.01.026>
- Dutcher, C. D., Vujanovic, A. A., Paulus, D. J., & Bartlett, B. A. (2017). Childhood maltreatment severity and alcohol use in adult psychiatric inpatients: The mediating role of emotion regulation difficulties. *General Hospital Psychiatry*, *48*, 42–50. <https://doi.org/10.1016/j.genhosppsych.2017.06.014>
- Evans, L. J., Harris, V., Newman, L., & Beck, A. (2017). Rapid and frequent psychiatric readmissions: Associated factors. *International Journal of Psychiatry in Clinical Practice*, *21*(4), 271–276. <https://doi.org/10.1080/13651501.2017.1324037>
- Feldman, R. (2020). What is resilience: An affiliative neuroscience approach. *World Psychiatry*, *19*(2), 132–150. <https://doi.org/10.1002/wps.20729>
- Flynn, S., Graney, J., Nyathi, T., Raphael, J., Abraham, S., Singh-Dernevik, S., Williams, A., Kapur, N., Appleby, L., & Shaw, J. (2020). Clinical characteristics and care pathways of patients with personality disorder who died by suicide. *BJPsych Open*, *6*(2), e29. <https://doi.org/10.1192/bjo.2020.11>
- Fok, M. L.-Y., Chang, C.-K., Broadbent, M., Stewart, R., & Moran, P. (2019). General hospital admission rates in people diagnosed with personality disorder. *Acta Psychiatrica Scandinavica*, *139*(3), 248–255. <https://doi.org/10.1111/acps.13004>
- Gårdvik, K. S., Torgersen, T., Rygg, M., Lydersen, S., & Indredavik, M. S. (2021). Suicidality, function and associated negative life events in an adolescent psychiatric population at 3-year follow-up. *BMC Psychiatry*, *21*(1), 109. <https://doi.org/10.1186/s12888-021-03100-w>
- Gawda, B. (2018). Cross-cultural studies on the prevalence of personality disorders. *Current Issues in Personality Psychology*, *6*(4), 318–329. <https://doi.org/10.5114/cipp.2018.80200>

- Gili, M., Comas, A., García-García, M., Monzón, S., Antoni, S.-B., & Roca, M. (2010). Comorbidity between common mental disorders and chronic somatic diseases in primary care patients. *General Hospital Psychiatry, 32*(3), 240–245. <https://doi.org/10.1016/j.genhosppsy.2010.01.013>
- Golay, P., Morandi, S., Conus, P., & Bonsack, C. (2019). Identifying patterns in psychiatric hospital stays with statistical methods: Towards a typology of post-deinstitutionalization hospitalization trajectories. *Social Psychiatry and Psychiatric Epidemiology, 54*(11), 1411–1417. <https://doi.org/10.1007/s00127-019-01717-7>
- Grant, B. F. (2009). *Prevalence, Correlates, Disability, and Comorbidity of DSM-IV Borderline Personality Disorder: Results from the Wave 2 National Epidemiologic Survey on Alcohol and Related Conditions*. 25.
- Grenyer, B. F. S., Lewis, K. L., Fanaian, M., & Kotze, B. (2018). Treatment of personality disorder using a whole of service stepped care approach: A cluster randomized controlled trial. *PLOS ONE, 13*(11), e0206472. <https://doi.org/10.1371/journal.pone.0206472>
- Gunnell, D., Hawton, K., Ho, D., Evans, J., O'Connor, S., Potokar, J., Donovan, J., & Kapur, N. (2008). Hospital admissions for self harm after discharge from psychiatric inpatient care: Cohort study. *BMJ, 337*(nov18 3), a2278–a2278. <https://doi.org/10.1136/bmj.a2278>
- Hosmer, D. W., Lemeshow, S., & Sturdivant, R. X. (2013). *Applied Logistic Regression* (1st ed.). Wiley. <https://doi.org/10.1002/9781118548387>
- Howell, D. C. (2010). *Statistical methods for psychology* (7th ed). Thomson Wadsworth.
- Iacovides, A., & Siamouli, M. (2008). Comorbid mental and somatic disorders: An epidemiological perspective. *Current Opinion in Psychiatry, 21*(4), 417–421. <https://doi.org/10.1097/YCO.0b013e328303ba42>
- James, R. K. (2016). Crisis intervention. In J. C. Norcross, G. R. VandenBos, D. K. Freedheim, & R. Krishnamurthy (Eds.), *APA handbook of clinical psychology: Applications and methods* (Vol. 3). (pp. 387–407). American Psychological Association. <https://doi.org/10.1037/14861-020>

- Kalisch, R., Müller, M. B., & Tüscher, O. (2015). A conceptual framework for the neurobiological study of resilience. *Behavioral and Brain Sciences*, 38, e92. <https://doi.org/10.1017/S0140525X1400082X>
- Kaplan, M. S., McFarland, B. H., Huguet, N., & Valenstein, M. (2012). Suicide Risk and Precipitating Circumstances Among Young, Middle-Aged, and Older Male Veterans. *American Journal of Public Health*, 102(S1), S131–S137. <https://doi.org/10.2105/AJPH.2011.300445>
- Kupek, E. (2006). Beyond logistic regression: Structural equations modelling for binary variables and its application to investigating unobserved confounders. *BMC Medical Research Methodology*, 6(1), 13. <https://doi.org/10.1186/1471-2288-6-13>
- Lakey, B., & Orehek, E. (2011). Relational regulation theory: A new approach to explain the link between perceived social support and mental health. *Psychological Review*, 118(3), 482–495. <https://doi.org/10.1037/a0023477>
- Lewis, K. L., Fanaian, M., Kotze, B., & Grenyer, B. F. S. (2019). Mental health presentations to acute psychiatric services: 3-year study of prevalence and readmission risk for personality disorders compared with psychotic, affective, substance or other disorders. *BJPsych Open*, 5(1), e1. <https://doi.org/10.1192/bjo.2018.72>
- Lin, C.-H., Chen, Y.-S., Lin, C.-Hsin., & Lin, K.-S. (2007). Factors affecting time to rehospitalization for patients with major depressive disorder. *Psychiatry and Clinical Neurosciences*, 61(3), 249–254. <https://doi.org/10.1111/j.1440-1819.2007.01662.x>
- Martin, C., Vujanovic, A., & Day, S. X. (2017). Posttraumatic Stress Disorder Symptom Severity and Emotion Regulation in Acute-Care Psychiatric Inpatients: Associations With Suicidality. *Journal of Aggression, Maltreatment & Trauma*, 26(9), 1024–1041. <https://doi.org/10.1080/10926771.2017.1341443>
- Milner, A., Spittal, M. J., Kapur, N., Witt, K., Pirkis, J., & Carter, G. (2016). Mechanisms of brief contact interventions in clinical populations: A systematic review. *BMC Psychiatry*, 16(1), 194. <https://doi.org/10.1186/s12888-016-0896-4>

- Moeller, F. G., Barratt, E. S., Dougherty, D. M., Schmitz, J. M., & Swann, A. C. (2001). Psychiatric Aspects of Impulsivity. *American Journal of Psychiatry*, *158*(11), 1783–1793. <https://doi.org/10.1176/appi.ajp.158.11.1783>
- Morel, D., Yu, K. C., Liu-Ferrara, A., Caceres-Suriel, A. J., Kurtz, S. G., & Tabak, Y. P. (2020). Predicting hospital readmission in patients with mental or substance use disorders: A machine learning approach. *International Journal of Medical Informatics*, *139*, 104136. <https://doi.org/10.1016/j.ijmedinf.2020.104136>
- Mushtaq, R. (2014). Relationship Between Loneliness, Psychiatric Disorders and Physical Health? A Review on the Psychological Aspects of Loneliness. *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH*. <https://doi.org/10.7860/JCDR/2014/10077.4828>
- Newton-Howes, G., Cunningham, R., & Atkinson, J. (2021). Personality disorder prevalence and correlates in a whole of nation dataset. *Social Psychiatry and Psychiatric Epidemiology*, *56*(4), 679–685. <https://doi.org/10.1007/s00127-020-01876-y>
- Oladottir, K., Wolf-Arehult, M., Ramklint, M., & Isaksson, M. (2022). Cluster analysis of personality traits in psychiatric patients with borderline personality disorder. *Borderline Personality Disorder and Emotion Dysregulation*, *9*(1), 7. <https://doi.org/10.1186/s40479-022-00178-w>
- Olfson, M., Marcus, S. C., & Bridge, J. A. (2013). Emergency Department Recognition of Mental Disorders and Short-Term Outcome of Deliberate Self-Harm. *American Journal of Psychiatry*, *170*(12), 1442–1450. <https://doi.org/10.1176/appi.ajp.2013.12121506>
- Perlman, C. M., Hirdes, J. P., & Vigod, S. (2015). Psychiatric Rehospitalization: Development of a Person-Level Indicator for Care Planning and Quality Assurance. *The Primary Care Companion For CNS Disorders*. <https://doi.org/10.4088/PCC.15m01784>
- Pernice-Duca, F. (2010). Family Network Support and Mental Health Recovery. *Journal of Marital and Family Therapy*, *36*(1), 13–27. <https://doi.org/10.1111/j.1752-0606.2009.00182.x>

- Polling, C., Bakolis, I., Hotopf, M., & Hatch, S. L. (2021). Variation in rates of self-harm hospital admission and re-admission by ethnicity in London: A population cohort study. *Social Psychiatry and Psychiatric Epidemiology*, *56*(11), 1967–1977. <https://doi.org/10.1007/s00127-021-02087-9>
- Puddephatt, J.-A., Jones, A., Gage, S. H., Fear, N. T., Field, M., McManus, S., McBride, O., & Goodwin, L. (2021). Associations of alcohol use, mental health and socioeconomic status in England: Findings from a representative population survey. *Drug and Alcohol Dependence*, *219*, 108463. <https://doi.org/10.1016/j.drugalcdep.2020.108463>
- Rajani, N. B., Giannakopoulos, G., & Filippidis, F. T. (2016). Job insecurity, financial difficulties and mental health in Europe. *Occupational Medicine*, *66*(8), 681–683. <https://doi.org/10.1093/occmed/kqw111>
- Reiland, S. A., & Clark, C. B. (2017). Relationship between event type and mental health outcomes: Event centrality as mediator. *Personality and Individual Differences*, *114*, 155–159. <https://doi.org/10.1016/j.paid.2017.04.009>
- Riblet, N. B. V., Shiner, B., Young-Xu, Y., & Watts, B. V. (2017). Strategies to prevent death by suicide: Meta-analysis of randomised controlled trials. *British Journal of Psychiatry*, *210*(6), 396–402. <https://doi.org/10.1192/bjp.bp.116.187799>
- Richard-Lepouriel, H., Weber, K., Baertschi, M., DiGiorgio, S., Sarasin, F., & Canuto, A. (2015). Predictors of Recurrent Use of Psychiatric Emergency Services. *Psychiatric Services*, *66*(5), 521–526. <https://doi.org/10.1176/appi.ps.201400097>
- Ronningstam, E. F., Keng, S.-L., Ridolfi, M. E., Arbabi, M., & Grenyer, B. F. S. (2018). Cultural Aspects in Symptomatology, Assessment, and Treatment of Personality Disorders. *Current Psychiatry Reports*, *20*(4), 22. <https://doi.org/10.1007/s11920-018-0889-8>
- Rosca, P., Bauer, A., Grinshpoon, A., Khawaled, R., Mester, R., & Ponizovsky, A. M. (2006). *Rehospitalizations Among Psychiatric Patients Whose First Admission was Involuntary: A 10-Year Follow-Up*. 9.
- Sansone, R. A., & Sansone, L. A. (2011). Gender patterns in Borderline Personality Disorder. *Innov Clin Neurosci.*, *8*(5), 16–20.

- Schmutte, T., Dunn, C. L., & Sledge, W. H. (2010). Predicting Time to Readmission in Patients With Recent Histories of Recurrent Psychiatric Hospitalization: A Matched-Control Survival Analysis. *Journal of Nervous & Mental Disease, 198*(12), 860–863. <https://doi.org/10.1097/NMD.0b013e3181fe726b>
- Schulte Holthausen, B., & Habel, U. (2018). Sex Differences in Personality Disorders. *Current Psychiatry Reports, 20*(12), 107. <https://doi.org/10.1007/s11920-018-0975-y>
- Scott, K. M., Al-Hamzawi, A. O., Andrade, L. H., Borges, G., Caldas-de-Almeida, J. M., Fiestas, F., Gureje, O., Hu, C., Karam, E. G., Kawakami, N., Lee, S., Levinson, D., Lim, C. C. W., Navarro-Mateu, F., Okoliyski, M., Posada-Villa, J., Torres, Y., Williams, D. R., Zakhosha, V., & Kessler, R. C. (2014). Associations Between Subjective Social Status and *DSM-IV* Mental Disorders: Results From the World Mental Health Surveys. *JAMA Psychiatry, 71*(12), 1400. <https://doi.org/10.1001/jamapsychiatry.2014.1337>
- Silberschmidt, A., Lee, S., Zanarini, M., & Schulz, S. C. (2015). Gender Differences in Borderline Personality Disorder: Results From a Multinational, Clinical Trial Sample. *Journal of Personality Disorders, 29*(6), 828–838. [https://doi.org/10.1521/pedi\\_2014\\_28\\_175](https://doi.org/10.1521/pedi_2014_28_175)
- Slade, K., & Forrester, A. (2013). Measuring IPDE-SQ personality disorder prevalence in pre-sentence and early-stage prison populations, with sub-type estimates. *International Journal of Law and Psychiatry, 36*(3–4), 207–212. <https://doi.org/10.1016/j.ijlp.2013.04.018>
- Stoltzfus, J. C. (2011). Logistic Regression: A Brief Primer: LOGISTIC REGRESSION: A BRIEF PRIMER. *Academic Emergency Medicine, 18*(10), 1099–1104. <https://doi.org/10.1111/j.1553-2712.2011.01185.x>
- Tabachnick, B. G., Fidell, L. S., & Ullman, J. B. (2019). *Using multivariate statistics* (Seventh edition). Pearson.
- Tulloch, A. D., David, A. S., & Thornicroft, G. (2016). Exploring the predictors of early readmission to psychiatric hospital. *Epidemiology and Psychiatric Sciences, 25*(2), 181–193. <https://doi.org/10.1017/S2045796015000128>



- Tyrer, P., Reed, G. M., & Crawford, M. J. (2015). Classification, assessment, prevalence, and effect of personality disorder. *The Lancet*, 385(9969), 717–726. [https://doi.org/10.1016/S0140-6736\(14\)61995-4](https://doi.org/10.1016/S0140-6736(14)61995-4)
- Volkert, J., Gablonski, T.-C., & Rabung, S. (2018). Prevalence of personality disorders in the general adult population in Western countries: Systematic review and meta-analysis. *The British Journal of Psychiatry*, 213(6), 709–715. <https://doi.org/10.1192/bjp.2018.202>
- Wamser-Nanney, R., Howell, K. H., Schwartz, L. E., & Hasselle, A. J. (2018). The moderating role of trauma type on the relationship between event centrality of the traumatic experience and mental health outcomes. *Psychological Trauma: Theory, Research, Practice, and Policy*, 10(5), 499–507. <https://doi.org/10.1037/tra0000344>
- Wang, J., Mann, F., Lloyd-Evans, B., Ma, R., & Johnson, S. (2018). Associations between loneliness and perceived social support and outcomes of mental health problems: A systematic review. *BMC Psychiatry*, 18(1), 156. <https://doi.org/10.1186/s12888-018-1736-5>
- Wilhelm, K., Kovess, V., Rios-Seidel, C., & Finch, A. (2004). Work and mental health. *Social Psychiatry and Psychiatric Epidemiology*, 39(11), 866–873. <https://doi.org/10.1007/s00127-004-0869-7>
- Winsper, C., Bilgin, A., Thompson, A., Marwaha, S., Chanen, A. M., Singh, S. P., Wang, A., & Furtado, V. (2020). The prevalence of personality disorders in the community: A global systematic review and meta-analysis. *The British Journal of Psychiatry*, 216(2), 69–78. <https://doi.org/10.1192/bjp.2019.166>
- Wolff, J. C., Davis, S., Liu, R. T., Cha, C. B., Cheek, S. M., Nestor, B. A., Frazier, E. A., Schaffer, M. M., & Spirito, A. (2018). Trajectories of Suicidal Ideation among Adolescents Following Psychiatric Hospitalization. *Journal of Abnormal Child Psychology*, 46(2), 355–363. <https://doi.org/10.1007/s10802-017-0293-6>