



OPEN ACCESS

Original research

Head-to-head trial of pegunigalsidase alfa versus agalsidase beta in patients with Fabry disease and deteriorating renal function: results from the 2-year randomised phase III BALANCE study

Eric L Wallace,¹ Ozlem Goker-Alpan,² William R Wilcox,³ Myrl Holida,⁴ John Bernat,⁴ Nicola Longo,⁵ Aleš Linhart ,⁶ Derralynn A Hughes ,⁷ Robert J Hopkin,⁸ Camilla Tøndel,^{9,10} Mirjam Langeveld,¹¹ Pilar Giraldo,^{12,13} Antonio Pisani,¹⁴ Dominique Paul Germain ,¹⁵ Ankit Mehta,¹⁶ Patrick B Deegan,¹⁷ Maria Judit Molnar,¹⁸ Damara Ortiz,¹⁹ Ana Jovanovic,²⁰ Michael Muriello,²¹ Bruce A Barshop,²² Virginia Kimonis,²³ Bojan Vujkovic,²⁴ Albina Nowak ,²⁵ Tarekegn Geberhiwot,²⁶ Ilkka Kantola,²⁷ Jasmine Knoll,²⁸ Stephen Waldek,²⁹ Khan Nedd,³⁰ Amel Karaa,³¹ Einat Brill-Almon,³² Sari Alon,³³ Raul Chertkoff,³² Rossana Rocco,³⁴ Anat Sakov,³⁵ David G Warnock ¹

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/jmg-2023-109445>).

For numbered affiliations see end of article.

Correspondence to

Dr David G Warnock, The University of Alabama at Birmingham, Birmingham, AL 35205, USA; dwarnock@uab.edu

Received 2 June 2023

Accepted 10 September 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Wallace EL, Goker-Alpan O, Wilcox WR, et al. *J Med Genet* Epub ahead of print: [please include Day Month Year]. doi:10.1136/jmg-2023-109445

ABSTRACT

Background Pegunigalsidase alfa is a PEGylated α -galactosidase A enzyme replacement therapy. BALANCE (NCT02795676) assessed non-inferiority of pegunigalsidase alfa versus agalsidase beta in adults with Fabry disease with an annualised estimated glomerular filtration rate (eGFR) slope more negative than -2 mL/min/1.73 m²/year who had received agalsidase beta for ≥ 1 year.

Methods Patients were randomly assigned 2:1 to receive 1 mg/kg pegunigalsidase alfa or agalsidase beta every 2 weeks for 2 years. The primary efficacy analysis assessed non-inferiority based on median annualised eGFR slope differences between treatment arms.

Results Seventy-seven patients received either pegunigalsidase alfa (n=52) or agalsidase beta (n=25). At baseline, mean (range) age was 44 (18–60) years, 47 (61%) patients were male, median eGFR was 74.5 mL/min/1.73 m² and median (range) eGFR slope was -7.3 (-30.5 , 6.3) mL/min/1.73 m²/year. At 2 years, the difference between median eGFR slopes was -0.36 mL/min/1.73 m²/year, meeting the prespecified non-inferiority margin. Minimal changes were observed in lyso-Gb3 concentrations in both treatment arms at 2 years. Proportions of patients experiencing treatment-related adverse events and mild or moderate infusion-related reactions were similar in both groups, yet exposure-adjusted rates were 3.6-fold and 7.8-fold higher, respectively, with agalsidase beta than pegunigalsidase alfa. At the end of the study, neutralising antibodies were detected in 7 out of 47 (15%) pegunigalsidase alfa-treated patients and 6 out of 23 (26%) agalsidase beta-treated patients. There were no deaths.

Conclusions Based on rate of eGFR decline over 2 years, pegunigalsidase alfa was non-inferior to agalsidase beta. Pegunigalsidase alfa had lower rates of treatment-emergent adverse events and mild or moderate infusion-related reactions.

Trial registration number NCT02795676.

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Currently available enzyme replacement therapies (ERTs) benefit patients with Fabry disease (FD) but are associated with infusion-related reactions and the development of antidrug antibodies.
- ⇒ Pegunigalsidase alfa is a novel, PEGylated α -galactosidase A ERT with prolonged half-life, improved tolerability and lower incidence of infusion-related reactions.

WHAT THIS STUDY ADDS

- ⇒ This was the first randomised, double-blind, head-to-head clinical trial of ERTs in FD and demonstrated comparable renal efficacy on estimated glomerular filtration rate slope of 2 years of treatment with pegunigalsidase alfa compared with agalsidase beta in adults with deteriorating renal function and history of long-term agalsidase beta treatment.
- ⇒ Pegunigalsidase alfa-treated patients experienced a lower rate of mild or moderate infusion-related reactions.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ Pegunigalsidase alfa provides patients with an additional treatment option for FD.

INTRODUCTION

Fabry disease (FD; OMIM #301500) is a rare, progressive X-linked lysosomal disorder caused by pathogenic variants in the *GLA* gene leading to deficiency of α -galactosidase A (α -Gal A) and associated accumulation of globotriaosylceramide (Gb3) and globotriaosylsphingosine (lyso-Gb3).^{1–5} FD involves many systems, including renal, cardiac,

neurologic and cerebrovascular.⁶ Phenotypic variation is marked in heterozygous females with higher plasma α -Gal A activity, random X-linked inactivation and deficient cross-correction.^{7–11}

At the time the BALANCE study was conducted, three FD treatment options were approved with varying availability by country, including two enzyme replacement therapies (ERTs; agalsidase alfa and agalsidase beta) and one oral pharmacological chaperone therapy (migalastat).^{12,13} Agalsidase alfa and agalsidase beta are ERT preparations with engineered α -Gal A, administered via intravenous infusion every 2 weeks (E2W) at 0.2 and 1 mg/kg doses, respectively.¹² In clinical trials, ERT reduced the rate of kidney function decline, improved cardiac structure, reduced neuropathic pain severity and improved gastrointestinal symptoms.^{14,15} Patients receiving agalsidase beta who initiated treatment at a younger age and with higher estimated glomerular filtration rate (eGFR) benefited most; baseline glomerular sclerosis and uncontrolled proteinuria were indicators for poor prognosis.¹⁶ ERTs can lead to clinically relevant improvements in natural disease course, although disease progression occurs in some cases.¹⁷ Antidrug antibodies (ADAs) occur in up to 83% of patients with FD receiving agalsidase beta in clinical trials¹⁸ and are more common in males.¹⁹ In real-world studies, ADAs negatively impact biomarker response to ERT (less robust lyso-Gb3 reduction) and clinical outcomes^{12,19–22} and are associated with infusion-related reactions.^{23–27}

Pegunigalsidase alfa, approved in the EU and the USA, is a novel PEGylated α -Gal A ERT with prolonged half-life, and designed to have reduced immunogenicity and potentially improved tolerability.^{13,28–30} It is chemically modified with 2 kDa homo-bifunctional polyethylene glycol (PEG) molecules cross-linking two plant cell-derived subunits of α -Gal A or bound to surface lysine residues by one end only,²⁹ resulting in PEGylated, covalently bound 114 kDa homodimer enzyme. Potential masking of some immune epitopes by PEGylation²⁹ may explain the lower immunogenicity.^{30,31} Comparing in vitro and in vivo properties of pegunigalsidase alfa versus agalsidase alfa and agalsidase beta demonstrated equivalent activity with longer in vivo plasma half-life, in vitro stability with plasma-like and lysosomal-like conditions, and different cellular uptake routes.²⁹ Elimination plasma half-life is approximately 90–110 min for agalsidase alfa and 80–120 min for agalsidase beta.^{18,32} Pegunigalsidase alfa has an elimination half-life of ~80 hours, effective Gb3 clearance from renal tissue and a favourable safety profile up to 12 months.^{30,33}

The phase III BALANCE study (NCT02795676) is the first randomised, double-blind, active-control, head-to-head clinical trial of ERTs in FD and is the first study to directly evaluate efficacy, safety and tolerability of pegunigalsidase alfa versus agalsidase beta in adult patients with previous agalsidase beta treatment and deteriorating renal function.

METHODS

Study design

BALANCE was conducted at 29 study centres in 12 countries from 22 August 2016 to 12 October 2021. Patients were randomly assigned 2:1 to receive pegunigalsidase alfa or agalsidase beta, 1 mg/kg intravenously E2W for 24 months. Randomisation was stratified by screening urine protein-to-creatinine ratio (UPCR) <1 or ≥ 1 g/g. The primary objective was to evaluate efficacy of pegunigalsidase alfa versus agalsidase beta; the primary analysis was to demonstrate non-inferiority of pegunigalsidase alfa with respect to annualised change in eGFR slope, based on a prespecified margin of median annualised eGFR slope difference and its CI between groups. The secondary efficacy

endpoint reported here is change in plasma lyso-Gb3 concentration. Safety endpoints included treatment-emergent adverse events (TEAEs), infusion-related reactions, premedication use, and pre-existing and on-study ADA status.

Pegunigalsidase alfa and agalsidase beta infusions were planned to be initially administered over 3 hours at the study centre. If well tolerated, infusion duration was gradually reduced to 1.5 hours, and patients could receive home infusions. If previously used, premedication was continued but gradually decreased over 3 months at investigator's discretion based on patient tolerability. After study completion, patients were invited to participate in an open-label extension of pegunigalsidase alfa (PB-102-F60; BRILLIANCE; NCT03566017).

Patients

Patients were symptomatic, aged 18–60 years, with ≥ 1 characteristic FD feature (neuropathic pain, cornea verticillata, clustered angiokeratomas), screening eGFR of 40–120 mL/min/1.73 m² calculated via 2009 Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation,³⁴ deteriorating renal function (linear eGFR slope more negative than -2 mL/min/1.73 m²/year based on ≥ 3 creatinine values over 9–18 months), and ≥ 1 year agalsidase beta treatment (1 mg/kg E2W). FD was confirmed in males by decreased plasma and/or leucocyte α -Gal A activity to $<30\%$ of mean normal levels and in females historically confirmed based on known pathogenic *GLA* variants or novel variants shared by a first-degree male relative with FD.

Key exclusion criteria included the following: anaphylaxis or type I hypersensitivity reactions to agalsidase beta; historical eGFR >120 mL/min/1.73 m² for 9–18 months prior; renal dialysis or transplantation; acute kidney injury within the last 12 months; angiotensin-converting enzyme inhibitor (ACEi) or angiotensin II receptor blocker (ARB) initiation or dose change within 4 weeks prior; UPCR >0.5 g/g and not ACEi/ARB-treated.³⁵

Estimated glomerular filtration rate

Patients provided pre-infusion blood samples for creatinine assessment. Analysis was performed centrally using an enzymatic assay; eGFR (2009 CKD-EPI equation³⁴) was calculated at screening and at least monthly for 30 visits during treatment (including baseline).

Plasma lyso-Gb3

Patients provided pre-infusion blood samples for plasma lyso-Gb3 assessment at baseline, 1.5 months, every 3 months the first year and every 6 months the second year. Quantification was performed centrally, including matrix lipid extraction and ultra-performance liquid chromatography–tandem mass spectrometry.^{36,37}

Safety

TEAEs were adverse events (AEs) occurring between treatment initiation and final infusion. TEAEs included new medical conditions and pre-existing ones that worsened during treatment. Infusion-related reactions were defined as TEAEs beginning during or within 2 hours of infusion whose causality was assessed as definitely, probably or possibly treatment-related; these excluded injection site reactions, which were considered procedure-related.

Antidrug antibodies

ADA assessment was performed using a multi-tiered approach based on a solid-phase ELISA using pegunigalsidase alfa or agalsidase beta for antibody capture.

Statistical methods

Intent-to-treat (ITT) population was the main set for efficacy analyses and included all randomly assigned patients who received at least one dose (including partial doses). Per protocol (PP) population included ITT patients who completed 24 months of treatment, with $\geq 80\%$ compliance and no major protocol violations potentially affecting the primary endpoint. In a non-inferiority study, PP and ITT analysis sets should be considered together for interpretation. Safety population included all randomly assigned patients who received any treatment dose (including partial doses). Treatment arm differences in rates of TEAEs and infusion-related reactions were analysed post hoc via Poisson regression with offset of treatment duration and number of infusions, respectively.

The primary efficacy endpoint was annualised eGFR slope. To determine non-inferiority, the lower limit of a 95% CI for the difference of median annualised eGFR slopes was prespecified to be $-3\text{ mL/min/1.73 m}^2\text{/year}$. Comparison of eGFR slopes was performed using quantile regression for the median, where individual slopes were estimated in the first stage using linear regression based on all eGFR assessments of each patient. In the second stage, quantile regression was used with treatment arm as the covariate.³⁸

A post hoc analysis adjusting for sex was performed for the primary endpoint. Lyso-Gb3 concentrations and change from baseline over time were compared between arms and stratified by sex using post hoc Wilcoxon rank test. Additional post hoc analyses were descriptive.

Refer to online supplemental material 1 for additional methods and CONSORT (Consolidated Standards of Reporting Trials) reporting guidelines.

RESULTS

Patients

Of the 127 screened patients, 49 failed screening and 78 met inclusion criteria and were randomly assigned (53 to pegunigalsidase alfa, 25 to agalsidase beta) (online supplemental figure 1). Of the 57 genetic variants identified, *GLA* c.679C>T (p.(Arg227Ter)) and c.680G>A (p.(Arg227Gln)) were the most common (online supplemental table 1).

Seventy-seven patients received either pegunigalsidase alfa (n=52) or agalsidase beta (n=25); one patient randomly assigned to pegunigalsidase alfa withdrew consent before receiving treatment (online supplemental figure 1). Forty-eight (90.6%) patients receiving pegunigalsidase alfa and 24 (96.0%) receiving agalsidase beta completed 24 months of treatment. Three patients on pegunigalsidase alfa (including the patient who did not receive treatment) and one patient on agalsidase beta voluntarily withdrew consent. Two patients on pegunigalsidase alfa discontinued due to AEs within the first year. In the ITT population (n=77), baseline characteristics were not significantly different between arms (table 1).

Estimated glomerular filtration rate

Median eGFR at baseline was nearly identical for patients on pegunigalsidase alfa ($73.5\text{ mL/min/1.73 m}^2$) and agalsidase beta ($74.9\text{ mL/min/1.73 m}^2$; $p=0.82$) (table 1 and figure 1). Ranges were broad: 30–126 and 34–108 mL/min/1.73 m^2 , respectively,

and changes were observed for individual patient values from screening to baseline. eGFR change from baseline showed a similar decline at 24 months in the two arms with a median change of $-2.39\text{ mL/min/1.73 m}^2$ for pegunigalsidase alfa and $-3.20\text{ mL/min/1.73 m}^2$ for agalsidase beta. Refer to online supplemental figure 2 for eGFR and eGFR change from baseline stratified by baseline eGFR.

Baseline median eGFR slope for the ITT population, based on historical patient data, was approximately $-7\text{ mL/min/1.73 m}^2\text{/year}$ overall and was similar between arms ($p=0.37$) (table 1). In males, baseline eGFR slope ranged from -30.5 to $6.3\text{ mL/min/1.73 m}^2\text{/year}$ with pegunigalsidase alfa and -20.3 to $-2.8\text{ mL/min/1.73 m}^2\text{/year}$ with agalsidase beta; in females, ranges were -19.2 to $-1.6\text{ mL/min/1.73 m}^2\text{/year}$ and -13.9 to $-6.9\text{ mL/min/1.73 m}^2\text{/year}$, respectively. Median (95%CI limits) eGFR slopes after 24 months of treatment were -2.51 ($-3.79, -1.24$) $\text{mL/min/1.73 m}^2\text{/year}$ with pegunigalsidase alfa and -2.16 ($-3.81, -0.51$) $\text{mL/min/1.73 m}^2\text{/year}$ with agalsidase beta (table 2). Difference in median eGFR slope for the ITT population between arms was $-0.36\text{ mL/min/1.73 m}^2\text{/year}$ (95%CI $-2.44, 1.73$). The lower limit of the CI was above the prespecified non-inferiority margin; hence, non-inferiority was achieved. The 95%CI included 0, with extensive overlap between individual CIs, indicating no significant difference between arms.

Subgroup analysis of eGFR slope showed median (95%CI) eGFR slope overlapped across arms for males: -3.44 ($-5.38, -1.50$) $\text{mL/min/1.73 m}^2\text{/year}$ with pegunigalsidase alfa and -2.01 ($-3.98, -0.04$) $\text{mL/min/1.73 m}^2\text{/year}$ with agalsidase beta, with a difference of -1.43 ($-3.96, 1.10$) $\text{mL/min/1.73 m}^2\text{/year}$ (table 2). High overlap was also observed across arms for females.

All ADA-positive patients at baseline were male (table 1). Median (95%CI) eGFR slope in ADA-positive males was similar between arms: -2.51 ($-5.28, 0.25$) $\text{mL/min/1.73 m}^2\text{/year}$ with pegunigalsidase alfa (n=18) and -2.16 ($-6.25, 1.93$) $\text{mL/min/1.73 m}^2\text{/year}$ with agalsidase beta (n=8), with a difference (95%CI) of -0.36 ($-5.16, 4.45$) $\text{mL/min/1.73 m}^2\text{/year}$ between arms (table 2). Median (95%CI) eGFR slope in ADA-negative patients in both treatment arms was similar: -2.22 ($-4.02, -0.43$) $\text{mL/min/1.73 m}^2\text{/year}$ with pegunigalsidase alfa and -2.16 ($-4.06, -0.26$) $\text{mL/min/1.73 m}^2\text{/year}$ with agalsidase beta; difference (95%CI) of -0.07 ($-2.41, 2.27$) $\text{mL/min/1.73 m}^2\text{/year}$.

Plasma lyso-Gb3

As expected, males had higher plasma lyso-Gb3 concentrations throughout the study compared with females for both arms (figure 2). At 24 months, median (range) plasma lyso-Gb3 change from baseline in males was 5.30 (-32.2 to 32.7) nM with pegunigalsidase alfa and -2.40 (-102.3 to 2.4) nM with agalsidase beta; ($p=0.0001$); in females, the change was minimal: 0.10 (-4.0 to 5.8) nM with pegunigalsidase alfa and -0.30 (-0.7 to 0.9) nM with agalsidase beta (table 3, online supplemental figure 3) ($p=0.54$). In the overall population, median plasma lyso-Gb3 remained relatively stable in each arm with <2 nM change from baseline to 24 months (1.15 nM for pegunigalsidase alfa and -1.50 nM for agalsidase beta). In a post hoc assessment of lyso-Gb3 dynamics over study duration, individual patient profiles were analysed for plasma lyso-Gb3 increases from baseline exceeding 20% and 10 nM. Results indicated that lyso-Gb3 increases of this magnitude likely occur in patients with baseline UPCr ≥ 1 g/g and ADA positive status.

Table 1 Patient demographics and baseline characteristics

Parameter	Pegunigalsidase alfa			Agalsidase beta			Overall (n=77)	P value*
	Males (n=29)	Females (n=23)	Overall (n=52)	Males (n=18)	Females (n=7)	Overall (n=25)		
Age, years								0.60
Mean±SD	42.6±11.5	45.6±8.3	43.9±10.2	46.5±6.9	41.7±14.5	45.2±9.6	44.3±10.0	
Sex, n (%)†	29 (56)	23 (44)	–	18 (72)	7 (28)	–	M 47 (61) F 30 (39)	0.19
eGFR, mL/min/1.73 m ² ‡								
Mean (SE)	71.6 (4.4)	75.8 (3.0)	73.5 (2.8)	69.2 (5.0)	86.9 (5.3)	74.2 (4.2)	73.7 (2.3)	0.82
Median	70.2	75.5	73.5	71.8	88.2	74.9	74.5	
Min, max	30.2, 125.9	47.2, 107.1	30.2, 125.9	34.1, 106.3	65.8, 107.6	34.1, 107.6	30.2, 125.9	
eGFR slope, mL/min/1.73 m ² /year§								0.37
Mean (SE)	–8.7 (1.5)	–7.2 (1.0)	–8.0 (0.9)	–7.8 (1.1)	–9.4 (1.0)	–8.3 (0.9)	–8.1 (0.7)	
Median	–7.3	–6.5	–6.7	–7.3	–8.3	–7.8	–7.3	
Min, max	–30.5, 6.3	–19.2, –1.6	–30.5, 6.3	–20.3, –2.8	–13.9, –6.9	–20.3, –2.8	–30.5, 6.3	
UPCR, n (%)								0.52
UPCR≤0.5 g/g	15 (52)	21 (91)	36 (69)	13 (72)	7 (100)	20 (80)	56 (73)	
0.5<UPCR<1 g/g	8 (28)	1 (4)	9 (17)	2 (11)	0	2 (8)	11 (14)	
UPCR≥1 g/g	6 (21)	1 (4)	7 (14)	3 (17)	0	3 (12)	10 (13)	
Treatment with ACEi or ARBs, n (%)	17 (59)	9 (39)	26 (50)	15 (83)	1 (14)	16 (64)	42 (55)	0.22
Positive ADA status¶, n (%)	18 (62)	0	18 (35)	8 (44)	0	8 (32)	26 (34)	0.82
Positive for neutralising antibodies, n (%)**	17 (59)	0	17 (33)	7 (39)	0	7 (28)	24 (31)	0.8
Length of previous agalsidase beta treatment, years††								0.25
Mean±SD	6.4±4.9	4.2±2.1	5.4±4.0	6.6±3.4	6.1±3.7	6.4±3.4	5.8±3.8	

*P values were calculated between treatment arms for age and length of previous agalsidase beta treatment with t-test; for sex, UPCR category, ACEi/ARB treatment, and ADA status by Pearson χ^2 test; for eGFR and eGFR slope by Wilcoxon; for neutralising antibodies by Fisher's exact test.
†Percentage calculated out of total number of patients per treatment arm.
‡Inclusion criteria specified patients have eGFR of 40–120 mL/min/1.73 m² at screening visit; eGFR at baseline visit (presented here) was outside of this range for some patients. Normal range 90–120 mL/min/1.73 m².⁴⁶
§eGFR slope at baseline was based on historical, screening, and baseline serum creatinine measurements and was more positive than –2 mL/min/1.73 m²/year at baseline for some patients. eGFR slope as negative as –1 mL/min/1.73 m²/year is considered normal for patients age ≥40 years.⁴⁶
¶All patients were evaluated for the presence of antidrug IgG antibodies to their assigned drug at baseline.
**Percentage calculated out of total number of patients of the respective sex per treatment arm.
††Last continuous agalsidase beta treatment.
ACEi, angiotensin-converting enzyme inhibitors; ADAs, antidrug antibodies; ARBs, angiotensin II receptor blockers; eGFR, estimated glomerular filtration rate; F, female; M, male; UPCR, urine protein creatinine ratio.

Patients with baseline UPCR ≥ 1 g/g and ADA positive status were all male and more frequently assigned to the pegunigalsidase alfa arm (of 18 ADA-positive patients, 6 had UPCR ≥ 1 g/g) than agalsidase beta (of 8 ADA-positive patients, 1 had UPCR ≥ 1 g/g) (data not shown).

Safety

Treatment-emergent adverse events

Most patients (90.4% of pegunigalsidase alfa-treated and 96.0% of agalsidase beta-treated) experienced ≥ 1 TEAE (table 4). The TEAE rate (events/100 exposure-years) was significantly lower with pegunigalsidase alfa (572) than agalsidase beta (817) (rate ratio (95% CI) of 0.70 (0.62, 0.80), $p < 0.0001$). Among males, rates were significantly lower with pegunigalsidase alfa (545) than agalsidase beta (922) (rate ratio (95% CI) of 0.59 (0.51, 0.69), $p < 0.0001$), but there was no significant difference among females (605 vs 549; rate ratio (95% CI) of 1.10 (0.86, 1.42), $p = 0.45$). Proportions of patients experiencing treatment-related TEAEs were similar (40% with pegunigalsidase alfa and 44% with agalsidase beta). Treatment-related TEAE rate (events/100 exposure-years) was 3.6-fold lower with pegunigalsidase alfa (43) than agalsidase beta (153).

On exploring whether all patients experienced relatively equal numbers of treatment-related TEAEs, it was

observed that 2 patients per treatment arm with the most treatment-related TEAEs constituted 26% (11 events) for pegunigalsidase alfa and 57% (43 events) for agalsidase beta. A male receiving pegunigalsidase alfa reported 6 related TEAEs and another male reported 5; the remaining 19 patients reported 1–4 events each (13 males, 6 females); one female receiving agalsidase beta reported 18 related TEAEs, and a male reported 25; the remaining 9 (8 males, 1 female) patients reported 1–8 events each.

Two pegunigalsidase alfa-treated patients experienced TEAEs leading to withdrawal. One experienced a hypersensitivity reaction during the first infusion which resolved that day; defined as an infusion-related reaction, and considered as serious, severe and treatment-related. This patient experienced another hypersensitivity reaction on rechallenge and withdrew. At first infusion, the patient was positive for anti-pegunigalsidase alfa IgE and IgG. Another patient was diagnosed with FD-related end-stage renal disease necessitating kidney transplant and withdrew. An additional pegunigalsidase alfa-treated patient had treatment-related immune complex-mediated membranoproliferative glomerulonephritis leading to cessation of treatment but not withdrawal from the study. A kidney biopsy confirmed the presence of 1+IgG subendothelial deposits and 1+kappa and lambda deposits; c3 was negative.




Figure 1 Median eGFR over time in (A) all patients and (B) by sex. (B) Number of female and male patients: pegunigalsidase alfa, n=23 and n=29, respectively; agalsidase beta, n=7 and n=18, respectively. ^aNumber of patients at baseline and 24 months: n=52 and n=47, respectively. ^bNumber of patients at baseline and 24 months: n=25 and n=24, respectively. eGFR, estimated glomerular filtration rate; ITT, intent-to-treat.

There were continued capillary cell and endothelial cell inclusions and numerous podocyte inclusions. Immunohistochemistry confirmed immune complexes collocated with α -Gal. TEAEs by system are presented in online supplemental table 2. There were no deaths.

Infusion-related reactions

With pegunigalsidase alfa, infusion-related reaction rate was 0.5 event/100 infusions, with 11 (21%) patients reporting 13 infusion-related reactions (table 4). Infusion-related reaction rate with agalsidase beta was significantly higher (3.9 events/100 infusions; rate ratio (95% CI) of 0.13 (0.07, 0.24), $p < 0.0001$), with six (24%) patients experiencing 51 infusion-related

reactions. The proportion of males reporting infusion-related reactions was numerically higher (31% with pegunigalsidase alfa and 28% with agalsidase beta) than that of females (9% with pegunigalsidase alfa and 14% with agalsidase beta) (difference between arms not significant). Infusion-related reaction rate was significantly lower in both males and females on pegunigalsidase alfa compared with males and females on agalsidase beta (males: 0.8 vs 3.5 events/100 infusion, respectively; rate ratio (95% CI) of 0.22 (0.11, 0.44); $p < 0.0001$) (females: 0.2 vs 4.9 events/100 infusions, respectively; rate ratio (95% CI) of 0.04 (0.01, 0.15); $p < 0.0001$).

In both arms, the proportion of ADA-positive patients reporting infusion-related reactions was higher (33% with

Table 2 Median eGFR slope and 95% CI model* (ITT population)—by treatment arm, sex and ADA status

ITT population median eGFR slope	Pegunigalsidase alfa (n=52)†	Agalsidase beta (n=25)‡	Difference between arms
Baseline, mL/min/1.73 m ² /year			
Overall	-6.70	-7.84	-
Male	-7.25	-7.25	-
Female	-6.45	-8.31	-
ADA-positive	-5.75	-6.08	-
ADA-negative	-7.10	-7.84	-
24 months, mL/min/1.73 m ² /year (95% CI)			
Overall	-2.51 (-3.79, -1.24)	-2.16 (-3.81, -0.51)	-0.36 (-2.44§, 1.73)
Male	-3.44 (-5.38, -1.50)	-2.01 (-3.98, -0.04)	-1.43 (-3.96, 1.10)
Female	-1.15 (-3.11, 0.81)	-2.79 (-6.28, 0.70)	1.64 (-2.56, 5.84)
ADA-positive	-2.51 (-5.28, 0.25)	-2.16 (-6.25, 1.93)	-0.36 (-5.16, 4.45)
ADA-negative	-2.22 (-4.02, -0.43)	-2.16 (-4.06, -0.26)	-0.07 (-2.41, 2.27)

*To determine non-inferiority, the annualised median eGFR slopes were analysed by quantile regression using SAS PROC QUANTREG to obtain the corresponding 95% CI; non-inferiority was declared if the lower limit of the CI for the treatment difference (pegunigalsidase alfa – agalsidase beta) was ≥ -3.0 mL/min/1.73 m²/year.

†Baseline: males (n=29), females (n=23), ADA-positive (n=18), ADA-negative (n=34); number of subjects considered in the model at 24 months: overall (n=51), males (n=28), females (n=23), ADA-positive (n=17), ADA-negative (n=34).

‡Baseline: males (n=18), females (n=7), ADA-positive (n=8), ADA-negative (n=17); number of subjects considered in the model at 24 months: overall (n=25), males (n=18), females (n=7), ADA-positive (n=8), ADA-negative (n=17).

§Value above the predefined non-inferiority margin.

ADA, antidrug antibody; eGFR, estimated glomerular filtration rate; ITT, intent-to-treat.

pegunigalsidase alfa and 50% with agalsidase beta) than ADA-negative patients (15% and 12%, respectively). Infusion-related reaction rate was lower with pegunigalsidase alfa than agalsidase beta among ADA-positive patients (0.9 and 7.5 events/100 infusions, respectively) and ADA-negative patients (0.3 and 2.2 events/100 infusions, respectively).

Premedications

Most patients who initially received premedications (based on prior agalsidase beta treatment regimen) successfully reduced premedication use. There was a notable drop in premedication use from baseline to 24 months, from 21 (40.4%) to 3 (6.4%) patients receiving pegunigalsidase alfa and from 16 (64.0%) to 3 (12.5%) patients receiving agalsidase beta.

Antidrug antibodies

For males receiving pegunigalsidase alfa, 18 out of 29 (62%) were ADA-positive at baseline, and 10 out of 25 (40%) were ADA-positive at study end. For females, none were ADA-positive at baseline, and 1 out of 22 (5%) females was ADA-positive at study end (with treatment-emergent ADA). Neutralising antibodies were present in 17 out of 52 (33%) patients at baseline, and 7 out of 47 (15%) patients at study end. Treatment-emergent ADAs were present in 6 out of 52 (12%) patients (3 ADA-negative at baseline who became positive during treatment, and three titre boosted by more than fourfold during treatment). All IgG-positive patients tested negative for antibodies recognising the plant glycans, and three patients tested positive for antibodies to the PEG moieties of pegunigalsidase alfa (transitory response) throughout the study.

With agalsidase beta, 8 out of 18 (44%) males were ADA-positive at baseline, and 6 out of 16 (38%) were ADA-positive at study end. Neutralising antibodies were present in 7 out of 25 (28%) patients at baseline, and 6 out of 23 (26%) patients at study end. Treatment-emergent ADAs were present in 5 out of 25 (20%) patients, 3 of whom had treatment-induced ADAs, and two titre boosted.

Infusion setting and duration

A mean (SD; median) of 22.8 (17.0; 30.0) pegunigalsidase alfa infusions/patient were administered at home (46.0% of total infusions). With pegunigalsidase alfa, mean (range) infusion duration of completed infusions decreased from 3.1 (2.0–4.9) hours at first infusion to 1.6 (1.4–2.1) hours at 24 months; with agalsidase beta, means were 3.0 (2.6–3.3) hours at first infusion and 1.7 (1.4–3.2) hours at 24 months (no significant difference between groups at 24 months).

DISCUSSION

BALANCE demonstrated non-inferior renal efficacy and the potential for improved tolerability with pegunigalsidase alfa compared with agalsidase beta in patients with FD and deteriorating renal function. Clinically, patients were heterogeneous with multisystem involvement (median (range) of 5 (2–7) organs; >83% had cardiac involvement, and >97% had neurological involvement, based on FD medical history). The study population had received agalsidase beta for 1 year at minimum and for 6 years on average. Importantly, these patients had severe disease relative to previous agalsidase beta trials, which included patients with either low or high renal involvement¹⁶ or ERT-naïve patients with decreased creatinine clearance and without advanced, serious cardiac and neurological problems.³⁹

The primary endpoint was achieved, showing pegunigalsidase alfa was comparable to agalsidase beta. The non-inferiority margin was based on natural history information available at study development,^{40 41} type of population enrolled (ie, progressive renal impairment based on historical eGFR slope), inherent variability of eGFR as an outcome and limited sample size with rare disease. Post hoc analysis indicated that the imbalance in sex distribution at randomisation (not statistically significant) did not influence the final results.

Median eGFR slope improved in both study arms: from -6.7 and -7.8 mL/min/1.73 m²/year at baseline with pegunigalsidase alfa and agalsidase beta, respectively, to -2.5 and -2.2 mL/min/1.73 m²/year, respectively, at 2 years. This difference in




Figure 2 Plasma lyso-Gb3 over time in (A) males and (B) females. Boxes and whiskers represent the median and quartiles, with outliers as circles; 'X' represents the mean. Lyso-Gb3, globotriaosylsphingosine.

pre-enrolment and on-study slope could have resulted from differences in how historical and on-study creatinine values were derived (site-specific vs centralised, non-uniform vs predefined time intervals, variable vs predefined number of assessments, different vs same laboratory methodology). Of note, patients were managed before enrolment by standards of care that may have varied across the 29 study centres in 12 countries. Renal function in both arms stabilised, despite declining renal function at baseline and the unchanged agalsidase beta regimen. Nonetheless, baseline kidney function was equivalent between arms, supporting the validity of the randomisation. Medication adjustments would likely not affect results, because the use of ACEi/ARBs remained stable in both arms. Another possible explanation for this observation is the Hawthorne effect, a known phenomenon whereby clinical study participants benefit by being more closely observed than with standard care.⁴²

Median plasma lyso-Gb3 remained stable over the 2-year study in both treatment arms. As expected, sex stratification revealed that lyso-Gb3 concentrations were higher in males. Further post hoc analysis of outliers suggests baseline ADAs and UPCR >1 g/g may relate to changes in lyso-Gb3; these patients were slightly over-represented in the pegunigalsidase alfa arm. Generally, the clinical significance of the magnitude of lyso-Gb3 changes in both groups should be interpreted with caution.

Overall, pegunigalsidase alfa was well tolerated, aligning with previous findings in ERT-naïve and other switch patients.³⁰ There were substantially fewer infusion-related reactions with pegunigalsidase alfa than agalsidase beta, with a 7.8-fold difference in rate of infusion-related reactions (0.5/100 vs 3.9/100 infusions), and most were mild or moderate. There was one serious infusion-related reaction, a hypersensitivity event in one pegunigalsidase alfa IgE-positive patient. In other studies, 59%

Table 3 Plasma lyso-Gb3 from baseline to 24 months by treatment arm and sex

Plasma lyso-Gb3 (nM)*	Pegunigalsidase alfa			Agalsidase beta		
	Male (n=29)	Female (n=23)	Overall (n=52)	Male (n=18)	Female (n=7)	Overall (n=25)
Baseline, n	29	23	52	18	7	25
Mean (SE)	40.40 (5.50)	8.35 (0.68)	26.22 (3.78)	42.43 (8.71)	5.69 (1.10)	32.14 (7.08)
Median	30.7	8.4	15.2	23.7	4.4	17.6
Min, max	0.8, 143.9	2.8, 16.2	0.8, 143.9	8.9, 142.0	2.1, 10.4	2.1, 142.0
24 months, n	25	21	46	15	7	22
Mean (SE)	46.88 (6.34)	8.19 (0.95)	29.22 (4.48)	26.17 (4.33)	5.66 (1.06)	19.65 (3.60)
Median	34.4	8.9	18.8	20.5	4.9	15.3
Min, max	3.2, 139.4	2.4, 22.0	2.4, 139.4	6.2, 71.2	1.5, 9.7	1.5, 71.2
Change from baseline to 24 months, n†	25	21	46	15	7	22
Mean (SE)	5.90 (2.41)	0.19 (0.46)	3.30 (1.38)	-12.80 (6.93)	-0.03 (0.27)	-8.74 (4.85)
Median	5.3	0.1	1.15	-2.40	-0.30	-1.50
Min, max	-32.2, 32.7	-4.0, 5.8	-32.2, 32.7	-102.3, 2.4	-0.7, 0.9	-102.3, 2.4

*Normal range ≤ 2.4 nM.

†Data from five patients in the pegunigalsidase alfa arm and one in the agalsidase beta arm are missing due to early termination; one and two patients, respectively, are missing data due to missed visits. Lyso-Gb3, globotriaosylsphingosine.

of agalsidase beta-treated patients experienced infusion-related reactions (adverse reactions occurring on the infusion day), some of which were severe.¹⁸ The lower proportion of patients reporting infusion-related reactions in BALANCE (21% with pegunigalsidase alfa; 24% with agalsidase beta) relative to what is reported in agalsidase beta's prescribing information¹⁸ may be due to the selection of patients who received long-term treatment with agalsidase beta (vs initial 2 years from ERT initiation) and/or exclusion of patients who might have discontinued agalsidase beta due to infusion-related reactions. No deaths were reported. Two patients on pegunigalsidase alfa withdrew due to AEs; one hypersensitivity reaction and the other not treatment-related.

The overall rate of TEAEs was lower with pegunigalsidase alfa than agalsidase beta (572 vs 817 events/100 exposure-years). The only serious AE was the above-reported infusion-related reaction of hypersensitivity with the first pegunigalsidase alfa treatment which resolved the same day.

At baseline, 26 patients had pre-existing ADAs. The proportion of ADA-positive patients decreased slightly in both arms, from 35% at baseline to 23% at study end with pegunigalsidase alfa and from 32% to 26% with agalsidase beta. The proportion of patients with neutralising antibodies declined with pegunigalsidase alfa, from 33% to 15% compared with a change from 28% to 26% with agalsidase beta. Baseline reactivity to pegunigalsidase

Table 4 Treatment-emergent adverse events and infusion-related reactions within 2 hours of infusion

Variable	Pegunigalsidase alfa			Agalsidase beta		
	Male (n=29)	Female (n=23)	Overall (n=52)	Male (n=18)	Female (n=7)	Overall (n=25)
Any TEAE*						
Patient, n (%)	25 (86)	22 (96)	47 (90)	18 (100)	6 (86)	24 (96)
Events, n (rate*)	294 (545)	267 (605)	561 (572)	329 (922)	77 (549)	406 (817)
TEAE related to drug						
Patient, n (%)	15 (52)	6 (26)	21 (40)	9 (50)	2 (29)	11 (44)
Events, n (rate*)	33 (61)	9 (20)	42 (43)	55 (154)	21 (150)	76 (153)
Serious TEAE related to drug						
Patient, n (%)	1 (3)	0 (0)	1 (2)	0 (0)	0 (0)	0 (0)
Events, n (rate*)	1 (2)	0 (0)	1 (1)	0 (0)	0 (0)	0 (0)
TEAE leading to withdrawal						
Patient, n (%)	2 (7)	0 (0)	2 (4)	0 (0)	0 (0)	0 (0)
Events, n (rate*)	2 (4)	0 (0)	2 (2)	0 (0)	0 (0)	0 (0)
Related TEAE leading to withdrawal						
Patient, n (%)	1 (3)	0 (0)	1 (2)	0 (0)	0 (0)	0 (0)
Events, n (rate*)	1 (2)	0 (0)	1 (1)	0 (0)	0 (0)	0 (0)
Any infusion-related reactions						
Patient, n (%)	9 (31)	2 (9)	11 (21)	5 (28)	1 (14)	6 (24)
Events, n (rate†)	11 (0.8)	2 (0.2)	13 (0.5)	33 (4)	18 (5)	51 (4)
Mild or moderate infusion-related reactions						
Patient, n (%)	9 (31)	2 (9)	11 (21)	5 (28)	1 (14)	6 (24)
Events, n (rate†)	10 (0.7)	2 (0.2)	12 (0.5)	33 (4)	18 (5)	51 (4)
Severe infusion-related reactions						
Patient, n (%)	1 (3)	0 (0)	1 (2)	0 (0)	0 (0)	0 (0)
Events, n (rate†)	1 (0.1)	0 (0)	1 (0)	0 (0)	0 (0)	0 (0)

TEAEs include infusion-related reactions (defined as TEAEs beginning during or within 2 hours of infusion whose causality was assessed as definitely, probably or possibly treatment-related; these excluded injection site reactions, which were considered procedure-related).

*Per 100 exposure-years.

†Per 100 infusions.

TEAE, treatment-emergent adverse event.

alfa is explained by cross-reactivity to the enzyme components of the amino acid sequence shared between pegunigalsidase alfa and agalsidase beta.³¹ Recent studies show that in some patients, pre-existing ADAs against agalsidase alfa and agalsidase beta have less affinity and enzymatic inhibitory effects against pegunigalsidase alfa³¹; however, it is currently not possible to predict which ADA-positive patients may benefit from ERT switch and additional analysis needs to be performed. In BALANCE, only 3 patients per arm (6% (3/52) with pegunigalsidase alfa; 12% (3/25) with agalsidase beta) showed treatment-induced de novo ADAs, all with long-term agalsidase beta exposure. These rates are lower than what has been described in trials of naïve patients, as de novo ADAs typically occur in the first months of ERT initiation. For example, treatment-induced ADAs developed in 19% of naïve patients treated with pegunigalsidase alfa³⁰ and 83% of naïve patients receiving agalsidase beta.¹⁸ Direct comparison of ADA incidence across trials is also challenging due to the use of different ADA assays. Overall, these ADA findings should be interpreted with caution and with consideration of the patients' long-term ERT exposure.

In many cases, premedications were successfully reduced or discontinued, and mean infusion duration was similar between arms at 24 months with a maximum infusion duration of approximately 2 hours with pegunigalsidase alfa versus over 3 hours with agalsidase beta. This indicates at least one patient required prolonged infusion time with agalsidase beta to achieve good tolerability. Altogether, these findings support the safety of pegunigalsidase alfa, with infusions found to be equally safe for both drugs when administered at home compared with the study site.

BALANCE inclusion criteria selected for advanced disease in both males and females with FD, and as such, the participants represent a relatively homogeneous subgroup of patients affected by the disease. Differences between arms in ADAs and infusion-related reactions could have been underestimated because patients were already treated with agalsidase beta for an average of 6 years, and these occur most commonly in the first years of treatment. Due to the methodological challenges of interpretation, the current analysis does not compare the level of ADAs (titres) between arms and is limited to describing the proportion and trends of patients with ADAs within arms over time. Furthermore, FD is a heterogeneous disease with remaining unmet needs; the availability of new therapies can contribute to personalising patient care and potentially establishing combination regimens.^{43–45}

BALANCE is the first clinical trial in FD to be conducted with a double-blind, active-control design. Pegunigalsidase alfa was comparable to agalsidase beta based on annualised eGFR slope, an accepted surrogate for progression to end-stage kidney disease. Results demonstrated the potential for improved tolerability with less infusion-related reactions in some patients. Further detailed analysis of pegunigalsidase alfa immunogenicity is warranted. Most patients who completed the study (96%) enrolled in the open-label extension study for up to 7 years of pegunigalsidase alfa treatment. Pegunigalsidase alfa is approved in the EU and the USA, providing an important new treatment option for patients with FD.

Refer to online supplemental material 2 for a plain language summary of the study results.

Author affiliations

¹Department of Medicine, Division of Nephrology, The University of Alabama at Birmingham, Birmingham, Alabama, USA

²Lysosomal and Rare Disorders Research and Treatment Center, Inc, Fairfax, Virginia, USA

³Department of Human Genetics, Emory University School of Medicine, Atlanta, Georgia, USA

⁴Department of Pediatrics, University of Iowa Hospitals and Clinics, Iowa City, Iowa, USA

⁵Department of Pediatrics, Division of Medical Genetics, University of Utah Health, Salt Lake City, Utah, USA

⁶Department of Internal Medicine, School of Medicine, Charles University and General University Hospital, Prague, Czech Republic

⁷Lysosomal Storage Disorders Unit, Royal Free London NHS Foundation Trust and University College London, London, UK

⁸Department of Human Genetics, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, USA

⁹Department of Clinical Science, University of Bergen, Bergen, Norway

¹⁰Department of Pediatrics, Haukeland University Hospital, Bergen, Norway

¹¹Academisch Medisch Centrum Universiteit van Amsterdam, Amsterdam, The Netherlands

¹²Unidad de Investigación Traslacional. Hospital Universitario Miguel Servet, Instituto de Investigación Sanitaria Aragón (IIS Aragón), Zaragoza, Spain

¹³Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER-ISCIII), Zaragoza, Spain

¹⁴Department of Public Health, Università degli Studi di Napoli Federico II, Napoli, Italy

¹⁵Division of Medical Genetics, University of Versailles, Garches, France

¹⁶Baylor University Medical Center at Dallas, Dallas, Texas, USA

¹⁷Lysosomal Disorders Unit, Department of Medicine, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK

¹⁸Institute of Genomic Medicine and Rare Disorders, Semmelweis University Clinical Center, Budapest, Hungary

¹⁹Department of Pediatrics, UPMC Children's Hospital of Pittsburgh, Pittsburgh, Pennsylvania, USA

²⁰Mark Holland Metabolic Unit, Northern Care Alliance NHS Foundation Trust, Greater Manchester, UK

²¹Department of Pediatrics, Medical College of Wisconsin, Milwaukee, Wisconsin, USA

²²Department of Pediatrics, University of California San Diego, La Jolla, California, USA

²³Department of Pediatrics, University of California Irvine, Irvine, California, USA

²⁴Department of Internal Medicine, General Hospital Slovenj Gradec, Slovenj Gradec, Slovenia

²⁵Department of Endocrinology and Clinical Nutrition, University Hospital Zurich and University of Zurich, Zurich, Switzerland

²⁶Department of Diabetes, Endocrinology and Metabolism, Queen Elizabeth Hospital Birmingham, Birmingham, UK

²⁷Division of Medicine, Turku University Hospital, Turku, Finland

²⁸Phoenix Children's Hospital, Phoenix, Arizona, USA

²⁹University of Sunderland, Sunderland, UK

³⁰Infusion Associates, Grand Rapids, Michigan, USA

³¹Massachusetts General Hospital for Children, Boston, Massachusetts, USA

³²Protalix Biotherapeutics, Carmiel, Israel

³³Product Development, Protalix Biotherapeutics, Carmiel, Israel

³⁴Chiesi Farmaceutici SpA, Parma, Italy

³⁵DataSights, Haifa, Israel

Correction notice The article has been corrected since it was published online first. Supplementary file 1 contained some display errors, which have been resolved.

Acknowledgements We thank all the patients, their families, the investigators and all study staff involved in this study. The authors would like to acknowledge the Bioanalytical Laboratory at Protalix for assistance with the ADA analysis, and the CHUS laboratory for assistance with the Lyso-Gb3 analysis. Medical writing support was provided by Marisa DeGuzman, PhD, of Oxford PharmaGenesis, Inc., Newtown, PA, USA, and was funded by Chiesi USA, Inc.

Contributors EB-A, SA, RC, RR, AS, DGW: Study design and data interpretation. EB-A, SA, RC, RR, AS: Data analysis. ELW, OG-A, WRW, MH, JAB, NL, DH, PG, MJM, DO, RJH, CT, AL, PD, AJ, MM, BAB, VK, BV, AN, TG, AP, DPG, IK, JK, AM, SW, ML, KN, AK: Acquisition and interpretation of data. All authors had full access to the data, participated fully in drafting and revising the manuscript, and approved the final version of the manuscript. All authors have agreed both to be accountable for all aspects of the work and in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. DGW, as corresponding author, also serves as the guarantor together with ELW. Both accept full responsibility for the work and/or the conduct of the study, had access to the data, and controlled the decision to publish.

Funding Protalix Biotherapeutics, Inc. provided financial and material support for the research and, with the assistance of the study investigators, monitored the

conduct of the study, collected data from the investigative centers, and analysed and interpreted the data. The authors confirm independence from the sponsors, the content of the article has not been influenced by the sponsors, and the decision to submit it for publication was made by the authors independently. Professional medical writing and editing support were funded by Chiesi USA, Inc.

Competing interests ELW has consulting agreements and/or grants with Sanofi, Protalix, Chiesi, Idorsia, 4DMT, Amicus and Natera. OG-A has conducted contracted research, received consulting fees and/or served on advisory boards with Amicus, Freeline, Genentech, Protalix, Sangamo, Sanofi, Takeda, 4DMT and Avrobio. WRW has been or is currently involved in clinical trials and/or registries with Alexion, Amicus, BioMarin, Chiesi, Freeline, Idorsia, Orphazyme, Pfizer, Protalix, Sanofi, Sangamo, Takeda and 4DMT. He has received honoraria from Alexion, Amicus, BioMarin, Sanofi, Spark and Takeda, and research funding from Amicus and Takeda. MH received speaker-related fees from Protalix and has been, or is currently, involved in clinical trials with Sanofi, Sangamo, Avrobio, Protalix and Idorsia (no direct funding received for these trials because they are institution directed). JB receives research support from Avrobio, BioMarin Pharmaceutical, Chiesi Farmaceutici, Idorsia Pharmaceuticals, Pfizer, Protalix Biotherapeutics, Sangamo Therapeutics, Sanofi, Takeda, Travere Therapeutics; and has received a speaker honorarium from the Fabry Support and Information Group; and has participated in advisory boards for Chiesi USA, Sanofi and Takeda. NL receives research support from and has participated in advisory boards for Amicus, Astellas, Avrobio, BioMarin Pharmaceutical, Homology, Horizon, Moderna, Pfizer, Protalix Biotherapeutics, PTC Biotherapeutics, Reneo, Sanofi, Takeda and Ultragenyx (no direct funding is received because they are institution directed). DH has received honoraria for speaking and consulting fees for advisory boards from Protalix, Takeda, Sanofi, Freeline and Sangamo, administered through University College London consultants and used in part to support research in lysosomal storage diseases. PG has been involved in premarketing studies with Genzyme, Protalix and Idorsia, and has received grants from Sanofi-Genzyme and Takeda; monies received for these activities have been deposited into the Spanish Foundation for the Study and Treatment of Gaucher Disease (FEETEG) to contribute to the development of research in lysosomal storage disorders. MJM, DO, MM, BAB, JK, TG and KN have no disclosures. RJH has received consulting fees from Alexion, Amicus Therapeutics, Inc., Avrobio, Chiesi, Sangamo, Sanofi/Genzyme, and Takeda; advisory fees from Alexion, Amicus Therapeutics, Inc., and Sanofi/Genzyme; speakers' bureau fees from Alexion and Sanofi/Genzyme and grants/research funding from Alexion, Amicus Therapeutics, Inc., Idorsia, Protalix, Sangamo, Sanofi/Genzyme and Takeda. CT has received honoraria, travel support, and/or participated as an investigator in clinical studies supported by Protalix, Sanofi, Idorsia, Takeda, Amicus, Freeline and Acelink. All received honoraria went to her institution Haukeland University Hospital. AL has received consultancy and speaker's honoraria from Amicus Therapeutics, Sanofi, Takeda, 4DMT and Chiesi. PD has been a paid consultant with Sanofi; received speaker honoraria from Sanofi and Takeda and participated in an advisory board with Protalix. AJ has received a grant from Amicus and consultancy and speaker's honoraria from Takeda, Sanofi and Amicus. VK is an advisory board member for the Sanofi-Genzyme North American Pompe Registry. She is the principal investigator for Sanofi-Genzyme Lysosomal Storage Disease registry at UC Irvine. She has received education grants, lecturing honoraria, and research support from Sanofi-Genzyme. She participates as an investigator in clinical studies supported by Protalix, Sanofi, Idorsia, Chiesi Farmaceutici and Sangamo. BV has received honoraria, travel and accommodation funding from Greenovation Biotech GmbH, Sanofi, Takeda, Amicus, Chiesi and Swixx, and is a member of the EU Advisory Board of Fabry Registry, sponsored by Sanofi. AN received lecturing honoraria and research support from Takeda, Amicus and Sanofi/Genzyme. AP received travel expenses and grants from Takeda, Sanofi, Amicus and Chiesi. DPG has received consulting honoraria from Chiesi, Idorsia Pharmaceuticals, Sanofi and Takeda, and speaker honoraria and travel support from Sanofi and Takeda. IK has received lecture, travel and consulting fees from Amicus, Chiesi, Bayer, Boehringer-Ingelheim, Sanofi-Genzyme and Takeda-Shire. AM has received non-financial support from Chiesi Pharmaceuticals, during the conduct of the study; personal fees from AstraZeneca, Bayer Pharmaceuticals and Janssen Pharmaceuticals, outside the submitted work. AK has received honoraria, travel support and/or participated as an investigator in clinical studies supported by Protalix, Sanofi and Idorsia and on advisory meetings for Protalix, Sanofi, Takeda, Amicus and Chiesi. SW has been a paid consultant to Protalix. ML is involved in premarketing studies with Sanofi-Genzyme, Protalix/Chiesi and Idorsia. Financial arrangements were made through AMC Research BV. No fees, travel support or grants were obtained from the pharmaceutical industry. EBA and RC were full-time employees of Protalix Biotherapeutics at the time of study conduct and analysis and are now consultants to Protalix Biotherapeutics. SA is a full-time employee of Protalix Biotherapeutics. RR is a full-time employee of Chiesi Farmaceutici S.p.A. AS was a paid consultant to Protalix at the time of study conduct and analysis and is currently a paid consultant to Chiesi USA, Inc. DGW is involved in clinical trials/registries/consulting with Amicus, Chiesi, Idorsia and Protalix.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. We will approve or deny data requests from external parties on a case-by-case basis. Chiesi reserves the right to deny requests for all legally appropriate reasons. Data requests that risk sharing participant-level data or proprietary information will not be approved.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Aleš Linhart <http://orcid.org/0000-0002-3372-7850>

Derralynn A Hughes <http://orcid.org/0000-0003-4531-9173>

Dominique Paul Germain <http://orcid.org/0000-0002-8355-007X>

Albina Nowak <http://orcid.org/0000-0002-4119-5516>

David G Warnock <http://orcid.org/0000-0003-4112-0677>

REFERENCES

- Vardarli I, Rischpler C, Herrmann K, *et al*. Diagnosis and screening of patients with fabry disease. *Ther Clin Risk Manag* 2020;16:551–8.
- Bokhari SRA, Zulfiqar H, Hariz A. *Fabry Disease*. Treasure Island (FL): StatPearls, 2022.
- Ezgu F, Alpsoy E, Bicik Bahcebasi Z, *et al*. Expert opinion on the recognition, diagnosis and management of children and adults with fabry disease: a multidisciplinary Turkey perspective. *Orphanet J Rare Dis* 2022;17:90.
- Arends M, Wanner C, Hughes D, *et al*. Characterization of classical and nonclassical fabry disease: a multicenter study. *J Am Soc Nephrol* 2017;28:1631–41.
- Oliveira JP, Ferreira S. Multiple phenotypic domains of fabry disease and their relevance for establishing genotype-phenotype correlations. *Appl Clin Genet* 2019;12:35–50.
- Mehta A, Hughes DA, *et al*. Fabry disease. In: Adam MP, Everman DB, Mirzaa GM, eds. *GeneReviews(R)*. Seattle (WA), 1993.
- Juchniewicz P, Kloska A, Tyłki-Szymańska A, *et al*. Female fabry disease patients and X-Chromosome inactivation. *Gene* 2018;641:259–64.
- Echevarria L, Benistan K, Toussaint A, *et al*. X-Chromosome inactivation in female patients with fabry disease. *Clin Genet* 2016;89:44–54.
- Řeboun M, Sikora J, Magner M, *et al*. Pitfalls of X-Chromosome inactivation testing in females with fabry disease. *Am J Med Genet A* 2022;188:1979–89.
- Pinto LLC, Vieira TA, Giugliani R, *et al*. Expression of the disease on female carriers of X-linked lysosomal disorders: a brief review. *Orphanet J Rare Dis* 2010;5:14.
- Fuller M, Mellett N, Hein LK, *et al*. Absence of A-Galactosidase cross-correction in Fabry heterozygote cultured skin fibroblasts. *Mol Genet Metab* 2015;114:268–73.
- Azevedo O, Gago MF, Miltenberger-Miltenyi G, *et al*. Fabry disease therapy: state-of-the-art and current challenges. *Int J Mol Sci* 2020;22:206.
- ELFABRIO (Pegunigalsidase Alfa-lwxj) injection, for intravenous use. ELFABRIO (Pegunigalsidase Alfa-lwxj) injection, for intravenous use Ed. Parma, Italy Chiesi Farmaceutici S.p.A.; 2023.
- Mehta A, West ML, Pintos-Morell G, *et al*. Therapeutic goals in the treatment of fabry disease. *Genet Med* 2010;12:713–20.
- Wanner C, Feldt-Rasmussen U, Jovanovic A, *et al*. Cardiomyopathy and kidney function in agalsidase beta-treated female fabry patients: a pre-treatment vs. post-treatment analysis. *ESC Heart Fail* 2020;7:825–34.
- Germain DP, Charrow J, Desnick RJ, *et al*. Ten-year outcome of enzyme replacement therapy with agalsidase beta in patients with fabry disease. *J Med Genet* 2015;52:353–8.
- Beck M, Ramaswami U, Hernberg-Ståhl E, *et al*. Twenty years of the Fabry outcome survey (FOS): insights, achievements, and lessons learned from a global patient registry. *Orphanet J Rare Dis* 2022;17:238.
- FABRAZYME® (Agalsidase beta) for injection, for intravenous use. FABRAZYME® (Agalsidase beta) for injection, for intravenous use ed. Cambridge, MA, USA Genzyme Corporation; 2021.
- van der Veen SJ, Vlietstra WJ, van Dussen L, *et al*. Predicting the development of anti-drug antibodies against recombinant alpha-galactosidase a in male patients with classical fabry disease. *IJMS* 2020;21:5784.

- 20 van der Veen SJ, van Kuilenburg ABP, Hollak CEM, *et al.* Antibodies against recombinant alpha-galactosidase a in fabry disease: subclass analysis and impact on response to treatment. *Mol Genet Metab* 2019;126:162–8.
- 21 Rombach SM, Aerts J, Poorthuis B, *et al.* Long-term effect of antibodies against infused alpha-galactosidase a in fabry disease on plasma and urinary (Lyso)Gb3 reduction and treatment outcome. *PLoS One* 2012;7:e47805.
- 22 Lenders M, Brand E. Fabry disease: the current treatment landscape. *Drugs* 2021;81:635–45.
- 23 Hughes DA, Elliott PM, Shah J, *et al.* Effects of enzyme replacement therapy on the cardiomyopathy of Anderson-fabry disease: a randomised, double-blind, placebo-controlled clinical trial of agalsidase alfa. *Heart* 2008;94:153–8.
- 24 Schiffmann R, Kopp JB, Austin HA, *et al.* Enzyme replacement therapy in fabry disease: a randomized controlled trial. *JAMA* 2001;285:2743–9.
- 25 Eng CM, Guffon N, Wilcox WR, *et al.* Safety and efficacy of recombinant human alpha-galactosidase a replacement therapy in fabry's disease. *N Engl J Med* 2001;345:9–16.
- 26 Wilcox WR, Banikazemi M, Guffon N, *et al.* Long-term safety and efficacy of enzyme replacement therapy for fabry disease. *Am J Hum Genet* 2004;75:65–74.
- 27 Nicholls K, Bleasel K, Becker G. Severe infusion reactions to fabry enzyme replacement therapy: rechallenge after tracheostomy. *JIMD Rep* 2012;5:109–12.
- 28 Ruderfer I, Shulman A, Kizhner T, *et al.* Development and analytical characterization of pegunigalsidase alfa, a chemically cross-linked plant recombinant human alpha-galactosidase-a for treatment of fabry disease. *Bioconjug Chem* 2018;29:1630–9.
- 29 Kizhner T, Azulay Y, Hainrichson M, *et al.* Characterization of a chemically modified plant cell culture expressed human α -galactosidase-a enzyme for treatment of fabry disease. *Mol Genet Metab* 2015;114:259–67.
- 30 Schiffmann R, Goker-Alpan O, Holida M, *et al.* Pegunigalsidase alfa, a novel pegylated enzyme replacement therapy for fabry disease, provides sustained plasma concentrations and favorable pharmacodynamics: a 1-year phase 1/2 clinical trial. *J Inherit Metab Dis* 2019;42:534–44.
- 31 Lenders M, Pollmann S, Terlinden M, *et al.* Pre-existing anti-drug antibodies in fabry disease show less affinity for pegunigalsidase alfa. *Mol Ther Methods Clin Dev* 2022;26:323–30.
- 32 Replagal® (agalsidase alfa). *Replagal*® (agalsidase alfa) ed. Dublin, Ireland: Takeda Pharmaceuticals, 2022.
- 33 Dostálová G, Hulková H, Linhart A. Anderson-Fabry disease: no histological signs of pathological accumulation in arterial and venous endothelium during pegunigalsidase alfa therapy. *Kardiol Pol* 2021;79:1385–6.
- 34 Levey AS, Stevens LA, Schmid CH, *et al.* A new equation to estimate glomerular filtration rate. *Ann Intern Med* 2009;150:604–12.
- 35 Warnock DG, Thomas CP, Vujkovic B, *et al.* Antiproteinuric therapy and fabry nephropathy: factors associated with preserved kidney function during agalsidase-beta therapy. *J Med Genet* 2015;52:860–6.
- 36 Boutin M, Auray-Blais C. Multiplex tandem mass spectrometry analysis of novel plasma lyso-gb₃-related analogues in fabry disease. *Anal Chem* 2014;86:3476–83.
- 37 Boutin M, Lavoie P, Abaoui M, *et al.* Tandem mass spectrometry quantitation of Lyso-Gb₃ and six related analogs in plasma for fabry disease patients. *Curr Protoc Hum Genet* 2016;90:17.
- 38 Ortiz A, Kanters S, Hamed A, *et al.* Agalsidase beta treatment SLOWS estimated glomerular filtration rate loss in classic fabry disease patients: results from an individual patient data meta-analysis. *Clin Kidney J* 2021;14:1136–46.
- 39 Banikazemi M, Bultas J, Waldek S, *et al.* Agalsidase-beta therapy for advanced fabry disease: a randomized trial. *Ann Intern Med* 2007;146:77–86.
- 40 Schiffmann R, Warnock DG, Banikazemi M, *et al.* Fabry disease: progression of nephropathy, and prevalence of cardiac and cerebrovascular events before enzyme replacement therapy. *Nephrol Dial Transplant* 2009;24:2102–11.
- 41 Branton M, Schiffmann R, Kopp JB. Natural history and treatment of renal involvement in fabry disease. *J Am Soc Nephrol* 2002;13 Suppl 2:S139–43.
- 42 Braunholtz DA, Edwards SJ, Lilford RJ. Are randomized clinical trials good for us (in the short term)? Evidence for a "trial effect." *J Clin Epidemiol* 2001;54:217–24.
- 43 Xu S, Lun Y, Brignol N, *et al.* Coformulation of a novel human α -galactosidase a with the pharmacological chaperone AT1001 leads to improved substrate reduction in fabry mice. *Mol Ther* 2015;23:1169–81.
- 44 Benjamin ER, Khanna R, Schilling A, *et al.* Co-administration with the pharmacological chaperone AT1001 increases recombinant human alpha-galactosidase a tissue uptake and improves substrate reduction in fabry mice. *Mol Ther* 2012;20:717–26.
- 45 Iacobucci I, Hay Mele B, Cozzolino F, *et al.* Enzyme replacement therapy for FABRY disease: possible strategies to improve its efficacy. *Int J Mol Sci* 2023;24:4548.
- 46 Wanner C, Arad M, Baron R, *et al.* European expert consensus statement on therapeutic goals in fabry disease. *Mol Genet Metab* 2018;124:189–203.