



Uniform negative immersions and the coherence of one-relator groups

Larsen Louder¹ · Henry Wilton²

Received: 6 September 2021 / Accepted: 9 February 2024
© Crown 2024

Abstract

Previously, the authors proved that the presentation complex of a one-relator group G satisfies a geometric condition called *negative immersions* if every two-generator, one-relator subgroup of G is free. Here, we prove that one-relator groups with negative immersions are coherent, answering a question of Baumslag in this case. Other strong constraints on the finitely generated subgroups also follow such as, for example, the co-Hopf property. The main new theorem strengthens negative immersions to *uniform negative immersions*, using a rationality theorem proved with linear-programming techniques.

1 Introduction

Definition 1.1 (Coherence) A group G is said to be *coherent* if all its finitely generated subgroups are finitely presentable.

A notorious question of Baumslag [3, p. 76] asks whether every one-relator group $G = F/\langle\langle w \rangle\rangle$ is coherent. It is a curious fact that the many known examples of one-relator groups with pathological properties — for instance, Baumslag–Solitar groups, the Baumslag–Gersten group [2], or the recent examples of Gardam–Woodhouse [14] — all have two generators. Our first theorem resolves Baumslag’s question, as long as such subgroups are excluded.

Theorem A *Let $G = F/\langle\langle w \rangle\rangle$ be a one-relator group. If every two-generator, one-relator subgroup of G is free, then G is coherent.*

✉ H. Wilton
h.wilton@maths.cam.ac.uk

L. Louder
l.louder@ucl.ac.uk

¹ Department of Mathematics, University College London, Gower Street, London WC1E 6BT, UK

² DPMMS, Centre for Mathematical Sciences, Wilberforce Road, Cambridge CB3 0WB, UK

The case where G has torsion was proved independently by the authors [26] and Wise [44]. Sapir and Spakulova showed that a generic one-relator group with at least three generators is coherent [33].

The hypothesis of Theorem A — that every two-generator, one-relator subgroup is free — appears on its face to be difficult to check. However, the results of [27] imply that it is equivalent to other, more effective, conditions. The following theorem, which follows from [27, Theorems 1.3 and 1.5], summarises these equivalences. (The relevant definitions are explained below.) By convention, the relator w is always assumed to be non-trivial.

Theorem 1.2 ([27]) *Let $G = F/\langle\langle w \rangle\rangle$ be a one-relator group, and let X be the natural presentation complex. The following conditions are equivalent:*

- (i) *the primitivity rank $\pi(w) > 2$;*
- (ii) *X has negative immersions;*
- (iii) *every two-generator subgroup of G is free;*
- (iv) *every two-generator, one-relator subgroup of G is free.*

Since it is the briefest to state, if any of these hold we say that X or G has *negative immersions*.

Remark 1.3 Puder's primitivity rank $\pi(w)$ is a positive integer (or infinite if w is a basis element) [30]. We will not need its definition here, but we make the following remarks.

- (i) The primitivity rank $\pi(w)$ can be computed algorithmically: see [27, Lemma 6.4] or [30, Appendix A] for algorithms, and see [9] for extensive computations.
- (ii) Puder also showed that a generic word w in a free group of rank r has $\pi(w) = r$ [31, Corollary 8.3], so generic one-relator groups with more than two generators have negative immersions. (A more refined result was later proved by Kapovich [21].)
- (iii) Combining [31, Corollary 8.3] with Theorems A and 1.2 recovers the result of Sapir–Spakulova that random one-relator groups with at least three generators are coherent. Instead of [31, Corollary 8.3], one can also appeal to the work of Arzhantseva–Olshanskii, who proved that a random group with at least 3 generators is 2-free [1].

Condition (ii) of Theorem 1.2 is a geometric condition on the 2-complex X . An *immersion* is a combinatorial map of 2-complexes that is locally injective. Roughly, a 2-complex X has *negative immersions* if, for any finite, connected 2-complex Y immersing into X , either

- (i) $\chi(Y) < 0$, or
- (ii) Y is homotopic to a graph.

See Definition 3.20 for a precise definition. As usual, $\chi(Y)$ denotes the Euler characteristic of Y . The negative immersions property is motivated by an analogy with the concept of *non-positive immersions*, which is defined similarly but with (i) replaced

by the inequality $\chi(Y) \leq 0$. The presentation complex of any torsion-free one-relator group has non-positive immersions [18, 25].

The proof of Theorem A leads to various other strong constraints on the subgroups of one-relator groups with negative immersions. The next theorem summarises these.

Theorem B *Let $G = F/\langle\langle w \rangle\rangle$ be a one-relator group with negative immersions.*

- (i) *Every finitely generated, one-ended subgroup H of G is co-Hopfian, i.e. H is not isomorphic to a proper subgroup of itself. In particular, if G itself is one-ended then it is co-Hopfian.*
- (ii) *For any integer r , there are only finitely many conjugacy classes of finitely generated, one-ended subgroups H of G such that the abelianisation of H has rational rank at most r .*
- (iii) *Every finitely generated non-cyclic subgroup H of G is large in the sense of Pride, i.e. there is a subgroup H_0 of finite index in H that surjects a non-abelian free group.*

In an earlier paper, the authors conjectured that every one-relator group G with negative immersions is hyperbolic [27, Conjecture 1.9]. Theorem B proves various consequences of this conjecture. Specifically, item (i) is consistent with Sela’s theorem that one-ended hyperbolic groups are co-Hopfian [35, Theorem 4.4], while item (ii) is a stronger version of a subgroup rigidity theorem of Gromov [15, 5.3.C’], Rips–Sela [32, Theorem 7.1] and Delzant [12]. Note, however, that items (ii) and (iii) are not implied simply by the fact that G is hyperbolic.

Recently, Linton has proved the authors’ conjecture [24, Theorem 8.2], making use of Theorem B in his proof. Linton’s theorem is especially remarkable in light of the recent discovery by Italiano–Martelli–Migliorini of higher-dimensional non-hyperbolic groups of finite type that do not contain Baumslag–Solitar subgroups [22, Corollary 3].

The main new technical ingredient in the proof of Theorem A is a uniform version of negative immersions, which provides a negative upper bound for a normalised version of the Euler characteristic. Roughly, a 2-complex X has *uniform negative immersions* if there is $\epsilon > 0$ such that for any finite, connected 2-complex Y immersing into X , either

(i)

$$\frac{\chi(Y)}{\#\{2\text{-cells of } Y\}} \leq -\epsilon,$$

or

- (ii) Y can be simplified by a homotopy. (In the language developed below, Y is *reducible*.)

The reader is referred to Definition 3.24 for a precise definition.

Remark 1.4 In Wise’s work on the coherence of one-relator groups with torsion [44] (and also implicitly in the authors’ [26]), a key role is played by a property that Wise also calls ‘negative immersions’, but that is closer in spirit to (indeed, stronger than)

the notion of *uniform* negative immersions used here. See Sect. 3.4 for a comparison between the notion of (uniform) negative immersions used in this paper and Wise's notion.

The main new technical ingredient towards the proof of Theorem A is the following result, which establishes uniform negative immersions for the 2-complexes of interest to us.

Theorem C *Let X be the presentation complex of a one-relator group. If X has negative immersions then it has uniform negative immersions.*

The deduction of Theorem A from Theorem C is similar to [26] and [44], with an additional lemma (Lemma 3.10) to account for the subtle nature of item (ii) in the definition.

In the terminology of [40], Theorem C asserts that one-relator presentation complexes with negative immersions have *negative irreducible curvature*.

Theorem C follows from a rationality theorem, Theorem 4.12, similar to those that appear in Calegari's proof of the rationality of stable commutator length in free groups [7] and in the second author's work on surface subgroups of graphs of free groups [39].

The first idea is to expand the category of maps considered from immersions $Y \looparrowright X$ to the more general class of *face-essential* maps $f : Y \rightarrow X$. This class of maps allows branching in the centres of the 2-cells, and thus each 2-cell C of Y comes with a well-defined *degree* $\deg_C(f)$. The degree of the map f is then the sum of the degrees of the 2-cells of Y , and we adjust the relevant quantities to take it into account. The role of Euler characteristic is taken by the *total curvature*

$$\tau(Y) := \deg(f) + \chi(Y_{(1)})$$

where $\chi(Y_{(1)})$ is the Euler characteristic of the 1-skeleton of Y , and the role of the number of 2-cells is taken by the degree $\deg(f)$. Note that, when f is an immersion, these reduce to the usual quantities: $\tau(Y) = \chi(Y)$, while $\deg(f)$ is the number of 2-cells of Y . With these adjustments, the supremum of the quantity

$$\frac{\tau(Y)}{\deg(f)}$$

over all face-essential maps $f : Y \rightarrow X$ with Y irreducible can be computed as a solution to a rational linear programming problem; this is the content of Theorem 4.12.

In particular, this supremum is in fact a maximum. The results of [27] then imply that this maximum is strictly negative, so provides the uniform bound $-\epsilon$ needed to prove Theorem C.

This proof provides no information about the value of ϵ , but we make the following conjecture. A face-essential map of 2-complexes $f : Y \rightarrow X$ is called *essential* if, for each component of Y , the induced map on 1-skeleta induces an injective homomorphism on fundamental groups.

Conjecture 1.5 *Let $G = F/\langle\langle w \rangle\rangle$ be a one-relator group and let X be the associated presentation complex. The supremum of the quantity*

$$\frac{\tau(Y)}{\deg(f)}$$

across all essential maps f from irreducible 2-complexes Y to X is equal to $2 - \pi(w)$.

In an earlier version of this paper, we stated a stronger version of Conjecture 1.5 for the supremum across all face-essential maps. The next example shows that this stronger conjecture is false.

Example 1.6 Let X be the presentation complex of

$$\langle a, b, c, d \mid a^4 b^2 c^2 d^2 a^3 b^2 c^2 d^2 \rangle$$

and let Y be the presentation complex of

$$\langle x, y, z \mid x^2 y z y x z \rangle.$$

Write w for the relator of X and u for the relator of Y . The homomorphism of free groups defined by

$$x, y \mapsto a, z \mapsto ab^2c^2d^2$$

sends u to w and so, after subdividing the 1-skeleton of Y appropriately, defines a combinatorial map of 2-complexes $f : Y \rightarrow X$. Since this morphism sends the single 2-cell of Y homeomorphically to the single 2-cell of X , f is face essential (see Definition 2.18) and $\deg(f) = 1$. Furthermore, the Whitehead graph of Y consists of two squares glued along an edge, so Y is (visibly) irreducible (see Definition 3.1).

However,

$$\frac{\tau(Y)}{\deg(f)} = \frac{\deg(f) + \chi(Y_{(1)})}{\deg(f)} = -1,$$

while a calculation using Puder’s algorithm shows that $\pi(w) = 4$ [8]. Thus, f provides an example of a face-essential map from an irreducible 2-complex Y such that

$$\frac{\tau(Y)}{\deg(f)} > 2 - \pi(w).$$

In particular, the ‘essential’ hypothesis in Conjecture 1.5 cannot be relaxed to ‘face-essential’.

Conjecture 1.5 has precursors in the literature. The following conjecture of Heuer posits a surprising relationship between stable commutator length and Puder’s primitivity rank [19, Conjecture 6.3.2]. (The same conjecture was made independently by Hanany and Puder [17, Conjecture 1.14].)

Conjecture 1.7 (Heuer's conjecture) *If w is a non-trivial element of the commutator subgroup of a free group F then*

$$2 \operatorname{scl}(w) \geq \pi(w) - 1.$$

Conjecture 1.5 implies Heuer's conjecture, and indeed Theorem 1.2, together with Calegari's rationality theorem for stable commutator length [7], unconditionally implies a consequence of Heuer's conjecture.

Corollary D *If w is an element of the commutator subgroup of a free group F and $\pi(w) > 2$ then*

$$2 \operatorname{scl}(w) > 1.$$

For context, Duncan–Howie proved that $\operatorname{scl}(w) \geq 1/2$ for any non-trivial element of $[F, F]$ [13]. The statement of Corollary D was noted by Heuer and Löh [20, Question 1.3(4)], but we will give a complete proof in §3.5 below. The proof of Corollary D and the deduction of Corollary 1.7 from Conjecture 1.5 both use Proposition 3.28, which summarises an argument of Calegari [5, Lemma 2.7].

The paper is structured as follows. Section 2 is devoted to setting up the framework in which we describe 2-complexes and maps between them. It is convenient to work with *pre-complexes*, which describe pieces that can be glued together to form complexes. Section 3 describes the notion of *irreducible complexes* that plays a crucial role in our definitions, and enables us to define (uniform) negative immersions. In Sect. 4, we set up the linear system needed to prove the rationality theorem, from which Theorem C follows. The brief Sect. 5 explains why one-relator groups with torsion also have uniform negative immersions. Finally, theorems about the subgroup structure of the fundamental groups of 2-complexes with uniform negative immersions are proved in Sect. 6, and Theorems A and B follow immediately.

2 Preliminaries

In this section, we give some foundational definitions in order to fix notation.

2.1 Graphs and pre-graphs

Graphs are our fundamental objects of study. A subgraph is always a closed subspace in any of the standard combinatorial frameworks, but it is also convenient to discuss open sub-objects of graphs. We therefore work with the slightly more general notion of *pre-graphs*.

Definition 2.1 (Pre-graph) *A pre-graph is a tuple*

$$G = (V_G, E_G, \iota_G, \tau_G) = (V, E, \iota, \tau)$$

where:

- (i) V and E are sets (the *vertices* and *edges* of G , respectively);
- (ii) $E_G^l = E^l$ and $E_G^r = E^r$ are subsets of E ; and
- (iii) $\iota: E^l \rightarrow V$ and $\tau: E^r \rightarrow V$ are maps.

The subscripts will be suppressed only when there is no danger of confusion. A *morphism* f from a pre-graph G to a pre-graph H consists of maps $f: V_G \rightarrow V_H$ and $f: E_G \rightarrow E_H$ such that $f(E_G^l) \subseteq E_H^l$ and $f(E_G^r) \subseteq E_H^r$, and such that $\iota_H \circ f = f \circ \iota_G$ and $\tau_H \circ f = f \circ \tau_G$.

The *boundary* of a pre-graph G is defined to be

$$\partial G = E \setminus E^l \cap E^r.$$

Definition 2.2 (Graph) An (oriented) *graph* is a pre-graph with empty boundary.

The reader is referred to the classic paper of Stallings [36] for the standard theory of the topology of graphs. Stallings phrases his arguments in terms of Serre graphs, which differ slightly from the model of graphs we adopt here, but his arguments are easily adapted to our context and we leave the details to the reader. Stallings does not use pre-graphs, but again the theory is easy to adapt.

Definition 2.3 (Immersion) A morphism $f: G \rightarrow H$ of pre-graphs is an *immersion* if, for all edges e and e' of G , if $\alpha(e) = \alpha(e')$ for some $\alpha \in \{\iota_G, \tau_G\}$ and $f(e) = f(e')$, then $e = e'$. An immersion $f: G \rightarrow H$ of pre-graphs is *closed* if $f(\partial G) \subseteq \partial H$.

The standard fiber product construction for graphs (see [36, 1.3]) also makes sense for pre-graphs.

Definition 2.4 Consider a pair of morphisms of pre-graphs

$$f_i: G_i \rightarrow H$$


where $i = 1, 2$. The *fiber product* $G_1 \times_H G_2$ has vertex (respectively edge) set defined to be the subset of $V_{G_1} \times V_{G_2}$ (respectively, $E_{G_1} \times E_{G_2}$) on which f_1 and f_2 agree. The attaching maps are then defined in the natural way wherever they make sense.

This definition coincides with the standard definition whenever G_1, G_2 and H are graphs. It enjoys various well-known properties, which are left as easy exercises: it is a pullback in the category of pre-graphs, the pullback of an immersion is an immersion, etc.

2.2 Complexes and pre-complexes

Our graphs will often be the 1-skeleta of 2-dimensional complexes, which can be formalised as immersions from disjoint unions of circles to graphs, where the circles correspond to the boundaries of the 2-cells (which we will always call *faces*). As in the case of graphs, we will sometimes want to discuss open sub-objects, and we therefore introduce *pre-complexes*. In this case, slightly more general pre-graphs can arise as the boundaries of faces.

Fig. 1 A pre-complex (G, S, w) : G is a theta graph with two half-edges sticking off, and S is the union of two open arcs and one cycle. The map w is a closed immersion, since it maps ∂S to ∂G



Definition 2.5 A connected finite pre-graph is a *pre-cycle* if every vertex has exactly two incident edges. A pre-cycle is a *cycle* if it has empty boundary, and is an *open arc* otherwise.

We can now define a pre-complex using immersions of pre-cycles.

Definition 2.6 A (2-dimensional) pre-complex X is a tuple

$$(G_X, S_X, w_X)$$

where

$$G_X = (V_X, E_X, \iota_X, \tau_X)$$

is a pre-graph, S_X is a union of pre-cycles, and $w_X : S_X \looparrowright G_X$ is a closed immersion. The pre-graph G_X is the 1-skeleton of X , S_X is the collection of (*pre-*)*faces* of X and w_X is the *attaching map*. For brevity, we will suppress the subscript when there is no danger of confusion.

A pre-complex X is a (2-dimensional) *complex* if the 1-skeleton G_X is a graph. Note that, in this case, every component of S_X is a cycle.

See Figure 1 for an example of a pre-complex.

Graphs and complexes have natural topological realisations, which we denote here with bold letters: the realisation of a graph G is a topological space denoted by \mathbf{G} and the realisation of a complex X is a topological space denoted by \mathbf{X} . In both cases, a morphism f functorially induces a continuous map of realisations \mathbf{f} .

2.3 Asterisks and regular neighbourhoods

It is often fruitful to study graphs and 2-complexes via the local structure of vertices. Pre-complexes provide a convenient framework for this.

Definition 2.7 An *asterisk* is a pre-graph with one vertex, such that $E = E^\iota \sqcup E^\tau$. If the 1-skeleton of a pre-complex X is an asterisk then X is called an *asterisk pre-complex*.

Every vertex of a pre-graph or pre-complex now has a regular neighbourhood, based on an asterisk.

Definition 2.8 Let G be a pre-graph and v be a vertex of G . The *regular neighbourhood* of v is an asterisk $N_G(v)$, with vertex set $\{v\}$ and edge set $\iota^{-1}(v) \sqcup \tau^{-1}(v)$.

(Note that the disjoint union here is important: an edge e with $\iota(e) = \tau(e) = v$ defines two edges in $N_G(v)$.) The attaching maps are the unique maps $\iota^{-1}(v) \rightarrow \{v\}$ and $\tau^{-1}(v) \rightarrow \{v\}$

Now let $X = (G, S, w)$ be a pre-complex and v be a vertex of X . The *regular neighbourhood* $N = N_X(v)$ of v is a pre-complex with 1-skeleton $G_N = N_G(v)$, i.e. the regular neighbourhood of v in the 1-skeleton G . The faces of $N_X(v)$ are pulled back from X , so $S_N = S \times_G G_N$, and the attaching map w_N is the pullback of w , i.e. the natural projection $S_N \rightarrow G_N$.

The regular neighbourhood of a vertex v carries the same information as the *link* or *Whitehead graph* of the vertex. Indeed, any asterisk pre-complex defines a Whitehead graph.

Definition 2.9 Let $X = (G, S, w)$ be an asterisk pre-complex with unique vertex v . The corresponding *Whitehead graph* has vertex-set equal to the edges of X and edge-set equal to $w^{-1}(v)$. Each $u \in w^{-1}(v)$ is incident at exactly two edges $\epsilon_+(v)$ and $\epsilon_-(v)$ of S , and the attaching maps of the Whitehead graph adjoin u to $w(\epsilon_+(v))$ and $w(\epsilon_-(v))$.

For v a vertex of a complex X , the Whitehead graph of the regular neighbourhood $N(v)$ coincides with the Whitehead graph $\text{Wh}(v)$ of [39]. This in turn generalises the graph used by Whitehead in his solution to the orbit problem for automorphism groups of free groups [38]. See also [10, 28] for related recent uses of Whitehead graphs.

Edges of pre-complexes also have regular neighbourhoods.

Definition 2.10 Let $X = (G, S, w)$ be a pre-complex and e be an edge of X . The *regular neighbourhood* of e is a pre-complex $N = N_X(e)$ with 1-skeleton G_N equal to the pre-graph that has no vertices and a single edge $\{e\}$. The faces S_N of $N_X(e)$ consists of the pre-image $w^{-1}(e)$, and the attaching map w is the restriction of w to S_N .

Remark 2.11 A morphism of pre-complexes or pre-graphs induces morphisms on regular neighbourhoods: $f_* : N(x) \rightarrow N(f(x))$, for x a vertex or edge of the domain.

2.4 Various kinds of maps

Using regular neighbourhoods, we can define the various kinds of maps that will play a role in the argument.

Definition 2.12 A morphism of pre-complexes $f : X \rightarrow Y$ is a *branched map* if the induced map

$$f_* : N_X(e) \rightarrow N_X(f(e))$$

is injective for every edge e of X . If the induced map

$$f_* : N_X(v) \rightarrow N_X(f(v))$$

is injective for every vertex v of X then f is a *branched immersion*.

Note that the realisation of a branched map of complexes $f : X \rightarrow Y$ is locally injective everywhere except possibly at the centres of the faces and the vertices, and a branched immersion is locally injective everywhere except possibly at the centres of the faces.

The number of times that a morphism f winds a face around its image is an important quantity, called the *degree*.

Definition 2.13 Let $f : X \rightarrow Y$ be a morphism of 2-complexes. Since $f : S_X \rightarrow S_Y$ is a local immersion it is a covering map; in particular, for every path-component $C \subseteq S_X$, the restriction $f : C \rightarrow S_Y$ has a well-defined *degree* $\text{deg}_C(f) \in \mathbb{N}$. The sum

$$\text{deg}(f) := \sum_{C \in \pi_0(S_X)} \text{deg}_C(f)$$

is the *degree of f*. If f is a branched immersion and $\text{deg}_C(f) = 1$ for every component C then f is called an *immersion*.

Remark 2.14 In [26], immersions were defined in the more general context of orbicomplexes, which allow cone points in the centres of 2-cells. In this paper, we only consider genuine 2-complexes, and the notion of immersions from [26] specialises exactly to the definition given in Definition 2.13.

Note that a morphism f is an immersion if and only if its realisation f is locally injective. Sometimes we need to be able to modify the 1-skeleton of a 2-complex without changing the faces. The next kind of map makes this possible.

Definition 2.15 A morphism of finite pre-complexes $f : Y \rightarrow X$ is called a *face-embedding* if the map of pre-graphs $f : S_Y \rightarrow S_X$ is injective, and a *face-equivalence* if it is a bijection.

Remark 2.16 If $f : X \rightarrow Y$ is a face-embedding of complexes then the restriction of f to the interiors of the two-cells of X is a homeomorphism onto its image, whence the term. Furthermore, if $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are essential face embeddings, then the composition $g \circ f : X \rightarrow Z$ is an essential face embedding.

Stallings famously popularised the folding operation for graphs [36]. In [26, 27], we made use of a folding operation on morphisms of 2-complexes that produces immersions. Here we will make use of a very natural folding operation on morphisms of 2-complexes that produces branched immersions.

Definition 2.17 Let $f : Y \rightarrow X$ be a morphism of pre-complexes. As in [36, §3.3], the induced map of 1-skeleta $f : G_Y \rightarrow G_X$ factors as

$$G_Y \xrightarrow{f_0} G_{\bar{Y}} \xrightarrow{f_1} G_X$$

where f_0 is a π_1 -surjective finite composition of folds and f_1 is an immersion. The map of faces $f : S_Y \rightarrow S_X$ factors through the fiber product $G_{\bar{Y}} \times_{G_X} S_X$ by the

universal property; set $S_{\bar{Y}}$ equal to its image, and let $w_{\bar{Y}} : S_{\bar{Y}} \rightarrow G_{\bar{Y}}$ be the natural projection. The data $(G_{\bar{Y}}, S_{\bar{Y}}, w_{\bar{Y}})$ define a pre-complex \bar{Y} and f factors as

$$Y \xrightarrow{f_0} \bar{Y} \xrightarrow{f_1} X$$

where f_0 is a π_1 -surjection and f_1 is a branched immersion. The pre-complex \bar{Y} is called the *folded representative* of Y , and the map f_1 is called the *folded representative* of f . The folded representative is characterised by its universal property: if f factors through a branched immersion $Z \rightarrow X$ then the folded representative f_1 also factors uniquely through $Z \rightarrow X$.

This enables us to define the last kind of map that we will need.

Definition 2.18 A morphism of finite pre-complexes $f : Y \rightarrow X$ is called *face essential* if the map to the folded representative $f_0 : Y \rightarrow \bar{Y}$ is a face-equivalence. If X and Y are both complexes, this is equivalent to requiring that $\deg(f) = \deg(f_1)$. If, in addition, the restriction of f to each connected component of Y induces a π_1 -injective map on 1-skeleta, then f is said to be *essential*.

Intuitively, a face-essential map is a map such that folding doesn't identify any faces.

Remark 2.19 If $Y \rightarrow X$ is a face-essential morphism of finite complexes and \bar{Y} is the folded representative of Y then $\chi(\bar{Y}) \geq \chi(Y)$, with equality if and only if $G_Y \rightarrow G_{\bar{Y}}$ is a homotopy equivalence.

We will be interested in local ways of certifying that a morphism f is face-essential. This turns out to be easier to do in a two-step process, encapsulated by the notion of a *face immersion*.

Definition 2.20 A *face immersion* for a map $f : Y \rightarrow X$ is a factorisation

$$Y \xrightarrow{f_0} Z \xrightarrow{f_1} X$$

where f_0 is a face-equivalence and f_1 is a branched immersion.

For brevity of notation, we will often abuse notation and use the map f to denote a face immersion for f . A face immersion certifies that f is face-essential, because of the following lemma.

Lemma 2.21 *Let $f : Y \rightarrow X$ be a morphism of finite 2-complexes. There is a face immersion for f if and only if f is face-essential.*

Proof If f is face essential then the factorisation through the folded representative \bar{Y} defines a face immersion. Conversely, the existence of a face immersion implies that f factors as

$$Y \rightarrow \bar{Y} \rightarrow Z \rightarrow X$$

by the universal property of the folded representative. Since $S_Y \rightarrow S_{\bar{Y}}$ is surjective and the bijection $S_Y \rightarrow S_Z$ factors through it, it is also bijective, so $Y \rightarrow \bar{Y}$ is a face equivalence and hence according to Definition 2.18 f is face essential. \square

2.5 Gluing pre-complexes

One advantage of working with pre-complexes is that they can be conveniently glued to create complexes.

Definition 2.22 Let $f : X \rightarrow Y$ be a branched map of pre-complexes. Suppose that e_1, e_2 are edges of G_X such that $f(e_1) = f(e_2)$. Since f is a branched map, it induces injective maps

$$f_* : w_X^{-1}(e_1) \rightarrow w_Y^{-1}(f(e_1))$$

and

$$f_* : w_X^{-1}(e_2) \rightarrow w_Y^{-1}(f(e_2)).$$

The edges e_1 and e_2 of X are said to be *gluable (along f)* if

- (i) $e_1 \notin E_X^l$ and $e_2 \notin E_X^r$, and
- (ii) $f_*(w_X^{-1}(e_1)) = f_*(w_X^{-1}(e_2))$.

In this case, f induces a well-defined bijection $w_X^{-1}(e_1) \rightarrow w_X^{-1}(e_2)$.

We can now define a new pre-complex X' from X by identifying e_1 and e_2 to a common edge e . Set $\iota(e) = \iota(e_2)$ (if it exists) and $\tau(e) = \tau(e_1)$ (if it exists). Each edge $w_X^{-1}(e_1)$ corresponds to a unique edge $e_2 \in w_X^{-1}(e_2)$ under the above bijection, and in X' these two edges are identified to a common edge $\epsilon \in w_{X'}^{-1}(e)$, with $\iota(\epsilon) = \iota(e_2)$ (if it exists) and $\tau(\epsilon) = \tau(e_1)$ (if it exists). Finally, note that f descends to a well-defined branched map $f : X' \rightarrow Y$.

The resulting pre-complex X' is said to be constructed from X by *gluing e_1 and e_2 (along f)*.

Gluing can be used to reassemble a complex from the regular neighbourhoods of its vertices.

Remark 2.23 Let X be a finite complex, and let N be the disjoint union of the regular neighbourhoods of the vertices of X . Then N is naturally equipped with a branched immersion $N \rightarrow X$, and X can be reconstructed from N by gluing the edges in pairs.

3 Irreducible complexes

The properties of non-positive immersions, negative immersions and uniform negative immersions have a common theme. Given an immersion of finite 2-complexes $Y \rightarrow X$, if the Euler characteristic $\chi(Y)$ is not negative enough, then Y can be simplified. The exact sense in which Y can be simplified is of crucial importance to the definition of uniform negative immersions. In the next subsection we define irreducible




Fig. 2 The central diagram illustrates the complex X from Example 3.7: the 1-skeleton is a barbell graph, and the single face is attached along the illustrated immersed curve. It admits two different unfoldings, whose realisations are tori

complexes, and with these definitions in hand we can give the correct definitions of non-positive, negative and uniformly negative immersions.

3.1 Unfolding

We are interested in ways that a finite 2-dimensional complex X might be homotopic to a point. The next definition summarises various ways in which X either is a point or can be simplified.

Definition 3.1 (Visibly reducible and irreducible pre-complexes) A finite pre-complex X is *visibly reducible* if any one of the following four conditions hold:

- (i) some component of X is a point;
- (ii) the 1-skeleton G has a vertex of valence 1;
- (iii) X has a *free face* — that is, an edge e of the 1-skeleton G with $\#w^{-1}(e) = 1$;
- (iv) a vertex v of X separates its regular neighbourhood $N_X(v)$, so $N_X(v) \setminus \{v\}$ is disconnected.

A fifth kind of reduction also plays an important role:

- (v) a vertex v of X has an incident edge e such that the union $v \cup e$ separates the regular neighbourhood $N_X(v)$.

If none of conditions (i)-(v) hold then X is called *visibly irreducible*. If condition (v) holds but none of conditions (i)-(iv) hold, then X is called *unfold-able*.¹ See Fig. 2 for a concrete example of an unfold-able 2-complex.

If neither of conditions (i) or (ii) hold then the 1-skeleton of X is a core graph. Recall that a graph is called *core* if every vertex is in the image of an immersed cycle. Henceforth, we will always implicitly assume that the 1-skeleton of any 2-complex we consider is a core graph.

The conditions of Definition 3.1 can also be phrased naturally in terms of Whitehead graphs: see for instance [39, Lemma 2.10] for a similar list of conditions. Recall from [36] that a morphism of graphs $f : G \rightarrow H$ is a *fold* if it identifies just a single pair of edges e_1, e_2 , incident at a common vertex.

¹The hyphen is to resolve an ambiguity: we mean that such a pre-complex can be unfolded, not that it cannot be folded.

Definition 3.2 A fold of graphs is called *essential* if it is injective on fundamental groups. A morphism of graphs $f : G \rightarrow H$ is called an *essential equivalence* if it is a composition of essential folds, or equivalently, it is an isomorphism on fundamental groups. A face-essential morphism of complexes $f : Y \rightarrow X$ is called an *essential equivalence* if the map of 1-skeleta $f : G_Y \rightarrow G_X$ is an essential equivalence of graphs. In this case, we also say that Y is an *essential unfolding* of X .

Note that, if $f : Y \rightarrow X$ is an essential equivalence of complexes, then the realisation f is a homotopy equivalence.

The next lemma, which is a restatement of [39, Lemma 2.8] in the terminology of this paper, explains the relationship between Definition 3.1(v) and (un)folding.

Lemma 3.3 *If a finite complex X is unfold-able then there is a finite complex X' and an essential fold $f : X' \rightarrow X$.*

Unfolding preserves the number of edges of the faces while increasing the number of vertices and edges of the 1-skeleton, so it is not hard to see that Lemma 3.3 can only be applied finitely many times before producing a complex which is either visibly reducible or visibly irreducible. The next lemma is a restatement of [39, Lemma 2.11] in the terminology of this paper.

Lemma 3.4 *For any finite complex X , there is a finite complex X' and an essential equivalence $f : X' \rightarrow X$ such that X' is either visibly reducible or visibly irreducible.*

We say that X *unfolds to X'* . To summarise, unfold-able complexes may be either reducible or irreducible, but their status can be determined by unfolding.

Definition 3.5 (Reducibility and irreducibility) If a finite 2-complex X unfolds to a visibly reducible complex X' (with the 1-skeleton still a core graph) then X is said to be *reducible*. Likewise, if X unfolds to a visibly irreducible complex X' then X is said to be *irreducible*.

Note that an irreducible 2-complex is not required to be connected.

Remark 3.6 The terminology ‘irreducible’ has been used in several papers related to this one, such as [25–27, 39]. Although the usage was always similar to its definition given here, it was not always identical, so the reader is advised to check the definition in the cited paper when a result is quoted.

It is not immediately obvious that a complex X cannot be both reducible and irreducible, since X may admit many different unfoldings. Indeed, a 2-complex may have several unfoldings, as the next example illustrates.

Example 3.7 Let X be a 2-complex defined as follows. The 1-skeleton of X is a ‘spectacles’ or ‘bar-bell’ graph, with two vertices u_1 and u_2 , and edges a_1, a_2, b , where each a_i is a loop starting and ending at u_i , and b joins u_1 to u_2 . The single face C

of X is attached along the loop labelled by the commutator of a_1 and ba_2b^{-1} . Now X admits two different unfoldings, each with realisation homeomorphic to the torus: one is obtained by dividing u_1 into two vertices, the other is obtained by dividing u_2 into two vertices.

Many further examples of this type can be constructed by taking any cellular decomposition X' of a surface and performing an essential fold to obtain a complex X ; such an X can always be unfolded in two ways.

Nevertheless, a famous lemma of Whitehead [38] implies that (ir)reducibility is equivalent to various group-theoretic properties of the conjugacy classes in the free group $F = \pi_1(G)$ represented by the components of S . Combined with the material of this section, Whitehead's lemma (see, for instance, [39, Lemma 2.10] for a suitable statement), gives the following proposition.

Proposition 3.8 *Let X be a finite connected complex. Let F be the fundamental group of the 1-skeleton G_X , and let $\{w_1, \dots, w_n\}$ be a finite set of representatives for the conjugacy classes of F determined by the components of S . The complex X is reducible if and only if one of the following three conditions hold:*

- (i) F is trivial and $n = 0$;
- (ii) F is infinite cyclic, $n \leq 1$ and, if $n = 1$, $F = \langle w_1 \rangle$;
- (iii) there is a non-trivial free splitting $F = A * B$ such that, for each i , w_i is conjugate into either A or B .

Since the conditions of Proposition 3.8 are group-theoretic and are not altered by homotopy-equivalences of the 1-skeleta (once we have ensured that they are core graphs), it follows that every finite complex is either reducible or irreducible.

3.2 Irreducible cores

Since this notion of irreducibility is slightly stronger than those used in [25, 27], we will need some lemmas that enable us to improve unfold-able complexes to irreducible ones, under suitable hypotheses on the fundamental group.

Definition 3.9 A group Γ is *freely decomposable* if Γ splits as a non-trivial free product $\Gamma = A * B$. Otherwise Γ is said to be *freely indecomposable*.

A 2-complex X can be replaced by a visibly irreducible one, as long as the fundamental group is complicated enough.

Lemma 3.10 *Let X be a finite, connected 2-complex, and suppose that $\pi_1(X)$ is neither freely decomposable nor free. Then there is a visibly irreducible 2-complex X' and an essential face-embedding $X' \rightarrow X$, which is also an isomorphism on fundamental groups.*

Proof The proof proceeds by induction on the number of faces of X . By Lemma 3.4, there is an essential equivalence $X' \rightarrow X$ such that X' is either visibly reducible or

visibly irreducible. If X' is visibly irreducible the result is immediate; otherwise, X' is visibly reducible. Since $\pi_1(X) \cong \pi_1(X')$ is non-trivial, X' is not a point, and we may assume that the 1-skeleton of X' is a core graph.

If X' has a free face, collapsing it exhibits a sub-complex $X'' \subseteq X'$ which is a deformation retract of X' with fewer faces. Note also that the inclusion of X'' into X' is an essential face-embedding. Therefore, the result holds for X'' , and hence for X' , by the inductive hypothesis.

Finally, suppose that X' has a locally separating vertex v . Since $\pi_1(X')$ is freely indecomposable and not infinite cyclic, v must be separating, and cutting X' along v realises X' as a wedge

$$X' = X_1 \vee X_2 .$$

Note that the inclusion maps $X_i \hookrightarrow X'$ are essential. Since every free splitting of $\pi_1(X')$ is trivial, X_2 is simply connected, without loss of generality. Since the 1-skeleton of X' is a core graph, X_2 is not a tree, and must therefore include at least one face. Hence X_1 has fewer faces than X , and so the lemma holds for X_1 by the inductive hypothesis. Since $X_1 \rightarrow X'$ is an isomorphism on fundamental groups, the result also holds for X , which completes the proof. \square

As in Example 3.7, the complex X' furnished by Lemma 3.10 may not be unique. However, we can make it unique by folding.

Definition 3.11 Let X be finite, connected 2-complex with $\pi_1(X)$ neither freely decomposable nor free. Let $f : X' \rightarrow X$ be provided by Lemma 3.10, and let $\widehat{X} \rightarrow X$ be the immersion provided by folding f . Then \widehat{X} is called an *irreducible core* for X .

Remark 3.12 Since $X' \rightarrow X$ is essential, X' is an essential unfolding of \widehat{X} . In particular, since X' is visibly irreducible, it follows that \widehat{X} is irreducible. Also, since $X' \rightarrow X$ is a π_1 -isomorphism and factors through the π_1 -surjection $X' \rightarrow \widehat{X}$, the immersion $\widehat{X} \rightarrow X$ is a π_1 -isomorphism.

We can also characterise irreducible cores group-theoretically. Let $X = (G_X, S_X, w_X)$ be a compact, connected 2-complex. The fundamental group of the 1-skeleton G_X is a finitely generated free group F . The images of the circles S_X under the attaching map w_X define a collection of conjugacy classes of cyclic subgroups $\{\langle w_1 \rangle, \dots, \langle w_n \rangle\}$ of F . Grushko’s theorem implies the existence of a maximal splitting

$$F = F_1 * \dots * F_k ,$$

such that each w_i is conjugate into some F_j . (If we combine the F_j that do not contain any of the w_i into a common free factor H , then this gives the *relative Grushko decomposition* of the pair $(F, \{\langle w_i \rangle\})$.) If Γ_j is the quotient of F_j by the conjugates of the w_i contained in F_j , then this gives a natural free-product decomposition

$$\Gamma = \Gamma_1 * \dots * \Gamma_k$$

of $\Gamma = \pi_1(X)$. If Γ is neither freely decomposable nor free, then exactly one of these factors – without loss of generality, Γ_1 – must be non-trivial. Now, \widehat{X} can be defined as follows: the 1-skeleton is the Stallings core of the subgroup F_1 of $F = \pi_1(G_X)$, and the faces correspond to those w_i that are conjugate into F_1 .

We shall see that irreducible cores enjoy a universal property. Before we prove that, we need to notice that unfolds can be pulled back along branched immersions.

Lemma 3.13 *Let $Y \rightarrow X$ be a branched immersion, and suppose that $X' \rightarrow X$ is an essential fold. Then there is an essential unfolding $Y' \rightarrow Y$ such that the composition $Y' \rightarrow X$ factors through $X' \rightarrow X$.*

Proof Such an unfolding of X means that there is an edge e with endpoints u and v such that e and v together separate the regular neighbourhood $N_X(v)$, so it can be written as a union

$$N_X(v) = N_1 \cup_{e \cup v} N_2.$$

Let v_1, \dots, v_n be the pre-images of v in Y and let e_i be the pre-images of e incident at v_i , if it exists. For each i , $N_Y(v_i)$ embeds in $N_X(v)$, and so either $N_Y(v_i)$ maps into one of N_1 or N_2 , or $N_Y(v_i)$ is separated by $e_i \cup v_i$. The unfolding Y' is now constructed by unfolding each of these separating e_i in turn. □

Using this, we can now prove the claimed universal property for irreducible cores.

Lemma 3.14 *Let $Y \rightarrow X$ be a branched immersion of finite, connected 2-complexes that is non-trivial on fundamental groups. Suppose that Y is irreducible and that $\pi_1(X)$ is neither freely decomposable nor free, and let $\widehat{X} \rightarrow X$ be an irreducible core of X . Then the morphism $Y \rightarrow X$ factors through the map $\widehat{X} \rightarrow X$.*

Proof Recall that the irreducible core \widehat{X} is constructed by folding an inductively constructed morphism $X' \rightarrow X$. The lemma is proved by arguing that there is an essential unfolding $Y' \rightarrow Y$ that lifts to X' .

Consider the inductive construction of X' from Lemma 3.10, which is a finite sequence of unfoldings and inclusions of sub-complexes. Lemma 3.13 handles the case of unfoldings. It remains to deal with the two cases of the argument that pass to subcomplexes of an unfolding X' of X .

In the first case, X' has a face C that is a free face, and $X'' \subseteq X'$ is the subcomplex obtained by collapsing that free face. Since $Y' \rightarrow X'$ is a branched immersion, if any face D of Y' maps to C then D is also a free face, so Y' is visibly reducible, contradicting the hypothesis that Y is irreducible. Therefore C is not in the image of Y' , and so the map $Y' \rightarrow X'$ restricts to X'' .

In the second case, X' has a locally separating vertex and splits as a wedge $X_1 \vee X_2$ with X_2 simply connected. Since $Y' \rightarrow X'$ is a branched immersion and no vertex of Y' is locally separating, it follows that the image of Y is contained entirely inside either X_1 or X_2 , and since $Y' \rightarrow X'$ is non-trivial on fundamental groups, it must be X_1 , as required. This completes the proof that Y unfolds to some Y' such that $Y' \rightarrow X$ lifts to X' .

To complete the proof, we fold Y' to construct a 2-complex \widehat{Y} . Since $\widehat{X} \rightarrow X$ and $Y \rightarrow X$ are branched immersions, there is a well-defined fibre-product 2-complex $Y \times_X \widehat{X}$, with 1-skeleton equal to the fibre product of graphs $G_Y \times_{G_X} G_{\widehat{X}}$ and faces given by pulling back the faces of Y and \widehat{X} . The universal property of the fibre product now provides a canonical map $Y' \rightarrow Y \times_X \widehat{X}$, and \widehat{Y} is the result of folding this map. By construction, the natural map $\widehat{Y} \rightarrow X$ factors through \widehat{X} .

Finally, since the identity map on Y is the folded representative of the map $Y' \rightarrow Y$, the universal property of folded representatives provides a canonical morphism $Y \rightarrow \widehat{Y}$, and since $\widehat{Y} \rightarrow X$ factors through \widehat{X} , the result follows. \square

Remark 3.15 The universal property of Lemma 3.14 implies in the usual way that irreducible cores are unique up to canonical isomorphism. We may therefore speak of *the* irreducible core of a finite, connected 2-complex X , as long as $\pi_1(X)$ is neither freely decomposable nor free.

Remark 3.16 The irreducible core of a connected 2-complex X can also be characterised group-theoretically. Let F be the fundamental group of the 1-skeleton, and let $\langle w_1 \rangle, \dots, \langle w_n \rangle$ be the cyclic subgroups of F defined (up to conjugacy) by the faces of X . The pair $(F, \{\langle w_1 \rangle, \dots, \langle w_n \rangle\})$ has a relative Grushko decomposition $F = F_1 * \dots * F_n$. Each w_i is conjugate into a unique $F_{j(i)}$ and so, after conjugating the w_i , the pair $(F, \{\langle w_i \rangle\})$ decomposes as a free product of pairs $(F_k, \{\langle w_i \rangle \mid j(i) = k\})$. If $\pi_1(X)$ is neither freely decomposable nor free, then there is a unique index k such that $F_k / \langle\langle w_i \mid j(i) = k \rangle\rangle$ is neither freely decomposable nor free. The 1-skeleton of the irreducible core \widehat{X} is given by the core graph associated to the subgroup F_k of F , and the 2-cells are given by the set $\{w_i \mid j(i) = k\}$.

The results of this section guarantee that X has an irreducible core if it is neither freely decomposable nor free. However, the converse is not true: X may be irreducible, and yet $\pi_1(X)$ may be trivial, free, freely indecomposable etc. For instance, the complex associated to the presentation

$$\langle a, b \mid baba^{-2}, abab^{-2} \rangle$$

of the trivial group is irreducible.

3.3 Curvature conditions

With the definitions of irreducibility in hand, we can now define non-positive and negative immersions. These definitions are inspired by Wise [41, 44], and similar definitions have played a role in several recent papers about one-relator groups, such as [18, 25, 27]. They are also naturally related to the work of Martínez-Pedroza and Wise on sectional curvature for 2-complexes [29, 42, 43].

It turns out that Wise’s original definitions are too strong to prove Theorem A. Here we will give slightly less restrictive definitions, using the notion of irreducible complexes from the last section. See Sect. 3.4 below for a comparison between our definitions and Wise’s. As usual, $\chi(X)$ denotes the Euler characteristic of a complex X .

Definition 3.17 A finite 2-complex X has *non-positive immersions* if, for every immersion of finite complexes $Y \looparrowright X$, either

- (i) $\chi(Y) \leq 0$, or
- (ii) Y is reducible.

The non-positive immersions property for presentation complexes of one-relator groups was proved in [18] and [25].

The next definition is motivated by the observation that, if $f : Y \rightarrow X$ is an immersion, then $\deg(f)$ is equal to the number of faces of Y , and can be thought of as a useful adaptation of Euler characteristic to the setting of branched maps.

Definition 3.18 Let $f : Y \rightarrow X$ be a branched map of 2-complexes, and let G be the 1-skeleton of Y . The quantity

$$\tau(f) := \deg(f) + \chi(G)$$

is called the *total curvature* of f . We will usually abuse notation and write $\tau(Y)$ for $\tau(f)$, since $\tau(f) = \chi(Y)$ when f is an immersion.

Indeed, the techniques of both papers [18, 25] prove something slightly stronger than non-positive immersions: the map can be taken to be a branched immersion, and the complex Y is only required to not be visibly reducible. For instance, [25, Theorem 1.2] can be stated as follows, after accounting for small differences in terminology.

Theorem 3.19 *Let X be the presentation complex of a torsion-free one-relator group. If Y is a finite complex and $Y \rightarrow X$ is a branched immersion then either*

- (i) $\tau(Y) \leq 0$, or
- (ii) Y has a free face, and in particular is visibly reducible.

The definition of negative immersions is a natural adaptation of Definition 3.17. Again, this definition assumes a slightly stronger notion of irreducibility than was used in [27].

Definition 3.20 A finite 2-complex X has *negative immersions* if, for every immersion of finite complexes $Y \rightarrow X$, either

- (i) $\chi(Y) < 0$, or
- (ii) Y is reducible.

The property of negative immersions for one-relator groups was characterised in [27], in which Theorem 1.2 was proved.

Remark 3.21 In [27, Definition 1.1], a 2-complex X is defined to have negative immersions if, for every immersion of finite complexes $Y \rightarrow X$, either $\chi(Y) < 0$ or Y Nielsen reduces to a graph. To say that Y Nielsen reduces to a graph means that there is a homotopy equivalence of 1-skeleta $Y_{(1)} \simeq Z_{(1)}$ so that the 2-complex Z obtained

by attaching the 2-cells of Y to $Z_{(1)}$ in the natural way is the result of wedging discs to a graph.

Let us briefly explain why Definition 3.20 agrees with [27, Definition 1.1]. To this end, let $Y \rightarrow X$ be an immersion of finite 2-complexes, and assume that $\chi(Y) \geq 0$. If X satisfies [27, Definition 1.1] then Y Nielsen reduces to a graph, and then Y is reducible by Proposition 3.8.

To prove the converse, assume that X satisfies Definition 3.20, and proceed by induction on the number of 2-cells of Y . Since $\chi(Y) \geq 0$, Y is reducible by assumption, and Proposition 3.8 applies. Items (i) and (ii) of the proposition correspond to the base cases of the induction where Y is a point, a circle or a disc. If item (iii) of Proposition 3.8 applies, then Y unfolds to a 2-complex Y' with a locally separating vertex v . Cutting along v yields a (possibly disconnected) new 2-complex Y'_1 , which folds to a 2-complex Y_1 immersing into Y with $\chi(Y_1) = \chi(Y) + 1$. Repeating this process n times leads to a sequence of immersed 2-complexes

$$Y_n \rightarrow Y_{n-1} \rightarrow \cdots \rightarrow Y_1 \rightarrow Y$$

with $\chi(Y_n) = \chi(Y) + n \geq n$, each with the same number of 2-cells as Y . Note that, for each n , Y is Nielsen equivalent to the result of wedging Y_n to a graph.

For large enough n , Y_n must have a free face. Indeed, let m be the number of 2-cells of Y and let l be their total edge length. Then, unless Y_n has a free face, it has at most $l/2$ edges and hence, since every vertex has valence at least 2, it has at most $l/2$ vertices, so $\chi(Y_n) \leq l/2 + m$. Therefore, Y_n must have a free face for $n > l/2 + m$, as claimed. Collapsing a free face in such a Y_n produces a new 2-complex Y' with $\chi(Y') \geq 0$ and with fewer 2-cells than Y . Now Y' Nielsen reduces to a graph by induction, and hence so does Y , as required.

Theorem 1.2 gives a useful statement, but the following more precise statement is an immediate consequence of [27, Lemma 6.11], after noticing that the proof of that result works equally well if immersions are replaced by branched immersions and χ is replaced by τ .

Theorem 3.22 *Let X be the presentation complex of a torsion-free one-relator group $G = F/\langle\langle w \rangle\rangle$ with negative immersions. If Y is a finite complex and $f : Y \rightarrow X$ is a branched immersion then either*

- (i) $\tau(Y) < 0$, or
- (ii) Y is reducible.

Below, we will need the following improvement of this statement to the context of face-essential maps,

Corollary 3.23 *Let X be the presentation complex of a torsion-free one-relator group $G = F/\langle\langle w \rangle\rangle$ with negative immersions. If Y is a finite complex and $f : Y \rightarrow X$ is a face-essential map then either*

- (i) $\tau(Y) < 0$, or
- (ii) Y is reducible.

Proof Consider the folded representative

$$Y \rightarrow \bar{Y} \rightarrow X$$

so $\bar{Y} \rightarrow X$ is a branched immersion. Since $Y \rightarrow X$ is face essential,

$$\tau(Y) \leq \tau(\bar{Y}),$$

because $\deg f$ is equal to the degree of its folded representative, with equality if and only if $Y \rightarrow \bar{Y}$ is an essential equivalence. Suppose now that $\tau(Y) = 0$, so $\tau(\bar{Y}) \geq 0$. There are two cases to consider: either $\tau(\bar{Y}) = 0$ or $\tau(\bar{Y}) > 0$.

In the first case, $\tau(\bar{Y}) = 0$ so $Y \rightarrow \bar{Y}$ is an essential equivalence. Theorem 3.22 then implies that \bar{Y} is reducible. Since reducibility is equivalent to the group-theoretic criteria of Proposition 3.8, it follows that Y is also reducible, because $Y \rightarrow \bar{Y}$ is an essential equivalence.

In the second case, $\tau(\bar{Y}) > 0$ so, by Theorem 3.19, \bar{Y} has a free face, whence Y also has a free face, and so is visibly reducible. \square

Finally, we are ready to give a definition of *uniform negative immersions*.

Definition 3.24 A finite 2-complex X has *uniform negative immersions* if there is $\epsilon > 0$ so that, for every immersion of finite complexes $Y \looparrowright X$, either

(i)

$$\frac{\chi(Y)}{\#\{2\text{-cells of } Y\}} \leq -\epsilon,$$

or

(ii) Y is reducible.

Theorem C asserts that presentation complexes of one-relator groups with negative immersions also have uniform negative immersions.

3.4 Wise's notion of negative immersions

We finish this section by contrasting Definition 3.24 with the following definition of Wise.

Definition 3.25 A finite 2-complex X has *negative immersions in the sense of Wise* if there is $\epsilon > 0$ so that, for every immersion of finite complexes $Y \looparrowright X$, either

(i)

$$\frac{\chi(Y)}{\#\{2\text{-cells of } Y\}} \leq -\epsilon,$$

or

(ii) Y has a free face.

In [44], Wise proves that every one-relator group with torsion G is coherent by showing that a presentation complex for (some finite-index subgroup of) G satisfies Definition 3.25, and then proving that this in turn implies coherence. The independent proof given in [26] can also be viewed this way.

Clearly, negative immersions in the sense of Wise implies our notions of negative immersions and uniform negative immersions. The following example shows that this implication is not reversible [44].

Example 3.26 Let X be the 2-complex associated to the presentation $G = F/\langle\langle w \rangle\rangle$, where F is the free group $\langle a, b, c \rangle$ and

$$w := ab^2c^2ab^2c^2b^2c^2.$$

By setting $a' = ab^2c^2$, this presentation can be seen to be Nielsen equivalent² to

$$\langle a', b, c \mid (a')^2b^2c^2 \rangle,$$

the standard presentation of the fundamental group of the surface of Euler characteristic -1 . In particular, G is 2-free and so X has negative immersions by Theorem 1.2, and indeed uniform negative immersions by Theorem C.

We now define a compact 2-complex Y and an immersion to X . The 1-skeleton of Y is a rose with two petals labelled α and β , and immerses into X via the assignment

$$f : \alpha \mapsto a, \beta \mapsto b^2c^2,$$

while a single face is attached along the word $u(\alpha, \beta) = \alpha\beta\alpha\beta^2$. Since $f(u(\alpha, \beta)) = w(a, b, c)$, the immersion of 1-skeleta extends to an immersion $Y \looparrowright X$. Clearly $\chi(Y) = 0$, but since both α and β appear at least twice in u , Y has no free face. Therefore, Y does not have negative immersions in the sense of Wise.

The immersion $Y \looparrowright X$ does not contradict our definitions of (uniform) negative immersions (Definitions 3.20 and 3.24), since u is a primitive element of the free group $\langle \alpha, \beta \rangle$, and so the complex Y is reducible by Proposition 3.8.

Example 3.26 shows that one cannot prove Theorem A by showing that all such complexes satisfy Definition 3.25. It also illustrates another important difference between the definitions: since the standard presentation complex for the surface of Euler characteristic -1 does have negative immersions in the sense of Definition 3.25, that notion is not invariant under Nielsen equivalence, whereas Definitions 3.20 and 3.24 are, by Remark 3.16.

3.5 Stable commutator length

Commutator length and its stabilisation provide natural quantifications of the complexity of an element of a commutator subgroup. See Calegari's monograph for a comprehensive treatment of stable commutator length [6].

²Recall that two presentations are *Nielsen equivalent* if one is taken to the other by applying a free-group automorphism to the generators.

Definition 3.27 The *commutator length* of an element g of the commutator subgroup of a group G is

$$\text{cl}(g) = \min\{n \in \mathbb{N} \mid g = \prod_{i=1}^n [x_i, y_i], x_i, y_i \in G\},$$

the minimal n such that g is a product of n commutators. The *stable commutator length* is defined to be

$$\text{scl}(g) = \inf_{n \in \mathbb{N}} \frac{\text{cl}(g^n)}{n}.$$

The purpose of this section is to prove Corollary D, and also to explain why Conjecture 1.5 implies Conjecture 1.7. Both follow from the next proposition, which is due to Calegari [5, Lemma 2.7]. Nevertheless, in order to clarify how the argument works in the setting of this paper, we give the proof here.

A 2-complex is called a *surface* if its realisation is homeomorphic to a closed, 2-dimensional surface.

Proposition 3.28 (Calegari) *Let X be the presentation complex of a one-relator group $F/\langle\langle w \rangle\rangle$ and suppose that $w \in [F, F]$. There is an essential branched map from an orientable surface $f : \widehat{\Sigma} \rightarrow X$ such that*

$$\frac{\tau(\widehat{\Sigma})}{\text{deg}(f)} = 1 - 2 \text{scl}(w).$$

Proof As above, the data of X consists of a map $w : S \looparrowright G$, where G is the 1-skeleton and w is the attaching map of the unique face. An *admissible surface* is a commutative diagram of continuous maps of topological spaces

$$\begin{array}{ccc} \partial \Sigma & \longrightarrow & \Sigma \\ \downarrow & & \downarrow \\ \mathcal{S} & \xrightarrow{w} & G \end{array}$$

such that Σ is a compact surface without spherical components, $\partial \Sigma \rightarrow \Sigma$ is the natural inclusion of the boundary and $\partial \Sigma \rightarrow \mathcal{S}$ is an orientation-preserving covering map. Calegari showed that

$$\text{scl}(w) \leq -\frac{\chi(\Sigma)}{2 \text{deg}(\partial \Sigma)}, \tag{1}$$

where $\text{deg}(\partial \Sigma)$ denotes the degree of the covering map $\partial \Sigma \rightarrow \mathcal{S}$, for any admissible surface [6, Proposition 2.10]. If equality is realised, so

$$\text{scl}(w) = -\frac{\chi(\Sigma)}{2 \text{deg}(\partial \Sigma)}, \tag{2}$$

then Σ is said to be *extremal*. Calegari's rationality theorem [7] (see also the account in [6, Theorem 4.24]) asserts that an extremal surface Σ exists whenever $w \in [F, F]$.

A lemma of Culler [11] (which is also implicit in the proof of Calegari's theorem) implies that Σ has a spine Σ_0 such that the map $\Sigma \rightarrow G$ factors as the composition of a deformation retraction $\Sigma \rightarrow \Sigma_0$ with a morphism of graphs $\Sigma_0 \rightarrow G$. Let $\widehat{\Sigma}$ be the closed surface obtained by gluing discs to the boundary components of Σ . Then $\widehat{\Sigma}$ naturally has the structure of a 2-complex $\widehat{\Sigma}$ with 1-skeleton Σ_0 , and the map $\Sigma \rightarrow G$ extends to a morphism of 2-complexes $f : \widehat{\Sigma} \rightarrow X$.

The extremality equation (2) can be rearranged to give the required relationship between total curvature and stable commutator length. Indeed, because the 1-skeleton of $\widehat{\Sigma}$ is a deformation retract of Σ ,

$$\tau(f) = \deg(f) + \chi(\Sigma) = \deg(f)(1 - 2 \operatorname{scl}(w))$$

which gives the required equation.

It now remains to show that f is essential. This follows from a lemma of Calegari [5, Lemma 2.7], which in turn relies on the subgroup separability of free groups, proved by Marshall Hall Jr [23]. For completeness, we outline the proof.

Suppose that f is not π_1 -injective on some connected component of the 1-skeleton. Then there is a homotopically essential closed curve γ on Σ such that $f \circ \gamma$ is homotopically trivial in G . Since $\pi_1(\Sigma)$ is subgroup separable there is, for some finite d , a d -sheeted covering space $\Sigma_1 \rightarrow \Sigma$ such that, after a homotopy, γ lifts to a simple closed curve γ_1 on Σ_1 . Let Σ_2 be the result of surgery on Σ_1 along γ_1 ; that is, Σ_2 is constructed by cutting Σ_1 along γ_1 and gluing in two discs along the resulting boundary components. Since γ_1 maps to a null-homotopic curve in G , f extends to a map $\Sigma_2 \rightarrow G$ that makes Σ_2 into an admissible surface. We now calculate:

$$\begin{aligned} -\frac{\chi(\Sigma_2)}{2 \operatorname{deg}(\partial \Sigma_2)} &= -\frac{\chi(\Sigma_1) + 2}{2 \operatorname{deg}(\partial \Sigma_1)} \\ &< -\frac{\chi(\Sigma_1)}{2 \operatorname{deg}(\partial \Sigma_1)} \\ &= -\frac{d\chi(\Sigma)}{2d \operatorname{deg}(\partial \Sigma)} \\ &= \operatorname{scl}(w) \end{aligned}$$

contradicting (1). Hence, f is π_1 -injective on each connected component.

Now suppose that f is not face-essential. Then the map from $\widehat{\Sigma}$ to its folded representative is not injective on faces, so there are distinct points $x, y \in \partial \Sigma$ with the same image in $\overline{\Sigma}_0$, the folded representative of the spine. Fix any path α in Σ from x to y . The image of α in $\overline{\Sigma}_0$ is a loop and so, since the map $\Sigma \rightarrow \overline{\Sigma}_0$ is surjective on fundamental groups, we may concatenate α with a loop in Σ so that its image in $\overline{\Sigma}_0$, and hence in G , is null-homotopic. That is to say, we may choose α so that $f \circ \alpha$ is a null-homotopic loop. Next, as above, modify α by a homotopy and pass to a d -sheeted cover $\Sigma_1 \rightarrow \Sigma$ so that α lifts to an embedded arc α_1 .

Let Σ_2 be constructed by attaching an interval β_1 to $\partial \Sigma_1$ at the endpoints of α_1 and thickening it to a 1-handle. The map f extends over this 1-handle and, after a homotopy to ensure that $\partial \Sigma_2 \rightarrow S$ is a covering map, this makes Σ_2 into an admissible

surface. Note that $\chi(\Sigma_2) = \chi(\Sigma_1) - 1$ and $\deg(\partial \Sigma_2) = \deg(\partial \Sigma_1)$. The concatenation of α_1 and β_1 defines an embedded loop γ_2 in Σ_2 such that $f \circ \gamma_2$ is null-homotopic so, as in the proof of π_1 -injectivity, we may perform surgery on γ_2 to obtain a new admissible surface Σ_3 . As before, we calculate:

$$\begin{aligned} -\frac{\chi(\Sigma_3)}{2 \deg(\partial \Sigma_3)} &= -\frac{\chi(\Sigma_2) + 2}{2 \deg(\partial \Sigma_2)} \\ &= -\frac{\chi(\Sigma_1) + 1}{2 \deg(\partial \Sigma_1)} \\ &< -\frac{\chi(\Sigma_1)}{2 \deg(\partial \Sigma_1)} \\ &= -\frac{d\chi(\Sigma)}{2d \deg(\partial \Sigma)} \\ &= \text{scl}(w) \end{aligned}$$

which, again, contradicts (1). Hence, f is face-essential as claimed. □

Corollary D now follows quickly from Corollary 3.23 and Proposition 3.28.

Proof of Corollary D By Theorem 1.2, the hypothesis that $\pi(w) > 2$ is equivalent to the statement that X has negative immersions. Consider the face-essential map $\widehat{\Sigma} \rightarrow X$ provided by Proposition 3.28. Since $\widehat{\Sigma}$ is a closed surface each condition of Definition 3.1 can be easily checked, and we see that $\widehat{\Sigma}$ is (visibly) irreducible. Combining Corollary 3.23 and Proposition 3.28 then gives

$$1 - 2 \text{scl}(w) = \frac{\tau(\widehat{\Sigma})}{\deg(f)} < 0$$

as required. □

The proposition also immediately implies that Heuer’s conjecture follows from our Conjecture 1.5.

Corollary 3.29 *Conjecture 1.5 implies Conjecture 1.7.*

Proof Let $w \in [F, F]$ be non-trivial and let X be the presentation complex of the associated one-relator group $F/\langle\langle w \rangle\rangle$. Applying Conjecture 1.5 to the essential map $f : \widehat{\Sigma} \rightarrow X$ provided by Proposition 3.28 gives

$$1 - 2 \text{scl}(w) = \frac{\tau(\widehat{\Sigma})}{\deg(f)} \leq 2 - \pi(w)$$

which rearranges to give $2 \text{scl}(w) \geq \pi(w) - 1$, as required. □




Fig. 3 An example of a vertex piece

4 A linear system

The principal extra ingredient in the proof of Theorem C is Theorem 4.12, a rationality theorem in the spirit of the main results of [7] and [39]. Fix a finite 2-complex X . The idea is to encode face immersions from visibly irreducible complexes to X as the integral points of a system of linear equations and inequations.

4.1 Vertex and edge pieces

Consider a face-essential morphism $f : Y \rightarrow X$ with Y visibly irreducible, and a face immersion

$$Y \xrightarrow{f_0} Z \xrightarrow{f_1} X$$

for f . The variables of the linear system count the combinatorial types of the vertices of Z and their preimages in Y . These are encoded as maps between (disjoint unions of) asterisk pre-complexes.

Definition 4.1 A vertex piece P over a 2-complex X is a face immersion

$$Y(P) \rightarrow Z(P) \rightarrow X$$

where $Y(P)$ is visibly irreducible and $Z(P)$ is an asterisk pre-complex. (In particular, $Y(P)$ is a disjoint union of asterisk pre-complexes.) Vertex pieces are considered up to the natural notion of isomorphism: if there are isomorphisms of pre-complexes $Y(P) \rightarrow Y(Q)$ and $Z(P) \rightarrow Z(Q)$ commuting with the maps $Y(\bullet) \rightarrow Z(\bullet) \rightarrow X$ then P and Q are isomorphic.

See Figure 3 for an example of a vertex piece.

For a 2-complex X , the set $\mathcal{P} = \mathcal{P}(X)$ of isomorphism types of vertex pieces over vertices of X indexes the variables of our linear system. It is of crucial importance that there are only finitely many variables.

Lemma 4.2 *If X is a finite complex then \mathcal{P} is finite.*

Proof Every regular neighbourhood $N(v)$ is finite. Since $Z(P) \rightarrow N_X(v)$ is a branched immersion from an asterisk pre-complex it is injective, so there are only finitely many possible combinatorial types for $Z(P)$ and the map to $N_X(v)$. Since $Y(P) \rightarrow Z(P)$ is a face-equivalence, the cardinality of $S_{Y(P)}$ is also bounded. Finally, since $Y(P)$ is visibly irreducible, each edge and vertex of $Y(P)$ has at least two pre-images in $S_{Y(P)}$, so the cardinality of $G_{Z(P)}$ is also bounded. So there are at most boundedly many different possible isomorphism types for $Y(P)$, and hence also only boundedly many possible maps $Y(P) \rightarrow Z(P)$. \square

The vertex pieces will define the variables of our linear system. Edge pieces will give the defining equations. An *edge pre-complex* is a pre-complex whose underlying pre-graph consists of a single edge and no vertices.

Definition 4.3 An *edge piece* R over an edge e of a 2-complex X consists of a face immersion

$$Y(R) \rightarrow Z(R) \rightarrow X$$

where $Z(R)$ is an edge pre-complex. Again, edge pieces are considered up to the natural notion of isomorphism.

Let $\mathcal{E} = \mathcal{E}(X)$ denote the set of isomorphism types of edge pieces over edges of X .

4.2 Weight vectors

Fix a finite 2-complex X . Again, consider a face immersion

$$Y \xrightarrow{f_0} Z \xrightarrow{f_1} X$$

for a face-essential morphism $f : Y \rightarrow X$ with Y visibly irreducible. Although this face immersion is our actual object of study, we will often abuse notation and just denote it by f , or even Y .

Definition 4.4 Each vertex v of Z defines an *induced vertex-piece* $P \equiv P_Y(v)$ over $f_1(v)$ as follows:

- (i) $Z(P) := N_Z(v)$;
- (ii) $Y(P) := \coprod_{f_0(u)=v} N_Y(u)$.

The morphisms $Y(P) \rightarrow Z(P) \rightarrow X$ are the natural induced morphisms. Each edge e of Z defines an *induced edge-piece* $R_Y(e)$ in a similar manner.

We translate face immersions into linear algebra by counting the number of times that each vertex-piece appears. This count naturally lives in the following vector space.

Definition 4.5 The *space of weight vectors* $\mathbb{R}[\mathcal{P}]$ is the real vector space of formal linear sums

$$v = \sum_{P \in \mathcal{P}} v_P P$$

of vertex-pieces $P \in \mathcal{P}$; the elements of this vector space are called *weight vectors*. We will also talk about the additive subgroups $\mathbb{Q}[\mathcal{P}]$ and $\mathbb{Z}[\mathcal{P}]$ consisting of rational and integral points respectively.

There is a map that associates to the face immersion

$$Y \xrightarrow{f_0} Z \xrightarrow{f_1} X$$

from a visibly irreducible complex Y a weight vector:

$$v(Y) := \sum_{y \in V_Z} P_Y(y),$$

recalling that V_Z is the set of vertices of the 1-skeleton of Z . Clearly, $v(Y)$ has non-negative integral coordinates, and, in fact, the possible weight vectors are exactly the points with non-negative integral coordinates that satisfy a certain system of linear equations, called the *gluing equations*. These gluing equations make use of *boundary maps*, which associate to each vertex piece the induced edge-pieces of the incident vertices. Let $\mathbb{R}[\mathcal{E}]$ be the real vector space of formal linear sums of edge pieces.

Definition 4.6 Recall from Definition 4.4 that, for any vertex piece

$$Y(P) \rightarrow Z(P) \rightarrow X$$

and any edge e of $Z(P)$, there is an induced edge piece $R_{Y(P)}(e)$.

The boundary map $\partial^t : \mathbb{R}[\mathcal{P}] \rightarrow \mathbb{R}[\mathcal{E}]$ is defined by extending

$$\partial^t(P) := \sum_{e \in E_{Z(P)}^t} R_{Y(P)}(e)$$

linearly. Similarly, the assignment

$$\partial^\tau(P) := \sum_{e \in E_{Z(P)}^\tau} R_{Y(P)}(e)$$

extends linearly to define the boundary map $\partial^\tau : \mathbb{R}[\mathcal{P}] \rightarrow \mathbb{R}[\mathcal{E}]$. We then set

$$\partial := \partial^t - \partial^\tau.$$

A vector $v \in \mathbb{R}[\mathcal{P}]$ is said to *satisfy the gluing equations* if $\partial(v) = 0$.

The domain of the optimisation problem of interest to us is the cone of non-negative vectors that satisfy the gluing equations. To this end, set

$$\mathbb{R}_{\geq 0}[\mathcal{P}] := \left\{ \sum_{P \in \mathcal{P}} v_P P \mid v_P \geq 0 \text{ for all } P \right\}.$$

We are interested in the cone

$$C(\mathbb{R}) := \ker \partial \cap \mathbb{R}_{\geq 0}[\mathcal{P}],$$

and also in the sets of integer points $C(\mathbb{Z}) := C(\mathbb{R}) \cap \mathbb{Z}[\mathcal{P}]$ and of rational points $C(\mathbb{Q}) := C(\mathbb{R}) \cap \mathbb{Q}[\mathcal{P}]$. The key fact here is that the map to weight vectors surjects the integer points of $C(\mathbb{R})$. It is also finite-to-one, although we will not make use of that here.

Proposition 4.7 *Let X be a finite 2-complex. If Y is a finite, visibly irreducible complex, Z is also a complex and*

$$Y \rightarrow Z \rightarrow X$$

is a face immersion then $v(Y) \in C(\mathbb{Z})$. Conversely, if $v \in C(\mathbb{Z})$ then there is a face immersion from a finite visibly irreducible complex

$$Y \rightarrow Z \rightarrow X$$

such that $v = v(Y)$.

Proof Consider an edge piece $R \in \mathcal{E}$ over X . The R -coefficient of $\partial^l(v(Y))$ counts the number of vertices z of Z for which there is an edge e with $\iota(e) = z$ such the induced edge piece $R_Y(e)$ is isomorphic to R . Since Z is a complex, such vertices z are in bijection with such edges e , and so this coordinate is equal to the number of edges e of Z such that $R_Y(e)$ is isomorphic to R : notice that the latter quantity does not mention ι . Similarly, the R -coefficient of $\partial^r(v(Y))$ also counts the number of such edges e , and so $\partial^l(v(Y)) = \partial^r(v(Y))$. Therefore, $v(Y) \in C(\mathbb{Z})$ as claimed.

To prove surjectivity, consider an integral weight vector $v \in C(\mathbb{Z})$, and define the pre-complexes

$$Z' := \coprod_{P \in \mathcal{P}} \coprod_{i_P=1}^{v_P} Z(P)$$

and

$$Y' := \coprod_{P \in \mathcal{P}} \coprod_{i_P=1}^{v_P} Y(P),$$

noting that these come naturally equipped with maps to define a face immersion

$$Y' \rightarrow Z' \rightarrow X$$

Furthermore, Y' is visibly irreducible by construction.

The gluing equations imply that for each edge piece R there is a bijection between the set of edges $e \in E_{Z'}^l$, such that $R_{Y'}(e) = R$ and the set of $e \in E_{Z'}^r$, such that $R_{Y'}(e) = R$. Therefore, we may choose a bijection $F : E_{Z'}^l \rightarrow E_{Z'}^r$, such that $R_{Y'}(F(e)) = R_{Y'}(e)$. Inductively applying the gluing construction from Definition 2.22, Z' can be glued up to a complex Z equipped with a branched immersion $Z \rightarrow X$ and a branched map $Y' \rightarrow Z$.

It remains to explain how to glue up Y' to create complex Y . For an edge e of Z , let $P(\iota(e))$ denote the piece used to construct the vertex $\iota(e)$ of Z , and let $P(\tau(e))$ the piece used to construct the vertex $\tau(e)$ of Z . The pre-complexes $Y(P(\iota(e)))$ and $Y(P(\tau(e)))$ each naturally induce partitions of $w_Z^{-1}(e)$, and by construction, these two partitions are equal. Therefore, there is a bijection $G : E_{Y'}^l \rightarrow E_{Y'}^r$, that respects the map $Y' \rightarrow Z$. Again, inductively applying the gluing construction from Definition 2.22, Y' can be glued up to a complex Y equipped with face-equivalence $Y \rightarrow Z$. By construction, Y is visibly irreducible. \square

4.3 Degree and curvature

The final ingredient is to notice that, given a face-essential map $f : Y \rightarrow X$, the total curvature and the degree are both realised by linear functions on $C(\mathbb{R})$.

Definition 4.8 Let $l_X(C)$ denote the number of vertices in each component C of S_X . Recall that a vertex-piece $P \in \mathcal{P}$ consists of a face immersion

$$Y(P) \rightarrow Z(P) \rightarrow X.$$

In particular, each vertex x of $S_{Y(P)}$ is sent to some component C_x of S_X . The degree map $\text{deg} : \mathbb{R}[\mathcal{P}] \rightarrow \mathbb{R}$ is now given by linearly extending the assignment

$$\text{deg}(P) := \sum_{x \in V_{S_{Y(P)}}} \frac{1}{l_X(C_x)}.$$

The point of this definition is to extend the definition of the degree of a face-essential map.

Lemma 4.9 *If*

$$Y \rightarrow Z \rightarrow X$$

is a face immersion for a face-essential map $f : Y \rightarrow X$, then

$$\text{deg}(v(Y)) = \text{deg}(f).$$

Proof For any component C of S_Y , $l_Y(C) = \text{deg}_C(f)l_X(f(C))$. Therefore,

$$\text{deg}(f) = \sum_{[C] \in \pi_0(S_Y)} \text{deg}_C(f)$$

$$\begin{aligned}
 &= \sum_{[C] \in \pi_0(S_Y)} \frac{l_Y(C)}{l_X(f(C))} \\
 &= \sum_{[C] \in \pi_0(S_Y)} \sum_{x \in V_C} \frac{1}{l_X(f(C))} \\
 &= \sum_{x \in V_{S_Y}} \frac{1}{l_X(C_x)} \\
 &= \sum_{P \in \mathcal{P}} \sum_{P_Y(y)=P} \text{deg}(P) \\
 &= \text{deg}(v(Y))
 \end{aligned}$$

as required. □

A similar observation applies to the total curvature.

Definition 4.10 For a vertex-piece $P \in \mathcal{P}$, set

$$\tau(P) := \text{deg}(P) + \#V_{Y(P)} - \#E_{Y(P)}/2$$

and extend this to a linear function $\tau : \mathbb{R}[\mathcal{P}] \rightarrow \mathbb{R}$.

Again, this extends the total curvature to a linear function on $C(\mathbb{R})$.

Lemma 4.11 *If*

$$Y \rightarrow Z \rightarrow X$$

is a face immersion for a face-essential map $f : Y \rightarrow X$, then $\tau(v(Y)) = \tau(Y)$.

Proof By definition, $\tau(Y) = \text{deg}(f) + \chi(G_Y)$, so it suffices to prove that

$$\chi(G_Y) = \sum_{v \in V_Z} (\#V_{Y(P(v))} - \#E_{Y(P(v))}/2).$$

Since $V_{Y(P(v))}$ bijects with the set of vertices of Y that map to $v \in Z$,

$$\sum_{v \in V_Z} \#V_{Y(P(v))} = \#V_Y.$$

Likewise, $E_{Y(P(v))}$ bijects with the set of edges of Y incident at a vertex mapping to v , so

$$\sum_{v \in V_Z} \#E_{Y(P(v))} = 2\#E_Y.$$

Therefore,

$$\sum_{v \in V_Z} (\#V_{Y(P(v))} - \#E_{Y(P(v))}/2) = \#V_Y - \#E_Y = \chi(G_Y)$$

as required. □

The following rationality theorem now follows by standard optimisation techniques.

Theorem 4.12 *Let X be a finite 2-complex. There is a finite, visibly irreducible complex Y_{\max} and a face-essential map $f_{\max} : Y_{\max} \rightarrow X$ such that*

$$\frac{\tau(Y_{\max})}{\deg(f_{\max})} \geq \frac{\tau(Y)}{\deg(f)}$$

for all face-essential maps $f : Y \rightarrow X$ with Y finite and visibly irreducible.

Proof Consider the rational polyhedron

$$\Delta := C(\mathbb{R}) \cap \{\deg(v) = 1\}.$$

Every coordinate vector in $\mathbb{R}[\mathcal{P}]$ has positive degree, and so

$$\mathbb{R}_{\geq 0}[\mathcal{P}] \cap \{\deg(v) = 1\}$$

is a simplex. Hence, Δ is compact.

By the simplex algorithm, the linear map τ is maximised at some vertex v_{\max} of Δ . Since Δ is rationally defined, $v_{\max} \in C(\mathbb{Q})$, and so has some multiple $u_{\max} \in C(\mathbb{Z})$. By Proposition 4.7, $u_{\max} = v(Y_{\max})$ for some face immersion

$$Y_{\max} \rightarrow Z_{\max} \rightarrow X,$$

with Y_{\max} finite and visibly irreducible. In particular, by Lemma 2.21, the morphism $f_{\max} : Y_{\max} \rightarrow X$ is face-essential.

For any face-essential map $f : Y \rightarrow X$, the factorisation through the folded representative defines a face immersion

$$Y \rightarrow \bar{Y} \rightarrow X$$

by Lemma 2.21, so $v(Y)/\deg(f) \in \Delta$ and

$$\frac{\tau(Y_{\max})}{\deg(f_{\max})} = \tau(v_{\max}) \geq \frac{\tau(Y)}{\deg(f)}$$

as required. □

Combined with Corollary 3.23, we can complete the proof of Theorem C.

Proof of Theorem C Let $f_{\max} : Y_{\max} \rightarrow X$ be the face-essential map provided by Theorem 4.12, and set

$$\epsilon := -\frac{\tau(Y_{\max})}{\deg(f_{\max})}.$$

Since X has negative immersions and Y_{\max} is visibly irreducible, Corollary 3.23 applies to show that $\tau(Y_{\max}) < 0$, whence $\epsilon > 0$ as required.

Suppose now that Y_0 is a finite, irreducible complex and that $f_0 : Y_0 \rightarrow X$ is an immersion; in particular, $\tau(Y_0) = \chi(Y_0)$ and $\deg(f)$ is the number of faces of Y_0 . By Lemma 3.4, Y_0 unfolds to a finite, visibly irreducible 2-complex $Y \rightarrow Y_0$, and the composition

$$f : Y \rightarrow Y_0 \xrightarrow{f_0} X$$

is face-essential. Therefore,

$$\begin{aligned} -\epsilon &\geq \frac{\tau(Y)}{\deg(f)} \\ &= \frac{\tau(Y_0)}{\deg(f_0)} \\ &= \frac{\chi(Y_0)}{\#\{2\text{-cells of } Y\}} \end{aligned}$$

as required. □

5 The case with torsion

We shall see that Theorem C leads to strong constraints on the subgroup structure of one-relator groups with negative immersions. As stated, it only applies to one-relator groups without torsion. Baumslag’s coherence conjecture was proved in the case with torsion independently by the authors [26] and Wise [44]. The purpose of this section is to note that one-relator groups with torsion also fit into the framework of this paper.

The presentation complex of a one-relator group with torsion also has uniform negative immersions, after passing to a finite-sheeted covering space.³

Theorem 5.1 *Consider a one-relator group with torsion $G = F/\langle\langle u^n \rangle\rangle$, where $n > 1$. There is a torsion-free subgroup G_0 of finite index in G that is the fundamental group of a finite 2-complex X_0 with uniform negative immersions.*

Proof By [26, Theorem 2.2], there is a torsion-free subgroup G_0 of G that is the fundamental group of a complex X_0 . As in the proof of [26, Theorem 2.2], X_0 can be obtained from the presentation complex X of G by passing to a finite-sheeted

³Alternatively, one might work with a presentation orbi-complex, which itself has uniform negative immersions for maps from complexes.

cover and collapsing duplicate faces. By [26, Corollary 3.1], any immersion $Y \looparrowright X_0$ satisfies

$$\chi(Y) + (n - 1)\#\{2\text{-cells of } Y\} \leq 0$$

as long as Y is finite and irreducible. Rearranging, we obtain that

$$\frac{\chi(Y)}{\#\{2\text{-cells of } Y\}} \leq 1 - n < 0,$$

so X_0 has uniform negative immersions as claimed. □

6 Uniform negative immersions and subgroups

In this section we complete the proof of Theorems A and B, by proving these properties for the fundamental group of any complex X with uniform negative immersions. Our strategy is similar to the proof of coherence for one-relator groups with torsion [26, 44]. There is, however, a small extra technicality, because we need to deal with irreducible 2-complexes, rather than non-visibly-reducible 2-complexes; this extra technicality is resolved by Lemmas 3.10 and 3.14.

Let G be a one-relator group with negative immersions, let X be its standard presentation complex, and let H be a finitely generated subgroup of G . The next lemma is the key consequence of uniform negative immersions that we will make use of. Two immersions of 2-complexes $Y_1 \rightarrow X$ and $Y_2 \rightarrow X$ are *isomorphic* if there is an isomorphism of 2-complexes $Y_1 \rightarrow Y_2$ such that the obvious diagram commutes.

Lemma 6.1 *Let X be a finite 2-complex with uniform negative immersions. For any integer r , there are only finitely many isomorphism classes of immersions $Y \rightarrow X$ such that Y is finite, connected, irreducible, and $b_1(Y) \leq r$ (where $b_1(Y)$ denotes the first Betti number of Y).*

Proof Consider such an immersion $Y \looparrowright X$. Rearranging the defining inequality of uniform negative immersions, we have that

$$\#\{2\text{-cells of } Y\} \leq \frac{-\chi(Y)}{\epsilon}.$$

But

$$\chi(Y) \geq 1 - b_1(Y) \geq 1 - r$$

since Y is a 2-complex, so $\#\{2\text{-cells of } Y\} \leq (r - 1)/\epsilon$. Thus, there is a uniform bound on the number of faces of Y . Because Y is irreducible, every vertex or edge of Y is incident at a face. Since $Y \looparrowright X$ is an immersion, there is a bound on the number of faces at which each vertex and edge of Y are incident, so the bound on the number of faces implies bounds on the numbers of vertices and edges. Therefore, there are only finitely many possible combinatorial types for Y , and also for immersions $Y \looparrowright X$. □

Note that Lemma 6.1 fails for 2-complexes with non-positive immersions. For instance, the torus has infinitely many non-isomorphic finite-sheeted covering spaces, all with fundamental groups generated by 2 elements.

Now consider a finitely generated subgroup H of $G = \pi_1(X)$. By Grushko’s theorem, we may assume that H is freely indecomposable. The next lemma is very similar to [26, Lemma 4.2]; however, the reader should beware that the definition of irreducible complex used here is more restrictive than the definition in [26].

Lemma 6.2 *Let X be a 2-complex, $G = \pi_1(X)$, and $H \leq G$ a finitely generated, non-cyclic, freely indecomposable subgroup. After replacing H by a conjugate, there is a sequence of π_1 -surjective immersions of compact, connected 2-complexes*

$$Y_0 \twoheadrightarrow Y_1 \twoheadrightarrow Y_2 \twoheadrightarrow \dots \twoheadrightarrow Y_i \twoheadrightarrow \dots X$$

with the following properties:

- (i) each Y_i is irreducible;
- (ii) $H = \varinjlim \pi_1 Y_i$.

Because of the stronger notion of irreducibility used in this paper, the proof of Lemma 6.2 requires one extra ingredient: the irreducible core of Lemma 3.10. With this in hand, the proof of Lemma 6.2 is very similar to the proof of [26, Lemma 4.3].

Proof of Lemma 6.2 [26, Lemma 4.3] (see also Remark 2.14) provides a sequence of π_1 -surjective immersions of 2-complexes

$$Z_i \rightarrow Z_{i+1} \xrightarrow{f_{i+1}} X$$

such that $H = \varinjlim \pi_1 Z_i$. It remains to improve this to make the terms irreducible.

Since H is non-cyclic, no Z_i has cyclic fundamental group. By Scott’s lemma (proved with extra hypotheses as [34, Lemma 2.2], proved as [37, Theorem 2.1]), since H is freely indecomposable, after throwing away finitely many terms from the sequence we may assume that each $\pi_1(Z_i)$ is freely indecomposable.

Applying Lemmas 3.10 and 3.14 inductively constructs a second sequence

$$\begin{array}{ccccccccccc} Y_0 & \longrightarrow & Y_1 & \longrightarrow & Y_2 & \longrightarrow & \dots & \longrightarrow & Y_i & \longrightarrow & \dots \\ \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \\ Z_0 & \longrightarrow & Z_1 & \longrightarrow & Z_2 & \longrightarrow & \dots & \longrightarrow & Z_i & \longrightarrow & \dots \longrightarrow X \end{array}$$

where each Y_i is the irreducible core \widehat{Z}_i of Z_i . Now choose a basepoint in Y_0 , if necessary moving the basepoints of the Z_i — this may involve conjugating H . The result now follows. □

The next theorem, which is an easy consequence of Lemmas 6.1 and 6.2, will imply Theorem A. Its proof is an elaboration of the arguments of [26] or [44].

Theorem 6.3 *Let X be a finite 2-complex with uniform negative immersions, and let $G = \pi_1(X)$. For any finitely generated, non-cyclic, freely indecomposable subgroup H of G , there is an immersion from a finite, irreducible 2-complex Y to X such that $\pi_1(Y) \cong H$ and $Y \looparrowright X$ induces the inclusion of H into G up to conjugacy.*

Proof After possibly conjugating H , Lemma 6.2 provides a sequence of π_1 -surjective immersions of compact, connected, irreducible complexes

$$Y_0 \looparrowright Y_1 \looparrowright Y_2 \looparrowright \cdots \looparrowright Y_i \looparrowright \cdots X$$

such that $H = \varinjlim \pi_1 Y_i$. Since the maps $Y_i \rightarrow Y_{i+1}$ are surjective on fundamental groups, there is a uniform bound

$$b_1(Y_0) \geq b_1(Y_i)$$

on the first Betti numbers. Therefore, by Lemma 6.1, infinitely many of the immersions $Y_i \looparrowright X$ are isomorphic, so passing to a subsequence we may assume that they are all equal to a fixed immersion $Y \looparrowright X$. By [42, Lemma 6.3], each immersion $Y_i \looparrowright Y_{i+1}$ is an isomorphism. In particular, since $H = \varinjlim \pi_1(Y_i)$, every $Y \rightarrow X$ represents H , as required. \square

Coherence is an immediate consequence.

Corollary 6.4 *If X is a finite 2-complex with uniformly negative immersions then $\pi_1(X)$ is coherent.*

Proof Let H be a finitely generated subgroup of G . By Grushko's theorem, H can be written as a free product

$$H = F * H_1 \dots * H_n$$

where F is free and each H_i is freely indecomposable. Each factor H_i is finitely presented by Theorem 6.3, so the result follows. \square

Theorem A now follows quickly.

Proof of Theorem A The theorem follows by combining Theorems 1.2 and C with Corollary 6.4. \square

Likewise, the conclusions of Theorem B also hold for all complexes with uniform negative immersions.

Theorem 6.5 *Let X be a finite 2-complex with uniform negative immersions, and let $G = \pi_1(X)$.*

- (i) *Every finitely generated one-ended subgroup H of G is co-Hopfian, i.e. H is not isomorphic to a proper subgroup of itself. In particular, G itself is co-Hopfian.*

- (ii) For any integer r , there are only finitely many conjugacy classes of finitely generated one-ended subgroups H of G such that the abelianisation of H has rational rank at most r .
- (iii) Every finitely generated non-cyclic subgroup H of G is large in the sense of Pride, i.e. there is a subgroup H_0 of finite index in H that surjects a non-abelian free group.

Proof Let H be a non-cyclic, finitely generated subgroup of G .

To prove item (i), assume that H is freely indecomposable, so is realised up to conjugacy by an immersion $Y = Y_0 \looparrowright X$, by Theorem 6.3. Given any injective homomorphism $H \rightarrow H$, applying Theorem 6.3 inductively gives rise to a sequence of immersions of finite, connected, irreducible complexes

$$\dots \looparrowright Y_i \looparrowright \dots \looparrowright Y_1 \looparrowright Y_0 \looparrowright X,$$

where $\pi_1(Y_i) \cong H$ for all i . By Lemma 6.1, there are finitely many isomorphism classes of complexes Y_i so, passing to a subsequence, we may assume that $Y_i = Y$ for some fixed Y and all n . Since Y is compact, every immersion $Y \looparrowright Y$ is an isomorphism [42, Lemma 6.3], so the homomorphism $H \rightarrow H$ is also surjective, as required.

Item (ii) is an immediate consequence of Theorem 6.3 and Lemma 6.1.

Finally, to prove item (iii), start by assuming that H is freely indecomposable. By Theorem 6.3, up to conjugacy, the inclusion of H into G is induced by an immersion $Y \looparrowright X$, where Y is a finite, connected, irreducible 2-complex. By negative immersions, $\chi(Y) \leq -1$. A choice of maximal tree in the 1-skeleton of Y leads to a presentation for H with m generators and n relators, where $\chi(Y) = 1 - m + n$, so the deficiency of this presentation is

$$m - n = 1 - \chi(Y) \geq 2.$$

Therefore, by the Baumslag–Pride theorem [4], H is large, as required. For general H , consider the Grushko decomposition

$$H = F * H_1 * \dots * H_k$$

where F is free and each H_i is non-cyclic and freely indecomposable. Since the factors H_i are all large, it follows that H is large unless H is cyclic. □

Theorem B follows immediately from Theorems C and 6.5.

We close by noting that Lemma 6.1 above can be made somewhat effective.

Proposition 6.6 *Let X be the presentation complex of a one-relator group $G = F/\langle\langle w \rangle\rangle$ with negative immersions. There is an algorithm that takes w as input and outputs a constant $C = C(w)$ such that every one-ended subgroup generated H of G with abelianisation of rational rank at most r is represented by an immersion $Y \looparrowright X$ with at most $C(r - 1)$ 2-cells.*

Proof Theorem C asserts that X has uniform negative immersions, meaning that there is an $\epsilon > 0$ such that

$$\frac{\chi(Y)}{\#\{2\text{-cells of } Y\}} \leq -\epsilon,$$

for every irreducible Y immersing into X . The constant ϵ is an extremal value of the linear-programming problem used in the proof of Theorem 4.12. This linear-programming problem is explicitly defined in terms of X , so ϵ can be computed from the data of X . The proof of Lemma 6.1 now shows that the number of 2-cells of Y is at most $(r - 1)/\epsilon$, so we may take $C = 1/\epsilon$. \square

Note that Proposition 6.6 does not immediately imply that there is an algorithm to compute presentations for finitely generated subgroups of one-relator groups. In [16], a coherent group in which there is an algorithm to compute a finite presentation for a given finitely generated subgroup is called *effectively coherent*. The following question remains open.

Question 6.7 Are one-relator groups with negative immersions effectively coherent?

Acknowledgements The authors would like to thank Chris Cashen for computing the primitivity rank of Example 1.6 and the anonymous referee for numerous suggested improvements to the exposition. Louder also thanks Mladen Bestvina for explaining the tower, Wise's ideas, and the Scott lemma to him back in Winter 2004.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Arzhantseva, G.N., Ol'shanskii, A.Y.: Generality of the class of groups in which subgroups with a lesser number of generators are free. *Mat. Zametki* **59**(4), 489–496, 638 (1996)
2. Baumslag, G.: A non-cyclic one-relator group all of whose finite quotients are cyclic. *J. Aust. Math. Soc.* **10**, 497–498 (1969)
3. Baumslag, G.: Some problems on one-relator groups. In: *Proceedings of the Second International Conference on the Theory of Groups* (Australian Nat. Univ., Canberra, 1973), Berlin. *Lecture Notes in Math.*, vol. 372, pp. 75–81. Springer, Berlin (1974)
4. Baumslag, B., Pride, S.J.: Groups with two more generators than relators. *J. Lond. Math. Soc.* (2) **17**(3), 425–426 (1978)
5. Calegari, D.: Surface subgroups from homology. *Geom. Topol.* **12**(4), 1995–2007 (2008)
6. Calegari, D.: *scl*. MSJ Memoirs, vol. 20. Math. Soc. Japan, Tokyo (2009)
7. Calegari, D.: Stable commutator length is rational in free groups. *J. Am. Math. Soc.* **22**(4), 941–961 (2009)
8. Cashen, C.H.: Personal Communication (2023)

9. Cashen, C.H., Hoffmann, C.: Short, highly imprimitive words yield hyperbolic one-relator groups. *Exp. Math.* **32**(4), 631–640 (2023)
10. Cashen, C.H., Macura, N.: Line patterns in free groups. *Geom. Topol.* **15**(3), 1419–1475 (2011)
11. Culler, M.: Using surfaces to solve equations in free groups. *Topology* **20**(2), 133–145 (1981)
12. Delzant, T.: L'image d'un groupe dans un groupe hyperbolique. *Comment. Math. Helv.* **70**(2), 267–284 (1995)
13. Duncan, A.J., Howie, J.: The genus problem for one-relator products of locally indicable groups. *Math. Z.* **208**(2), 225–237 (1991)
14. Gardam, G., Woodhouse, D.J.: The geometry of one-relator groups satisfying a polynomial isoperimetric inequality. *Proc. Am. Math. Soc.* **147**(1), 125–129 (2019)
15. Gromov, M.: Hyperbolic groups. In: *Essays in Group Theory*. Math. Sci. Res. Inst. Publ., vol. 8, pp. 75–263. Springer, New York (1987)
16. Groves, D., Wilton, H.: Enumerating limit groups. *Groups Geom. Dyn.* **3**(3), 389–399 (2009)
17. Hanany, L., Puder, D.: Word measures on symmetric groups. *Sémin. Lothar. Comb.* **84B**, 79 (2020)
18. Helfer, J., Wise, D.T.: Counting cycles in labeled graphs: the nonpositive immersion property for one-relator groups. *Int. Math. Res. Not.* **9**, 2813–2827 (2016)
19. Heuer, N.: Constructions in stable commutator length and bounded cohomology. Ph.D. thesis, University of Oxford (2019)
20. Heuer, N., Löh, C.: Simplicial volume of one-relator groups and stable commutator length. *Algebraic Geom. Topol.* **22**(4), 1615–1661 (2022)
21. Ilya, K.: Primitivity rank for random elements in free groups. *J. Group Theory* **25**(5), 823–835 (2022)
22. Italiano, G., Martelli, B., Migliorini, M.: Hyperbolic 5-manifolds that fiber over S^1 . *Invent. Math.* **231**(1), 1–38 (2023)
23. Hall, M. Jr.: Subgroups of finite index in free groups. *Can. J. Math.* **1**, 187–190 (1949)
24. Linton, M.: One-relator hierarchies (2022). [arXiv:2202.11324](https://arxiv.org/abs/2202.11324)
25. Louder, L., Wilton, H.: Stackings and the W -cycles conjecture. *Can. Math. Bull.* **60**(3), 604–612 (2017)
26. Louder, L., Wilton, H.: One-relator groups with torsion are coherent. *Math. Res. Lett.* **27**(5), 1499–1512 (2020)
27. Louder, L., Wilton, H.: Negative immersions for one-relator groups. *Duke Math. J.* **171**(3), 547–594 (2022)
28. Manning, J.F.: Virtually geometric words and Whitehead's algorithm. *Math. Res. Lett.* **17**(5), 917–925 (2010)
29. Martínez-Pedroza, E., Wise, D.T.: Coherence and negative sectional curvature in complexes of groups. *Mich. Math. J.* **62**(3), 507–536 (2013)
30. Puder, D.: Primitive words, free factors and measure preservation. *Isr. J. Math.* **201**(1), 25–73 (2014)
31. Puder, D.: Expansion of random graphs: new proofs, new results. *Invent. Math.* **201**(3), 845–908 (2015)
32. Rips, E., Sela, Z.: Structure and rigidity in hyperbolic groups. I. *Geom. Funct. Anal.* **4**(3), 337–371 (1994)
33. Sapir, M., Špakulová, I.: Almost all one-relator groups with at least three generators are residually finite. *J. Eur. Math. Soc.* **13**(2), 331–343 (2011)
34. Scott, G.P.: Finitely generated 3-manifold groups are finitely presented. *J. Lond. Math. Soc. (2)* **6**, 437–440 (1973)
35. Sela, Z.: Structure and rigidity in (Gromov) hyperbolic groups and discrete groups in rank 1 Lie groups. II. *Geom. Funct. Anal.* **7**(3), 561–593 (1997)
36. Stallings, J.R.: Topology of finite graphs. *Invent. Math.* **71**(3), 551–565 (1983)
37. Swarup, G.A.: Delzant's variation on Scott complexity (2004). [arXiv:math/0401308](https://arxiv.org/abs/math/0401308)
38. Whitehead, J.H.C.: On equivalent sets of elements in a free group. *Ann. Math. (2)* **37**(4), 782–800 (1936)
39. Wilton, H.: Essential surfaces in graph pairs. *J. Am. Math. Soc.* **31**(4), 893–919 (2018)
40. Wilton, H.: Rational curvature invariants for 2-complexes (2022). [arXiv:2210.09853](https://arxiv.org/abs/2210.09853)
41. Wise, D.T.: Coherence, local-indicability and non-positive immersions (2003). <http://www.gidon.com/dani/tl.cgi?athe=pspapers/NonPositiveCoherence.ps>. Preprint
42. Wise, D.T.: Sectional curvature, compact cores, and local quasiconvexity. *Geom. Funct. Anal.* **14**(2), 433–468 (2004)

43. Wise, D.T.: Nonpositive sectional curvature for (p, q, r) -complexes. Proc. Am. Math. Soc. **136**(1), 41–48 (2008)
44. Wise, D.T.: Coherence, local-indicability and nonpositive immersions. J. Inst. Math. Jussieu **First view**, 1–16 (2020)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.