

Climate-related health impact indicators for public health surveillance in a changing climate: a systematic review and local suitability analysis

Yasna Palmeiro-Silva,^{a,*} Raúl Aravena-Contreras,^b José Izcue Gana,^c Rocío González Tapia,^d and Ilan Kelman^{a,e,f}

^aInstitute for Global Health, University College London, London, United Kingdom

^bDepartment of Chemical Engineering, Imperial College London, United Kingdom

^cInstitute for Global Prosperity, University College London, London, United Kingdom

^dSociedad Chilena de Salud Planetaria, Puerto Varas, Chile

^eInstitute for Risk and Disaster Reduction, University College London, London, United Kingdom

^fUniversity of Agder, Norway



Summary

Climate change challenges public health. Effective management of climate-related health risks relies on robust public health surveillance (PHS) and population health indicators. Despite existing global and country-specific indicators, their integration into local PHS systems is limited, impacting decision-making. We conducted a systematic review examining population health indicators relevant to climate change impacts and their suitability for national PHS systems. Guided by a registered protocol, we searched multiple databases and included 41 articles. Of these, 35 reported morbidity indicators, and 39 reported mortality indicators. Using Chile as a case study, we identified three sets of indicators for the Chilean PHS. The high-priority set included vector-, food-, and water-borne diseases, as well as temperature-related health outcomes indicators due to their easy integration into existing PHS systems. This review highlights the importance of population health indicators in monitoring climate-related health impacts, emphasising the need for local contextual factors to guide indicator selection.

Funding This research project was partly funded by ANID Chile and University College London. None of these sources had any involvement in the research conceptualisation, design, or interpretation of the results.

Copyright © 2024 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).

Keywords: Climate change; Population health; Surveillance; Indicators; Health impacts

Introduction

Although climate-related health outcomes have always occurred throughout the history of humanity, current anthropogenic climate change (hereafter climate change) is changing the pattern and exerting significant pressure on public health and health systems.^{1,2} With global warming currently at 1.2 °C above pre-industrial levels, the world is witnessing record-breaking climate hazards³ that are severely impacting people's health and wellbeing.⁴ Under current climate change policies, global surface temperatures are likely to continue rising and even exceed 2 °C above pre-industrial levels by 2100, intensifying climate hazards and weather events, hence further affecting people's health, wellbeing, and livelihoods.^{5,6} In this evolving scenario, a comprehensive understanding of population exposure to these hazards, vulnerability factors, and potential climate-related health outcomes

is vital for effective and proactive health adaptation, hence mitigating associated health risks. Recently, this issue has been captured by the UEA Framework for Global Climate Resilience agreed at COP28, which established health as a key sector for climate change adaptation.⁷

Public health surveillance (PHS), is an essential function of public health,⁸ and refers to the ongoing systematic collection, analysis, and interpretation of data that can be used for planning and implementation of health interventions, programmes, and policies according to social contexts.^{9–11} Nonetheless, several challenges exist for PHS to effectively serve its purpose, such as integrating diverse data sources, ensuring complete high-quality data, and employing comprehensive and policy-relevant indicators.^{11–13}

Historically, PHS systems have been primarily focused on infectious diseases,^{13,14} but have expanded their scope given demographic and epidemiological population changes and the emergence of different hazards and technologies. Some PHS systems now include information on injuries, noncommunicable

*Corresponding author. Institute for Global Health, University College London, London, United Kingdom.

E-mail address: y.palmeiro@ucl.ac.uk (Y. Palmeiro-Silva).

The Lancet Regional Health - Americas 2024;38: 100854

Published Online xxx
<https://doi.org/10.1016/j.lana.2024.100854>

diseases, lifestyle factors, and environmental hazards.¹² From an environmental perspective, the most common hazards and exposures under surveillance have included chemicals, pesticides, radiation, and metalloids, as well as air, water, and soil pollution.^{15–19} However, the inclusion of climatic and meteorological factors (e.g., heatwaves) is still in its infancy, and few public health agencies, mostly from High-Income countries, have considered them as part of PHS.²⁰

Global²¹ and country-specific^{22–25} indicators at the intersection of climate and health have been developed, such as those from The *Lancet* Countdown, providing valuable information to other countries or locations. However, some challenges impede their seamless integration into national or sub-national PHS systems, including local relevance, context-specific factors that partly determine public health responses (e.g., socio-political context and health inequities), and PHS systems characteristics (e.g., local resources and capacities). In consequence, it is not quite straightforward to determine which indicators can be used as part of PHS at national and sub-national levels to track the impacts of climate-related hazards on population health under a changing climate.

From a practical perspective and before proposing new and potentially reiterative climate-related health impact indicators for PHS, it seems sensible to explore the evidence already published as well as its potential use in a local context. Therefore, we systematically reviewed the evidence on population health indicators related to the impacts of climate hazards under a changing climate. We then created and applied a guide that assesses the suitability of the indicators for their potential integration into a national PHS system. As an example, we present the case of Chile; however, this guide can be applied in other national or sub-national settings, especially in Latin America.

Methods

Study design, research question, and protocol

We conducted a systematic review following our previous published protocol (PROSPERO [CRD42021253704](https://doi.org/10.1186/CRD42021253704)),²⁶ and adhered to PRISMA guidelines for reporting.²⁷ The research question was “*What are the population indicators that monitor the health effects/impacts of climate change?*”, which was broad and open in order to comprehensively search the evidence on the topic.

Information sources

To cover most of the scientific evidence, we searched seven multidisciplinary and specialised databases: OVID/MEDLINE, Scopus, Web of Science Core Collection, PubMed, Cochrane Library, SciELO, and BIREME/LILACS. Complementary, we included grey

literature from the World Health Organisation (WHO), the United Nations (UN), the European Centre for Disease Prevention and Control (ECDC), and Centre for Disease Prevention and Control (CDC) of the United States of America (USA). These sources were chosen for their role in informing public policy and their focus on climate change and population health.

Eligibility criteria

Studies were included if they i) were available in English, Spanish, or Portuguese; ii) focused on human populations; and iii) presented, summarised, discussed, or analysed population health indicators related to the impacts of climate change on population health. We imposed no restrictions on publication date, study type, or geographical scope. Unpublished work, news, or opinion articles were excluded due to the potential for redundant information and high risk of bias.

Search strategies

A generic search strategy consisted of: (*indicator* OR measure* OR metric* OR index OR indices*) AND (*health OR well?being*) AND (*monitor OR surveillance*) AND (*“climate* change*” OR “global warming” OR “climate crisis” OR “climate* varia*”*). The words “*effects*” or “*impacts*” were not included to make the search as broad as possible; however, any article unrelated to effects/impacts of climate change was excluded based on the eligibility criteria described above. Depending on the number and relevance of articles retrieved, the filter “*humans*” or keyword “*human*” was considered. The searches were conducted from the 15th to 24th of October 2021. [Supplementary Tables S1 and S2](#) in the [Supplementary Material](#) present related terms and detailed search strategies for each database.

Two search updates were done: one in August 2022, which was conducted via Google Scholar using the syntax (“*climate change*” AND *indicators* AND *impacts* AND “*human health*”). The second update took place on the 12th of June 2024 and considered a revision of all seven multidisciplinary databases mentioned above and followed the same search strategies as in [Supplementary Tables S1 and S2](#), plus a Google Scholar search. We also screened the reference lists of selected studies to identify any additional relevant articles not captured in the previous searches.

Study selection process

The retrieved articles were imported into Covidence.²⁸ After deduplication, two researchers (YPS and RAC or JIG) independently screened titles and abstracts, achieving an agreement rate of 90%. Full texts were also independently screened by two researchers, resulting in a 74% agreement rate. Titles, abstracts, and articles in disagreement were discussed with a third reviewer (RGT). Screening of the reference lists and grey

literature was performed by one reviewer (YPS) in August 2022 and June 2024.

Data extraction process

Information on the title; year of publication; authors; type of publication; region or country; brief context; aim; overview of the methods; population considered; eligibility criteria; funding; conflict of interest; and main findings was extracted using a form designed in Covidence. The data extraction process was piloted using one article and necessary adjustments were made thereafter. Two reviewers independently extracted the data from each study (YPS and RAC or JIG), resolving any discrepancies through thorough re-analysis and discussion of the article in reference until complete agreement was reached.

Quality assessment

Given the broad scope of the research question, a diverse range of study designs was anticipated. Consequently, two complementary critical appraisal tools were employed: the Quality Assessment for Diverse Studies (QuADS), which facilitates the evaluation of various study designs in systematic reviews,²⁹ and the Joanna Briggs Institute (JBI) critical appraisal tools.³⁰ Each article, excluding grey literature, underwent appraisal by two researchers (YPS and RAC or JIG), with any discrepancies resolved through joint re-analysis and discussion. The critical appraisal was not used as a basis for inclusion/exclusion of articles, but rather to assess the overall quality of evidence and inform the suitability analysis. However, a qualitative sensitivity analysis was conducted in order to evaluate the potential influence of low quality/high risk of bias articles in the results, and thus, ensuring rigour and transparency of this review.

Data synthesis

We synthesised the evidence using a thematic approach with no pre-imposed themes.^{31,32} Although we anticipated groups of all-cause and cause-specific morbidity and mortality indicators, we remained open to other categories emerging from the data. The indicators are presented as they were published to accurately reflect the breadth of evidence and mitigate potential interpretation biases.

Methods for the suitability analysis

A suitability analysis is essential for identifying scientifically sound and policy-relevant indicators that can be integrated into PHS systems. However, there is no standard process for this purpose; therefore, we developed a guide that comprises four dimensions and 13 sub-dimensions (Table 1; Spanish and Portuguese versions in Supplementary Tables S3 and S4). The development of this guide was performed in parallel to the current systematic review and consisted in a systematic approach of the following steps:

- 1) A literature review of the principles of PHS systems was conducted and included information from scientific literature, international frameworks (e.g., CDC and NHS England), and measurement theory.^{33–44}
- 2) Information from step 1 was synthesised and analysed by one researcher (YPS) with feedback from a second senior researcher (IK). Several indicators' attributes or sub-dimensions were identified, which were grouped into four broad dimensions: scientific foundations that represent epidemiological attributes of the indicators; measurability, which refers to the capacity of the indicators of being measured; relevance to public health, which is one of the main purposes of the indicators in public health surveillance; and contextual to climate change, which refers to the indicators' attributes related to inherent climate change characteristics (e.g., long-term changes) rather than to any indicator for public health surveillance.
- 3) The first draft of the guide was then analysed with two policymakers in Chile and two external researchers to ascertain its precision and usefulness in guiding the suitability analysis of indicators. Experts looked at: definitions, words, assigned scores, and overall completeness of the guide. Feedback was integrated into the final guide.

By employing a simple 3-level scoring system for each sub-dimension, the guide facilitates indicator analysis and provides users with insights into the overall suitability of indicators for addressing population health and climate change concerns within a local context. Scores are not numerically combinable as they only represent categories.

Results

Findings from the systematic review

Searches in 2021 yielded 1587 references in scientific databases, of which 1068 remained after de-duplication, and 68 after screening titles and abstracts. After full-text review, 16 articles met eligibility criteria, while 52 were excluded: 30 did not include impact indicators^{45–74}; seven were unrelated to human health^{75–81}; eleven did not include indicators related to climate change^{82–92}; three were inaccessible^{93–95}; and one did not meet language criteria.⁹⁶ Three additional articles were included from reference lists,^{19,97,98} and six from grey literature.^{99–104} From the search update in August 2022, 80 references were identified, with four meeting eligibility criteria.^{105–108}

The search update on the 12th of June 2024 yielded 963 references, of which 750 remained after de-duplication, and 11 after screening titles and abstracts. After full-text review, only five met eligibility criteria and six were excluded because did not explicitly include

Dimension I: Scientific foundations	
Sub-dimension	Definition and scores
Ia) Validity	<p><i>Degree to which a measurement measures what it purports to measure.</i>³³</p> <ul style="list-style-type: none"> Score 0: Indicator does not measure what it intends to measure Score 1: Indicator unclearly measure what intents to measure Score 2: Indicator clearly measures what intents to measure
Ib) Reliability	<p><i>Degree of stability when a measurement is repeated under identical or similar conditions, either over time, different populations, or different datasets.</i>³⁴</p> <ul style="list-style-type: none"> Score 0: Indicator is not stable under identical or similar conditions Score 1: Indicator is to some extent stable under identical or similar conditions Score 2: Indicator is stable under identical or similar conditions
Ic) Scientific rationale	<p><i>Degree to which an indicator is well-defined and supported by scientific evidence. Overall, scientific evidence is consistent and coherent regarding the indicator.</i></p> <ul style="list-style-type: none"> Score 0: Indicator is not well-defined <u>nor</u> supported by scientific evidence Score 1: Indicator is confusingly defined, <u>or</u> scientific evidence is not clear respect to the scientific rationale of the indicator Score 2: Indicator is well-defined <u>and</u> scientific evidence is consistent and coherent regarding the indicator
Dimension II: Measurability (capacity of being measured)	
Sub-dimension	Definition and scores
Ila) Data availability	<p><i>Degree of availability of complete, valid, and reliable data.</i></p> <ul style="list-style-type: none"> Score 0: Data are not available, <u>and</u> it is difficult to collect data within a short period of time (e.g., 1–2 years) Score 1: Data are available, but national and local authorities have restricted access. Data are available, but data collection is not frequently available (e.g., surveys every 2–3 years) Score 2: Data are frequently collected (every week, month, and/or year) <u>and</u> are already available to national and local authorities
Ilb) Data analysis	<p><i>Process of cleaning, calculating, and modelling data to obtain useful information.</i></p> <ul style="list-style-type: none"> Score 0: Data analysis is not easy to perform and needs specialised training Score 1: Data analysis is easily performed because calculation methods are already standardised and widely available Score 2: Data analysis is not needed because indicators are already calculated, analysed, and available, including modelled indicators
Dimension III: Relevance to public health	
Sub-dimension	Definition and scores
IIIa) Policy-relevance	<p><i>Degree to which an indicator is relevant to policy and informs public health policies or key policy issues according to local context.</i></p> <ul style="list-style-type: none"> Score 0: Indicator is not relevant to population health <u>and</u> does not inform public health policies or key policy issues Score 1: Indicator is relevant to a limited proportion of a target population <u>and</u> may partially inform public health policies or key policy issues Score 2: Indicator is relevant to population health, covers the whole target population, and inform public health policies or key policy issues
IIIb) Cost-effectiveness	<p><i>Degree to which an indicator achieves desired aims at reasonable investments, including time and other resources.</i></p> <ul style="list-style-type: none"> Score 0: Investments of producing and analysing the indicator exceed the value of using the indicator Score 1: Investments of producing and analysing the indicator are similar in comparison to the value of using the indicator Score 2: Investments of producing and analysing the indicator are well-below the value of using the indicator
IIIc) Simplicity	<p><i>Degree to which an indicator is easy to interpret and understand by users (i.e., decision- or policymakers and the general public).</i></p> <ul style="list-style-type: none"> Score 0: The indicator is not easy to interpret and understand by users <u>and</u> requires specialised training to do it Score 1: The indicator has standardised guidelines to be interpreted and understood by users Score 2: The indicator is easy to interpret and understand by a wide range of users
IIId) Actionability	<p><i>Degree to which an indicator allows the planning and implementation of public health policies</i></p> <ul style="list-style-type: none"> Score 0: The indicator does not allow action Score 1: The indicator allows action, but more information or other indicators are needed Score 2: The indicator allows action by itself
IIIe) Timeliness	<p><i>Degree to which an indicator can be measured and used in an appropriate period of time to inform public health policies.</i></p> <ul style="list-style-type: none"> Score 0: The indicator needs a long period of time to be measured and is not ready when it is needed Score 1: The indicator can be measured and used within the required period of time, but requires specialised skills to achieve it Score 2: The indicator is measured and ready to be used within a short period of time (e.g., real-time, hours, or days)
Dimension IV: Contextual to climate change	
Sub-dimension	Definition and scores
IVa) Specificity	<p><i>Degree to which an indicator represents health impacts associated with changes in the climate or climate-related hazards.</i>³⁸</p> <ul style="list-style-type: none"> Score 0: The indicator does not represent health impacts associated with climate-related hazards under climate change. There are important influences of other non-climatic factors Score 1: The indicator represents the association with climate-related hazards under climate change, but further analyses are required in order to clarify influences of other non-climatic factors Score 2: The indicator represents the association with climate-related hazards under climate change, and non-climatic factors are under control to some extent
IVb) Sensitivity	<p><i>Degree to which an indicator captures changes in health outcomes associated with changes in the climate or climate-related hazards.</i></p> <ul style="list-style-type: none"> Score 0: The indicator does not capture changes in health outcomes associated with climate-related hazards under climate change Score 1: The indicator partially captures changes in health outcomes associated with climate-related hazards under climate change, but further analyses are required in order to clarify potential aberrations Score 2: The indicator captures changes in health outcomes associated with climate-related hazards under climate change
IVc) Comparability	<p><i>Degree to which an indicator can be comparable over time and across different populations, including different regions and countries.</i></p> <ul style="list-style-type: none"> Score 0: The indicator is not comparable over time or across population groups due to different methods Score 1: The indicator is comparable over time and some other populations Score 2: The indicator is fully comparable over time and across populations at regional or international levels

Table 1: Definitions and scores of dimensions for analysing indicators on climate change and population health.

impact indicators.^{109–114} Seven references were found through a Google Scholar search.^{1,2,4,115–118}

A total of 41 references were included in this systematic review. Fig. 1 shows the PRISMA flow diagram.²⁷

Overall study characteristics

Articles were published between 1993 and 2024. Geographically, 13 articles had a worldwide scope^{1,4,19,97,100,102,104,107,108,119–122}; six were focused on the USA^{99,101,116,123–125}; five on Australia^{22,98,105,117,126}; three on Europe^{103,127,128}; two on Canada^{38,129}; two on China^{106,130}; and one article on: Asia,¹³¹ Korea,¹³² Ireland,³⁶ Ghana,¹³³ Latin America,¹¹⁸ South America,² Spain,¹³⁴ the UK¹³⁵; and Vietnam.¹³⁶ Regarding the grey literature, three had a worldwide scope,^{100,102,104} two were focused on the USA,^{99,101} one on Europe,¹⁰³ and one on the UK.¹¹⁵ A total of 13 articles were a literature review, either systematic or non-systematic^{36,38,115,116,119,121–123,125,127,129,131,135}; 16 were ecological studies^{1,2,4,19,22,38,97,98,105–108,117,118,120,128,130}; five were case-studies^{126,132–134,136}; and one was a cross-sectional study.¹²⁴ Tables 2 and 3 summarise the articles included in this review.

The critical appraisal based on QuADS and JBI tools showed consistent results. A total of 23 articles were considered of acceptable quality (“include”),^{1,2,4,19,22,38,97,98,105–108,117,118,120,123,127–132,135} while 11 articles were flagged as “further information is needed”.^{36,116,119,121,122,124–126,133,134,136} Overall, the weakest aspects of the articles were related to methods description, and strengths and limitations (Supplementary Table S5). These results underscore the need for caution when utilising the articles’ findings in the review and suitability analysis.

Types of indicators

A total of 35 studies reported indicators for morbidity,^{1,4,19,22,36,38,97,99–105,107,115–121,123–131,133–136} including cases of, incidence of, or prevalence of certain diseases, hospitalisations, and medical visits. A total of 39 studies reported indicators for mortality.^{1,2,4,19,22,36,38,97–108,115,116,118–135} These indicators may also be classified according to specific exposures or causes (Table 4).

Some articles also highlighted analyses of indicators by subgroups, including different age categories,^{104,126,131} sex,¹²⁶ aboriginal groups,¹²⁶ and seasons.³⁶

Findings from the suitability analysis

The suitability analysis of indicators was performed based on the dimensions in Table 1 and the Chilean context, which is extensively described in Supplementary Box S1 in the Appendix. Briefly, there are different official PHS systems implemented in Chile, which mainly cover communicable and non-communicable diseases. Regarding communicable diseases, specific ones are subject to mandatory

notification, including immediate and daily notification, as well as sentinel notification. Additionally, there are laboratory-based surveillance of specific and pre-defined microbiology agents, as well as surveillance for antimicrobial resistance.¹³⁷ A few non-communicable chronic and acute health conditions are under surveillance, including cancers,¹³⁸ pesticide poisoning,¹³⁹ and risk factors (e.g., physical inactivity, tobacco use).¹⁴⁰ There are other specific surveillance programmes focused on occupational health.¹⁴¹ All these data are collected by the Ministry of Health and analysed considering different variables. The information that emerges is then used to inform national or local health programmes. Based on this background information, Supplementary Tables S6–S8 show the collective scores for each sub-dimension and provide insights into the suitability of each indicator for the Chilean PHS system. A brief example on how the scoring process was performed is provided.

Overall, general indicators with broad disease description received lower scores in scientific foundations and contextuality to climate change, while specific indicators scored higher in these areas. Most temperature-related indicators scored high in terms of scientific foundations; however, indicators requiring advanced statistical modelling may be relevant for climate change but have limited measurability. As a result, three primary prioritised sets of climate-related health impact indicators are proposed for PHS in Chile. Table 5 presents high-priority indicators, many of which are already under surveillance and/or data are already available for analysis, but they are not currently and explicitly considered for climate change and public health surveillance. Medium- and low-priority indicators may be relevant for PHS, but their measurability and contextuality to climate change are important challenges that might limit their use in the short-term (Supplementary Tables S9 and S10).

The sensitivity analysis showed that the results were robust and low quality/high risk of bias articles did not influence the primary prioritised indicators. Supplementary Table S11 shows all indicators that were identified from low quality/high risk of bias articles only if the indicators were not identified by high quality-low risk of bias articles. Some indicators, such as cases of leishmaniasis, chikungunya, zika fever, and shellfish poisoning are in Supplementary Table S11 and in Table 5 because these indicators are already part of the PHS system in Chile and were considered as primary prioritised indicators. In the case of number of cases and deaths due to heatstroke, these indicators are covered by temperature-related diseases indicators in Table 5, which were also primarily prioritised.

Discussion

This systematic review identified population health indicators that help monitor the health impacts of climate-




Fig. 1: PRISMA flow diagram of the systematic review.

related hazards in the context of climate change. Based on the revised literature, it is possible that current PHS systems already include several of the identified indicators; however, two main challenges remain. First, to better understand these indicators in a context of climate change, it is necessary to analyse whether the PHS systems and these indicators are compatible and adaptable. Second, although there are relevant indicators that can be integrated into PHS systems, the moderate quality of the literature may limit their use; therefore, specific analyses are needed.

In terms of geographical distribution, most of the evidence comes from High-Income countries, such as Canada, the USA, and the UK. Therefore, the use and adaptation of indicators from these countries to other countries may have some limitations since, for example, data collection methods, indicators definition, and spatial and temporal coverage are not quite compatible. This issue is relevant for PHS systems because one of the main characteristics of indicators for PHS is that

they should represent and be aligned with characteristics of the local context, resulting in better use and acceptability by users.^{12,142}

Regarding critical appraisal, several articles combined literature review processes and expert consultations; however, it was not always possible to analyse the associated methods due to limited description. Although literature reviews do not follow the same systematicity as systematic reviews, unclear methodology and methods undermine their reproducibility, replicability, and overall quality.¹⁴³ It is arguable whether true reproducibility and replicability can be achieved¹⁴⁴ and sometimes these qualities are not the main goal of literature reviews; however, clear methods would facilitate their critical assessment by providing a rationale for their structure and line of argument; and therefore, increasing their use in other contexts. Complementary, expert and stakeholder engagement is an important step when translating scientific evidence into public policies^{145,146}; however, the methodology of expert

Reference	Setting & population	Brief methods	Main findings
Akearok et al. (2019) ¹²⁹	Canada, Nunavut territory	Scoping review and consultation to identify human health indicators of climate change relevant to the Canadian Arctic	i) number of injuries or mortality from extreme weather events or sea ice instability; ii) human cases of environmental infectious diseases/positive tests results in reservoirs/sentinels/vectors; iii) respiratory/allergic disease and mortality related to increased air pollution and pollens; iv) reports of depression, anxiety related to climate change.
Beggs et al. (2019) ^{98,a}	Australia	Review of scientific evidence and analysis of indicators in Australia	i) suicide rate associated with ambient temperature
Beggs et al. (2021) ^{105,a}	Australia	Review of scientific evidence and analysis of indicators in Australia	i) rates of cardiovascular and respiratory disease associated with heat stress; ii) heat stress risk of people participating in physical and sporting activities; iii) cold- and heat-related mortality; iv) dengue incidence; v) Ross River virus cases
Beggs et al. (2024) ^{117,b}	Australia	Review of scientific evidence and analysis of indicators in Australia	i) heat stress risk of people participating in physical and sporting activities
Benson et al. (2022) ^{122,b}	Worldwide	Literature review	i) Suicide mortality rate
Brown (1993) ¹²⁶	Australia	Analysis of evidence on social and environmental indicators to monitor health effects of climate change and application to Australia	i) rates of hospitalisation for circulatory diseases, respiratory diseases, and accidents, poisoning and violence; ii) infants death rates among aboriginal groups; iii) number of deaths of young men from accidents, poisoning, violence; iv) deaths before 65 years old.
Cai et al. (2021) ^{106,a}	China	Review of scientific evidence, expert consensus, and analysis of indicators in China	i) heatwaves-related mortality
Cheng et al. (2013) ³⁸	Canada	Literature review and analysis of key indicators to quantify health impacts of climate change on Canadians	Selected modelled indicators: i) excess daily all-cause mortality due to heat; ii) premature deaths due to air pollution (ozone and particulate matter 2.5); iii) preventable deaths from climate change; iv) disability-adjusted life years (DALYs) lost from climate change. Selected non-modelled indicators: i) daily all-cause mortality; ii) daily non-accidental mortality; iii) West Nile disease incidence (in humans); iv) Lyme borreliosis incidence (in humans).
Donnelly et al. (2004) ³⁶	Ireland	Review of evidence on indicators of the impact of climate change suitable for Ireland.	i) skin cancers; ii) seasonal pattern of mortality
Doubleday et al. (2020) ¹²⁵	United States of America, Pacific Northwest	Narrative review of literature and consultation with experts to develop indicators	Examples of indicators related to impacts are i) weather-related morbidity and mortality, including cold- and heat-related illness and mortality; ii) asthma exacerbations, and other respiratory hospitalisations related to wildfire events; iii) shellfish poisoning outbreaks, and iv) utilisation of mental health services.
Dovie et al. (2017) ¹³³	Ghana	Review of evidence, focus groups, and analysis as a case report.	i) mortality; ii) morbidity; iii) history of occurrence; iv) reported/confirmed cases
Driscoll et al. (2013) ¹²⁴	United States of America, Alaska	Participatory process to identify health risks related to climate change.	i) food security; ii) food security; iii) cold-related injuries and fatalities; iv) cases of allergic asthma; and v) paralytic shellfish poisoning cases.
Ebi et al. (2018) ¹²¹	Worldwide	Review of evidence and expert consultation. Case report taking Cambodia as an example.	i) excess mortality associated with exposure to high ambient temperatures; ii) all-cause and cause-specific morbidity and mortality associated with other extreme weather events; iii) respiratory disease mortality from exposure to air pollutants such as ozone and particulate matter; iv) changes in the incidence and geographic range of climate-sensitive infectious diseases; v) undernutrition. Aggregated indicators: i) disability adjusted life years or years of life lost from climate variability and climate change. Other indicators: i) injuries, illnesses, and deaths attributed to wildfires; ii) the number of asthmatic episodes associated with high pollen events.
English et al. (2009) ¹²³	United States of America	Review of scientific evidence and expert consultation.	i) excess mortality due to extreme heat; ii) excess morbidity due to extreme heat; iii) number of injuries/mortality from extreme weather events; v) human cases of environmental infectious disease/positive tests results in reservoirs/sentinels/vectors; vi) respiratory/allergic disease and mortality related to increased air pollution and pollens.
Hartinger et al. (2022) ^{2,b}	South America	Review of scientific evidence and analysis of indicators in South America	i) heat-related mortality
Hartinger et al. (2023) ^{118,b}	Latin America	Review of scientific evidence and analysis of indicators in Latin America	i) heat-related mortality; ii) heat stress risk related to physical activity
Heo et al. (2019) ¹³²	South Korea (Seoul and Busan)	Analysis of heatwaves using different thermal indices from 2011 to 2015	i) daily all-cause mortality and excess mortality associated with heat index, humidex, apparent temperature, effective temperature, WetBulb Globe Temperature (WBGT).

(Table 2 continues on next page)

Reference	Setting & population	Brief methods	Main findings
(Continued from previous page)			
Jung et al. (2017) ¹³¹	Asia, Low-and Middle-Income Countries	Systematic review and analysis of indicators that can be applied in Low- and Middle-Income countries	Indicators associated with respiratory diseases: i) incidence of morbidity due to respiratory diseases in children aged under 5 years; ii) annual mortality rate due to respiratory diseases in children aged under 5 years; iii) prevalence of chronic respiratory illnesses in children aged 0–14 years. Indicators associated with diarrhoeal diseases: i) incidence of outbreaks of water-borne diseases; ii) diarrhoea morbidity rate in children aged under 5 years; iii) diarrhoea mortality rate in children aged under 5 years; iv) recurrence rate of outbreaks of diarrhoeal disease among children aged under 5 years. Indicators associated with insect-borne diseases: i) prevalence of insect-borne diseases in children aged 0–14 years; ii) mortality rate of children aged under 5 years due to insect-borne diseases.
Liu et al. (2021) ^{116,b}	USA	Literature review	Allergies and other respiratory conditions: i) asthma mortality; ii) asthma and allergic disease-related hospitalisations; iii) asthma and allergic disease-related emergency room visits Vector-borne diseases: i) West Nile Virus neuroinvasive disease cases Vibriosis: i) <i>Vibrio</i> illnesses
Lowe et al. (2011) ¹²⁷	Europe	Scoping review of heatwave early warning system in Europe	i) real time mortality; ii) daily mortality; iii) number of calls to emergency services; iii) number of visits to general practitioner (GP); iv) heatwave-related morbidity and mortality
Mercuriali et al. (2022) ^{134,b}	Spain	Case study (presented as a field note) that included a literature review and expert consultations	Morbimortality related to heat: i) cases and deaths due to heatstroke; ii) daily deaths during heatwaves (extracted from funeral and mortality registries); iii) births, premature births, and low-weight births during heatwaves; iv) mortality attributable to heatwaves (all-cause and cause specific, hospital discharges, emergency visits, and primary healthcare services); v) deaths and mortality rate due to suicide and homicide. Vector related diseases: i) cases of dengue, Chikungunya, Zika, West Nile, malaria, leishmaniasis among residents; ii) viremic cases of dengue, Chikungunya, and Zika in periods of vector activity; iii) viremic cases of dengue, Chikungunya, and Zika with <i>Aedes albopictus</i> activity detected in the area (and proportion of the total viremic cases). Air quality: i) mortality and morbidity attributable to air pollution; ii) mortality due to asthma Water quality and water related diseases: i) Number of outbreaks related to drinkable water, leisure water, and food; ii) Notified cases of <i>Salmonella</i> , <i>Campylobacter</i> , and <i>Escherichia coli</i> .
Murage et al. (2024) ^{135,b}	United Kingdom	Three stage study that included: 1) consultation with key stakeholders, 2) literature review, and 3) expert consultations	i) annual heat-related attributable deaths; ii) heat illness indicator from syndromic surveillance data
National Research Council (2010) ¹¹⁹	Worldwide	Workshop and technical review based on scientific evidence.	Epidemics/Pandemics: i) morbidity and mortality data; ii) disability-adjusted life years; iii) human cases of environmental infectious disease/positive tests results in reservoirs/sentinels/vectors Incidence of respiratory disease: i) respiratory/allergic disease and mortality related to increased air pollution and pollens; ii) general morbidity and mortality data; iii) disability-adjusted life years; iv) cancer rates Harmful Algal Blooms (HAB): human shellfish poisonings
Nguyen et al. (2018) ¹³⁶	Vietnam	Applications of remote sensing on health and vector-borne diseases using Vietnam as a case report.	i) number of human cases of malaria
Romanello et al. (2021) ^{107,a}	Worldwide	Review of scientific evidence and analysis of indicators worldwide	i) number of hours available for safe physical activity per day; ii) general sentiment of expressions (from Twitter) associated with heatwaves; iii) heat-related excess mortality; iv) number of deaths and affected people associated with weather-related disasters
Romanello et al. (2022) ^{1,b}	Worldwide	Review of scientific evidence and analysis of indicators worldwide	i) heat stress risk related to physical activity; ii) heat-related mortality; iii) sentiment of expressions (from Twitter) associated with heatwaves
Romanello et al. (2023) ^{4,b}	Worldwide	Review of scientific evidence and analysis of indicators worldwide	i) heat stress risk related to physical activity; ii) heat-related mortality; iii) sentiment of expressions (from Twitter) associated with heatwaves
Van Daalen et al. (2022) ^{128,b}	Europe	Review of scientific evidence and analysis of indicators in Europe	i) heat stress risk related to physical activity; ii) heat-related mortality
Watts et al. (2017) ¹²⁰	Worldwide	Review of scientific evidence and analysis of indicators worldwide	i) incidence of climate-sensitive infectious diseases; ii) incidence of food-borne diseases, vector-borne diseases, parasitic diseases or zoonotic diseases; iii) mortality attributed to heat.
Watts et al. (2018) ^{19,a}	Worldwide	Review of scientific evidence and analysis of indicators worldwide	i) number of deaths associated with weather-related disasters; ii) number of deaths due to: all causes, dengue, diarrhoeal disease, heat and cold exposure, malaria, malignant melanoma, and protein-energy malnutrition; iii) number of undernourished people

(Table 2 continues on next page)

Reference	Setting & population	Brief methods	Main findings
(Continued from previous page)			
Watts et al. (2019) ^{97,a}	Worldwide	Review of scientific evidence and analysis of indicators worldwide	i) number of deaths associated with weather-related disasters; ii) number of deaths due to: all causes, dengue, diarrhoeal disease, forces of nature, heat and cold exposure, malaria, malignant melanoma, and protein-energy malnutrition; iii) number of undernourished people
Watts et al. (2020) ^{108,a}	Worldwide	Review of scientific evidence and analysis of indicators worldwide	i) heat-related excess mortality; ii) number of deaths associated with weather-related disasters
Zhang et al. (2018) ²²	Australia	Review of scientific evidence and analysis of indicators in Australia	i) daily non-accidental deaths associated with temperature; ii) daily non-accidental deaths associated with heatwaves; iii) trends in overall burden of climate-sensitive diseases (Ross River, Barmah Forest virus, dengue incidence, and salmonella cases); iv) prevalence of malnutrition; v) suicide rates associated with mean annual maximum temperatures
Zhang et al. (2023) ^{130,b}	China	Review of scientific evidence and analysis of indicators in China	i) number of hours available for safe physical activity per day; ii) heatwave-related mortality

^aFrom reference lists and updated searches in 2022. ^bFrom updated searches in 2024.

Table 2: Summary of study characteristics and main findings related to indicators from literature.

Reference	Setting & population	Main findings
CDC (2021) ¹⁰¹	United States, general population	i) heat-related emergency department visits for heat stress; ii) heat-related hospitalisations for heat stress; iii) heat-related mortality
Climate-ADAPT (2021) ¹⁰³	Europe, general population	i) deaths related to flooding; ii) cases of vibriosis infections, salmonellosis, campylobacteriosis, cryptosporidiosis; iii) number of heat-related deaths.
UKHSA (2023) ^{115,a}	United Kingdom, general population	Heatwaves and heat risk to health: annual heat-related mortality; annual heat illness; health impacts of wildfires Cold and cold risks to health: i) annual cold-related mortality and morbidity Flooding and flood risks to health: i) death or injury from flood events; ii) estimated number of people suffering flood-related adverse mental health impacts Vector-borne disease: i) number (rate) of Lyme disease cases; ii) autochthonous cases of vector-borne disease Food systems and health impacts: i) incidence of foodborne diseases
United Nations (2017) ¹⁰⁰	Worldwide, general population	Indicator related to Sustainable Development Goal 13 (Take urgent action to combat climate change and its impacts): Number of deaths, missing persons, and directly affected persons attributed to disasters per 100,000 populations.
United Nations Statistics Division (2022) ¹⁰²	Worldwide, general population	i) incidence of cases of climate-related disease: airborne diseases and water-related diseases; ii) incidence of climate-related vector-borne diseases: Lyme disease, malaria, West Nile virus, yellow fever, and dengue; iii) incidence of heat- and cold-related illnesses or excess mortality: excess mortality related to heat, excess mortality related to cold
US EPA (2016) ⁹⁹	United States, general population	i) heat-related deaths; ii) heat-related illnesses; iii) cold-related deaths; iv) cases of Lyme disease; iv) cases of West Nile virus
World Health Organization (2022) ¹⁰⁴	Worldwide, general population	i) climate change attributable deaths; ii) climate change attributable DALYs; iii) climate change attributable deaths in children under 5 years; iv) climate change attributable DALYs in children under 5 years

^aFrom updated searches in 2024.

Table 3: Summary of reports characteristics and main findings related to indicators from grey literature.

consultations was vaguely explained in some articles, limiting their quality and guiding role for other studies.

Despite of these challenges, several indicators related to morbidity and mortality outcomes were identified. Some articles clearly identified and defined indicators associated with specific climate-related hazards and exposures; however, the level of complexity in the exposure-response analysis and risk attribution varied. Based on the historical development of PHS systems, it is expected that the most common indicators are related to infectious and communicable diseases, which have been well-defined, and international and national supporting structures are relatively well-established.^{10,14,147,148} The next group of relatively well-developed indicators – although not exempt from challenges– covers chronic

non-communicable diseases and those related with occupational health.^{149,150} However, the use of indicators related to mental health is less straightforward¹⁵¹ mainly due to the challenges associated with population measurement and data collection,^{152,153} and potential underdiagnosis of mental health issues in some populations (e.g., anxiety, depression, and post-traumatic stress disorder),¹⁵⁴ which affects non-modelled and modelled indicators. For example, suicide rates associated with ambient temperature,⁹⁸ although possible of modelling, are sometimes less reliable due to underreporting and misclassification of the cause of death.^{155–158} Therefore, analysing the relationship with climate change can be complex, particularly when estimating the fraction of these outcomes attributable to climate change.¹⁵⁹

Exposure ^a	Morbidity, mortality, wellbeing, other	Specific diseases or causes	Indicator	
No specific exposure	Morbidity	No specific disease or cause	Morbidity rate ^{119,133}	
		Accidents	Rates of hospitalisations ¹²⁶	
		Cancer	Skin cancers ³⁶ Cancer rates ¹¹⁹	
		Circulatory	Rates of hospitalisations ¹²⁶	
		Nutrition	Number of undernourished people ^{19,97} Prevalence of malnutrition ²²	
		Poisoning	Rates of hospitalisations ¹²⁶	
		Respiratory	Rates of hospitalisations ¹²⁶ Asthma hospitalisations ¹¹⁶ Allergic disease-related hospitalisations ¹¹⁶ Asthma emergency room visits ¹¹⁶ Allergic disease-related emergency room visits ¹¹⁶	
		Violence	Rates of hospitalisations ¹²⁶	
		Mortality	No specific disease or cause	Number of deaths ^{36,38,97,119,126,127,133} Premature deaths ¹²⁶
			Accidents	Number of deaths ¹²⁶
	Cancer		Number of deaths due to malignant melanoma ^{19,97}	
	Non-accidental		Number of deaths ³⁸	
	Nutrition		Undernutrition mortality rates ¹²¹ Malnutrition mortality rates ^{19,97}	
	Other	Poisoning	Number of deaths ¹²⁶	
		Respiratory	Asthma ¹¹⁶	
Violence		Number of deaths ¹²⁶		
No specific disease or cause		Number of calls to emergency services ¹²⁷ Number of visits to general practitioner (GP) ¹²⁷ Number of hours available for safe physical activity per day ¹⁰⁷ Disability-adjusted life years ¹¹⁹		
Mental health		Utilisation of mental health services ¹²⁵		
Air pollution (i.e., ozone, particulate matter) and pollens	Morbidity	No specific disease or cause	Morbidity rate ¹³⁴	
		Respiratory/allergic	Cases of respiratory diseases ^{119,123,129,131} Cases of allergic diseases ^{124,129} Number of asthmatic episodes ¹²¹ Cases of chronic respiratory diseases ¹³¹	
		No specific disease or cause	Premature deaths ³⁸	
	Mortality	No specific disease or cause	Mortality rate ¹³⁴	
		Respiratory	Respiratory mortality ^{119,121,123,129,131,134}	
		Mental health	Reports of depression, anxiety related to climate change ¹²⁹	
Climate change	Morbidity	Mental health	Reports of depression, anxiety related to climate change ¹²⁹	
	Mortality	No specific disease or cause	Attributable deaths ¹⁰⁴ Preventable deaths ³⁸	
	Other	No specific disease or cause	Attributable disability-adjusted life years (DALYs) ^{38,104,121}	
Climate sensitive pathogens	Morbidity	Air-borne diseases	Airborne diseases ¹⁰²	
		Food-borne and water-borne diseases	Incidence of food-borne diseases ^{115,120} Incidence of water-borne diseases ¹⁰² Incidence of outbreaks of water-borne diseases ^{131,134} Shellfish poisoning outbreaks ¹²⁵ Shellfish poisoning cases ^{119,124} Cases of vibriosis infections ^{103,116} Cases of salmonellosis ^{103,134}	

(Table 4 continues on next page)

Exposure ^a	Morbidity, mortality, wellbeing, other	Specific diseases or causes	Indicator
(Continued from previous page)			
			Cases of campylobacteriosis ^{103,134} Cases of cryptosporidiosis ¹⁰³
		Gastrointestinal: diarrhoeal diseases	Number of diarrhoeal cases ¹³¹
		Infectious diseases	Recurrence rate of outbreaks of diarrhoeal disease ¹³¹ Incidence of climate-sensitive infectious diseases ^{119-121,123,129}
		Insect-borne diseases	Prevalence of insect-borne diseases ¹³¹
		Mosquito-borne diseases	Leishmaniasis cases ¹³⁴ Dengue incidence ^{102,105,134} Chikungunya cases ¹³⁴ Zika cases ¹³⁴ Ross River virus cases ¹⁰⁵ West Nile disease incidence ^{38,99,102,116} Malaria cases ^{102,134,136} Yellow fever cases ¹⁰²
	Mortality	Parasitic diseases	Parasitic diseases ¹²⁰
		Tick-borne diseases	Lyme disease cases ^{38,99,102,115}
		Vector-borne diseases	Vector-borne diseases ^{115,120}
		Zoonotic diseases	Zoonotic diseases ¹²⁰
		Gastrointestinal: diarrhoeal diseases	Mortality rate diarrhoeal ^{19,131}
		Insect-borne diseases	Mortality rate of children ¹³¹
		Mosquito-borne diseases	Mortality due to dengue ^{19,97} Mortality due to malaria ^{19,97}
Climatic variables	Morbidity	No specific disease or cause	Incidence of climate-related disease ¹⁰²
Extreme ambient temperatures (i.e., heat stress, heatwaves, cold waves)	Morbidity	No specific disease or cause	Rates of diseases ^{99,115,125,127,135}
			Excess morbidity ¹²³
			Hospitalisations ^{101,134}
		Cardiovascular	Rates of diseases ¹⁰⁵
		Heatstroke	Number of deaths ¹³⁴
		Injuries	Number of injuries ¹²⁴
	Mortality	Pregnancy	Births ¹³⁴ Premature births ¹³⁴ Low-weight births ¹³⁴
		Respiratory	Rates of diseases ¹⁰⁵
		No specific disease or cause	Mortality rate ^{1,2,4,19,97,99,101,103,105,106,115,118,120,124,125,127,128,130,132,134,135}
			Excess mortality ^{38,102,107,108,121,123,132}
		Heatstroke	Number of cases ¹³⁴
		Mental health	Suicide rate ^{22,98,122,134} Homicide rate ¹³⁴
Wellbeing	Non-accidental	Mortality rate ²²	
	No specific disease or cause	General sentiment of expressions ^{1,4,107}	
Other		Heat stress risk of people participating in physical and sporting activities ^{1,4,105,117,118,128,130}	
		Emergency department visits ¹⁰¹	
Extreme weather events	Morbidity	No specific disease or cause	Morbidity ^{121,125}
	Mortality	Injuries	Number of injuries ^{123,129}
		No specific disease or cause	Mortality ^{19,97,107,108,121,123,125}

(Table 4 continues on next page)

Exposure ^a	Morbidity, mortality, wellbeing, other	Specific diseases or causes	Indicator
(Continued from previous page)			
	Other	No specific disease or cause	Number of people affected ¹⁰⁷
Flooding	Morbidity	Injuries Mental health	Number of injuries ¹¹⁵ Illnesses ¹¹⁵
	Mortality	No specific disease or cause	Number of deaths ^{103,115}
Sea ice instability	Morbidity	Injuries	Number of injuries ¹²⁹
	Mortality	No specific disease or cause	Mortality ¹²⁹
Wildfires	Morbidity	No specific disease or cause	Illnesses ^{115,121}
		Respiratory	Asthma exacerbations ¹²⁵ Hospitalisations ¹²⁵ Injuries ¹²¹
	Mortality	No specific disease or cause	Number of deaths ¹²¹
Other	Mortality	No specific disease or cause	Number of deaths attributed to disasters per 100,000 population ¹⁰⁰
	Other	No specific disease or cause	Food security ¹²⁴ Number missing persons and directly affected persons attributed to disasters per 100,000 population ¹⁰⁰ Water security ¹²⁴

^aNo specific exposure" category included indicators that were identified as being associated with climate change; however, no specific climate-related exposure category was implicitly established.

Table 4: Indicators of morbidity, mortality, wellbeing, or other by exposure and causes.

Additionally, the potential use of indicators by national and sub-national agencies varies depending on the technical capacities at these levels. Most of the articles proposed general and non-modelled population health indicators, such as all-cause mortality or incidence of respiratory diseases.^{36,125,129,131,133} Although these indicators are relatively simple to measure and are already part of PHS systems, they need more advanced analyses in order to establish an association with climate and climate change. The reasons lie in that they are highly influenced by other social factors; for example, malaria is a parasitic disease transmitted by mosquitoes of the genus *Anopheles*, which are affected by climatic conditions and conditions in the built environment that are modified by human actions.¹⁶⁰ Similarly, food-borne diseases, such as salmonellosis, depend on multiple factors, including ambient temperature and human manipulation, such as the food chain quality.¹⁶¹ Overall, these simple indicators may alert about population health issues; however, the link with climate-related hazards and climate change is less straightforward. To provide more targeted information on climate-sensitive health outcomes, it is usually necessary to consider multiple climatic and non-climatic factors to isolate the associations of primary interest. This is often done using statistical models leading to model-based indicators, where the quality of the modelling depends on the quality and management of the data, as well as the

modelling assumptions. Additionally, model-based indicators might be more complex to analyse depending on the technical capacities at local levels.

One of the last areas to note is that all indicators in this review correspond to health outcomes related to the exposure to climate-related hazards, not directly to climate change. Although climate change affects climate-related hazards, it is important to differentiate their temporal scope. The assessment of health outcomes and climate change generally needs long-term analyses to capture long-term changes in the climate (i.e., climate signal)¹⁶² and thus capture the effect of that climate signal on health outcomes. Complementary, it is important to consider the different drivers of climate-related hazards, including natural or anthropogenic drivers, as well as other non-climatic hazards. For example, there are several articles that included health outcomes associated with air pollution.^{121,129} Although changes in the climate closely affect air quality and *vice versa*, most of the pollution (e.g., PM_{2.5}) comes from anthropogenic sources, such as wood burning, transport, and industry production.¹⁶³ It is important to identify where the health hazard comes in order to inform cost-effective interventions that act on the primary driver. Another example includes the analysis of population health outcomes associated with El Niño-Southern Oscillation (ENSO).⁹² Although this natural phenomenon affect ecosystems (e.g., making them

Broad health outcome	Incidence and Mortality of (unless otherwise stated)	Observations
Vector-borne diseases and health outcomes	<p>Malaria</p> <p>Zika virus disease</p> <p>West Nile virus disease</p> <p>Yellow fever</p> <p>Dengue fever</p> <p>Chikungunya fever</p> <p>Leishmaniasis</p> <p>Schistosomiasis</p> <p>Onchocerciasis</p> <p>Lyme disease</p> <p>Chagas disease</p> <p>Hantavirus pulmonary syndrome</p>	<ul style="list-style-type: none"> • These indicators are part of the PHS systems in Chile • Historically, mosquitoes that transmit these diseases have not been present in (mainland) Chile. However, given their presence in the region and potential changes in ecological niches due to climate change, it is relevant to include them into PHS. • Although some of these indicators did not explicitly appear in the systematic review, they are part of the PHS systems in Chile. • <i>Triatoma infestans</i>, the vector of <i>Trypanosoma cruzi</i> that causes chagas disease, is climate-sensitive; therefore, it is relevant to be integrated into PHS. • <i>Oligoryzomys longicaudatus</i> (i.e., long-tailed rat “colilargo”), the vector of Hantavirus that causes hantavirus pulmonary syndrome, may be climate-sensitive; therefore, it is relevant to be integrated into PHS.
Food- and water-borne diseases and health outcomes	<p>Diarrhoea</p> <p>Vibriosis</p> <p>Salmonellosis</p> <p>Campylobacteriosis</p> <p>Cryptosporidiosis</p> <p>Shellfish poisoning</p> <p>Harmful algal blooms</p> <p>Acute gastrointestinal illnesses</p> <p>Cholera</p> <p>Hepatitis A and C</p> <p>Brucellosis</p> <p>Leptospirosis</p> <p>Typhoid fever</p> <p><i>Escherichia coli</i> infections</p>	<ul style="list-style-type: none"> • These indicators are part of the PHS systems in Chile. • The indicators on specific pathogens include all cases independently of the causing microorganisms (e.g., <i>Vibrio</i> spp, <i>Salmonella</i> spp.). • Although these indicators did not explicitly appear in the systematic review, they are part of the PHS systems in Chile. • Diseases in these indicators are climate-sensitive; therefore, they are relevant for PHS
Temperature-related health outcomes (including heat and cold)	<p>Temperature-related diseases</p> <p>Number (or incidence) of emergency department visits due to temperature-related diseases</p> <p>Number (or incidence) of hospitalisations due to temperature-related diseases</p> <p>Number (or incidence) of emergency department visits associated with extreme temperatures</p> <p>Number (or incidence) of hospitalisations associated with extreme temperatures</p> <p>Mortality associated with extreme temperatures</p> <p>Excess of emergency department visits due to temperature-related causes</p> <p>Excess of hospitalisations due to temperature-related causes</p> <p>Excess mortality due to temperature-related causes</p> <p>Cardiovascular diseases associated with temperature-related causes</p> <p>Respiratory diseases associated with temperature-related causes</p>	<ul style="list-style-type: none"> • Health data for these indicators can be obtained from databases that are compiled and published by the Department of Statistics and Health Information (DEIS). • Environmental data for these indicators can be obtained from reanalysis datasets. • These indicators are based on the International Classification of Disease (ICD); therefore, they should be used with caution due to potential underdiagnosis. • Health data for these indicators can be obtained from databases that are compiled and published by the Department of Statistics and Health Information (DEIS). • Environmental data for these indicators can be obtained from reanalysis datasets. • Extreme temperatures may be defined locally based on percentiles or another climate index, such as warm days, heatwaves, or cold spell. • These indicators need further analyses based on statistical models; however, data are already available.

Table 5: Suggested high-priority indicators for PHS in Chile.

prone to mosquitoes), the evidence still shows uncertainty between ENSO and climate change^{164,165}; therefore, indicators analysed in the context of ENSO should be carefully analysed.

The selection and analysis of indicators that can be integrated into PHS systems involve several steps and requirements that need complementary scientific and policy perspectives. As it was shown in this review, it was not necessary to create new indicators as many are already proposed; however, their direct application to local contexts is complex and should be discussed with decision-makers at the corresponding levels. In this sense, the guide presented in this study informed the identification of indicators relevant for the Chilean context, resulting in three sets of prioritised indicators that can be considered for PHS under a climate change and health framework. Nonetheless, all indicators are related to the most severe impacts of climate-related hazards, leaving aside potential less severe outcomes but that affect a higher proportion of the population. Fig. 2 shows the outcome/data iceberg for indicators on climate change and population health in Chile. All elements that are above the horizontal line represent measurable outcomes/data; however, currently only some of them are reliably and systematically collected and available for analysis (i.e., those marked with **) while others need careful analysis if used (i.e., those marked with *). Up to date, elements below the horizontal line (i.e., unknown and unmeasured) may encompass a high proportion of the impacts of climate change and climate-related hazards on population health, but it is not yet possible to understand their actual magnitude due to limitations related to measurement, data collection, and reporting/diagnosis.

In this regard, it would be desirable to integrate other indicators that track upstream climate-related exposures and outcomes, which would allow anticipated and preventive public health measures. The DPSEEA (Driving, Pressure, State, Exposure, Effect, and Action) or the MEME (Multiple Exposures-Multiple Effects) frameworks may help identify and organise indicators that inform public health decision-making according to different stages of exposures and outcomes in the population, including intermediate indicators (i.e., indicators on changes in the environment) and indicators of health impacts.^{43,166} These frameworks may be complemented by others, such as the social determinants of health framework,¹⁶⁷ which can expand the perspective of social vulnerabilities to climate change and climate-related hazards, including indicators from other health-determining sectors (e.g., urban planning, energy, transport, and housing) and different levels of disaggregation. The rationale behind the use of complementary frameworks relies on the issue that several public agencies, not only public health agencies, are already collecting data and evaluating indicators that can be integrated into PHS systems. All these indicators may be evaluated with the suitability guide presented in this study to facilitate the selection for local PHS systems, which may maximise resources and strengthen climate adaptation and population health policies.

This study has several strengths. First, it is one of the first that systematically searched and assessed scientific and grey literature in relation to population health indicators related to the impacts of climate change and climate-related hazards for PHS. As it was shown, several literature reviews proposed indicators based on expert opinion rather than a more comprehensive




Fig. 2: Data iceberg for indicators of impacts of climate change on population health in Chile. NB: ** valid and reliable sources of data; * sources with limited use. Source: Own elaboration.

analysis. Second, the systematic use of two critical appraisal tools (i.e., QuADS and JBI) allowed a thorough evaluation of the evidence with different designs of studies. Third, the suitability analysis provided a comprehensive and systematic evaluation of indicators according to a national scenario, in this case Chile. This approach, in contrast to several indicators developed with limited application to PHS systems, maximises the probability of evidence translation into public policy.

This work supports public health surveillance systems in Chile, and potentially other Latin American countries, helping decision- and policymaking processes monitor the impacts of climate-related hazards on population health in a changing climate by analysing the proposed high-priority indicators, and adapt a set of indicators that can be used at local levels by implementing the suitability analysis guide. Additionally, the findings of this study also contribute to the two-year UAE–Belém work programme, which was established at COP28 and aims at developing indicators for measuring progress of targets outlined in the UEA Framework for Global Climate Resilience, including health and climate change-related impacts.⁷

Nonetheless, this study has also several limitations. First, the systematic review only included three languages relevant to the Chilean context (and other Spanish and Portuguese speaking contexts) and it is possible that a proportion of the literature was not included. Additionally, the quality of the articles varied and showed that current evidence lacks methodological systematicity when analysing and proposing indicators. This, together with the overrepresentation of evidence coming from High-Income settings, may introduce some biases in the conclusions, affecting the uptake of this evidence by end-users at different levels. Second, the tool for the suitability analysis needs further developments, including a validation of it and the results obtained by its application. The dimensions, sub-dimensions, and criteria are potentially subjective; therefore, it would be desirable to perform a co-construction process (e.g., analyses based on Delphi method) with other interested parties (e.g., experts, decision- and policymakers, researchers, public health practitioners) to determine their robustness, acceptability, and feasibility. Nonetheless, the two interviews with officials at the Ministry of Health and Ministry of the Environment in Chile revealed that the tool captures most of the relevant criteria and is useful for public health decision-making. Finally, the criteria for validity, reliability, sensitivity, specificity, and comparability from the suitability analysis need to be further evaluated using actual data. In this case, these criteria were only evaluated based on theoretical assumptions, which may be different when applying different datasets.

Finally, this study is not intended to be definitive and invites all public health experts to continue developing and discussing the evidence presented here. It is

imperative that PHS systems are adaptive and dynamic because changes in the climate will continue as the 2023 and 2024 have already demonstrated.

Conclusion

There are several population health indicators that help monitor the health impacts of climate-related hazards in the context of climate change; however, local analyses are needed based on scientific foundations, measurability, relevance to public health, and contextual appropriateness. The guide for suitability analysis in this article seems useful for determining which indicator can be used in a local context for tracking the impacts of climate change and climate change-related hazards on population health. The process and information presented here may contribute to local public health surveillance systems and the two-year UAE–Belém work programme.

Contributors

YPS conceptualised the idea, wrote the protocol, planned and carried out the review, led the project, interpreted and prepared the results for publication, trained researchers, wrote the draft of the manuscript, and wrote the final manuscript. RAC, JIG, and RGT contributed to the screening and appraisal of references and reviewed the final manuscript. IK gave feedback during the review, writing process, and final manuscript. All authors agreed on the final manuscript and the decision to submit the study for publication. YPS was responsible for submitting it to the journal.

Declaration of interests

YPS was funded by the National Agency for Research and Development (ANID)/Scholarship Program/DOCTORADO BECAS CHILE/2018–72190338. YPS thanks University College London for the economic support in the publication process. The authors declare no conflict of interests.

Acknowledgements

We thank Richard E. Chandler for his support and feedback throughout the research process; and Julio Monreal and Priscilla Ulloa for their time and important feedback and suggestions on the suitability tool.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.lana.2024.100854>.

References

- Romanello M, Napoli CD, Drummond P, et al. The 2022 report of the Lancet Countdown on health and climate change: health at the mercy of fossil fuels. *Lancet*. 2022. [https://doi.org/10.1016/S0140-6736\(22\)01540-9](https://doi.org/10.1016/S0140-6736(22)01540-9). published online Oct 25.
- Hartinger SM, Yglesias-González M, Blanco-Villafuerte L, et al. The 2022 South America report of the Lancet Countdown on health and climate change: trust the science. Now that we know, we must act. *Lancet Reg Health Am*. 2023. <https://doi.org/10.1016/j.lana.2023.100470>. published online Mar 28.
- World Meteorological Organization. *State of the global climate 2023*. Geneva, Switzerland: WMO; 2024. <https://wmo.int/publication-series/state-of-global-climate-2023>. Accessed May 16, 2024.
- Romanello M, Napoli C di, Green C, et al. The 2023 report of the Lancet Countdown on health and climate change: the imperative for a health-centred response in a world facing irreversible harms. *Lancet*. 2023;402:2346–2394.
- IPCC. *Climate change 2021: the physical science basis. Contribution of working group I to the sixth assessment report of the intergovernmental panel on climate change*. Cambridge, United Kingdom and New

- York, NY, USA: Cambridge University Press; 2021. <https://doi.org/10.1017/9781009157896>.
- 6 IPCC. *Climate change 2022: impacts, adaptation, and vulnerability. Contribution of working group II to the sixth assessment report of the intergovernmental panel on climate change*. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press; 2022. <https://doi.org/10.1017/9781009325844>.
 - 7 UNFCCC. *Glasgow–Sharm el-Sheikh work programme on the global goal on adaptation referred to in decision 7/CMA.3*; 2023. https://unfccc.int/sites/default/files/resource/cma5_auv_8a_gga.pdf. Accessed April 8, 2024.
 - 8 Pan American Health Organization. *The essential public health functions in the Americas: a renewal for the 21st century. Conceptual framework and description*. Washington, D.C.; 2020. <https://iris.paho.org/handle/10665.2/53124>. Accessed November 2, 2022.
 - 9 Smith PF, Hadler JL, Stanbury M, Rolfs RT, Hopkins RS, CSTE Surveillance Strategy Group. 'Blueprint version 2.0': updating public health surveillance for the 21st century. *J Public Health Manag Pract JPHMP*. 2013;19:231–239.
 - 10 Thacker SB, Stroup DF. Origins and progress in surveillance systems. In: *Infectious disease surveillance*. John Wiley & Sons, Ltd; 2013:21–31.
 - 11 Detels R, Karim Q, Baum F, Li L, Leyland A, eds. *Oxford textbook of global public health*. 7th ed. Oxford, United Kingdom: Oxford University Press; 2021. <https://doi.org/10.1093/med/9780198816805.001.0001>.
 - 12 Groseclose SL, Buckeridge DL. Public health surveillance systems: recent advances in their use and evaluation. *Annu Rev Public Health*. 2017;38:57–79.
 - 13 Lash T, VanderWeele T, Haneuse S, Rothman K. *Modern epidemiology*. 4th ed. Lippincott Williams and Wilkins; 2021.
 - 14 World Health Organization. *International health regulations*. 3rd ed. WHO; 2016.
 - 15 Joas A, Schöpel M, David M, et al. Environmental health surveillance in a future European health information system. *Arch Public Health*. 2018;76:27. <https://doi.org/10.1186/s13690-018-0272-6>.
 - 16 Briggs D. *Making a difference: indicators to improve children's environmental health*. Geneva, Switzerland: WHO; 2003. <https://apps.who.int/iris/handle/10665/43721>. Accessed March 2, 2023.
 - 17 Waheed B, Khan F, Veitch B. Linkage-based frameworks for sustainability assessment: making a case for driving force-pressure-state-exposure-effect-action (DPSEEA) frameworks. *Sustainability*. 2009;1:441–463.
 - 18 Sly JL, Moore SE, Gore F, et al. Children's environmental health indicators in Australia. *Ann Glob Health*. 2016;82:156–168.
 - 19 Watts N, Amann M, Ayeb-Karlsson S, et al. The Lancet Countdown on health and climate change: from 25 years of inaction to a global transformation for public health. *Lancet*. 2018;391:581–630. [https://doi.org/10.1016/S0140-6736\(17\)32464-9](https://doi.org/10.1016/S0140-6736(17)32464-9).
 - 20 UK Health Security Agency. Environmental public health surveillance system (EPHSS). Report for 2020 to 2021. <https://www.gov.uk/government/publications/environmental-public-health-surveillance-system>; 2022. Accessed March 3, 2023.
 - 21 Watts N, Adger WN, Agnolucci P, et al. Health and climate change: policy responses to protect public health. *Lancet*. 2015;386:1861–1914.
 - 22 Zhang Y, Beggs PJ, Bambrick H, et al. The MJA-Lancet Countdown on health and climate change: Australian policy inaction threatens lives. *Med J Aust*. 2018;209:474. <https://doi.org/10.5694/mja18.00789>.
 - 23 Environmental health intelligence New Zealand. *Environ Health Indic Clim Change*; 2022. <http://www.ehinz.ac.nz/indicators/climate-change/>. Accessed March 7, 2023.
 - 24 Ffoulkes C, Hockridge B, Illman H, Holmes G, Manning F, Wilson L. *Research to review and update indicators of climate-related risks and actions in England*. England, UK: CCC; 2021. <https://www.theccc.org.uk/wp-content/uploads/2021/06/Research-to-review-and-update-indicators-of-climate-related-risks-and-actions-in-England-ADAS.pdf>. Accessed March 7, 2023.
 - 25 Centres for Disease Control and Prevention. Climate change | tracking | NCEH | CDC. *Clim Change*; 2020. <https://www.cdc.gov/nceh/tracking/topics/ClimateChange.htm>. Accessed March 7, 2023.
 - 26 Palmeiro-Silva Y, Kelman I. A systematic review of population health indicators that monitor the health effects of climate change. *PROSPERO 2021 CRD42021253704*; 2021. https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42021253704. Accessed August 23, 2022.
 - 27 Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71. <https://doi.org/10.1136/bmj.n71>.
 - 28 Covidence. Covidence systematic review software. www.covidence.org; 2022. Accessed January 13, 2021.
 - 29 Harrison R, Jones B, Gardner P, Lawton R. Quality assessment with diverse studies (QuADS): an appraisal tool for methodological and reporting quality in systematic reviews of mixed- or multi-method studies. *BMC Health Serv Res*. 2021;21:144. <https://doi.org/10.1186/s12913-021-06122-y>.
 - 30 JBI. *Critical appraisal tools | JBI. Crit Apprais Tools*; 2020. <https://jbi.github.io/critical-appraisal-tools>. Accessed May 1, 2021.
 - 31 Hsieh H-F, Shannon SE. Three approaches to qualitative content analysis. *Qual Health Res*. 2005;15:1277–1288.
 - 32 Wilson V. Research methods: content analysis. *Evid Based Libr Inf Pract*. 2016;11:41–43.
 - 33 Validity, measurement. In: Porta M, ed. *A dictionary of epidemiology*. 6th ed. New York, USA: Oxford University Press; 2016. <https://www.oxfordreference.com/display/10.1093/acref/9780199976720.001.0001/acref-9780199976720-e-1930>. Accessed March 27, 2023.
 - 34 Reliability. In: Porta M, ed. *A dictionary of epidemiology*. 6th ed. Oxford University Press; 2016. <https://www.oxfordreference.com/display/10.1093/acref/9780199976720.001.0001/acref-9780199976720-e-1630>. Accessed March 27, 2023.
 - 35 PAHO Special Program for health Analysis. Health indicators: building blocks for health analysis. *Epidemiol Bull*. 2001;22. https://www.paho.org/english/sha/be_v22n4-indicators.htm.
 - 36 Donnelly A, Jones MB, Sweeney J. A review of indicators of climate change for use in Ireland. *Int J Biometeorol*. 2004;49:1–12.
 - 37 de Vet HCW, Terwee CB, Mokkink LB, Knol DL. *Measurement in medicine: a practical guide*. Cambridge: Cambridge University Press; 2011. <https://doi.org/10.1017/CBO9780511996214>.
 - 38 Cheng JJ, Berry P. Development of key indicators to quantify the health impacts of climate change on Canadians. *Int J Public Health*. 2013;58:765–775.
 - 39 Health indicator. In: Porta M, ed. *A dictionary of epidemiology*. 6th ed. New York, USA: Oxford University Press; 2016. <https://www.oxfordreference.com/display/10.1093/acref/9780199976720.001.0001/acref-9780199976720-e-904>. Accessed February 23, 2023.
 - 40 Pan American Health Organization. *Health Indicators. Conceptual and operational considerations*. Washington, D.C.: PAHO; 2018. <https://iris.paho.org/handle/10665.2/49056>. Accessed March 6, 2023.
 - 41 Pencheon D. *The good indicators guide: understanding how to use and choose indicators*. England, UK: NHS England; 2007. <https://www.england.nhs.uk/improvement-hub/publication/the-good-indicators-guide-understanding-how-to-use-and-choose-indicators/>. Accessed March 6, 2023.
 - 42 Larson C, Mercer A. Global health indicators: an overview. *CMAJ Can Med Assoc J*. 2004;171:1199–1200.
 - 43 Briggs D. *Environmental health indicators: framework and methodologies*. Geneva, Switzerland: WHO; 1999. <https://www.who.int/publications/i/item/WHO-SDE-OEH-99.10>. Accessed March 8, 2023.
 - 44 Institute of Medicine (US) Committee on Leading Health Indicators for Healthy People 2010. *Leading Health Indicators for Healthy People 2010: Final Report*. Washington (DC): National Academies Press (US); 1999. <http://www.ncbi.nlm.nih.gov/books/NBK231607/>. Accessed March 6, 2023.
 - 45 Kovats RS, Haines A, Stanwell-Smith R, Martens P, Menne B, Bertollini R. Climate change and human health in Europe. *BMJ*. 1999;318:1682–1685.
 - 46 Warren JA, Berner JE, Curtis T. Climate change and human health: infrastructure impacts to small remote communities in the north. *Int J Circumpolar Health*. 2005;64:487–497.
 - 47 Costello A, Abbas M, Allen A, et al. Managing the health effects of climate change: Lancet and university College London Institute for global health commission. *Lancet*. 2009;373:1693–1733. [https://doi.org/10.1016/S0140-6736\(09\)60935-1](https://doi.org/10.1016/S0140-6736(09)60935-1).
 - 48 Hambling T, Weinstein P, Slaney D. A review of frameworks for developing environmental health indicators for climate change and health. *Int J Environ Res Public Health*. 2011;8:2854–2875.
 - 49 Kan H. Climate change and human health in China. *Environ Health Perspect*. 2011;119:A60–A61.
 - 50 Toutant S, Gosselin P, Bélanger D, Bustintza R, Rivest S. An open source web application for the surveillance and prevention of the impacts on public health of extreme meteorological events: the

- SUPREME system. *Int J Health Geogr.* 2011;10:39. <https://doi.org/10.1186/1476-072X-10-39>.
- 51 Armstrong B, Hajat S, Kovats S, Lloyd S, Scovronick N, Wilkinson P. Climate change: how can epidemiology best inform policy? *Epidemiol Camb Mass.* 2012;23:780–784.
 - 52 Pascal M, Viso AC, Medina S, Delmas MC, Beaudreau P. How can a climate change perspective be integrated into public health surveillance? *Publ Health.* 2012;126:660–667.
 - 53 Bustinza R, Lebel G, Gosselin P, Bélanger D, Chebana F. Health impacts of the July 2010 heat wave in Québec, Canada. *BMC Public Health.* 2013;13:56. <https://doi.org/10.1186/1471-2458-13-56>.
 - 54 Nairn JR, Fawcett RJB. The excess heat factor: a metric for heatwave intensity and its use in classifying heatwave severity. *Int J Environ Res Public Health.* 2014;12:227–253.
 - 55 Upperman CR, Parker J, Jiang C, He X, Murtugudde R, Sapkota A. Frequency of extreme heat event as a surrogate exposure metric for examining the human health effects of climate change. *PLoS One.* 2015;10:e0144202. <https://doi.org/10.1371/journal.pone.0144202>.
 - 56 Weber S, Sadoff N, Zell E, de Sherbinin A. Policy-relevant indicators for mapping the vulnerability of urban populations to extreme heat events: a case study of Philadelphia. *Appl Geogr.* 2015;63:231–243.
 - 57 Barcellos C, Roux E, Ceccato P, et al. An observatory to gather and disseminate information on the health-related effects of environmental and climate change. *Rev Panam Salud Publica Pan Am J Public Health.* 2016;40:167–173.
 - 58 Ban J, Huang L, Chen C, Guo Y, He MZ, Li T. Integrating new indicators of predictors that shape the public's perception of local extreme temperature in China. *Sci Total Environ.* 2017;579:529–536.
 - 59 Navi M, Hansen A, Nitschke M, Hanson-Easey S, Pisanelli D. Developing health-related indicators of climate change: Australian stakeholder perspectives. *Int J Environ Res Public Health.* 2017;14:552. <https://doi.org/10.3390/ijerph14050552>.
 - 60 Alexander KA, Heaney AK, Shaman J. Hydrometeorology and flood pulse dynamics drive diarrheal disease outbreaks and increase vulnerability to climate change in surface-water-dependent populations: a retrospective analysis. *PLoS Med.* 2018;15:e1002688. <https://doi.org/10.1371/journal.pmed.1002688>.
 - 61 Chen Q, Ding M, Yang X, Hu K, Qi J. Spatially explicit assessment of heat health risk by using multi-sensor remote sensing images and socioeconomic data in Yangtze River Delta, China. *Int J Health Geogr.* 2018;17:15. <https://doi.org/10.1186/s12942-018-0135-y>.
 - 62 Gao C, Kuklane K, Östergren P-O, Kjellstrom T. Occupational heat stress assessment and protective strategies in the context of climate change. *Int J Biometeorol.* 2018;62:359–371.
 - 63 Jolliet O, Antón A, Boulay A-M, et al. Global guidance on environmental life cycle impact assessment indicators: impacts of climate change, fine particulate matter formation, water consumption and land use. *Int J Life Cycle Assess.* 2018;23:2189–2207.
 - 64 Nissan H, Conway D. From advocacy to action: projecting the health impacts of climate change. *PLoS Med.* 2018;15:e1002624. <https://doi.org/10.1371/journal.pmed.1002624>.
 - 65 Patz JA, Thomson MC. Climate change and health: moving from theory to practice. *PLoS Med.* 2018;15:e1002628. <https://doi.org/10.1371/journal.pmed.1002628>.
 - 66 Weyant C, Brandeau ML, Burke M, Lobell DB, Bendavid E, Basu S. Anticipated burden and mitigation of carbon-dioxide-induced nutritional deficiencies and related diseases: a simulation modeling study. *PLoS Med.* 2018;15:e1002586. <https://doi.org/10.1371/journal.pmed.1002586>.
 - 67 Oliveira BFAD, Silveira IH, Feitosa RC, Horta MAP, Junger WL, Hacon S. Human heat stress risk prediction in the Brazilian semiarid region based on the wet-bulb globe temperature. *An Acad Bras Cienc.* 2019;91:e20180748. <https://doi.org/10.1590/0001-3765201920180748>.
 - 68 Sadiq LS, Hashim Z, Osman M. The impact of heat on health and productivity among maize farmers in a tropical climate area. *J Environ Public Health.* 2019;2019:e9896410. <https://doi.org/10.1155/2019/9896410>.
 - 69 Valois P, Caron M, Gousse-Lessard A-S, Talbot D, Renaud J-S. Development and validation of five behavioral indices of flood adaptation. *BMC Public Health.* 2019;19:245. <https://doi.org/10.1186/s12889-019-6564-0>.
 - 70 Campbell AM, Racault M-F, Goult S, Laurenson A. Cholera risk: a machine learning approach applied to essential climate variables. *Int J Environ Res Public Health.* 2020;17:9378. <https://doi.org/10.3390/ijerph17249378>.
 - 71 Turek-Hankins LL, Hino M, Mach KJ. Risk screening methods for extreme heat: implications for equity-oriented adaptation. *PLoS One.* 2020;15:e0240841. <https://doi.org/10.1371/journal.pone.0240841>.
 - 72 Nathwani J, Lind N, Renn O, Schellhuber HJ. Balancing health, economy and climate risk in a multi-crisis. *Energies.* 2021;14:4067. <https://doi.org/10.3390/en14144067>.
 - 73 Ranadive N, Desai J, Sathish LM, et al. Climate change adaptation: prehospital data facilitate the detection of acute heat illness in India. *West J Emerg Med.* 2021;22:739–749.
 - 74 Wang Q, Zhang Y, Ban J, Zhu H, Xu H, Li T. The relationship between population heat vulnerability and urbanization levels: a county-level modeling study across China. *Environ Int.* 2021;156:106742. <https://doi.org/10.1016/j.envint.2021.106742>.
 - 75 Quincey PG. A relationship between black smoke index and black carbon concentration. *Atmos Env.* 2007;41:7964–7968.
 - 76 Liu G, Yang Z, Chen B, Ulgiati S. Monitoring trends of urban development and environmental impact of Beijing, 1999–2006. *Sci Total Environ.* 2011;409:3295–3308.
 - 77 Sunbury T, Driscoll D. Characteristics of a community-based sentinel surveillance system: lessons learned from toolkit development and implementation. *Int Public Health J.* 2013;5:105–114.
 - 78 Tseng C-H, Lu L-C, Lan S-H, Hsieh Y-P, Lan S-J. Relationship between emergency care utilization, ambient temperature, and the pollution standard index in Taiwan. *Int J Environ Health Res.* 2017;27:344–354.
 - 79 Blake R, Jacob K, Yohe G, et al. New York city panel on climate change 2019 report chapter 8: indicators and monitoring. *Ann N Y Acad Sci.* 2019;1439:230–279.
 - 80 Madden DL, McLean M, Brennan M, Moore A. Why use indicators to measure and monitor the inclusion of climate change and environmental sustainability in health professions' education? *Med Teach.* 2020;42:1119–1122.
 - 81 Yadav K, Geli HME, Cibils AF, et al. An integrated food, energy, and water nexus, human well-being, and resilience (FEW-WISE) framework: New Mexico. *Front Environ Sci.* 2021;9:667018. <https://www.frontiersin.org/articles/10.3389/fenvs.2021.667018>. Accessed August 30, 2022.
 - 82 Lam LT. The association between climatic factors and childhood illnesses presented to hospital emergency among young children. *Int J Environ Health Res.* 2007;17:1–8.
 - 83 Holmes SH, Phillips T, Wilson A. Overheating and passive habitability: indoor health and heat indices. *Build Res Inf.* 2016;44:1–19.
 - 84 Lal A, Hashizume M, Hales S. Indian ocean dipole and cryptosporidiosis in Australia: short-term and nonlinear associations. *Environ Sci Technol.* 2017;51:8119–8127.
 - 85 Andrianou XD, Makris KC. The framework of urban exposome: application of the exposome concept in urban health studies. *Sci Total Environ.* 2018;636:963–967.
 - 86 Green HK, Edgheere O, Elliot AJ, et al. Google search patterns monitoring the daily health impact of heatwaves in England: how do the findings compare to established syndromic surveillance systems from 2013 to 2017? *Environ Res.* 2018;166:707–712.
 - 87 Dushkova DO, Tikunov VS, Cheresnyia OYU. Methodology for assessment of public health at the municipalities level (A case study of arkhangel'sk oblast). *Geogr Nat Resour.* 2019;40:74–81.
 - 88 Valois P, Tessier M, Bouchard D, et al. Monitoring the evolution of individuals' flood-related adaptive behaviors over time: two cross-sectional surveys conducted in the Province of Quebec, Canada. *BMC Public Health.* 2020;20:1643. <https://doi.org/10.1186/s12889-020-09763-6>.
 - 89 Beauchamp E, Sainsbury NC, Greene S, Chaigneau T. Aligning resilience and wellbeing outcomes for locally-led adaptation in Tanzania. *Sustainability.* 2021;13:8976. <https://doi.org/10.3390/su13168976>.
 - 90 Di Maria F, Mastrantonio M, Uccelli R. The life cycle approach for assessing the impact of municipal solid waste incineration on the environment and on human health. *Sci Total Environ.* 2021;776:145785. <https://doi.org/10.1016/j.scitotenv.2021.145785>.
 - 91 Yin Y, Grundstein A, Mishra DR, Ramaswamy L, Hashemi Tonekaboni N, Dowd J. DTEX: a dynamic urban thermal exposure index based on human mobility patterns. *Environ Int.* 2021;155:106573. <https://doi.org/10.1016/j.envint.2021.106573>.
 - 92 Hurtado-Díaz M, Riojas-Rodríguez H, Rothenberg SJ, Gomez-Dantés H, Cifuentes E. Short communication: impact of climate variability on the incidence of dengue in Mexico. *Trop Med Int Health TM IH.* 2007;12:1327–1337.

- 93 Regens J, Hodges D. GIS and remote sensing applications to evaluate public health effects of global warming: world Resource Review. *World Resour Rev*. 2000;12:458–464.
- 94 Derne B, Fearnley E, Goater S, Carter K, Weinstein P. Ciguatera fish poisoning and environmental change: a case for strengthening health surveillance in the Pacific? *Pac Health Dialog*. 2010;16:99–108.
- 95 Simane B, Deressa W, Kumie A, et al. Health vulnerability and adaptation strategies to climate change in Ethiopia. In: *Climate change adaptation in africa*. Springer International Publishing; 2017.
- 96 Qian Y-J, Li S-Z, Xu J-F, Zhang L, Fu Q, Zhou X-N. [Establishment of policy indicators of adaptation to the impact of climate change on the transmission of schistosomiasis and malaria in China]. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi*. 2013;31:433–437.
- 97 Watts N, Amann M, Arnell N, et al. The 2019 report of the lancet countdown on health and climate change: ensuring that the health of a child born today is not defined by a changing climate. *Lancet*. 2019;394:1836–1878.
- 98 Beggs PJ, Zhang Y, Bambrick H, et al. The 2019 report of the MJA–Lancet Countdown on health and climate change: a turbulent year with mixed progress. *Med J Aust*. 2019;211. <https://www.mja.com.au/journal/2019/211/11/2019-report-mja-lancet-countdown-health-and-climate-change-turbulent-year-mixed>. Accessed August 30, 2022.
- 99 US EPA O. *Climate change indicators: health and society*; 2016. published online July 1 <https://www.epa.gov/climate-indicators/health-society>. Accessed August 30, 2022.
- 100 United Nations. *Resolution adopted by the general assembly on 6 July 2017. Work of the statistical commission pertaining to the 2030 agenda for sustainable development*. United Nations; 2017. https://ggim.un.org/documents/a_res_71_313.pdf. Accessed August 26, 2022.
- 101 CDC. Climate change | tracking | NCEH | CDC. *Natl Environ Public Health Track*; 2021. published online Aug 31 <https://www.cdc.gov/nceh/tracking/topics/ClimateChange.htm>. Accessed August 26, 2022.
- 102 United Nations Statistics Division. *Background document to the report of the secretary-general on climate change statistics (E/CN.3/2022/17)*; 2022. <https://unstats.un.org/unsd/statcom/53rd-session/documents/BG-3m-Globalsetandmetadata-E.pdf>. Accessed August 26, 2022.
- 103 Climate-ADAPT. Indicators on climate change and health — European climate and health observatory. *Indic Clim Change Health*; 2021. https://climate-adapt.eea.europa.eu/observatory/evidence/indicators_intro. Accessed August 26, 2022.
- 104 World Health Organization. The global health observatory - climate change. *Glob Health Obs Explore World Health Data*; 2022. <https://www.who.int/data/gho/data/themes/topics/topic-details/GHO/climate-change>. Accessed August 26, 2022.
- 105 Beggs PJ, Zhang Y, McGushin A, et al. The 2021 report of the MJA–Lancet Countdown on health and climate change: Australia increasingly out on a limb. *Med J Aust*. 2021;215:390–392.e22. <https://doi.org/10.5694/mja2.51302>.
- 106 Cai W, Zhang C, Suen HP, et al. The 2020 China report of the Lancet Countdown on health and climate change. *Lancet Public Health*. 2021;6:e64–e81. [https://doi.org/10.1016/S2468-2667\(20\)30256-5](https://doi.org/10.1016/S2468-2667(20)30256-5).
- 107 Romanello M, McGushin A, Napoli CD, et al. The 2021 report of the Lancet Countdown on health and climate change: code red for a healthy future. *Lancet*. 2021;398:1619–1662.
- 108 Watts N, Amann M, Arnell N, et al. The 2020 report of the Lancet Countdown on health and climate change: responding to converging crises. *Lancet*. 2021;397:129–170.
- 109 Romanello M, Daalen K van, Anto JM, et al. Tracking progress on health and climate change in Europe. *Lancet Public Health*. 2021;6:e858–e865. [https://doi.org/10.1016/S2468-2667\(21\)00207-3](https://doi.org/10.1016/S2468-2667(21)00207-3).
- 110 Requejo J, Strong K, Agweyu A, et al. Measuring and monitoring child health and wellbeing: recommendations for tracking progress with a core set of indicators in the Sustainable Development Goals era. *Lancet Child Adolesc Health*. 2022;6:345–352.
- 111 Sinno-Tellier S, Abadie E, Guillotin S, Bossée A, Nicolas M, Delcourt N. Human shellfish poisoning: implementation of a national surveillance program in France. *Front Mar Sci*. 2023;9:1089585. <https://doi.org/10.3389/fmars.2022.1089585>.
- 112 Barteit S, Colmar D, Nellis S, et al. Developing climate change and health impact monitoring with eHealth at the South east Asia community observatory and health and demographic surveillance site, Malaysia (CHIMES). *Front Public Health*. 2023;11:1153149. <https://doi.org/10.3389/fpubh.2023.1153149>.
- 113 Alfano R, De' Donato F, Vineis P, Romanello M. Lancet countdown indicators for Italy: tracking progress on climate change and health. *Epidemiol Prev*. 2023;47:6–21.
- 114 Di Napoli C, McGushin A, Romanello M, et al. Tracking the impacts of climate change on human health via indicators: lessons from the Lancet Countdown. *BMC Public Health*. 2022;22:663. <https://doi.org/10.1186/s12889-022-13055-6>.
- 115 UK Health Security Agency. Climate change and public health indicators: scoping review. <https://www.gov.uk/government/publications/climate-change-and-public-health-indicators-scoping-review>; 2023. Accessed June 16, 2024.
- 116 Liu AY, Trtanj JM, Lipp EK, Balbus JM. Toward an integrated system of climate change and human health indicators: a conceptual framework. *Clim Change*. 2021;166:49. <https://doi.org/10.1007/s10584-021-03125-w>.
- 117 Beggs PJ, Trueck S, Linnenluecke MK, et al. The 2023 report of the MJA–Lancet Countdown on health and climate change: sustainability needed in Australia's health care sector. *Med J Aust*; 2024. Online first <https://www.mja.com.au/journal/2024/220/6/2023-report-mja-lancet-countdown-health-and-climate-change-sustainability-needed>. Accessed June 16, 2024.
- 118 Hartinger SM, Palmeiro-Silva YK, Ilerena-Cayo C, et al. The 2023 Latin America report of the Lancet Countdown on health and climate change: the imperative for health-centred climate-resilient development. *Lancet Reg Health Am*. 2024;33. <https://doi.org/10.1016/j.lana.2024.100746>.
- 119 National Research Council. *Monitoring climate change impacts: metrics at the intersection of the human and Earth Systems*. Washington DC: The National Academy Press; 2010. <https://nap.nationalacademies.org/read/12965>. Accessed August 25, 2021.
- 120 Watts N, Adger WN, Ayeb-Karlsson S, et al. The lancet countdown: tracking progress on health and climate change. *Lancet*. 2017;389:1151–1164.
- 121 Ebi KL, Boyer C, Bowen KJ, Frumkin H, Hess J. Monitoring and evaluation indicators for climate change-related health impacts, risks, adaptation, and resilience. *Int J Environ Res Public Health*. 2018;15:E1943. <https://doi.org/10.3390/ijerph15091943>.
- 122 Benson R, Brunson C, Rigby J, et al. Real-time suicide surveillance supporting policy and practice. *Glob Ment Health*. 2022;9:384–388.
- 123 English PB, Sinclair AH, Ross Z, et al. Environmental health indicators of climate change for the United States: findings from the state environmental health indicator collaborative. *Environ Health Perspect*. 2009;117:1673–1681.
- 124 Driscoll DL, Sunbury T, Johnston J, Renes S. Initial findings from the implementation of a community-based sentinel surveillance system to assess the health effects of climate change in Alaska. *Int J Circumpolar Health*. 2013;72. <https://doi.org/10.3402/ijch.v72i0.21405>.
- 125 Doubleday A, Errett NA, Ebi KL, Hess JJ. Indicators to guide and monitor climate change adaptation in the US Pacific Northwest. *Am J Public Health*. 2020;110:180–188.
- 126 Brown VA. The uses of social and environmental health indicators in monitoring the effects of climate change. *Clim Change*. 1993;25:389–403.
- 127 Lowe D, Ebi KL, Forsberg B. Heatwave Early warning systems and adaptation advice to reduce human health consequences of heatwaves. *Int J Environ Res Public Health*. 2011;8:4623–4648.
- 128 Daalen KR van, Romanello M, Rocklöv J, et al. The 2022 Europe report of the Lancet Countdown on health and climate change: towards a climate resilient future. *Lancet Public Health*. 2022;7:e942–e965. [https://doi.org/10.1016/S2468-2667\(22\)00197-9](https://doi.org/10.1016/S2468-2667(22)00197-9).
- 129 Akearok GH, Holzman S, Kunnuk J, et al. Identifying and achieving consensus on health-related indicators of climate change in nunavut. *Arctic*. 2019;72:289–299.
- 130 Zhang S, Zhang C, Cai W, et al. The 2023 China report of the Lancet Countdown on health and climate change: taking stock for a thriving future. *Lancet Public Health*. 2023;8:e978–e995. [https://doi.org/10.1016/S2468-2667\(23\)00245-1](https://doi.org/10.1016/S2468-2667(23)00245-1).
- 131 Jung EM, Kim EM, Kang M, et al. Children's environmental health indicators for low- and middle-income countries in Asia. *Ann Glob Health*. 2017;83:530–540.
- 132 Heo S, Bell ML. Heat waves in South Korea: differences of heat wave characteristics by thermal indices. *J Expo Sci Environ Epidemiol*. 2019;29:790–805.
- 133 Dovie DBK, Dzodzomenyo M, Ogunseitan OA. Sensitivity of health sector indicators' response to climate change in Ghana. *Sci Total Environ*. 2017;574:837–846.

- 134 Mercuriali L, Oliveras L, Gómez A, et al. Un sistema de vigilancia de salud pública para el cambio climático en las ciudades. *Gac Sanit.* 2022;36:283–286.
- 135 Murage P, Hajat S, Macintyre HL, et al. Indicators to support local public health to reduce the impacts of heat on health. *Environ Int.* 2024;183:108391.
- 136 Nguyen AK, Liou Y-A. An approach for risk maps of vector-borne infectious diseases: ecological and adaptive capacity indicators. In: *Igarss 2018 - 2018 IEEE international geoscience and remote sensing symposium.* 2018:1164–1167.
- 137 Ministerio de Salud - Chile. *Reglamento sobre Notificación de Enfermedades Transmisibles de Declaración Obligatoria DTO N°158/04.* 2004.
- 138 Ministerio de Salud - Chile. *Registro Poblacional del Cáncer DTO vol. 5;* 2002. <https://www.bcn.cl/leychile/navegar?i=196649&f=2002-04-12>.
- 139 Ministerio de Salud - Chile. *Reglamento de Notificación Obligatoria de las Intoxicaciones Agudas con Pesticidas. DTO 88;* 2004. <https://www.bcn.cl/leychile/navegar?idNorma=231067>. Accessed February 20, 2023.
- 140 Ministerio de Salud - Chile. *Norma Técnica de Vigilancia de Enfermedades No Transmisibles y sus Factores de Riesgo;* 2011. <http://epi.minsal.cl/wp-content/uploads/2016/03/NormaTecnicaVENT.pdf>.
- 141 Ministerio de Salud - Chile. *Condiciones Sanitarias y Ambientales Básicas en los Lugares de Trabajo. DTO 594 [Basic Sanitary and Environmental Conditions in Workplaces DTO 594];* 1999. <https://www.bcn.cl/leychile/navegar?idNorma=167766>. Accessed February 20, 2023.
- 142 Gilbert R, Cliffe SJ. Public health surveillance. In: *Public Health Surveillance.* 2016:91–110.
- 143 Shokraneh F. Reproducibility and replicability of systematic reviews. *World J Meta Anal.* 2019;7:66–71.
- 144 Puljak L, Pieper D. Replicability in the context of systematic reviews: a call for a framework with considerations regarding duplication, overlap, and intentionality. *J Clin Epidemiol.* 2022;142:313–314.
- 145 Bordier M, Goutard FL, Antoine-Moussiaux N, Pham-Duc P, Lailier R, Binot A. Engaging stakeholders in the design of one health surveillance systems: a participatory approach. *Front Vet Sci.* 2021;8:646458. <https://doi.org/10.3389/fvets.2021.646458>.
- 146 Vargas C, Whelan J, Brimblecombe J, Allender S. Co-creation, co-design, co-production for public health – a perspective on definition and distinctions. *Public Health Res Pract.* 2022;32. <https://doi.org/10.17061/phrp3222211>.
- 147 Choi BCK. The past, present, and future of public health surveillance. *Scientifica.* 2012;2012:e875253. <https://doi.org/10.6064/2012/875253>.
- 148 Lombardo JS, Buckeridge DL. *Disease surveillance: a public health informatics approach.* John Wiley & Sons; 2012. ISBN: 978-1-118-56905-4.
- 149 Thacker SB, Stroup DF, Parrish RG, Anderson HA. Surveillance in environmental public health: issues, systems, and sources. *Am J Public Health.* 1996;86:633–638.
- 150 Koh D, Aw T-C. Surveillance in occupational health. *Occup Environ Med.* 2003;60:705–710.
- 151 Clayton S. Climate change and mental health. *Curr Environ Health Rep.* 2021;8:1–6.
- 152 Wainberg ML, Scorza P, Shultz JM, et al. Challenges and opportunities in global mental health: a research-to-practice perspective. *Curr Psychiatry Rep.* 2017;19:28. <https://doi.org/10.1007/s11920-017-0780-z>.
- 153 Hwong AR, Wang M, Khan H, et al. Climate change and mental health research methods, gaps, and priorities: a scoping review. *Lancet Planet Health.* 2022;6:e281–e291. [https://doi.org/10.1016/S2542-5196\(22\)00012-2](https://doi.org/10.1016/S2542-5196(22)00012-2).
- 154 Faisal-Cury A, Ziebold C, Rodrigues DM de O, Matijasevich A. Depression underdiagnosis: prevalence and associated factors. A population-based study. *J Psychiatr Res.* 2022;151:157–165.
- 155 Katz C, Bolton J, Sareen J. The prevalence rates of suicide are likely underestimated worldwide: why it matters. *Soc Psychiatry Psychiatr Epidemiol.* 2016;51:125–127. <https://doi.org/10.1007/s00127-015-1158-3>.
- 156 Tøllefsen IM, Hem E, Ekeberg Ø. The reliability of suicide statistics: a systematic review. *BMC Psychiatr.* 2012;12:9. <https://doi.org/10.1186/1471-244X-12-9>.
- 157 King CA, Merchant CR. Social and interpersonal factors relating to adolescent suicidality: a review of the literature. *Arch Suicide Res Off J Int Acad Suicide Res.* 2008;12:181–196.
- 158 Stack S. Contributing factors to suicide: political, social, cultural and economic. *Prev Med.* 2021;152:106498. <https://doi.org/10.1016/j.ypmed.2021.106498>.
- 159 Ebi KL, Ogden NH, Semenza JC, Woodward A. Detecting and attributing health burdens to climate change. *Environ Health Perspect.* 2017;125:085004. <https://doi.org/10.1289/EHP1509>.
- 160 Rossati A, Bargiacchi O, Kroumova V, Zaramella M, Caputo A, Garavelli PL. Climate, environment and transmission of malaria. *Infez Med.* 2016;24:93–104.
- 161 Chlebicz A, Śliżewska K. Campylobacteriosis, salmonellosis, yersiniosis, and listeriosis as zoonotic foodborne diseases: a review. *Int J Environ Res Public Health.* 2018;15:863. <https://doi.org/10.3390/ijerph15050863>.
- 162 Hawkins E, Sutton R. Time of emergence of climate signals. *Geophys Res Lett.* 2012;39. <https://doi.org/10.1029/2011GL050087>.
- 163 Orru H, Ebi KL, Forsberg B. The interplay of climate change and air pollution on health. *Curr Environ Health Rep.* 2017;4:504–513.
- 164 Latif M, Keenlyside NS. El Niño/Southern Oscillation response to global warming. *Proc Natl Acad Sci.* 2009;106:20578–20583.
- 165 Cai W, Santoso A, Collins M, et al. Changing el niño–southern oscillation in a warming climate. *Nat Rev Earth Environ.* 2021;2:628–644.
- 166 Kjellström T, Corvalán C. Framework for the development of environmental health indicators. *World Health Stat Q Rapp Trimest Stat Sanit Mond.* 1995;48:144–154.
- 167 Marmot M, Friel S, Bell R, Houweling TA, Taylor S. Closing the gap in a generation: health equity through action on the social determinants of health. *Lancet.* 2008;372:1661–1669.