ADDITIONAL FILE 3. Health Survey for England 1994-2008 sample size (blood sample), by gender, age and deprivation quintiles

uepri	vation qu					
	1994	1998	2003	2006	2008	Total
Men 16-54						
Q1	772	709	503	429	418	2831
×-	(21.2%)	(21.7%)	(19.6%)	(19.1%)	(20.5%)	(20.5%)
02	751	695	492	378	386	(20.370) 2702
Q2	(20.6%)	(21.3%)	(20.5%)	(18.2%)	(19.3%)	
~ •				· · · · ·		(20.1%)
Q3	824	653	453	441	376	2747
	(22.6%)	(20.0%)	(19.8%)	(23.4%)	(19.5%)	(21.1%)
Q4	701	639	507	379	375	2601
	(19.2%)	(19.6%)	(22.4%)	(20.4%)	(20.4%)	(20.3%)
Q5	602	567	371	340	368	2248
ו	(16.5%)	(17.4%)	(17.7%)	(18.8%)	(20.4%)	(18.0%)
Maria	<i></i>					(10.070)
Men ≥		277	410	252	204	1000
Q1	358 (21.1%)	377 (21.7%)	412 (26.1%)	352 (23.5%)	384	1883
					(26.0%)	(23.4%)
Q2	369	391	341	326	313	1740
	(21.8%)	(22.5%)	(22.5%)	(21.6%)	(21.6%)	(22.1%)
Q3	380	358	294	343	280	1655
	(22.4%)	(20.6%)	(20.0%)	(23.9%)	(19.9%)	(21.4%)
Q4	283	337	250	236	248	1354
Q+	(16.7%)	(19.4%)	(17.4%)	(17.3%)	(18.0%)	
05						(17.8%)
Q5	304 (18.0%)	272 (15.7%)	191 (14.1%)	186 (13.6%)	201 (14.5%)	1154
	(10.070)	(15.770)	(14.170)	(15.0%)	(14.570)	(15.4%)
Wome	n 16-54					
Q1	764	801	575	499	530	3169
	(20.3%)	(22.4%)	(19.9%)	(19.1%)	(21.7%)	(20.8%)
Q2	762	748	541	496	428	2975
<b>X</b> -	(20.3%)	(20.9%)	(20.1%)	(19.7%)	(18.4%)	(20.0%)
Q3	815	714	485	520	415	(20.070) 2949
QS	(21.7%)	(20.0%)	(18.4%)	(22.1%)	(18.2%)	
<u>.</u>						(20.2%)
Q4	753 (20.1%)	689 (19.3%)	580 (22.6%)	434 (19.2%)	431 (19.7%)	2887
						(20.1%)
Q5	662	625	477	424	444	2632
	(17.6%)	(17.5%)	(19.1%)	(19.9%)	(21.9%)	(18.9%)
Women ≥ 55						
Q1	401	417	495	415	437	2165
<b>X</b> <sup>1</sup>	(19.5%)	(21.0%)	(26.6%)	(23.3%)	(25.0%)	(22.7%)
01	442	435	402	391	379	
Q2	(21.5%)	(21.9%)	(22.0%)	(22.5%)	(22.3%)	2049
0.5						(22.0%)
Q3	470	414	380	413	352	2029
	(22.8%)	(20.8%)	(21.1%)	(24.3%)	(21.2%)	(22.0%)
Q4	355	372	301	262	286	1576
	(17.2%)	(18.7%)	(17.4%)	(16.5%)	(17.5%)	(17.5%)
Q5	393	350	224	207	223	1397
τ.	(19.1%)	(17.6%)	(12.9%)	(13.4%)	(14.0%)	(15.9%)

*Notes:* Respondents with a valid cholesterol measurement (including those taking lipid lowering drugs). Respondent totals are presented unweighted; percentages show distribution across the deprivation fifths (unweighted 1994 and 1998, weighted for non-response from 2003 onwards). Totals may not sum due to rounding. (Q1 = most affluent, Q5 = most deprived).