

Review

# Single deletions in mitochondrial DNA – Molecular mechanisms and disease phenotypes in clinical practice

R.D.S. Pitceathly<sup>a</sup>, S. Rahman<sup>a,b,c</sup>, M.G. Hanna<sup>a,d,\*</sup>

<sup>a</sup> MRC Centre for Neuromuscular Diseases, UCL Institute of Neurology and National Hospital for Neurology and Neurosurgery, Queen Square, London WC1N 3BG, UK

<sup>b</sup> Mitochondrial Research Group, Clinical and Molecular Genetics Unit, UCL Institute of Child Health, London WC1N 1EH, UK

<sup>c</sup> Metabolic Unit, Great Ormond Street Hospital for Children, London WC1N 3JH, UK

<sup>d</sup> Department of Molecular Neuroscience, UCL Institute of Neurology, Queen Square, London WC1N 3BG, UK

Received 9 November 2011; received in revised form 26 February 2012; accepted 21 March 2012

## Abstract

Over 20 years ago single clonal deletions were the first mitochondrial DNA (mtDNA) genetic defects described in association with human disease. Since then very large numbers of children and adults harbouring such deletions have been described and it is clear they are an important cause of human mitochondrial disease. However, there still remain many important challenges in relation to our understanding of mechanisms leading to deletion formation and propagation and in relation to the factors determining the complex and varying relationship between genotype and clinical phenotype. Although multidisciplinary team care is essential and can improve quality of life and outcomes for patients, a definitive molecular treatment for single mtDNA deletions remains an important translational research goal. Patients with mtDNA deletions exhibit a very wide range of different clinical phenotypes with marked variation in age at onset and disease severity. Single mtDNA deletions may enter into the differential diagnosis of many different paediatric and adult presentations across a wide range of medical specialties, although neurological presentations are amongst the most common. In this review, we examine the molecular mechanisms underpinning mtDNA replication and we consider the hypotheses proposed to explain the formation and propagation of single large-scale mtDNA deletions. We also describe the range of clinical features associated with single mtDNA deletions, outline a molecular diagnostic approach and discuss current management including the role of aerobic and resistance exercise training programmes.

© 2012 Elsevier B.V. All rights reserved.

**Keywords:** Single deletion; Mitochondrial DNA; Mitochondrial disease

## 1. Introduction

Mitochondria are double membrane-bound intracellular organelles that form large branching reticular networks inside all eukaryotic cells except erythrocytes. Their primary function is to convert food substrates in the presence of oxygen to energy, in the form of ATP, by aerobic

respiration. This oxidative phosphorylation (OXPHOS) reaction is performed in the mitochondrial inner membrane by five enzyme complexes (so-called complexes I, II, III, IV and V). Electrons are provided by NADH and FADH<sub>2</sub>, which are reducing agents supplied by the oxidation of carbohydrates and fatty acids via the Krebs cycle and  $\beta$ -oxidation, respectively. Electrons are passed between a series of donors and increasingly electronegative acceptors until the final reduction of oxygen to water. Each step releases energy which is used to actively pump protons (via complex I, III and IV) from within the mitochondrial matrix into the intermembrane space, thereby creating an electrochemical gradient ( $\Delta\psi_m$ ) across the inner mitochondrial membrane

\* Corresponding author at: MRC Centre for Neuromuscular Diseases, UCL Institute of Neurology and National Hospital for Neurology and Neurosurgery, Queen Square, London WC1N 3BG, UK. Tel.: +44 203 448 8014; fax: +44 203 448 3633.

E-mail address: [m.hanna@ucl.ac.uk](mailto:m.hanna@ucl.ac.uk) (M.G. Hanna).

[1]. Some electrons leak directly to oxygen and form reactive oxygen species (ROS) and these molecules have long since been implicated in a number of disease states and ageing [2–4]. The proton gradient enables the terminal OXPHOS enzyme, complex V (ATP synthase), to generate ATP from ADP and inorganic phosphate, by allowing protons to flow passively through it along the electrochemical gradient back into the mitochondrial matrix.

Mitochondria are unique amongst intracellular organelles in that they contain their own DNA [5,6]. The mitochondrial genome is a double-stranded circular structure consisting of 16,659 base-pairs (bp). Genes are located on both strands; the so-called heavy (H) and light (L) strands due to their relative composition of either mainly purines (adenine and guanine) or pyrimidines (thymine and cytosine), respectively, with the former having greater molecular weight than the latter. Mitochondrial DNA (mtDNA) is highly compact with no introns and only one major non-coding region (NCR), known as the control region (consisting of ~900 bp in mammals), which contains the origin of replication for the H-strand ( $O_H$ ), the origins of transcription for both H- and L-strands and the Displacement loop (D-loop). The D-loop is a triple-stranded region whose third strand (or 7S DNA) spans approximately 650 bp from nucleotide numbers 16,111 to 191 [7]. MtDNA codes for 37 genes which enable synthesis of 13 OXPHOS subunits required for aerobic respiration. The mtDNA is susceptible to oxidative DNA damage due to its close proximity to ROS at the inner mitochondrial membrane, lack of protective histone proteins, and limited mitochondrial DNA proof-reading and repair systems. As a result, mtDNA has a mutation rate 10–20 times that of nDNA [8,9].

*In vivo*, nuclear DNA (nDNA) adopts a condensed structure associated with histones and other proteins [10,11] called nucleosomes which form the basic repeating units of eukaryotic chromatin. These protein–DNA complexes compact, order, and protect the DNA, and create a degree of regulatory control to ensure its correct expression. They also facilitate co-ordinated copying and segregation of DNA molecules. MtDNA is also packaged as protein–DNA complexes called nucleoids, which may be the units of mtDNA transmission and inheritance [12,13]. Each mitochondrial nucleoid appears to contain a single mtDNA molecule. Nucleoids are associated with the inner mitochondrial membrane and are evenly spaced along the mitochondrial reticulum [14–17]. Interestingly, the nucleoid distribution process appears to be reliant on protein–protein interactions and is independent from the DNA itself [18]. More than 25 nucleoid-interacting proteins have been identified, but their precise role in mtDNA maintenance is not fully established. In fact, many of these proteins appear to have functions unrelated to mtDNA maintenance suggesting they are bifunctional [18]. Some are important for packaging, division and sorting of nucleoids, whereas others are involved in mtDNA replication, repair and transcription [17]. A detailed review of

mitochondrial nucleoid-associated proteins and their function is given elsewhere [19].

Although the mtDNA encodes 13 essential OXPHOS subunits, all other proteins needed for mitochondrial OXPHOS function are encoded by nuclear genes. These include the remaining 77 OXPHOS subunits, and proteins involved with transcription and translation of mtDNA, assembly of the OXPHOS enzyme complexes, biosynthesis of mobile electron carriers in the inner mitochondrial membrane, import of solutes and proteins into the mitochondrion, regulation of mitochondrial membrane dynamics and composition, and mitochondrial fission and fusion. More than 80 nuclear genes have already been linked to human mitochondrial disease; however, an estimated 1500 nuclear-encoded proteins are targeted to mitochondria [20]. Defects in any of these genes could theoretically cause a mitochondrial disorder.

Single large-scale deletions of mtDNA were the first mtDNA genetic defects to be described and associated with human mitochondrial disorders [21]. In 1988 Holt et al. identified a deleted mtDNA subpopulation in skeletal muscle but not blood in nine patients with mitochondrial myopathy. They found that the deleted population coexisted with wild-type (WT) mtDNA in muscle, a state known as heteroplasmy. It has subsequently been shown that such single deletions account for a quarter of all adult patients with mtDNA-related disease [22]; data consistent with the authors' own clinical practice (unpublished data). They are also an important cause of paediatric mtDNA-related mitochondrial disease. Single mtDNA deletions are considered sporadic events with low inheritance risk, whereas multiple mtDNA deletions are the result of primary nuclear defects in genes responsible for mtDNA maintenance or nucleoside metabolism and follow Mendelian inheritance patterns [23].

Here we review current understanding of the molecular mechanisms underlying single large-scale mtDNA deletion formation. We will also consider the clinical phenotypes, diagnostic work-up and management of this subgroup of mitochondrial disorders.

## 2. Mitochondrial DNA replication

mtDNA encodes only 13 of the 90 essential OXPHOS subunits. The remaining subunits, along with all the proteins required for their transcription, translation, post-translational modification and assembly, are nuclear-encoded. The nuclear encoded proteins include those essential for mtDNA maintenance and repair.

Accurate mtDNA replication is essential to maintain the  $\Delta\psi_m$  electrochemical potential, established across the inner mitochondrial membrane by respiration, and is necessary for energy production in the form of ATP. Although the mtDNA replication machinery is nuclear-encoded, it is completely separate from its nuclear counterpart, and consists of just 4 basic enzyme components: DNA polymerase gamma (POL $\gamma$ ); DNA helicase Twinkle; mitochondrial

single-stranded DNA-binding protein (mtSSB); and mitochondrial RNA polymerase (POLRMT). Unlike nuclear DNA which replicates once during cell division, mtDNA replicates continuously throughout the cell cycle in dividing and post-mitotic cells.

Although the replication apparatus is relatively well characterised, the precise mode of mtDNA replication has been intensely debated [24–26]. For many years mtDNA replication was thought to occur exclusively by a strand-displacement model, where a leading strand is synthesised from  $O_H$  and a lagging strand from the origin of L-strand replication ( $O_L$ ) in an asymmetric and unidirectional pattern [27–30]. In 2000, Holt et al. proposed an alternative model in which leading- and lagging-strand syntheses were coupled (so-called synchronous or coupled replication), on the basis of Brewer and Fangman 2D-agarose gel electrophoresis (2D-AGE) [31]. The strand-coupled mechanism was suggested to initiate simultaneously in both directions from a broad zone, OriZ [32]. Later studies demonstrated that some replication intermediates were RNase H-sensitive, which led to the assumption that they were derived from ribonucleotide-rich DNA synthesis initiated unidirectionally from the non-coding region [33–35]. According to this replication mode, referred to as the ribonucleotide incorporation throughout the lagging strand (RITOLS) model, the lagging strand is initially laid down as RNA before being converted to DNA. In a similar manner to the strand-displacement model, RITOLS predicts delayed lagging strand DNA synthesis. Although the precise mode of mtDNA replication is contentious, the weight of experimental data in support of all three models makes their combined existence likely.

### 3. Molecular mechanisms underlying formation and timing of single mitochondrial DNA deletions

Most deletions (~85%) occur within the major arc between  $O_H$  and  $O_L$  and are flanked by runs of direct homologous base-pair repeats. It is generally considered that mtDNA deletions arise during replication through slipped-strand mispairing between direct homologous base-pairs (Fig. 1A–D) [36]. This mechanism assumes the asynchronous strand displacement model of mtDNA replication, since it is reliant on the presence of long stretches of single-stranded DNA which do not exist in the coupled and RITOLS replication models. During replication a single stranded repeat of the H-strand misanneals with a newly exposed L-strand repeat (Fig. 1C). This forms a downstream loop of single-stranded parental H-strand which is degraded, with subsequent ligation of the free H-strands and continued replication. The end result is one WT and one deleted mtDNA molecule (Fig. 1D). A recent alternative hypothesis proposed that deletions form during repair of mtDNA at double-stranded breaks, rather than during replication [37]. During repair of a double-stranded break (a commonly occurring transition state needed to restore the replication fork after a stalling event) single-stranded

regions are generated by exonuclease activity. The single strands would be free to anneal with microhomologous sequences [38]. Subsequent repair, ligation and degradation of the exposed single strands would lead to mtDNA molecules with a deleted portion. Krishnan et al. suggest that if deletion formation only occurs during replication, higher levels of deleted molecules should be present in mitotic rather than post-mitotic cells (where mtDNA replication is less frequent); however, this has been shown not to be the case in rapidly dividing colonic tissue [39]. The slipped-strand model also predicts conservation of the 3' repeat sequence in the deleted daughter molecule, which does not always occur [40,41].

Uncertainty surrounding the pathophysiological mechanism underlying deletion formation in mtDNA includes the precise timing of the deletional event itself, which may occur during early embryonic development [42,43] or oogenesis [37]. Deleted species have been demonstrated in tissue of mesodermal (skeletal muscle) and ectodermal (brain) origin, suggesting that the deletion is likely to occur prior to differentiation of these two embryonic layers [43]. However, single large-scale mtDNA deletions have also been shown to exist in the oocyte prior to fertilisation [44,45]. In the case of progressive external ophthalmoplegia (PEO), where muscle is the only tissue clinically involved, the deletion may arise later in development, for example during differentiation of the muscle precursor cells. An earlier deletion event during germ cell maturation followed by mitotic segregation of the mutant mtDNA might also explain the tissue-specific phenotype of this disorder [46].

The case for deletion formation during germline maturation is compelling. Human oocytes contain around 100,000 mtDNA molecules [44]. It is estimated that half of mature oocytes in healthy women each contain 5–20 molecules with the same “common” 4977 bp mtDNA deletion found in patients with mtDNA deletion disorders [44,47,48]. Chinnery et al. have hypothesised that the inherited deletion arises in the grandmother's oocyte [46]. As mtDNA is maintained in a mature oocyte for many years without replicating, Krishnan et al. [37] suggested that the original mtDNA deletion event occurs in the grandmother's oocyte during repair of damaged mtDNA rather than during mtDNA replication. The lack of any correlation between increasing age of the maternal grandmother and risk of having a grandchild with a single mtDNA deletion disorder, however, does not appear to support this hypothesis [49]. So, if an oocyte containing deleted mtDNA molecules due to aberrant replication or repair from the maternal grandmother is fertilised, the zygote is destined to become the unaffected mother, and it is possible a proportion of primordial germ cells contain low levels of deleted mtDNA. The deleted mtDNA levels must, however, reach a critical biochemical threshold for disease to develop in the grandchild.

This rapid shift of mtDNA heteroplasmy between generations is attributed to the so-called genetic bottleneck theory of mtDNA transmission [50]. Although the precise




Fig. 1. Diagram illustrating the consensus slipped-strand model of mitochondrial DNA deletion formation (adapted from Shoffner et al. [36]). (A) A mitochondrial DNA (mtDNA) molecule prior to replication containing two direct homologous base-pair (bp) repeats (a/a1 and b/b1). The origin of heavy strand ( $O_H$ ) and light strand ( $O_L$ ) replication are shown. (B) Replication of the mtDNA is initiated at  $O_H$ . During replication the daughter heavy strand (brown arrow) uses the parent light strand as a template, thereby displacing the parent heavy strand. (C) The first direct homologous bp repeat on the parent heavy strand (a) misanneals with the second newly exposed direct bp repeat on the parent light strand (b1), forming a downstream loop of heavy strand. The single-stranded downstream loop is susceptible to damage and is subsequently degraded until double-stranded DNA is reached, with loss of the second parent heavy strand bp repeat (b). The free ends of the parent heavy strand are then ligated and replication of the daughter heavy strand continues. When heavy strand synthesis reaches  $O_L$  (two-thirds around the circumference of the mtDNA molecule), replication of the daughter light strand begins using the parent heavy strand as a template, in the opposite direction. As such, mtDNA replication occurs in an asynchronous manner. (D) Slipped-strand replication results in the synthesis of one wild-type (consisting of the parent light strand and the daughter heavy strand) and one deleted (consisting of the parent heavy strand and the daughter light strand) mtDNA molecule.

mechanism and timing of the bottleneck is widely discussed in the literature, there are currently three main biological mechanisms that might potentially explain the phenomenon: (1) amplification of a reduced number of mtDNA molecules during early embryonic primordial germ cell development (approximately 7–9 days post-fertilisation) [51]; (2) selective replication and transmission of a subpopulation of mtDNA molecules at a later stage of development during post-natal oocyte maturation [52]; (3) segregation of identical heteroplasmic mtDNA molecules thereby forming homoplasmic units, with effective restriction on the pool available for mtDNA transmission without any significant reduction in mtDNA content in the germ cells [53,54].

If a deleted mtDNA molecule or homoplasmic deleted unit is selected for clonal expansion after the bottleneck, an accumulation of abnormally high levels of deleted mtDNA molecules might occur in a small number of

mature oocytes in the unaffected mother. This would predict a 5/100,000 risk of offspring inheriting high levels of the pathogenic mtDNA deletion, which roughly corresponds to the incidence of mtDNA deletion disorders in the general population [22]. Clinically unaffected mothers are very unlikely to have more than one affected child. However, there is an estimated 4% risk of subsequent transmission from an affected woman [46].

#### 4. Size and location of single mitochondrial DNA deletions

The size and location of large-scale deletions may vary. However, one third of patients harbour a common 4977 bp mtDNA deletion [44,47,48]. The deleted molecule exists as a single species and is heteroplasmic, with variable mutant levels detectable in different tissues. The majority of deletions arise within the long arc between  $O_H$  and  $O_L$  but retain these replication sites. The likely explanation for this

is that for mtDNA deletions to accumulate within a cell, the deleted molecules must retain the ability to replicate. The deletion removes a variable number of essential protein-encoding subunits for complexes I, III, IV and V and tRNA-encoding genes required for translation of the mtDNA-encoded OXPHOS complexes, with resulting dysfunction in OXPHOS.

Early work found no correlation between clinical severity and the type, size or location of the large-scale rearrangement [55]. However, Yamashita et al. reported that a larger deleted segment, and consequently a greater number of deleted tRNAs, correlated with an earlier age of disease onset. Patients with large numbers of deleted tRNAs also appear to be more likely to develop Kearns–Sayre syndrome (KSS), suggesting that the quality of the deletion is somehow linked either to the development of the KSS phenotype or increased propensity for multi-systemic involvement [56]. Since patients with the same “common” deletion show different clinical symptoms, other factors are postulated to contribute to the observed phenotypic heterogeneity, such as systemic distribution and mutant load of deleted mtDNA molecules. The time-point at which the deletion arises has also been suggested to determine the affected tissues, clinical phenotype, percentage and location of the deletion [57].

Despite the probable importance of deletion size and location, many other factors are likely to contribute to the varied clinical phenotype and severity seen in this patient group. Deletions are heteroplasmic and mutant load may vary between tissue types and can shift over time. This may explain the clinical features (dependent on the tissue with the greatest energy demand and level of mutant mtDNA) and delayed onset that is sometimes seen in patients with the disease. Random mitotic segregation of mutant and WT mtDNA molecules, relaxed replication, tissue-specific energy requirements, contrasting selection bias of mitotic and post-mitotic cells, together with the mtDNA and nuclear haplotypes, are all thought to influence the ultimate spectrum of disease. There is marked tissue specificity seen with mtDNA deletions when compared with other mtDNA mutations [58–61]. This observation, taken together with ongoing research examining the role the mitochondrial nucleoid plays in mtDNA distribution, is likely to be important in unravelling the cause of extreme tissue segregation seen in mtDNA deletion disorders.

## 5. Clinical phenotypes associated with single mitochondrial DNA deletions

Three sporadic clinical syndromes are classically associated with large-scale deletions of mtDNA. However, it is increasingly recognised that there is a continuum of clinical phenotypes associated with single mtDNA deletions, and a large number of variants exist. The less common phenotypes associated with single mtDNA deletions are summarised in Table 1, but for purposes of this review we will describe each of the classical phenotypes in detail.

Table 1  
Clinical syndromes and associated features of single mtDNA deletion disorders.

	Reference
<i>Clinical syndromes associated with single mtDNA deletions</i>	
Pearson marrow-pancreas syndrome	[62–64]
Kearns–Sayre syndrome	[76–80]
Progressive external ophthalmoplegia	[109]
Isolated anaemia ± lactic acidosis	[65,70]
Pancreatic exocrine insufficiency without marrow involvement	[72]
Pancytopenia and tubulopathy	[73]
Tubulopathy and hypoparathyroidism ± encephalopathy	[74,75]
Mitochondrial myopathy, encephalopathy, lactic acidosis and stroke-like episodes	[110]
Myoclonic epilepsy with ragged red fibres	[111]
Leigh syndrome	[66–68]
Diabetes and deafness	[112]
<i>Other clinical features associated with single mtDNA deletions<sup>a</sup></i>	
Myopathy	[109]
Endocrinopathy: multiple endocrine disturbance; hypoparathyroidism; Addison’s disease; adrenal insufficiency; short stature	[113–116]
Gastrointestinal disturbance: constipation; diarrhoea; pseudo-obstruction; cyclical vomiting; gastro-oesophageal reflux	[116,117]
Organic acidopathy	[116,118]
Seizures	[116]
Cardiomyopathy and cardiac conduction defects	[69,117,119]
Neuropsychiatric symptoms and dementia	[120]

<sup>a</sup> These may occur in isolation, but are more usually seen in the context of an early-onset multisystem disorder.

### 5.1. Pearson marrow-pancreas syndrome (PMPS)

This is a severe and often rapidly fatal disorder of infancy/early childhood characterised by sideroblastic anaemia and variable involvement of other blood cells, together with exocrine pancreatic dysfunction. Onset is usually in early infancy with transfusion-dependent sideroblastic anaemia, neutropaenia, thrombocytopenia or pancytopenia, poor growth and lactic acidosis. Pancreatic dysfunction may be present at disease onset or it may develop later. Other clinical features include an enteropathy that may need parenteral nutrition support, and severe renal tubulopathy. Bone marrow examination typically reveals high levels of ringed sideroblasts and vacuolisation of myeloid precursors [62–64]. The disorder is frequently fatal in early childhood because of overwhelming lactic acidosis or liver failure. It is estimated that one out of three neonatal cases of PMPS surviving infancy develop features compatible with Kearns–Sayre syndrome (KSS) [65], including cardiac conduction defects, sensorineural hearing loss, pigmentary retinopathy and progressive neurological impairment which can sometimes resemble Leigh syndrome [66–69]. In PMPS deleted mtDNA molecules are usually present at higher levels in blood than other tissues; hence haematological manifestations are such a prominent feature of the syndrome. Due to the high replication rate of blood cells, mtDNA deletions can be cleared from these

cells with time, leading to resolution of the haematological manifestations (for example, the need for transfusions has usually resolved by 2 years). In contrast, deleted mtDNA molecules tend to progressively accumulate in other tissues (especially non-dividing tissues), leading to the multisystem and relentlessly progressive nature of this disorder. Despite the classical description of PMPS, however, it is important to be aware of the more atypical presentations. Neonates can present with anaemia as the only onset symptom in one third of cases [65,70], and hepatopathy may be the most prominent feature [71]. The diagnosis should also be considered when there is isolated pancreatic exocrine insufficiency without marrow involvement [72]. Pancytopenia and tubulopathy can occur without exocrine pancreatic dysfunction [73], and there are case reports of milder, childhood onset, isolated tubulopathy and hypoparathyroidism [74], which has also been reported in association with recurrent encephalopathy [75].

### 5.2. Kearns–Sayre syndrome (KSS)

KSS is a multisystem disorder defined clinically by the triad of PEO and pigmentary retinopathy with onset before the age of 20 years plus at least one of: cardiac conduction block; cerebrospinal fluid (CSF) protein concentration greater than 0.1 g/L; cerebellar ataxia [76–80]. Frequent additional signs include sensorineural hearing loss, renal tubular acidosis, dementia, seizures, short stature and endocrine disturbance (diabetes mellitus, hypoparathyroidism and growth hormone deficiency). A progressive skeletal myopathy is also frequently seen. The most common magnetic resonance imaging (MRI) findings are cerebral and cerebellar atrophy with bilateral, often symmetrical, hyperintense lesions in the subcortical white matter, thalamus, basal ganglia and brainstem [81–83]. Interestingly, there appears to be little correlation between neurological deficits and the severity of the MRI features [84]. KSS is most commonly associated with single large-scale deletions of mtDNA [56]. We recently identified a case of KSS caused by nuclear genomic dysfunction due to mutations in *RRM2B* in a patient with multiple deletions of the mtDNA [85]. It is, therefore, possible for KSS to follow a Mendelian pattern of inheritance, although the majority of cases are sporadic.

### 5.3. Progressive external ophthalmoplegia (PEO)

PEO is considered the mildest of the three syndromes most commonly associated with single mtDNA deletions. It is characterised by slowly progressive bilateral ptosis and limitation of eye movements (often with reduced upgaze early in the disease process) secondary to extra-ocular muscle (EOM) weakness. There is often no diplopia as the ophthalmoparesis is bilateral, relatively symmetrical, and has gradual onset. Additional clinical features may include oropharyngeal dysfunction causing dysphagia, proximal limb weakness and exercise intolerance. Isolated

PEO is usually associated with a normal lifespan. The reason for selective involvement of the EOMs in the single deletion disorders is not understood. It may relate to distinct structural, functional, biochemical and immunological properties of this muscle group compared to other skeletal muscles [86]. The EOMs have also been reported to have a lower biochemical threshold for COX deficiency than limb muscles [87].

## 6. Investigations and diagnosis of single mitochondrial DNA deletion disorders

Table 1 outlines the potentially wide range of different clinical settings in which mitochondrial disease caused by single deletions of mtDNA enters into the differential diagnosis. The diagnosis of single mtDNA deletion disorders involves careful clinical assessment and tailored investigations that are often guided by the age of the patient.

### 6.1. Mitochondrial DNA analysis

Children, unlike adults, often have single large-scale rearrangements detectable in blood, and muscle biopsy is seldom required. In PMPS the mtDNA deletions are usually present at higher levels in blood than other tissues, so diagnosis is reliably made using leucocyte mtDNA. The deleted mtDNA molecules are selectively shed from the rapidly dividing leucocytes with increasing age [88–91]. The consequence is that the deleted mtDNA molecules are often undetectable in blood by early adolescence. Deleted mtDNA molecules, however, appear more readily detectable in cells extracted from urine, and a diagnosis can occasionally be achieved by analysing mtDNA from uroepithelial cells (extracted from 30 to 50 mls of urine). Unfortunately, in contrast to the m.3243A > G point mutation, where uroepithelial cell mutant levels have been shown to closely resemble mutant levels present in muscle tissue [92], urine levels of mtDNA deletions are not always representative of the most clinically affected tissues [93]. As a consequence, a diagnostic muscle biopsy is still often ultimately required. MtDNA is analysed using long-range polymerase chain reaction (PCR) and Southern blot genetic techniques to demonstrate large-scale rearrangements (deletions and/or duplications) of the mitochondrial genome.

### 6.2. Muscle histochemical analysis

The muscle tissue itself can be examined using light microscopy for features that support the diagnosis of mitochondrial disease. These include subsarcolemmal mitochondrial accumulation demonstrated on routine histochemical staining as “ragged-red fibres” (RRF) in the modified Gomori trichrome stain, or as succinate dehydrogenase (SDH)-positive fibres. Cytochrome *c* oxidase (COX) staining may reveal fibres deficient in complex IV, and is particularly useful when combined with sequential SDH staining since COX-deficient SDH-positive fibres appear

as “ragged-blue” fibres. Both RRF and COX-negative fibres are a common and expected finding in patients with single mtDNA deletions.

### 6.3. Respiratory chain enzymology

Biochemical analysis of the respiratory chain enzymes can be performed on fresh muscle by measuring substrate oxidation using polarography. Specific enzyme activities can also be measured using spectrophotometric assays using either fresh muscle or muscle immediately frozen in liquid nitrogen or isopentane and stored at  $-80^{\circ}\text{C}$ . Single mtDNA deletions usually involve several genes, including at least one tRNA, so that global mitochondrial protein synthesis is affected. As a result, activities of OXPHOS complexes containing mtDNA-encoded subunits (complexes I, III, IV and V) can potentially be reduced [94]. However, normal OXPHOS enzyme activities may be observed in the muscle of patients with single mtDNA deletions; therefore, genetic analysis should be performed if there is clinical suspicion of a single mtDNA disorder, regardless of the biochemical findings.

## 7. Management of single mitochondrial DNA deletion disorders

This review does not provide a comprehensive discussion of all current treatment regimens for mitochondrial disorders; these may be found in detail elsewhere [94–96]. The following is a summary of treatment strategies aimed specifically at patients with single mtDNA deletions.

### 7.1. Importance of multidisciplinary team care to optimise quality of life and improve outcomes

Since no curative therapy currently exists for patients with mitochondrial disease, symptom management is critical to maintain quality of life. Awareness of potential disease complications allows early intervention, thereby reducing morbidity and mortality. All patients require support from a dedicated multidisciplinary team to facilitate careful consideration of both medical and non-medical needs.

Clinical issues related to single mtDNA deletion disorders require input from doctors with specialist expertise in mitochondrial disorders. Patients with PMPS typically present during childhood to paediatricians and need supportive measures, including regular blood transfusions, pancreatic enzyme replacement, fat-soluble vitamin supplementation, hormone replacement for endocrine disease, and enteral feeding. Epilepsy, movement disorders and neuromuscular disease are common manifestations of mitochondrial disease in later life. As a consequence, patients usually present to the neurologist during adulthood. Ophthalmic review is often necessary to assess and monitor retinopathy, exclude cataracts, and provide prosthetic lid props or perform surgery for severe ptosis.

A cardiology opinion is mandatory even if cardiac symptoms are absent. This is particularly important in patients with KSS in whom cardiac conduction defects are common and a permanent pacemaker or implantable defibrillator may be required. Audiological assessment for aids and cochlear implants for hearing loss, and referral to endocrinology, gastroenterology, renal or palliative care may all be appropriate. Finally, advice should be sought prior to general anaesthesia so that perioperative complications related to cardiac arrhythmia and administration of potentially mitotoxic agents are avoided.

A specialist mitochondrial disease nurse is crucial and provides the patient with a point of contact so that medical needs are addressed between out-patient clinic appointments. They also offer invaluable practical advice concerning available social and income support. Speech and language therapy and dietetic review are required if there is bulbar involvement or problems surrounding adequate nutrition, and occupational therapy, physiotherapy and orthotics can maintain and improve mobility to enable an independent lifestyle.

### 7.2. Pharmacotherapy

Pharmacotherapy, including Co-enzyme Q<sub>10</sub> (Ubiquinone), creatine and dichloroacetate have all been studied in randomised controlled double-blind clinical trials of patients with single mtDNA deletion disorders [97–101]. However, a Cochrane systematic review concluded there was no clear evidence supporting their routine use in mitochondrial disease [102]. Cerebral folate deficiency can occur in KSS [103,104], and significant clinical improvement and resolution of white matter changes on MRI have been reported in patients with KSS treated with folinic acid which, unlike folate, is able to cross the blood brain barrier [105]. The frequency of cerebral folate deficiency in KSS is unknown, but CSF measurement and folinic acid supplementation should be considered, particularly when leukoencephalopathy is present.

### 7.3. Exercise therapy to reduce deleted mitochondrial DNA molecules

Finally, exercise training has provided promising results in patients with single mtDNA deletion disorders. Activation of muscle satellite cells by specific exercise training regimens allows WT mtDNA to repopulate muscle tissue and effectively reduce mutant load below the biochemical threshold needed to impair OXPHOS [106]. Endurance exercise training has been shown to increase muscle aerobic capacity [107], whilst resistance training improves muscle strength and oxidative ability [108]. The merits of resistance versus endurance exercise training are currently being evaluated in patients with single mtDNA deletions, with the aim of providing tailored exercise regimes suitable for individual patients in the near future.

## 8. Conclusions

Although single mtDNA deletions were the first mtDNA mutations associated with human disease [21], there are still many unanswered questions surrounding their origin and propagation. The variable clinical severity and heterogeneity of disease phenotype remains incompletely understood. Single mtDNA deletion disorders continue to provide a diagnostic challenge to paediatric and adult neurologists. Although primarily sporadic, a Mendelian inheritance pattern should not be used as an exclusion criterion for KSS, especially in the context of multiple mtDNA deletions [85]. Diagnosis is usually achieved using DNA extracted from blood in children. However, use of uroepithelial cells is increasingly being recognised as a non-invasive alternative to muscle biopsy, particularly in children with cardiac conduction disorders in whom a general anaesthetic poses significant risks. Although treatment is primarily supportive, a multidisciplinary team complemented by a tailored exercise regimen significantly improves overall function and life quality for patients and their families.

## Acknowledgements

R.D.S.P. is funded by MRC Grant Number G0800674. M.G.H. is supported by an MRC Centre grant G0601943 and The Myositis Support Group; S.R. is supported by Great Ormond Street Hospital Children's Charity. This review was supported by the NIHR UCLH/UCL Comprehensive Biomedical Research Centre and undertaken at University College London Hospitals/University College London, which received a proportion of funding from the Department of Health's National Institute for Health Research Biomedical Research Centres funding scheme.

## References

- [1] Mitchell P. Coupling of phosphorylation to electron and hydrogen transfer by a chemi-osmotic type of mechanism. *Nature* 1961;191:144–8.
- [2] Pieczek SR, Neustadt J. Mitochondrial dysfunction and molecular pathways of disease. *Exp Mol Pathol* 2007;83:84–92.
- [3] Harman D. Aging: a theory based on free radical and radiation chemistry. *J Gerontol* 1956;11:298–300.
- [4] Harman D. Free radical theory of aging. *Mutat Res* 1992;275:257–66.
- [5] Nass MM, Nass S. Intramitochondrial fibers with DNA characteristics. I. Fixation and electron staining reactions. *J Cell Biol* 1963;19:593–611.
- [6] Schatz G, Haslbrunner E, Tuppy H. Deoxyribonucleic acid associated with yeast mitochondria. *Biochem Biophys Res Commun* 1964;15:127–32.
- [7] Mao C-C, Holt IJ. Clinical and molecular aspects of diseases of mitochondrial DNA instability. *Chang Gung Med J* 2009;32:354–69.
- [8] Richter C, Park JW, Ames BN. Normal oxidative damage to mitochondrial and nuclear DNA is extensive. *Proc Natl Acad Sci USA* 1988;85:6465–7.
- [9] Lynch M, Koskella B, Schaack S. Mutation pressure and the evolution of organelle genomic architecture. *Science* 2006;311:1727–30.
- [10] Kornberg RD. Chromatin structure: a repeating unit of histones and DNA. *Science* 1974;184:868–71.
- [11] Olins AL, Olins DE. Spheroid chromatin units (v bodies). *Science* 1974;183:330–2.
- [12] Malka F, Lombès A, Rojo M. Organization, dynamics and transmission of mitochondrial DNA: focus on vertebrate nucleoids. *Biochim Biophys Acta* 2006;1763:463–72.
- [13] Holt IJ, He J, Mao C-C, et al. Mammalian mitochondrial nucleoids: organizing an independently minded genome. *Mitochondrion* 2007;7:311–21.
- [14] Legros F, Malka F, Frachon P, Lombes A, Rojo M. Organization and dynamics of human mitochondrial DNA. *J Cell Sci* 2004;117:2653.
- [15] Iborra FJ, Kimura H, Cook PR. The functional organization of mitochondrial genomes in human cells. *BMC Biol* 2004;2:9.
- [16] Garrido N, Griparic L, Jokitalo E, Wartiovaara J, van der Bliek AM, Spelbrink JN. Composition and dynamics of human mitochondrial nucleoids. *Mol Biol Cell* 2003;14:1583–96.
- [17] Chen XJ, Butow RA. The organization and inheritance of the mitochondrial genome. *Nat Rev Genet* 2005;6:815–25.
- [18] Kucej M, Butow RA. Evolutionary tinkering with mitochondrial nucleoids. *Trends Cell Biol* 2007;17:586–92.
- [19] Spelbrink JN. Functional organization of mammalian mitochondrial DNA in nucleoids: history, recent developments, and future challenges. *IUBMB Life* 2010;62:19–32.
- [20] Calvo S, Jain M, Xie X, et al. Systematic identification of human mitochondrial disease genes through integrative genomics. *Nat Genet* 2006;38:576–82.
- [21] Holt IJ, Harding AE, Morgan-Hughes JA. Deletions of muscle mitochondrial DNA in patients with mitochondrial myopathies. *Nature* 1988;331:717–9.
- [22] Chinnery PF, Johnson MA, Wardell TM, et al. The epidemiology of pathogenic mitochondrial DNA mutations. *Ann Neurol* 2000;48:188–93.
- [23] Suomalainen A, Kaukonen J. Diseases caused by nuclear genes affecting mtDNA stability. *Am J Med Genet* 2001;106:53–61.
- [24] Bogenhagen DF, Clayton DA. The mitochondrial DNA replication bubble has not burst. *Trends Biochem Sci* 2003;28:357–60.
- [25] Holt IJ, Jacobs HT. Response: the mitochondrial DNA replication bubble has not burst. *Trends Biochem Sci* 2003;28:355–6.
- [26] Bogenhagen DF, Clayton DA. Concluding remarks: the mitochondrial DNA replication bubble has not burst. *Trends Biochem Sci* 2003;28:404–5.
- [27] Robberson DL, Clayton DA. Replication of mitochondrial DNA in mouse L cells and their thymidine kinase-derivatives: displacement replication on a covalently-closed circular template. *Proc Natl Acad Sci USA* 1972;69:3810–4.
- [28] Robberson DL, Kasamatsu H, Vinograd J. Replication of mitochondrial DNA. Circular replicative intermediates in mouse L cells. *Proc Natl Acad Sci USA* 1972;69:737–41.
- [29] Clayton DA. Replication of animal mitochondrial DNA. *Cell* 1982;28:693–705.
- [30] Brown TA, Cecconi C, Tkachuk AN, Bustamante C, Clayton DA. Replication of mitochondrial DNA occurs by strand displacement with alternative light-strand origins, not via a strand-coupled mechanism. *Genes Dev* 2005;19:2466–76.
- [31] Holt IJ, Lorimer HE, Jacobs HT. Coupled leading- and lagging-strand synthesis of mammalian mitochondrial DNA. *Cell* 2000;100:515–24.
- [32] Bowmaker M, Yang MY, Yasukawa T, et al. Mammalian mitochondrial DNA replicates bidirectionally from an initiation zone. *J Biol Chem* 2003;278:50961–9.
- [33] Yang MY, Bowmaker M, Reyes A, et al. Biased incorporation of ribonucleotides on the mitochondrial L-strand accounts for apparent strand-asymmetric DNA replication. *Cell* 2002;111:495–505.
- [34] Yasukawa T, Yang M-Y, Jacobs HT, Holt IJ. A bidirectional origin of replication maps to the major noncoding region of human mitochondrial DNA. *Mol Cell* 2005;18:651–62.
- [35] Yasukawa T, Reyes A, Cluett TJ, et al. Replication of vertebrate mitochondrial DNA entails transient ribonucleotide incorporation throughout the lagging strand. *EMBO J* 2006;25:5358–71.



- [36] Shoffner JM, Lott MT, Voljavec AS, Soueidan SA, Costigan DA, Wallace DC. Spontaneous Kearns–Sayre/chronic external ophthalmoplegia plus syndrome associated with a mitochondrial DNA deletion: a slip-replication model and metabolic therapy. *Proc Nat Acad Sci USA* 1989;86:7952–6.
- [37] Krishnan KJ, Reeve AK, Samuels DC, et al. What causes mitochondrial DNA deletions in human cells? *Nat Genet* 2008;40:275–9.
- [38] Haber JE. Partners and pathways repairing a double-strand break. *Trends Genet* 2000;16:259–64.
- [39] Taylor RW, Barron MJ, Borthwick GM, et al. Mitochondrial DNA mutations in human colonic crypt stem cells. *J Clin Invest* 2003;112:1351–60.
- [40] Bua E, Johnson J, Herbst A, et al. Mitochondrial DNA-deletion mutations accumulate intracellularly to detrimental levels in aged human skeletal muscle fibers. *Am J Hum Genet* 2006;79:469–80.
- [41] Reeve AK, Krishnan KJ, Elson JL, et al. Nature of mitochondrial DNA deletions in substantia nigra neurons. *Am J Hum Genet* 2008;82:228–35.
- [42] Zeviani M, Gellera C, Pannacci M, et al. Tissue distribution and transmission of mitochondrial DNA deletions in mitochondrial myopathies. *Ann Neurol* 1990;28:94–7.
- [43] Marzuki S, Berkovic SF, Saifuddin Noer A, et al. Developmental genetics of deleted mtDNA in mitochondrial oculomyopathy. *J Neurol Sci* 1997;145:155–62.
- [44] Chen X, Prosser R, Simonetti S, Sadlock J, Jagiello G, Schon EA. Rearranged mitochondrial genomes are present in human oocytes. *Am J Hum Genet* 1995;57:239–47.
- [45] Blakely EL, He L, Taylor RW, et al. Mitochondrial DNA deletion in “identical” twin brothers. *J Med Genet* 2004;41:e19.
- [46] Chinnery PF, DiMauro S, Shanske S, et al. Risk of developing a mitochondrial DNA deletion disorder. *Lancet* 2004;364:592–6.
- [47] Barritt JA, Brenner CA, Cohen J, Matt DW. Mitochondrial DNA rearrangements in human oocytes and embryos. *Mol Hum Reprod* 1999;5:927–33.
- [48] Chan CCW, Liu VWS, Lau EYL, Yeung WSB, Ng EHY, Ho PC. Mitochondrial DNA content and 4977 bp deletion in unfertilized oocytes. *Mol Hum Reprod* 2005;11:843–6.
- [49] Elson JL, Apabhahi S, Gorman G, Whittaker RG, Krishnan KJ. Older mothers are not at risk of having grandchildren with sporadic mtDNA deletions. *Genet Med* 2010;12:313–4.
- [50] Hauswirth WW, Lapis PJ. Mitochondrial DNA polymorphism in a maternal lineage of Holstein cows. *Proc Nat Acad Sci USA* 1982;79:4686–90.
- [51] Cree LM, Samuels DC, De Sousa Lopes SC, et al. A reduction of mitochondrial DNA molecules during embryogenesis explains the rapid segregation of genotypes. *Nat Genet* 2008;40:249–54.
- [52] Wai T, Teoli D, Shoubridge EA. The mitochondrial DNA genetic bottleneck results from replication of a subpopulation of genomes. *Nat Genet* 2008;40:1484–8.
- [53] Cao L, Shitara H, Horii T, et al. The mitochondrial bottleneck occurs without reduction of mtDNA content in female mouse germ cells. *Nat Genet* 2007;39:386–90.
- [54] Cao L, Shitara H, Sugimoto M, Hayashi J-I, Abe K, Yonekawa H. New evidence confirms that the mitochondrial bottleneck is generated without reduction of mitochondrial DNA content in early primordial germ cells of mice. *PLoS Genet* 2009;5:e1000756.
- [55] Rötig A, Bourgeron T, Chretien D, Rustin P, Munnich A. Spectrum of mitochondrial DNA rearrangements in the Pearson marrow-pancreas syndrome. *Hum Mol Genet* 1995;4:1327–30.
- [56] Yamashita S, Nishino I, Nonaka I, Goto Y-I. Genotype and phenotype analyses in 136 patients with single large-scale mitochondrial DNA deletions. *J Hum Genet* 2008;53:598–606.
- [57] López-Gallardo E, López-Pérez MJ, Montoya J, Ruiz-Pesini E. CPEO and KSS differ in the percentage and location of the mtDNA deletion. *Mitochondrion* 2009;9:314–7.
- [58] Macmillan C, Lach B, Shoubridge EA. Variable distribution of mutant mitochondrial DNAs (tRNA[Leu<sup>3243</sup>]) in tissues of symptomatic relatives with MELAS: the role of mitotic segregation. *Neurology* 1993;43:1586–90.
- [59] Larsson NG, Tulinius MH, Holme E, et al. Segregation and manifestations of the mtDNA tRNA(Lys) A → G(8344) mutation of myoclonus epilepsy and ragged-red fibers (MERRF) syndrome. *Am J Hum Genet* 1992;51:1201–12.
- [60] Zeviani M, Moraes CT, DiMauro S, et al. Deletions of mitochondrial DNA in Kearns–Sayre syndrome. 1988. *Neurology* 1998;38:1339–46.
- [61] Moraes CT, DiMauro S, Zeviani M, et al. Mitochondrial DNA deletions in progressive external ophthalmoplegia and Kearns–Sayre syndrome. *N Engl J Med* 1989;320:1293–9.
- [62] Pearson HA, Lobel JS, Kocoshis SA, et al. A new syndrome of refractory sideroblastic anemia with vacuolization of marrow precursors and exocrine pancreatic dysfunction. *J Pediatr* 1979;95:976–84.
- [63] Rötig A, Colonna M, Bonnefont JP, et al. Mitochondrial DNA deletion in Pearson’s marrow/pancreas syndrome. *Lancet* 1989;1:902–3.
- [64] Rötig A, Cormier V, Blanche S, et al. Pearson’s marrow-pancreas syndrome. A multisystem mitochondrial disorder in infancy. *J Clin Invest* 1990;86:1601–8.
- [65] Manea EM, Leverger G, Bellmann F, et al. Pearson syndrome in the neonatal period: two case reports and review of the literature. *J Pediatr Hematol Oncol* 2009;31:947–51.
- [66] McShane MA, Hammans SR, Sweeney M, et al. Pearson syndrome and mitochondrial encephalomyopathy in a patient with a deletion of mtDNA. *Am J Hum Genet* 1991;48:39–42.
- [67] Santorelli FM, Barmada MA, Pons R, Zhang LL, DiMauro S. Leigh-type neuropathology in Pearson syndrome associated with impaired ATP production and a novel mtDNA deletion. *Neurology* 1996;47:1320–3.
- [68] Lee H-F, Lee H-J, Chi C-S, Tsai C-R, Chang T-K, Wang C-J. The neurological evolution of Pearson syndrome: case report and literature review. *Eur J Paediatr Neurol* 2007;11:208–14.
- [69] Rahman S, Leonard JV. Early onset of complete heart block in Pearson syndrome. *J Inherit Metab Dis* 2000;23:753–4.
- [70] Binder V, Steenpass L, Laws H-J, Ruebo J, Borkhardt A. A novel mtDNA large-scale mutation clinically exclusively presenting with refractory anemia: is there a chance to predict disease progression? *J Pediatr Hematol/Oncol*. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22246159>; 2012 [accessed 10.02.2012].
- [71] Bianchi M, Rizza T, Verrigni D, et al. Novel large-range mitochondrial DNA deletions and fatal multisystemic disorder with prominent hepatopathy. *Biochem Biophys Res Commun* 2011;415:300–4.
- [72] Morris AA, Lamont PJ, Clayton PT. Pearson’s syndrome without marrow involvement. *Arch Dis Child* 1997;77:56–7.
- [73] Atale A, Bonneau-Amati P, Rötig A, et al. Tubulopathy and pancytopenia with normal pancreatic function: a variant of Pearson syndrome. *Eur J Med Genet* 2009;52:23–6.
- [74] Lee YS, Yap HK, Barshop BA, Lee YS, Rajalingam S, Loke KY. Mitochondrial tubulopathy: the many faces of mitochondrial disorders. *Pediatr Nephrol* 2001;16:710–2.
- [75] Chae J-H, Lim BC, Cheong HI, Hwang YS, Kim KJ, Hwang H. A single large-scale deletion of mtDNA in a child with recurrent encephalopathy and tubulopathy. *J Neurol Sci* 2010;292:104–6.
- [76] Kearns TP, Sayre GP. Retinitis pigmentosa, external ophthalmoplegia, and complete heart block: unusual syndrome with histologic study in one of two cases. *AMA Arch Ophthalmol* 1958;60:280–9.
- [77] Kearns TP. External ophthalmoplegia, pigmentary degeneration of the retina, and cardiomyopathy: a newly recognized syndrome. *Trans Am Ophthalmol Soc* 1965;63:559–625.
- [78] Rowland LP. Progressive external ophthalmoplegia. In: *Handbook of clinical neurology. Part II: System disorders and atrophies*. New York: American Elsevier Publishing; 1975. p. 177–202.

- [79] Berenberg RA, Pellock JM, DiMauro S, et al. Lumping or splitting? “Ophthalmoplegia-plus” or Kearns–Sayre syndrome? *Ann Neurol* 1977;1:37–54.
- [80] Rowland LP, Hays AP, DiMauro S, DeVivo DC, Behrens M. Diverse clinical disorders associated with morphological abnormalities of mitochondria. In: *Mitochondrial pathology in muscle disease*. Padua: Piccin Medical Books; 1983. p. 141–58.
- [81] Lerman-Sagie T, Leshinsky-Silver E, Waternberg N, Luckman Y, Lev D. White matter involvement in mitochondrial diseases. *Mol Genet Metab* 2005;84:127–36.
- [82] Wray SH, Provenzale JM, Johns DR, Thulborn KR. MR of the brain in mitochondrial myopathy. *AJNR Am J Neuroradiol* 1995;16:1167–73.
- [83] Saneto RP, Friedman SD, Shaw DW. Neuroimaging of mitochondrial disease. *Mitochondrion* 2008;8:396–413.
- [84] Chu BC, Terae S, Takahashi C, et al. MRI of the brain in the Kearns–Sayre syndrome: report of four cases and a review. *Neuroradiology* 1999;41:759–64.
- [85] Pitceathly RDS, Fassone E, Taanman J-W, et al. Kearns–Sayre syndrome caused by defective R1/p53R2 assembly. *J Med Genet* 2011;48:610–7.
- [86] Yu Wai Man CY, Chinnery PF, Griffiths PG. Extraocular muscles have fundamentally distinct properties that make them selectively vulnerable to certain disorders. *Neuromuscul Disord* 2005;15:17–23.
- [87] Greaves LC, Yu-Wai-Man P, Blakely EL, et al. Mitochondrial DNA defects and selective extraocular muscle involvement in CPEO. *Invest Ophthalmol Vis Sci* 2010;51:3340–6.
- [88] Hammans SR, Sweeney MG, Hanna MG, Brockington M, Morgan-Hughes JA, Harding AE. The mitochondrial DNA transfer RNA-Leu(UUR) A → G(3243) mutation. A clinical and genetic study. *Brain* 1995;118(Pt. 3):721–34.
- [89] t Hart LM, Jansen JJ, Lemkes HH, de Knijff P, Maassen JA. Heteroplasmy levels of a mitochondrial gene mutation associated with diabetes mellitus decrease in leucocyte DNA upon aging. *Hum Mutat* 1996;7:193–7.
- [90] Rahman S, Poulton J, Marchington D, Suomalainen A. Decrease of 3243 A → G mtDNA mutation from blood in MELAS syndrome: a longitudinal study. *Am J Hum Genet* 2001;68:238–40.
- [91] Pyle A, Taylor RW, Durham SE, et al. Depletion of mitochondrial DNA in leucocytes harbouring the 3243A → G mtDNA mutation. *J Med Genet* 2007;44:69–74.
- [92] Whittaker RG, Blackwood JK, Alston CL, et al. Urine heteroplasmy is the best predictor of clinical outcome in the m.3243A > G mtDNA mutation. *Neurology* 2009;72:568–9.
- [93] Blackwood JK, Whittaker RG, Blakely EL, Alston CL, Turnbull DM, Taylor RW. The investigation and diagnosis of pathogenic mitochondrial DNA mutations in human urothelial cells. *Biochem Biophys Res Commun* 2010;393:740–5.
- [94] Rahman S, Hanna MG. Diagnosis and therapy in neuromuscular disorders: diagnosis and new treatments in mitochondrial diseases. *J Neurol Neurosurg Psychiatr* 2009;80:943–53.
- [95] DiMauro S, Hirano M. Pathogenesis and treatment of mitochondrial disorders. *Adv Exp Med Biol* 2009;652:139–70.
- [96] DiMauro S. Pathogenesis and treatment of mitochondrial myopathies: recent advances. *Acta Myol* 2010;29:333–8.
- [97] Muller W, Reimers CD, Berninger T, et al. Coenzyme Q10 in ophthalmoplegia plus – a double blind cross over therapeutic trial. *J Neurol Sci* 1990;98(Suppl.):442.
- [98] Chen RS, Huang CC, Chu NS. Coenzyme Q10 treatment in mitochondrial encephalomyopathies. Short-term double-blind, crossover study. *Eur Neurol* 1997;37:212–8.
- [99] Glover EI, Martin J, Maher A, Thornhill RE, Moran GR, Tarnopolsky MA. A randomized trial of coenzyme Q10 in mitochondrial disorders. *Muscle Nerve* 2010;42:739–48.
- [100] Klopstock T, Querner V, Schmidt F, et al. A placebo-controlled crossover trial of creatine in mitochondrial diseases. *Neurology* 2000;55:1748–51.
- [101] De Stefano N, Matthews PM, Ford B, Genge A, Karpati G, Arnold DL. Short-term dichloroacetate treatment improves indices of cerebral metabolism in patients with mitochondrial disorders. *Neurology* 1995;45:1193–8.
- [102] Chinnery P, Majamaa K, Turnbull D, Thorburn D. Treatment for mitochondrial disorders. *Cochrane Database Syst Rev* 2006: CD004426.
- [103] Allen RJ, DiMauro S, Coulter DL, Papadimitriou A, Rothenberg SP. Kearns–Sayre syndrome with reduced plasma and cerebrospinal fluid folate. *Ann Neurol* 1983;13:679–82.
- [104] Serrano M, Garcia-Silva MT, Martin-Hernandez E, et al. Kearns–Sayre syndrome: cerebral folate deficiency, MRI findings and new cerebrospinal fluid biochemical features. *Mitochondrion* 2010;10:429–32.
- [105] Pineda M, Ormazabal A, López-Gallardo E, et al. Cerebral folate deficiency and leukoencephalopathy caused by a mitochondrial DNA deletion. *Ann Neurol* 2006;59:394–8.
- [106] Clark KM, Bindoff LA, Lightowers RN, et al. Reversal of a mitochondrial DNA defect in human skeletal muscle. *Nat Genet* 1997;16:222–4.
- [107] Taivassalo T, Gardner JL, Taylor RW, et al. Endurance training and detraining in mitochondrial myopathies due to single large-scale mtDNA deletions. *Brain* 2006;129:3391–401.
- [108] Murphy JL, Blakely EL, Schaefer AM, et al. Resistance training in patients with single, large-scale deletions of mitochondrial DNA. *Brain* 2008;131:2832–40.
- [109] Holt IJ, Harding AE, Cooper JM, et al. Mitochondrial myopathies: clinical and biochemical features of 30 patients with major deletions of muscle mitochondrial DNA. *Ann Neurol* 1989;26:699–708.
- [110] Campos Y, Garcia-Silva T, Barrionuevo CR, Cabello A, Muley R, Arenas J. Mitochondrial DNA deletion in a patient with mitochondrial myopathy, lactic acidosis, and stroke-like episodes (MELAS) and Fanconi’s syndrome. *Pediatr Neurol* 1995;13:69–72.
- [111] Pang CY, Huang CC, Yen MY, et al. Molecular epidemiologic study of mitochondrial DNA mutations in patients with mitochondrial diseases in Taiwan. *J Formos Med Assoc* 1999;98:326–34.
- [112] Ballinger SW, Shoffner JM, Hedaya EV, et al. Maternally transmitted diabetes and deafness associated with a 10.4 kb mitochondrial DNA deletion. *Nat Genet* 1992;1:11–5.
- [113] Isotani H, Fukumoto Y, Kawamura H, et al. Hypoparathyroidism and insulin-dependent diabetes mellitus in a patient with Kearns–Sayre syndrome harbouring a mitochondrial DNA deletion. *Clin Endocrinol (Oxf)* 1996;45:637–41.
- [114] Boles RG, Roe T, Senadheera D, Mahnovski V, Wong LJ. Mitochondrial DNA deletion with Kearns Sayre syndrome in a child with Addison disease. *Eur J Pediatr* 1998;157:643–7.
- [115] Bruno C, Minetti C, Tang Y, et al. Primary adrenal insufficiency in a child with a mitochondrial DNA deletion. *J Inherit Metab Dis* 1998;21:155–61.
- [116] Wong LJ. Recognition of mitochondrial DNA deletion syndrome with non-neuromuscular multisystemic manifestation. *Genet Med* 2001;3:399–404.
- [117] Campos Y, Martín MA, Caballero C, et al. Single large-scale mitochondrial DNA deletion in a patient with encephalopathy, cardiomyopathy, and prominent intestinal pseudo-obstruction. *Neuromuscul Disord* 2000;10:56–8.
- [118] Barshop BA, Nyhan WL, Naviaux RK, McGowan KA, Friedlander M, Haas RH. Kearns–Sayre syndrome presenting as 2-oxoadipic aciduria. *Mol Genet Metab* 2000;69:64–8.
- [119] Moslemi AR, Selimovic N, Bergh CH, Oldfors A. Fatal dilated cardiomyopathy associated with a mitochondrial DNA deletion. *Cardiology* 2000;94:68–71.
- [120] Kato M, Nakamura M, Ichiba M, et al. Mitochondrial DNA deletion mutations in patients with neuropsychiatric symptoms. *Neurosci Res* 2011;69:331–6.