

A uniformly derived catalogue of exoplanets from radial velocities

Morgan D. J. Hollis,^{1★} Sreekumar T. Balan,^{2★} Greg Lever^{1★} and Ofer Lahav^{1★}

¹*Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT*

²*Astrophysics Group, Cavendish Laboratory, J J Thomson Avenue, Cambridge CB3 0HE*

Accepted 2012 April 11. Received 2012 April 5; in original form 2011 December 21

ABSTRACT

A new catalogue of extrasolar planets is presented by re-analysing a selection of published radial velocity data sets using EXOFAST. All objects are treated on an equal footing within a Bayesian framework to give orbital parameters for 94 exoplanetary systems. Model selection (between one- and two-planet solutions) is then performed using both a visual flagging method and a standard chi-square analysis, with agreement between the two methods for 99 per cent of the systems considered. The catalogue is to be made available online, and this ‘proof of concept’ study may be maintained and extended in the future to incorporate all systems with publicly available radial velocity data, as well as transit and microlensing data.

Key words: methods: data analysis – astronomical data bases: miscellaneous – stars: individual – planetary systems.

1 INTRODUCTION

Since the discovery of the first extrasolar planet in 1995 (Mayor & Queloz 1995), research on extrasolar planets has undergone exponential expansion. A wide range of search methods have been developed during this period, resulting in the discovery of more than 700 planets to date, the majority of which have been found using the radial velocity (RV) method. Traditional data reduction methods use a periodogram (Lomb 1976; Scargle 1982) to fix the orbital period and then the Levenberg–Marquardt minimization (Levenberg 1944; Marquardt 1963) to fit the other orbital parameters. A catalogue of exoplanets has already been published by Butler et al. (2006) using this method to extract the orbital parameters of exoplanets. Recently, Bayesian Markov chain Monte Carlo (MCMC) methods have been introduced by Gregory (2005), Ford (2005) and Ford & Gregory (2007) as a replacement for the traditional data reduction pipeline. EXOFAST (Balan & Lahav 2009) is a freely available tool for estimating orbital parameters of extrasolar planets from RV data using a Bayesian framework. Here we analyse 94 previously published data sets using EXOFAST, forming a new, uniformly derived catalogue of exoplanets from a Bayesian perspective.

Statistical properties of the distributions of orbital parameters are critical for explaining the planetary formation process. It has been argued that there is now a statistically significant number of known companions to make inferences about the correlations between orbital elements. Early discussions on this subject can be found in a series of articles on the statistical properties of exoplanets by Udry, Mayor & Santos (2003), Santos et al. (2003), Eggenberger, Udry & Mayor (2004) and Halbwegs, Mayor & Udry (2005). The

statistical discussion in this article is informed by the comparison to the published catalogues at <http://www.exoplanet.eu> (Schneider et al. 2011) and <http://exoplanets.org> (Wright et al. 2011).

The rest of this article is structured as follows. In Sections 2 and 3 the Bayesian framework and EXOFAST software package are introduced. The data analysis pipeline is described in Section 4, model selection is discussed in Section 5 and details of the prior distributions and boundaries used are shown in Tables 1–3. The catalogue is presented in Section 6 and in Tables A1–A3. The statistical properties of the distributions of various orbital parameters are discussed in Section 7 and the results are summarized in Section 8.

2 BAYESIAN FRAMEWORK

The Bayesian framework provides a transparent way of making probabilistic inferences from data. It is based on Bayes’ theorem, which states that for a given model H with a set of parameters Θ and data D , the posterior probability distribution of parameters $\Pr(\Theta|D, H)$ is proportional to the prior probability distribution $\Pr(\Theta|H)$ times the likelihood of data, $\Pr(D|\Theta, H)$. Using standard mathematical notation, one can write

$$\Pr(\Theta|D, H) = \frac{\Pr(D|\Theta, H) \Pr(\Theta|H)}{\Pr(D|H)}. \quad (1)$$

The denominator of the right-hand side of the above equation is called the Bayesian Evidence. Since it is the estimation of parameters that is of interest here, this term can be considered as a normalizing constant and equation (1) can be written as

$$\Pr(\Theta|D, H) \propto \Pr(D|\Theta, H) \Pr(\Theta|H). \quad (2)$$

The key step in the Bayesian approach is to obtain the posterior distribution of parameters accurately. The inferences are then derived from the posterior distribution. The MCMC method is a

*E-mail: mdjh@star.ucl.ac.uk (MDJH); st452@mrao.cam.ac.uk (STB); gl319@cam.ac.uk (GL); lahav@star.ucl.ac.uk (OL)

widely employed technique for simulating the posterior distribution (the left-hand side of equation 2). The basic steps in Bayesian parameter estimation can be summarized as follows:

- (i) model the observed data, i.e. construct the likelihood function;
- (ii) choose the prior probability distributions of parameters;
- (iii) obtain the posterior probability distribution;
- (iv) make inferences based on the posterior probability distribution.

EXOFIT is a software package that estimates the orbital parameters of extrasolar planets, following the steps outlined above. It should be noted that EXOFIT does not perform any Bayesian model selection – for a discussion of the relation of this aspect of the Bayesian framework to this study, the reader is directed to Section 5.

3 EXOFIT

EXOFIT (Balan & Lahav 2009) is a publicly available tool for extracting orbital parameters of exoplanets from RV measurements, using the MCMC method to simulate the posterior probability distribution of the orbital parameters.

The observed RV data are modelled by (Gregory 2005)

$$d_i = v_i + \epsilon_i + \delta, \quad (3)$$

where v_i is the true RV of the star and ϵ_i is the accounted measurement uncertainty. The term δ explains any unknown error and is chosen to be Gaussian with finite variance s^2 . Thus the combination of uncertainties $\epsilon_i + \delta$ has a Gaussian distribution with variance equal to $\sigma_i^2 + s^2$.

The likelihood of data $\Pr(\mathbf{D}|\boldsymbol{\Theta}, H)$ in equation (2) then connects the mathematical model to the observed data – this is assumed to be Gaussian, and is given as in Gregory (2005) by

$$\Pr(\mathbf{D}|\boldsymbol{\Theta}, H) = A \exp \left[- \sum_{i=1}^N \frac{(d_i - v_i)^2}{2(\sigma_i^2 + s^2)} \right], \quad (4)$$

where

$$A = (2\pi)^{-N/2} \left[\prod_{i=1}^N (\sigma_i^2 + s^2)^{-1/2} \right]. \quad (5)$$

The prior probabilities are as used by Ford & Gregory (2007).

EXOFIT then generates samples from the posterior distributions of the orbital parameters in the mathematical model, which can be analysed with the aid of any statistical software. Details of the algorithmic structure of the code, including methods of controlling chain mixing and assessing convergence, can be found in Balan & Lahav (2009). For further information, the reader is directed to the EXOFIT user's guide.

4 DATA ANALYSIS

As of 2009 August 21, when the data sets were extracted from the literature (Butler et al. 2006, and references therein), there were 295 planetary systems detected using the RV method according to <http://www.exoplanet.eu>, with 346 individual planets in total. Some RV data sets from the literature have fewer than 10 entries and as such are not appropriate for use with Bayesian inference methods, as a RV data set will need to include at least half an orbital period of a potential planetary companion. Hence data sets with not enough measurements to give accurate orbital solutions were not included. Also, the RV data of any systems with more than two confirmed

planets were ignored since at present EXOFIT can only search for either one or two planets.

Many more and different RV data sets and stellar mass estimates are now available (though some not publicly), but for the sake of uniformity the original RV data were used (i.e. those publicly available, frozen as of 2009 August 21 when the original data were collected). At a later stage the results can be improved by updating the original data sets to those which are now available, as well as incorporating the data from the many hundreds of additional planets that have been detected since the start of this study.

To enable accurate calculation of the derivable orbital parameters, the masses as well as the RVs of the associated stars were needed. These values were all taken from the published literature at <http://www.exoplanet.eu>, frozen as of 2011 March 1. The input for EXOFIT is in the form of a simple text file with RV, uncertainty and the time of observation (in Julian Date format), where the RV values must be in m s^{-1} . The Julian Date of the observation is offset to zero within EXOFIT.

The publicly available statistical data analysis package, `R`, from the R Project for Statistical Computing (<http://www.r-project.org>), was used to analyse the output of EXOFIT. The output from `R` includes the mean, median and standard deviation of the orbital parameters extracted from the posterior distribution samples produced by EXOFIT. The modal values are also produced, but will only have significance in the event of the posterior having more than one peak. Posterior distribution plots can also be produced with `R`, and the marginal distributions of each parameter can be found by plotting a histogram of the samples from the posterior. The full posterior distribution is helpful in analysing correlations between various parameters. Even though parameter degeneracy is present in the orbital solutions, highly degenerate solutions are less common.

The calculation time required for EXOFIT depends on computational resources available to the user. It scales linearly with the number of RV entries input to the code, and also depends significantly on the ease with which EXOFIT can converge the data. If EXOFIT is presented with data from a non-converging posterior distribution, it will take much longer than a larger data set with convergent orbital solutions. In technical terms, the mean average calculation time using EXOFIT on 26 RV data sets ranging from 10 to 50 data entries for a one-planet search was 44 s per data entry. For a two-planet search using 30 data sets with between 11 and 256 entries, the calculation time increased to 3 min and 40 s per data entry. These times were achieved on a 2.80-GHz dual core Linux system. Multiple runs were performed in order to confirm the orbital solutions for each system – these analyses were carried out using the UCL Legion High Performance Computing Facility, details of which can be found at <http://www.ucl.ac.uk/isd/common/research-computing/services/legion-upgrade>.

5 MODEL SELECTION

One of the most challenging aspects of the statistical inference procedure is the model selection problem. For the analysis of the RV data, the question of model selection refers to the selection of the correct number of planets to fit the observed data. Ford & Gregory (2007) and Gregory (2007) employed thermodynamic integration for calculating the Evidence and selecting the optimal number of planets that fit the data. On the other hand, Feroz, Balan & Hobson (2011a,b) approached the situation as an object detection problem.

One of the most commonly employed model selection procedures makes use of the chi-square statistic. This is one of the most prominent methods for estimating the goodness of fit, and it has

been applied to many astronomical problems including the analysis of RV data (see e.g. Butler et al. 2006). Bayesian inference also offers a straightforward way of performing statistical model selection, based on equation (1) and the Evidence. Even though this approach is conceptually simple, its implementation is in general computationally expensive, and EXOFAST does not currently have the functionality to perform such Bayesian model selection.

Hence in this analysis, the traditional chi-square statistic was used as well as a visual flagging approach, and the relationship between the two is discussed in Section 8. The study was limited to one or two planets, as per the current capabilities of the code, but this may of course be extended in later studies. The rationale behind the visual flagging is that one can identify the poor fits to the data by comparing the predicted RV curves for the one-planet and the two-planet solutions. The method involves assigning a ‘visual quality flag’ by eye to each system, where ‘1’ signifies that the one-planet fit is best, ‘2’ means that the two-planet fit is best and ‘3’ means that both one- and two-planet solutions provide equally good (or equally bad) fits. The results of this classification are shown in Table A3, next to the number of planets currently confirmed to exist in that system, taken by comparing the values on both <http://www.exoplanet.eu> and <http://exoplanets.org> (as these catalogues do not agree with each other in some cases).

Table A3 also shows the log likelihood ratio of the reduced chi-square value of the one-planet fit to that of the two-planet fit, where the log likelihood ratio is defined as

$$R \equiv -\frac{1}{2} \left(\chi_{1p}^2 - \chi_{2p}^2 \right). \quad (6)$$

Hence a value of $R > 0$ indicates that the one-planet fit was best (had a smaller reduced chi-square value), and $R < 0$ indicates that the two-planet model provided the best fit to the data. For all but one system (HD 8574), every ‘1’ and ‘2’ quality flag assigned to the fits by eye was in agreement with the calculated value of R , endorsing this method of assignment by visual inspection (see Fig. 5). For only a few systems there were not sufficient degrees of freedom to calculate a value for R (due to e.g. only having 11 data points for the 12 parameters), denoted by ‘-’ in the table.

6 CATALOGUE OF EXTRASOLAR PLANETS

In this paper the catalogue of extrasolar planets generated using EXOFAST is presented in Tables A1 and A2. These contain the best estimates of the orbital parameters for both one- and two-planet fits for all systems analysed. The orbital parameters used to fit to the model were the systematic velocity offset of the data V , the orbital period of the planet T , the RV semi-amplitude K , the orbital eccentricity e , the argument of periastron ω and u , the fraction of the orbit from the point of periastron passage through which the planet has travelled up to the start time of the data.

The final parameter, s , as described in Section 3, is a measure of all extra signal and/or noise in the data after the planetary fits have been accounted for. Hence a high value could indicate the presence of an additional planet or noisy data due to stellar activity or the combined noise from all sources. The reader is referred to Balan & Lahav (2009) for a more complete description of this parameter, which is not considered in any more detail in this study.

The direct output values from EXOFAST shown in the tables are the medians of the parameter posterior distributions and the associated 68.3 per cent confidence regions. The other displayed derivable parameters of the systems (minimum mass and semimajor axis) were calculated by transforming the orbital parameter posteriors using

the standard relations

$$a_p = \frac{m_* a_* \sin i}{m_p \sin i} \quad (7)$$

and

$$m_p \sin i \approx \frac{K_* m_*^{2/3} T^{1/3} \sqrt{1-e^2}}{(2\pi G)^{1/3}}, \quad (8)$$

assuming $m_p \ll m_*$. The final values for these derivable quantities were again taken to be the medians with 68.3 per cent confidence regions and are also displayed in the tables.

6.1 Choice of priors

The prior distributions and ranges used for the initial analysis were as shown in Tables 1 and 2. The prior for the systematic velocity

Table 1. The assumed orbital parameter prior distributions and their boundaries for a one-planet model. The minimum and maximum values for the systematic velocity parameter were the mean value of the raw RVs for that data file minus 5000 m s^{-1} and plus 5000 m s^{-1} , respectively.

Parameter	Prior	Mathematical form	Min	Max
$V \text{ (m s}^{-1}\text{)}$	Uniform	$\frac{1}{V_{\max} - V_{\min}}$	-	-
$T_1 \text{ (d)}$	Jeffreys	$\frac{1}{T_1 \ln\left(\frac{T_{1,\max}}{T_{1,\min}}\right)}$	0.2	15 000
$K_1 \text{ (m s}^{-1}\text{)}$	Mod. Jeffreys	$\frac{(K_1 + K_{1,0})^{-1}}{\ln\left(\frac{K_{1,0} + K_{1,\max}}{K_{1,0}}\right)}$	0.0	2000
e_1	Uniform	1	0	1
ω_1	Uniform	$\frac{1}{2\pi}$	0	2π
u_1	Uniform	1	0	1
$s \text{ (m s}^{-1}\text{)}$	Mod. Jeffreys	$\frac{(s+s_0)^{-1}}{\ln\left(\frac{s_0+s_{\max}}{s_0}\right)}$	0	2000

Table 2. The assumed orbital parameter prior distributions and their boundaries for a two-planet model. The boundaries for V were as detailed previously.

Para.	Prior	Mathematical form	Min	Max
$V \text{ (m s}^{-1}\text{)}$	Uniform	$\frac{1}{V_{\max} - V_{\min}}$	-	-
$T_1 \text{ (d)}$	Jeffreys	$\frac{1}{T_1 \ln\left(\frac{T_{1,\max}}{T_{1,\min}}\right)}$	0.2	15 000
$K_1 \text{ (m s}^{-1}\text{)}$	Mod. Jeffreys	$\frac{(K_1 + K_{1,0})^{-1}}{\ln\left(\frac{K_{1,0} + K_{1,\max}}{K_{1,0}}\right)}$	0.0	2000
e_1	Uniform	1	0	1
ω_1	Uniform	$\frac{1}{2\pi}$	0	2π
u_1	Uniform	1	0	1
$T_2 \text{ (d)}$	Jeffreys	$\frac{1}{T_2 \ln\left(\frac{T_{2,\max}}{T_{2,\min}}\right)}$	0.2	15 000
$K_2 \text{ (m s}^{-1}\text{)}$	Mod. Jeffreys	$\frac{(K_2 + K_{2,0})^{-1}}{\ln\left(\frac{K_{2,0} + K_{2,\max}}{K_{2,0}}\right)}$	0.0	2000
e_2	Uniform	1	0	1
ω_2	Uniform	$\frac{1}{2\pi}$	0	2π
u_2	Uniform	1	0	1
$s \text{ (m s}^{-1}\text{)}$	Mod. Jeffreys	$\frac{(s+s_0)^{-1}}{\ln\left(\frac{s_0+s_{\max}}{s_0}\right)}$	0	2000

Table 3. RV data sets analysed with different period prior boundaries, as explained in Section 6.1.

System	Initial period value	Min period	Max period
ϵ Eri	2500	1800	3200
γ Cep	906	899.84	912.1
GJ 849	1900	1400	2400
GJ 86	15.7649	15.764 12	15.765 68
HAT-P-9	3.922 89	3.922 81	3.922 97
HD 118203	6.1335	6.1323	6.1347
HD 12661	262.71	262.54	262.88
HD 128311	924	913.4	934.6
HD 131664	1950	1868	2032
HD 142	350	342.8	357.2
HD 149143	4.072	4.058	4.086
HD 162020	8.428 20	8.428 088	8.428 312
HD 168443	58.1121	58.111 142	58.113 058
HD 169830	225.6	225.16	226.04
HD 183263	627	624.8	629.2
HD 187123	3.096 583	3.096 567 32	3.096 598 68
HD 189733	2.218 5757	2.218 5754	2.218 5760
HD 190360	2920	2862.2	2977.8
HD 202206	255.87	255.75	255.99
HD 20868	380.85	380.67	381.03
HD 209458	3.524 7486	3.524 747 84	3.524 749 36
HD 217107	7.126 82	7.126 7318	7.126 8882
HD 219828	3.833	3.807	3.859
HD 28185	379	375	383
HD 330075	3.387 73	3.387 57	3.387 89
HD 33636	2128	2111.6	2144.4
HD 38529	2146	2134.98	2157.02
HD 46375	3.023 57	3.023 44	3.023 7
HD 47536	430	0.2	860
HD 50499	2460	2384.2	2535.8
HD 5319	670	636	704
HD 68988	6.2771	6.276 68	6.277 52
HD 73267	1260	1246	1274
HD 74156	2520	2490	2550
HD 80606	111.4367	111.4359	111.4375
HD 86081	2.1375	2.1371	2.1379
HD 89307	2170	2094	2246
τ Boo	3.312 46	3.312 432	3.312 488
TrES-3	1.31	0.2	2.62
WASP-2	2.152 226	2.152 218	2.152 234
WASP-3	1.846 834	1.846 83	1.846 838
XO-1	3.941 53	3.941 476	3.941 584
XO-2	2.615 838	2.615 822	2.615 854
XO-4	4.125 083	4.125 075	4.125 091

was dependent on the system – the mean of the input RV data was calculated and used as the initial value, and the allowed range was 10 km s^{-1} symmetrically about this.

For some systems, different sets of prior boundaries were used in a second round of analysis – these stars and the prior boundaries applied are listed in Table 3, and systems which did not return good fits using the normal prior boundaries were rerun with these ‘tight’ priors. The initial value was set to the published value of the period, the maximum value to the initial value plus twice the published error and the minimum value to the initial value minus twice the published error; hence the period of the planet was constrained to be within a range given by

$$T \in [T_{\text{pub}} - 2\sigma_{\text{pub}}, T_{\text{pub}} + 2\sigma_{\text{pub}}], \quad (9)$$




Figure 1. The resulting marginal posterior distributions of the orbital parameters for a one-planet fit to the BD-17 63 data (a), and the corresponding RV curve (b).

where T_{pub} is the published value of the period and σ_{pub} is the published error on the period, both taken from <http://exoplanets.org> on 2011 August 1. The only two exceptions were HD 47536 and TrES-3, for which no errors were obtainable from the literature, and hence a range of $[0.2, 2T_{\text{pub}}]$ was used.

This approach was generally necessary for those systems (e.g. WASP and XO data sets) where the number of data points available at the time of selecting the data was low, thus requiring tighter priors to adequately constrain the solution. Further constraints may also be applied; for example, systems with near zero eccentricities require tight priors on the orbital parameters ω and u in order to avoid multimodal distributions (see Section 8), whilst systems with eccentricities close to unity need tight priors on the orbital period T in order to achieve convergence of the MCMC chains. Examples of the output of EXOFIT are shown in Figs 1 and 2.

6.2 Ambiguous systems

For some of the systems analysed, there is a clear trend in the RVs indicating the possibility of a second planet, but the data are not informative enough to properly constrain the orbital parameters of such an object. Plotting the resulting RV curve and judging by eye can help us to assess and distinguish between the one- and two-planet fits and evaluate the validity of the orbital solution, though





Figure 2. The resulting marginal posterior distributions of the orbital parameters for a two-planet fit to the HD 108874 data (a), and the corresponding RV curve (b).

such poorly constrained orbits will lead to large error bars on the estimates of the orbital parameters.

This did not always lead to a clear classification though, and many systems were then rerun with the tight priors on the period given in Table 3. Some of these ambiguities were caused by data which were poor or less accurate due to age or too noisy due to stellar jitter. Others were simply due to the correlation between ω and u or data not good enough to constrain these two parameters. This resulted in near-uniform posteriors for ω and u , and hence fits that match few of the measured data points as a result of being shifted in time. Estimates for minimum masses and semimajor axes derived from these results are still valid, however (providing reliable estimates for T , K and e are obtained, which was generally the case), as these values have no dependence on mean anomaly at epoch and the time evolution of the Keplerian orbit.

The class 3 (both one- and two-planet fits equally good or equally bad) systems, as introduced in Section 5, are those which were considered to be somewhat ambiguous even after being analysed with tighter priors. This category was subdivided further – in some





Figure 3. The resulting RV curve fits using the derived orbital parameters for HD 89307. Imposing a two-planet model on data with only one planet can have the result shown here, where the two-planet fit (b) is the same as the one-planet fit (a) with a superimposed artificial small-amplitude periodic variation.

cases these are distinct RV solutions which provide plausible fits for both one- and two-planet models, and are classified as ‘3a’. However, there are also systems where the ‘second planet’ fit just produces small-amplitude variations on the one-planet solution (see Fig. 3 for an example), or where the one- and two-planet fits are identical but the ‘second planet’ posteriors are peaks at very small values for T and K , and uniform for e , ω and u (i.e. there is no single solution for a second planet from these data). From this, an ‘Occam’s razor’ approach could be taken and the assumption made that the correct model for most of these ‘3b’ class systems is in fact the single planet one. In a few cases though there may truly be a second planet present, and the data used are simply not good enough to change the likelihoods of the parameters from the initial ‘no-knowledge’ (uniform prior) situation. Therefore, for all class 3 systems, better (or at least more up-to-date) data and more complete analyses (such as using the log likelihood ratio in more




Figure 4. Orbital parameter values taken from Butler et al. (2006), plotted against values yielded using EXOFIT. Plotted systems are only those that occur in both catalogues, and where EXOFIT gave unambiguous (either class 1 or 2) results. (a) Period (in d). (b) Semimajor axis (in au). (c) Minimum mass (in M_{Jup}). (d) Eccentricity.




Figure 5. The log likelihood ratio for each planetary system assigned to visual class 1 or 2 is shown in panel (a). Panel (b) shows the same data on a smaller scale, around the threshold at $R = 0$. Class 2 systems (open circles) are all below the chi-square ambiguity threshold, and class 1 systems (filled circles) are all above, with the single exception of HD 8574 (shown as a triangle), class 1 but located just below the threshold with a value of $R = -0.03$.

detail to narrow down the classification) are required to accurately determine the correct orbital solutions.

7 COMPARISON OF ORBITAL PARAMETERS

Fig. 4 shows values for specified orbital elements from the literature against values yielded using EXOFIT a sample of systems. The minimum mass, semimajor axis and period values all exhibit good correlations in general between the independently derived values and those in the published literature – this is unsurprising for the period as it has not been derived from other quantities. The eccen-

tricity, however, shows a greater spread than expected – whereas this uniformly derived catalogue is consistent in taking the median, the published values use, in general, different statistical measures to extract the final parameter values from varying analysis techniques. This highlights the value of using a consistent technique to build up reliable data bases of orbital parameters.

As minimum mass and semimajor axis values are themselves derived from period and eccentricity, any inaccuracies in algorithms used will propagate, and also present discrepancies in the values yielded using EXOFIT and are likely to amplify outliers in these plots. These outliers will be investigated in the future in order to assess the validity of the solutions.




Figure 6. Examples of bimodal and ambiguous posterior densities obtained from HD 49674 and HD 190228. (a) Posterior densities for a one-planet fit to the HD 49674 data, exhibiting some bimodality in the ω and u orbital parameter values. (b) Posterior densities for a two-planet fit to the HD 190228 data, showing ambiguity in the estimates for the ω , u and e values for one of the planets.

There are some discrepancies in the global distribution of parameter values between this catalogue and the published literature, especially for the eccentricity parameter. This may be partly due to poor or outdated data, and is almost certainly affected by the ubiquitous effects of certain parameter correlations (as explained in Section 8). These should be analysed in more detail in the future, and techniques developed to explore the parameter space more efficiently and minimize or eradicate such dependencies.

8 DISCUSSION

The primary objective of this article is to analyse RV data sets uniformly, using a single platform for the data analysis. Butler et al. (2006) produced a catalogue of extrasolar planets using traditional methods (using periodograms and Levenberg–Marquardt minimization). Here are analysed a selection of RV data sets using a Bayesian parameter estimation technique for extrasolar planets. However, a model selection criterion is required for completion of the statistical inference process, and for this purpose, as described in Section 5, a chi-square statistic was employed as well as a visual flagging technique. Inconclusive results are obtained for a few data sets, but from those analysed here it can be seen that both model selection methods perform well, agreeing in 99 per cent of the cases, as demonstrated in Fig. 5.

Investigating further, the chi-square values were found to depend on the point estimates of the orbital parameters used to construct the predicted RV curve. If the posterior distribution is unimodal, such an approach will work flawlessly. However, posterior distributions of the orbital parameters exhibit multimodality on many occasions. For example, the parameters ω and u are extremely correlated and their posterior distributions are bimodal for many data sets (an example of this is shown in Fig. 6), especially for planets with $e \approx 0$. This problem has also been noted by Gregory (2007), who proposed reparametrization of the problem as a possible way of dealing with this situation.

Many data sets contain planetary signals whose period is greater than the span of the observations, and so obtaining constraints on the orbital parameters of these objects is an extremely difficult task.

There are several data sets where it was possible to obtain estimates for the orbital parameters for one of the planets, but then the second signal could not be constrained due to weak signal-to-noise ratio. In most cases these signals appear to be a linear or quadratic trend in the RV data. Therefore, it becomes extremely difficult to classify these objects as planets, and this is one of the reasons why a visual flagging method was employed. One example of this is shown in Fig. 6, in the results of the two-planet fit to the data of the system HD 190228. The strongest signal is picked up and well constrained, as can be seen from the error bars in Table A2, and in addition the values for this planet match well those from the fit for a single planet. Thus the parameters of the first planet are reasonably certain, but those of the second, shown as HD 190228c in Table A2, are significantly less secure.

Additionally, sharp prior boundaries were used on the orbital period for several data sets. In these cases, either the planetary signal is very weak or the signal from a systematic trend from an additional companion in the RVs masks the weaker planetary signal. In addition, the aliasing effects (see e.g. Dawson & Fabrycky 2010) in observations can produce additional peaks in the posterior distributions, necessitating the use of the sharp prior on the period.

In summary, a brief overview of the Bayesian theory has been given here, along with a description of the MCMC approach used in order to estimate the orbital parameters of extrasolar planets, more details of which can be found in Balan & Lahav (2009). A new catalogue of extrasolar planets is presented from the re-analysis of published RV data sets, giving both one- and two-planet orbital solutions for 94 systems derived on a uniform basis. An attempt is made to distinguish between the solutions for each system by using both a visual categorization method and a standard reduced chi-square technique, giving good agreement in 99 per cent of the cases presented here. Improvements in this ‘model selection’ area of the analysis may be made by taking into account Bayesian Evidence, as seen in Gregory (2007) and Feroz et al. (2011a,b); more rigorous approaches such as these are outside the scope of this ‘proof of concept’ study, but may be looked into in the future. Other further work will include updating this catalogue to incorporate the most up-to-date data, as well as extending EXOFAST to be able to use transit

and microlensing results, to search for an arbitrary number of planets, and to look into the possibility of accounting for interactions between planetary bodies.

ACKNOWLEDGMENTS

The authors would like to thank Paul Gorman, Robert Michaelides, Lisa Menahem, Andrew Strang, Robert C. Clouth and Milroy Travasso for their help in analysing planetary data sets, and David Kipping for useful discussions and feedback. MDJH is supported by an Impact/Perren studentship; STB acknowledges support from the Isaac Newton Studentship and a studentship from the Astrophysics Group, Cavendish Laboratory; and OL acknowledges a Royal Society Wolfson Research Merit Award. The authors acknowledge the use of the UCL Legion High Performance Computing Facility, and associated support services, in the completion of this work. This research has made use of the Exoplanet Orbit Database and the Exoplanet Data Explorer at <http://exoplanets.org>, and the Extrasolar Planets Encyclopaedia at <http://www.exoplanet.eu>.

The catalogue will be made available for public viewing at <http://www.ucl.ac.uk/exoplanets/exocat>.

REFERENCES

- Balan S. T., Lahav O., 2009, MNRAS, 394, 1936
Butler R. P. et al., 2006, ApJ, 646, 505
Dawson R. I., Fabrycky D. C., 2010, ApJ, 722, 937

- EGGENBERGER A., UDRY S., MAYOR M., 2004, A&A, 417, 353
FEROZ F., BALAN S. T., HOBSON M. P., 2011a, MNRAS, 415, 3462
FEROZ F., BALAN S. T., HOBSON M. P., 2011b, MNRAS, 416, L104
FORD E. B., 2005, AJ, 129, 1706
FORD E. B., GREGORY P. C., 2007, in Babu G. J., Feigelson E. D., eds, ASP Conf. Ser. Vol. 371, Statistical Challenges in Modern Astronomy IV. Astron. Soc. Pac., San Francisco, p. 189
GREGORY P. C., 2005, Bayesian Logical Data Analysis for the Physical Sciences: A Comparative Approach with ‘Mathematica’ Support. Cambridge Univ. Press, Cambridge
GREGORY P. C., 2007, MNRAS, 381, 1607
HALBWACHS J. L., MAYOR M., UDRY S., 2005, A&A, 431, 1129
LEVENBERG K., 1944, Q. Applied Math., 2, 164
LOMB N. R., 1976, Ap&SS, 39, 447
MARQUARDT D., 1963, SIAM J. Applied Math., 11, 431
MAYOR M., QUELOZ D., 1995, Nat, 378, 355
SANTOS N. C., ISRAELIAN G., MAYOR M., REBOLLO R., UDRY S., 2003, A&A, 398, 363
SCARGLE J. D., 1982, ApJ, 263, 835
SCHNEIDER J., DEDIEU C., LE SIDANER P., SAVALLE R., ZOLOTUKHIN I., 2011, A&A, 532, A79
UDRY S., MAYOR M., SANTOS N. C., 2003, A&A, 407, 369
WRIGHT J. T. et al., 2011, PASP, 123, 412

APPENDIX A: A CATALOGUE OF EXOPLANET ORBITAL PARAMETERS FROM RADIAL VELOCITIES, UNIFORMLY DERIVED IN A BAYESIAN FRAMEWORK

Table A1. Table of the orbital parameters for a one-planet fit, both directly output from EXOFT and thence derived. The values of the parameters T , K , e and s (generated from EXOFT) are the medians of the parameter posterior distributions, with the associated 68.3 per cent confidence regions. The other parameters were calculated using these values and stellar masses taken from the published literature. Note that some parameters are extremely well constrained; hence, the errors on the parameter estimates are so small as to appear to be zero to the two decimal places shown in this table. A full table in machine-readable format will also be provided at <http://www.ucl.ac.uk/exoplanets/exocat>, and the reader is directed there if such data are required.

Planet	m_* (M_\odot)	T (d)	K (m s^{-1})	e	s	$m_p \sin(i)$ (M_{Jup})	a (au)
BD-17 63 b	0.74	655.49 ^{+0.59} _{-0.62}	172.44 ^{+0.62} _{-1.61}	0.54 ^{+0.01} _{-0.01}	4.59 ^{+0.95} _{-0.73}	5.06 ^{+0.05} _{-0.05}	1.34 ^{+0.00} _{-0.00}
ChaHa8 b	0.10	304.59 ^{+1.81} _{-1.79}	1221.89 ^{+186.78} _{-128.02}	0.15 ^{+0.15} _{-0.10}	32.67 ^{+130.79} _{-30.48}	8.60 ^{+1.12} _{-0.89}	0.41 ^{+0.00} _{-0.00}
ϵ Eri b	0.86	2503.68 ^{+57.36} _{-52.69}	17.83 ^{+1.93} _{-1.81}	0.16 ^{+0.16} _{-0.11}	9.44 ^{+0.91} _{-0.82}	1.05 ^{+0.10} _{-0.10}	3.43 ^{+0.05} _{-0.05}
ϵ Tau b	2.70	597.53 ^{+12.02} _{-11.52}	96.16 ^{+3.93} _{-3.85}	0.13 ^{+0.04} _{-0.04}	8.85 ^{+2.48} _{-1.95}	7.66 ^{+0.30} _{-0.30}	1.93 ^{+0.03} _{-0.02}
γ Cep b	1.59	905.03 ^{+4.52} _{-3.68}	317.34 ^{+77.57} _{-71.25}	0.51 ^{+0.14} _{-0.16}	225.74 ^{+34.05} _{-26.78}	17.44 ^{+4.02} _{-4.15}	2.14 ^{+0.01} _{-0.01}
GJ 3021 b	0.90	133.70 ^{+0.20} _{-0.20}	167.02 ^{+3.87} _{-3.95}	0.51 ^{+0.02} _{-0.02}	15.86 ^{+2.34} _{-2.05}	3.36 ^{+0.08} _{-0.08}	0.49 ^{+0.00} _{-0.00}
GJ 317 b	0.24	672.33 ^{+8.26} _{-7.27}	90.96 ^{+46.14} _{-12.10}	0.45 ^{+0.20} _{-0.10}	15.63 ^{+4.38} _{-3.16}	1.36 ^{+0.40} _{-0.16}	0.93 ^{+0.01} _{-0.01}
GJ 674 b	0.35	4.69 ^{+0.00} _{-0.00}	9.50 ^{+0.99} _{-1.02}	0.11 ^{+0.10} _{-0.08}	3.55 ^{+0.58} _{-0.46}	0.04 ^{+0.00} _{-0.00}	0.04 ^{+0.00} _{-0.00}
GJ 849 b	0.49	2014.09 ^{+60.32} _{-61.27}	26.68 ^{+9.48} _{-4.46}	0.68 ^{+0.10} _{-0.09}	7.14 ^{+1.40} _{-1.08}	0.77 ^{+0.19} _{-0.11}	2.46 ^{+0.05} _{-0.05}
GJ 86 b	0.80	15.77 ^{+0.00} _{-0.00}	431.19 ^{+61.11} _{-59.29}	0.23 ^{+0.11} _{-0.12}	204.72 ^{+26.76} _{-21.68}	4.44 ^{+0.59} _{-0.59}	0.11 ^{+0.00} _{-0.00}
HAT-P-6 b	1.29	3.85 ^{+0.00} _{-0.00}	115.69 ^{+3.99} _{-4.17}	0.04 ^{+0.04} _{-0.03}	8.73 ^{+3.25} _{-2.46}	1.06 ^{+0.04} _{-0.04}	0.05 ^{+0.00} _{-0.00}
HAT-P-8 b	1.28	3.09 ^{+0.00} _{-0.00}	162.59 ^{+7.36} _{-6.46}	0.05 ^{+0.05} _{-0.03}	6.56 ^{+4.14} _{-3.02}	1.37 ^{+0.06} _{-0.05}	0.05 ^{+0.00} _{-0.00}
HAT-P-9 b	1.30	3.92 ^{+0.00} _{-0.00}	84.50 ^{+10.56} _{-9.37}	0.12 ^{+0.14} _{-0.09}	4.09 ^{+9.61} _{-3.40}	0.77 ^{+0.09} _{-0.09}	0.05 ^{+0.00} _{-0.00}
HD 101930 b	0.74	70.58 ^{+0.40} _{-0.37}	17.99 ^{+0.89} _{-0.91}	0.08 ^{+0.05} _{-0.05}	1.92 ^{+0.65} _{-0.46}	0.30 ^{+0.01} _{-0.02}	0.30 ^{+0.00} _{-0.00}
HD 108874 b	0.95	395.16 ^{+5.60} _{-4.43}	34.93 ^{+3.82} _{-3.57}	0.05 ^{+0.08} _{-0.04}	13.00 ^{+1.65} _{-1.38}	1.21 ^{+0.13} _{-0.12}	1.04 ^{+0.01} _{-0.01}
HD 11506 b	1.19	1456.01 ^{+136.42} _{-85.10}	81.49 ^{+13.56} _{-4.62}	0.37 ^{+0.16} _{-0.10}	10.60 ^{+2.06} _{-1.60}	4.76 ^{+0.46} _{-0.23}	2.66 ^{+0.16} _{-0.10}
HD 118203 b	1.23	6.13 ^{+0.00} _{-0.00}	213.96 ^{+6.49} _{-6.40}	0.30 ^{+0.03} _{-0.03}	22.83 ^{+3.88} _{-3.28}	2.11 ^{+0.06} _{-0.06}	0.07 ^{+0.00} _{-0.00}
HD 12661 b	1.14	262.75 ^{+0.09} _{-0.13}	77.37 ^{+2.52} _{-2.55}	0.27 ^{+0.03} _{-0.03}	17.60 ^{+1.38} _{-1.24}	2.56 ^{+0.08} _{-0.09}	0.84 ^{+0.00} _{-0.00}
HD 128311 c	0.83	921.18 ^{+6.65} _{-5.13}	93.88 ^{+7.75} _{-7.27}	0.46 ^{+0.05} _{-0.05}	30.34 ^{+2.80} _{-2.42}	3.51 ^{+0.24} _{-0.24}	1.74 ^{+0.01} _{-0.01}
HD 131664 b	1.10	1976.18 ^{+32.94} _{-41.05}	356.10 ^{+24.90} _{-18.59}	0.64 ^{+0.02} _{-0.02}	5.11 ^{+0.79} _{-0.66}	18.03 ^{+0.85} _{-0.65}	3.18 ^{+0.04} _{-0.04}

Table A1 – *continued*

Planet	m_* (M_\odot)	T (d)	K (m s^{-1})	e	s	$m_p \sin(i)$ (M_{Jup})	a (au)
HD 132406 b	1.09	1172.21 ^{+75.55} _{-49.55}	122.19 ^{+157.18} _{-32.90}	0.34 ^{+0.28} _{-0.19}	17.04 ^{+4.70} _{-3.61}	6.31 ^{+5.80} _{-1.47}	2.24 ^{+0.10} _{-0.06}
HD 142 b	1.23	344.05 ^{+2.12} _{-0.93}	32.00 ^{+7.12} _{-6.14}	0.19 ^{+0.16} _{-0.13}	20.29 ^{+2.53} _{-2.15}	1.24 ^{+0.24} _{-0.23}	1.03 ^{+0.00} _{-0.00}
HD 142022 b	0.90	1861.66 ^{+14.86} _{-13.47}	140.10 ^{+112.02} _{-39.74}	0.64 ^{+0.12} _{-0.09}	3.00 ^{+1.67} _{-1.29}	6.10 ^{+3.21} _{-1.36}	2.86 ^{+0.02} _{-0.01}
HD 149143 b	1.20	4.07 ^{+0.00} _{-0.00}	149.71 ^{+1.67} _{-1.61}	0.01 ^{+0.01} _{-0.01}	1.21 ^{+1.69} _{-0.92}	1.33 ^{+0.01} _{-0.01}	0.05 ^{+0.00} _{-0.00}
HD 154345 b	0.89	3332.50 ^{+84.05} _{-74.54}	14.10 ^{+0.84} _{-0.85}	0.05 ^{+0.05} _{-0.04}	2.84 ^{+0.37} _{-0.32}	0.96 ^{+0.06} _{-0.06}	4.20 ^{+0.07} _{-0.06}
HD 155358 b	0.87	194.26 ^{+0.88} _{-0.80}	31.86 ^{+1.98} _{-1.97}	0.21 ^{+0.06} _{-0.06}	9.69 ^{+1.03} _{-0.89}	0.81 ^{+0.05} _{-0.05}	0.63 ^{+0.00} _{-0.00}
HD 162020 b	0.80	8.43 ^{+0.00} _{-0.00}	1808.97 ^{+5.15} _{-5.13}	0.28 ^{+0.00} _{-0.00}	11.13 ^{+2.65} _{-2.39}	15.01 ^{+0.04} _{-0.04}	0.08 ^{+0.00} _{-0.00}
HD 168443 b	1.01	58.11 ^{+0.00} _{-0.00}	510.46 ^{+252.18} _{-117.22}	0.52 ^{+0.20} _{-0.18}	220.77 ^{+46.40} _{-34.29}	8.28 ^{+2.91} _{-1.94}	0.29 ^{+0.00} _{-0.00}
HD 169830 b	1.41	225.62 ^{+0.29} _{-0.31}	83.07 ^{+3.05} _{-3.09}	0.37 ^{+0.03} _{-0.03}	1.52 ^{+2.40} _{-1.18}	2.91 ^{+0.10} _{-0.10}	0.81 ^{+0.00} _{-0.00}
HD 171028 b	0.99	545.13 ^{+10.10} _{-12.19}	59.75 ^{+3.04} _{-2.05}	0.59 ^{+0.02} _{-0.02}	2.55 ^{+0.70} _{-0.51}	1.92 ^{+0.13} _{-0.10}	1.30 ^{+0.02} _{-0.02}
HD 183263 b	1.12	627.80 ^{+1.03} _{-1.64}	89.99 ^{+13.01} _{-11.63}	0.42 ^{+0.08} _{-0.09}	26.59 ^{+3.48} _{-2.91}	3.72 ^{+0.42} _{-0.41}	1.49 ^{+0.00} _{-0.00}
HD 185269 b	1.30	6.84 ^{+0.00} _{-0.00}	89.57 ^{+4.12} _{-4.02}	0.28 ^{+0.03} _{-0.04}	7.72 ^{+1.92} _{-1.61}	0.96 ^{+0.04} _{-0.04}	0.08 ^{+0.00} _{-0.00}
HD 187123 b	1.04	3.10 ^{+0.00} _{-0.00}	65.68 ^{+3.34} _{-3.35}	0.05 ^{+0.06} _{-0.04}	18.33 ^{+1.83} _{-1.58}	0.48 ^{+0.02} _{-0.03}	0.04 ^{+0.00} _{-0.00}
HD 189733 b	0.81	2.22 ^{+0.00} _{-0.00}	204.58 ^{+5.15} _{-5.10}	0.01 ^{+0.01} _{-0.01}	15.65 ^{+1.33} _{-1.17}	1.14 ^{+0.03} _{-0.03}	0.03 ^{+0.00} _{-0.00}
HD 190228 b	1.82	1141.21 ^{+15.40} _{-14.66}	92.26 ^{+4.58} _{-3.48}	0.52 ^{+0.04} _{-0.04}	1.23 ^{+1.87} _{-0.94}	6.07 ^{+0.17} _{-0.15}	2.61 ^{+0.02} _{-0.02}
HD 190360 b	0.98	2925.83 ^{+36.05} _{-41.53}	19.38 ^{+2.67} _{-2.28}	0.33 ^{+0.11} _{-0.10}	5.92 ^{+1.47} _{-1.47}	1.27 ^{+0.14} _{-0.13}	3.98 ^{+0.03} _{-0.04}
HD 190647 b	1.10	1038.09 ^{+5.27} _{-5.38}	36.78 ^{+1.19} _{-1.17}	0.17 ^{+0.02} _{-0.02}	0.97 ^{+0.70} _{-0.65}	1.92 ^{+0.06} _{-0.06}	2.07 ^{+0.01} _{-0.01}
HD 195019 b	1.02	18.20 ^{+0.00} _{-0.00}	270.12 ^{+1.54} _{-1.55}	0.01 ^{+0.01} _{-0.01}	10.42 ^{+1.19} _{-1.14}	3.54 ^{+0.02} _{-0.02}	0.14 ^{+0.00} _{-0.00}
HD 202206 b	1.07	255.90 ^{+0.06} _{-0.09}	585.94 ^{+6.24} _{-6.13}	0.42 ^{+0.01} _{-0.01}	30.18 ^{+2.61} _{-2.31}	17.41 ^{+0.16} _{-0.16}	0.81 ^{+0.00} _{-0.00}
HD 20868 b	0.78	380.79 ^{+0.13} _{-0.09}	97.02 ^{+7.97} _{-7.95}	0.61 ^{+0.04} _{-0.04}	32.81 ^{+4.04} _{-3.35}	2.31 ^{+0.18} _{-0.18}	0.95 ^{+0.00} _{-0.00}
HD 209458 b	1.13	3.52 ^{+0.00} _{-0.00}	84.33 ^{+0.87} _{-0.87}	0.01 ^{+0.01} _{-0.01}	3.34 ^{+0.69} _{-0.67}	0.69 ^{+0.01} _{-0.01}	0.05 ^{+0.00} _{-0.00}
HD 212301 b	1.05	2.27 ^{+0.00} _{-0.00}	56.31 ^{+5.83} _{-5.94}	0.08 ^{+0.08} _{-0.05}	17.70 ^{+3.65} _{-2.73}	0.37 ^{+0.04} _{-0.04}	0.03 ^{+0.00} _{-0.00}
HD 217107 b	1.11	7.13 ^{+0.00} _{-0.00}	140.71 ^{+2.35} _{-2.41}	0.15 ^{+0.02} _{-0.02}	22.70 ^{+1.23} _{-1.12}	1.41 ^{+0.02} _{-0.02}	0.08 ^{+0.00} _{-0.00}
HD 219828 b	1.24	3.84 ^{+0.01} _{-0.02}	3.11 ^{+7.83} _{-2.55}	0.58 ^{+0.32} _{-0.39}	15.98 ^{+2.97} _{-2.27}	0.02 ^{+0.05} _{-0.02}	0.05 ^{+0.00} _{-0.00}
HD 221287 b	1.30	458.77 ^{+8.13} _{-6.20}	69.77 ^{+8.66} _{-6.77}	0.11 ^{+0.10} _{-0.07}	10.16 ^{+1.97} _{-1.50}	3.14 ^{+0.38} _{-0.32}	1.27 ^{+0.01} _{-0.01}
HD 224693 b	1.30	26.73 ^{+0.03} _{-0.03}	39.92 ^{+1.52} _{-1.53}	0.04 ^{+0.04} _{-0.03}	1.92 ^{+1.07} _{-1.10}	0.70 ^{+0.03} _{-0.03}	0.19 ^{+0.00} _{-0.00}
HD 23127 b	1.13	1226.63 ^{+21.59} _{-21.71}	27.75 ^{+3.08} _{-2.84}	0.44 ^{+0.09} _{-0.10}	10.89 ^{+2.03} _{-1.67}	1.42 ^{+0.17} _{-0.16}	2.34 ^{+0.03} _{-0.03}
HD 2638 b	0.93	3.44 ^{+0.00} _{-0.00}	67.59 ^{+1.06} _{-1.02}	0.01 ^{+0.01} _{-0.01}	3.31 ^{+0.70} _{-0.57}	0.48 ^{+0.01} _{-0.01}	0.04 ^{+0.00} _{-0.00}
HD 27442 b	1.20	415.32 ^{+6.25} _{-5.74}	32.48 ^{+1.79} _{-1.76}	0.07 ^{+0.06} _{-0.04}	2.98 ^{+1.41} _{-1.13}	1.34 ^{+0.07} _{-0.07}	1.16 ^{+0.01} _{-0.01}
HD 27894 b	0.75	18.01 ^{+0.02} _{-0.01}	57.01 ^{+1.61} _{-1.66}	0.04 ^{+0.03} _{-0.02}	4.39 ^{+1.08} _{-0.83}	0.61 ^{+0.02} _{-0.02}	0.12 ^{+0.00} _{-0.00}
HD 28185 b	0.99	381.81 ^{+0.83} _{-1.32}	174.72 ^{+12.09} _{-7.75}	0.05 ^{+0.02} _{-0.02}	7.82 ^{+1.76} _{-1.53}	6.19 ^{+0.43} _{-0.28}	1.03 ^{+0.00} _{-0.00}
HD 285968 b	0.49	10.23 ^{+0.00} _{-0.00}	11.88 ^{+2.24} _{-1.79}	0.25 ^{+0.20} _{-0.17}	2.47 ^{+1.77} _{-1.74}	0.08 ^{+0.01} _{-0.01}	0.07 ^{+0.00} _{-0.00}
HD 330075 b	0.70	3.39 ^{+0.00} _{-0.00}	107.34 ^{+1.00} _{-1.03}	0.01 ^{+0.01} _{-0.00}	2.02 ^{+0.86} _{-0.74}	0.63 ^{+0.01} _{-0.01}	0.04 ^{+0.00} _{-0.00}
HD 33636 b	1.02	2127.74 ^{+11.40} _{-11.04}	389.98 ^{+156.81} _{-152.41}	0.90 ^{+0.03} _{-0.11}	0.57 ^{+0.69} _{-0.42}	11.02 ^{+2.19} _{-1.89}	3.26 ^{+0.01} _{-0.01}
HD 3651 b	0.88	60.36 ^{+0.04} _{-0.05}	9.60 ^{+1.91} _{-1.60}	0.54 ^{+0.15} _{-0.16}	7.73 ^{+0.68} _{-0.61}	0.14 ^{+0.02} _{-0.02}	0.29 ^{+0.00} _{-0.00}
HD 38529 c	1.48	2143.62 ^{+8.24} _{-6.24}	177.12 ^{+6.26} _{-6.00}	0.35 ^{+0.03} _{-0.03}	40.24 ^{+2.46} _{-2.22}	13.65 ^{+0.42} _{-0.42}	3.71 ^{+0.01} _{-0.01}
HD 4203 b	1.13	434.21 ^{+2.65} _{-1.97}	74.24 ^{+88.38} _{-18.24}	0.71 ^{+0.14} _{-0.09}	5.63 ^{+1.49} _{-1.18}	2.15 ^{+1.42} _{-0.39}	1.17 ^{+0.00} _{-0.00}
HD 4208 b	0.88	828.90 ^{+8.03} _{-7.83}	19.01 ^{+0.68} _{-0.70}	0.05 ^{+0.04} _{-0.03}	1.26 ^{+0.86} _{-0.85}	0.81 ^{+0.03} _{-0.03}	1.65 ^{+0.01} _{-0.01}
HD 43691 b	1.38	36.93 ^{+0.04} _{-0.04}	123.78 ^{+4.76} _{-4.65}	0.11 ^{+0.05} _{-0.05}	14.47 ^{+3.07} _{-2.51}	2.50 ^{+0.10} _{-0.10}	0.24 ^{+0.00} _{-0.00}
HD 43848 b	0.93	2390.59 ^{+121.57} _{-85.08}	888.17 ^{+447.92} _{-286.53}	0.81 ^{+0.06} _{-0.08}	5.58 ^{+4.19} _{-2.90}	33.07 ^{+9.51} _{-7.29}	3.42 ^{+0.11} _{-0.08}
HD 46375 b	0.93	3.02 ^{+0.00} _{-0.00}	33.67 ^{+0.79} _{-0.80}	0.06 ^{+0.03} _{-0.03}	3.30 ^{+0.59} _{-0.52}	0.23 ^{+0.01} _{-0.01}	0.04 ^{+0.00} _{-0.00}
HD 47536 b	0.94	717.27 ^{+15.06} _{-13.33}	108.71 ^{+12.96} _{-12.35}	0.15 ^{+0.10} _{-0.09}	6.65 ^{+9.40} _{-5.62}	4.53 ^{+0.57} _{-0.56}	1.54 ^{+0.02} _{-0.02}
HD 49674 b	1.01	4.95 ^{+0.00} _{-0.00}	11.76 ^{+1.09} _{-1.13}	0.05 ^{+0.07} _{-0.04}	3.65 ^{+0.78} _{-0.69}	0.10 ^{+0.01} _{-0.01}	0.06 ^{+0.00} _{-0.00}
HD 50499 b	1.28	2460.46 ^{+51.66} _{-51.79}	25.77 ^{+5.54} _{-4.56}	0.29 ^{+0.21} _{-0.19}	13.38 ^{+2.08} _{-1.71}	1.91 ^{+0.33} _{-0.32}	3.87 ^{+0.05} _{-0.05}
HD 5319 b	1.60	684.51 ^{+12.52} _{-16.78}	36.23 ^{+17.09} _{-5.87}	0.10 ^{+0.20} _{-0.07}	10.44 ^{+1.73} _{-1.37}	2.13 ^{+0.91} _{-0.35}	1.78 ^{+0.02} _{-0.03}
HD 63454 b	0.80	2.82 ^{+0.00} _{-0.00}	63.32 ^{+1.76} _{-1.81}	0.02 ^{+0.03} _{-0.02}	5.80 ^{+1.25} _{-0.98}	0.38 ^{+0.01} _{-0.01}	0.04 ^{+0.00} _{-0.00}
HD 68988 b	1.12	6.28 ^{+0.00} _{-0.00}	183.95 ^{+13.27} _{-13.39}	0.09 ^{+0.06} _{-0.06}	42.29 ^{+7.33} _{-5.71}	1.79 ^{+0.12} _{-0.13}	0.07 ^{+0.00} _{-0.00}
HD 73267 b	0.89	1259.62 ^{+6.47} _{-6.80}	64.28 ^{+0.44} _{-0.46}	0.26 ^{+0.01} _{-0.01}	0.72 ^{+0.50} _{-0.48}	3.06 ^{+0.02} _{-0.02}	2.20 ^{+0.01} _{-0.01}
HD 73526 b	1.01	193.32 ^{+0.75} _{-1.05}	114.79 ^{+22.33} _{-18.22}	0.57 ^{+0.07} _{-0.09}	25.52 ^{+4.75} _{-3.69}	2.70 ^{+0.40} _{-0.35}	0.66 ^{+0.00} _{-0.00}
HD 74156 c	1.24	2519.62 ^{+20.55} _{-20.33}	120.88 ^{+139.67} _{-42.57}	0.89 ^{+0.08} _{-0.13}	54.09 ^{+4.69} _{-4.09}	4.31 ^{+1.34} _{-0.68}	3.89 ^{+0.02} _{-0.02}

Table A1 – continued

Planet	m_* (M_\odot)	T (d)	K (m s^{-1})	e	s	$m_p \sin(i)$ (M_{Jup})	a (au)
HD 75289 b	1.19	3.51 ^{+0.00} _{-0.00}	53.90 ^{+1.31} _{-1.31}	0.01 ^{+0.02} _{-0.01}	0.73 ^{+1.04} _{-0.54}	0.45 ^{+0.01} _{-0.01}	0.05 ^{+0.00} _{-0.00}
HD 75898 b	1.30	421.64 ^{+7.49} _{-7.35}	74.99 ^{+3.83} _{-3.66}	0.05 ^{+0.07} _{-0.04}	9.56 ^{+2.16} _{-1.59}	3.29 ^{+0.17} _{-0.16}	1.20 ^{+0.01} _{-0.01}
HD 76700 b	1.13	3.97 ^{+0.00} _{-0.00}	27.30 ^{+1.35} _{-1.29}	0.08 ^{+0.06} _{-0.05}	2.79 ^{+1.46} _{-1.58}	0.23 ^{+0.01} _{-0.01}	0.05 ^{+0.00} _{-0.00}
HD 80606 b	0.96	111.44 ^{+0.00} _{-0.00}	560.69 ^{+187.15} _{-113.26}	0.95 ^{+0.01} _{-0.02}	12.47 ^{+2.65} _{-2.53}	3.84 ^{+0.51} _{-0.30}	0.45 ^{+0.00} _{-0.00}
HD 81040 b	0.96	1108.32 ^{+7.45} _{-4.65}	169.71 ^{+10.03} _{-10.03}	0.42 ^{+0.07} _{-0.14}	25.51 ^{+6.90} _{-5.16}	7.62 ^{+0.56} _{-0.47}	2.07 ^{+0.01} _{-0.01}
HD 82943 b	1.13	4030.24 ^{+7058.52} _{-3812.23}	88.77 ^{+347.92} _{-55.10}	0.49 ^{+0.30} _{-0.33}	34.40 ^{+8.27} _{-6.74}	5.58 ^{+28.20} _{-4.69}	5.16 ^{+4.97} _{-4.42}
HD 8574 b	1.12	227.45 ^{+0.82} _{-0.79}	64.92 ^{+4.36} _{-4.37}	0.28 ^{+0.05} _{-0.05}	9.70 ^{+2.53} _{-2.39}	2.02 ^{+0.13} _{-0.13}	0.76 ^{+0.00} _{-0.00}
HD 86081 b	1.21	2.14 ^{+0.00} _{-0.00}	207.49 ^{+0.86} _{-0.84}	0.01 ^{+0.01} _{-0.01}	0.76 ^{+0.92} _{-0.56}	1.49 ^{+0.01} _{-0.01}	0.03 ^{+0.00} _{-0.00}
HD 89307	0.99	2174.23 ^{+49.69} _{-53.83}	38.05 ^{+26.97} _{-7.05}	0.29 ^{+0.36} _{-0.20}	3.31 ^{+5.42} _{-2.68}	2.29 ^{+0.83} _{-0.41}	3.27 ^{+0.05} _{-0.05}
HD 93083 b	0.70	143.99 ^{+1.54} _{-1.47}	18.68 ^{+1.26} _{-1.20}	0.12 ^{+0.07} _{-0.06}	2.28 ^{+0.75} _{-0.54}	0.38 ^{+0.03} _{-0.03}	0.48 ^{+0.00} _{-0.00}
HR 810 b	1.11	302.94 ^{+2.27} _{-2.18}	57.17 ^{+5.60} _{-5.43}	0.12 ^{+0.10} _{-0.08}	17.92 ^{+3.70} _{-2.91}	2.00 ^{+0.20} _{-0.20}	0.91 ^{+0.00} _{-0.00}
κ CrB b	1.80	1218.78 ^{+35.22} _{-28.41}	24.06 ^{+1.39} _{-1.41}	0.12 ^{+0.08} _{-0.07}	4.73 ^{+0.86} _{-0.75}	1.86 ^{+0.11} _{-0.11}	2.72 ^{+0.05} _{-0.04}
NGC 2423 3 b	2.40	713.95 ^{+4.99} _{-5.20}	133.14 ^{+7.95} _{-7.28}	0.18 ^{+0.06} _{-0.06}	18.05 ^{+2.72} _{-2.32}	10.32 ^{+0.56} _{-0.55}	2.09 ^{+0.01} _{-0.01}
NGC 4349 127 b	3.90	676.93 ^{+4.37} _{-4.48}	189.46 ^{+10.22} _{-9.44}	0.19 ^{+0.05} _{-0.05}	14.59 ^{+3.43} _{-2.50}	19.89 ^{+1.06} _{-1.02}	2.38 ^{+0.01} _{-0.01}
τ Boo b	1.34	3.31 ^{+10.38} _{-10.09}	469.05 ^{+14.76} _{-14.87}	0.07 ^{+0.03} _{-0.04}	96.65 ^{+8.42} _{-7.39}	4.17 ^{+0.13} _{-0.13}	0.05 ^{+0.00} _{-0.00}
TrES-3 b	0.92	1.34 ^{+0.04} _{-0.04}	352.25 ^{+17.08} _{-21.08}	0.06 ^{+0.09} _{-0.04}	10.70 ^{+24.41} _{-9.07}	1.80 ^{+0.09} _{-0.12}	0.02 ^{+0.00} _{-0.00}
WASP-2 b	0.88	2.15 ^{+0.00} _{-0.00}	159.43 ^{+8.00} _{-6.39}	0.25 ^{+0.10} _{-0.11}	2.32 ^{+5.14} _{-1.87}	0.90 ^{+0.04} _{-0.03}	0.03 ^{+0.00} _{-0.00}
WASP-3 b	1.22	1.85 ^{+0.00} _{-0.00}	250.29 ^{+11.08} _{-11.16}	0.07 ^{+0.05} _{-0.04}	3.85 ^{+11.11} _{-3.21}	1.72 ^{+0.08} _{-0.08}	0.03 ^{+0.00} _{-0.00}
WASP-4 b	0.91	1.34 ^{+0.00} _{-0.00}	238.69 ^{+10.83} _{-11.83}	0.03 ^{+0.04} _{-0.02}	11.37 ^{+11.28} _{-9.52}	1.21 ^{+0.06} _{-0.06}	0.02 ^{+0.00} _{-0.00}
WASP-5 b	1.01	1.63 ^{+0.00} _{-0.00}	282.92 ^{+9.11} _{-9.08}	0.07 ^{+0.04} _{-0.03}	2.77 ^{+6.48} _{-2.23}	1.64 ^{+0.05} _{-0.05}	0.03 ^{+0.00} _{-0.00}
XO-1 b	1.03	3.94 ^{+0.00} _{-0.00}	119.02 ^{+12.73} _{-11.60}	0.10 ^{+0.11} _{-0.07}	3.09 ^{+8.03} _{-2.54}	0.94 ^{+0.10} _{-0.09}	0.05 ^{+0.00} _{-0.00}
XO-2 b	0.97	2.62 ^{+0.00} _{-0.00}	86.00 ^{+10.51} _{-10.07}	0.19 ^{+0.15} _{-0.13}	3.58 ^{+9.40} _{-2.94}	0.56 ^{+0.06} _{-0.07}	0.04 ^{+0.00} _{-0.00}
XO-3 b	1.41	3.19 ^{+0.00} _{-0.00}	1501.73 ^{+16.33} _{-16.28}	0.29 ^{+0.01} _{-0.01}	39.89 ^{+7.29} _{-6.39}	13.09 ^{+0.14} _{-0.14}	0.05 ^{+0.00} _{-0.00}
XO-4 b	1.32	4.13 ^{+0.00} _{-0.00}	200.12 ^{+92.23} _{-34.80}	0.35 ^{+0.22} _{-0.20}	4.34 ^{+13.06} _{-3.64}	1.79 ^{+0.50} _{-0.27}	0.06 ^{+0.00} _{-0.00}

Table A2. Table of the orbital parameters for a two-planet fit, both directly output from EXOFAST and thence derived. The values of the parameters T , K , e and s (generated from EXOFAST) are the medians of the parameter posterior distributions, with the associated 68.3 per cent confidence regions. The other parameters were calculated using these values and stellar masses taken from the published literature. Note that some parameters are extremely well constrained; hence, the errors on the parameter estimates are so small as to appear to be zero to the two decimal places shown in this table. A full table in machine-readable format will also be provided at <http://www.ucl.ac.uk/exoplanets/exocat>, and the reader is directed there if such data are required.

Planet	m_* (M_\odot)	T (d)	K (m s^{-1})	e	s	$m_p \sin(i)$ (M_{Jup})	a (au)
BD-17 63 b	0.74	1976.15 ^{+226.40} _{-182.28}	297.62 ^{+389.34} _{-93.47}	0.87 ^{+0.07} _{-0.07}	9.46 ^{+1.96} _{-1.47}	7.48 ^{+4.67} _{-1.37}	2.79 ^{+0.21} _{-0.17}
c		2829.26 ^{+303.62} _{-183.09}	182.40 ^{+7.20} _{-5.52}	0.84 ^{+0.01} _{-0.01}		5.68 ^{+0.17} _{-0.14}	3.54 ^{+0.25} _{-0.15}
ChaHa8 b	0.10	1184.29 ^{+619.30} _{-1170.75}	870.23 ^{+660.70} _{-855.84}	0.43 ^{+0.30} _{-0.29}	29.42 ^{+159.46} _{-27.44}	6.15 ^{+10.65} _{-6.11}	1.02 ^{+0.33} _{-0.97}
c		504.54 ^{+1270.37} _{-493.52}	965.92 ^{+484.80} _{-952.10}	0.39 ^{+0.32} _{-0.27}		6.84 ^{+9.21} _{-6.80}	0.58 ^{+0.76} _{-0.53}
ϵ Eri b	0.86	2443.29 ^{+45.64} _{-44.45}	13.91 ^{+1.51} _{-1.49}	0.05 ^{+0.07} _{-0.04}	7.72 ^{+0.84} _{-0.76}	0.83 ^{+0.09} _{-0.09}	3.37 ^{+0.04} _{-0.04}
c		541.54 ^{+2.09} _{-1.05}	14.86 ^{+8.97} _{-4.11}	0.82 ^{+0.09} _{-0.10}		0.30 ^{+0.08} _{-0.06}	1.23 ^{+0.00} _{-0.00}
ϵ Tau b	2.70	598.14 ^{+11.23} _{-10.56}	97.38 ^{+4.08} _{-3.98}	0.13 ^{+0.06} _{-0.05}	6.17 ^{+3.70} _{-5.02}	7.76 ^{+0.30} _{-0.31}	1.93 ^{+0.02} _{-0.02}
c		71.16 ^{+113.78} _{-70.55}	12.48 ^{+13.93} _{-10.77}	0.66 ^{+0.25} _{-0.37}		0.18 ^{+0.90} _{-0.14}	0.47 ^{+2.58} _{-0.45}
γ Cep b	1.59	905.43 ^{+4.49} _{-3.95}	37.97 ^{+4.81} _{-4.42}	0.04 ^{+0.07} _{-0.03}	1.51 ^{+2.27} _{-1.16}	2.46 ^{+0.31} _{-0.29}	2.14 ^{+0.01} _{-0.01}
c		3648.72 ^{+2293.49} _{-661.35}	1761.05 ^{+176.92} _{-243.13}	0.75 ^{+0.06} _{-0.09}		117.32 ^{+51.84} _{-22.41}	5.41 ^{+2.08} _{-0.68}
GJ 3021 b	0.90	133.71 ^{+0.20} _{-0.21}	166.78 ^{+3.96} _{-3.91}	0.51 ^{+0.02} _{-0.02}	15.40 ^{+2.49} _{-2.32}	3.36 ^{+0.08} _{-0.08}	0.49 ^{+0.00} _{-0.00}
c		33.23 ^{+3748.49} _{-32.08}	5.00 ^{+14.75} _{-4.26}	0.56 ^{+0.32} _{-0.38}		0.05 ^{+0.26} _{-0.05}	0.20 ^{+4.39} _{-0.17}
GJ 317 b	0.24	682.64 ^{+4.93} _{-4.68}	82.68 ^{+4.15} _{-3.87}	0.27 ^{+0.05} _{-0.06}	4.51 ^{+4.39} _{-2.90}	1.33 ^{+0.06} _{-0.06}	0.94 ^{+0.00} _{-0.00}
c		4602.69 ^{+3780.22} _{-1555.08}	33.00 ^{+22.89} _{-7.52}	0.35 ^{+0.23} _{-0.19}		0.95 ^{+0.86} _{-0.26}	3.37 ^{+1.65} _{-0.81}
GJ 674 b	0.35	4.69 ^{+0.00} _{-0.00}	8.59 ^{+0.39} _{-0.39}	0.14 ^{+0.04} _{-0.05}	1.17 ^{+0.28} _{-0.22}	0.03 ^{+0.00} _{-0.00}	0.04 ^{+0.00} _{-0.00}
c		34.82 ^{+0.07} _{-0.08}	4.84 ^{+0.40} _{-0.42}	0.23 ^{+0.08} _{-0.08}		0.04 ^{+0.00} _{-0.00}	0.15 ^{+0.00} _{-0.00}

Table A2 – *continued*

Planet	m_* (M_\odot)	T (d)	K (m s^{-1})	e	s	$m_p \sin(i)$ (M_{Jup})	a (au)
GJ 849 b	0.49	$1971.53^{+178.78}_{-104.48}$	$21.41^{+1.79}_{-1.68}$	$0.11^{+0.07}_{-0.07}$	$2.89^{+0.92}_{-0.85}$	$0.81^{+0.08}_{-0.07}$	$2.43^{+0.14}_{-0.09}$
c		$9065.80^{+3863.21}_{-3507.96}$	$65.32^{+84.73}_{-38.59}$	$0.59^{+0.31}_{-0.37}$		$3.06^{+2.95}_{-1.70}$	$6.71^{+1.79}_{-1.87}$
GJ 86 b	0.80	$15.77^{+0.00}_{-0.00}$	$378.50^{+1.12}_{-1.11}$	$0.05^{+0.00}_{-0.00}$	$0.98^{+1.25}_{-0.73}$	$4.02^{+0.01}_{-0.01}$	$0.11^{+0.00}_{-0.00}$
c		$6771.07^{+1223.25}_{-972.87}$	$1695.71^{+186.49}_{-272.94}$	$0.80^{+0.04}_{-0.03}$		$80.82^{+7.66}_{-7.35}$	$6.50^{+0.76}_{-0.64}$
HAT-P-6 b	1.29	$3.85^{+1126.88}_{-85.42}$	$107.11^{+13.37}_{-101.50}$	$0.11^{+0.60}_{-0.09}$	$48.20^{+37.86}_{-35.69}$	$1.00^{+0.14}_{-0.91}$	$0.05^{+2.26}_{-0.00}$
c		$87.07^{+4320.01}_{-0.01}$	$8.92^{+63.07}_{-7.84}$	$0.55^{+0.33}_{-0.38}$		$0.14^{+1.66}_{-0.12}$	$0.42^{+5.31}_{-0.39}$
HAT-P-8 b	1.28	$3.08^{+0.00}_{-0.01}$	$160.51^{+8.09}_{-6.77}$	$0.05^{+0.06}_{-0.03}$	$4.94^{+4.87}_{-3.95}$	$1.35^{+0.07}_{-0.06}$	$0.05^{+0.00}_{-0.00}$
c		$24.39^{+2317.94}_{-23.15}$	$11.23^{+49.73}_{-9.61}$	$0.54^{+0.33}_{-0.37}$		$0.13^{+1.02}_{-0.11}$	$0.18^{+3.57}_{-0.15}$
HAT-P-9 b	1.30	$3.92^{+0.00}_{-0.00}$	$84.33^{+11.66}_{-9.75}$	$0.17^{+0.14}_{-0.10}$	$3.28^{+8.16}_{-2.70}$	$0.77^{+0.10}_{-0.09}$	$0.05^{+0.00}_{-0.00}$
c		$138.54^{+4330.13}_{-136.58}$	$8.19^{+35.70}_{-7.18}$	$0.53^{+0.33}_{-0.37}$		$0.13^{+1.31}_{-0.12}$	$0.57^{+5.22}_{-0.54}$
HD 101930 b	0.74	$70.64^{+0.48}_{-0.44}$	$18.14^{+0.93}_{-0.92}$	$0.08^{+0.05}_{-0.05}$	$1.62^{+0.71}_{-0.64}$	$0.30^{+0.02}_{-0.02}$	$0.30^{+0.00}_{-0.00}$
c		$362.82^{+3814.57}_{-357.99}$	$2.85^{+7.90}_{-2.19}$	$0.55^{+0.33}_{-0.36}$		$0.05^{+0.26}_{-0.04}$	$0.90^{+3.69}_{-0.85}$
HD 108874 b	0.95	$395.33^{+0.97}_{-0.94}$	$37.76^{+1.14}_{-1.14}$	$0.07^{+0.03}_{-0.03}$	$0.81^{+0.91}_{-0.60}$	$1.31^{+0.04}_{-0.04}$	$1.04^{+0.00}_{-0.00}$
c		$1608.44^{+41.21}_{-33.38}$	$18.43^{+0.95}_{-0.90}$	$0.25^{+0.04}_{-0.04}$		$0.99^{+0.06}_{-0.05}$	$2.64^{+0.04}_{-0.04}$
HD 11506 b	1.19	$1337.48^{+247.15}_{-64.76}$	$65.38^{+18.91}_{-6.35}$	$0.30^{+0.16}_{-0.08}$	$4.08^{+1.63}_{-1.35}$	$3.78^{+1.15}_{-0.35}$	$2.52^{+0.30}_{-0.08}$
c		$170.36^{+1.82}_{-89.93}$	$25.54^{+6.16}_{-7.51}$	$0.36^{+0.13}_{-0.17}$		$0.73^{+0.16}_{-0.32}$	$0.64^{+0.00}_{-0.25}$
HD 118203 b	1.23	$6.13^{+0.00}_{-0.00}$	$217.20^{+4.95}_{-4.82}$	$0.31^{+0.02}_{-0.02}$	$14.34^{+3.29}_{-2.96}$	$2.14^{+0.04}_{-0.04}$	$0.07^{+0.00}_{-0.00}$
c		$6753.13^{+5243.74}_{-3967.75}$	$397.12^{+386.81}_{-294.44}$	$0.51^{+0.24}_{-0.33}$		$31.82^{+40.68}_{-24.02}$	$7.49^{+3.50}_{-3.34}$
HD 12661 b	1.14	$262.67^{+0.12}_{-0.09}$	$74.33^{+0.73}_{-0.72}$	$0.36^{+0.01}_{-0.01}$	$3.02^{+0.67}_{-0.64}$	$2.39^{+0.02}_{-0.02}$	$0.84^{+0.00}_{-0.00}$
c		$1681.47^{+29.15}_{-26.29}$	$29.16^{+0.79}_{-0.83}$	$0.02^{+0.02}_{-0.01}$		$1.86^{+0.05}_{-0.05}$	$2.89^{+0.03}_{-0.03}$
HD 128311 b	0.83	$458.32^{+3.00}_{-2.95}$	$53.71^{+5.30}_{-5.55}$	$0.33^{+0.06}_{-0.06}$	$16.15^{+1.55}_{-1.37}$	$1.69^{+0.18}_{-0.19}$	$1.09^{+0.00}_{-0.00}$
c		$924.69^{+6.43}_{-6.83}$	$75.43^{+3.28}_{-3.15}$	$0.14^{+0.10}_{-0.10}$		$3.14^{+0.12}_{-0.12}$	$1.74^{+0.01}_{-0.01}$
HD 131664 b	1.10	$1964.51^{+35.83}_{-45.84}$	$363.85^{+31.98}_{-23.96}$	$0.64^{+0.03}_{-0.03}$	$4.67^{+0.86}_{-0.74}$	$18.28^{+1.10}_{-0.86}$	$3.17^{+0.04}_{-0.05}$
c		$561.36^{+1790.01}_{-558.23}$	$5.40^{+10.32}_{-4.29}$	$0.46^{+0.39}_{-0.31}$		$0.14^{+0.53}_{-0.13}$	$1.37^{+2.20}_{-1.33}$
HD 132406 b	1.09	$1156.94^{+381.48}_{-815.09}$	$89.58^{+49.38}_{-63.98}$	$0.39^{+0.29}_{-0.25}$	$6.90^{+9.73}_{-5.67}$	$4.75^{+2.15}_{-3.97}$	$2.22^{+0.46}_{-1.23}$
c		$540.91^{+892.18}_{-326.23}$	$49.33^{+68.99}_{-27.88}$	$0.39^{+0.31}_{-0.24}$		$1.74^{+4.10}_{-1.19}$	$1.34^{+1.22}_{-0.62}$
HD 142 b	1.23	$349.57^{+3.40}_{-3.58}$	$29.66^{+4.66}_{-4.22}$	$0.20^{+0.11}_{-0.11}$	$8.74^{+1.54}_{-1.34}$	$1.15^{+0.16}_{-0.15}$	$1.04^{+0.01}_{-0.01}$
c		$9822.95^{+3348.44}_{-3208.17}$	$53.42^{+53.99}_{-17.59}$	$0.18^{+0.20}_{-0.13}$		$6.27^{+7.33}_{-2.53}$	$9.62^{+2.08}_{-2.23}$
HD 142022 b	0.90	$1877.47^{+35.74}_{-20.20}$	$127.61^{+84.60}_{-35.05}$	$0.62^{+0.11}_{-0.10}$	$2.84^{+1.76}_{-1.47}$	$5.70^{+2.52}_{-1.21}$	$2.88^{+0.04}_{-0.02}$
c		$244.04^{+6905.04}_{-242.28}$	$3.41^{+8.61}_{-2.65}$	$0.52^{+0.35}_{-0.36}$		$0.05^{+6.27}_{-0.05}$	$0.74^{+6.27}_{-0.71}$
HD 149143 b	1.20	$4.07^{+0.00}_{-0.00}$	$149.85^{+1.79}_{-1.66}$	$0.01^{+0.01}_{-0.00}$	$1.33^{+1.83}_{-1.02}$	$1.33^{+0.02}_{-0.01}$	$0.05^{+0.00}_{-0.00}$
c		$248.47^{+3300.06}_{-245.40}$	$3.73^{+13.28}_{-3.00}$	$0.55^{+0.33}_{-0.37}$		$0.07^{+0.55}_{-0.06}$	$0.82^{+4.02}_{-0.78}$
HD 154345 b	0.89	$3216.13^{+150.82}_{-3170.93}$	$13.11^{+1.63}_{-10.59}$	$0.10^{+0.47}_{-0.08}$	$2.38^{+0.47}_{-0.41}$	$0.87^{+0.12}_{-0.84}$	$4.10^{+0.13}_{-3.86}$
c		$2695.22^{+697.19}_{-2649.86}$	$6.21^{+8.30}_{-4.22}$	$0.15^{+0.55}_{-0.11}$		$0.15^{+0.83}_{-0.13}$	$3.65^{+0.60}_{-3.41}$
HD 155358 b	0.87	$193.24^{+0.98}_{-0.97}$	$32.88^{+1.51}_{-1.50}$	$0.13^{+0.04}_{-0.04}$	$5.59^{+0.72}_{-0.65}$	$0.84^{+0.04}_{-0.04}$	$0.62^{+0.00}_{-0.00}$
c		$314.82^{+4.92}_{-3.95}$	$33.63^{+41.74}_{-12.77}$	$0.83^{+0.12}_{-0.17}$		$0.57^{+0.23}_{-0.11}$	$0.86^{+0.01}_{-0.01}$
HD 162020 b	0.80	$8.43^{+0.00}_{-0.00}$	$1808.83^{+5.13}_{-5.09}$	$0.28^{+0.00}_{-0.00}$	$10.45^{+2.84}_{-2.75}$	$15.00^{+0.04}_{-0.04}$	$0.08^{+0.00}_{-0.00}$
c		$180.71^{+3271.59}_{-175.94}$	$6.73^{+21.75}_{-5.73}$	$0.57^{+0.33}_{-0.39}$		$0.10^{+0.41}_{-0.09}$	$0.58^{+3.57}_{-0.53}$
HD 168443 b	1.01	$58.11^{+0.00}_{-0.00}$	$529.04^{+82.29}_{-52.08}$	$0.57^{+0.04}_{-0.04}$	$7.80^{+3.15}_{-2.74}$	$8.37^{+0.93}_{-0.61}$	$0.29^{+0.00}_{-0.00}$
c		$1755.58^{+6.65}_{-7.25}$	$302.53^{+3.86}_{-3.76}$	$0.23^{+0.02}_{-0.02}$		$17.61^{+0.21}_{-0.21}$	$2.86^{+0.01}_{-0.01}$
HD 169830 b	1.41	$225.61^{+0.29}_{-0.31}$	$83.01^{+3.29}_{-3.25}$	$0.37^{+0.03}_{-0.03}$	$1.46^{+2.36}_{-1.13}$	$2.91^{+0.11}_{-0.11}$	$0.81^{+0.00}_{-0.00}$
c		$139.53^{+4188.83}_{-137.98}$	$3.39^{+9.48}_{-2.78}$	$0.55^{+0.33}_{-0.37}$		$0.05^{+0.43}_{-0.05}$	$0.59^{+5.24}_{-0.56}$
HD 171028 b	0.99	$267.54^{+7.12}_{-4.33}$	$47.44^{+5.43}_{-2.78}$	$0.57^{+0.13}_{-0.08}$	$1.77^{+0.66}_{-0.47}$	$1.23^{+0.18}_{-0.19}$	$0.81^{+0.01}_{-0.01}$
c		$498.38^{+29.74}_{-41.02}$	$72.24^{+80.06}_{-23.28}$	$0.61^{+0.15}_{-0.11}$		$2.27^{+1.75}_{-0.69}$	$1.23^{+0.05}_{-0.07}$
HD 183263 b	1.12	$625.64^{+0.96}_{-0.59}$	$86.82^{+1.28}_{-1.28}$	$0.38^{+0.01}_{-0.01}$	$3.54^{+0.57}_{-0.48}$	$3.65^{+0.05}_{-0.05}$	$1.49^{+0.00}_{-0.00}$
c		$4630.04^{+1456.78}_{-1072.15}$	$74.41^{+42.59}_{-19.98}$	$0.07^{+0.09}_{-0.05}$		$6.57^{+4.72}_{-2.18}$	$5.65^{+1.13}_{-0.91}$
HD 185269 b	1.30	$6.84^{+0.00}_{-0.00}$	$89.81^{+4.33}_{-4.08}$	$0.28^{+0.03}_{-0.04}$	$7.14^{+2.10}_{-2.25}$	$0.96^{+0.04}_{-0.04}$	$0.08^{+0.00}_{-0.00}$
c		$32.98^{+2639.27}_{-31.94}$	$4.17^{+9.25}_{-3.46}$	$0.55^{+0.32}_{-0.36}$		$0.05^{+0.19}_{-0.04}$	$0.22^{+3.89}_{-0.20}$
HD 187123 b	1.04	$3.10^{+0.00}_{-0.00}$	$69.57^{+0.49}_{-0.49}$	$0.01^{+0.01}_{-0.01}$	$0.49^{+0.53}_{-0.35}$	$0.51^{+0.00}_{-0.00}$	$0.04^{+0.00}_{-0.00}$
c		$5502.18^{+2809.80}_{-1322.20}$	$28.50^{+5.09}_{-3.06}$	$0.32^{+0.12}_{-0.08}$		$2.42^{+0.67}_{-0.40}$	$6.18^{+1.96}_{-1.03}$

Table A2 – continued

Planet	m_* (M_\odot)	T (d)	K (m s^{-1})	e	s	$m_p \sin(i)$ (M_{Jup})	a (au)
HD 189733 b	0.81	$2.22^{+0.00}_{-0.00}$	$198.36^{+4.00}_{-4.07}$	$0.02^{+0.02}_{-0.02}$	$11.26^{+0.98}_{-0.87}$	$1.11^{+0.02}_{-0.02}$	$0.03^{+0.00}_{-0.00}$
c		$3.92^{+0.00}_{-0.00}$	$25.58^{+4.20}_{-3.99}$	$0.96^{+0.01}_{-0.01}$		$0.05^{+0.01}_{-0.01}$	$0.05^{+0.00}_{-0.00}$
HD 190228 b	1.82	$1140.04^{+16.47}_{-15.62}$	$92.35^{+5.40}_{-3.72}$	$0.52^{+0.05}_{-0.04}$	$1.20^{+1.82}_{-0.91}$	$6.07^{+4.92}_{-6.64}$	$2.61^{+0.03}_{-0.02}$
c		$149.82^{+3434.51}_{-147.94}$	$2.23^{+5.57}_{-1.79}$	$0.55^{+0.33}_{-0.37}$		$0.05^{+0.30}_{-0.04}$	$0.67^{+4.92}_{-0.64}$
HD 190360 b	0.98	$2925.88^{+36.43}_{-41.38}$	$18.84^{+2.95}_{-3.16}$	$0.34^{+0.13}_{-0.12}$	$5.31^{+1.64}_{-2.14}$	$1.23^{+0.16}_{-0.20}$	$3.98^{+0.03}_{-0.04}$
c		$184.70^{+3848.83}_{-181.67}$	$5.10^{+10.61}_{-4.14}$	$0.55^{+0.32}_{-0.35}$		$0.09^{+0.49}_{-0.08}$	$0.63^{+4.30}_{-0.59}$
HD 190647 b	1.10	$1036.28^{+8.27}_{-14.95}$	$36.17^{+1.89}_{-1.75}$	$0.17^{+0.03}_{-0.04}$	$0.59^{+0.68}_{-0.43}$	$1.89^{+0.10}_{-0.10}$	$2.07^{+0.01}_{-0.02}$
c		$422.54^{+3134.94}_{-411.93}$	$2.88^{+13.42}_{-2.23}$	$0.55^{+0.30}_{-0.36}$		$0.08^{+0.77}_{-0.07}$	$1.14^{+3.57}_{-1.04}$
HD 195019 b	1.02	$45.87^{+11.86}_{-0.76}$	$176.87^{+70.77}_{-43.26}$	$0.79^{+0.11}_{-0.17}$	$149.90^{+9.71}_{-8.67}$	$1.96^{+0.64}_{-0.42}$	$0.25^{+0.04}_{-0.00}$
c		$51.93^{+0.08}_{-0.10}$	$159.88^{+33.08}_{-34.85}$	$0.74^{+0.07}_{-0.08}$		$1.95^{+0.52}_{-0.45}$	$0.27^{+0.00}_{-0.00}$
HD 202206 b	1.07	$255.87^{+0.08}_{-0.08}$	$706.93^{+119.27}_{-65.64}$	$0.37^{+0.02}_{-0.02}$	$23.09^{+2.18}_{-1.91}$	$21.46^{+3.75}_{-2.07}$	$0.81^{+0.00}_{-0.00}$
c		$258.20^{+1.14}_{-1.02}$	$291.17^{+164.81}_{-66.27}$	$0.26^{+0.06}_{-0.06}$		$9.23^{+4.97}_{-2.03}$	$0.81^{+0.00}_{-0.00}$
HD 20868 b	0.78	$380.86^{+0.08}_{-0.08}$	$100.22^{+0.45}_{-0.46}$	$0.76^{+0.00}_{-0.00}$	$0.74^{+0.43}_{-0.48}$	$1.99^{+0.01}_{-0.01}$	$0.95^{+0.00}_{-0.00}$
c		$111.05^{+1942.40}_{-110.42}$	$1.59^{+2.19}_{-1.07}$	$0.60^{+0.30}_{-0.38}$		$0.02^{+0.07}_{-0.01}$	$0.42^{+2.49}_{-0.40}$
HD 209458 b	1.13	$3.52^{+0.00}_{-0.00}$	$84.27^{+0.85}_{-0.84}$	$0.01^{+0.01}_{-0.01}$	$2.91^{+0.79}_{-0.88}$	$0.68^{+0.01}_{-0.01}$	$0.05^{+0.00}_{-0.00}$
c		$264.63^{+1050.15}_{-259.91}$	$2.85^{+3.04}_{-2.03}$	$0.54^{+0.34}_{-0.37}$		$0.07^{+0.11}_{-0.06}$	$0.84^{+1.61}_{-0.78}$
HD 212301 b	1.05	$2.25^{+0.00}_{-0.00}$	$56.85^{+1.92}_{-1.93}$	$0.09^{+0.04}_{-0.04}$	$5.42^{+1.58}_{-1.26}$	$0.38^{+0.01}_{-0.01}$	$0.03^{+0.00}_{-0.00}$
c		$2356.61^{+6274.66}_{-2104.70}$	$26.81^{+76.37}_{-17.28}$	$0.52^{+0.35}_{-0.36}$		$1.22^{+4.93}_{-0.89}$	$3.52^{+4.85}_{-2.73}$
HD 217107 b	1.11	$7.13^{+0.00}_{-0.00}$	$138.34^{+1.11}_{-1.11}$	$0.13^{+0.01}_{-0.01}$	$10.19^{+0.64}_{-0.59}$	$1.39^{+0.01}_{-0.01}$	$0.08^{+0.00}_{-0.00}$
c		$4106.23^{+248.05}_{-113.88}$	$39.48^{+13.92}_{-3.49}$	$0.56^{+0.04}_{-0.03}$		$2.77^{+1.00}_{-0.27}$	$5.20^{+0.21}_{-0.10}$
HD 219828 b	1.24	$3.83^{+0.00}_{-0.00}$	$7.22^{+0.59}_{-0.62}$	$0.09^{+0.09}_{-0.06}$	$1.78^{+0.55}_{-0.39}$	$0.06^{+0.01}_{-0.01}$	$0.05^{+0.00}_{-0.00}$
c		$956.74^{+1301.58}_{-366.96}$	$78.15^{+131.30}_{-45.22}$	$0.53^{+0.17}_{-0.20}$		$3.55^{+10.00}_{-2.29}$	$2.04^{+1.58}_{-0.56}$
HD 221287 b	1.30	$455.12^{+6.45}_{-4.50}$	$71.62^{+16.56}_{-7.17}$	$0.13^{+0.12}_{-0.08}$	$8.00^{+2.88}_{-2.87}$	$3.21^{+0.65}_{-0.32}$	$1.26^{+0.01}_{-0.01}$
c		$0.60^{+525.67}_{-0.00}$	$9.66^{+6.31}_{-8.16}$	$0.41^{+0.39}_{-0.28}$		$0.05^{+0.11}_{-0.04}$	$0.02^{+1.38}_{-0.00}$
HD 224693 b	1.30	$26.75^{+0.03}_{-0.03}$	$38.90^{+1.77}_{-1.72}$	$0.04^{+0.04}_{-0.03}$	$1.81^{+1.07}_{-1.11}$	$0.68^{+0.03}_{-0.03}$	$0.19^{+0.00}_{-0.00}$
c		$7882.48^{+4298.55}_{-3049.23}$	$1849.37^{+94.81}_{-501.16}$	$0.68^{+0.12}_{-0.11}$		$153.48^{+42.92}_{-58.00}$	$8.46^{+2.85}_{-2.35}$
HD 23127 b	1.13	$1219.14^{+36.54}_{-1099.09}$	$25.84^{+4.15}_{-22.15}$	$0.44^{+0.17}_{-0.14}$	$10.66^{+2.22}_{-1.81}$	$1.31^{+0.23}_{-1.26}$	$2.33^{+0.05}_{-1.83}$
c		$1124.66^{+841.10}_{-1121.44}$	$8.38^{+19.55}_{-7.33}$	$0.46^{+0.34}_{-0.24}$		$0.15^{+1.28}_{-0.14}$	$2.20^{+0.99}_{-2.16}$
HD 2638 b	0.93	$3.44^{+0.00}_{-0.00}$	$67.54^{+0.87}_{-0.84}$	$0.01^{+0.01}_{-0.01}$	$2.52^{+0.98}_{-0.94}$	$0.48^{+0.01}_{-0.01}$	$0.04^{+0.00}_{-0.00}$
c		$26.04^{+1978.12}_{-25.00}$	$4.45^{+9.20}_{-2.36}$	$0.41^{+0.39}_{-0.27}$		$0.05^{+0.34}_{-0.03}$	$0.17^{+2.87}_{-0.15}$
HD 27442 b	1.20	$417.14^{+945.92}_{-9.45}$	$31.26^{+2.69}_{-28.78}$	$0.10^{+0.49}_{-0.06}$	$2.57^{+1.53}_{-1.47}$	$1.29^{+0.11}_{-1.22}$	$1.16^{+1.40}_{-0.02}$
c		$410.58^{+866.97}_{-407.42}$	$6.33^{+26.90}_{-5.26}$	$0.30^{+0.52}_{-0.26}$		$0.14^{+1.23}_{-0.12}$	$1.15^{+1.30}_{-1.10}$
HD 27894 b	0.75	$17.98^{+0.03}_{-0.01}$	$56.34^{+2.56}_{-1.29}$	$0.05^{+0.03}_{-0.03}$	$0.71^{+1.02}_{-0.53}$	$0.60^{+0.03}_{-0.01}$	$0.12^{+0.00}_{-0.00}$
c		$23.31^{+9.65}_{-0.23}$	$10.37^{+2.85}_{-1.71}$	$0.28^{+0.44}_{-0.22}$		$0.12^{+0.02}_{-0.02}$	$0.15^{+0.04}_{-0.00}$
HD 28185 b	0.99	$381.37^{+1.22}_{-2.47}$	$161.15^{+8.85}_{-10.53}$	$0.04^{+0.03}_{-0.03}$	$4.00^{+2.23}_{-2.54}$	$5.71^{+0.32}_{-0.38}$	$1.03^{+0.00}_{-0.00}$
c		$508.09^{+1160.41}_{-198.94}$	$22.84^{+26.97}_{-11.21}$	$0.37^{+0.36}_{-0.21}$		$0.86^{+1.01}_{-0.49}$	$1.24^{+1.50}_{-0.35}$
HD 285968 b	0.49	$10.23^{+0.01}_{-1.62}$	$10.62^{+2.09}_{-5.78}$	$0.19^{+0.29}_{-0.14}$	$1.91^{+1.90}_{-1.45}$	$0.07^{+0.01}_{-0.04}$	$0.07^{+0.00}_{-0.01}$
c		$10.25^{+1457.68}_{-8.28}$	$3.84^{+7.55}_{-3.20}$	$0.43^{+0.41}_{-0.32}$		$0.03^{+0.05}_{-0.02}$	$0.07^{+1.92}_{-0.05}$
HD 330075 b	0.70	$3.39^{+0.00}_{-0.00}$	$106.96^{+1.00}_{-1.00}$	$0.01^{+0.01}_{-0.00}$	$1.65^{+0.78}_{-0.68}$	$0.62^{+0.01}_{-0.01}$	$0.04^{+0.00}_{-0.00}$
c		$655.66^{+6970.11}_{-589.87}$	$10.51^{+98.32}_{-8.53}$	$0.42^{+0.40}_{-0.33}$		$0.27^{+5.27}_{-0.24}$	$1.31^{+5.42}_{-1.03}$
HD 33636 b	1.02	$2127.71^{+11.43}_{-10.98}$	$773.84^{+123.39}_{-762.03}$	$0.89^{+0.03}_{-0.22}$	$0.64^{+0.71}_{-0.47}$	$22.86^{+3.33}_{-22.32}$	$3.26^{+0.01}_{-0.01}$
c		$7841.32^{+3695.12}_{-3270.74}$	$1923.73^{+58.92}_{-174.57}$	$0.61^{+0.03}_{-0.01}$		$149.03^{+17.99}_{-27.71}$	$7.78^{+2.28}_{-2.35}$
HD 3651 b	0.88	$62.25^{+0.03}_{-0.03}$	$16.14^{+1.46}_{-1.48}$	$0.60^{+0.05}_{-0.06}$	$4.41^{+0.52}_{-0.46}$	$0.23^{+0.01}_{-0.02}$	$0.29^{+0.00}_{-0.00}$
c		$294.67^{+61.62}_{-195.79}$	$3.49^{+1.88}_{-1.60}$	$0.32^{+0.41}_{-0.23}$		$0.10^{+0.05}_{-0.06}$	$0.83^{+0.11}_{-0.43}$
HD 38529 b	1.48	$14.31^{+0.00}_{-0.00}$	$54.97^{+1.70}_{-1.66}$	$0.17^{+0.03}_{-0.03}$	$13.11^{+0.93}_{-0.86}$	$0.84^{+0.03}_{-0.02}$	$0.13^{+0.00}_{-0.00}$
c		$2148.41^{+5.95}_{-7.87}$	$170.83^{+1.92}_{-1.86}$	$0.34^{+0.01}_{-0.01}$		$13.23^{+0.14}_{-0.14}$	$3.71^{+0.01}_{-0.01}$
HD 4203 b	1.13	$438.04^{+7.20}_{-4.69}$	$56.24^{+29.91}_{-9.58}$	$0.69^{+0.13}_{-0.08}$	$1.39^{+1.65}_{-1.04}$	$1.72^{+0.53}_{-0.38}$	$1.18^{+0.01}_{-0.01}$
c		$391.98^{+202.96}_{-200.21}$	$10.70^{+11.08}_{-3.76}$	$0.32^{+0.33}_{-0.23}$		$0.37^{+0.42}_{-0.20}$	$1.09^{+0.35}_{-0.41}$
HD 4208 b	0.88	$829.27^{+9.57}_{-9.59}$	$18.79^{+0.82}_{-0.95}$	$0.06^{+0.05}_{-0.04}$	$0.80^{+0.92}_{-0.58}$	$0.80^{+0.04}_{-0.04}$	$1.66^{+0.01}_{-0.01}$
c		$129.04^{+708.37}_{-121.12}$	$2.62^{+3.03}_{-1.80}$	$0.53^{+0.34}_{-0.39}$		$0.05^{+0.07}_{-0.04}$	$0.48^{+1.19}_{-0.40}$

Downloaded from <http://mnras.oxfordjournals.org/> at University College London on April 22, 2013

Table A2 – continued

Planet	m_* (M_\odot)	T (d)	K (m s^{-1})	e	s	$m_p \sin(i)$ (M_{Jup})	a (au)
HD 43691 b	1.38	$36.96^{+0.04}_{-0.05}$	$125.00^{+4.49}_{-4.59}$	$0.12^{+0.04}_{-0.04}$	$11.09^{+3.45}_{-2.98}$	$2.52^{+0.10}_{-0.10}$	$0.24^{+0.00}_{-0.00}$
c		$516.72^{+5417.79}_{-471.62}$	$19.26^{+70.84}_{-11.52}$	$0.45^{+0.36}_{-0.31}$		$0.57^{+5.63}_{-0.39}$	$1.40^{+5.74}_{-1.13}$
HD 43848 b	0.93	$2305.82^{+158.97}_{-2261.64}$	$476.62^{+555.76}_{-471.61}$	$0.73^{+0.12}_{-0.35}$	$4.49^{+4.32}_{-3.44}$	$22.16^{+14.50}_{-22.11}$	$3.33^{+0.15}_{-3.10}$
c		$2261.82^{+513.95}_{-2252.19}$	$44.31^{+612.06}_{-41.42}$	$0.67^{+0.16}_{-0.37}$		$1.41^{+26.25}_{-1.38}$	$3.29^{+0.48}_{-3.21}$
HD 46375 b	0.93	$3.02^{+0.00}_{-0.00}$	$33.79^{+0.79}_{-0.81}$	$0.05^{+0.03}_{-0.03}$	$2.92^{+0.70}_{-0.71}$	$0.23^{+0.01}_{-0.01}$	$0.04^{+0.00}_{-0.00}$
c		$30.41^{+931.01}_{-29.23}$	$2.93^{+4.85}_{-2.28}$	$0.65^{+0.25}_{-0.42}$		$0.03^{+0.08}_{-0.02}$	$0.19^{+1.67}_{-0.16}$
HD 47536 b	0.94	$713.17^{+4061.39}_{-662.35}$	$87.58^{+41.26}_{-81.24}$	$0.30^{+0.47}_{-0.14}$	$4.58^{+9.07}_{-3.83}$	$3.76^{+1.67}_{-3.69}$	$1.53^{+3.91}_{-1.27}$
c		$698.09^{+24.29}_{-686.96}$	$105.12^{+22.10}_{-98.90}$	$0.18^{+0.40}_{-0.14}$		$4.33^{+0.95}_{-4.28}$	$1.51^{+0.03}_{-1.41}$
HD 49674 b	1.01	$4.95^{+0.00}_{-0.00}$	$11.88^{+1.04}_{-1.00}$	$0.09^{+0.08}_{-0.06}$	$2.56^{+1.04}_{-1.11}$	$0.10^{+0.01}_{-0.01}$	$0.06^{+0.00}_{-0.00}$
c		$275.09^{+2808.10}_{-248.28}$	$4.23^{+4.57}_{-2.19}$	$0.64^{+0.30}_{-0.42}$		$0.10^{+0.10}_{-0.07}$	$0.83^{+3.33}_{-0.65}$
HD 50499 b	1.28	$2450.67^{+54.08}_{-45.96}$	$21.82^{+1.91}_{-1.90}$	$0.25^{+0.07}_{-0.07}$	$2.95^{+1.02}_{-1.08}$	$1.65^{+0.15}_{-0.15}$	$3.86^{+0.06}_{-0.05}$
c		$9734.20^{+3449.64}_{-3466.65}$	$57.00^{+109.73}_{-32.17}$	$0.54^{+0.30}_{-0.34}$		$5.58^{+6.81}_{-2.96}$	$9.69^{+2.17}_{-2.46}$
HD 5319 b	1.60	$684.85^{+13.55}_{-20.85}$	$37.35^{+9.92}_{-4.77}$	$0.11^{+0.12}_{-0.07}$	$5.43^{+1.03}_{-0.82}$	$2.20^{+0.54}_{-0.28}$	$1.78^{+0.02}_{-0.04}$
c		$1541.64^{+1673.87}_{-412.57}$	$15.24^{+7.88}_{-3.29}$	$0.41^{+0.29}_{-0.28}$		$1.08^{+0.56}_{-0.23}$	$3.05^{+1.93}_{-0.57}$
HD 63454 b	0.80	$2.82^{+0.00}_{-0.00}$	$63.26^{+1.78}_{-1.79}$	$0.02^{+0.03}_{-0.02}$	$5.54^{+1.33}_{-1.18}$	$0.38^{+0.01}_{-0.01}$	$0.04^{+0.00}_{-0.00}$
c		$33.11^{+3825.01}_{-32.19}$	$3.07^{+7.74}_{-2.50}$	$0.57^{+0.33}_{-0.39}$		$0.03^{+0.15}_{-0.02}$	$0.19^{+4.28}_{-0.17}$
HD 68988 b	1.12	$6.28^{+0.00}_{-0.00}$	$189.57^{+1.57}_{-1.59}$	$0.16^{+0.01}_{-0.01}$	$3.36^{+1.19}_{-1.11}$	$1.83^{+0.01}_{-0.01}$	$0.07^{+0.00}_{-0.00}$
c		$4053.58^{+1411.19}_{-588.92}$	$68.13^{+13.26}_{-6.01}$	$0.16^{+0.09}_{-0.07}$		$5.69^{+1.68}_{-0.72}$	$5.17^{+1.14}_{-0.51}$
HD 73267 b	0.89	$1278.04^{+16.28}_{-23.90}$	$73.21^{+19.97}_{-5.33}$	$0.28^{+0.03}_{-0.09}$	$0.99^{+0.49}_{-0.55}$	$3.46^{+1.00}_{-0.23}$	$2.22^{+0.02}_{-0.03}$
c		$1265.34^{+6.17}_{-9.40}$	$18.63^{+19.36}_{-8.74}$	$0.56^{+0.23}_{-0.53}$		$0.72^{+1.12}_{-0.43}$	$2.20^{+0.01}_{-0.01}$
HD 73526 b	1.01	$193.42^{+0.47}_{-0.56}$	$114.30^{+8.71}_{-8.35}$	$0.52^{+0.04}_{-0.05}$	$9.51^{+2.84}_{-2.23}$	$2.79^{+0.16}_{-0.16}$	$0.66^{+0.00}_{-0.00}$
c		$178.17^{+0.53}_{-1.50}$	$45.30^{+9.09}_{-6.41}$	$0.70^{+0.08}_{-0.09}$		$0.91^{+0.13}_{-0.11}$	$0.62^{+0.00}_{-0.00}$
HD 74156 b	1.24	$51.65^{+0.01}_{-0.01}$	$116.23^{+3.58}_{-3.46}$	$0.65^{+0.01}_{-0.01}$	$8.55^{+0.85}_{-0.74}$	$1.88^{+0.04}_{-0.04}$	$0.29^{+0.00}_{-0.00}$
c		$2519.02^{+20.76}_{-20.00}$	$109.95^{+13.05}_{-9.31}$	$0.41^{+0.05}_{-0.05}$		$7.74^{+1.12}_{-0.85}$	$3.89^{+0.02}_{-0.02}$
HD 75289 b	1.19	$3.51^{+0.00}_{-0.00}$	$53.94^{+1.37}_{-1.33}$	$0.02^{+0.02}_{-0.01}$	$0.73^{+1.03}_{-0.55}$	$0.45^{+0.01}_{-0.01}$	$0.05^{+0.00}_{-0.00}$
c		$112.58^{+5535.62}_{-110.72}$	$2.51^{+10.42}_{-2.03}$	$0.54^{+0.34}_{-0.37}$		$0.04^{+0.61}_{-0.04}$	$0.48^{+6.09}_{-0.45}$
HD 75898 b	1.30	$419.62^{+10.28}_{-8.47}$	$69.03^{+9.74}_{-8.65}$	$0.10^{+0.08}_{-0.07}$	$5.21^{+1.49}_{-1.04}$	$3.00^{+0.43}_{-0.34}$	$1.20^{+0.02}_{-0.02}$
c		$368.23^{+6020.36}_{-59.73}$	$32.11^{+86.77}_{-16.18}$	$0.51^{+0.32}_{-0.34}$		$1.34^{+5.41}_{-0.77}$	$1.10^{+6.26}_{-0.12}$
HD 76700 b	1.13	$4.44^{+368.07}_{-0.47}$	$12.00^{+15.68}_{-10.10}$	$0.23^{+0.54}_{-0.19}$	$7.53^{+3.00}_{-4.24}$	$0.22^{+0.07}_{-0.19}$	$0.06^{+1.00}_{-0.00}$
c		$1.33^{+0.00}_{-0.01}$	$21.74^{+4.32}_{-18.71}$	$0.14^{+0.43}_{-0.10}$		$0.13^{+0.03}_{-0.11}$	$0.02^{+0.00}_{-0.00}$
HD 80606 b	0.96	$111.44^{+0.00}_{-0.00}$	$857.60^{+577.43}_{-312.73}$	$0.97^{+0.02}_{-0.02}$	$2.84^{+4.11}_{-2.27}$	$4.30^{+0.86}_{-0.56}$	$0.45^{+0.00}_{-0.00}$
c		$0.49^{+0.00}_{-0.00}$	$18.43^{+4.06}_{-3.77}$	$0.41^{+0.20}_{-0.29}$		$0.06^{+0.01}_{-0.01}$	$0.01^{+0.00}_{-0.00}$
HD 81040 b	0.96	$1100.01^{+7.14}_{-8.03}$	$178.02^{+11.03}_{-12.52}$	$0.48^{+0.08}_{-0.06}$	$8.20^{+12.12}_{-6.91}$	$7.75^{+0.37}_{-0.58}$	$2.06^{+0.01}_{-0.01}$
c		$207.07^{+265.81}_{-177.01}$	$33.11^{+13.01}_{-12.96}$	$0.25^{+0.44}_{-0.18}$		$0.87^{+0.34}_{-0.63}$	$0.68^{+0.50}_{-0.49}$
HD 82943 b	1.13	$221.52^{+1.41}_{-1.59}$	$62.85^{+9.22}_{-7.35}$	$0.35^{+0.05}_{-0.05}$	$3.83^{+1.51}_{-1.36}$	$1.90^{+0.32}_{-0.25}$	$0.75^{+0.00}_{-0.00}$
c		$445.99^{+4.32}_{-4.96}$	$37.75^{+3.92}_{-3.57}$	$0.25^{+0.14}_{-0.17}$		$1.48^{+0.10}_{-0.11}$	$1.19^{+0.01}_{-0.01}$
HD 8574 b	1.12	$227.24^{+0.70}_{-0.69}$	$65.11^{+3.48}_{-3.41}$	$0.30^{+0.04}_{-0.04}$	$3.86^{+3.52}_{-3.02}$	$2.01^{+0.10}_{-0.10}$	$0.76^{+0.00}_{-0.00}$
c		$7931.89^{+4630.89}_{-4011.00}$	$59.61^{+114.58}_{-41.65}$	$0.52^{+0.30}_{-0.33}$		$4.80^{+8.94}_{-3.38}$	$8.08^{+2.90}_{-3.03}$
HD 86081 b	1.21	$2.14^{+0.00}_{-0.00}$	$207.62^{+0.84}_{-0.84}$	$0.01^{+0.01}_{-0.00}$	$0.68^{+0.84}_{-0.50}$	$1.49^{+0.01}_{-0.01}$	$0.03^{+0.00}_{-0.00}$
c		$554.39^{+5438.59}_{-549.17}$	$2.78^{+22.00}_{-2.24}$	$0.54^{+0.34}_{-0.37}$		$0.06^{+1.29}_{-0.06}$	$1.41^{+5.47}_{-1.34}$
HD 89307 b	0.99	$2173.22^{+50.00}_{-53.20}$	$34.25^{+21.54}_{-26.30}$	$0.33^{+0.41}_{-0.23}$	$3.05^{+5.33}_{-2.46}$	$2.05^{+0.83}_{-1.67}$	$3.27^{+0.05}_{-0.05}$
c		$318.89^{+7411.65}_{-317.49}$	$9.35^{+85.84}_{-8.33}$	$0.49^{+0.36}_{-0.33}$		$0.12^{+6.16}_{-0.11}$	$0.91^{+6.72}_{-0.89}$
HD 93083 b	0.70	$143.85^{+1.68}_{-1.71}$	$17.87^{+1.46}_{-1.47}$	$0.12^{+0.07}_{-0.07}$	$1.56^{+0.89}_{-0.83}$	$0.36^{+0.03}_{-0.03}$	$0.48^{+0.00}_{-0.00}$
c		$241.82^{+3377.10}_{-225.82}$	$4.26^{+9.78}_{-2.78}$	$0.51^{+0.34}_{-0.34}$		$0.09^{+0.40}_{-0.08}$	$0.67^{+3.42}_{-0.56}$
HR 810 b	1.11	$302.15^{+2.91}_{-281.12}$	$55.27^{+7.21}_{-47.77}$	$0.14^{+0.34}_{-0.10}$	$16.54^{+4.06}_{-3.58}$	$1.93^{+0.26}_{-1.86}$	$0.91^{+0.01}_{-0.76}$
c		$302.20^{+1507.67}_{-299.19}$	$18.06^{+39.95}_{-16.02}$	$0.38^{+0.46}_{-0.29}$		$0.48^{+1.53}_{-0.45}$	$0.91^{+2.10}_{-0.87}$
κ CrB b	1.80	$1185.12^{+158.87}_{-1182.23}$	$15.83^{+9.29}_{-14.64}$	$0.18^{+0.60}_{-0.15}$	$4.75^{+0.90}_{-0.84}$	$0.95^{+1.00}_{-0.93}$	$2.67^{+0.23}_{-2.62}$
c		$1198.66^{+128.88}_{-1181.54}$	$22.27^{+2.93}_{-21.04}$	$0.14^{+0.57}_{-0.11}$		$1.71^{+0.25}_{-1.69}$	$2.69^{+0.19}_{-2.53}$
NGC 2423 3 b	2.40	$357.54^{+4.89}_{-5.52}$	$107.45^{+56.48}_{-16.98}$	$0.47^{+0.13}_{-0.13}$	$31.88^{+4.64}_{-3.78}$	$5.91^{+3.13}_{-0.96}$	$1.32^{+0.01}_{-0.01}$
c		$234.98^{+3.77}_{-3.09}$	$55.95^{+7.84}_{-7.69}$	$0.09^{+0.11}_{-0.06}$		$3.02^{+0.42}_{-0.42}$	$1.00^{+0.01}_{-0.01}$

Table A2 – continued

Planet	m_* (M_\odot)	T (d)	K (m s^{-1})	e	s	$m_p \sin(i)$ (M_{Jup})	a (au)
NGC 4349 127 b	3.90	$678.17^{+7.23}_{-7.34}$	$188.39^{+15.39}_{-12.26}$	$0.14^{+0.05}_{-0.05}$	$14.64^{+4.03}_{-3.24}$	$19.97^{+1.73}_{-1.42}$	$2.38^{+0.02}_{-0.02}$
c		$213.34^{+3126.70}_{-210.52}$	$10.89^{+49.78}_{-9.60}$	$0.53^{+0.32}_{-0.34}$		$0.37^{+6.09}_{-0.34}$	$1.10^{+5.78}_{-1.04}$
τ Boo b	1.34	$3.31^{+0.00}_{-0.00}$	$467.29^{+5.35}_{-5.14}$	$0.02^{+0.01}_{-0.01}$	$27.05^{+5.09}_{-4.69}$	$4.17^{+0.05}_{-0.05}$	$0.05^{+0.00}_{-0.00}$
c		$11695.40^{+2209.03}_{-2416.40}$	$142.19^{+14.64}_{-13.08}$	$0.39^{+0.08}_{-0.11}$		$17.61^{+1.84}_{-1.68}$	$11.12^{+1.36}_{-1.59}$
TrES-3 b	0.92	$1.34^{+0.09}_{-0.05}$	$348.72^{+23.24}_{-75.06}$	$0.08^{+0.20}_{-0.06}$	$9.34^{+23.48}_{-8.01}$	$1.77^{+0.12}_{-0.45}$	$0.02^{+0.00}_{-0.00}$
c		$50.79^{+2572.53}_{-49.53}$	$87.40^{+641.98}_{-83.99}$	$0.46^{+0.35}_{-0.34}$		$1.26^{+11.74}_{-1.22}$	$0.26^{+3.35}_{-0.24}$
WASP-2 b	0.88	$2.15^{+0.00}_{-0.00}$	$159.33^{+8.66}_{-7.16}$	$0.22^{+0.12}_{-0.13}$	$2.63^{+6.47}_{-2.12}$	$0.90^{+0.04}_{-0.04}$	$0.03^{+0.00}_{-0.00}$
c		$265.16^{+4448.69}_{-263.05}$	$10.34^{+144.62}_{-9.20}$	$0.53^{+0.34}_{-0.35}$		$0.15^{+5.19}_{-0.14}$	$0.77^{+4.50}_{-0.74}$
WASP-3 b	1.22	$1.85^{+0.00}_{-0.00}$	$249.49^{+11.68}_{-11.41}$	$0.08^{+0.05}_{-0.05}$	$3.58^{+10.43}_{-2.98}$	$1.71^{+0.08}_{-0.08}$	$0.03^{+0.00}_{-0.00}$
c		$230.52^{+4532.94}_{-227.26}$	$10.38^{+92.25}_{-9.17}$	$0.54^{+0.33}_{-0.36}$		$0.19^{+3.82}_{-0.18}$	$0.79^{+5.13}_{-0.74}$
WASP-4 b	0.91	$1.34^{+0.00}_{-0.00}$	$243.72^{+9.77}_{-10.23}$	$0.04^{+0.03}_{-0.03}$	$3.70^{+8.51}_{-3.07}$	$1.24^{+0.05}_{-0.05}$	$0.02^{+0.00}_{-0.00}$
c		$710.90^{+2716.90}_{-618.39}$	$174.77^{+810.23}_{-163.03}$	$0.54^{+0.28}_{-0.36}$		$5.46^{+36.64}_{-5.32}$	$1.51^{+2.80}_{-1.12}$
WASP-5 b	1.01	$1.63^{+0.00}_{-0.00}$	$282.05^{+9.68}_{-9.28}$	$0.07^{+0.04}_{-0.04}$	$2.89^{+7.23}_{-2.35}$	$1.64^{+0.05}_{-0.05}$	$0.03^{+0.00}_{-0.00}$
c		$338.95^{+4577.80}_{-334.17}$	$10.05^{+107.77}_{-8.95}$	$0.53^{+0.33}_{-0.37}$		$0.17^{+4.54}_{-0.16}$	$0.95^{+4.72}_{-0.90}$
XO-1 b	1.03	$3.94^{+0.00}_{-0.00}$	$118.26^{+13.15}_{-11.76}$	$0.10^{+0.12}_{-0.07}$	$3.26^{+8.68}_{-2.68}$	$0.93^{+0.10}_{-0.09}$	$0.05^{+0.00}_{-0.00}$
c		$253.27^{+4507.93}_{-251.52}$	$9.60^{+86.27}_{-8.36}$	$0.56^{+0.32}_{-0.37}$		$0.14^{+3.70}_{-0.13}$	$0.79^{+4.80}_{-0.76}$
XO-2 b	0.97	$2.62^{+0.00}_{-0.00}$	$85.83^{+10.59}_{-10.43}$	$0.18^{+0.15}_{-0.13}$	$3.52^{+9.57}_{-2.90}$	$0.55^{+0.07}_{-0.07}$	$0.04^{+0.00}_{-0.00}$
c		$188.83^{+4420.68}_{-186.36}$	$9.80^{+95.98}_{-8.70}$	$0.52^{+0.35}_{-0.35}$		$0.14^{+3.68}_{-0.13}$	$0.64^{+4.73}_{-0.60}$
XO-3 b	1.41	$3.60^{+0.01}_{-0.01}$	$1133.58^{+83.26}_{-78.68}$	$0.20^{+0.06}_{-0.06}$	$203.12^{+24.39}_{-20.46}$	$10.51^{+0.82}_{-0.77}$	$0.05^{+0.00}_{-0.00}$
c		$3.38^{+0.01}_{-0.01}$	$2272.98^{+98.15}_{-97.47}$	$0.03^{+0.03}_{-0.02}$		$21.09^{+0.91}_{-0.91}$	$0.05^{+0.00}_{-0.00}$
XO-4 b	1.32	$4.13^{+0.00}_{-0.00}$	$191.72^{+89.73}_{-32.29}$	$0.32^{+0.25}_{-0.21}$	$4.44^{+13.91}_{-3.74}$	$1.74^{+0.51}_{-0.26}$	$0.06^{+0.00}_{-0.00}$
c		$235.76^{+4051.59}_{-233.46}$	$14.93^{+186.58}_{-13.58}$	$0.52^{+0.34}_{-0.35}$		$0.27^{+8.47}_{-0.25}$	$0.82^{+4.85}_{-0.78}$

Table A3. The number of published planets compared with the best-fitting model from this analysis (i.e. the flags and reduced chi-square ratios for the results for each system). The ‘candidates’ column shows the current number of confirmed planets (from <http://www.exoplanet.eu> and <http://exoplanets.org>, as of 2011 August 01), and the ‘visual quality flag’ (assigned by eye) is the best EXOFAST model, where ‘1’ signifies that the one-planet fit is best and ‘2’ means that the two-planet fit is best. ‘3’ means that both one- and two-planet solutions provide equally good or bad fits, and this class is again subdivided into ‘3a’ and ‘3b’, as explained in Section 6.2. Also shown is the log likelihood ratio of the chi-square values, R , as defined in Section 5. The visual flag assignments are validated somewhat by noting that in 99 per cent of systems the visual flag and chi-square results agree (or at least are not contradictory, for the class 3 cases). Those systems denoted by ‘–’ are those where there were not sufficient degrees of freedom to calculate a value for the log likelihood ratio. The prior flag is also shown, where flag N indicates that the analysis was performed using the normal priors shown in Tables 1 and 2, and flag D indicates an analysis with different priors as shown in Table 3.

System	Number of candidates from literature	Visual quality flag	R	Flag for period prior used
BD-17 63	1	1	25.06	N
ChaHa8	1	1	–	N
ϵ Eri	1	3a	–0.84	D
ϵ Tau	1	1	4.62	N
γ Cep	1	2	–398.83	D
GJ 3021	1	3b	0.28	N
GJ 317	1	1	47.75	N
GJ 674	1	3a	–8.44	N
GJ 849	1	3a	8.20	D
GJ 86	1	2	–1554.76	D
HAT-P-6	1	1	1450.29	N

Table A3 – continued

System	Number of candidates from literature	Visual quality flag	R	Flag for period prior used
HAT-P-8	1	3b	–	N
HAT-P-9	1	3b	0.95	D
HD 101930	1	3b	6.12	N
HD 108874	2	2	–8.23	N
HD 11506	2	2	–3.36	N
HD 118203	1	1	80.12	D
HD 12661	2	2	–15.22	D
HD 128311	2	2	–32.30	D
HD 131664	1	1	3.34	D
HD 132406	1	1	234.41	N
HD 142	1	3a	–5.87	D
HD 142022	1	3b	0.43	N
HD 149143	1	3b	1.49	D
HD 154345	1	1	44.33	N
HD 155358	2	2	–2.90	N
HD 162020	1	3b	102.59	D
HD 168443	2	2	–994.46	D
HD 169830	2	3b	0.25	D
HD 171028	1	3a	18.72	N
HD 183263	2	2	–187.35	D
HD 185269	1	3b	0.83	N
HD 187123	2	3b	–29.06	D
HD 189733	1	3a	–140.98	D
HD 190228	1	3b	0.08	N
HD 190360	2	3b	0.04	D
HD 190647	1	3b	5.01	N
HD 195019	1	1	771.62	N
HD 202206	2	3b	–4.59	D
HD 20868	1	2	–252.94	D

Table A3 – *continued*

System	Number of candidates from literature	Visual quality flag	R	Flag for period prior used
HD 209458	1	3b	−0.03	D
HD 212301	1	1	5.15	N
HD 217107	2	3a	−59.88	D
HD 219828	1	3a	−352.73	D
HD 221287	1	1	141.08	N
HD 224693	1	3b	20.92	N
HD 23127	1	1	69.47	N
HD 2638	1	3b	−0.97	N
HD 27442	1	3b	10.37	N
HD 27894	1	1	53.60	N
HD 28185	1	1	10.27	D
HD 285968	1	1	0.42	N
HD 330075	1	3b	745.43	D
HD 33636	1	1	104.05	D
HD 3651	1	3b	−0.06	N
HD 38529	2	2	−41.83	D
HD 4203	1	3a	4.92	N
HD 4208	1	3b	0.65	N
HD 43691	1	3b	5.07	N
HD 43848	1	1	−	N
HD 46375	1	3b	−0.04	D
HD 47536	2	1	17.15	D
HD 49674	1	3b	0.24	N
HD 50499	1	2	−6.74	D
HD 5319	1	2	−12.72	D
HD 63454	1	3b	2.57	N
HD 68988	1	2	−100.51	D
HD 73267	1	3b	7.60	D
HD 73526	2	2	−7.43	N
HD 74156	2	2	−241.14	D

Table A3 – *continued*

System	Number of candidates from literature	Visual quality flag	R	Flag for period prior used
HD 75289	1	1	0.14	N
HD 75898	1	3a	28.79	N
HD 76700	1	1	14.28	N
HD 80606	1	1	4.22	D
HD 81040	1	3b	−1.13	N
HD 82943	2	2	−319.74	N
HD 8574	1	1	−0.03	N
HD 86081	1	1	0.19	D
HD 89307	1	3b	−	D
HD 93083	1	3b	32.00	N
HR 810	1	3b	8.71	N
κ CrB	1	1	8.87	N
NGC 2423 3	1	3b	52.75	N
NGC4 349 127	1	3b	24.85	N
τ Boo	1	3b	−8.90	D
TrES-3	1	2	−	D
WASP-2	1	3b	−	D
WASP-3	1	3b	−	D
WASP-4	1	3b	−33.89	N
WASP-5	1	3b	−	N
XO-1	1	1	−	D
XO-2	1	3b	−	D
XO-3	1	1	20.24	N
XO-4	1	3b	−	D

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.